

# How creating one additional well can generate Bose-Einstein condensation

Mihály Máté,<sup>1,2</sup> Örs Legeza,<sup>1</sup> Rolf Schilling,<sup>3</sup> Mason Yousif,<sup>4</sup> and Christian Schilling<sup>5,4,\*</sup>

<sup>1</sup>*Wigner Research Centre for Physics, Hungarian Academy of Sciences,  
Konkoly-Thege Miklós út 29-33, 1121 Budapest, Hungary*

<sup>2</sup>*Department of Physics of Complex Systems, Eötvös Loránd University, Pf. 32, H-1518 Budapest, Hungary*

<sup>3</sup>*Institut für Physik, Johannes Gutenberg-Universität, D-55099 Mainz, Germany*

<sup>4</sup>*Clarendon Laboratory, University of Oxford, Parks Road, Oxford OX1 3PU, United Kingdom*

<sup>5</sup>*Department of Physics, Arnold Sommerfeld Center for Theoretical Physics,  
Ludwig-Maximilians-Universität München, Theresienstrasse 37, 80333 München, Germany*

(Dated: July 26, 2021)

## Abstract

The realization of Bose-Einstein condensation in ultracold trapped gases has led to a revival of interest in that fascinating quantum phenomenon. This experimental achievement necessitated both extremely low temperatures and sufficiently weak interactions. Particularly in reduced spatial dimensionality even an infinitesimal interaction immediately leads to a departure to quasi-condensation. We propose a system of strongly interacting bosons which overcomes those obstacles by exhibiting a number of intriguing related features: (i) The tuning of just a single control parameter drives a transition from quasi-condensation to complete condensation, (ii) the destructive influence of strong interactions is compensated by the respective increased mobility, (iii) topology plays a crucial role since a crossover from one- to ‘infinite’-dimensionality is simulated, (iv) a ground state gap opens which makes the condensation robust to thermal noise. Remarkably, all these features can be derived by analytical and exact numerical means despite the non-perturbative character of the system.

## Introduction

Bose-Einstein condensation (BEC) is one of the most striking quantum phenomena in nature [1–4]. While its theoretical prediction dates back almost one hundred years ago it has more recently seen a revival of interest due to its realization in trapped gases [5–7]. The accurate study of BEC by theoretical and computational approaches particularly for systems with strong quantum correlations is rather challenging. This has been the reason why most studies of BEC so far were concerned with weakly interacting bosons (corresponding to the experimental situation for ultracold gases) or even ideal Bose gases, eventually allowing for feasible mean-field approaches. Prime examples are the Bogoliubov theory [8] for uniform systems, Gross-Pitaevskii theory [9–11] for general inhomogeneous systems, and perturbation theoretical approaches [12–18]. Although these widely used approaches have led to a deeper understanding of BEC, their range of validity is limited. To go beyond that limitation, various methods were developed [19–22].

Since the experimental realization of BEC, the respective field of ultracold gases has become one of the most exciting fields of research with a fruitful interplay between theory and experiment. It allowed for the experimental verification of numerous other theoretical predictions as well, stimulated further theoretical investigations of trapped particles [19] and even revealed phenomena not observed before such as the crossover from BEC-superfluidity to BCS-superconductivity [23–26]. One of the most promising recent avenues has been the study of effectively one-dimensional quantum systems [27–33]. Their most striking difference to three-dimensional systems is probably the absence of BEC: Already an infinitesimally weak interaction between the  $N$  bosons leads to a

“sublinear” behaviour of the number of condensed bosons,  $N_0(N) \sim N^\alpha$  [34, 35], even at zero temperature, for homogeneous gases as well for gases in a harmonic trap and regardless of the form of the interaction [35–42]. A prominent system giving rise to this phase called “quasi-condensation” [35] is the Lieb-Liniger model [43, 44], a ring system with  $N$  spinless bosons interacting via a  $\delta$ -potential. Tuning the coupling constant to infinity leads to impenetrable bosons (Tonk-Girardeau gas) [45] with the proven scaling  $N_0(N) \sim \sqrt{N}$  [34].

Thermodynamic phase transitions (at finite temperatures) in  $D = 3$  dimensions have been studied for more than a century. However, the study of quantum phase transitions (at zero temperature) [46], and particularly of the entanglement close to that transition [47, 48] have attracted much attention in recent years, only. The latter studies were performed mostly for low-dimensional lattice models. They have revealed a striking similarity between the behavior of the order parameter and of quantum informational quantities, like entanglement entropy. As discussed above, at zero-temperature an interacting Bose gas exhibits two qualitatively different phases, a quasi-condensate in  $D = 1$  and a true BE-condensate in  $D \geq 3$ . Therefore, it is of interest to search for a model which exhibits a transition (or a crossover) between these two phases, and in particular allows to check whether this special transition has common properties with general quantum phase transitions.

BEC was explored in cylindrical or toroidal trap geometries, both experimentally [27–33] and theoretically [49–51]. But, changing the radial dimension of the confinement, neither the transition from the sublinear  $N$ -dependence of  $N_0(N)$  of the quasi-condensate to the linear dependence of the true condensate, nor its entanglement properties have been investigated. The only systematic study of such a transition was performed for a one-dimensional Bose gas in a harmonic trap [52]. However, that transition occurs only at temperatures  $T > 0$ .

\* c.schilling@physik.uni-muenchen.de

It is the challenge of the present work to propose and investigate a lattice model for strongly interacting bosons which allows one to drive such a transition by changing just a single parameter,  $s/t$ , which is the ratio of the model's two hopping rates  $s$  and  $t$ , as explained below. One of our major results is to establish by this model a mechanism which can generate ‘infinite’ range hopping by increasing  $s/t$ . This is important since enhancing the boson's mobility allows to overcome the destructive effects of the repulsive interactions, leading finally to maximal possible condensation, despite infinitely strong repulsion. A further important feature of our model is the generation of an excitation gap in the  $N$ -particle spectrum for  $s/t > 0$ . This makes BEC even robust to thermal noise and quantum fluctuations and thus may allow experimentalists to overcome the typical obstacles faced while realizing BEC. The other important result concerns the application of tools from quantum information theory. We show that the mutual information possesses the qualitatively similar dependence on  $s/t$  as the number  $N_0(N)$  of condensed bosons. This supports the connection between the behavior of an order parameter and of entanglement at a quantum phase transition even for the transition(or crossover) from a quasi-condensate to a true one.

All these key findings will be derived by analytical or exact numerical means despite the nonperturbative character of our system.

## Results

**Model Hamiltonian.** To motivate our model, let us first recall that the possible presence of BEC depends in general not only on the spatial dimensionality and temperature but also on the ratio between kinetic and interaction energy. In case of systems which are inhomogeneous, e.g., due to the presence of an external field or disorder, the occurrence of BEC will also depend on these quantities. Concerning the ratio between kinetic and interaction energy, lattice systems have the great advantage that the kinetic energy can be manipulated by varying the hopping range between the lattice sites. The most prominent lattice model for bosons is the widely studied Bose-Hubbard model [53]

$$\hat{H} = - \sum_{i,j} t_{ij} b_i^\dagger b_j + U \sum_{i=1} \hat{n}_i (\hat{n}_i - 1), \quad (1)$$

where  $b_i^\dagger, b_i$  creates/annihilates a spinless boson at site  $i$ ,  $\hat{n}_i \equiv b_i^\dagger b_i$  and  $t_{ij}$  is the hopping rate between sites  $i$  and  $j$ . It was shown that the Bose-Hubbard model can be experimentally realized by ultracold bosonic atoms in an optical lattice [54].

The conflict between interaction and mobility is maximized in the limit of strong interactions  $U \rightarrow \infty$  in which the bosons become hard-core[55, 56]. By employing respective hard-core boson(HCB) creation( $h_i^\dagger$ ) and annihilation operators( $h_i$ ) (1) takes the compact form  $\hat{H}_{hc} = - \sum_{i,j} t_{ij} h_i^\dagger h_j$ . Particularly the case of HCBs makes clear the important role of the hopping range, since for infinite-range hopping(a kind of mean field limit [53]) HCBs exhibit BEC even at finite temperatures, despite their infinitely strong repulsion [57–59].

Moreover, the effect of the interaction on BEC is distinctively destructive in one-dimensional systems. At zero tem-




FIG. 1. **Topology of the lattice.** An interpolation between the 1D regime and the ‘star’ through the ‘wheel’ .

perature even an infinitesimally weak interaction already leads to a departure from BEC to the phase of quasi-condensation. This raises a fundamental question which our work shall answer in an affirmative and constructive way: After having confined a 3D Bose gas to one dimension, is it possible to tweak in an experimentally feasible way this one-dimensional system with the effect of enhancing the mobility of the interacting bosons to reintroduce BEC? From a general point of view, one is immediately tempted to negate this question. The hopping amplitudes  $t_{ij}$  namely resemble the overlap of Wannier orbitals at sites  $i, j$  which in turn decays exponentially as function of the spatial separation  $|i - j|$ . Screening effects reduce the hopping even further and eventually motivate the common restriction of  $t_{ij}$  in the Bose-Hubbard model to just nearest neighbors. The potential physical significance of long-range hopping has motivated experimentalists in recent years to realize at least effectively hopping terms beyond nearest neighbors. Despite a remarkable effort, the regime of infinite-range hopping has been out of reach but only the typical decay of  $t_{ij}$  could be slowed down to an algebraic dipolar- and van der Waals-type one [60, 61]. It will be one of our key achievements to propose a model which eventually would allow one to enhance the mobility even to infinite-range.

In contrast to the rather involved experimental realisation of algebraically decaying hopping rates our proposal to realize ‘infinite’-range hopping will be surprisingly simple. As it is illustrated in Fig. 1, we consider  $N$  HCBs on a lattice consisting of a ring with  $d$  sites, lattice constant  $a$  and *one* additional site at its center. The ring gives rise to a hopping between nearest neighbors at a rate  $t > 0$ . The crucial point is now that the ring's topology allows hopping between the central site and any ring site at a rate  $s \geq 0$ . Accordingly, the central site has an effect similar to an impurity, making the lattice inhomogeneous.

We remind the reader that proposing and studying this model shall be seen as one of our key achievements. It is also worth noticing that various other studies of BEC for inhomogeneous lattices differ significantly from ours. They either consider the rather trivial case of ideal bosons [62–68] or restrict to the mean field regime [69, 70]. At the same time, our model could be particularly appealing to experimentalists since the underlying graph emerges from a Mexican hat potential(see below) and HCBs can be realized experimentally [31, 71] by tuning the interactions at the Feshbach resonance [72–75].




FIG. 2. **Schematic representation of the unperturbed and perturbed spectrum for  $n \leq 1/2$ .** (a) The unperturbed band spectrum of  $(N - 1)$  hard-core bosons (red dashed lines) and  $N$  hard-core bosons (blue solid lines). The crosses mark those levels which disappear under the perturbation. (b) The perturbed spectrum consisting of the shifted unperturbed levels (dashed red and solid blue lines) and the crosses mark those levels of the unperturbed  $(N - 1)$ -particle spectrum which have disappeared. The open circles are the new levels generated by the perturbation.

Accordingly, the Hamiltonian of our proposed model of bosons with hard-core interaction reads

$$\hat{H} = -t \sum_{i=1}^d [h_i^\dagger h_{i+1} + h_{i+1}^\dagger h_i] - s \sum_{i=1}^d [h_i^\dagger h_c + h_c^\dagger h_i], \quad (2)$$

where  $h_c^\dagger, h_c$  denote the corresponding operators for the central site. For  $s/t \rightarrow 0$ ,  $\hat{H}$  reduces to the pure ring-model (left of Fig. 1) and the limit  $s/t \rightarrow \infty$  leads to the star-model (right of Fig. 1). The solution of the eigenvalue problem for these two limiting cases is known. For  $s = 0$  it follows from the solution for impenetrable bosons [43–45] which only exhibits quasi-condensation, and  $s = \infty$  was solved in Ref. [76] proving the existence of true BEC with maximal possible number  $N_0(N, d) = N(d - N + 1)/d$  of condensed bosons. For finite values of  $s/t$  the Hamiltonian (2) interpolates between the ring-lattice and the star-lattice (cf. Fig. 1). Hence, changing the single parameter  $s/t$  allows us to investigate in a systematic way the crossover from the regime of quasi-condensation to maximally possible condensation, eventually leading to a number of remarkable insights.

**Spectral properties, BEC and entanglement.** The present section contains only the crucial steps. Technical details can be found in the section ‘Methods’ and particularly in the ‘Supplementary Methods’.

Since the central site couples to the  $(N - 1)$ - and  $N$ -particle state space on the ring, a simple and fully analytic solution does not exist. Yet, after implementing a number of steps, the eigenvalue problem for Hamiltonian (2) can be rewritten as

$$1 = s^2 F_d^{(N)}(E; \{A_\nu\}), \quad (3)$$

where  $E$  is the eigenvalue and  $\{A_\nu\}$  are amplitudes of the unperturbed (i.e., corresponding to  $s = 0$ )  $N$ -particle eigenstates  $|\psi_\nu^0(N)\rangle$  on the ring. Although this equation cannot

be solved analytically for the entire regime of  $s$ , it allows us to derive in a nontrivial way important qualitatively correct features of the spectrum. The unperturbed  $(N - 1)$ - and  $N$ -particle spectrum forms a band of discrete levels (see Fig. 2a) which becomes continuous for  $d \rightarrow \infty$ . The hopping between the central site and the ring introduces a ‘hybridization’ of these two spectra leading on one hand to a shift of order  $1/d$  of the unperturbed band-levels. On the other hand, some energy levels (marked by crosses) of the smaller  $(N - 1)$ -particle band (assuming  $n = N/d < 1/2$ , which is not a restriction due to the particle-hole duality) are found to disappear. These levels, however, reappear as new discrete eigenvalues symmetrically below and above the perturbed  $N$ -particle band (see open circles in Fig. 2b). The larger  $s$  and  $N$ , the more of those new discrete energy levels occur. As a matter of fact, they follow from the eigenvalues of an effective Hamiltonian for  $N$  HCBs with ‘infinite’-range hopping:  $\hat{H}^{\text{eff}} = \tilde{s}^2 (1/d) \sum_{i,j=1}^d h_i^\dagger h_j$ . Here, the parameter  $\tilde{s} = (s/t)\sqrt{d}$  is a scaled dimensionless hopping rate. This mapping of the original model to an effective one holds for  $\tilde{s} \gg 2\sqrt{2} \pi/\sqrt{d}$  for the diluted gas ( $n \ll 1$ ) and in case of finite  $n$  for  $\tilde{s} \gg (4/\pi)\sqrt{d} \sin \pi n/\sqrt{n(1-n)}$ .

Most importantly, these findings imply also the opening of an energy gap  $\Delta E = E_{\text{low}}^0 - E_0$  between the perturbed ground state energy  $E_0$  and  $E_{\text{low}}^0$ , the lower edge of the  $N$ -particle band:

$$\Delta E \simeq |E_F/2| \left[ \sqrt{(1 + \tilde{s}^2 d n(1-n)/(E_F/2t)^2) - 1} \right]. \quad (4)$$

Also the number  $N_0$  of condensed HCBs can be derived analytically since it is related to the largest eigenvalue of  $H^{\text{eff}}$ . We obtain

$$N_0 \simeq N \left[ (1-n) - |\beta|^2 (1-2n)N^{-1} \right], \quad (5)$$

where the prefactor  $|\beta|^2$  of the  $1/N$ -correction is given in the Supplementary Eq. S30.

In order to support these analytical results and to extend those for finite  $d$  to small and intermediate values of  $\tilde{s}$  we have performed large-scale density matrix renormalization group computations (DMRG) [77–79]. The corresponding results together with the analytical ones are presented in Figs. 3 and 4. The log-log representation of the gap  $\Delta E(\tilde{s})$  in Fig. 3 reveals a distinctive crossover from a  $\tilde{s}^2$ -dependence for  $\tilde{s} \ll 1$  to the linear dependence on  $\tilde{s}$  for  $\tilde{s} \gg 1$ . For the diluted gas, i.e.  $n \ll 1$ , the analytical and DMRG results in the  $\tilde{s}^2$ - and  $\tilde{s}$ -regime are in good agreement. When the density is increased this agreement remains excellent in the linear regime while it gets worse in the complementary range. Fig. 4 illustrates clearly for the diluted gas (Fig. 4a) and for higher densities (Fig. 4b) the crossover from a quasi-condensate with  $N_0 \sim \sqrt{N}$  to the maximally possible condensation  $N_0(N, n) \simeq N(1-n)$ . The deviations from the  $\sqrt{N}$ -dependence for small  $\tilde{s}$  and higher densities (see lower panel) is an effect of the lattice-discreteness. In the regime in which the mapping to the effective Hamiltonian is valid (see above) the analytical and DMRG results agree well.

To explore a possible relation between BEC and the entanglement structure of the ground state we have used DMRG for




FIG. 3. **Excitation gap.** log-log plot for the excitation gap as a function of  $\tilde{s}$  for  $d = 199$  and various filling factors  $n$ . Results from density matrix renormalization group calculations (symbols) and analytical result (Eq.(4)) (solid lines). Dashed-dotted and dashed line represent slope two and one, respectively.




FIG. 4. **Number of condensed bosons.** (a)  $N_0$  as a function of  $\log(\tilde{s})$  for fixed low density  $n \simeq 0.05$  and various site numbers  $d$ . (b)  $N_0$  as a function of  $\log(\tilde{s})$ , various numbers of particles  $N$ , and fixed number of sites  $d = 199$ . The dots on the vertical axis represent  $N(1 - n)$ . Symbols: Results from density matrix renormalization group calculations, dashed lines: guide for the eye, solid lines: analytical result (Supplementary equation S28). The dotted lines mark the asymptotic values for finite  $N$ ,  $s = 0$ ,  $d \rightarrow \infty$  obtained by the exact numerical calculation of a Toeplitz determinant [39].

calculating the mutual information between the central site  $c$  and any ring site  $i$  ( $I_{i|c}$ ) and between two  $\ell$ -th nearest neighbor ring sites ( $I_{i|i+\ell}$ ) (see Methods). The corresponding results for  $d = 199$  and  $n \simeq 0.05$  are shown in Fig. 5. The change in the respective pattern related to the crossover from quasi-BEC to genuine BEC is clearly visible through the mutual information, as well. The correlation between the central and any ring site,  $I_{i|c}$ , vanishes for  $\tilde{s}$  small while it saturates to a finite value in the limit of large  $\tilde{s}$  when the model exhibits ‘infinite’-range hopping.  $I_{i|i+\ell}$  saturates also with increasing  $\tilde{s}$  to a constant value for all  $\ell$  demonstrating the growth of long range correlations. This relates to the generation of BEC. For  $\tilde{s} = 0$ ,  $I_{i|i+\ell}$  decays algebraically with increasing  $\ell$  which reflects the al-




FIG. 5. **Quantum informational quantity.** Two-site correlation measured in terms of the mutual information  $I_{i|c}$  between the central site and a ring site (red stars), and  $I_{i|i+\ell}$  for two sites on the ring separated by distance  $\ell = 1, 3, 10, 99$  for  $d = 199$  and  $n \simeq 0.05$  (other symbols). The dashed and dotted lines are a guide for the eye.

gebraic dependence of the quasi-condensate on  $N$ . Whereas for finite values of  $\tilde{s}$  its decay becomes exponential as the gap opens, and saturates to finite value for very large  $\ell$  values.

**Potential experimental realization.** As a possible experimental realization of our model (2) we propose in a first step to confine  $N$  ultracold bosonic atoms into two dimensions subject to a Mexican-hat-type potential  $V(x, y)$  with  $d$  local wells (Fig. 6a) in complete analogy to several recent years’ experiments [80–84]. Then, one may tune the interaction at the Feshbach resonance to realize HCBs in the same way as reported in Ref. [31] for a cigar-shaped confinement to realize quasi-condensation of HCB with  $N_0(N) \propto N^{1/2}$ . Next, creation of a local well at the hat’s center (Fig. 6b) and increasing its depth more and more would strongly enhance the mobility of the HCBs due to their possible transitions back and forth between any ring-well and the central one. This would significantly change the physical behavior and BEC would occur with  $N_0(N) \sim N$ . In order for this to happen already for *finite*  $d$  it must be  $s/t \gg 2\sqrt{2}\pi/d$  in case of a diluted gas (see previous section) which is the regime relevant for ultracold gases. The hopping occurs due to tunneling between the corresponding wells. Let  $(V_r, l_r = a)$  and  $(V_c, l_c = ad/(2\pi))$  denote the potential barrier and tunneling distance, respectively, between two adjacent ring-wells and between a ring-well and the central one. Use of the WKB tunneling rate yields the estimate  $s/t \approx (\gamma_c/\gamma_r) \exp[-\sqrt{ma^2/\hbar^2}(\sqrt{V_c d}/(2\pi) - \sqrt{V_r})]$  with  $m$  the particle’s mass and  $\gamma_\alpha, \alpha = c, r$  the so-called attempt frequency related to the zero-point oscillation frequency in the corresponding well. For instance, if  $d = 79$  and  $N = 4$  (one data set in Fig. 4(a)) ‘BEC’-like behavior should occur for  $s/t > 1$ . This can be satisfied if  $V_c/V_r \approx (2\pi/d)^2$  or if  $a$  compared to  $\hbar/\sqrt{m} \max\{V_c, V_r\}$  is small enough, provided  $\gamma_c/\gamma_r \approx 1$ .

If the trap potential in Fig. 6 is chosen such that it represents a good experimental realization of the ‘wheel’ lattice (cf. Fig. 1) there is true condensation for sufficiently




FIG. 6. **Mexican-hat-type trap potential.** (a) Realization of the ring lattice for  $d = 10$  by a Mexican-hat-potential. (b) Realization of the wheel lattice for  $d = 10$  by a Mexican-hat-like potential with a local well at its center. Loading hard-core bosons into the potential landscape in (a) and creating a local well as shown in (b) generates a crossover from quasi to complete Bose-Einstein condensation.

large  $s/t$ . In particular, since only a single one-particle state (zero-momentum state) is macroscopically occupied, no fragmented condensation exists per definition. This is consistent with the expectation that homogeneous bosonic systems with purely repelling pair interactions do not exhibit fragmented condensation [4]. Although the presence of the central well (central site) makes the system inhomogeneous it can not generate fragmentation, because it accommodates maximally one HCB, only. But increasing the width of the central well in Fig. 6b such that it can accommodate a “macroscopic number of bosons of an ultracold gas, a situation similar to the double-well-like trap potential in one dimension occurs [85]. As shown in that work, fragmented condensation may then occur if the barrier height of the double well is high enough.

It is worth noticing that according to the DMRG results (see also Fig. 4) one would not need to realize a macroscopically large ring to observe our crossover. Yet, in case experimentalists could even realise our model with a huge number  $d$  of sites on a ring of fixed size (i.e., the limit  $d \rightarrow \infty, a \rightarrow 0$  with  $ad$  fixed) this would generate a *true* Mexican-hat potential with *continuous* rotational invariance and the HCBs would become a Tonks-Girardeau gas. Again, creating a central well would generate genuine Bose-Einstein condensation.

## Discussions

We proposed and comprehensively studied a physical model of strongly interacting bosons which allows one to drive a non-trivial transition from quasi-condensation to maximal Bose-Einstein condensation (BEC). It is particularly appealing that this necessitates the tuning of just a single control parameter which changes the underlying topology in such a distinctive way that the ‘infinite’ range hopping model is simulated. The enhanced mobility of the bosons then compensates for the destructive effects of the strong interaction to generate BEC. Without solving the model’s eigenvalue equation exactly, our kind of analytical approach (see the section “Spectral properties, BEC and entanglement” above and also the ‘Supplementary Methods’) allows us to show on a qualitative level why an excitation gap occurs in the  $N$ -particle spectrum, which usually is highly demanding. Similarly to, e.g., superconductivity, the quantum Hall effect and the Haldane phase the exist-

tence of such a gap has an enormous influence on the physical behavior, e.g., making the BE-condensate robust to thermal noise and perturbations in general.

It is worth highlighting the striking potential of our mechanism for generating BEC. As a matter of fact, it is conceptually quite different to the well-known generation of BEC at finite temperatures for non-interacting bosons. The latter is either merely due to the opening of a gap in the “one-particle” spectrum or a deformation of the density of states (in analogy to the transition from  $D \leq 2$  to  $D = 3$ ) [62–68]. The same effectively applies to the experimental [27–33] and theoretical studies [49–51] in which the cylindrical or toroidal confinement is relaxed to reach the mean-field regime. In our system, however, it is the interplay between mobility and interaction within the “non-perturbative” regime which generates genuine BEC (see the rather involved derivation in the Supplementary Methods). The non-trivial influence of the interaction is also well-illustrated by the analytical result for the ground state gap (Eq. (4)) which in the regime of maximal BEC differs from the one of non-interacting bosons by the crucial factor  $\sqrt{1-n}$ . Remarkably,  $1-n$  is nothing else than the universal reduction of the maximal possible degree of condensation due to the hard-core constraint [76], which is the quantum depletion. In case of *finite* on-site interactions, this depletion factor  $\nu(n)$  is expected to interpolate between both extremal cases of hard-core and ideal bosons,  $1-n \leq \nu(n) \leq 1$ . This would provide a remarkable exact relation between the ground state gap, quantum depletion and the interaction strength of the ultracold atoms. Since the latter can systematically be tuned at the Feshbach resonance [72–75] this would open an avenue for steering ground state gaps and controlling the number of bosons in BEC.

Finally, inspired by the fruitful interplay of theory and experiments in the field of ultracold gases our work based on analytical and exact large scale DMRG calculations shall be understood as a proposal to the experimentalists as well. Our model could be particularly appealing since the underlying graph emerges from a Mexican-hat-type potential and the entire transition can be driven by tuning just a single control parameter. It is then exactly the respective central site which can be probed to confirm that transition. At the same time, this would also exploit the fruitful link [47, 48] between quantum phase transitions and entanglement or related promising quantum informational theoretical concepts, as illustrated in Fig. 5.

## Methods

**Eigenvalue problem.** The central site generates for the  $N$ -particle state a superposition  $|\Psi_N\rangle = \alpha|\phi_N\rangle_r \otimes |0\rangle_c + \beta|\varphi_{N-1}\rangle_r \otimes |1\rangle_c$  of a  $N$ - and  $(N-1)$ -particle ring-state. Expansion of these states with respect to the unperturbed  $(N-1)$ - and  $N$ -particle ring-states allows to decouple the original eigenvalue problem. This leads to a nonintegrable eigenvalue problem on the ring itself. Straightforward manipulation allows to derive Eq. (3). For details see the Supplementary Methods.

**Density matrix renormalization group.** The DMRG calculations were performed for  $d \leq 199$  and  $N \leq 98$ . In the DMRG procedure we have performed calculations using the dynamic block state selection approach [86]. We have set a tight error bound on the diagonalization procedure, i.e., we set the residual error of the Davidson method to  $10^{-9}$  and used ten DMRG sweeps. We have checked that the various quantities of interest are practically insensitive on the bond dimension being larger than 1024.

Besides calculating energy eigenvalues and the one- $(\rho_i)$  and two-site $(\rho_{ij})$  reduced density matrices we have also determined one- and two-site von Neumann entropies  $s_i$  and  $s_{ij}$ , respectively, as well as the two-site mutual information,  $I_{i|j}$ , given as  $I_{i|j} = s_i + s_j - s_{ij}$  [87, 88]. Here  $s_i = -\text{Tr}\rho_i \ln \rho_i$  and  $s_{ij} = -\text{Tr}\rho_{ij} \ln \rho_{ij}$ .

## Acknowledgements

We gratefully acknowledge critical comments on the present manuscript by P. van Dongen, F. Gebhard, J. Marino and L. Pollet. We also would like to thank P.J. Forrester and T.M. Garoni for providing the exact results obtained directly from the Toeplitz determinant for the number of condensed particles in one dimension for small  $N$ . This work has been supported in part by the Hungarian National Research, Development and Innovation Office (grant no. K120569 and K134983), and the Hungarian Quantum Technology National Excellence Program (project no. 2017-1.2.1-NKP-2017-00001). Ö.L. acknowledges financial support from the Alexander von Humboldt foundation. M.M. has been supported by the ÚNKP-19-3 Hungarian New National Excellence Program of the Ministry for Innovation and Technology. The development of the DMRG libraries was supported by the Center for Scalable and Predictive methods for Excitation and Correlated phenomena (SPEC), which is funded from the Computational Chemical Sciences Program by the U.S. Department of Energy (DOE), at Pacific Northwest National Laboratory. C.S. acknowledges financial support from the UK Engineering and Physical Sciences Research Council (Grant EP/P007155/1) and Deutsche Forschungsgemeinschaft (Grant SCHI 1476/1-1).

## References

- [1] A. Griffin, D. W. Snoke, and S. Stringari, *Bose-Einstein Condensation* (Cambridge University Press, Cambridge, 1995).
- [2] C. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases* (Cambridge University Press, Cambridge, 2002).
- [3] A. Leggett, *Quantum Liquid: Bose Condensation and Cooper Pairing in Condensed-Matter Systems* (Oxford University Press, Oxford, 2006).
- [4] L. P. Pitaevskii and S. Stringari, *Bose-Einstein Condensation and Superfluidity* (Oxford University Press, Oxford, 2016).
- [5] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, Observation of Bose-Einstein condensation in a dilute atomic vapor, *Science* **269**, 198 (1995).
- [6] C. C. Bradley, C. A. Sackett, J. J. Tollett, and R. G. Hulet, Evidence of Bose-Einstein Condensation in an Atomic Gas with Attractive Interactions, *Phys. Rev. Lett.* **75**, 1687 (1995).
- [7] K. B. Davis, M.-O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, Bose-Einstein condensation in a gas of Sodium atoms, *Phys. Rev. Lett.* **75**, 3969 (1995).
- [8] N. N. Bogoliubov, On the theory of superfluidity, *J. Phys. USSR* **11**, 23 (1947).
- [9] E. Gross, Structure of a quantized vortex in boson systems, *Nuovo Cimento* **20**, 454 (1961).
- [10] E. Gross, Hydrodynamics of a superfluid condensate, *J. Math. Phys.* **4**, 195 (1963).
- [11] P. L. P., Vortex lines in an imperfect bose gas, *Sov. Phys. JETP* **13**, 451 (1961).
- [12] T. D. Lee and C. N. Yang, Many-Body Problem in Quantum Mechanics and Quantum Statistical Mechanics, *Phys. Rev.* **105**, 1119 (1957).
- [13] T. D. Lee, K. Huang, and C. N. Yang, Eigenvalues and Eigenfunctions of a Bose System of Hard Spheres and Its Low-Temperature Properties, *Phys. Rev.* **106**, 1135 (1957).
- [14] K. A. Brueckner and K. Sawada, Bose-Einstein Gas with Repulsive Interactions: General Theory, *Phys. Rev.* **106**, 1117 (1957).
- [15] K. A. Brueckner and K. Sawada, Bose-Einstein Gas with Repulsive Interactions: Hard Spheres at High Density, *Phys. Rev.* **106**, 1128 (1957).
- [16] S. T. Beliaev, Energy-Spectrum of a Non-ideal Bose gas, *Sov. Phys. JETP* **34**, 299 (1958).
- [17] N. M. Hugenholtz and D. Pines, Ground-State Energy and Excitation Spectrum of a System of Interacting Bosons, *Phys. Rev.* **116**, 489 (1959).
- [18] E. H. Lieb, Simplified Approach to the Ground-State Energy of an Imperfect Bose Gas, *Phys. Rev.* **130**, 2518 (1963).
- [19] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari, Theory of Bose-Einstein condensation in trapped gases, *Rev. Mod. Phys.* **71**, 463 (1999).
- [20] J. O. Andersen, Theory of the weakly interacting Bose gas, *Rev. Mod. Phys.* **76**, 599 (2004).
- [21] M. A. Cazalilla, R. Citro, T. Giamarchi, E. Orignac, and M. Rigol, One dimensional bosons: From condensed matter systems to ultracold gases, *Rev. Mod. Phys.* **83**, 1405 (2011).
- [22] A. U. J. Lode, C. Lévêque, L. Bojer Madsen, A. I. Streltsov, and O. E. Alon, Multiconfigurational time-dependent Hartree approaches for indistinguishable particles, *Rev. Mod. Phys.* **92**, 011001 (2020).
- [23] M. Greiner, C. A. Regal, and D. S. Jin, Emergence of a molecular Bose-Einstein condensate from a Fermi gas, *Nature* **426**, 537 (2003).
- [24] M. Bartenstein, A. Altmeyer, S. Riedl, S. Jochim, C. Chin, J. Hecker Denschlag, and R. Grimm, Crossover from a Molecular Bose-Einstein Condensate to a Degenerate Fermi Gas, *Phys. Rev. Lett.* **92**, 120401 (2004).
- [25] M. W. Zwierlein, C. A. Stan, C. H. Schunck, S. M. F. Raupach, A. J. Kerman, and W. Ketterle, Condensation of Pairs of Fermionic Atoms near a Feshbach Resonance, *Phys. Rev. Lett.* **92**, 120403 (2004).
- [26] T. Bourdel, L. Khaykovich, J. Cubizolles, J. Zhang, F. Chevy, M. Teichmann, L. Tarruell, S. J. J. M. F. Kokkelmans, and C. Salomon, Experimental Study of the BEC-BCS Crossover Region in Lithium 6, *Phys. Rev. Lett.* **93**, 050401 (2004).
- [27] M. Greiner, I. Bloch, O. Mandel, T. W. Hänsch, and T. Esslinger, Exploring Phase Coherence in a 2D Lattice of Bose-Einstein Condensates, *Phys. Rev. Lett.* **87**, 160405 (2001).
- [28] S. Dettmer, D. Hellweg, P. Ryytty, J. J. Arlt, W. Ertmer, K. Sengstock, D. S. Petrov, G. V. Shlyapnikov, H. Kreutzmann, L. Santos, and M. Lewenstein, Observation of Phase Fluctuations in Elongated Bose-Einstein Condensates, *Phys. Rev. Lett.*

- 87**, 160406 (2001).
- [29] A. Görlitz, J. M. Vogels, A. E. Leanhardt, C. Raman, T. L. Gustavson, J. R. Abo-Shaeer, A. P. Chikkatur, S. Gupta, S. Inouye, T. Rosenband, and W. Ketterle, Realization of Bose-Einstein Condensates in Lower Dimensions, *Phys. Rev. Lett.* **87**, 130402 (2001).
- [30] C. Orzel, A. K. Tuchman, M. L. Fenselau, M. Yasuda, and M. A. Kasevich, Squeezed States in a Bose-Einstein Condensate, *Science* **291**, 2386 (2001).
- [31] B. Paredes, A. Widera, V. Murg, O. Mandel, S. Fölling, I. Cirac, G. V. Shlyapnikov, T. W. Hänsch, and I. Bloch, Tonks-Girardeau gas of ultracold atoms in an optical lattice, *Nature* **429**, 277 (2004).
- [32] T. Kinoshita, T. Wenger, and D. S. Weiss, Observation of a One-Dimensional Tonks-Girardeau Gas, *Science* **305**, 1125 (2004).
- [33] T. Stöferle, H. Moritz, C. Schori, M. Köhl, and T. Esslinger, Transition from a Strongly Interacting 1D Superfluid to a Mott insulator, *Phys. Rev. Lett.* **92**, 130403 (2004).
- [34] A. Lenard, Momentum distribution in the ground state of the one-dimensional systems of impenetrable bosons, *J. Math. Phys.* **5**, 930 (1964).
- [35] V. N. Popov, On the theory of the superfluidity of two- and one-dimensional bose systems, *Theor. Math. Phys.* **11**, 565 (1972).
- [36] H. Widom, Toeplitz Determinants with Singular Generating Functions, *Am. J. Math.* **95**, 333 (1973).
- [37] M. Schwartz, Off-diagonal long-range behavior of interacting Bose systems, *Phys. Rev. B* **15**, 1399 (1977).
- [38] M. D. Girardeau and E. M. Wright, Bose-Fermi Variational Theory of the Bose-Einstein Condensate Crossover to the Tonks Gas, *Phys. Rev. Lett.* **87**, 210401 (2001).
- [39] P. J. Forrester, N. E. Frankel, T. M. Garoni, and N. S. Witte, Finite one-dimensional impenetrable bose systems: Occupation numbers, *Phys. Rev. A* **67**, 043607 (2003).
- [40] D. M. Gangardt, Universal correlations of trapped one-dimensional impenetrable bosons, *J. Phys. A: Math. Gen.* **37**, 9335 (2004).
- [41] M. Rigol and A. Muramatsu, Universal properties of hard-core bosons confined on one-dimensional lattices, *Phys. Rev. A* **70**, 031603R (2004).
- [42] M. Rigol and A. Muramatsu, Emergence of quasicondensates of hard-core bosons at finite momentum, *Phys. Rev. Lett.* **93**, 230404 (2004).
- [43] E. H. Lieb and W. Liniger, Exact Analysis of an Interacting Bose Gas. I. The General Solution and the Ground State, *Phys. Rev.* **130**, 1605 (1963).
- [44] E. H. Lieb, Exact Analysis of an Interacting Bose Gas. II. The Excitation Spectrum, *Phys. Rev.* **130**, 1616 (1963).
- [45] M. Girardeau, Relationship between systems of impenetrable bosons and fermions in one dimension, *J. Math. Phys.* **1**, 516 (1960).
- [46] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, Cambridge, 1999).
- [47] A. Osterloh, L. Amico, and R. Fazio, Scaling of entanglement close to a quantum phase transition, *Nature* **416**, 608 (2002).
- [48] T. J. Osborne and M. A. Nielsen, Entanglement in a simple quantum phase transition, *Phys. Rev. A* **66**, 032110 (2002).
- [49] K. K. Das, M. D. Girardeau, and E. M. Wright, Crossover from One to Three Dimensions for a Gas of Hard-Core Bosons, *Phys. Rev. Lett.* **89**, 110402 (2002).
- [50] L. Salasnich, A. Parola, and L. Reatto, Transition from three dimensions to one dimension in Bose gases at zero temperature, *Phys. Rev. A* **70**, 013606 (2004).
- [51] L. Salasnich, A. Parola, and L. Reatto, Quasi-one-dimensional bosons in three-dimensional traps: From strong-coupling to weak-coupling regime, *Phys. Rev. A* **72**, 025602 (2005).
- [52] D. S. Petrov, G. V. Shlyapnikov, and J. T. M. Walraven, Regimes of Quantum Degeneracy in Trapped 1D Gases, *Phys. Rev. Lett.* **85**, 3745 (2000).
- [53] M. P. A. Fisher, P. B. Weichman, G. Grinstein, and D. S. Fisher, Boson localization and the superfluid-insulator transition, *Phys. Rev. B* **40**, 546 (1989).
- [54] D. Jaksch, C. Bruder, J. I. Cirac, C. W. Gardiner, and P. Zoller, Cold Bosonic Atoms in Optical Lattices, *Phys. Rev. Lett.* **81**, 3108 (1998).
- [55] T. Matsubara and H. Matsuda, A lattice model of liquid Helium, I, *Prog. Theor. Phys.* **16**, 569 (1956).
- [56] H. Matsuda and T. Matsubara, A lattice model of liquid Helium, II, *Prog. Theor. Phys.* **17**, 19 (1957).
- [57] B. Tóth, Phase transitions in an interacting bose system. an application of the theory of Venttsel' and Freidlin, *J. Stat. Phys.* **61**, 749 (1990).
- [58] O. Penrose, Bose-Einstein condensation in an exactly soluble system of interacting particles, *J. Stat. Phys.* **63**, 761 (1991).
- [59] M. W. Kirson, Bose-Einstein condensation in an exactly solvable model for strongly interacting bosons, *J. Phys. A: Math. Gen.* **33**, 731 (2000).
- [60] G. Günter, H. Schempp, M. Robert-de Saint-Vincent, V. Gavryusev, S. Helmrich, C. S. Hofmann, S. Whitlock, and M. Weidemüller, Observing the Dynamics of Dipole-Mediated Energy Transport by Interaction-Enhanced Imaging, *Science* **342**, 954 (2013).
- [61] H. Schempp, G. Günter, S. Wüster, M. Weidemüller, and S. Whitlock, Correlated Exciton Transport in Rydberg-Dressed-Atom Spin Chains, *Phys. Rev. Lett.* **115**, 093002 (2015).
- [62] R. Burioni, D. Cassi, I. Meccoli, M. Rasetti, S. Regina, P. Sodano, and A. Vezzani, Bose-Einstein condensation in inhomogeneous Josephson arrays, *Europhys. Lett.* **52**, 251 (2000).
- [63] R. Burioni, D. Cassi, M. Rasetti, P. Sodano, and A. Vezzani, Bose-Einstein condensation on inhomogeneous complex networks, *J. Phys. B* **34**, 4697 (2001).
- [64] P. Buonsante, R. Burioni, D. Cassi, and A. Vezzani, Bose-Einstein condensation on inhomogeneous networks: Mesoscopic aspects versus thermodynamic limit, *Phys. Rev. B* **66**, 094207 (2002).
- [65] I. Brunelli, G. Giusiano, F. Mancini, P. Sodano, and A. Trombettoni, Topology-induced spatial Bose-Einstein condensation for bosons on star-shaped optical networks, *J. Phys. B* **37**, S275 (2004).
- [66] E. J. G. Vidal, R. P. A. Lima, and M. L. Lyra, Bose-Einstein condensation in the infinitely ramified star and wheel graphs, *Phys. Rev. E* **83**, 061137 (2011).
- [67] I. N. de Oliveira, T. B. dos Santos, F. A. B. F. de Moura, M. L. Lyra, and M. Serva, Critical behavior of the ideal-gas Bose-Einstein condensation in the Apollonian network, *Phys. Rev. E* **88**, 022139 (2013).
- [68] M. L. Lyra, F. A. B. F. de Moura, I. N. de Oliveira, and M. Serva, Bose-Einstein condensation in diamond hierarchical lattices, *Phys. Rev. E* **89**, 052133 (2014).
- [69] P. Buonsante, R. Burioni, D. Cassi, V. Penna, and A. Vezzani, Topology-induced confined superfluidity in inhomogeneous arrays, *Phys. Rev. B* **70**, 224510 (2004).
- [70] A. Halu, L. Ferretti, A. Vezzani, and G. Bianconi, Phase diagram of the Bose-Hubbard model on complex networks, *Europhys. Lett.* **99**, 18001 (2012).
- [71] M. T. DePue, C. McCormick, S. L. Winoto, S. Oliver, and D. S. Weiss, Unity Occupation of Sites in a 3D Optical Lattice, *Phys. Rev. Lett.* **82**, 2262 (1999).
- [72] I. Bloch, J. Dalibard, and W. Zwerger, Many-body physics with

- ultracold gases, *Rev. Mod. Phys.* **80**, 885 (2008).
- [73] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, Feshbach resonances in ultracold gases, *Rev. Mod. Phys.* **82**, 1225 (2010).
- [74] M. Weidemüller and C. Zimmermann, *Interactions in ultracold gases: from atoms to molecules* (John Wiley & Sons, 2011).
- [75] G. Zürn, F. Serwane, T. Lompe, A. N. Wenz, M. G. Ries, J. E. Bohn, and S. Jochim, Fermionization of two distinguishable fermions, *Phys. Rev. Lett.* **108**, 075303 (2012).
- [76] F. Tennie, V. Vedral, and C. Schilling, Universal upper bounds on the Bose-Einstein condensate and the Hubbard star, *Phys. Rev. B* **96**, 064502 (2017).
- [77] S. R. White, Density matrix formulation for quantum renormalization groups, *Phys. Rev. Lett.* **69**, 2863 (1992).
- [78] S. R. White, Density-matrix algorithms for quantum renormalization groups, *Phys. Rev. B* **48**, 10345 (1993).
- [79] U. Schollwöck, The density-matrix renormalization group, *Rev. Mod. Phys.* **77**, 259 (2005).
- [80] L. Amico, A. Osterloh, and F. Cataliotti, Quantum Many Particle Systems in Ring-Shaped Optical Lattices, *Phys. Rev. Lett.* **95**, 063201 (2005).
- [81] S. Franke-Arnold, J. Leach, M. J. Padgett, V. E. Lembessis, D. Ellinas, A. J. Wright, J. M. Girkin, P. Öhberg, and A. S. Arnold, Optical ferris wheel for ultracold atoms, *Opt. Express* **15**, 8619 (2007).
- [82] A. Ramanathan, K. C. Wright, S. R. Muniz, M. Zelan, W. T. Hill, C. J. Lobb, K. Helmerson, W. D. Phillips, and G. K. Campbell, Superflow in a toroidal Bose-Einstein condensate: An atom circuit with a tunable weak link, *Phys. Rev. Lett.* **106**, 130401 (2011).
- [83] L. Amico, D. Aghamalyan, F. Auksztol, H. Crepaz, R. Dumke, and L. C. Kwek, Superfluid qubit systems with ring shaped optical lattices, *Sci. Rep.* **4**, 4298 (2014).
- [84] T. A. Bell, J. A. P. Glidden, L. Humbert, M. W. J. Bromley, S. A. Haine, M. J. Davis, T. W. Neely, M. A. Baker, and H. Rubinsztein-Dunlop, Bose-Einstein condensation in large time-averaged optical ring potentials, *New J. Phys.* **18**, 035003 (2016).
- [85] K. Sakmann, A. I. Streltsov, O. E. Alon, and L. S. Cederbaum, Reduced density matrices and coherence of trapped interacting bosons, *Phys. Rev. A* **78**, 023615 (2008).
- [86] O. Legeza and J. Sólyom, Quantum data compression, quantum information generation, and the density-matrix renormalization-group method, *Phys. Rev. B* **70**, 205118 (2004).
- [87] J. Rissler, R. M. Noack, and S. R. White, Measuring orbital interaction using quantum information theory, *Chemical Physics* **323**, 519 (2006).
- [88] S. Szalay, M. Pfeffer, V. Murg, G. Barcza, F. Verstraete, R. Schneider, and O. Legeza, Tensor product methods and entanglement optimization for ab initio quantum chemistry, *Int. J. Quant. Chem.* **115**, 1342 (2015).
- [89] H. Bethe, Zur Theorie der Metalle, *Z. Phys.* **71**, 1703 (1931).
- [90] W. Nie, H. Katsura, and M. Oshikawa, Ground State Energies of Spinless Free Fermions and Hard Core Bosons, *Phys. Rev. Lett.* **111**, 100402 (2013).
- [91] W. Nie, H. Katsura, and M. Oshikawa, Particle statistics, frustration, and ground state energy, *Phys. Rev. B* **97**, 125153 (2018).
- [92] C. J. Pethick and L. P. Pitaevskii, Criterion for Bose-Einstein condensation for particles in traps, *Phys. Rev. A* **62**, 033609 (2000).

## SUPPLEMENTARY METHODS

### Derivation of the effective Hamiltonian

The Hamiltonian  $\hat{H}$  is given by Eq. (1). Using  $\hat{H} = \hat{H}_0 + \hat{H}_1$  (with  $\hat{H}_0$  the ring-Hamiltonian) and

$$|\Psi_N\rangle = \alpha|\phi_N\rangle \otimes |0\rangle_c + \beta|\varphi_{N-1}\rangle \otimes |1\rangle_c, \quad (\text{S1})$$

the eigenvalue equation  $\hat{H}|\Psi_N\rangle = E|\Psi_N\rangle$  becomes

$$\begin{aligned} \alpha\hat{H}_0|\phi_N\rangle - \beta s \sum_{i=1}^d h_i^\dagger |\varphi_{N-1}\rangle &= \alpha E|\phi_N\rangle \\ -\alpha s \sum_{i=1}^d h_i |\phi_N\rangle + \beta H_0|\varphi_{N-1}\rangle &= \beta E|\varphi_{N-1}\rangle. \end{aligned} \quad (\text{S2})$$

The unperturbed eigenstates (i.e.,  $s=0$ ) for  $N$  HCBs can be represented as

$$|\psi_\mu^0(N)\rangle = \sum_{1 \leq n_1 < \dots < n_N \leq d} \psi_\mu^0(n_1, \dots, n_N) h_{n_1}^\dagger \dots h_{n_N}^\dagger |0\rangle. \quad (\text{S3})$$

The normalized, totally symmetric ‘wave functions’  $\psi_\mu^0(n_1, \dots, n_N)$  are given for  $1 \leq n_1 < \dots < n_N \leq d$  by the determinant constructed from the one-particle states  $\exp(iq_{\mu_k} n_i)$  [43, 89]

$$\psi_\mu^0(n_1, \dots, n_N) = \mathcal{N} \sum_{P \in S_N} \text{sgn}(P) \exp\left(i \sum_{k=1}^N q_{\mu_{P(k)}} n_k\right), \quad (\text{S4})$$

with  $\mathcal{N} = d^{-N/2}$ .  $S_N$  denotes the permutation group of the integers  $(1, 2, \dots, N)$  and  $\text{sgn}(P)$  its signature. The form (S4) highlights the well-known equivalence of spinless fermions and hard-core bosons in 1d and one has  $\mu_1 < \mu_2 < \dots < \mu_d$ . The unperturbed eigenstates are labelled by  $\boldsymbol{\mu} = (\mu_1, \dots, \mu_N)$  and  $\mu$  determines the wave number  $q_\mu = (\pi/d)(2\mu + 1)$  for  $N$  even and  $q_\mu = (\pi/d)2\mu$  for  $N$  odd [43, 90, 91]. Since  $q_\mu$  is restricted to the first Brillouin zone  $\mu$  takes the values  $-d/2 + 1, -d/2 + 2, \dots, -1, 0, 1, \dots, d/2 - 1, d/2$  for  $d$  even and  $-(d-1)/2, -(d-1)/2 + 1, \dots, -1, 0, 1, \dots, (d-1)/2 - 1, (d-1)/2$  for  $d$  odd. The corresponding unperturbed eigenvalues are given by

$$E_\mu^0(N, d) = -2 \sum_{k=1}^N \cos(q_{\mu_k}). \quad (\text{S5})$$

The unperturbed ground state energy,  $E_{low}^0(N, d)$ , is easily calculated. Using  $\cos(x) = [\exp(ix) + \exp(-ix)]/2$  it is straightforward to calculate the sum in Eq. (S5). As a result one gets

$$E_{low}^0(N, d) = -2 \sin(\frac{\pi}{d}N) / \sin(\frac{\pi}{d}) \simeq -2 \frac{d}{\pi} \sin(\frac{\pi}{d}N), \quad (\text{S6})$$

and the Fermi energy  $E_F(N, d) = E_0^0(N, d) - E_0^0(N-1, d)$  becomes

$$E_F(N, d) = -2 \left[ \tan(\frac{\pi}{2d}) \sin(\frac{\pi}{d}N) + \cos(\frac{\pi}{d}N) \right] \simeq -2 \cos(\pi n) \quad (\text{S7})$$

with  $n = N/d$  the particle density. For  $d \rightarrow \infty, N \rightarrow \infty$  with  $n = N/d$  fixed the unperturbed  $N$ -particle spectrum is a single band with lower band edge  $E_{low}^0(N, n) \simeq -2N \sin(\pi n) / (\pi n)$  and band width  $W(N, n) = 4N \sin(\pi n) / (\pi n) \propto 4N$ . First we note that the result for  $E_{low}^0(N, n)$  holds for  $N$  even and odd and second that the ground state lies in the subspace with total momentum  $Q = \sum_{k=1}^N q_{\mu_k}$  equal to zero.  $Q$  is a good quantum number due to the invariance of  $\hat{H}$  under lattice translations on the ring.

Since the unperturbed eigenstates are complete we have(cf. also Eq. (3) of the main text)

$$\begin{aligned} |\phi_N\rangle &= \sum_{\boldsymbol{\nu}} A_{\boldsymbol{\nu}} |\psi_{\boldsymbol{\nu}}^0(N)\rangle \\ |\varphi_{N-1}\rangle &= \sum_{\boldsymbol{\mu}} a_{\boldsymbol{\mu}} |\psi_{\boldsymbol{\mu}}^0(N-1)\rangle. \end{aligned} \quad (\text{S8})$$

with  $\sum_{\nu} |A_{\nu}|^2 = 1$  and  $\sum_{\mu} |a_{\mu}|^2 = 1$ . The summations in Eq. (S8) are restricted such that  $Q = \sum_{i=1}^N q_{\nu_i} = \sum_{j=1}^{N-1} q_{\mu_j}$  is fixed (mod  $2\pi$ ). In the following we choose  $Q = 0$ , i.e., we consider the HCBs in the frame where the center of mass of the HCBs is at rest (see also [92]). Substituting the ansatz (S8) into Eq. (S2) leads to a decoupling of the  $(N-1)$ -particle and the  $N$ -particle sector:

$$\begin{aligned} [E - E_{\nu}^0(N, d)] A_{\nu} &= s^2 \sum_{\nu'} M_{\nu\nu'}(E) A_{\nu'} \\ [E - E_{\mu}^0(N-1, d)] a_{\mu} &= s^2 \sum_{\mu'} m_{\mu\mu'}(E) a_{\mu'} . \end{aligned} \quad (\text{S9})$$

Here we used that  $E_{\mu}^0(N-1, d)$  and  $E_{\nu}^0(N, d)$  are the corresponding unperturbed eigenvalues of  $|\psi_{\mu}^0(N-1)\rangle$  and  $|\psi_{\nu}^0(N)\rangle$ , respectively. The matrix elements  $M_{\nu\nu'}(E)$  and  $m_{\mu\mu'}(E)$  depend only on the unperturbed eigenstates and eigenvalues and are given by

$$\begin{aligned} M_{\nu\nu'}(E) &= \sum_{\mu'} (b^{\dagger})_{\nu\mu'} [E - E_{\mu'}^0(N-1, d)]^{-1} b_{\mu'\nu'} \\ m_{\mu\mu'}(E) &= \sum_{\nu'} b_{\mu\nu'} [E - E_{\nu'}^0(N, d)]^{-1} b_{\nu'\mu'}^{\dagger} . \end{aligned} \quad (\text{S10})$$

The crucial quantity is the matrix  $\mathbf{b}$  with elements

$$b_{\mu\nu} = \langle \psi_{\mu}^0(N-1) | \sum_{i=1}^d h_i | \psi_{\nu}^0(N) \rangle . \quad (\text{S11})$$

Having solved Eq. (S9) one obtains from Eq. (S2) with Eq. (S8) the coefficients  $\alpha$  and  $\beta$ .

Operating with  $\sum_{\nu} A_{\nu}^* [E - E_{\nu}^0(N, d)]^{-1}$  and  $\sum_{\mu} a_{\mu}^* [E - E_{\mu}^0(N-1, d)]^{-1}$ , respectively, on the 1st and 2nd line of Eq. (S9) and taking the normalization of  $\{A_{\nu}\}$  and  $\{a_{\mu}\}$  into account the eigenvalue equations take the form

$$\begin{aligned} 1 &= s^2 f_d^{(N-1)}(E; \{a_{\mu}\}) \\ 1 &= s^2 F_d^{(N)}(E; \{A_{\nu}\}) , \end{aligned} \quad (\text{S12})$$

with

$$f_d^{(N-1)}(E; \{a_{\mu}\}) = \sum_{\mu\mu'\nu'} a_{\mu}^* [E - E_{\mu}^0(N-1, d)]^{-1} b_{\mu\nu'} [E - E_{\nu'}^0(N, d)]^{-1} (b^{\dagger})_{\nu'\mu'} a_{\mu'} . \quad (\text{S13})$$

and

$$F_d^{(N)}(E; \{A_{\nu}\}) = \sum_{\nu\nu'\mu'} A_{\nu}^* [E - E_{\nu}^0(N, d)]^{-1} (b^{\dagger})_{\nu\mu'} [E - E_{\mu'}^0(N-1, d)]^{-1} b_{\mu'\nu'} A_{\nu'} \quad (\text{S14})$$

The unperturbed eigenfunctions can always be chosen to be real, since the unperturbed Hamiltonian is real. Therefore  $b_{\mu\nu}$  and  $(b^{\dagger})_{\nu\mu}$  are real. Furthermore,  $\{a_{\mu}\}$  and  $\{A_{\nu}\}$  can also be chosen to be real since the Hamiltonian  $\hat{H}$  is real as well. Therefore the functions  $f_d^{(N-1)}(E; \{a_{\mu}\})$  and  $F_d^{(N)}(E; \{A_{\nu}\})$  are real.

Eq. (S12) together with Eqs. (S13) and (S14) already allows to obtain some qualitative information on the low-energy part of the perturbed  $N$ -particle spectrum. A crucial observation is that  $f_d^{(N-1)}(E; \{a_{\mu}\})$  and  $F_d^{(N)}(E; \{A_{\nu}\})$  have *poles* at the unperturbed  $(N-1)$ - and  $N$ -particle eigenvalues. As discussed above the unperturbed spectrum of  $(N-1)$  and  $N$  HCBs form a band with lower band edge  $E_{low}^0(N-1, d)$  and  $E_{low}^0(N, d)$ , respectively. It is  $E_{low}^0(N, d) = E_{low}^0(N-1, d) + E_F(N, d)$  with the Fermi energy from Eq. (S7). For  $n < 1/2$  it follows  $E_F(N, d) < 0$ . Note, this is not a restriction due to the particle-hole duality. Therefore,  $E_{low}^0(N, d) < E_{low}^0(N-1, d)$ . Now, we will show that the low-lying perturbed eigenvalues form a discrete spectrum below  $E_{low}^0(N, d)$ , and the band between  $E_{low}^0(N, d)$  and  $E_{low}^0(N-1, d)$  persists. The discrete spectrum exhibits an excitation gap even for  $d = \infty$ .

Let us choose  $E$  between  $E_{low}^0(N, d)$  and  $E_{low}^0(N-1, d)$  and let us denote the increasingly ordered unperturbed eigenvalues  $\{E_{\nu}^0(N, d)\}$  in this interval by  $E_{\nu_k}^0(N, d)$ ,  $k \geq 1$ . Due to  $E < E_{low}^0(N-1, d)$  the denominators  $[E - E_{\mu}^0(N-1, d)]^{-1}$  in Eq. (S13) are negative for all  $\mu$ , i.e., they do not change sign. Then, under variation of  $E$  between  $E_{\nu_k}^0(N, d)$  and  $E_{\nu_{k+1}}^0(N, d)$ , the function  $f_d^{(N-1)}(E; \{a_{\mu}\})$  varies continuously from  $\pm\infty$  at  $E = E_{\nu_k}^0(N, d)$  to  $\mp\infty$  at  $E = E_{\nu_{k+1}}^0(N, d)$ , independent of  $\{a_{\mu}\}$ . Accordingly, for arbitrary  $s \neq 0$  the first equation of (S12) has always a solution  $E_{\nu_k}(N, d; s, \{a_{\nu}\})$  which is between

$E_{\nu_k}^0(N, d)$  and  $E_{\nu_{k+1}}^0(N, d)$ . Substituting  $E_{\nu_k}(N, d; s, \{a_\nu\})$  into the second line of Eq. (S9) yields  $\{a_{\nu^{(\nu_k)}}\}$  which in turn leads to the perturbed eigenvalues  $E_{\nu_k}(N, d; s) = E_{\nu_k}(N, d; s, \{a_{\nu^{(\nu_k)}}\})$ ,  $k \geq 1$ . For  $d \rightarrow \infty$  these perturbed eigenvalues  $\{E_{\nu_k}(N, d; s)\}$  form a band with lower band edge  $E_{low}^0(N, d)$  and upper edge  $E_{low}^0(N-1, d)$ .

For  $E \geq E_{low}^0(N-1, d)$  there exist pairs  $\nu', \mu$  such that there is no unperturbed eigenvalue between  $E_{\nu'}^0(N, d)$  and  $E_\mu^0(N-1, d)$ . In that case  $[E - E_{\nu'}^0(N, d)]^{-1}[E - E_\mu^0(N-1, d)]^{-1}$  in Eq. (S13) changes from  $\pm\infty$  at  $E_{\nu'}^0(N, d)$  to  $\pm\infty$  at  $E_\mu^0(N-1, d)$  under varying  $E$  between  $E_{\nu'}^0(N, d)$  and  $E_\mu^0(N-1, d)$ . Therefore,  $f_d^{(N-1)}(E; \{a_\nu\})$  does not necessarily change sign and the first equation of (S12) may only have a solution for  $s^2$  small enough. In case that the solution between  $E_{\nu'}^0(N, d)$  and  $E_\mu^0(N-1, d)$  disappears if  $s^2$  becomes large enough, a perturbed eigenvalue must appear below(or above) the lower(upper) band edge  $-|E_{low}^0(N, d)|(|E_{low}^0(N-1, d)|)$ , since the total number of eigenvalues does not depend on  $s$ .

Finally, let us discuss  $E < E_{low}^0(N, d)$ . In that case the product of both denominators in Eq. (S13) is always positive. For  $E \rightarrow E_{low}^0(N, d)$  from below  $f_d^{(N-1)}(E; \{a_\mu\})$  will diverge to  $+\infty$ . Because  $f_d^{(N-1)}(E; \{a_\mu\}) \rightarrow 0$  for  $E \rightarrow -\infty$  there must exist at least one solution  $E(N, d; s, \{a_\mu\})$  of the first equation of Eq. (S12) for all  $s \neq 0$ .  $E(N, d; s, \{a_\mu\})$  will have a gap to the lower band edge  $E_{low}^0(N, d)$ . Depending on  $s^2$  and  $\{a_\mu\}$  there may exist more than one solution. Substitution them into the first line of Eq. (S9) yields a discrete spectrum. The same qualitative discussion can be done for  $F_d^{(N)}(E; \{A_\nu\})$  (Eq. (S14)) in combination with the second equation in (S12). We have checked the correctness of these qualitative results on the perturbed spectrum for  $N = 2$ . There is little doubt that they become incorrect for  $N > 2$ .

Now we describe how the discrete part of the perturbed spectrum and the corresponding eigenstates below  $E_{low}^0(N, d)$  can be obtained exactly in two limiting cases. In these two cases  $[E - E_{\mu'}^0(N-1, d)]^{-1}$  and  $[E - E_{\nu'}^0(N, d)]^{-1}$  can be replaced by  $[E - E_{low}^0(N, d) + E_F(N, d)]^{-1}$  and  $[E - E_{low}^0(N, d)]^{-1}$ , respectively. Then, the matrices  $(M_{\nu\nu'}(E))$  and  $(m_{\mu\mu'}(E))$  strongly simplify since the sums in Eq. (S10) can be performed using Eq. (S11) and the completeness relations  $\sum_\mu |\psi_\mu^0(N-1)\rangle\langle\psi_\mu^0(N-1)| = \mathbb{1}|_{N-1}$ ,  $\sum_\nu |\psi_\nu^0(N)\rangle\langle\psi_\nu^0(N)| = \mathbb{1}|_N$ .  $\mathbb{1}|_{N-1}$  and  $\mathbb{1}|_N$  denote the identity operator, respectively, in the  $(N-1)$  and  $N$  particle subspace. Then it follows

$$\begin{aligned} M_{\nu\nu'}(E) &\simeq [E - E_{low}^0(N, d) + E_F(N, d)]^{-1} \langle\psi_\nu^0(N)| \sum_{ij} h_i^\dagger h_j |\psi_{\nu'}^0(N)\rangle \\ m_{\mu\mu'}(E) &\simeq [E - E_{low}^0(N, d)]^{-1} \langle\psi_\mu^0(N-1)| \sum_{ij} h_i h_j^\dagger |\psi_{\mu'}^0(N-1)\rangle. \end{aligned}$$

and Eq. (S9) simplifies to

$$\begin{aligned} E^{eff} A_\nu &\simeq s^2 \sum_{\nu'} \langle\psi_\nu^0(N)| \sum_{ij} h_i^\dagger h_j |\psi_{\nu'}^0(N)\rangle A_{\nu'} \\ E^{eff} a_\mu &\simeq s^2 \sum_{\mu'} \langle\psi_\mu^0(N-1)| \sum_{ij} h_i h_j^\dagger |\psi_{\mu'}^0(N-1)\rangle a_{\mu'}, \end{aligned} \quad (\text{S15})$$

with

$$E^{eff}(N, d) = [E - E_{low}^0(N, d) + E_F(N, d)][E - E_{low}^0(N, d)]. \quad (\text{S16})$$

The eigenvalue equations (S15) are identical to the eigenvalue equations following from

$$\begin{aligned} \hat{H}_N^{eff} |\phi_N\rangle &= E^{eff}(N, d) |\phi_N\rangle \\ \hat{h}_{N-1}^{eff} |\varphi_{N-1}\rangle &= E^{eff}(N, d) |\varphi_{N-1}\rangle. \end{aligned} \quad (\text{S17})$$

with the effective Hamiltonians

$$\begin{aligned} \hat{H}_N^{eff} &= \tilde{s}^2 \frac{1}{d} \sum_{i,j=1}^d h_i^\dagger h_j \\ \hat{h}_{N-1}^{eff} &= \tilde{s}^2 \frac{1}{d} \sum_{i,j=1}^d h_i h_j^\dagger, \end{aligned} \quad (\text{S18})$$

and  $|\phi_N\rangle, |\varphi_{N-1}\rangle$  from Eq. (S8). Note, the eigenvalue  $E^{eff}(N, d)$  is identical for  $\hat{H}_N^{eff}$  and  $\hat{h}_{N-1}^{eff}$ .  $\tilde{s} = (s/t)\sqrt{d}$  denotes the scaled dimensionless coupling constant. We remind the reader that we used  $t = 1$ .

In the following it is more convenient to use the equivalence  $h_i^\dagger = S_i^-$ ,  $h_i = S_i^+$ ,  $(1 - 2h_i^\dagger h_i) = 2S_i^z$  between the hard-core Bose operators and the spin-one-half operators. The commutation relations of the latter read

$$[S_i^+, S_j^-] = 2\delta_{ij}S_i^z, \quad [S_j^z, S_i^\pm] = \pm\delta_{ij}S_i^\pm. \quad (\text{S19})$$

The effective Hamiltonians become

$$\hat{H}_N^{eff} = \tilde{s}^2 \frac{1}{d} \sum_{i,j=1}^d S_i^- S_j^+ \quad (\text{S20})$$

$$\hat{h}_{N-1}^{eff} = \tilde{s}^2 \frac{1}{d} \sum_{i,j=1}^d S_i^+ S_j^- , \quad (\text{S21})$$

Let  $\vec{S} = \sum_{i=1}^d \vec{S}_i$  be the spin operator of the total spin. Because  $\hat{H}_N^{eff}$  (and  $\hat{h}_{N-1}^{eff}$ ) commutes with  $\vec{S}^2$  and  $S^z$  all its eigenstates can be chosen such that they are also eigenstates of  $\vec{S}^2$  and  $S^z$  with eigenvalues  $S(S+1)$  and  $M$ , respectively. They will be denoted by  $|S, M\rangle$ .  $M$  is related to the particle number by  $M = d/2 - N$  and for fixed  $N$  the total spin quantum number takes the values  $S = d/2 - N, d/2 - N + 1, \dots, d/2$ . The corresponding eigenvalues of  $\hat{H}_N^{eff}$  are given by

$$E^{eff}(S, M; \tilde{s}) = \tilde{s}^2 \frac{1}{d} [S(S+1) - M(M+1)]. \quad (\text{S22})$$

The ground state eigenvalue  $E_0(N, d; \tilde{s})$  of  $\hat{H}$  follows from the largest eigenvalue  $E_{max}^{eff}(N, d; \tilde{s})$  of  $\hat{H}_N^{eff}$  which corresponds to  $S_{max} = d/2$ . Then we obtain from Eq. (S22) in the thermodynamic limit  $N \rightarrow \infty, d \rightarrow \infty$  with density  $n = N/d$  fixed

$$E_{max}^{eff}(N, d; \tilde{s}) \simeq \tilde{s}^2 N(1 - n). \quad (\text{S23})$$

The corresponding eigenstates are given by

$$\begin{aligned} |\phi_N^{eff}\rangle &= \binom{d}{N}^{-1/2} \left( \sum_{i=1}^d h_i^\dagger \right)^N |0\rangle \\ |\varphi_{N-1}^{eff}\rangle &= \binom{d}{N-1}^{-1/2} \left( \sum_{i=1}^d h_i^\dagger \right)^{N-1} |0\rangle. \end{aligned} \quad (\text{S24})$$

Using the spin analogy it is  $|\phi_N^{eff}\rangle = |d/2, d/2 - N\rangle$  and  $|\varphi_{N-1}^{eff}\rangle = |d/2, d/2 - (N-1)\rangle$ . Note, these eigenstates belong to the subspace with total momentum  $Q = 0$ .

Substitution of  $E_{max}^{eff}(N, d; \tilde{s})$  from Eq. (S23) into Eq. (S16) leads to the perturbed ground state eigenvalue

$$E_0(N, d; \tilde{s}) \simeq E_{low}^0(N, d) - E_F(N, d)/2 - \sqrt{(E_F(N, d)/2)^2 + \tilde{s}^2 d n(1 - n)}. \quad (\text{S25})$$

We remind the reader that Eq. (S7) implies  $E_F(N, d) \leq 0$  for  $0 \leq n \leq 1/2$ . Due to the particle hole duality  $n \leq 1/2$  does not restrict generality.

The energy,  $E_1(N, d; \tilde{s})$ , of the first excitation follows from Eq. (S16) for the second largest effective eigenvalue. Using the spin analogy the latter corresponds to  $S = S_{max} - 1 \equiv d/2 - 1$  which yields  $E^{eff}(N, d; \tilde{s}) = E_{max}^{eff}(N, d; \tilde{s}) - \tilde{s}^2$ . Accordingly we obtain

$$E_1(N, d; \tilde{s}) \simeq E_{low}^0(N, d) - E_F(N, d)/2 - \sqrt{(E_F(N, d)/2)^2 + \tilde{s}^2 [d n(1 - n) - 1]}. \quad (\text{S26})$$

It is easy to see that the excitation gap  $\Delta E(N, d; \tilde{s}) = E_1(N, d; \tilde{s}) - E_0(N, d; \tilde{s})$  is finite for  $\tilde{s} \neq 0$ . The higher excitation energies  $E_n(N, d; \tilde{s})$  for  $n = 2, \dots, N-1$  follow similarly using  $S = d/2 - n$ . Therefore, the eigenvalues of  $\hat{H}$  below  $E_{low}^0(N, d)$  form a discrete spectrum of  $N$  eigenvalues.

Let us summarize: The unperturbed spectrum in the subspace  $Q = 0$  consists of two bands. One band,  $B_{N-1}^0$ , of  $K_{N-1}$  eigenvalues,  $\{E_\mu^0(N-1, d)\}$ , and the other band,  $B_N^0$ , with  $K_N$  eigenvalues,  $\{E_\nu^0(N, d)\}$ . These two bands correspond to  $(N-1)$  and  $N$  HCBs on the ring-lattice. The band edges of  $B_{N-1}^0$  are at  $\pm E_{low}^0(N-1, d)$  and those of  $B_N^0$  at  $\pm E_{low}^0(N, d)$ . For  $d$  finite, both sets  $\{E_\mu^0(N-1, d)\}$  and  $\{E_\nu^0(N, d)\}$  are disjoint, and for density  $n = N/d \leq 1/2$   $B_{N-1}^0$  is a subset within the interval  $[-|E_{low}^0(N, d)|, +|E_{low}^0(N, d)|]$ . Turning on  $s$  leads to a coupling between these two bands. Part of these two bands persist. The lower band edge of the perturbed band coincides for  $d = \infty$  with the lower band edge of the unperturbed band.

Below that band a discrete spectrum of maximally  $N$  eigenvalues occurs exhibiting an excitation gap. The number of discrete eigenvalues may change with  $\tilde{s}$ .

The number  $N_0 = (1/d)\langle\Psi_N|\sum_{i,j=1}^d h_i^\dagger h_j|\Psi_N\rangle$  of condensed particles in the state  $|\Psi_N\rangle$  is easily obtained since  $(1/d)\langle\Psi_N|\sum_{i,j=1}^d h_i^\dagger h_j|\Psi_N\rangle \simeq \langle\Psi_N|\hat{H}_N^{eff}|\Psi_N\rangle/\tilde{s}^2$ . Substitution of  $|\Psi_N\rangle$  from Eq. (S1) leads to  $N_0 = [|\alpha|^2\langle\phi_N^{eff}|\hat{H}_N^{eff}|\phi_N^{eff}\rangle + |\beta|^2\langle\varphi_{N-1}^{eff}|\hat{H}_N^{eff}|\varphi_{N-1}^{eff}\rangle]/\tilde{s}^2$ . Making use of the commutation relations of  $\{h_i^\dagger\}, \{h_j\}$  one can express  $\hat{H}_N^{eff}$  in the second summand by  $\hat{h}_{N-1}^{eff}$ :

$$\hat{H}_N^{eff} = \hat{h}_{N-1}^{eff} - \frac{\tilde{s}^2}{d} \sum_{i=1}^d (1 - 2\hat{n}_i). \quad (\text{S27})$$

With  $|\alpha|^2 + |\beta|^2 = 1$ ,  $\sum_{i=1}^d \hat{n}_i |\varphi_{N-1}^{eff}\rangle = (N-1)|\varphi_{N-1}^{eff}\rangle$ , Eqs. (S17), (S18) and (S23) we obtain for  $d \rightarrow \infty$  the final result

$$N_0(N; \tilde{s}) \simeq N[(1-n) - |\beta|^2(1-2n)N^{-1}]. \quad (\text{S28})$$

Since  $|\beta|^2 \leq 1$  the second term on the r.h.s. of Eq. (S28) is a negative correction (for  $n < 1/2$ ) to the leading order term  $(1-n)$  which is of order  $\mathcal{O}(1/N)$ .

As discussed above the mapping of the original model to an effective one is valid if one is allowed to replace  $E_{\nu'}^0(N, d)$  by the unperturbed ground state energy eigenvalue  $E_{low}^0(N, d)$ . This is equivalent to the replacement of  $\hat{H}_0$  in the first line of Eq. (S2) by  $E_{low}^0(N, d)$ . With this replacement and that of  $(|\phi_N\rangle, |\varphi_{N-1}\rangle)$  by  $(|\phi_N^{eff}\rangle, |\varphi_{N-1}^{eff}\rangle)$  from Eq. (S24) one can solve the linear equation for  $\alpha, \beta$ . With use of  $[E_0(N, d; \tilde{s}) - E_{low}^0(N, d)]$  from Eq. (S25) and the normalization condition  $|\alpha|^2 + |\beta|^2 = 1$  one obtains for the ground state

$$|\alpha|^2 \simeq 1 - \frac{1}{2}\tilde{s}^2 N(1-n) \left\{ [(E_F/2)^2 + \tilde{s}^2 N(1-n)] - (E_F/2)\sqrt{(E_F/2)^2 + \tilde{s}^2 N(1-n)} \right\}^{-1} \quad (\text{S29})$$

$$|\beta|^2 \simeq \frac{1}{2}\tilde{s}^2 N(1-n) \left\{ [(E_F/2)^2 + \tilde{s}^2 N(1-n)] - (E_F/2)\sqrt{(E_F/2)^2 + \tilde{s}^2 N(1-n)} \right\}^{-1}. \quad (\text{S30})$$

Now we discuss the validity of the above mapping of the original eigenvalue problem to an effective one. The simplest limiting case under which the mapping becomes exact is the strong coupling limit  $s \rightarrow \infty$ . For the ground state energy  $E = E_0(N, d; \tilde{s})$  the denominators in Eq. (S10) can be rewritten as follows

$$[E_0(N, d; \tilde{s}) - E_{\nu'}^0(N, d)] = [E_0(N, d; \tilde{s}) - E_{low}^0(N, d)] \{1 + [E_{\nu'}^0(N, d) - E_{low}^0(N, d)]/[E_{low}^0(N, d) - E_0(N, d; \tilde{s})]\}^{-1}. \quad (\text{S31})$$

and similar for  $[E_0(N, d; \tilde{s}) - E_{\mu'}^0(N-1, d)]$ . Accordingly, the mapping becomes exact if

$$[E_{\nu'}^0(N, d) - E_{low}^0(N, d)]/[E_{low}^0(N, d) - E_0(N, d; \tilde{s})] \rightarrow 0. \quad (\text{S32})$$

Since the upper band edge fulfils  $E_{up}^0(N, d) = -E_{low}^0(N, d)$  it follows  $|[E_{\nu'}^0(N, d) - E_{low}^0(N, d)]| \leq 2|E_{low}^0(N, d)|$  for all  $\nu'$  and the condition (S32) becomes  $2|E_{low}^0(N, d)|/[E_{low}^0(N, d) - E_0(N, d; \tilde{s})] \ll 1$ . Using Eqs. (S6), (S7) and (S25) we obtain the condition

$$s \gg \frac{4}{\pi} \frac{\sin \pi n}{\sqrt{n(1-n)}}. \quad (\text{S33})$$

The reason why the mapping becomes exact in the scaling limit  $s \rightarrow 0, d \rightarrow \infty$  with  $\tilde{s} = s\sqrt{d}$  and  $N$  fixed, is more subtle. In that case the variation with  $\mu'$  of the numerator  $b_{\nu\mu}^\dagger, b_{\mu'\nu'}$  and of the denominator  $[E - E_{\mu'}^0(N-1, d)]$  in the first line of Eq. (S10) plays the essential role. In Appendix B we prove that  $b_{\mu\nu} \sim \sqrt{d}$  for  $\mu, \nu$  fixed and  $d \rightarrow \infty$  whereas  $b_{\mu\nu} = \mathcal{O}(1)$  if  $\mu_k \in \mu$  and  $\nu_k \in \nu$  are of  $\mathcal{O}(d)$ . This means that  $b_{\mu\nu}$  decreases fast with increasing  $\mu_k$  and  $\nu_k$ . Therefore the main contributions in the sums in Eq. (S10) for  $N$  arbitrary large but fixed comes from  $\mu', \nu$  with  $\mu'_k$  and  $\nu_k$  arbitrary large but fixed. Therefore restricting the sums in Eq. (S10), e.g., over  $\mu'$  to  $|\mu'_k| \leq \sqrt{d}$  for all  $k$  does not change the result if  $d$  becomes very large. Due to this restriction of  $\{\mu_k\}$  we obtain with  $E_{low}^0(N, d) \simeq -2N$  and  $\cos(2\pi\mu_k/d) \leq 1 - 2\pi^2/d$  from (S5) the upper bound  $[E_{\nu'}^0(N, d) - E_{low}^0(N, d)] \ll 4\pi^2 N/d$  for the numerator in Eq. (S32). Substituting this upper bound and  $[E_{low}^0(N, d) - E_0(N, d; \tilde{s})]$  from Eq. (S25) into Eq. (S32) leads for  $N$  fixed and  $d \gg 1$  to the condition

$$s \gg 2\sqrt{2}\pi/d. \quad (\text{S34})$$

Here we also used  $E_F(N, d) \simeq -2$  because  $n \approx 0$  for  $N$  fixed and  $d \gg 1$  (cf. Eq. (S7)).

**Behavior of  $b_{\mu,\nu}$  for  $N$  fixed and  $d \rightarrow \infty$**

To study the behavior of  $b_{\mu\nu}$  for  $N$  fixed and  $d \rightarrow \infty$  we first observe that the translational invariance on the ring implies that Eq. (S11) becomes  $b_{\mu\nu} = d \langle \psi_{\mu}^0(N-1) | h_1 | \psi_{\nu}^0(N) \rangle$ . Substituting  $|\psi_{\mu}^0(N-1)\rangle, |\psi_{\nu}^0(N)\rangle$  from Eq. (S3) and taking advantage of the ordering  $1 \leq n_1 < \dots < n_N \leq d$  one arrives at

$$b_{\mu\nu} = d \sum_{2 \leq m_2 < \dots < m_N \leq d} \psi_{\mu}^0(m_2, \dots, m_N)^* \psi_{\nu}^0(1, m_2, \dots, m_N). \quad (\text{S35})$$

Introducing new variables  $n_i = m_{i+1} - 1$  and taking the translational invariance into account this yields

$$b_{\mu\nu} = d \sum_{1 \leq n_1 < \dots < n_{N-1} \leq d-1} \psi_{\mu}^0(n_1, \dots, n_{N-1})^* \psi_{\nu}^0(0, n_1, \dots, n_{N-1}). \quad (\text{S36})$$

Substituting the normalized ‘wave functions’ from Eq. (S4) leads to

$$b_{\mu\nu} = dd^{-(N-\frac{1}{2})} \sum_{P \in S_{N-1}} \sum_{P' \in S_N} \text{sgn}(P) \text{sgn}(P') \sum_{1 \leq n_1 < \dots < n_{N-1} \leq (d-1)} \exp \left[ -i \sum_{k=1}^{N-1} (q_{\mu P(k)} - q_{\nu P'(k+1)}) n_k \right]. \quad (\text{S37})$$

The crucial quantity is the 2nd line of Eq. (S37). This sum can be written as  $\sum_{n_1=1}^{d-N+1} \sum_{n_2=n_1+1}^{d-N+2} \dots \sum_{n_{N-1}=n_{N-2}+1}^{d-1} (\dots)$ . Each single sum generates a denominator of the form  $(1 - \exp[-i(\sum_{j=1}^{N-1} a_j q_{\mu_j} - \sum_{j'=1}^N b_{j'} q_{\nu_{j'}})])$  where the integers  $\{a_j\}$  and  $\{b_{j'}\}$  take values  $0, \pm 1$ . There is a product of  $(N-1)$  such denominators. For  $\mu, \nu$  fixed and for  $d \rightarrow \infty$  this product is proportional to  $d^{N-1}$ . Performing the sums in the 2nd line of Eq. (S37) also generates numerators of the form  $(1 - \exp[-i(\sum_{j=1}^{N-1} a'_j q_{\mu_j} - \sum_{j'=1}^N b'_{j'} q_{\nu_{j'}})d])$  where  $\{a'_j\}$  and  $\{b'_{j'}\}$  take values  $0, \pm 1$ .  $\exp[iq_{\mu_j}d] = +1(-1)$  and  $\exp[iq_{\nu_j}d] = -1(+1)$  for  $N$  even(odd). Therefore some of the numerators vanish and some do not. The latter take the value 2. Accordingly, for  $d \rightarrow \infty$  the contribution of these terms in the 2nd line of Eq. (S37) is of order  $d^{N-1}$ . The contribution of all the other terms are of  $\mathcal{O}(d^{N-2})$ . Taking the prefactor  $dd^{-(N-\frac{1}{2})}$  on the r.h.s. of Eq. (S37) into account one obtains for  $(\mu, \nu)$  arbitrary but fixed and  $d \rightarrow \infty$  in leading order in  $d$

$$b_{\mu\nu} \sim \sqrt{d}, \quad (\text{S38})$$

which we wanted to prove.