

ARITHMETIC COMPLEXITY REVISITED

IGOR V. NIKOLAEV¹

ABSTRACT. The arithmetic complexity $c(\mathcal{A}_\theta)$ of a noncommutative torus \mathcal{A}_θ measures the rank r of a rational elliptic curve $\mathcal{E}(K) \cong \mathbf{Z}^r \oplus \mathcal{E}_{tors}$ via the formula $r = c(\mathcal{A}_\theta) - 1$. The number $c(\mathcal{A}_\theta)$ is equal to the dimension of a connected component $V_{N,k}^0$ of the Brock-Elkies-Jordan variety associated to a periodic continued fraction $\theta = [b_1, \dots, b_N, \overline{a_1, \dots, a_k}]$ of the period (a_1, \dots, a_k) . We prove that the component $V_{N,k}^0$ is a fiber bundle over the Fermat-Pell conic \mathcal{Q} with the structure group \mathcal{E}_{tors} and the fiber an r -dimensional affine space. As an application, we evaluate the Tate-Shafarevich group $\text{III}(\mathcal{E}(K))$ of elliptic curve $\mathcal{E}(K)$ in terms of the group $\text{III}(\mathcal{Q})$ calculated by Lemmermeyer.

1. INTRODUCTION

Noncommutative geometry deals with an interplay between spatial forms and algebras with non-commutative multiplication. To illustrate the idea, consider the noncommutative torus \mathcal{A}_θ , i.e. a C^* -algebra generated by the unitary operators U and V satisfying the relation $VU = e^{2\pi i\theta}UV$, where θ is a real constant [Rieffel 1990] [7]. The algebra \mathcal{A}_θ is the coordinate ring of a complex elliptic curve $\mathcal{E}(\mathbf{C})$, but no longer a commutative ring. Namely, there exists a covariant functor F mapping isomorphic elliptic curves $\mathcal{E}(\mathbf{C})$ to the Morita equivalent noncommutative tori \mathcal{A}_θ [5, Section 1.3]. A restriction of F to the curves $\mathcal{E}(K)$ over a number field $K \subset \mathbf{C}$ corresponds to θ being the irrational root of a quadratic polynomial with integer coefficients. Such a root unfolds into a periodic continued fraction of the period (a_1, \dots, a_k) , i.e.

$$\theta = [b_1, \dots, b_N, \overline{a_1, \dots, a_k}]. \quad (1.1)$$

Let $\mathcal{E}(K) \cong \mathbf{Z}^r \oplus \mathcal{E}_{tors}$ be the Mordell-Weil group of the K -points of $\mathcal{E}(\mathbf{C})$. To recast the rank $r \geq 0$ of $\mathcal{E}(K)$ in terms of the continued fraction (1.1), an arithmetic complexity $c(\mathcal{A}_\theta)$ was used [5, Section 6.2.1]. Roughly speaking, the $c(\mathcal{A}_\theta)$ counts the independent entries a_i and b_j in (1.1). Namely, the $c(\mathcal{A}_\theta)$ is equal to the Krull dimension of a projective variety \mathcal{V}_E associated to the continued fraction (1.1). The equation of \mathcal{V}_E for $\theta = \{\sqrt{D} \mid D > 0 \text{ square-free integer}\}$ was first considered by [Euler 1765] [2], hence the notation. The rank of elliptic curve $\mathcal{E}(K)$ and the arithmetic complexity of $\mathcal{A}_\theta = F(\mathcal{E}(K))$ are linked by a simple formula:

$$r = c(F(\mathcal{E}(K))) - 1. \quad (1.2)$$

An affine variety $V_{N,k}(\mathbf{C})$ associated to (1.1) has been studied by [Brock, Elkies & Jordan 2019] [1]. In this remarkable paper, the authors prove that the $V_{N,k}(\mathbf{C})$ fibers over the Fermat-Pell conic \mathcal{Q} [Brock, Elkies & Jordan 2019] [1, Theorem 3.5].

2010 *Mathematics Subject Classification.* Primary 11G05; Secondary 46L85.

Key words and phrases. elliptic curve, noncommutative torus, Brock-Elkies-Jordan variety.

The \mathbf{Z} -points of $V_{N,k}(\mathbf{C})$ correspond to the continued fractions (1.1) and the reader is referred to Section 2.1 for other details. We call the $V_{N,k}(\mathbf{C})$ a *Brock-Elkies-Jordan variety*.

The aim of our note is to relate geometry of the variety $V_{N,k}(\mathbf{C})$ with the rank r , torsion \mathcal{E}_{tors} and the arithmetic complexity $c(F(\mathcal{E}(K)))$ of the elliptic curve $\mathcal{E}(K)$. The maximal connected component of the Brock-Elkies-Jordan variety containing a point $(b_1, \dots, b_N; a_1, \dots, a_k)$ will be denoted by $V_{N,k}^0$. (We refer the reader to Section 4.2 for quick examples of the $V_{1,2}^0$.) Let \mathbb{A}^r be the r -dimensional affine space. Our main results can be formulated as follows.

Theorem 1.1. *The $V_{N,k}^0$ is a fiber bundle over the Fermat-Pell conic \mathcal{Q} with the fibers \mathbb{A}^r and the structure group \mathcal{E}_{tors} . In particular, the arithmetic complexity is given by the formula*

$$c(F(\mathcal{E}(K))) = \dim_{\mathbf{C}} V_{N,k}^0. \quad (1.3)$$

Remark 1.2. Theorem 1.1 can be used to evaluate ranks of the elliptic curves via formula (1.2). As an example, we calculate such ranks in Section 4 using explicit formulas for the continued fractions (1.1) with $N + k \leq 3$ developed by [Brock, Elkies & Jordan 2019] [1, Sections 5-8].

Recall that the Tate-Shafarevich group $\text{III}(\mathcal{E}(K))$ measures the failure of the Hasse principle for the elliptic curve $\mathcal{E}(K)$. Denote by $Cl(\Lambda)$ the (narrow) class group of the order $\Lambda := \mathbf{Z} + \mathbf{Z}\theta$ in the ring of integers of the real quadratic field $k = \mathbf{Q}(\theta)$. If Λ is the maximal order, then $Cl(\Lambda)$ is the class group of the field k . If \mathcal{Q} is a conic, then $\text{III}(\mathcal{Q}) \cong Cl(\Lambda) \oplus Cl(\Lambda)$ [Lemmermeyer 2003] [4, Theorem 11]. Applying this formula to the fiber bundle $V_{N,k}^0$ over the Fermat-Pell conic \mathcal{Q} , one gets the following result.

Corollary 1.3. $\text{III}(\mathcal{E}(K)) \cong Cl(\Lambda) \oplus Cl(\Lambda)$.

The article is organized as follows. Preliminary facts can be found in Section 2. The proofs of theorem 1.1 and corollary 1.3 are given in Section 3. Two examples illustrating theorem 1.1 are considered in Section 4.

2. PRELIMINARIES

We briefly review the Brock-Elkies-Jordan varieties and arithmetic complexity of the noncommutative tori. For a detailed account we refer the reader to [Brock, Elkies & Jordan 2019] [1] and [5, Section 6.2.1], respectively.

2.1. Brock-Elkies-Jordan variety. By an infinite continued fraction one understands an expression of the form:

$$[c_1, c_2, c_3, \dots] := c_1 + \frac{1}{c_2 + \frac{1}{c_3 + \dots}}, \quad (2.1)$$

where c_1 is an integer and c_2, c_3, \dots are positive integers. The continued fraction (2.1) converges to an irrational number and each irrational number has a unique representation by (2.1). The expression (2.1) is called *k-periodic*, if $c_{i+k} = c_i$ for all $i \geq N$ and a minimal index $k \geq 1$. We shall denote the k -periodic continued fraction by

$$[b_1, \dots, b_N, \overline{a_1, \dots, a_k}], \quad (2.2)$$

where (a_1, \dots, a_k) is the minimal period of (2.1). The continued fraction (2.2) converges to the irrational root of a quadratic polynomial

$$Ax^2 + Bx + C \in \mathbf{Z}[x]. \quad (2.3)$$

Conversely, the irrational root of any quadratic polynomial (2.3) has a representation by the continued fraction (2.2). Notice that the following two continued fractions define the same irrational number:

$$[b_1, \dots, b_N, \overline{a_1, \dots, a_k}] = [b_1, \dots, b_N, a_1, \dots, a_k, \overline{a_1, \dots, a_k}]. \quad (2.4)$$

But it is well known, that two infinite continued fraction with at most finite number of distinct entries must be related by the linear fractional transformation given by a matrix $\mathcal{E} \in GL_2(\mathbf{Z})$. Therefore equation (2.4) can be written in the form

$$x = \frac{E_{11}x + E_{12}}{E_{21}x + E_{22}}, \quad (2.5)$$

where $\mathcal{E} = (E_{ij}) \in GL_2(\mathbf{Z})$ and $x = [b_1, \dots, b_N, \overline{a_1, \dots, a_k}]$.

Remark 2.1. It is easy to see, that x in (2.5) is the root of quadratic polynomial (2.3) with $A = E_{21}$, $B = E_{22} - E_{11}$ and $C = -E_{12}$.

Definition 2.2. The Brock-Elkies-Jordan variety $V_{N,k}(\mathbf{C}) \subset \mathbb{A}^{N+k}$ is an affine variety over \mathbf{Z} defined by the three equations:

$$\begin{cases} A[E_{22} - E_{11}](y_1, \dots, y_N, x_1, \dots, x_k) = BE_{21}(y_1, \dots, y_N, x_1, \dots, x_k) \\ -AE_{12}(y_1, \dots, y_N, x_1, \dots, x_k) = CE_{21}(y_1, \dots, y_N, x_1, \dots, x_k) \\ -BE_{12}(y_1, \dots, y_N, x_1, \dots, x_k) = C[E_{22} - E_{11}](y_1, \dots, y_N, x_1, \dots, x_k). \end{cases}$$

It is verified directly from remark 2.1 and the equality $E_{11}E_{22} - E_{12}E_{21} = (-1)^k$, that

$$CE_{21}^2 - BE_{21}E_{22} + AE_{22}^2 = (-1)^k A. \quad (2.6)$$

Definition 2.3. By the Fermat-Pell conic \mathcal{Q} one understands the plane curve:

$$Cx^2 - Bxy + Ay^2 = (-1)^k A. \quad (2.7)$$

Theorem 2.4. (Brock, Elkies & Jordan [1]) *The affine variety $V_{N,k}(\mathbf{C})$ fibers over the Fermat-Pell conic \mathcal{Q} , i.e. there exists a map $\pi : V_{N,k}(\mathbf{C}) \rightarrow \mathcal{Q}$, such that*

$$\pi(y_1, \dots, y_N, x_1, \dots, x_k) = (E_{21}, E_{22}). \quad (2.8)$$

2.2. Arithmetic complexity. The noncommutative torus \mathcal{A}_θ is said to have real multiplication, if θ is the irrational root of a quadratic polynomial (2.3). To distinguish this case from the other values of θ , we shall write \mathcal{A}_{RM} . It is clear, that we can write θ as:

$$\theta_d = \frac{a + b\sqrt{d}}{c}, \quad (2.9)$$

where $a, b, c \in \mathbf{Z}$ and $d > 0$ is a square-free integer. Consider a family of the irrational numbers $\{\theta_x \mid x > 0\}$ of the form:

$$\theta_x := \left\{ \frac{a + b\sqrt{x}}{c} \mid x \in U_d; \quad a, b, c = Const \right\}, \quad (2.10)$$

where U_d is a set of the square-free integers containing d . A system of the polynomial equations in the ring $\mathbf{Z}[y_1, \dots, y_N, x_1, \dots, x_k]$ is called Euler's, if each $\{\theta_x \mid x \in U_d\}$ can be written as

$$\theta_x = [b_1(x), \dots, b_N(x), \overline{a_1(x), \dots, a_k(x)}], \quad (2.11)$$

where $a_i(x), b_j(x) \in \mathbf{Z}[x]$. The Euler equations define an algebraic set in the affine space \mathbb{A}^{N+k} . The Euler variety \mathcal{V}_E is defined as the projective closure of an irreducible affine variety containing the point $x = d$ of this set.

Remark 2.5. The case $a = 0$ and $c = 1$ of (2.10) was first studied by [Euler 1765] [2]; hence the name.

Definition 2.6. By an arithmetic complexity $c(\mathcal{A}_{RM})$ of the noncommutative torus \mathcal{A}_{RM} one understands the Krull dimension of the Euler variety \mathcal{V}_E .

Remark 2.7. The $c(\mathcal{A}_{RM})$ counts the number of the algebraically independent entries in the continued fraction $\theta_d = [b_1, \dots, b_N, \overline{a_1, \dots, a_k}]$.

Theorem 2.8. ([5, Theorem 6.2.1]) *The rank r of the elliptic curve $\mathcal{E}(K)$ is given by the formula:*

$$r = c(\mathcal{A}_{RM}) - 1, \quad (2.12)$$

where $\mathcal{A}_{RM} = F(\mathcal{E}(K))$.

3. PROOFS

3.1. Proof of theorem 1.1. We shall split the proof in a series of lemmas.

Lemma 3.1. *The connected component $V_{N,k}^0$ of the Brock-Elkies-Jordan variety $V_{N,k}(\mathbf{C})$ is a fiber bundle over the Fermat-Pell conic \mathcal{Q} :*

$$\pi : V_{N,k}^0 \rightarrow \mathcal{Q}, \quad (3.1)$$

such that each fiber $\{\pi^{-1}(q) \mid q \in \mathcal{Q}\}$ is an r -dimensional affine space \mathbb{A}^r .

Proof. Roughly speaking, the idea is to take the projective closure of (3.1) and apply [5, Lemma 6.2.2]. Such a lemma says the Euler variety \mathcal{V}_E is fiber bundle over the projective line $\mathbf{C}P^1$ with the fiber an abelian variety A_E .

(i) Let us show that

$$\text{Proj } V_{N,k}^0 \cong \mathcal{V}_E, \quad (3.2)$$

where *Proj* is the projective closure of the affine variety $V_{N,k}^0$. Indeed, recall that the $V_{N,k}^0$ is defined by the coefficients A_0, B_0 and C_0 of the quadratic polynomial (2.3). Using these coefficients, one can write (2.9) in the form:

$$\theta_d = \frac{-B_0 + b\sqrt{d}}{2A_0}, \quad \text{where } b^2d = B_0^2 - 4A_0C_0. \quad (3.3)$$

Thus the $V_{N,k}^0$ defines the Euler equations for the family θ_x and the Euler variety \mathcal{V}_E , see Section 2.2. Moreover, the equations in Definition 2.2 must coincide with the Euler equations for the family (2.11). We conclude from (3.3) that the $V_{N,k}^0$ is an irreducible affine variety containing the point $x = d$. Thus, $\text{Proj } V_{N,k}^0 \cong \mathcal{V}_E$ by the definition of the Euler variety \mathcal{V}_E in Section 2.2. Item (i) is proved.

(ii) To prove that $\{\pi^{-1}(q) \cong \mathbb{A}^r \mid q \in \mathcal{Q}\}$, let $(\mathcal{V}_E, \mathbf{CP}^1, \pi', A_E)$ be the fiber bundle constructed in [5, Lemma 6.2.2]. Consider a morphism of the following fiber bundles:

$$(V_{N,k}^0, \mathcal{Q}, \pi, X) \rightarrow (\mathcal{V}_E, \mathbf{CP}^1, \pi', A_E), \quad (3.4)$$

where X is a fiber $\pi^{-1}(q)$ over $q \in \mathcal{Q}$. Recall from item (i), that we have an isomorphism $\text{Proj } V_{N,k}^0 \cong \mathcal{V}_E$. Since the genus of the Fermat-Pell conic \mathcal{Q} is equal to zero, we conclude that $\text{Proj } \mathcal{Q} \cong \mathbf{CP}^1$. The fiber map π' is the projective extension of the map π .

To calculate the fiber X in (3.4), recall that the A_E is an abelian variety of dimension r [5, Lemma 6.2.4]. In particular, the A_E is a compact algebraic group (group variety). Thus one gets a group morphism

$$h : X \rightarrow A_E, \quad (3.5)$$

where X is an affine group variety. It is not hard to see, that $\ker(h) \cong \mathbf{Z}^r$, where \mathbf{Z}^r is a discrete subgroup of X . Thus $X \cong \mathbb{A}^r$ is the r -dimensional affine space. Item (ii) is proved.

To finish the proof of lemma 3.1, we apply [5, Lemma 6.2.2]. A pullback of the $(\mathcal{V}_E, \mathbf{CP}^1, \pi', A_E)$ along the morphism (3.4) defines a fiber bundle $(V_{N,k}^0, \mathcal{Q}, \pi, \mathbb{A}^r)$. Lemma 3.1 is proved. \square

Lemma 3.2. *The structure group of the fiber bundle $(V_{N,k}^0, \mathcal{Q}, \pi, \mathbb{A}^r)$ is isomorphic to the abelian group \mathcal{E}_{tors} .*

Proof. Let \mathcal{Q} be the Fermat-Pell conic defined by the equation (2.7). Since θ_d in formula (3.3) is a real number, the coefficients A, B, C must satisfy the inequality $B^2 - 4AC > 0$. Thus equation (2.7) defines a hyperbola. The completion of \mathcal{Q} by the point at infinity is homeomorphic to the unit circle $S^1 = \mathcal{Q} \cup \{\infty\}$. Denote by $(\tilde{V}_{N,k}^0, S^1, \tilde{\pi}, \mathbb{A}^r)$ the corresponding one-point completion of $(V_{N,k}^0, \mathcal{Q}, \pi, \mathbb{A}^r)$ by a fiber \mathbb{A}^r at the infinity.

Consider the universal cover \mathbf{R} of S^1 given by the formula

$$t \mapsto e^{2\pi it}, \quad t \in \mathbf{R}. \quad (3.6)$$

We denote by $(\mathbb{A}^r \times \mathbf{R}, \tilde{\pi}, \mathbb{A}^r)$ the pullback of the fiber bundle $(\tilde{V}_{N,k}^0, S^1, \tilde{\pi}, \mathbb{A}^r)$ along the map (3.6). Consider the integer points \mathbf{Z}^r of the affine space \mathbb{A}^r . It follows from [5, Lemma 6.2.2], that $\mathcal{E}(K)$ is embedded into $\mathbb{A}^r \times \mathbf{R}$ according to the formula:

$$\mathcal{E}(K) \cong \mathbf{Z}^r \oplus \mathcal{E}_{tors} \hookrightarrow \mathbb{A}^r \times \mathbf{R}/m\mathbf{Z}, \quad (3.7)$$

where $m = |\mathcal{E}_{tors}|$. Using formula (3.7) and the map

$$t \mapsto e^{2\pi imt}, \quad t \in \mathbf{R}, \quad (3.8)$$

we conclude that the fiber bundle $(\tilde{V}_{N,k}^0, S^1, \tilde{\pi}, \mathbb{A}^r)$ has the structure group

$$\mathbf{Z}/m\mathbf{Z} \cong \mathcal{E}_{tors}. \quad (3.9)$$

The same is true of the fiber bundle $(V_{N,k}^0, \mathcal{Q}, \pi, \mathbb{A}^r)$, provided the point ∞ of conic \mathcal{Q} is endowed with the fiber \mathbb{A}^r . Lemma 3.2 is proved. \square

Lemma 3.3. $c(\mathcal{A}_\theta) = \dim_{\mathbf{C}} V_{N,k}^0$.

Proof. Consider the fiber bundle $(V_{N,k}^0, \mathcal{Q}, \pi, \mathbb{A}^r)$. The dimension of a fiber bundle is the sum of dimensions of the base space and the fibers, i.e.

$$\dim_{\mathbf{C}} V_{N,k}^0 = \dim_{\mathbf{C}} \mathbb{A}^r + \dim_{\mathbf{C}} \mathcal{Q} = r + 1. \quad (3.10)$$

Comparing (3.10) with (2.12), we conclude that $c(\mathcal{A}_\theta) = \dim_{\mathbf{C}} V_{N,k}^0$. Lemma 3.3 is proved. \square

Theorem 1.1 follows from lemmas 3.1-3.3.

3.2. Proof of corollary 1.3. We shall split the proof in two lemmas.

Lemma 3.4. $\text{III}(\mathcal{Q}) \cong Cl(\Lambda) \oplus Cl(\Lambda)$.

Proof. We adapt the argument of [Lemmermeyer 2003] [4, Theorem 11]. Consider the order $\Lambda = \mathbf{Z} + \mathbf{Z}\theta$ in the ring of integers in the real quadratic field $k = \mathbf{Q}(\sqrt{D})$. Since θ is the root of quadratic equation (2.3), we conclude that $A = 1, B = 0$ and $C = -D$. Using (2.7) one can write the Fermat-Pell conic \mathcal{Q} in the form:

$$\mathbf{y}^2 - D\mathbf{x}^2 = (-1)^k. \quad (3.11)$$

According to [Lemmermeyer 2003] [4, Theorem 11], the Tate-Shafarevich group $\text{III}(\mathcal{Q})$ of the conic (2.7) is given by the formula:

$$\text{III}(\mathcal{Q}) \cong Cl(\mathcal{Q}) \oplus Cl(\mathcal{Q}), \quad (3.12)$$

where $Cl(\mathcal{Q})$ is the class group of \mathcal{Q} . But $Cl(\mathcal{Q})$ is equal to the narrow class group of the order Λ . Thus $\text{III}(\mathcal{Q}) \cong Cl(\Lambda) \oplus Cl(\Lambda)$. Lemma 3.4 is proved. \square

Lemma 3.5. $\text{III}(\mathcal{E}(K)) \cong Cl(\Lambda) \oplus Cl(\Lambda)$.

Proof. Using formula (3.7), one can identify the K -rational points of the elliptic curve $\mathcal{E}(K) \cong \mathbf{Z}^r \oplus \mathcal{E}_{tors}$ with the integer points of the fiber bundle $(V_{N,k}^0, \mathcal{Q}, \pi, \mathbb{A}^r)$. It is not hard to see, that the group $\text{III}(\mathbb{A}^r)$ of the fiber \mathbb{A}^r is trivial. We conclude therefore, that the failure of the Hasse principle for $\mathcal{E}(K)$ occurs only in the base space \mathcal{Q} . On the other hand, lemma 3.4 says that $\text{III}(\mathcal{Q}) \cong Cl(\Lambda) \oplus Cl(\Lambda)$. Thus $\text{III}(\mathcal{E}(K)) \cong Cl(\Lambda) \oplus Cl(\Lambda)$. Lemma 3.5 is proved. \square

Corollary 1.3 follows from lemma 3.5.

Remark 3.6. The formula $\text{III}(\mathcal{E}(K)) \cong Cl(\Lambda) \oplus Cl(\Lambda)$ can be proved in purely algebraic terms [6, Corollary 1.3]. However, the approach based on the Lemmermeyer's Lemma 3.4 and geometry of the fiber bundle $(V_{N,k}^0, \mathcal{Q}, \pi, \mathbb{A}^r)$ seems to be more elegant.

4. EXAMPLES

We shall consider two examples illustrating theorem 1.1. They correspond to the variety $V_{1,2}(\mathbf{C})$ [Brock, Elkies & Jordan 2019] [1, Section 8].

4.1. Brock-Elkies-Jordan variety $V_{1,2}(\mathbf{C})$. Consider the variety $V_{N,k}(\mathbf{C})$ with $N = 1$ and $k = 2$. According to [Brock, Elkies & Jordan 2019] [1, Section 8], in this case the variety $V_{1,2}(\mathbf{C})$ is defined by the system of equations:

$$\begin{cases} A(x_1x_2 - 2y_1x_1) = Bx_1 \\ A(y_1^2x_1 - y_1x_1x_2 - x_2) = Cx_1 \\ B(y_1^2x_1 - y_1x_1x_2 - x_2) = C(x_1x_2 - 2y_1x_1). \end{cases} \quad (4.1)$$

Theorem 4.1. ([Brock, Elkies & Jordan 2019] [1, Section 8]) *The Brock-Elkies-Jordan variety $V_{1,2}(\mathbf{C})$ has:*

- (i) one component $[y_1, \overline{0}, \overline{0}]$, if $A = 0 \neq B$ or $[y_1, \overline{0}, \overline{x_2}]$, if $A = B = 0$;
- (ii) two components $[y_1, \overline{0}, \overline{0}]$ and

$$\left[y_1, -\frac{2Ay_1 + B}{Ay_1^2 + By_1 + C}, \frac{2Ay_1 + B}{A} \right], \quad \text{if } A \neq 0; \quad (4.2)$$

- (iii) three components $[y_1, \overline{0}, \overline{0}]$, (4.2) and $[-\frac{B}{2A}, \overline{x_1}, \overline{0}]$, if $B^2 = 4AC$ and $A \neq 0$.

4.2. Arithmetic complexity $c(F(\mathcal{E}(K)))$. Consider the following two families of elliptic curves $\mathcal{E}(K)$.

Example 4.2. ([5, Section 1.4]) Let $b \geq 1$ be an integer. Denote by \mathcal{E}_{CM} an elliptic curve with complex multiplication by the integers of the imaginary quadratic field $\mathbf{Q}(i\sqrt{b^2 + 2})$. Then $F(\mathcal{E}_{CM}) = \mathcal{A}_{\sqrt{b^2+2}}$ [5, Theorem 1.4.1]. It is easy to see, that

$$\sqrt{b^2 + 2} = [b, \overline{b}, \overline{2b}]. \quad (4.3)$$

Therefore $N = 1$ and $k = 2$, i.e. the $V_{1,2}(\mathbf{C})$ is the Brock-Elkies-Jordan variety corresponding to the continued fraction (4.3). Recall that a classification of the connected components of the variety $V_{1,2}(\mathbf{C})$ is provided by Theorem 4.1. It is clear, that the connected components $[y_1, \overline{0}, \overline{0}]$, $[y_1, \overline{0}, \overline{x_2}]$ or $[-\frac{B}{2A}, \overline{x_1}, \overline{0}]$ in Theorem 4.1 must be excluded. Thus the component $V_{1,2}^0$ containing continued fraction (4.3) is given by the formulas (4.2). To calculate dimension of the component $V_{1,2}^0$, notice that the substitution

$$\begin{cases} y_1 = uv \\ A = 1 \\ B = 0 \\ C = -v(u^2v + 1), \end{cases} \quad (4.4)$$

brings component (4.2) to the form:

$$[uv, \overline{2u}, \overline{2uv}] = \sqrt{v(u^2v + 1)}. \quad (4.5)$$

The restriction $2u = b$, $v = 2$ shows that our fraction (4.3) is contained in the component $V_{1,2}^0$ described by (4.5). Moreover, the $V_{1,2}^0$ is the maximal component with such a property. Since the $V_{1,2}^0$ is parametrized by two complex variables u and v , we conclude that

$$\dim_{\mathbf{C}} V_{1,2}^0 = 2 = c(F(\mathcal{E}_{CM})). \quad (4.6)$$

Example 4.3. ([5, Section 6.2.4.2]) Let $b \geq 3$ be an integer. Denote by $\mathcal{E}(\mathbf{Q})$ an elliptic curve defined by the affine equation

$$y^2 = x(x-1) \left(x - \frac{b-2}{b+2} \right). \quad (4.7)$$

It is known ([5, Section 6.2.4.2]) that $F(\mathcal{E}(\mathbf{Q})) = \mathcal{A}_{\frac{1}{2}(b+\sqrt{b^2-4})}$. The reader can verify that

$$\frac{b + \sqrt{b^2 - 4}}{2} = [b - 1, \overline{1, b - 2}]. \quad (4.8)$$

Thus the $V_{1,2}(\mathbf{C})$ is the Brock-Elkies-Jordan variety corresponding to the continued fraction (4.8). Clearly, the connected components $[y_1, \overline{0, 0}]$, $[y_1, \overline{0, x_2}]$ or $[-\frac{B}{2A}, \overline{x_1, 0}]$ in Theorem 4.1 are excluded. Therefore the connected component $V_{1,2}^0$ containing periodic continued fraction (4.8) is given by formula (4.2). Using the substitution

$$\begin{cases} y_1 &= u - 1 \\ A &= 1 \\ B &= -u \\ C &= 1, \end{cases} \quad (4.9)$$

one can write (4.2) in the form:

$$[u - 1, \overline{1, u - 2}] = \frac{u + \sqrt{u^2 - 4}}{2}. \quad (4.10)$$

The continued fractions (4.8) and (4.10) are identical after the substitution $u = b$. Thus (4.8) parametrizes the component $V_{1,2}^0$. Moreover, the $V_{1,2}^0$ is the maximal component with such a property. Since the $V_{1,2}^0$ depends on one complex variable u , we conclude that

$$\dim_{\mathbf{C}} V_{1,2}^0 = 1 = c(F(\mathcal{E}(\mathbf{Q}))). \quad (4.11)$$

Remark 4.4. Formula (4.11) is valid only for the odd values of $b \geq 3$ in (4.8). Indeed, the substitution $b = 2k$ brings the LHS of (4.8) to the form $k + \sqrt{k^2 - 1}$. The latter is a special case of the RHS of the family (4.5) with $u = k$ and $v = -1$. Thus the even values of $b \geq 3$ must be excluded.

4.3. Ranks of elliptic curves $\mathcal{E}(K)$. The ranks of elliptic curves \mathcal{E}_{CM} and $\mathcal{E}(\mathbf{Q})$ can be evaluated using formulas (1.2), (4.6) and (4.11).

Example 4.5. Let the \mathcal{E}_{CM} be as in example 4.2. The formulas (1.2) and (4.6) say that

$$r(\mathcal{E}_{CM}) = c(F(\mathcal{E}_{CM})) - 1 = 2 - 1 = 1. \quad (4.12)$$

Remark 4.6. The arithmetic of elliptic curves \mathcal{E}_{CM} has been studied by [Gross 1980] [3]. In particular, it was proved that $r(\mathcal{E}_{CM}) = 1$, if \mathcal{E}_{CM} is an elliptic curve with complex multiplication by the integers of the field $\mathbf{Q}(\sqrt{-p})$, where $p \equiv 3 \pmod{8}$ is a prime number [Gross 1980] [3, Theorem 22.4.2]. The reader can verify, that letting $b = 2k + 1$ in formula (4.3) implies $b \equiv 3 \pmod{8}$. Therefore formula (4.12) is a generalization of the result of [Gross 1980] [3, Theorem 22.4.2]. We refer the reader to [5, Table 1.1] for more examples.

Example 4.7. Let the $\mathcal{E}(\mathbf{Q})$ be as in example 4.3. One gets using (1.2), (4.11) and remark 4.4, that

$$r(\mathcal{E}(\mathbf{Q})) = c(F(\mathcal{E}(\mathbf{Q}))) - 1 = 1 - 1 = 0. \quad (4.13)$$

Thus the rank of elliptic curve (4.7) must be zero for each odd integer $b \geq 3$.

REFERENCES

1. B. W. Brock, N. D. Elkies and B. W. Jordan, *Periodic continued fractions over S -integers in number fields and Skolem's p -adic method*, arXiv:1909.11214v2
2. L. Euler, *De usu novi algorithmi in Problemate Pelliano solvendo*, Novi Commentarii Academiae Scientiarum Imperialis Petropolitanae **11**, 1765.
3. B. H. Gross, *Arithmetic on Elliptic Curves with Complex Multiplication*, Lecture Notes Math. **776** (1980), Springer.
4. F. Lemmermeyer, *Conics – a poor man's elliptic curves*, arXiv:math/0311306
5. I. V. Nikolaev, *Noncommutative Geometry*, De Gruyter Studies in Math. **66**, Berlin, 2017.
6. I. V. Nikolaev, *Quantum dynamics of elliptic curves*, arXiv:1808.03493
7. M. A. Rieffel, *Non-commutative tori – a case study of non-commutative differentiable manifolds*, Contemp. Math. **105** (1990), 191-211.

¹ DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, ST. JOHN'S UNIVERSITY, 8000 UTOPIA PARKWAY, NEW YORK, NY 11439, UNITED STATES.

E-mail address: igor.v.nikolaev@gmail.com