

NON-DENSITY OF POINTS OF SMALL ARITHMETIC DEGREES

YOSUKE MATSUZAWA, SHENG MENG, TAKAHIRO SHIBATA, AND DE-QI ZHANG

ABSTRACT. Given a surjective endomorphism $f : X \rightarrow X$ on a projective variety over a number field, one can define the arithmetic degree $\alpha_f(x)$ of f at a point x in X . The Kawaguchi–Silverman Conjecture (KSC) predicts that any forward f -orbit of a point x in X at which the arithmetic degree $\alpha_f(x)$ is strictly smaller than the first dynamical degree δ_f of f is not Zariski dense. We extend the KSC to sAND (= small Arithmetic Non-Density) Conjecture that the locus $Z_f(d)$ of all points of small arithmetic degree is not Zariski dense, and verify this sAND Conjecture for endomorphisms on projective varieties including surfaces, HyperKähler varieties, abelian varieties, Mori dream spaces, simply connected smooth varieties admitting int-amplified endomorphisms, smooth threefolds admitting int-amplified endomorphisms, and some fiber spaces. We also show close relations between our sAND Conjecture and the Uniform Boundedness Conjecture of Morton and Silverman on endomorphisms of projective spaces and another long standing conjecture on Uniform Boundedness of torsion points in abelian varieties.

CONTENTS

1. Introduction	2
2. Preliminaries	5
3. Polarized endomorphisms, curves and Mori dream spaces	12
4. Surfaces and Hyperkähler varieties	13
5. Abelian varieties	15
6. Varieties admitting int-amplified endomorphisms	20
7. Fiber spaces; UBC vs sAND; Proofs of Theorems 1.10 and 1.13	32
8. Further Generalization	40
References	41

2010 *Mathematics Subject Classification.* Primary: 37P55, Secondary: 14G05, 37B40.

Key words and phrases. Dynamical degree, Arithmetic degree, Kawaguchi-Silverman conjecture, small Arithmetic Non-Density conjecture, Uniform boundedness conjectures for \mathbb{P}^n and abelian varieties.

1. INTRODUCTION

Let $f : X \rightarrow X$ be a surjective endomorphism of a geometrically irreducible projective variety X defined over a number field K . Fix an algebraic closure \overline{K} of K . One may consider two dynamical invariants: the (first) *dynamical degree* δ_f of f and the *arithmetic degree* $\alpha_f(x)$ of f at $x \in X(\overline{K})$. The dynamical degree reflects the geometric complexity of iterations of f . On the other hand, the arithmetic degree reflects the arithmetic complexity of the given f -orbit $O_f(x) := \{x, f(x), f^2(x), \dots\}$. For definitions, see 2.1.

One naturally likes to compare these two invariants. By Kawaguchi–Silverman [KS16a], the arithmetic degree at any point is less than or equal to the dynamical degree (see also [Mat16]). Then, one wonders when the arithmetic degree at a point attains (its upper bound) the dynamical degree. Kawaguchi and Silverman proposed (see [KS16a]):

Conjecture 1.1. (KSC) *Let $f : X \rightarrow X$ be a surjective endomorphism of a projective variety X defined over a number field K . Let $x \in X(\overline{K})$ such that $\alpha_f(x) < \delta_f$. Then the orbit $O_f(x)$ is not Zariski dense in $X_{\overline{K}} := X \times_K \overline{K}$.*

Although the original Conjecture in [KS16a] has also been formulated for rational maps, we consider only endomorphisms in this paper (see Remark 1.7).

Given a surjective endomorphism $f : X \rightarrow X$ on a projective variety X over a number field K , let $Z_f \subseteq X(\overline{K})$ be the set of points $x \in X(\overline{K})$ where $\alpha_f(x) < \delta_f$. Conjecture 1.1 predicts that the orbit $O_f(x)$ is not dense for any $x \in Z_f$. So one wonders whether the set Z_f is also non-dense. Unfortunately, the answer is negative by the following:

Example 1.2. Let the morphism $f : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ be given by $f(x) = x^2$ in affine coordinate. Then the set

$$f^{-\infty}(1) := \bigcup_{n=0}^{\infty} f^{-n}(1) = \bigcup_{n=0}^{\infty} \{x \in \overline{\mathbb{Q}} \mid x^{2^n} = 1\}$$

consists of f -preperiodic points and is infinite. Clearly any f -preperiodic point has arithmetic degree one while $\delta_f = 2$, so $f^{-\infty}(1) \subseteq Z_f$. Hence Z_f is a dense subset of \mathbb{P}^1 .

We now recall the Northcott finiteness property. Let X be a projective variety over a number field K and H an ample Cartier divisor on X . Take a height function h_H associated to H . Then the Northcott finiteness property asserts that the set

$$\{x \in X(\overline{K}) \mid [K(x) : K] \leq d, h_H(x) \leq M\}$$

is finite for any given $d > 0$ and $M > 0$. This suggests the non-density of a subset of Z_f when we impose a bound on the degree $[K(x) : K]$ for points x in Z_f . In view of this observation, we are going to propose Conjecture 1.4 as follows.

Definition 1.3. Let X be a projective variety and f a surjective endomorphism on X over a number field K . Let $d > 0$. Set

$$Z_f = \{x \in X(\overline{K}) \mid \alpha_f(x) < \delta_f\},$$

$$Z_f(d) = \{x \in X(L) \mid K \subseteq L \subseteq \overline{K}, [L : K] \leq d, \alpha_f(x) < \delta_f\}.$$

Conjecture 1.4. (sAND = small Arithmetic Non-Density) *Let $f : X \rightarrow X$ be a surjective endomorphism of a projective variety X defined over a number field K . Then the set $Z_f(d)$ is not Zariski dense in $X_{\overline{K}} = X \times_K \overline{K}$ for any positive constant $d > 0$.*

Remark 1.5. In [Shi19], the third author defined a similar notion $Z_f(K)$ and Conjectured the non-density of it ([Shi19, Conjecture 1.3]). It is the set of points at which the ample height growth does not grow maximally for iterations of f . Clearly, it is larger than the locus we defined in Definition 1.3, so Conjecture 1.4 is weaker than [Shi19, Conjecture 1.3]. However, it is easier for us to handle the arithmetic degree than the height growth. In addition, we can consider arithmetic degrees for rational self-maps or endomorphisms on non-projective varieties. So we propose the present formulation in this paper.

Remark 1.6. Let $K \subseteq L (\subseteq \overline{K})$ be a finite extension. Then Conjecture 1.4 for $f : X \rightarrow X$ is equivalent to that of $f_L : X_L \rightarrow X_L$.

The sAND Conjecture 1.4 implies KSC Conjecture 1.1 for endomorphisms. Indeed, for any $x \in Z_f$, we can take a suitable $d > 0$ such that the orbit $O_f(x) \subseteq Z_f(d)$. Hence the sAND Conjecture 1.4 is a generalization of the KSC Conjecture 1.1.

Remark 1.7. In sAND Conjecture 1.4, it is important to assume that f is a well defined morphism. Indeed, Call and Silverman [CS16, Proposition 13 (a)] mentioned Junyi Xie's construction of a family $f = (f_t)_{t \in \mathbb{P}^1}$ of rational self-maps on \mathbb{P}^2 such that the dynamical degree drops when $t = 1, 2, 3, \dots \in \mathbb{A}^1(\mathbb{Q})$, and hence $Z_f(d)$ is Zariski dense.

Another motivation for us to extend KSC Conjecture 1.1 to sAND Conjecture 1.4 is in order to cover the case of a endomorphism on a fiber space which fixes fibers set-theoretically. Such a surjective endomorphism has no dense orbit, so the KSC Conjecture 1.1 vacuously holds. But the sAND Conjecture 1.4 is still non-trivial in this case. In fact, we will prove sAND Conjecture 1.4 for some special fibrations in Section 7.

Theorems 1.8, 1.10, and 1.13 below are our main results; see also Theorem 3.1 for polarized endomorphisms, Theorem 7.4 for endomorphisms of projective bundles fixing the base pointwise, and Theorem 7.11 for non-invertible surjective endomorphisms on smooth projective threefold of non-negative Kodaira dimension, as well as §8 for further generalization of Conjecture 1.4 to Conjecture 8.1.

From the classification theory of algebraic varieties and the typical fibrations: Albanese map, Iitaka fibration, and Maximal rationally connected fibration, which all can be chosen to be f -equivariant, we see that the building blocks of varieties are abelian varieties, rationally connected varieties and varieties of Kodaira dimension zero; further, each variety of the last type, if smooth and minimal, has an (equivariant) Beauville–Bogomolov étale cover by a product of Abelian varieties, Hyperkähler varieties and Calabi–Yau varieties. Our result below shows that Conjecture 1.4 holds in many cases, and it covers the three types of building blocks: Abelian varieties, Hyperkähler varieties and Rationally connected varieties (but not Calabi–Yau varieties).

Theorem 1.8. *The sAND Conjecture 1.4 holds for every surjective endomorphism on any projective variety X which fits one of the following cases.*

- (1) $X_{\overline{K}}$ is a Mori dream space (cf. Theorem 3.2).
- (2) $\dim(X) \leq 2$ (cf. Theorems 4.1 and 4.3).
- (3) $X_{\overline{K}}$ is a Hyperkähler variety (cf. Theorem 4.2).
- (4) $X_{\overline{K}}$ is a (Q) -abelian variety (cf. Corollary 5.9).
- (5) $X_{\overline{K}}$ is a smooth rationally connected projective variety admitting an int-amplified endomorphism (cf. Theorem 6.3, Corollary 6.19).
- (6) $X_{\overline{K}}$ is a smooth projective threefold admitting an int-amplified endomorphism (cf. Theorem 6.6, Corollary 6.19).

Let us recall the uniform boundedness conjecture due to Morton–Silverman.

Conjecture 1.9. (UBC = Uniform Boundedness Conjecture for \mathbb{P}^N , [MS94]) *Fix $r \in \mathbb{Z}_{\geq 2}$, $N \in \mathbb{Z}_{\geq 1}$, and $d \in \mathbb{Z}_{\geq 1}$. Then there exists a positive constant $C = C(r, N, d) > 0$ such that the following holds: for any finite extension $K \subseteq L (\subseteq \overline{K})$ with $[L : K] \leq d$ and any morphism $\phi : \mathbb{P}^N \rightarrow \mathbb{P}^N$ which is defined over L with $\deg \phi = r$, the number of ϕ -preperiodic L -rational points of \mathbb{P}^N is at most C .*

We show the equivalence of the above Uniform Boundedness Conjecture and a special case of a relative version of the sAND Conjecture 7.1: the non-density of $Z_f(d)$ for a family of polarized endomorphisms, which suggests the importance and naturality of conjecturing the non-density of the locus $Z_f(d)$ of small arithmetic degree.

Theorem 1.10. *The following statements are equivalent.*

- (1) *The Uniform Boundedness Conjecture 1.9.*
- (2) *Let $\pi : X \rightarrow S$ be a projective morphism of varieties and $f : X \rightarrow X$ a surjective morphism satisfying $\pi \circ f = \pi$. Assume that there is a π -ample divisor H on X such that $f^*H \sim_{\pi} rH$ for some $r > 1$. Let $d > 0$. Then there is a constant $C > 0$ (depending on d) such that $|Z_{f|_{X_s}}(d)| < C$ for any $s \in S(d)$.*

- (3) Let $\pi : X \rightarrow S$ be a projective morphism of quasi-projective varieties and $f : X \rightarrow X$ a surjective morphism satisfying $\pi \circ f = \pi$. Assume that there is a π -ample divisor H on X such that $f^*H \sim_\pi rH$ for some $r > 1$. Let $d > 0$. Then $Z_f(d)$ is not dense.

Our sAND conjecture is also related to another long-standing conjecture which is completely solved only in dimension one by Merel [Mer96, Corollaire].

Conjecture 1.11. (UBC = Uniform Boundedness Conjecture for Torsion Points) Take any $N \in \mathbb{Z}_{\geq 1}$ and $d > 0$. Then there is a constant $C = C(N, d) > 0$ such that the number of torsion points in $A(d)$ is at most C for any abelian variety A of dimension N defined over K .

Remark 1.12. (1) Conjecture 1.9 implies Conjecture 1.11 (cf. [Fak03, Corollary 2.4]).
 (2) Conjecture 1.11 is usually stated as follows:

(*) Take any $N \in \mathbb{Z}_{\geq 1}$ and $d > 0$. Then there is a constant $C = C(N, d) > 0$ such that the order of $\text{Tor}(A(L))$ is at most C for any abelian variety A of dimension N over K and any finite extension $K \subseteq L (\subseteq \overline{K})$ with $[L : K] \leq d$.

In fact, (*) is equivalent to Conjecture 1.11. For, assuming (*), a sufficiently large integer m depending only on N and d annihilates all torsion points of $A(L)$ for any abelian variety A of dimension N and any finite extension $K \subseteq L (\subseteq \overline{K})$ with $[L : K] \leq d$. So m annihilates all torsion points contained in $A(d)$. This implies that the number of torsion points in $A(d)$ is at most m^{2N} .

As an evidence of our sAND Conjecture (1.4 or 7.1), we show that Conjecture 1.11 implies Conjecture 7.1 for abelian fibrations.

Theorem 1.13. Let $\pi : X \rightarrow Y$ be an abelian fibration of quasi-projective varieties and $f : X \rightarrow X$ a surjective morphism such that $\delta_f > 1$ and $\pi \circ f = \pi$. Assume that there is a Zariski dense open subset $U \subseteq Y$ such that for any $d > 0$ and $d' > 0$, there is a constant $C = C(d, d') > 0$ such that the number of torsion points in $X_y(d)$, where X_y is the fiber over $y \in U(d')$, is at most C . Then for any $d > 0$, $Z_f(d)$ is not dense in $X_{\overline{K}}$.

Acknowledgments. The first, second, third and last authors are supported by a JSPS Overseas Research Fellowship, a Research Fellowship of KIAS, a Research Fellowship of NUS, and an ARF of NUS, respectively. The authors would like to thank Prof. Kawaguchi for kindly pointing out Remark 1.12 (1).

2. PRELIMINARIES

In this section, we prepare some notions and lemmas needed.

2.1. Notation, Convention and Terminology

- Throughout this article, we work over a fixed number field K unless otherwise stated. We fix an algebraic closure \overline{K} of K . The base change to \overline{K} of some object A defined over K is denoted by $A_{\overline{K}}$.
- A *variety* means a geometrically integral separated scheme of finite type over K .
- An *endomorphism on X* is a morphism over K from X to itself.
- Let X be a projective variety and f a surjective endomorphism on X .
 - (1) The (*forward*) f -*orbit* of a point x in $X(\overline{K})$ is the set $O_f(x) = \{f^s(x) \mid s \geq 0\}$.
 - (2) A point x in $X(\overline{K})$ is f -*periodic* if $f^n(x) = x$ for a positive integer n . $\text{Per}(f)$ denotes the set of f -periodic \overline{K} -rational points of X .
 - (3) A point x in $X(\overline{K})$ is f -*preperiodic* if $f^k(x)$ is f -periodic for a positive integer k . $\text{Prep}(f)$ denotes the set of f -preperiodic \overline{K} -rational points of X .
We remark that x is f -preperiodic if and only if $O_f(x)$ is finite. Further, if f is an automorphism, then x is f -preperiodic if and only if x is f -periodic.
 - (4) A closed subset $V \subseteq X$ is f -*invariant* (resp. f^{-1} -*invariant*) if $f(V) \subseteq V$ (resp. $f^{-1}(V) \subseteq V$). Similarly V is f -*periodic* if $f^n(V) \subseteq V$ for some $n > 0$ and f -*preperiodic* if $f^{n+k}(V) \subseteq f^k(V)$ for some $n, k > 0$.
- Let X be a projective variety and f a surjective endomorphism on X . The (*first*) *dynamical degree* δ_f of f is defined to be the (first) dynamical degree of $f_{\overline{K}}$. Namely, it is the following limit, with H any ample divisor

$$\delta_f = \lim_{n \rightarrow +\infty} ((f^n)^* H \cdot H^{\dim(X)-1})^{1/n}.$$

- Let $f : X \rightarrow X$ be a surjective endomorphism on a projective variety. Let $Z \subseteq X_{\overline{K}}$ be an f -preperiodic subvariety.
 - (1) Suppose $f^{m+k}(Z) = f^k(Z)$ for some $m, k > 0$. Define the *dynamical degree of f along Z* as

$$\delta_{f|_Z} = (\delta_{f^m|_{f^k(Z)}})^{\frac{1}{m}}.$$
 Clearly this definition is independent of m and k . If Z is f -invariant, then this invariant coincides with the dynamical degree of $f|_Z$.
 - (2) We call Z a f -*preperiodic subvariety of small dynamical degree* if $\delta_{f|_Z} < \delta_f$. Moreover we say Z is a *maximal f -preperiodic subvariety of small dynamical degree* if Z is maximal among all of the f -preperiodic subvarieties of small dynamical degree.
- The symbols \sim (resp. $\sim_{\mathbb{Q}}$, $\sim_{\mathbb{R}}$) and \equiv (resp. \equiv_w) mean the linear equivalence (resp. \mathbb{Q} -linear equivalence, \mathbb{R} -linear equivalence) and the numerical equivalence (resp. weak numerical equivalence) on divisors (cf. [MZ18, Definition 2.2]).

- A surjective morphism $f : X \rightarrow X$ is q -polarized (or just polarized) if $f^*H \sim qH$ for some ample divisor H and integer $q > 1$ (or equivalently if $f^*B \equiv qB$ for some big \mathbb{R} -divisor B ; cf. [MZ18, Proposition 3.6]).
- A surjective morphism $f : X \rightarrow X$ is *int-amplified* if $f^*L - L = H$ for some ample divisors L and H (or equivalently if $f^*L - L = H$ for some big \mathbb{R} -divisors L and H ; cf. [Men17, Theorem 1.1]).
- A finite surjective morphism is *quasi-étale* if it is étale in codimension one.
- A normal projective variety X over \overline{K} is Q -abelian, if there is a finite quasi-étale morphism $A \rightarrow X$ from an abelian variety A . A normal projective variety X over K is Q -abelian if so is $X_{\overline{K}}$.
- For a projective variety X over \overline{K} , $\text{NS}(X) := \text{Pic}(X)/\text{Pic}^0(X)$ denotes the *Néron-Severi* group of X . Let $\mathbb{K} = \mathbb{Q}, \mathbb{R}$ or \mathbb{C} . Set $\text{NS}(X)_{\mathbb{K}} := \text{NS}(X) \otimes_{\mathbb{Z}} \mathbb{K}$. Set $\text{N}^1(X) := \text{NS}(X)_{\mathbb{R}}$ and $\rho(X) = \dim_{\mathbb{R}} \text{N}^1(X)$. The number $\rho(X)$ is called the *Picard number of X* . Denote by $\text{Nef}(X)$ the cone of nef divisors in $\text{N}^1(X)$. Denote by $\text{PE}^1(X)$ the cone of pseudo-effective divisors in $\text{N}^1(X)$.
- Let $\mathbb{K} = \mathbb{Q}, \mathbb{R}$ or \mathbb{C} . For a \mathbb{K} -linear endomorphism $\phi : V \rightarrow V$ on a \mathbb{K} -vector space V , $\rho(\phi)$ denotes the *spectral radius* of f , that is, the maximum of absolute values of eigenvalues (in \mathbb{C}) of ϕ .
- Let $f : X \rightarrow X$ be a surjective endomorphism of a projective variety X . Then it is known that the first dynamical degree $\delta(f)$ is equal to $\rho(f^*|_{\text{NS}(X)_{\mathbb{C}}})$.
- Let f, g and h be \mathbb{R} -valued functions on a domain. Denote $f = g + O(h)$ if there is a positive constant C such that $|f - g| \leq C|h|$. In particular, denote $f = g + O(1)$ if there is a positive constant C such that $|f - g| \leq C$.
- Let X be a projective variety. For an \mathbb{R} -Cartier \mathbb{R} -divisor D on X , we can attach a function $h_D : X(\overline{K}) \rightarrow \mathbb{R}$ called logarithmic Weil height function. The function h_D is called the *height function associated to D* . Note that to construct h_D from D , we need additional information, for example an adelic metric on the associate line bundle if D is integral, but the resulting function is unique up to bounded functions. For definition and properties of height functions, see e.g. [HS00, Part B] or [Lan83, Chapter 3].
- Let X be a normal projective variety and f a surjective endomorphism on X . Fix an ample height function $h_H \geq 1$. Then, for every $x \in X(\overline{K})$, the limit

$$\alpha_f(x) = \lim_{n \rightarrow +\infty} h_H(f^n(x))^{1/n}$$

exists and is independent of the choice of ample height function; we call $\alpha_f(x)$ the *arithmetic degree of f at x* . (cf. [KS16a], [KS16b, Theorem 3 (a)] for details).

- Let $f : X \rightarrow X$ be a surjective endomorphism on a projective variety X and D an \mathbb{R} -Cartier divisor on X . Take a height function h_D associated to D . Fix an ample Cartier divisor H on X and an associated height function $h_H \geq 1$.

(1) Assume that $f^*D \sim_{\mathbb{R}} \lambda D$ with $\lambda > 1$. Then, for any $x \in X(\overline{K})$, the limit

$$\hat{h}_{D,f}(x) = \lim_{n \rightarrow +\infty} \frac{h_D(f^n(x))}{\lambda^n}$$

converges and satisfies $\hat{h}_{D,f} = h_D + O(1)$ (cf. [CS93, Theorem 1.1]).

(2) Assume that $f^*D \equiv \lambda D$ with $\lambda > \sqrt{\delta_f}$. Then, for any $x \in X(\overline{K})$, the limit

$$\hat{h}_{D,f}(x) = \lim_{n \rightarrow +\infty} \frac{h_D(f^n(x))}{\lambda^n}$$

converges and satisfies $\hat{h}_{D,f} = h_D + O(\sqrt{h_H})$ (cf. [KS16a, Theorem 5]).

- Given a variety X and $d > 0$, set

$$X(d) = \bigcup_{K \subseteq L \subseteq \overline{K}, [L:K] \leq d} X(L).$$

- For an abelian group G , $\text{Tor}(G)$ denotes the set of torsion elements of G .
- Let A be an abelian variety over \overline{K} .

(1) For a Cartier divisor D on A , let $\varphi_D : A \rightarrow \hat{A}; x \mapsto [T_x^*D - D]$, where \hat{A} is the dual abelian variety and T_x is the translation by x . Note that $D \in \text{Pic}^0(A)$ if and only if $\varphi_D = 0$ (this is also adopted by Mumford as the definition of $\text{Pic}^0(A)$ in his book). We get an \mathbb{R} -linear map $\text{NS}(A)_{\mathbb{R}} \rightarrow \text{Hom}(A, \hat{A}) \otimes_{\mathbb{Z}} \mathbb{R}$. Note that if D is ample, then φ_D is an isogeny.

(2) For an ample divisor H on A , set

$$\Phi^H : \text{NS}(A)_{\mathbb{R}} \rightarrow \text{End}(A)_{\mathbb{R}}; D \mapsto \Phi_D^H = \varphi_H^{-1} \circ \varphi_D.$$

(3) Denote the *Rosati involution* with respect to an ample divisor H by

$$i_H : \text{End}(A)_{\mathbb{R}} \rightarrow \text{End}(A)_{\mathbb{R}}; \alpha \mapsto \varphi_H^{-1} \circ \hat{\alpha} \circ \varphi_H$$

where $\hat{\alpha}$ is the dual of α in $\text{End}(\hat{A})_{\mathbb{R}}$.

- For a generically finite surjective morphism $\pi : X \rightarrow Y$ between varieties over \overline{K} , denote by $\text{Exc}(\pi)$ the π -*exceptional locus*, i.e., the union of curves in X contracted by π .

The following lemma follows from the product formula (cf. e.g. [Tru15]).

Lemma 2.2. *Let $\pi : X \dashrightarrow Y$ be a dominant rational map between projective varieties, and $f : X \rightarrow X$, $g : Y \rightarrow Y$ surjective morphisms such that $\pi \circ f = g \circ \pi$. Then we have:*

- (1) $\delta_f \geq \delta_g$.
- (2) If $\dim(X) = \dim(Y)$, then $\delta_f = \delta_g$.

The following two lemmas enable us to replace a given endomorphism by its positive power or direct summands, respectively.

Lemma 2.3. *Let $f : X \rightarrow X$ be a surjective endomorphism of a projective variety. Take a positive integer N . Then $Z_f = Z_{f^N}$. In particular, Conjecture 1.4 holds for f if and only if it holds for f^N .*

Proof. The lemma follows from that $\delta_{f^N} = \delta_f^N$ and $\alpha_{f^N}(x) = \alpha_f^N(x)$. \square

Lemma 2.4. *Let X, Y be projective varieties and f, g surjective endomorphisms on X, Y , respectively. Then*

$$Z_{f \times g} = \begin{cases} Z_f \times Y & \text{if } \delta_f > \delta_g, \\ Z_f \times Z_g & \text{if } \delta_f = \delta_g, \\ X \times Z_g & \text{if } \delta_f < \delta_g. \end{cases}$$

In particular, Conjecture 1.4 holds if it holds for f and g .

Proof. We may assume that $\delta_f \geq \delta_g$, without loss of generality. We can easily show (e.g. by the product formula) that $\delta_{f \times g} = \max(\delta_f, \delta_g) = \delta_f$.

Let $p : X \times Y \rightarrow X$, $q : X \times Y \rightarrow Y$ be the projections. Take ample divisors H_X, H_Y on X, Y , respectively. Set $H = p^*H_X + q^*H_Y$. Then

$$\begin{aligned} \alpha_{f \times g}(x, y) &= \lim_{n \rightarrow +\infty} h_H((f \times g)^n(x, y))^{1/n} \\ &= \lim_{n \rightarrow +\infty} \{h_{H_X}(f^n(x)) + h_{H_Y}(g^n(y))\}^{1/n} \\ &= \max(\alpha_f(x), \alpha_g(y)). \end{aligned}$$

For the last equality, see [San16, Lemma 3.1]. So $(x, y) \in Z_{f \times g}$ if and only if $\max(\alpha_f(x), \alpha_g(y)) < \max(\delta_f, \delta_g)$.

If $\delta_f > \delta_g$, then $(x, y) \in Z_{f \times g}$ if and only if $\alpha_f(x) < \delta_f$. Hence $Z_{f \times g} = Z_f \times Y$. If $\delta_f = \delta_g$, then $(x, y) \in Z_{f \times g}$ if and only if $\alpha_f(x) < \delta_f$ and $\alpha_g(y) < \delta_g$. So $Z_{f \times g} = Z_f \times Z_g$. \square

The following is easy but fundamental for the reduction to invariant subvarieties.

Lemma 2.5. *Let $f : X \rightarrow X$ be a surjective endomorphism on a projective variety X and $W \subseteq X$ an f -invariant closed subvariety. Then $\alpha_{f|_W}(x) = \alpha_f(x)$ for any $x \in W(\overline{K})$.*

Proof. Let $\iota : W \rightarrow X$ be the inclusion. For an ample divisor H on X , the restriction $H|_W$ is also ample and $h_{H|_W} = h_H \circ \iota$. So the assertion holds. \square

The following lemma is useful when dealing with equivariant fibrations or covers.

Lemma 2.6. *Let X, Y be projective varieties and f, g surjective endomorphisms on X, Y , respectively. Let $\pi : X \rightarrow Y$ be a surjective morphism such that $\pi \circ f = g \circ \pi$. Then:*

- (1) $\alpha_f(x) \geq \alpha_g(\pi(x))$ for any $x \in X(\overline{K})$.
- (2) Assume that $\delta_f = \delta_g$. Then $\pi(Z_f) \subseteq Z_g$. Moreover, Conjecture 1.4 holds for f if it holds for g .
- (3) Assume that π is finite. Then $\delta_f = \delta_g$, $\alpha_f(x) = \alpha_g(\pi(x))$, and $\pi(Z_f) = Z_g$. Moreover, Conjecture 1.4 holds for f if and only if it holds for g .

Proof. (1) Take ample divisors H, A on X, Y respectively such that $H - \pi^*A$ is ample. Then we can take associated height functions $h_H, h_A \geq 1$ as satisfying $Ch_H \geq h_A \circ \pi$ for some $C > 0$. We have

$$h_A(g^n(\pi(x)))^{1/n} = h_A(\pi(f^n(x)))^{1/n} \leq (Ch_H(f^n(x)))^{1/n}.$$

Letting $n \rightarrow +\infty$, we obtain $\alpha_g(\pi(x)) \leq \alpha_f(x)$.

(2) Clearly $\pi(Z_f) \subseteq Z_g$ by (1). Take any $d > 0$. Then $\pi(Z_f(d)) \subseteq Z_g(d)$. So the non-density of $Z_g(d)$ implies the non-density of $Z_f(d)$.

(3) By Lemma 2.2, $\delta_f = \delta_g$. Since π is finite, π^*A is also ample. This implies that $\alpha_f(x) = \alpha_g(\pi(x))$ for any $x \in X(\overline{K})$. So $\pi(Z_f) = Z_g$.

Take any $d > 0$. Since π is a finite cover, there is some $d' > 0$ such that $\pi^{-1}(Y(d)) \subseteq X(d')$. So $\pi^{-1}(Z_g(d)) \subseteq Z_f(d')$. Therefore the non-density of $Z_f(d')$ implies the non-density of $Z_g(d)$. \square

Below is a description of the closure of an orbit.

Lemma 2.7. *Let $f : X \rightarrow X$ be a surjective endomorphism on a projective variety X over \overline{K} and $x \in X(\overline{K})$. Then for some $s \geq 1$ and $t \geq 1$,*

$$\overline{O_f(x)} = \{x\} \cup \{f(x)\} \cup \cdots \cup \{f^{t-1}(x)\} \cup \bigcup_{i=0}^{s-1} f^i(\overline{O_{f^s}(f^t(x))}),$$

which is a union of irreducible closed subsets and $\overline{O_{f^s}(f^t(x))}$ is f^s -invariant.

Proof. Let $Z := \overline{O_f(x)}$ which is f -invariant. Since Z has finitely many irreducible components, we may assume $f^s(Z_1) = Z_1$ for some $s > 0$ and irreducible component Z_1 of Z . Note that $y = f^t(x) \in Z_1$ for some $t > 0$ (minimal). Indeed, otherwise, we will have $Z = \overline{Z \setminus Z_1}$, a contradiction. So $\overline{O_{f^s}(f^t(x))} \subseteq Z_1$. Note that

$$O_f(x) = \{x\} \cup \{f(x)\} \cup \cdots \cup \{f^{t-1}(x)\} \cup \bigcup_{i=0}^{s-1} f^i(O_{f^s}(f^t(x))).$$

Since f is a closed map,

$$Z = \{x\} \cup \{f(x)\} \cup \cdots \cup \{f^{t-1}(x)\} \cup \bigcup_{i=0}^{s-1} f^i(\overline{O_{f^s}(f^t(x))}).$$

Thus $\dim(\overline{O_{f^s}(f^t(x))}) = \dim(Z_1)$, so $\overline{O_{f^s}(f^t(x))} = Z_1$ which is irreducible. \square

Lemma 2.8. *Let X, Y be projective varieties over \overline{K} and f, g surjective endomorphisms on X, Y , respectively. Let $\pi : X \rightarrow Y$ be a generically finite surjective morphism such that $\pi \circ f = g \circ \pi$. Then $\delta_f = \delta_g$ and $\alpha_f(x) = \alpha_g(\pi(x))$ for any $x \in X \setminus \text{Exc}(\pi)$.*

Proof. By Lemma 2.2, $\delta_f = \delta_g$. Taking normalization, we may assume X and Y are normal by Lemma 2.6(3). Taking the Stein factorization of π (cf. [CMZ17, Lemma 5.2]), we may assume that π is a birational morphism. Then $\text{Exc}(\pi)$ is f^{-1} -invariant by [CMZ17, Lemma 7.3].

Let $x \in X \setminus \text{Exc}(\pi)$. By Lemma 2.7, replacing x by an f -iteration and f by a positive power, we may assume that $W = \overline{O_f(x)}$ is an irreducible closed f -invariant subset. Note that W is not contained in $\text{Exc}(\pi)$. So $\pi|_W : W \rightarrow \pi(W)$ gives a birational morphism. Since $\alpha_f(x) = \alpha_{f|_W}(x)$ by Lemma 2.5, replacing $\pi : X \rightarrow Y$ by $\pi|_W : W \rightarrow \pi(W)$, we may assume that $O_f(x)$ is Zariski dense in X .

Take an ample divisor A on Y . Then π^*A is big, so $\pi^*A \sim H + E$ for some ample divisor H and an effective divisor E on X . Take height functions $h_H, h_A \geq 1$. Then we have $h_H \leq h_A \circ \pi + O(1)$ on $X \setminus \text{Supp}(E)$. So $h_H(f^n(x)) \leq h_A(g^n(\pi(x))) + O(1)$ if $f^n(x) \notin \text{Supp}(E)$. We can take a subsequence $\{f^{n_k}(x)\}_k \subseteq X \setminus \text{Supp}(E)$ since $O_f(x)$ is dense. Then

$$\alpha_f(x) = \lim_{k \rightarrow +\infty} h_H(f^{n_k}(x))^{1/n_k} \leq \lim_{k \rightarrow +\infty} h_A(g^{n_k}(\pi(x)))^{1/n_k} = \alpha_g(\pi(x)).$$

This together with Lemma 2.6 imply $\alpha_f(x) = \alpha_g(\pi(x))$. \square

We can extend Lemma 2.6 to rational fibrations.

Lemma 2.9. *Let X, Y be projective varieties and f, g surjective endomorphisms on X, Y , respectively. Let $\pi : X \dashrightarrow Y$ be a dominant rational map such that $\pi \circ f = g \circ \pi$.*

- (1) *Assume $\delta_f = \delta_g$. If Conjecture 1.4 holds for g , then so does for f .*
- (2) *Assume that $\dim(X) = \dim(Y)$. Then $\delta_f = \delta_g$. Moreover, Conjecture 1.4 holds for f if and only if it holds for g .*

Proof. Let Γ be the graph of π , with the projections $p_1 : \Gamma \rightarrow X, p_2 : \Gamma \rightarrow Y$. Let $h : \Gamma \rightarrow \Gamma$ be the restriction of $f \times g : X \times Y \rightarrow X \times Y$ to Γ . Note that $\delta_h = \delta_f$ by Lemma 2.2. Note that p_1 is birational and p_2 is generically finite surjective when $\dim(X) = \dim(Y)$. By Lemma 2.8, we have $p_1^{-1}(Z_f) \setminus \text{Exc}(p_1) = Z_h \setminus \text{Exc}(p_1)$ and $p_2^{-1}(Z_g) \setminus \text{Exc}(p_2) = Z_h \setminus \text{Exc}(p_2)$ when $\dim(X) = \dim(Y)$. Hence,

$$(*) \quad p_1^{-1}(Z_f(d)) \subseteq Z_h(d'), \quad p_2^{-1}(Z_g(d)) \subseteq Z_h(d') \quad (\text{when } \dim(X) = \dim(Y))$$

with d' depending on d .

- (1) If Conjecture 1.4 holds for g , it holds for h by Lemma 2.6, and also for f by (*).

(2) We have $\delta_f = \delta_g$ by Lemma 2.2. Assume f satisfies Conjecture 1.4. Then h satisfies Conjecture 1.4 by Lemma 2.6. So does g by (*) above. \square

The following lemmas are used for the study of abelian fibrations in Section 7.

Lemma 2.10 (cf. [KS16b, Proposition 26]). *Let A be an abelian variety over an algebraically closed field of characteristic zero, H an ample divisor and D a nef \mathbb{R} -divisor. Then there exists some $\alpha \in \text{End}(A)_{\mathbb{R}}$ such that $\Phi_D^H = i_H(\alpha) \circ \alpha$ and $i_H(\alpha) = \alpha$ (see 2.1).*

Lemma 2.11. (cf. [FAG05]) *Let Y be a Noetherian scheme. Let $\pi: X \rightarrow Y$ be a smooth projective group scheme with geometrically integral fibers. Let D be a Cartier divisor on X . Then the natural transformation*

$$X(S) \longrightarrow \text{Pic}_{(X/Y)_{(et)}}(S); \quad x \mapsto [T_x^* D_S - D_S]$$

(here S runs over all locally Noetherian Y -schemes) defines a morphism $X \rightarrow \mathbf{Pic}_{X/Y}$ which factors through $\mathbf{Pic}_{X/Y}^0$. Denote this morphism as $\varphi_D: X \rightarrow \mathbf{Pic}_{X/Y}^0$. This construction commutes with base changes of Y , i.e., for any morphism $Z \rightarrow Y$ from a Noetherian scheme Z , we have the following commutative diagram:

$$\begin{array}{ccc} X \times_Y Z & \xrightarrow{\varphi_{D \times_Y Z}} & \mathbf{Pic}_{X/Y}^0 \times_Y Z \\ \text{id} \downarrow & & \downarrow \simeq \\ X \times_Y Z & \xrightarrow{\varphi_{D_Z}} & \mathbf{Pic}_{X \times_Y Z/Z}^0. \end{array}$$

Remark 2.12. For Picard functors and Picard schemes, see [FAG05, Part 5]. For relative Pic^0 , see [FAG05, Part 5, Proposition 9.5.20].

Remark 2.13. In the setting of the above lemma, if H is a π -ample divisor on X , then φ_H is a finite surjective morphism. Thus we can define $\Phi_D^H \in \text{End}(X/Y)_{\mathbb{R}}$ for an \mathbb{R} -Cartier divisor D on X , and also Rosati involution with respect to H , as in 2.1.

3. POLARIZED ENDOMORPHISMS, CURVES AND MORI DREAM SPACES

Theorem 3.1 (cf. [CS93], [KS14]). *Let $f: X \rightarrow X$ be a polarized endomorphism. Then:*

- (1) $Z_f = \text{Prep}(f)$.
- (2) For any $d > 0$, $X(d) \cap \text{Prep}(f)$ is a finite set.

Hence Conjecture 1.4 holds for every polarized endomorphism of any projective variety and every surjective endomorphism of any projective curve.

Proof. By assumption, $f^*H \sim dH$ for some ample divisor H and $d > 1$. Then we have a canonical height function

$$\hat{h}_{H,f}(x) = \lim_{n \rightarrow +\infty} \frac{h_H(f^n(x))}{d^n}$$

which is a height function associated to H (cf. [CS93]). The Northcott property of $\hat{h}_{H,f}$ implies that $\hat{h}_{H,f}(x) = 0$ if and only if $x \in \text{Prep}(f)$ for $x \in X(\overline{K})$. So $Z_f = (\hat{h}_{H,f} = 0) = \text{Prep}(f)$. It is easy to see that $\hat{h}_{H,f}(x) > 0$ if and only if $\alpha_f(x) = d (= \delta_f)$. So $Z_f = \text{Prep}(f)$. In addition, $\text{Prep}(f) \cap X(d)$ is finite for any $d > 0$ because of the Northcott finiteness property of $\hat{h}_{H,f}$. \square

Theorem 3.2 (cf. [Mat19, Theorem 4.1]). *Let X be a normal projective variety such that $N^1(X_{\overline{K}})_{\mathbb{Q}} = \text{Pic}(X_{\overline{K}})_{\mathbb{Q}}$ and the nef cone of $X_{\overline{K}}$ is generated by finitely many semi-ample Cartier divisors. Then $Z_f(d)$ is not Zariski dense in $X_{\overline{K}}$ for every surjective endomorphism f on X and any $d > 0$. In particular, Conjecture 1.4 holds for all surjective endomorphisms on Mori dream spaces.*

Proof. We may assume that $\delta_f > 1$. By the assumption, replacing f by a positive power, $f^*D \sim \delta_f D$ for some base point free effective Cartier divisor $D \neq 0$. Then $\hat{h}_{D,f}|_{Z_f} = 0$ and hence $h_D|_{Z_f} = O(1)$. For any $d > 0$, we have $Z_f(d) \subseteq \{x \in X(d) \mid h_D(x) \leq M\}$ for some $M > 0$. By [Shi19, Corollary 2.3]), the set of right hand side and hence $Z_f(d)$ are not Zariski dense in $X_{\overline{K}}$. \square

4. SURFACES AND HYPERKÄHLER VARIETIES

Theorem 4.1 (cf. [Kaw08]). *Let $f : X \rightarrow X$ be an automorphism of positive entropy on a projective surface. Then we have:*

- (1) *The number of f -periodic irreducible curves on $X_{\overline{K}}$ is finite.*
- (2) *Let C_i ($1 \leq i \leq r$) be the f -periodic irreducible curves on $X_{\overline{K}}$. Then $Z_f = \bigcup_{i=1}^r C_i \cup \text{Per}(f)$.*
- (3) *For any $d > 0$, $X(d) \cap (\text{Per}(f) \setminus (\bigcup_{i=1}^r C_i))$ is a finite set.*

Hence Conjecture 1.4 holds for every automorphism of any projective surface.

Proof. Note that $\delta := \delta_f (> 1)$ is known to be a Salem number with $\delta_{f^{-1}} = \delta$.

First we consider the case when $X_{\overline{K}}$ is smooth. We can find nef \mathbb{R} -divisors $D^+, D^- \neq 0$ on $X_{\overline{K}}$ such that $f^*D^+ \sim_{\mathbb{R}} \delta D^+$ and $f_*D^- \sim_{\mathbb{R}} \delta D^-$ (cf. [Kaw08, Lemma 3.8] or [San17, Remark 5.11]). Then $D = D^+ + D^-$ is nef and big by the Hodge index theorem. Let $\{C_\lambda\}_{\lambda \in \Lambda}$ be the family of all f -periodic irreducible curves. Then Λ is finite and we can take $a_\lambda > 0$ for each $\lambda \in \Lambda$ such that $D - \sum_{\lambda} a_\lambda C_\lambda$ is ample (cf. [Kaw08, Proposition 1.3]).

Set $E = \sum_{\lambda} a_{\lambda} C_{\lambda}$ and $A = D - E$. Take canonical height functions

$$\hat{h}_{D^+,f}(x) = \lim_{n \rightarrow +\infty} \frac{h_{D^+}(f^n(x))}{\delta^n}, \quad \hat{h}_{D^-,f^{-1}}(x) = \lim_{n \rightarrow +\infty} \frac{h_{D^-}(f^{-n}(x))}{\delta^n}$$

associated to D^+, D^- , respectively. Then

$$(*) \quad h_A + h_E = \hat{h}_{D^+,f} + \hat{h}_{D^-,f^{-1}} + O(1).$$

Substitute $f^n(x)$ to $(*)$, we have

$$(**) \quad h_A(f^n(x)) + h_E(f^n(x)) = \delta^n \hat{h}_{D^+,f}(x) + \delta^{-n} \hat{h}_{D^-,f^{-1}}(x) + O(1).$$

If $x \in Z_f \setminus \text{Supp}(E)$, then $\hat{h}_{D^+,f}(x)$ must be zero and so $\{h_A(f^n(x))\}_{n=0}^{+\infty}$ is upper bounded. The Northcott property for h_A implies $x \in \text{Per}(f)$ since f is an automorphism. If $x \in \text{Supp}(E)$, then $\alpha_f(x) = 1$ since f gives an automorphism on $\text{Supp}(E)$ which is a union of curves and any automorphism on a curve has first dynamical degree one. Therefore,

$$Z_f = \text{Supp}(E) \cup \text{Per}(f) = \bigcup_{\lambda \in \Lambda} C_{\lambda} \cup \text{Per}(f).$$

Thus the set $X(d) \cap (\text{Per}(f) \setminus \bigcup_{\lambda \in \Lambda} C_{\lambda})$ is finite for any $d > 0$ by the Northcott property of h_A . This proves the theorem for the case of smooth surfaces.

Now we consider the general case. Take an f -equivariant resolution $\pi : \tilde{X}_{\overline{K}} \rightarrow X_{\overline{K}}$ with an automorphism $\tilde{f} : \tilde{X}_{\overline{K}} \rightarrow \tilde{X}_{\overline{K}}$ such that $\pi \circ \tilde{f} = f \circ \pi$. Extending K , we may assume that \tilde{X} , π and \tilde{f} are defined over K . Lemma 2.6(2) and Lemma 2.8 imply that $\pi(Z_{\tilde{f}}) \subseteq Z_f \subseteq \pi(Z_{\tilde{f}} \cup \text{Exc}(\pi))$. Now $\text{Exc}(\pi)$ is f -invariant, so it is a union of f -periodic curves. Thus $Z_{\tilde{f}} \cup \text{Exc}(\pi) = Z_{\tilde{f}}$ by Step 1. Hence $\pi(Z_{\tilde{f}}) = Z_f$.

Let $\tilde{C}_1, \dots, \tilde{C}_r$ be the \tilde{f} -periodic irreducible curves on $\tilde{X}_{\overline{K}}$. Note that $\pi(\text{Exc}(\pi)) \subseteq \text{Per}(f)$. Then $\pi(\tilde{C}_1), \dots, \pi(\tilde{C}_r)$ exhaust the f -periodic irreducible curves on $X_{\overline{K}}$ and

$$Z_f = \pi(Z_{\tilde{f}}) = \bigcup_{i=1}^r \pi(\tilde{C}_i) \cup \text{Per}(f).$$

Fix $d > 0$. By the case for smooth surfaces, $\text{Per}(\tilde{f}) \setminus (\bigcup_{i=1}^r \tilde{C}_i)$ is finite in $\tilde{X}(d)$. So $\text{Per}(f) \setminus (\bigcup_{i=1}^r \pi(\tilde{C}_i))$ (the image of $\text{Per}(\tilde{f}) \setminus (\bigcup_{i=1}^r \tilde{C}_i)$) is also finite in $X(d)$. \square

A projective variety X over K is *Hyperkähler* if so is $X_{\overline{K}}$.

Theorem 4.2 (cf. [LS18, Theorem 1.2]). *Let $f : X \rightarrow X$ be a surjective endomorphism on a projective Hyperkähler variety X . Then $Z_f(d)$ is not Zariski dense in $X_{\overline{K}}$ for any $d > 0$.*

Proof. We may assume $\delta_f > 1$. By the ramification divisor formula, f is étale and hence an automorphism since a Hyperkähler variety has trivial $\pi_1^{\text{alg}}(X_{\overline{K}})$. The proof is almost the same as that of Theorem 4.1. Indeed, take nef \mathbb{R} -divisors $D^+, D^- \neq 0$ on X such

that $f^*D^+ \sim_{\mathbb{R}} \delta_f D^+$ and $f_*D^- \sim_{\mathbb{R}} \delta_{f^{-1}} D^-$. Then the Beauville–Bogomolov–Fujiki form enables us to show that $D^+ + D^-$ is nef and big (cf. [LS18, Lemma 3.8]). Write $D^+ + D^- \sim_{\mathbb{R}} A + E$ where A is ample and E is effective. Then we have $Z_f \subseteq \text{Supp } E \cup \text{Per}(f)$. The set $X(d) \cap (\text{Per}(f) \setminus \text{Supp } E)$ is finite for any $d > 0$ by the Northcott property of h_A . So $Z_f(d)$ is not Zariski dense in $X_{\overline{K}}$. \square

Theorem 4.3 (cf. [MSS18], [MZ19b]). *Let $f : X \rightarrow X$ be a non-invertible surjective endomorphism on a projective surface. Then $Z_f(d)$ is not Zariski dense in $X_{\overline{K}}$ for any $d > 0$.*

Proof. We may assume $X_{\overline{K}}$ is normal by Lemma 2.6. Since $\deg f > 1$, $X_{\overline{K}}$ is log canonical by [War90, Theorem 2.8].

If $K_{X_{\overline{K}}}$ is pseudo-effective, then by [Nak08, Theorem 7.1.1], after replacing f by a suitable power and extending K , there is a quasi-étale finite morphism $\phi : X' \rightarrow X$ with a lift $f' : X' \rightarrow X'$ of f (all are defined over K) such that

- (1) X' is an abelian surface; or
- (2) X' is a product of an elliptic curve and a curve of genus ≥ 2 .

The first case is a special case of Corollary 5.9, and the second case is a special case of Theorem 7.8, whose proofs are independent of the results of this theorem.

Assume that $K_{X_{\overline{K}}}$ is not pseudo-effective. Then, after replacing f by a suitable power and extending K , it is proved in [MZ19b, Theorem 5.4] that

- (3) f is a polarized endomorphism; or
- (4) There is a Fano contraction $\pi : X \rightarrow Y$ to a curve and f induces $g : Y \rightarrow Y$ with $\delta_f = \delta_g$; or
- (5) There is a finite surjective morphism $\tau : X \rightarrow \mathbb{P}^1 \times Y$ to the product of \mathbb{P}^1 and a curve Y with endomorphisms $g : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ and $h : Y \rightarrow Y$ such that $\tau \circ f = (g \times h) \circ \tau$.

These cases are verified by Theorem 3.1, Lemma 2.6, Lemma 2.4, respectively. \square

Combining Theorems 4.1 and 4.3, we obtain the following.

Theorem 4.4. *Let X be a projective surface and $f : X \rightarrow X$ a surjective endomorphism. Then $Z_f(d)$ is not Zariski dense in $X_{\overline{K}}$ for any $d > 0$.*

5. ABELIAN VARIETIES

In this section, we first prove:

Theorem 5.1 (cf. [MS18, Section 5]). *Let X be an abelian variety and $f : X \rightarrow X$ a surjective endomorphism (which is not necessarily an isogeny) with the first dynamical*

degree $\delta_f > 1$. Then there is a proper abelian subvariety $B \subset X_{\overline{K}}$ and a point $p \in X(\overline{K})$ such that $B + p$ is f -invariant and $Z_f = B + p + \text{Tor}(X_{\overline{K}})$.

To prove Theorem 5.1, we use the following results.

Lemma 5.2 (cf. [Sil17, Proof of Theorem 2]). *Let $f : A \rightarrow A$ be an isogeny on an abelian variety over \overline{K} . Then there are abelian subvarieties A_1, A_2 of A such that we have:*

- (1) *The addition map $m : A_1 \times A_2 \rightarrow A$ is an isogeny.*
- (2) *A_i is f -invariant for $i = 1, 2$. Set $f_i = f|_{A_i}$.*
- (3) *The map $1_{A_1} - f_1 : A_1 \rightarrow A_1$ is surjective.*
- (4) $\delta_{f_2} = 1$.

Remark 5.3. Let $F : A \rightarrow A$ be a surjective endomorphism. Then we can write $F = \tau_a \circ f$ where f is an isogeny and τ_a is a translation by $a \in A$. Apply Lemma 5.2 to f . Take $(a_1, a_2) \in A_1 \times A_2$ such that $a_1 + a_2 = a$. Then we have $F \circ m = m \circ ((\tau_{a_1} \circ f_1) \times (\tau_{a_2} \circ f_2))$, $\tau_{a_1} \circ f_1$ is conjugate to an isogeny by a translation, and $\delta_{\tau_{a_2} \circ f_2} = 1$ (cf. [MS18, §5]).

Theorem 5.4 ([KS16b, Theorem 1]). *Let $f : A \rightarrow A$ be an isogeny on an abelian variety with $\delta_f > 1$. Take a symmetric nef \mathbb{R} -divisor D such that D is not numerically trivial and $f^*D \equiv \delta_f D$. Take the canonical height function $\hat{h}_{D,f}$ associated to D :*

$$\hat{h}_{D,f}(x) = \lim_{n \rightarrow \infty} h_D(f^n(x)) / \delta_f^n.$$

Then there is an f -invariant proper abelian subvariety $B \subset A_{\overline{K}}$ such that the zero locus of $\hat{h}_{D,f}$ is equal to $B + \text{Tor}(A_{\overline{K}})$.

Proof of Theorem 5.1. First we assume that f is an isogeny, and show:

Claim 5.5. Let $f : X \rightarrow X$ be an isogeny on an abelian variety with $\delta_f > 1$. Then there is an f -invariant proper abelian subvariety $B \subset X_{\overline{K}}$ such that $Z_f = B + \text{Tor}(X_{\overline{K}})$.

Proof. We prove the claim by induction on the dimension. If $\dim(X) = 1$, then $Z_f = \text{Prep}(f)$. So we have to show $\text{Prep}(f) = \text{Tor}(X_{\overline{K}})$. Since f is an isogeny, $f(X_{\overline{K}}[N]) \subseteq X_{\overline{K}}[N]$ for any $N > 0$, where $X_{\overline{K}}[N]$ denotes the (finite) set of N -torsion points of $X_{\overline{K}}$. So $\text{Tor}(X_{\overline{K}}) \subseteq \text{Prep}(f)$. Conversely, take any $x \in \text{Prep}(f)$. Then $f^{n+k}(x) = f^n(x)$ for some $n, k \in \mathbb{Z}_{>0}$. This means that x is contained in $(1_X - f^k)^{-1}(\text{Ker}(f^n))$, which is a finite group since f^n and $1_X - f^k$ are isogenies. Here we used the assumption $\delta_f > 1$ to guarantee $f^k \neq 1_X$. Therefore, $x \in \text{Tor}(X_{\overline{K}})$. Hence $Z_f = \text{Tor}(X_{\overline{K}})$.

Assume $\dim(X) \geq 2$ and the claim holds for abelian varieties of smaller dimensions. Take a nef symmetric \mathbb{R} -Cartier divisor D on X such that D is not numerically trivial and $f^*D \equiv \delta_f D$. Then Theorem 5.4 implies that the zero locus of $\hat{h}_{D,f}$ is equal to $B + \text{Tor}(X_{\overline{K}})$ for some proper f -invariant abelian subvariety $B \subset X_{\overline{K}}$. For any point

$x \in X_{\overline{K}}$, $\hat{h}_{D,f}(x) > 0$ implies that $\alpha_f(x) = \delta_f$. So $Z_f \subseteq B + \text{Tor}(X_{\overline{K}})$. Note that $\hat{h}_{D,f}$ is non-negative (cf. [KS16b, Section 6]).

Set $f_B = f|_B : B \rightarrow B$. Then $\delta_{f_B} \leq \delta_f$ by [NZ09, Proposition A.11]. If $\delta_{f_B} < \delta_f$, then the arithmetic degree of any point of B is smaller than δ_f . Then $Z_f = B + \text{Tor}(X_{\overline{K}})$. Next, assume that $\delta_{f_B} = \delta_f$. Then the induction hypothesis implies that there is a proper f_B -invariant abelian subvariety $C \subset B$ such that $Z_{f_B} = C + \text{Tor}(B)$. Now we have $C + \text{Tor}(X_{\overline{K}}) \subseteq Z_f$. Conversely, take any $x \in Z_f$. Then mx is in B (and also in $m_A(Z_f) = Z_f$; see Lemma 2.6) for some $m \in \mathbb{Z}_{>0}$. Now $mx \in Z_f \cap B = Z_{f_B}$, so $kmx \in C$ for some $k \in \mathbb{Z}_{>0}$. Thus $Z_f = C + \text{Tor}(X_{\overline{K}})$. This proves the claim. \square

Next we consider a general endomorphism f . Write $f = \tau_a \circ \phi$ such that ϕ is an isogeny and τ_a is the translation map by $a \in X$. By Lemma 5.2, there are abelian subvarieties A_1, A_2 of X such that we have:

- (1) The addition map $m : A_1 \times A_2 \rightarrow A$ is an isogeny.
- (2) A_i is ϕ -invariant for $i = 1, 2$. Set $\phi_i = \phi|_{A_i}$.
- (3) The map $1_{A_1} - \phi_1 : A_1 \rightarrow A_1$ is surjective.
- (4) $\delta_{\phi_2} = 1$.

Take $a_i \in A_i$ for $i = 1, 2$ such that $a = a_1 + a_2$ and set $f_i = \tau_{a_i} \circ \phi_i$. Take $p \in A_1$ satisfying $p - \phi(p) = a_1$. Since translations act trivially on the Neron-Severi lattice and the map $\phi_1 \times \phi_2$ descends to ϕ via the map m , Lemma 2.2 and the product formula imply $\delta_f = \delta_\phi = \delta_{\phi_1 \times \phi_2} = \delta_{\phi_1} = \delta_{f_1}$. Now we have the commutative diagram:

$$\begin{array}{ccc}
 A_1 \times A_2 & \xrightarrow{\phi_1 \times \phi_2} & A_1 \times A_2 \\
 \tau_p \times 1_{A_2} \downarrow & & \downarrow \tau_p \times 1_{A_2} \\
 A_1 \times A_2 & \xrightarrow{f_1 \times f_2} & A_1 \times A_2 \\
 m \downarrow & & \downarrow m \\
 X & \xrightarrow{f} & X
 \end{array}$$

Applying Claim 5.5 to ϕ_1 , there is a proper ϕ_1 -invariant abelian subvariety $B_1 \subset (A_1)_{\overline{K}}$ such that $Z_{\phi_1} = B_1 + \text{Tor}((A_1)_{\overline{K}})$. Since $\phi_1 \times \phi_2$ descends to f via a finite morphism, the first equality below follows from Lemma 2.6, while the second follows from Lemma 2.4:

$$\begin{aligned}
 Z_f &= (m \circ (\tau_p \times 1_{A_2}))(Z_{\phi_1 \times \phi_2}) \\
 &= (m \circ (\tau_p \times 1_{A_2}))((B_1 + \text{Tor}((A_1)_{\overline{K}})) \times (A_2)_{\overline{K}}) \\
 &= B_1 + \text{Tor}((A_1)_{\overline{K}}) + p + (A_2)_{\overline{K}}.
 \end{aligned}$$

Set $B = B_1 + (A_2)_{\overline{K}}$. Then we have $Z_f = B + p + \text{Tor}(X_{\overline{K}})$. We compute

$$\begin{aligned}
f(B + p) &= \phi(B_1 + (A_2)_{\overline{K}} + p) + a \\
&= \phi(B_1) + \phi((A_2)_{\overline{K}}) + \phi(p) + a_1 + a_2 \\
&= B_1 + (A_2)_{\overline{K}} + \phi(p) + a_1 + a_2 \\
&= (B + a_2) + (\phi(p) + a_1) \\
&= B + p.
\end{aligned}$$

So $B + p$ is f -invariant. □

Theorem 5.6. *Notation is as in Theorem 5.1. Write $Z_f = B + p + \text{Tor}(X_{\overline{K}})$. Then:*

- (1) $\{B + p + t\}_{t \in \text{Tor}(X_{\overline{K}})}$ is the family of all maximal f -preperiodic subvarieties of small dynamical degree in $X_{\overline{K}}$.
- (2) There are only finitely many maximal f -invariant subvarieties of small dynamical degree in $X_{\overline{K}}$ (cf. Question 8.4(3)).
- (3) For any $d > 0$, $Z_f(d) \subseteq \bigcup_{i=1}^r (B + p + t_i)$ for some $t_1, \dots, t_r \in \text{Tor}(X_{\overline{K}})$. In particular, $Z_f(d)$ is not Zariski dense.

Proof. (1) Write $f = \tau_a \circ \phi$ such that ϕ is an isogeny and τ_a is the translation map by $a \in X$. Pick $t \in \text{Tor}(X_{\overline{K}})$. Since $f(B + p) = B + p$, inductively, for any $s \geq 1$, we get

$$\begin{aligned}
f^s(B + p + t) &= f^{s-1}(\phi(B + p + t) + a) \\
&= f^{s-1}(\phi(B + p) + \phi(t) + a) \\
&= f^{s-1}(f(B + p) + \phi(t)) \\
&= f^{s-1}(B + p + \phi(t)) \\
&= \dots = B + p + \phi^s(t).
\end{aligned}$$

Since ϕ is an isogeny, if t is in the set of N -torsion points (which is a finite set), then so is $\phi^s(t)$. Thus $B + p + t$ is an f -preperiodic subvariety of small dynamical degree. The maximality of $B + p + t$ is clear, since $Z_f = \bigcup_{t \in \text{Tor}(X_{\overline{K}})} (B + p + t)$.

(2) Extending K , we may assume that B and p are defined over K . Fix $d > 0$. Take the quotient $\pi : X \rightarrow Y := X/B$. Then π is f -equivariant. Set $g := f|_Y$. If $B + p + t$ is f -invariant, then $g(\pi(p+t)) = g(\pi(B+p+t)) = \pi(f(B+p+t)) = \pi(B+p+t) = \pi(p+t)$. By (1), it suffices to show that there are only finitely many g -fixed points in $Y_{\overline{K}}$. Suppose the contrary. Then $g|_Z = \text{id}$ for some positive dimensional irreducible closed subvariety $Z \subseteq Y_{\overline{K}}$. By the product formula, $\delta_{f|_{\pi^{-1}(Z)}} = \delta_{f|_{B+p}} < \delta_f$. However, $\pi^{-1}(Z) \supseteq B + p + t$ for some $t \in \text{Tor}(X_{\overline{K}})$, a contradiction with the maximality.

(3) Fix $d > 0$. Then $\pi(Z_f(d) - p) \subseteq \text{Tor}(Y_{\overline{K}}) \cap Y(d)$ and $\text{Tor}(Y_{\overline{K}}) \cap Y(d)$ is a finite group. So

$$\pi([N]_X(Z_f(d) - p)) = [N]_Y(\pi(Z_f(d) - p)) = 0$$

for some $N > 0$. Hence $Z_f(d) - p \subseteq [N]_X^{-1}(B_{\overline{K}}) = B_{\overline{K}} + X_{\overline{K}}[N]$, where $X_{\overline{K}}[N]$ denotes the set of N -torsion points of $X_{\overline{K}}$. As a consequence, $Z_f(d)$ is contained in a finite union of subvarieties of the form $B_{\overline{K}} + p + t$ with t being a torion point. \square

We can use the above precise description of Z_f to prove Conjecture 1.4 for varieties of maximal Albanese dimension.

Definition 5.7. Let X be a normal projective variety over an algebraically closed field. Then there is an abelian variety $\text{Alb}(X)$, called the *Albanese variety* of X , and a dominant rational (Albanese) map $\text{alb}_X : X \dashrightarrow \text{Alb}(X)$, such that every rational map from X to another abelian variety factors through alb_X .

We say that a variety X is of *maximal Albanese dimension* if $\dim(\text{alb}_X(X)) = \dim(X)$. We say that a variety X defined over K is of *maximal Albanese dimension* if so is $X_{\overline{K}}$.

Corollary 5.8. *Conjecture 1.4 holds for surjective endomorphisms on normal projective varieties of maximal Albanese dimension.*

Proof. Let X be a normal projective variety of maximal Albanese dimension and $f : X \rightarrow X$ a surjective endomorphism with $\delta_f > 1$. By Lemma 2.9 and replacing X and alb_X by the graph, we may assume that the albanese map $\text{alb}_X : X \rightarrow \text{Alb}(X) =: A$ is a well-defined morphism, and also defined over K after extending K . Then f induces a surjective endomorphism $g : A \rightarrow A$. By Lemma 2.6 or 2.9 and replacing (X, f) by $(\text{alb}_X(X), g|_{\text{alb}_X(X)})$, we may assume that X is a g -invariant subvariety of A such that $0 \in X$ and X is not contained in any proper abelian subvariety of A .

Note that $\delta_f \leq \delta_g$ (cf. [NZ09, Proposition A.11]). We prove the assertion by induction on $\dim(A)$. If $\dim(A) = 1$, then X is either a point or A , so the assertion follows from Theorem 5.6.

Consider the general case. By Theorem 5.1, $Z_g = B + p + \text{Tor}(A_{\overline{K}})$, where $B + p$ is a proper g -invariant translated abelian subvariety of $A_{\overline{K}}$. If $\delta_f < \delta_g$, then $X_{\overline{K}} \subseteq Z_g$. This implies that $X_{\overline{K}} \subseteq B + p + t$ for some $t \in \text{Tor}(A_{\overline{K}})$. Then $B + p + t = B$ since $0 \in X_{\overline{K}} \subseteq B + p + t$, but this contradicts that X is not contained in any proper abelian subvariety of A . So we have $\delta_f = \delta_g$.

Take any $d > 0$. By Theorem 5.6, $Z_g(d) \subseteq \bigcup_{i=1}^r (B + p + t_i)$ for some torsion points $t_1, \dots, t_r \in A_{\overline{K}}$. Then

$$Z_f(d) = X_{\overline{K}} \cap Z_g(d) \subseteq X_{\overline{K}} \cap \left(\bigcup_{i=1}^r (B + p + t_i) \right) = \bigcup_{i=1}^r (X_{\overline{K}} \cap (B + p + t_i)).$$

Moreover, $X_{\overline{K}} \not\subseteq B + p + t_i$ for any i . So $\bigcup_{i=1}^r (X_{\overline{K}} \cap (B + p + t_i))$ is a proper closed subset of $X_{\overline{K}}$. \square

A variety X over K is \mathbb{Q} -abelian if so is $X_{\overline{K}}$. We can also prove Conjecture 1.4 for \mathbb{Q} -abelian varieties.

Corollary 5.9 (cf. [MZ19b, Theorem 2.8]). *Conjecture 1.4 holds for surjective endomorphisms on \mathbb{Q} -abelian varieties.*

Proof. Let $f : X \rightarrow X$ be a surjective endomorphism on a \mathbb{Q} -abelian variety. After enlarging the ground field if necessary, we can take a finite surjective morphism $\pi : A \rightarrow X$ from an abelian variety and a surjective endomorphism $g : A \rightarrow A$ such that $\pi \circ g = f \circ \pi$ (cf. [NZ10] or [CMZ17, Corollary 8.2]). Then the assertion follows from Lemma 2.6.

test123 test123 \square

6. VARIETIES ADMITTING INT-AMPLIFIED ENDOMORPHISMS

In this section, the ground field is \overline{K} (or any algebraically closed field of characteristic zero) unless otherwise stated. We focus on a \mathbb{Q} -factorial klt projective variety X admitting an int-amplified endomorphism \mathcal{I} , i.e., $\mathcal{I}^*L - L = H$ for some ample Cartier divisors L and H . Alternatively, it is equivalent to saying that all the eigenvalues of $\mathcal{I}^*|_{\mathbb{N}^1(X)}$ are of modulus greater than 1; see [Men17, Theorem 1.1]. Given any (not necessarily int-amplified) surjective endomorphism f of X , we may run the f -equivariant minimal model program (after some iteration of f) by [MZ19a, Theorem 1.2]. In this way, it suffices for us to focus on the following Case TIR when studying the Kawaguchi–Silverman Conjecture 1.1; see [MZ19b, Theorem 1.7].

Case TIR_n (Totally Invariant Ramification case). Let X be a normal projective variety of dimension $n \geq 1$, which has only \mathbb{Q} -factorial Kawamata log terminal (klt) singularities and admits an int-amplified endomorphism. Let $f : X \rightarrow X$ be a surjective endomorphism. Moreover, we impose the following conditions.

- (A1) The anti-Iitaka dimension $\kappa(X, -K_X) = 0$; $-K_X$ is nef, whose class is extremal in both the *nef cone* $\text{Nef}(X)$ and the *pseudo-effective divisors cone* $\text{PE}^1(X)$.
- (A2) $f^*D = \delta_f D$ for some prime divisor $D \sim_{\mathbb{Q}} -K_X$.
- (A3) The ramification divisor of f satisfies $\text{Supp } R_f = D$.
- (A4) There is an f -equivariant Fano contraction $\pi : X \rightarrow Y$ with $\delta_f > \delta_{f|_Y} (\geq 1)$.
- (A5) $\dim(X) \geq \dim(Y) + 2 \geq 3$.

For the sAND Conjecture 1.4, we show that the same strategy works.

Theorem 6.1. (cf. [MZ19b, Theorem 1.7]) *Let X be a normal projective variety over K such that $X_{\overline{K}}$ has only \mathbb{Q} -factorial Kawamata log terminal (klt) singularities and admitting an int-amplified endomorphism. Then we have:*

- (1) *If $K_{X_{\overline{K}}}$ is pseudo-effective, then Conjecture 1.4 holds for any surjective endomorphism of X .*
- (2) *Suppose that Conjecture 1.4 holds for Case TIR (for those $f|_{X_i} : X_i \rightarrow X_i$ appearing in any equivariant MMP starting from $X_{\overline{K}}$). Then Conjecture 1.4 holds for any surjective endomorphism on X .*

Proof. We apply almost the same proof of [MZ19b, Theorem 1.7]. Indeed, we only need to replace [MZ19b, Lemma 2.5 and Theorem 2.8] for Conjecture 1.1 by our Lemma 2.6 and Corollary 5.9 for Conjecture 1.4. \square

In the rest of this section, we show that Case TIR will not occur in many cases.

We first extend [MY19, Theorem 4.4] a bit to make all surjective endomorphisms liftable (after suitable iteration).

Theorem 6.2. (cf. [MY19, Theorem 4.4]) *In Case TIR, there exists the following commutative diagram:*

$$\begin{array}{ccc} X & \xleftarrow{\mu_X} & \tilde{X} \\ \pi \downarrow & & \downarrow \tilde{\pi} \\ Y & \xleftarrow{\mu_Y} & A \end{array}$$

where A is an abelian variety, μ_Y is finite surjective, \tilde{X} is the normalization of the main component of $X \times_Y A$, and μ_X is quasi-étale finite surjective, such that any commutative diagram of surjective morphisms

$$h_X \circlearrowleft X \xrightarrow{\pi} Y \circlearrowright h_Y$$

can be lifted equivariantly to

$$\begin{array}{ccc} h_X \circlearrowleft X & \xleftarrow{\mu_X} & \tilde{X} \circlearrowright h_{\tilde{X}} \\ \pi \downarrow & & \downarrow \tilde{\pi} \\ h_Y \circlearrowleft Y & \xleftarrow{\mu_Y} & A \circlearrowright h_A \end{array}$$

Proof. Note that in Case TIR, $\kappa(X, -K_X) = 0$ and $-K_X \sim_{\mathbb{Q}} D$ for some reduced effective divisor D . By [MZ19b, Theorem 6.2], $\text{Supp } R_{h_X} \subseteq D$ for any surjective endomorphism h_X of X . Next, we only consider those h_X which can descend equivariantly to h_Y via π . Since D is π -ample, $\pi(D) = Y$. Note also that D is irreducible in our assumption.

Using the same notation in the proof of [MY19, Theorem 4.4], we can define a positive integer m_E for any prime divisor E on Y so that $\pi^*E = m_E F$ for some prime divisor F on X . We set $\Delta = \sum \frac{m_E-1}{m_E} E$ which is a finite sum. By the same computation in the proof of [MY19, Theorem 4.4] and the key reason $\pi(D) = Y$, we have $h_Y^*(K_Y + \Delta) \sim K_Y + \Delta$. Since the int-amplified endomorphism on X descends to an int-amplified endomorphism on Y , $K_Y + \Delta \sim_{\mathbb{Q}} 0$ by [Men17, Theorem 1.1] and [MZ19b, Proposition 3.4].

Applying [MY19, Lemma 4.9] to the pair (Y, Δ) , we obtain the commutative diagram:

$$\begin{array}{ccc} Y_1 & \xrightarrow{h_{Y_1}} & Y_1 \\ \mu_{Y_1} \downarrow & & \downarrow \mu_{Y_1} \\ Y & \xrightarrow{h_Y} & Y \end{array}$$

where $K_{Y_1} \sim 0$ and μ_{Y_1} is finite surjective which does not depend on h_Y . In the proof of [MY19, Theorem 4.4], it is further shown that Y_1 is klt. Note that Y_1 admits an int-amplified endomorphism. So Y_1 is \mathbb{Q} -abelian by [Men17, Theorem 1.9]. Further, we can find abelian variety A and the commutative diagram:

$$\begin{array}{ccc} A & \xrightarrow{h_A} & A \\ \mu_{Y_2} \downarrow & & \downarrow \mu_{Y_2} \\ Y_1 & \xrightarrow{h_{Y_1}} & Y_1 \end{array}$$

where μ_{Y_2} is quasi-étale, finite and surjective (cf. [CMZ17, Corollary 8.2]).

Now h_A is the lifting of h_Y via $\mu_Y := \mu_{Y_1} \circ \mu_{Y_2}$. The induced cover μ_X is quasi-étale by [MY19, Lemma 4.5]. \square

Let X be a normal projective variety. Denote by $\pi_1^{\text{alg}}(X_{\text{reg}})$ the *algebraic fundamental group* of the smooth locus X_{reg} of X . Conjecture 1.1 for smooth rationally connected projective varieties admitting int-amplified endomorphisms has been solved in [MZ19b] and [MY19]. Indeed, the key point of the proof is the following, noting that a smooth rationally connected projective variety has trivial algebraic fundamental group.

Theorem 6.3. (cf. [MZ19b], [MY19]) *Let X be a \mathbb{Q} -factorial klt projective variety with the group $\pi_1^{\text{alg}}(X_{\text{reg}})$ being finite. Suppose X admits an int-amplified endomorphism. Then Case TIR will not occur during any MMP starting from X .*

Proof. We argue along the lines of [MY19, Section 4.3] (while [MZ19b] is effective in low dimension or relative dimension). Suppose the contrary that there exists some f -equivariant (after iteration) MMP

$$X_1 \dashrightarrow \cdots \dashrightarrow X_r \rightarrow Y,$$

such that $(X_r, Y, f|_{X_r}, f|_Y)$ satisfies Case TIR. Let $\mathcal{I} : X \rightarrow X$ be an int-amplified endomorphism such that the MMP is also \mathcal{I} -equivariant. Note that $(X_r, Y, \mathcal{I}|_{X_r}, \mathcal{I}|_Y)$ may not satisfy Case TIR. Nevertheless, by Theorem 6.2, $\mathcal{I}|_{X_r}$ satisfies $(*)$ in [MY19, Definition 4.11]. By [MY19, Lemmas 4.12 and 4.13], \mathcal{I} also satisfies $(*)$ in [MY19, Definition 4.11]. In particular, without considering any dynamic property, we have the commutative diagram:

$$\begin{array}{ccc} X & \xleftarrow{\mu_X} & \tilde{X} \\ \pi \downarrow & & \downarrow \tilde{\pi} \\ Y & \xleftarrow{\mu_Y} & A \end{array}$$

where A is an abelian variety, μ_Y is finite surjective, and μ_X is quasi-étale finite surjective. Note that $\dim(A) = \dim(Y) > 0$. So \tilde{X} admits an étale cover of arbitrary large degree. In particular, $\pi_1^{\text{alg}}(X_{\text{reg}})$ is infinite, a contradiction. \square

In [MZ19b, Theorem 8.6], the relative dimensional one case has been ruled out in Case TIR. Based on this, we will deal with the relative dimensional two case.

We will show that Case TIR is equivalent to the following simpler Case TIR'.

Case TIR'_n (Totally Invariant Ramification case). Let X be a normal projective variety of dimension $n \geq 1$, which has only \mathbb{Q} -factorial Kawamata log terminal (klt) singularities and admits an int-amplified endomorphism. Let $f : X \rightarrow X$ be a surjective endomorphism. Moreover, we impose the following conditions.

- (A1') The anti-Iitaka dimension $\kappa(X, -K_X) = 0$.
- (A2') $f^*D = \delta_f D$ for some reduced effective Weil divisor $D \sim_{\mathbb{Q}} -K_X$.
- (A4) There is an f -equivariant Fano contraction $\pi : X \rightarrow Y$ with $\delta_f > \delta_{f|_Y} (\geq 1)$.

Lemma 6.4. *Case TIR_n is equivalent to Case TIR'_n.*

Proof. Assume that Case TIR'_n holds. We first show A2 that D is irreducible. Let P and Q be two different irreducible components of D . If P is not π -ample, then $P \equiv \pi^*P_Y$ for some $P_Y \in N^1(Y) \setminus 0$ by the cone theorem (cf. [KM98, Theorem 3.7]). Note that $g^*P_Y = \delta_f P_Y$. This is a contradiction with the assumption A4. So we may assume P and Q are π -ample. Then $P - tQ$ is π -trivial for some rational number $t > 0$ and the same argument shows that $P - tQ \equiv 0$. Note that $f^*(P - tQ) = \delta_f(P - tQ)$ and the Albanese morphism of X is surjective by [Men17, Theorem 1.8]. Then $P - tQ \sim_{\mathbb{Q}} 0$ by [MZ19b, Proposition 3.3]. In particular, $\kappa(X, D) \geq \kappa(X, P) > 0$, a contradiction with A1'.

Note that $N^1(X)/\pi^*N^1(Y)$ is 1-dimensional and $f^*|_{N^1(X)/\pi^*N^1(Y)} = q \text{ id}$ for some integer $q > 0$. Then $q = \delta_f$ is the only eigenvalue of $f^*|_{N^1(X)}$ with modulus δ_f and the q -eigenspace is 1-dimensional. On the other hand $f^*H \equiv \delta_f H$ for some nef divisor H by a version of

the Perron-Frobenius theorem (cf. [Birk]). So the rays $R_H = R_D = R_{-K_X}$ in $\text{Nef}(X)$ and $\text{PE}^1(X)$, and the second part of A1 holds, by [MZ19b, Lemma 9.1].

By [MZ19b, Theorem 6.2], $\text{Supp } R_f \subseteq D$. On the other hand, A2' and $\delta_f > 1$ imply A3 that $\text{Supp } R_f = D$. A5 follows from [MZ19b, Theorem 8.6]. \square

Proposition 6.5. *Let $\pi : X \rightarrow Y$ be the Fano contraction of a K_X -negative extremal ray R_C on a \mathbb{Q} -factorial klt projective variety X to an abelian variety Y with $\dim(X) = \dim(Y) + 1$. Let $f : X \rightarrow X$ and $g : Y \rightarrow Y$ be int-amplified endomorphisms such that $\pi \circ f = g \circ \pi$ and $\delta_f > \delta_g$. Then we have:*

- (1) R_C is the only K_X -negative extremal ray.
- (2) The anti-Iitaka dimension $\kappa(X, -K_X) > 0$, and $f^*K_X \equiv \delta_f K_X$.
- (3) There is an f -equivariant surjective morphism $\tau : X \rightarrow Z \cong \mathbb{P}^1$ such that $f|_Z$ is δ_f -polarized.
- (4) P is semi-ample for any f^{-1} -invariant prime divisor P .

Proof. (1) Let R_{C_2} be a K_X -negative extremal ray and $\pi_2 : X \rightarrow X_2$ the contraction of R_{C_2} . By [KM98, Theorem 3.7], we may take C_2 as a rational curve. Then $\pi(C_2)$ is a point in the abelian variety Y . Let F be a (connected) fiber of π_2 . Note that $\pi(C'_2)$ is a point for any curve $C'_2 \subseteq F$ since $C'_2 \in R_{C_2}$. Hence, $\pi(F)$ is a point. Then the rigidity lemma [Deb01, Lemma 1.15] implies that π factors through π_2 . Since π itself is a contraction of R_C , $R_{C_2} = R_C$ and $X_2 = Y$. This proves (1).

(2) By a version of the Perron-Frobenius theorem in [Birk], there exists a nef eigenvector D of $f^*|_{\text{N}^1(X)}$ with eigenvalue δ_f . Note that the eigenvalues of $f^*|_{\text{N}^1(X)}$ consist of one δ_f and eigenvalues of $g^*|_{\text{N}^1(Y)}$ (with modulus $< \delta_f$). So D is π -ample and extremal in $\text{PE}^1(X)$ by [MZ19b, Lemma 9.1].

Let $a > 0$ be the rational number such that $B := aK_X + D$ is π -trivial, i.e., $B \cdot C = 0$. We claim that B is pseudo-effective. Suppose the contrary. For a small effective ample \mathbb{Q} -Cartier divisor E , $(1/a)B + E$ is not pseudo-effective. Denote by $A := E + (1/a)D$ which is ample since D is nef. Thus $K_X + A = (1/a)B + E$ is not pseudo-effective. Note that (X, A) is a klt pair by a suitable choice of A . By the cone theorem (cf. [KM98, Theorem 3.7]), there exists some $(K_X + A)$ -negative extremal ray $R_{C'}$. Note that $(K_X + A) \cdot C = E \cdot C > 0$. So $R_{C'} \neq R_C$, contradicting (1). This proves the claim.

Since $-K_X$ is effective by [MZ19b, Theorem 1.5], the claim above and $D = B + (-aK_X)$ imply $R_D = R_{-K_X}$ in $\text{PE}^1(X)$ and hence $B \equiv 0$ and $D \equiv -aK_X$. We assert that $\kappa(X, -K_X) > 0$. Suppose the contrary. Then $-K_X \sim_{\mathbb{Q}} D'$ for some reduced Weil divisor D' such that $f^{-1}(D') = D'$ and $\text{Supp } R_f = D'$ by [MZ19b, Theorem 1.5]. Note that $f^*D' \equiv f^*D/a = (\delta_f/a)D \equiv \delta_f D'$ and $\delta_f D' - f^*D' \geq 0$. Then $f^*D' = \delta_f D'$. In

particular, Case TIR' (and hence Case TIR) is satisfied. However, this contradicts the condition A5 of Case TIR. So $\kappa(X, -K_X) > 0$ as asserted. This also proves (2).

(3) By [MZ19b, Theorem 7.8], there exists an f -equivariant dominant rational map $\tau : X \dashrightarrow Z$ to a normal projective variety Z (with connected general fibers) such that $\dim(Z) > 0$ and $f|_Z$ is δ_f -polarized. Note that the indeterminacy locus of τ is an f^{-1} -invariant closed subset of codimension ≥ 2 in X . By [CMZ17, Lemma 7.5] and [Men17, Lemma 4.1], τ is then a morphism since $\dim(A) = \dim(X) - 1$.

We claim that $Z \cong \mathbb{P}^1$. Let $h := f|_Z$. Write $h^*H_Z \sim \delta_f H_Z$ for some very ample divisor H_Z . Let $H_X := \tau^*H_Z$. Then $f^*H_X \sim \delta_f H_X$. So $H_X \equiv bD$ for some $b > 0$. By [MZ19b, Proposition 8.4], $H_X^2 \equiv_w b^2 D^2 \equiv_w 0$. This is possible only when $\dim(Z) = 1$. Since $H_X \equiv bD$ is π -ample, the general fiber F of π (which is a smooth rational curve) is not contracted by τ and hence dominates Z . So the claim and hence (3) are proved.

(4) Let P be an f^{-1} -invariant prime divisor. By [CMZ17, Lemma 7.5] and [Men17, Lemma 4.1], P dominates Y . Then $f^*P = \delta_f P$ and $R_P = R_D$ in $\text{PE}^1(X)$. Take $H_Z = z$ for some general point of Z and let X_z be the fiber of τ . Note that $X_z \cdot P \equiv_w 0$. So either $X_z \cap P = \emptyset$ or $X_z \cap P = P$. In particular, this implies that $\tau(P)$ is a point. We claim that $P = \tau^{-1}(\tau(P))$. If the claim is false, then $\tau^{-1}(\tau(P))$ is reducible; hence there exists some irreducible component $Q \neq P$ of $\tau^{-1}(\tau(P))$ such that $P \cap Q \neq \emptyset$, since τ has connected fibers. Note that $\tau(P)$ is h^{-1} -periodic by [CMZ17, Lemma 7.5]. So Q and hence $P \cap Q$ are f^{-1} -periodic. By [CMZ17, Lemma 7.5], $\pi(P \cap Q)$ is g^{-1} -periodic and hence equal to Y by [Men17, Lemma 4.1], contradicting that $\dim(P \cap Q) < \dim(Y)$. So the claim is true. In particular, P is semi-ample and (4) is proved. \square

Theorem 6.6. *Suppose that X has only terminal singularities in Case TIR. Then $\dim(X) \geq \dim(Y) + 3 \geq 4$.*

Proof. The proof consists of Claims 6.7 to 6.18. We prove by contrapositive, so suppose the contrary that $\dim(X) = \dim(Y) + 2$. Let $\mathcal{I} : X \rightarrow X$ be an int-amplified endomorphism such that $\pi : X \rightarrow Y$ is \mathcal{I} -equivariant.

Since X has terminal singularities, $\text{Sing}(X)$ has dimension $\leq \dim(X) - 3 < \dim(Y)$, so it does not dominate Y . By [Fak03, Theorem 5.1], there is a Zariski dense subset Y_0 of Y such that the fiber $X_y = \pi^{-1}(y)$ is smooth and \mathcal{I} -periodic for any $y \in Y_0$. Note that for each $y \in Y_0$, X_y is \mathcal{I}^s -invariant for some $s > 0$; and $\mathcal{I}^s|_{X \setminus D}$ is quasi-étale. Shrinking Y_0 a bit, we may assume that $\mathcal{I}^s|_{X_y}$ is étale outside $D \cap X_y$ by the purity of branch loci.

Let $y \in Y_0$ be a general point. Denote by

$$m := \text{the number of irreducible components (curves) of } D \cap X_y.$$

Since π is a Fano contraction, X_y is a smooth Fano surface and $\mathcal{I}^s|_{X_y}$ is polarized. By [MZ19, Corollary 1.4], $(X_y, D \cap X_y)$ is then a toric pair. Moreover, by the classification of smooth toric Fano surfaces, the boundary $D \cap X_y$ is a simple normal crossing loop of m smooth rational curves ℓ_i with $m \geq 3$. Note that ℓ_i and $\ell_i \cap \ell_j$ are $(\mathcal{I}^s|_{X_y})^{-1}$ -periodic. Since $\mathcal{I}^s|_{X_y \setminus D \cap X_y}$ is étale, any $(\mathcal{I}^s|_{X_y})^{-1}$ -periodic point x lies in $D \cap X_y$ by noting that $\mathcal{I}^s|_{X_y}$ is ramified at x . Since each $\ell_i \cong \mathbb{P}^1$ has (exactly) two $(\mathcal{I}^s|_{X_y})^{-1}$ -periodic points which are the intersecting points with some other two ℓ_j and ℓ_k , x further lies in $\text{Sing}(D \cap X_y)$ by the same argument on $\mathcal{I}^{sm}|_{\ell_i}$. In particular, $\text{Sing}(D \cap X_y)$ is exactly the set of $(\mathcal{I}^s|_{X_y})^{-1}$ -periodic points in X_y . Note that $\#\text{Sing}(D \cap X_y) = m \geq 3$.

Claim 6.7. Let Z be the union of irreducible components of dimension $\dim(X) - 2$ in $\text{Sing}(D)$. Then $h^{-1}(Z) = Z$ for any surjective endomorphism h of X .

Let $\nu : \overline{D} \rightarrow D \subseteq X$ be the normalization of D and \mathfrak{c} the conductor of D , regarded as a Weil divisor on \overline{D} . Since X is smooth in codimension 2, the adjunction formula gives

$$K_{\overline{D}} + \mathfrak{c} = \nu^*(K_X + D)$$

where $\nu^*(K_X + D)$ is regarded as the pullback of a divisorial sheaf. There is an endomorphism $\overline{h} : \overline{D} \rightarrow \overline{D}$ such that $\nu \circ \overline{h} = h \circ \nu$ and its ramification divisor $R_{\overline{h}}$ is $\overline{h}^* \mathfrak{c} - \mathfrak{c}$. In fact, $K_X + D = h^*(K_X + D)$ implies $K_{\overline{D}} + \mathfrak{c} = \overline{h}^*(K_{\overline{D}} + \mathfrak{c})$. By choosing an int-amplified $h = \mathcal{I}$, \mathfrak{c} is reduced and \overline{h}^{-1} -invariant for any h (cf. [Men17, Theorem 3.3], [NZ10, Lemma 5.3, the arxiv version]). Then $\mathfrak{c} = \nu^{-1}(Z)$ and $h^{-1}(Z) = Z$. So the claim is proved.

6.8. Some notation. By Theorem 6.2, we have the commutative diagrams:

$$\begin{array}{ccc} f \circlearrowleft X & \xleftarrow{\mu_X} & \tilde{X} \circlearrowleft \tilde{f} \\ \pi \downarrow & & \downarrow \tilde{\pi} \\ g \circlearrowleft Y & \xleftarrow{\mu_Y} & A \circlearrowleft g_A \end{array}$$

$$\begin{array}{ccc} \mathcal{I} \circlearrowleft X & \xleftarrow{\mu_X} & \tilde{X} \circlearrowleft \tilde{\mathcal{I}} \\ \pi \downarrow & & \downarrow \tilde{\pi} \\ \mathcal{I}|_Y \circlearrowleft Y & \xleftarrow{\mu_Y} & A \circlearrowleft \mathcal{I}_A \end{array}$$

where \tilde{X} is a \mathbb{Q} -Gorenstein klt normal projective variety, A is an abelian variety, μ_Y is a finite surjective morphism, and μ_X is a finite surjective quasi-étale morphism. Note that $\mathcal{I}|_Y, \mathcal{I}_A, \tilde{\mathcal{I}}$ are still int-amplified; see [Men17, Section 3]. Note that \tilde{X} still has terminal singularities by [KM98, Proposition 5.20]. Then \tilde{X} is \mathbb{Q} -factorial by Proposition 6.22.

Claim 6.9. $\deg \pi|_Z = m \geq 3$.

By the generic smoothness, $X_y \cap Z = X_y \cap \text{Sing}(D) = \text{Sing}(D \cap X_y)$ for general $y \in Y_0$. Then $\sharp Z \cap X_y = m \geq 3$ for general $y \in Y_0$. So the claim is proved.

Claim 6.10. $\pi|_C : C \rightarrow Y$ is a finite surjective morphism for any \mathcal{I}^{-1} -periodic irreducible closed subvariety C of codimension 2 in X .

Since π has connected fibers, $\pi(C)$ is $\mathcal{I}|_Y^{-1}$ -periodic by [CMZ17, Lemma 7.5] and hence $\mu_Y^{-1}(\pi(C))$ is \mathcal{I}_A^{-1} -periodic. By [Men17, Lemma 4.1], $\mu_Y^{-1}(\pi(C)) = A$ and hence $\pi(C) = Y$. Note that $\dim(C) = \dim(Y)$ and $\pi|_C$ is equidimensional (cf. [Men17, Lemma 4.3]). So $\pi|_C$ is finite surjective and the claim is proved.

Claim 6.11. Z is the union of \mathcal{I}^{-1} -periodic irreducible closed subvarieties of codimension 2 in X .

First note that Z is \mathcal{I}^{-1} -invariant by Claim 6.7. Let C be an \mathcal{I}^{-1} -periodic irreducible closed subvariety of codimension 2 in X . For any $y \in Y_0$, X_y is \mathcal{I}^s -invariant for some $s > 0$. Then $C \cap X_y$ is $(\mathcal{I}^s|_{X_y})^{-1}$ -invariant. Recall that $Z \cap X_y$ is exactly the set of $(\mathcal{I}^s|_{X_y})^{-1}$ -periodic points. By Claim 6.10, π_C is finite surjective and hence $C \cap X_y \subseteq Z \cap X_y$. Then $C = \overline{\bigcup_{y \in Y_0} C \cap X_y} \subseteq Z$. So the claim is proved.

Claim 6.12. Z has exactly one or two irreducible components.

Note that the general fiber of $\pi|_D : D \rightarrow Y$ has $m \geq 3$ irreducible components. Recall that $\nu : \overline{D} \rightarrow D$ is the normalization. Note that the general fiber $\pi|_D \circ \nu$ has m connected (and irreducible) components.

Let B be an irreducible component of $\overline{D}_y := (\pi|_D \circ \nu)^{-1}(y)$ for some general $y \in Y_0$. Since \overline{D} is normal, $\text{Sing}(\overline{D})$ does not dominate Y and hence \overline{D}_y and B are smooth. Note that $\nu|_B : B \rightarrow \nu(B)$ is a finite surjective birational morphism since D is smooth at $\nu(B) \setminus \text{Sing}(D) \neq \emptyset$. Recall that $\nu(B)$ is a smooth rational curve lying on the boundary divisor of the smooth toric Fano surface X_y . So $\nu|_B$ is isomorphic. In particular,

$$\nu^{-1}(Z) \cap \overline{D}_y = \bigsqcup_{B \subseteq \overline{D}_y} (\nu^{-1}(Z) \cap B) = \bigsqcup_{B \subseteq \overline{D}_y} (\nu^{-1}(Z \cap \nu(B)) \cap B).$$

Note that $\sharp Z \cap \nu(B) = 2$. So $\sharp \nu^{-1}(Z) \cap \overline{D}_y = 2m$ and hence $\deg(\pi|_D \circ \nu)|_{\nu^{-1}(Z)} = 2m$.

Let $p_1 : \overline{D} \rightarrow \overline{Y}$ and $p_2 : \overline{Y} \rightarrow Y$ be the Stein factorization of $\pi|_D \circ \nu$. Note that $\deg p_2 = m$. Then $\deg(p_1)|_{\nu^{-1}(Z)} = 2m/m = 2$. Since \overline{Y} is irreducible, $\nu^{-1}(Z)$ has at most two irreducible components. Since $Z \neq \emptyset$, the claim is proved.

Claim 6.13. $\tilde{Z} := \mu_X^{-1}(Z)$ is the union of $\tilde{\mathcal{I}}^{-1}$ -periodic irreducible closed subvarieties of codimension 2 in \tilde{X} .

Let $y \in Y_0$ be a general point such that μ_X is quasi-étale over X_y . Since X_y is a smooth toric Fano surface, X_y has no non-isomorphic quasi-étale cover. Then $\tilde{X}_y := \mu_X^{-1}(X_y)$ is a disjoint union of smooth toric Fano surfaces mapping isomorphically to X_y via μ_X . Assume X_y is \mathcal{I}^s -invariant for some $s > 0$. Then $\mu_X^{-1}(Z \cap X_y)$ is exactly the set of $(\tilde{\mathcal{I}}^s|_{\tilde{X}_y})^{-1}$ -periodic points in \tilde{X}_y . So the claim follows from the same proof of Claim 6.11.

Claim 6.14. By choosing μ_Y and μ_X suitably in Theorem 6.2, we may assume further $\deg \tilde{\pi}|_{\tilde{C}} = 1$ for any $\tilde{\mathcal{I}}^{-1}$ -periodic irreducible closed subvariety \tilde{C} of codimension 2 in \tilde{X} .

Let \tilde{C} be an irreducible component of \tilde{Z} . As reasoned in Claim 6.10, $\tilde{\pi}|_{\tilde{C}} : \tilde{C} \rightarrow A$ is finite surjective. Let $\overline{\tilde{C}}$ be the normalization of \tilde{C} . Note that $\overline{\tilde{C}}$ admits an int-amplified endomorphism. By the ramification divisor formula and [Men17, Theorem 1.5], the induced finite surjective morphism $\overline{\tilde{C}} \rightarrow A$ is quasi-étale and hence étale. Then $\overline{\tilde{C}}$ is an abelian variety by [Mum74, Section 18, Theorem]. If $\deg \tilde{\pi}|_{\tilde{C}} > 1$, then we replace \tilde{X} by $\tilde{X} \times_A \overline{\tilde{C}}$ and A by $\overline{\tilde{C}}$. Note that \tilde{f} is also liftable after some iteration. We show that such replacements can be done in only finitely many steps. After each replacement, $\deg \tilde{\pi}|_{\tilde{Z}}$ remains unchanged, while the number of irreducible components of \tilde{Z} strictly increases. This is because the natural embedding $\tilde{C} \hookrightarrow \tilde{C} \times_A \tilde{C}$ and $\deg \tilde{\pi}|_{\tilde{C}} > 1$ imply that the preimage of \tilde{C} in $\tilde{X} \times_A \overline{\tilde{C}}$ splits into at least two irreducible components. Finally, Claim 6.13 implies that after each replacement, \tilde{Z} is still the union of $\tilde{\mathcal{I}}^{-1}$ -periodic irreducible closed subvariety \tilde{C} of codimension 2 in \tilde{X} . In particular, \tilde{Z} has at most (fixed) $\deg \tilde{\pi}|_{\tilde{Z}}$ irreducible components. So we can only do at most finitely many replacements till the claim is satisfied.

In the following, we choose the \tilde{X} in Claim 6.14. Denote by $\tilde{D} := \mu_X^{-1}(D)$. Since $\mu_X : \tilde{X} \rightarrow X$ is quasi-étale, we have $K_{\tilde{X}} = \mu_X^* K_X \sim_{\mathbb{Q}} -\mu_X^* D = -\tilde{D}$. Since D is nef, so is \tilde{D} . By [Uen, Theorem 5.13], $\kappa(\tilde{X}, \tilde{D}) = \kappa(X, D) = 0$. Note that $K_{\tilde{X}}$ is not pseudo-effective. By [MZ19a, Theorem 1.2], replacing \tilde{f} and $\tilde{\mathcal{I}}$ by a positive power, there is an $(\tilde{f}, \tilde{\mathcal{I}})$ -equivariant birational MMP $\varphi : \tilde{X} \dashrightarrow \tilde{X}_r$ and a Fano contraction $\tilde{\pi}_r : \tilde{X}_r \rightarrow \tilde{Y}$. Denote by $\tilde{f}_r := \tilde{f}|_{\tilde{X}_r}$, $\tilde{g}_r := \tilde{f}|_{\tilde{Y}}$ and $\tilde{\mathcal{I}}_r := \tilde{\mathcal{I}}|_{\tilde{X}_r}$. Since A is an abelian variety, this MMP is over A ; see the first paragraph of the proof of Proposition 6.5.

Claim 6.15. $\varphi : \tilde{X} \dashrightarrow \tilde{X}_r$ is a composition of divisorial contractions; and \tilde{X}_r has \mathbb{Q} -factorial terminal singularities.

Note that $K_{\tilde{X}} = \mu_X^* K_X$. Recall that \tilde{X} has only \mathbb{Q} -factorial terminal singularities (cf. 6.8). Since the MMP is over A , whenever a flipping contraction occurs, one can find some $(\tilde{\mathcal{I}}|_{\tilde{Y}})^{-1}$ -invariant subvariety of dimension at most $\dim(\tilde{X}) - 3$ by [CMZ17, Lemma 7.5]. Note that $\tilde{\mathcal{I}}|_{\tilde{Y}}$ is int-amplified and $\dim(A) > \dim(\tilde{X}) - 3$. Then we get a

contradiction by [Men17, Lemma 4.1]. The second assertion of the claim follows from [KM98, Proposition 3.36, Corollary 3.43]. The claim is proved.

Claim 6.16. $\tilde{D}_r := \varphi(\tilde{D})$ is a (nonzero) reduced divisor containing all the $(\tilde{\mathcal{I}}_r)^{-1}$ -periodic irreducible closed subvarieties of codimension 2 in \tilde{X}_r . Further, $\varphi^{-1}(\tilde{D}_r) = \tilde{D}$, $-K_{\tilde{X}_r} \sim_{\mathbb{Q}} \tilde{D}_r$, $\kappa(\tilde{X}_r, \tilde{D}_r) = 0$, and $\tilde{f}_r^* \tilde{D}_r = \delta_{\tilde{f}_r} \tilde{D}_r$.

Note that $\tilde{\mathcal{I}}|_{\tilde{X} \setminus \tilde{D}}$ is quasi-étale. So the φ -exceptional divisor (being \tilde{I}^{-1} -invariant) is contained in \tilde{D} . Denote by \tilde{D}_r^1 the union of divisors contained in \tilde{D}_r . Then $\tilde{\mathcal{I}}_r|_{\tilde{X}_r \setminus \tilde{D}_r^1}$ is quasi-étale and $\text{Supp } R_{\tilde{\mathcal{I}}_r} = \tilde{D}_r^1$. Let \tilde{C}_r be some $(\tilde{\mathcal{I}}_r)^{-1}$ -invariant irreducible closed subvariety of codimension 2 in \tilde{X}_r . By [CMZ17, Lemma 7.5] and [Men17, Lemma 4.1], \tilde{C}_r dominates A and $\deg \tilde{\mathcal{I}}_r|_{\tilde{C}_r} = \deg \tilde{\mathcal{I}}_A$. By the projection formula, $\tilde{\mathcal{I}}_r^* \tilde{C}_r = (\deg \tilde{\mathcal{I}}_r / \deg \mathcal{I}_A) \tilde{C}_r$. Note that $\tilde{\mathcal{I}}_r|_F$ is polarized (after iteration) for the general $\tilde{\mathcal{I}}_r$ -periodic fiber F of $\tilde{X}_r \rightarrow A$ (cf. [Fak03, Theorem 5.1]). So $\deg \tilde{\mathcal{I}}_r / \deg \mathcal{I}_A > 1$. Since $\text{Sing}(\tilde{X}_r)$ is of codimension ≥ 3 in \tilde{X}_r , $\tilde{C}_r \subseteq \tilde{D}_r^1$ by the purity of branch loci. Let E be any prime exceptional divisor of φ . Then $\varphi(E)$ is $\tilde{\mathcal{I}}_r^{-1}$ -periodic and dominates A by [CMZ17, Lemma 7.5] and [Men17, Lemma 4.1]. So $\varphi(E)$ is of codimension 2 in \tilde{X}_r and $\varphi(E) \subseteq \tilde{D}_r^1$. In particular, $\tilde{D}_r = \tilde{D}_r^1$ and $\varphi^{-1}(\tilde{D}_r) = \tilde{D}$. Therefore, $\kappa(\tilde{X}_r, \tilde{D}_r) = \kappa(\tilde{X}, \tilde{D}) = 0$ by [Uen, Theorem 5.13].

By the ramification divisor formula, we have

$$K_{\tilde{X}_r} + \tilde{D}_r = \tilde{\mathcal{I}}_r^*(K_{\tilde{X}_r} + \tilde{D}_r).$$

Therefore, $-K_{\tilde{X}_r} \sim_{\mathbb{Q}} \tilde{D}_r$ by applying [Men17, Theorem 1.1] and [MZ19b, Proposition 3.3]. Note that $\tilde{f}_r^* \tilde{D} = \delta_{\tilde{f}_r} \tilde{D}$ and hence $\tilde{f}_r^* \tilde{D}_r = \delta_{\tilde{f}_r} \tilde{D}_r$. So the claim is proved.

Claim 6.17. $(\tilde{\pi}_r : \tilde{X}_r \rightarrow \tilde{Y}, \tilde{f}_r, \tilde{g})$ satisfies Case TIR with $\dim(\tilde{X}_r) = \dim(\tilde{Y}) + 2$ and $\tilde{Y} = A$.

If $\delta_{\tilde{f}_r} > \delta_{\tilde{g}}$, then Case TIR' and hence Case TIR are satisfied, so $\tilde{Y} = A$ by the condition (A5) of Case TIR and since $\tilde{\pi} : \tilde{X} \rightarrow A$ has connected fibers. Suppose next that $\delta_{\tilde{f}_r} = \delta_{\tilde{g}}$. If $\dim(\tilde{Y}) = \dim(A)$, then $\tilde{Y} = A$ and hence $\delta_f = \delta_{\tilde{g}} = \delta_{g_A} < \delta_f$, a contradiction.

We are only left with the case where $\delta_{\tilde{f}_r} = \delta_{\tilde{g}} > \delta_{g_A}$ and $\dim(\tilde{Y}) = \dim(A) + 1$. Denote by $\tau : \tilde{Y} \rightarrow A$ the induced morphism. Note that \tilde{Y} is \mathbb{Q} -factorial and klt. Since $\text{Sing}(\tilde{Y})$ does not dominate A and the general fiber of $\tilde{\pi} : \tilde{X} \rightarrow A$ is a rational surface, the general fiber of τ is a smooth rational curve. By the adjunction, the restriction of $K_{\tilde{Y}}$ on the general fiber of τ is anti-ample. Hence $K_{\tilde{Y}}$ is not pseudo-effective. Then we may run MMP starting from \tilde{Y} which ends up with A . Since $\dim(\tilde{Y}) = \dim(A) + 1$ and τ has connected fibers, any appearance of divisorial contraction or flip will create some proper $(\tilde{\mathcal{I}}|_A)^{-1}$ -periodic closed subvariety in A by [CMZ17, Lemma 7.5]. So τ has to be a Fano contraction by [Men17, Lemma 4.1]. Note that $D_{\tilde{Y}} := (\tilde{\pi}_r \circ \varphi)(\tilde{Z})$ is an $(\tilde{\mathcal{I}}|_{\tilde{Y}})^{-1}$ -periodic

divisor in \tilde{Y} since $\tilde{\pi}|_{\tilde{Z}} : \tilde{Z} \rightarrow A$ is finite surjective. By Proposition 6.5, $\kappa(\tilde{Y}, D_{\tilde{Y}}) > 0$. Note that $\tilde{\pi}_r^{-1}(D_{\tilde{Y}})$ is $\tilde{\mathcal{I}}_r^{-1}$ -periodic. So $\tilde{\pi}_r^{-1}(D_{\tilde{Y}}) \subseteq \tilde{D}_r$ and hence $\kappa(\tilde{X}_r, \tilde{D}_r) > 0$, a contradiction to Claim 6.16. The claim is proved.

Claim 6.18. $\deg(\tilde{\pi}_r|_{\tilde{C}_r} : \tilde{C}_r \rightarrow \tilde{Y} = A) = 1$ for any $\tilde{\mathcal{I}}_r^{-1}$ -periodic irreducible closed subvariety \tilde{C}_r of codimension 2 in \tilde{X}_r .

By Claim 6.14, it suffices for us to show that \tilde{C}_r is dominated by some $\tilde{\mathcal{I}}^{-1}$ -periodic irreducible closed subvariety of codimension 2 in \tilde{X} . Note that $\tilde{C}_r \subseteq \tilde{D}_r = \varphi(\tilde{D})$ by Claim 6.16. If $\dim(\varphi^{-1}(\tilde{C}_r)) = \dim(A)$, then we are done. If $\dim(\varphi^{-1}(\tilde{C}_r)) = \dim(A) + 1$, then $\varphi(E) = \tilde{C}_r$ for some (φ -exceptional) irreducible component E of \tilde{D} . Recall that E is an irreducible component of $\tilde{D} = \mu_X^{-1}(D)$. So $\mu_X|_E : E \rightarrow D$ is finite surjective and there is at least one $\tilde{\mathcal{I}}^{-1}$ -periodic irreducible closed subvariety $\tilde{C} \subseteq (\mu_X|_E)^{-1}(Z)$ of codimension 2 in \tilde{X} dominating A . Note that $\varphi(E) = \varphi(\tilde{C}) = \tilde{C}_r$. So the claim is proved.

Completion of the proof of Theorem 6.6. By Claim 6.17, we may apply Claims 6.7 and 6.9 to \tilde{X}_r . Then there are at least three $\tilde{\mathcal{I}}_r^{-1}$ -periodic irreducible closed subvarieties of codimension 2 in \tilde{X}_r by Claim 6.18. So we get a contradiction by applying Claim 6.12 to \tilde{X}_r again. This completes the proof of Theorem 6.6. \square

Corollary 6.19. *Let X be a normal projective variety over K such that $X_{\overline{K}}$ has only \mathbb{Q} -factorial Kawamata log terminal (klt) singularities and admits an int-amplified endomorphism. Then Conjecture 1.4 holds for any surjective endomorphism on X in the following cases:*

- (1) *The group $\pi_1^{\text{alg}}((X_{\overline{K}})_{\text{reg}})$ is finite (e.g. when X is smooth and rationally connected).*
- (2) *$X_{\overline{K}}$ has only terminal singularities and $\dim(X) = 3$.*

Proof. It follows from Theorems 6.1, 6.3 and 6.6. Note that for (2), if $X_{\overline{K}}$ has terminal singularities, then the birational MMP involves only with the terminal singularities (cf. [KM98, Corollary 3.43]). \square

In the rest of this section, we show the \mathbb{Q} -factorial property of \tilde{X} (Proposition 6.22) used in the proof of Theorem 6.6 (i.e., in 6.8). A local ring (R, \mathfrak{m}) is \mathbb{Q} -factorial if for any prime ideal \mathfrak{p} of height one, $\mathfrak{p} = \sqrt{(f)}$ for some $f \in \mathfrak{p}$.

Lemma 6.20. *Let $\pi : X \rightarrow Y$ be a finite surjective morphism of normal varieties. Let $x \in X$ be a closed point and $y = \pi(x)$. Suppose the (completion) local ring $\hat{\mathcal{O}}_{X,x}$ is \mathbb{Q} -factorial. Then so is $\hat{\mathcal{O}}_{Y,y}$.*

Proof. Set $A := \hat{\mathcal{O}}_{Y,y}$ and $B := \hat{\mathcal{O}}_{X,x}$. Let $\varphi : A \rightarrow B$ be the induced local homomorphism, which is an injective finite morphism (cf. [AM69, Proposition 10.13]). We may regard A

as a subring of B . Let \mathfrak{p} be any prime ideal of height 1 on A . Take a prime ideal \mathfrak{q} in B of height 1 such that $\mathfrak{p} = \mathfrak{q} \cap A$. Since B is \mathbb{Q} -factorial, we can take $\beta \in \mathfrak{q}$ such that $\mathfrak{q} = \sqrt{\beta B}$. Take any $\alpha \in \mathfrak{p}$. Now $\alpha \in \mathfrak{q} = \sqrt{\beta B}$. So $\alpha^k = c\beta$ for some $k > 0$ and $c \in B$. Taking norm, we have $\alpha^{kr} = N(c)N(\beta)$ where $r = \dim_{Q(A)} Q(B)$. Note that $N(B) \subseteq A$ since φ is finite. Hence we have $\mathfrak{p} \subseteq \sqrt{(N(\beta))A}$. Conversely, $N(\beta) \in \mathfrak{p}$ since $N(\beta) \in \mathfrak{q} \cap A = \mathfrak{p}$. So we have $\mathfrak{p} = \sqrt{N(\beta)A}$. \square

Lemma 6.21. *Let X be a normal variety. Then we have the following:*

- (1) *Let $x \in X$ be a closed point. If $\hat{\mathcal{O}}_{X,x}$ is \mathbb{Q} -factorial, then so is $\mathcal{O}_{X,x}$.*
- (2) *If $\mathcal{O}_{X,x}$ is \mathbb{Q} -factorial for any closed point $x \in X$, then X is \mathbb{Q} -factorial.*

Proof. (1) Let $S = \text{Spec } \mathcal{O}_{X,x}$ and $\hat{S} = \text{Spec } \hat{\mathcal{O}}_{X,x}$. Let $\pi : \hat{S} \rightarrow S$ be the completion morphism which is faithfully flat. Since S is normal, so is \hat{S} . Take a prime divisor D on S . Since π is flat, $\pi^*\mathcal{O}_S(D)$ is a reflexive sheaf of rank 1 on \hat{S} . So $\pi^*\mathcal{O}_S(D) = \mathcal{O}_{\hat{S}}(\hat{D})$ for some Weil divisor \hat{D} on \hat{S} . If $\hat{\mathcal{O}}_{X,x}$ is \mathbb{Q} -factorial, then $n\hat{D}$ is Cartier for some $n > 0$. Then $\pi^*\mathcal{O}_S(nD) \cong \pi^*((\mathcal{O}_S(D)^{\otimes n})^{\vee\vee}) \cong (\pi^*\mathcal{O}_S(D)^{\otimes n})^{\vee\vee} \cong \mathcal{O}_{\hat{S}}(n\hat{D})$ is invertible. By the fpqc descent along π , $\mathcal{O}_S(nD)$ is invertible.

(2) Take any prime divisor D of X . Take any closed point $x \in D$. Take an affine neighbourhood $U = \text{Spec } A \subseteq X$ of x . Let $\mathfrak{p} \subseteq A$ be the prime ideal of height 1 representing D and $\mathfrak{m} \subseteq A$ the maximal ideal representing x . By assumption, $\mathfrak{p}A_{\mathfrak{m}} = \sqrt{\alpha A_{\mathfrak{m}}}$ for some $\alpha \in \mathfrak{p}$. Write $\mathfrak{p} = (f_1, \dots, f_r)$. Then there exists $n > 0$ such that $f_i^n \in \alpha A_{\mathfrak{m}}$ for all i . Write $f_i^n = \alpha a_i / b_i$ for some $a_i \in A$ and $b_i \in A \setminus \mathfrak{m}$. Then $f_i \in \sqrt{\alpha A_b}$ where $b = b_1 \cdots b_r \in A \setminus \mathfrak{m}$. In particular, $\mathfrak{p}A_b = \sqrt{\alpha A_b}$ and $\text{Spec } A_b$ is an affine open neighbourhood of x . \square

Proposition 6.22. *Let $\pi : X \rightarrow Y$ be a surjective morphism of normal projective varieties with connected fibers. Let $f : X \rightarrow X$ and $g : Y \rightarrow Y$ be int-amplified endomorphisms such that $g \circ \pi = \pi \circ f$. Suppose Y is an abelian variety and X has only terminal singularities. Suppose further $\dim(X) \leq \dim(Y) + 2$. Then X is \mathbb{Q} -factorial.*

Proof. Denote by S the set of closed points $x \in X$ such that $\hat{\mathcal{O}}_{X,x}$ is not \mathbb{Q} -factorial. By Lemma 6.20, $f^{-1}(S) \subseteq S$. Then $f^{-1}(\overline{S}) \subseteq \overline{S}$ by [CMZ17, Lemma 7.2]. Hence $f^{-1}(\overline{S}) = \overline{S}$. By [CMZ17, Lemma 7.5], $\pi(\overline{S})$ is g^{-1} -periodic. Note that $S \subseteq \overline{S} \subseteq \text{Sing}(X)$ and $\dim(\text{Sing}(X)) \leq \dim(X) - 3 < \dim(Y)$ since X has terminal singularities. So $\pi(\overline{S}) = \emptyset$ and hence $S = \emptyset$ by [Men17, Lemma 4.1]. By Lemma 6.21, X is \mathbb{Q} -factorial. \square

7. FIBER SPACES; UBC VS sAND; PROOFS OF THEOREMS 1.10 AND 1.13

In this section, we study the set Z_f for endomorphisms f on a fibration which fixes fibers set-theoretically, i.e. a family of endomorphisms. First we formulate a relative version of the sAND Conjecture.

Let X, S be quasi-projective varieties over K and $\pi : X \rightarrow S$ a projective morphism. Let $f : X \rightarrow X$ be a surjective morphism over S . Take any immersion $X \rightarrow P$ into a projective variety P , any ample divisor H on P , and height $h_H \geq 1$ associated with H . For any point $x \in X(\overline{K})$, setting $X_{\pi(x)} := \pi^{-1}(\pi(x))$, we define

$$\alpha_f(x) := \lim_{n \rightarrow +\infty} h_H(f^n(x))^{1/n} = \lim_{n \rightarrow +\infty} h_{H|_{X_{\pi(x)}}}((f|_{X_{\pi(x)}})^n(x))^{1/n} = \alpha_{f|_{X_{\pi(x)}}}(x)$$

which is well-defined in this relative setting, since the arithmetic degree is well-defined and independent of the choice of ample height on any projective scheme over a number field. We define Z_f and $Z_f(d)$ in the same way as in Definition 1.3.

Conjecture 7.1 (Relative sAND Conjecture). *Let X, S be quasi-projective varieties over K and $\pi : X \rightarrow S$ a projective morphism. Let $f : X \rightarrow X$ be a surjective morphism with $\pi \circ f = \pi$. Then the set $Z_f(d)$ is not Zariski dense in $X_{\overline{K}}$ for any constant $d > 0$.*

Remark 7.2. Suppose π has geometrically irreducible fibers. Then the dynamical degree of $f_{\overline{K}} : X_{\overline{K}} \rightarrow X_{\overline{K}}$ is equal to the dynamical degree of $f|_{X_s} : X_s \rightarrow X_s$ for general $s \in S(\overline{K})$, by the product formula involving the relative dynamical degree (cf. [Tru15]).

Remark 7.3. If X is projective, then Conjecture 7.1 is a special case of Conjecture 1.4.

First we consider projective bundles over projective bases.

Theorem 7.4 (cf. [Ame03]). *Let f be a surjective endomorphism on a projective bundle $\mathbb{P}_Y(E) \xrightarrow{\pi} Y$ over a projective variety with $\pi \circ f = \pi$. Then $Z_f(d)$ is not Zariski dense for any $d > 0$.*

Proof. We may assume that $m := \delta_f > 1$. Replacing f by a power if necessary, we may further assume that $m > r + 1$. Take a Zariski open cover $Y = \bigcup_{\lambda} U_{\lambda}$ with local trivializations $\psi_{\lambda} : \pi^{-1}(U_{\lambda}) \rightarrow U_{\lambda} \times \mathbb{P}^r$ and set $f_{\lambda} = \psi_{\lambda} \circ f \circ \psi_{\lambda}^{-1} : U_{\lambda} \times \mathbb{P}^r \rightarrow U_{\lambda} \times \mathbb{P}^r$. Using local trivializations, we obtain a morphism $\Phi : Y \rightarrow R^m(\mathbb{P}^r, \mathbb{P}^r)/\mathrm{PGL}(r+1)$ defined by $\Phi(y) = f_y$, where $R^m(\mathbb{P}^r, \mathbb{P}^r)$ is the space of endomorphisms given by degree m polynomials on \mathbb{P}^r and $\mathrm{PGL}(r+1)$ acts on $R^m(\mathbb{P}^r, \mathbb{P}^r)$ by conjugation. By Amerik [Ame03, Proposition 1.1], $R^m(\mathbb{P}^r, \mathbb{P}^r)/\mathrm{PGL}(r+1)$ is affine if $m > r + 1$. So Φ is a constant map since Y is projective.

Let $t \in R^m(\mathbb{P}^r, \mathbb{P}^r)$ be an element whose class in $R^m(\mathbb{P}^r, \mathbb{P}^r)/\mathrm{PGL}(r+1)$ is the image of Φ . Set $f_{\lambda, y} = f_{\lambda}|_{\{y\} \times \mathbb{P}^r} \in R^m(\mathbb{P}^r, \mathbb{P}^r)$. Then there exists $h_{\lambda} : U_{\lambda} \rightarrow \mathrm{PGL}(r+1)$ such that

$f_{\lambda,y} = h_\lambda(y) \cdot t$, where $h_\lambda(y) \cdot t = h_\lambda(y) \circ t \circ h_\lambda(y)^{-1}$. Set $G_{\lambda\mu} = \psi_\lambda \circ \psi_\mu^{-1} : U_{\lambda\mu} \times \mathbb{P}^r \rightarrow U_{\lambda\mu} \times \mathbb{P}^r$. Then $G_{\lambda\mu}$ induces $g_{\lambda\mu} : U_{\lambda\mu} \rightarrow \mathrm{PGL}(r+1)$ which satisfies $f_{\lambda,y} = g_{\lambda\mu}(y) \cdot f_{\mu,y}$ for any $y \in U_{\lambda\mu}$. Therefore we have $(h_\lambda^{-1}(y) \circ g_{\lambda\mu}(y) \circ h_\mu(y)) \cdot t = t$ for any $y \in U_{\lambda\mu}$.

Set $\tilde{g}_{\lambda\mu} = h_\lambda^{-1} \circ g_{\lambda\mu} \circ h_\mu$ and $\tilde{f}_{\lambda,y} = h_\lambda^{-1}(y) \cdot f_{\lambda,y} (= t)$. Let $\tilde{G}_{\lambda\mu} : U_{\lambda\mu} \times \mathbb{P}^r \rightarrow U_{\lambda\mu} \times \mathbb{P}^r$ and $H_\lambda : U_\lambda \times \mathbb{P}^r \rightarrow U_\lambda \times \mathbb{P}^r$ be morphisms corresponding to $\tilde{g}_{\lambda\mu}$, h_λ , respectively.

$$\begin{array}{ccc} U_{\lambda\mu} \times \mathbb{P}^r & \xrightarrow{H_\mu} & U_{\lambda\mu} \times \mathbb{P}^r \\ \tilde{G}_{\lambda\mu} \downarrow & & \downarrow G_{\lambda\mu} \\ U_{\lambda\mu} \times \mathbb{P}^r & \xrightarrow{H_\lambda} & U_{\lambda\mu} \times \mathbb{P}^r \end{array}$$

This commutative diagram shows that $(h_\lambda)_\lambda$ defines an isomorphism $h : \tilde{X} \rightarrow X$, where \tilde{X} is the projective bundle glued by $(\tilde{G}_{\lambda\mu})_{\lambda,\mu}$. Now we have

$$\begin{aligned} \tilde{g}_{\lambda\mu}(y) \cdot \tilde{f}_{\mu,y} &= (h_\lambda^{-1}(y) \circ g_{\lambda\mu}(y) \circ h_\mu(y)) \cdot \tilde{f}_{\mu,y} \\ &= (h_\lambda^{-1}(y) \circ g_{\lambda\mu}(y)) \cdot f_{\mu,y} \\ &= h_\lambda^{-1}(y) \cdot f_{\lambda,y} \\ &= \tilde{f}_{\lambda,y}. \end{aligned}$$

So $(\tilde{f}_\lambda)_\lambda$ defines a surjective endomorphism $\tilde{f} : \tilde{X} \rightarrow \tilde{X}$. Note that $Z_{\tilde{f}}(d)$ is not Zariski dense for any $d > 0$ since $(\tilde{f})_y$ is invariant when y moves on local trivializations. Moreover, $h \circ \tilde{f} = f \circ h$ by definition of \tilde{f} . So the assertion follows from Lemma 2.6. \square

The next result gives the structure of the small arithmetic degree set Z_f in the case of abelian fibrations, and is the key in proving Theorem 1.13.

Theorem 7.5. *Let $\pi : X \rightarrow Y$ be an abelian fibration, that is, a projective surjective morphism of quasi-projective normal varieties such that $\pi_* \mathcal{O}_X = \mathcal{O}_Y$ and its generic fiber X_η is an abelian variety. Let $f : X \rightarrow X$ be a surjective endomorphism with $\delta_f > 1$ and $\pi \circ f = \pi$. Then, after enlarging the ground field by a finite extension and taking base change along a finite cover of Y , there is a Zariski open dense subset $V \subset Y$ such that $X_V = \pi^{-1}(V) \rightarrow V$ has an abelian scheme structure compatible with the abelian variety structure of the generic fibre X_η , and there are a non-trivial V -group homomorphism $\beta : X_V \rightarrow X_V$ and a section $M \subset X_V$ of $X_V \rightarrow V$ such that, for any $d > 0$, we have:*

$$Z_f(d) \subseteq \pi^{-1}(Y \setminus V) \cup (M(d) + \beta^{-1}(\mathrm{Tor}(X_V(d)))) .$$

Here $\mathrm{Tor}(X_V(d))$ is the set of rational points in $X(d)$ which are torsion points of the fibers over V and $M(d) + \beta^{-1}(\mathrm{Tor}(X_V(d))) = \{a + b \mid \pi(a) = \pi(b), a \in M(d), b \in \beta^{-1}(\mathrm{Tor}(X_V(d)))\}$.

Consider first the case when the endomorphism induces an isogeny on the generic fiber.

Lemma 7.6. *Let $\pi : X \rightarrow Y$ be an abelian fibration and $f : X \rightarrow X$ a surjective endomorphism such that $\delta_f > 1$, $\pi \circ f = \pi$, and the induced endomorphism $f_\eta : X_\eta \rightarrow X_\eta$ on the generic fiber of π is an isogeny. Then, after enlarging the ground field by a finite extension and taking base change along a finite cover of Y , there are a Zariski open dense subset $V \subset Y$ over which $X_V = \pi^{-1}(V) \rightarrow V$ has an abelian scheme structure restricting to the abelian variety structure of the generic fibre X_η and a non-trivial V -group homomorphism $\beta : X_V \rightarrow X_V$ such that, for any $d > 0$, we have:*

$$Z_f(d) \subseteq \pi^{-1}(Y \setminus V) \cup \beta^{-1}(\text{Tor}(X_V(d))).$$

Proof. Let $X_{\bar{\eta}}$ be the geometric generic fiber, and $f_{\bar{\eta}}$ the induced endomorphism. Then, by a calculation of intersection numbers, we have $\delta_{f_{\bar{\eta}}} = \delta_f = \delta_{f_y}$ where y is a general closed point of Y . Denote this number as δ .

We can find nef \mathbb{R} -divisors $\bar{D} \neq 0$ on $X_{\bar{\eta}}$ such that $f_{\bar{\eta}}^* \bar{D} \sim_{\mathbb{R}} \delta \bar{D}$ (cf. [San17, Remark 5.11]). Fix an ample divisor H on X . By Lemma 2.10, there exists some $\bar{\alpha} \in \text{End}(X_{\bar{\eta}})_{\mathbb{R}}$ such that $\Phi_{\bar{D}}^{H_{\bar{\eta}}} = i_{H_{\bar{\eta}}}(\bar{\alpha}) \circ \bar{\alpha}$ and $i_{H_{\bar{\eta}}}(\bar{\alpha}) = \bar{\alpha}$. Note that $\bar{\alpha}$ is not zero since $\bar{D} \neq 0$.

Take a sufficiently large finite extension F of $k(\eta)$ so that \bar{D} and $\bar{\alpha}$ descend to an \mathbb{R} -Cartier divisor D_F on X_F and $\alpha_F \in \text{End}(X_F)_{\mathbb{R}}$ respectively. Moreover, we can take F so that $f_F^* D_F \sim_{\mathbb{R}} \delta D_F$, $\Phi_{D_F}^{H_F} = i_{H_F}(\alpha_F) \circ \alpha_F$, and $i_{H_F}(\alpha_F) = \alpha_F$. (For the definition of $\Phi_{D_F}^{H_F}$, see Lemma 2.11 and Remark 2.13). Represent α_F as follows:

$$\alpha_F = \sum_{i=1}^r \beta_{i,F} \otimes c_i$$

where $\beta_{i,F} \in \text{End}(X_F)$ and $c_i \in \mathbb{R}$ are linearly independent over \mathbb{Q} .

Replacing Y with its normalization in F and X with the base change, we may assume $F = k(\eta)$. (Here we also extend the ground field so that Y is geometrically irreducible.) Moreover, by shrinking Y , we may assume $\pi : X \rightarrow Y$ has an abelian scheme structure compatible with the one on the generic fiber. Further, one can find:

- an \mathbb{R} -Cartier divisor D on X which restricts to D_F ; and
- $\beta_i \in \text{End}(X/Y)$, group endomorphisms of abelian group scheme X over Y restricting to $\beta_{i,F}$

such that $f^* D \sim_{\mathbb{R}} \delta D$, $\varphi_H^{-1} \circ \varphi_D = i_H(\alpha) \circ \alpha$ in $\text{End}(X/Y)_{\mathbb{R}}$ where $\alpha = \sum_{i=1}^r \beta_i \otimes c_i$, and $i_H(\alpha) = \alpha$.

Take any $y \in Y(\bar{K})$. On X_y we have

- $f_y^* D_y \sim_{\mathbb{R}} \delta D_y$;
- $\varphi_{H_y}^{-1} \circ \varphi_{D_y} = i_{H_y}(\alpha_y) \circ \alpha_y$, $i_{H_y}(\alpha_y) = \alpha_y$ in $\text{End}(X_y)_{\mathbb{R}}$; and
- $\alpha_y = \sum_{i=1}^r \beta_{i,y} \otimes c_i$.

Now, since the isogeny f_y commutes with the multiplication map $[m]$ (with $m \geq 2$) on X_y , by [HS00, Exercise B.5], we have

$$\hat{h}_{D_y}(x) = \lim_{n \rightarrow +\infty} \frac{h_{D_y}(f_y^n(x))}{\delta^n} = \lim_{n'=m^n \rightarrow +\infty} \frac{h_{D_y}(n'x)}{(n')^2}$$

for all $x \in X_y(\overline{K})$. Then as in the proof of [KS16b, Theorem 29], we have

$$\hat{h}_{D_y}(x) = \frac{1}{2} \langle \alpha_y(x), \alpha_y(x) \rangle_{H_y}$$

where \langle, \rangle_{H_y} is the height pairing associated with the quadratic part of the canonical height associated with H_y .

Fix some $d > 0$. Since \langle, \rangle_{H_y} is positive definite on $X_y(d) \otimes_{\mathbb{Z}} \mathbb{R}$, for $x \in X_y(d)$, we have

$$\begin{aligned} \alpha_f(x) < \delta &\implies \alpha_{f_y}(x) < \delta \implies \hat{h}_{D_y}(x) = 0 \\ &\implies \alpha_y(x) = 0 \text{ in } X_y(d) \otimes_{\mathbb{Z}} \mathbb{R} \\ &\implies \sum_{i=1}^r \beta_{i,y}(x) \otimes c_i = 0 \text{ in } X_y(d) \otimes_{\mathbb{Z}} \mathbb{R}. \end{aligned}$$

Since c_i are linearly independent over \mathbb{Q} , the last equality implies

$$\beta_{i,y}(x) = 0 \text{ in } X_y(d) \otimes_{\mathbb{Z}} \mathbb{Q} \text{ for all } i.$$

There is at least one $\beta := \beta_i$ which is a non-zero Y -group endomorphism of the abelian scheme $X \rightarrow Y$. Then we have $Z_f(d) \subseteq \beta^{-1}(\text{Tor}(X(d)))$. \square

Proof of Theorem 7.5. First we apply Lemma 5.2 and Remark 5.3 to the endomorphism $f_{\overline{\eta}} : X_{\overline{\eta}} \rightarrow X_{\overline{\eta}}$ on the geometric generic fiber of π . Then, enlarging the ground field by a finite extension and Y with a Zariski open dense subset of its finite cover, we may assume:

- $X = X_1 \times_Y X_2$ where $\pi_i : X_i \rightarrow Y$ are abelian scheme over Y ($i = 1, 2$);
- $f = f_1 \times_Y f_2$ where f_i are surjective endomorphism of X_i over Y ($i = 1, 2$);
- there is a section $p_1 : Y \rightarrow X_1$ such that $g_1 = \tau_{p_1}^{-1} \circ f_1 \circ \tau_{p_1} : X_1 \rightarrow X_1$ is a Y -group endomorphism of the abelian scheme $\pi_1 : X_1 \rightarrow Y$; and
- $\delta_{f_2} = 1$.

By Lemma 7.6, after enlarging the ground field by a finite extension and replacing Y with a Zariski open dense subset of its finite cover again, there exists a non-trivial Y -group homomorphism $\beta_1 : X_1 \rightarrow X_1$ such that

$$Z_{g_1}(d) \subseteq \beta_1^{-1}(\text{Tor}(X_1(d))).$$

Then we have $Z_{f_1}(d) \subseteq M_1(d) + \beta_1^{-1}(\text{Tor}(X_1(d)))$, where $M_1 \subset X_1$ is the image of p_1 .

Take a general point $y \in Y(d)$. Then

$$Z_{f_y}(d) \subseteq (p_1(y) + \beta_1^{-1}(\text{Tor}((X_1)_y(d)))) \times (X_2)_y(d).$$

Extend p_1, β_1 to $p : Y \rightarrow X$, $\beta : X \rightarrow X$ by $p(y) = (p_1(y), 0)$, $\beta(x) = (\beta_1(x), 0)$. Let $M \subset X$ be the image of p . Then we have $Z_f(d) \subseteq M(d) + \beta^{-1}(\text{Tor}(X(d)))$. \square

We remark that the uniform boundedness conjecture 1.11 is proved for elliptic curves.

Theorem 7.7 (Merel, [Mer96, Corollaire]). *Take any $d > 0$. Then there is a constant $C = C(d) > 0$ such that the number of torsion points in $E(d)$ is at most C for any elliptic curve E defined over K .*

Now we are ready to prove Theorem 1.13.

Proof of Theorem 1.13. By Theorem 7.5, after enlarging the ground field by a finite extension and replacing Y with a Zariski open dense subset of its finite cover, there are a section $M \subset X$ of π and a non-trivial Y -group homomorphism $\beta : X \rightarrow X$ such that $Z_f(d) \subseteq M(d) + \beta^{-1}(\text{Tor}(X(d)))$. By assumption, the multiplication by a sufficiently divisible integer N maps all points of $\text{Tor}(X(d))$ to the zero section E . So we have $Z_f(d) \subseteq M(d) + (\beta^{-1}([N]^{-1}E))$. Thus $Z_f(d)$ is not Zariski dense in $X_{\overline{K}}$. \square

As corollaries, Conjecture 7.1 for elliptic fibrations and trivial abelian fibrations follow.

Theorem 7.8. *Let f be a surjective endomorphism on an elliptic fibration $X \xrightarrow{\pi} Y$ over a quasi-projective variety with $\pi \circ f = \pi$. Then $Z_f(d)$ is not Zariski dense in $X_{\overline{K}}$ for any $d > 0$.*

Theorem 7.9. *Let f be a surjective endomorphism on a trivial abelian fibration $X = A \times Y \xrightarrow{\pi} Y$ over a quasi-projective variety with $\pi \circ f = \pi$. Then $Z_f(d)$ is not Zariski dense in $X_{\overline{K}}$ for any $d > 0$.*

Next, we consider endomorphisms on projective varieties of non-negative Kodaira dimension. The monoid $\text{SEnd}(X)$ of surjective endomorphisms of a variety of general type is a finite set, so Conjecture 1.4 holds for them. For a smooth projective variety X with $\kappa(X) = \dim(X) - 1$, we can prove Conjecture 1.4 as an application of Theorem 7.8.

Theorem 7.10. *Let X be a smooth projective variety with $\kappa(X) = \dim(X) - 1$ and $f : X \rightarrow X$ a surjective endomorphism. Then $Z_f(d)$ is not Zariski dense in $X_{\overline{K}}$ for any $d > 0$.*

Proof. By [NZ09, Theorem A], there is an automorphism g of finite order on the base W of an Iitaka fibration $\phi : X \dashrightarrow W$, such that $\phi \circ f = g \circ \phi$. Replacing f by a positive power, we may assume that $g = \text{id}$ and hence $\phi \circ f = \phi$. By Lemma 2.9, we may replace

X with the normalization Γ of the graph of ϕ and f with its lifting to Γ . Thus we may assume that ϕ is an elliptic fibration. Then apply Theorem 7.8. \square

For a non-invertible endomorphism on a threefold X with $\kappa(X) \geq 0$, we can show the following weak version of non-density thanks to the classification results by Fujimoto and Nakayama ([Fuj02], [FN07]).

Theorem 7.11. *Let X be a smooth projective threefold with Kodaira dimension $\kappa(X) \geq 0$ and $f : X \rightarrow X$ a non-invertible surjective endomorphism. Then $Z_f(L) = \{x \in X(L) \mid \alpha_f(x) < \delta_f\}$ is not Zariski dense in $X_{\overline{K}}$ for any finite extension field $(\overline{K} \supseteq) L \supseteq K$.*

Proof. The proof is in steps.

Step 1. By [Fuj02], we can run an f -equivariant MMP

$$X = X_0 \rightarrow X_1 \rightarrow \cdots \rightarrow X_r = Y$$

such that each step is a blowdown of a smooth surface to an elliptic curve (so each X_i is smooth) and K_Y is nef and hence semi-ample by the known abundance for threefolds. By Lemma 2.6 and replacing X by Y , we may assume that K_X is semi-ample. Take the Iitaka fibration $\phi : X \rightarrow C$. By [NZ09, Theorem A], replacing f by a positive power, we may assume that $\phi \circ f = \phi$. Let F be a general fiber of ϕ . Then $K_F \sim_{\mathbb{Q}} 0$.

If $\kappa(X) = 3$, then f is an automorphism of finite order, contradicting the assumption. The case $\kappa(X) = 2$ is a special case of Theorem 7.10.

Assume $\kappa(X) = 0$. We use the Beauville–Bogomolov decomposition theorem over \overline{K} ([BGRS17, Proposition 3.1]). Then there is a finite étale cover $\phi : \tilde{X} \rightarrow X$ where \tilde{X} is either an abelian threefold or the product of an elliptic curve and a K3 surface, and f lifts to an endomorphism \tilde{f} on \tilde{X} (cf. [LS18, Section 4]). Moreover, if \tilde{X} is the product of an elliptic curve and a K3 surface, then \tilde{f} splits. So the assertion follows from Lemma 2.6, Lemma 2.4, and Theorem 5.6.

From now on, we assume $\kappa(X) = 1$. A general fiber F has a non-invertible (necessarily étale) endomorphism $f|_F$, so it is either an abelian or a hyperelliptic surface, by surface theory or [Fuj02, Theorem 3.2].

Step 2. Assume that F is a hyperelliptic surface. By Fujimoto [Fuj02, Theorem 5.10], ϕ decomposes as $\phi : X \xrightarrow{\pi} T \xrightarrow{q} C$ where T is a normal projective surface, π is an elliptic fibration, and q is a \mathbb{P}^1 -fibration. Moreover, f descends to a morphism $g : T \rightarrow T$. Take a Zariski open dense subset $U \subset C$ such that $\phi_U = \phi|_{X_U} : X_U = \phi^{-1}(U) \rightarrow U$ is smooth. Then we can take the relative Albanese morphism $a : X_U \rightarrow A_U$ over U (cf. [FAG05, Proposition 9.5.20]). Let $h_U : A_U \rightarrow A_U$ be the morphism induced from f . Setting

$T_U = q^{-1}(U)$, the natural morphism $\theta : X_U \rightarrow T_U \times_U A_U$ is induced. Taking the fibers at a general point $c \in U$, we can see that $\theta_c : X_c \rightarrow T_c \times A_c$ is a finite surjective morphism because of the hyperelliptic surface structure of X_c . So θ is a finite surjective morphism.

Consider the endomorphism $g_U \times_U h_U$ on $T_U \times_U A_U$. Since $f|_C = \text{id}$ and by the product formula in [Tru15], the dynamical degree of g (resp. $h_U, g_U \times_U h_U$) is equal to the dynamical degree of the restriction g_c (resp. $h_c, g_c \times h_c$) of g (resp. $h_U, g_U \times_U h_U$) to the fiber at a general point $c \in U$. Further, taking a general point $c \in U$, we have

$$\delta_{g_U \times_U h_U} = \delta_{g_c \times h_c} = \max\{\delta_{g_c}, \delta_{h_c}\} = \max\{\delta_g, \delta_{h_U}\}.$$

Now $\delta_f = \delta_{g_U \times_U h_U}$ since θ is finite surjective. Therefore $\delta_f = \max\{\delta_g, \delta_{h_U}\}$.

Note that $Z_g(d)$ and $Z_{h_U}(d)$ are not Zariski dense (Theorem 4.4 and Theorem 7.8). If $\delta_f = \delta_g$, then $Z_f(d) \subset \pi^{-1}(Z_g(d))$ is not Zariski dense. Similarly if $\delta_f = \delta_{h_U}$, then by applying Lemma 2.6 fiberwise, we get $Z_f(d) \subseteq \phi^{-1}(C \setminus U) \cup a^{-1}(Z_{h_U}(d))$ and again $Z_f(d)$ is not Zariski dense.

Step 3. Assume that F is an abelian surface. If F is a simple abelian surface or ϕ is primitive, then ϕ is a Seifert abelian fibration in the sense of [FN07, Definition 2.3] (cf. [FN07, Theorem 4.1], [FN07, Theorem 4.3 (1)]). Then X admits an finite étale cover $W \rightarrow X$ from a smooth abelian fibration $W \rightarrow T$. Now $\kappa(T) = \kappa(W) = \kappa(X) = 1$ (cf. e.g. [FN07, Lemma 2.1]). Thus T is a curve of genus ≥ 2 , so it is not potentially dense by Faltings theorem. So W is not potentially dense, too. The Chevalley–Weil theorem implies that X is not potentially dense. So $Z_f(L)$ is not Zariski dense for any finite extension field $L \supseteq K$.

Finally assume that F is an abelian surface and ϕ is imprimitive. By [FN07, Theorem 5.6 and Corollary 5.8], ϕ decomposes as $X \xrightarrow{\pi} T \xrightarrow{q} C$ where T is a normal projective surface and π, q are elliptic fibrations. [FN07, Proposition 5.7] implies the following commutative diagram:

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\alpha} & X \\ \tilde{\pi} \downarrow & & \downarrow \pi \\ \tilde{T} & \xrightarrow{\beta} & T \\ \tilde{q} \downarrow & & \downarrow q \\ \tilde{C} & \xrightarrow{\gamma} & C \end{array}$$

where γ is a finite Galois cover, \tilde{T}, \tilde{X} are the normalizations of $T \times_C \tilde{C}, X \times_C \tilde{C}$, respectively, and \tilde{T} is isomorphic to $E \times \tilde{C}$ for some elliptic curve E . Let $\rho : \tilde{X} \rightarrow E$ be the composition of $\tilde{\pi} : \tilde{X} \rightarrow \tilde{T}$ with the projection $\tilde{T} \cong E \times \tilde{C} \rightarrow E$. By [FN07, Theorem

5.10], there is a finite étale cover $\nu : E' \rightarrow E$ such that $\tilde{X}' = \tilde{X} \times_E E'$ is isomorphic to $E' \times S$ for some projective surface S and f on X lifts to endomorphism $\tilde{f}' : E' \times S \rightarrow E' \times S$ which splits as $\tilde{f}' = \phi' \times v$ where ϕ' is an automorphism on S and v is an endomorphism on E . Then $Z_{\tilde{f}'}(d)$ is not Zariski dense by Theorem 3.1 and Theorem 4.1. Now \tilde{X}' is an equivariant étale cover of X , so the theorem follows from Lemma 2.6. \square

Finally, we show Theorem 1.10, the equivalence of the uniform boundedness conjecture and the relative sAND conjecture for a family of polarized endomorphisms.

Proof of Theorem 1.10. “(2) \Rightarrow (1)” We can take a quasi-projective variety U with a morphism $\phi : \mathbb{P}_U^N \rightarrow \mathbb{P}_U^N$ over U which parametrizes all the endomorphisms on \mathbb{P}^N of degree d . So the implication is clear.

“(1) \Rightarrow (3)” Let $\eta \in S$ be the generic point. Now $f_\eta : X_\eta \rightarrow X_\eta$ is a polarized endomorphism. So X_η can be embedded into a projective space over f_η -equivariantly (cf. [Fak03, Proposition 2.1]). After shrinking S , we may assume that X is a closed subvariety of \mathbb{P}_S^N and f extends to a surjective endomorphism $\phi : \mathbb{P}_S^N \rightarrow \mathbb{P}_S^N$ over S .

By assumption, $Z_\phi(d)$ is contained in the closed subset

$$W = \{x \in \mathbb{P}_S^N \mid f^m(x) = f^n(x)\}$$

for some $m > n \geq 0$. Since $f = \phi|_X$ is of infinite order, $X \not\subseteq W$. Hence $Z_f(d) = Z_\phi(d) \cap X \subseteq W \cap X$, a proper closed subset of X . So $Z_f(d)$ is not Zariski dense in $X_{\overline{K}}$.

“(3) \Rightarrow (2)” We argue by induction on $\dim(X)$. By assumption, $W = \overline{Z_f(d)}$ is a proper f -invariant closed subset of X . Take the irreducible decomposition $W = W_1 \cup \dots \cup W_p \cup W'_1 \cup \dots \cup W'_q$ such that W_1, \dots, W_p are f -periodic and W'_j maps to $\bigcup_{i=1}^p W_i$ by iteration of f for $1 \leq j \leq q$. Then $Z_f(d) \subseteq f^{-k}(\bigcup_{i=1}^p W_i)$ for some $k \geq 0$.

Take a sufficiently large $m > 0$ such that W_i is f^m -invariant for $1 \leq i \leq p$. Set $g_i = f^m|_{W_i}$. We may assume that $\dim(W_i) = \dim(\pi(W_i))$ for $1 \leq i \leq r$ and $\dim(W_i) > \dim(\pi(W_i))$ for $r + 1 \leq i \leq p$. Fix i with $1 \leq i \leq r$. Then g_i is an automorphism of finite order on W_i since g_i induces an endomorphism on a general fiber of $\pi|_{W_i}$ which is a finite set. Take some $M > 0$ such that $g_i^M = \text{id}_{W_i}$ for each i . If there is a curve $C \subseteq W_i$ contracted by π , then $0 < (C \cdot H) = (C \cdot (g_i^M)^* H) = r^{mM} (C \cdot H)$, which is a contradiction. So $\pi|_{W_i}$ is finite for $1 \leq i \leq r$.

Now we have

$$Z_f(d) \subseteq f^{-k} \left(\bigcup_{i=1}^r W_i(d) \cup \bigcup_{j=r+1}^p Z_{g_j}(d) \right).$$

By induction hypothesis, the number of the points in $Z_{g_j}(d) \cap X_s$ is uniformly bounded for $r + 1 \leq j \leq p$. Moreover the number of the points of $W_i(d) \cap X_s$ is also uniformly bounded for $1 \leq i \leq r$. So the assertion (2) follows. \square

8. FURTHER GENERALIZATION

We can generalize Conjecture 1.4 to a more precise one:

Conjecture 8.1. *Let $f : X \rightarrow X$ be a surjective endomorphism on a projective variety with the first dynamical degree $\delta_f > 1$. Let $\{Z_\lambda\}_{\lambda \in \Lambda}$ be the collection of all f -preperiodic proper subvarieties of small dynamical degree in $X_{\overline{\mathbb{K}}}$. Then, for any $d > 0$, $Z_f(d) \subseteq Z_{\lambda_1} \cup \cdots \cup Z_{\lambda_m}$ for some $\lambda_1, \dots, \lambda_m \in \Lambda$.*

Remark 8.2. Conjecture 8.1 holds for the following cases.

- (1) Polarized endomorphisms (cf. Theorem 3.1).
- (2) Surjective endomorphisms on abelian varieties (cf. Theorem 5.6).
- (3) Automorphisms on projective surfaces (cf. Theorem 4.1).

In fact, in these cases, the respective theorems have given precise descriptions of Z_f .

Clearly Conjecture 8.1 implies Conjecture 1.4. For the converse, we have:

Proposition 8.3. *Fix a positive integer n . If Conjecture 1.4 holds in dimension $\leq n$, then Conjecture 8.1 holds in dimension n .*

Proof. Let $f : X \rightarrow X$ be a surjective endomorphism on a projective variety X of dimension n with $\delta_f > 1$. Take any $d > 0$. Then $Z_f(d)$ is not Zariski dense by assumption. Let W be the Zariski closure of $Z_f(d)$ in $X_{\overline{\mathbb{K}}}$. We can take the irreducible decomposition of W as $W = \bigcup_i W_i \cup \bigcup_j W'_j$ where each W_i is f -periodic and each W'_j is mapped to $\bigcup_i W_i$ by iteration of f . Then we have $Z_f(d) \subseteq f^{-s}(\bigcup_i W_i)$ for some $s > 0$.

Replacing f by a positive power, we may assume that W_i is f -invariant for each i . Set $g = f|_{W_i}$. By assumption, $Z_f(d) \cap W_i$ is Zariski dense in W_i . So Conjecture 1.4 for g implies $\delta_g < \delta_f$. Therefore $W_i \subseteq Z_f$. Now we have $Z_f(d) \subseteq f^{-s}(\bigcup_i W_i)$, and each irreducible component of $f^{-s}(\bigcup_i W_i)$ is an f -preperiodic subvariety of small dynamical degree. So we are done. \square

Conjecture 8.1 entails that studying f -preperiodic subvarieties (especially those of small dynamical degree) is important in order to understand Z_f . We conclude this section with some questions on f -preperiodic subvarieties.

Question 8.4. *Let $f : X \rightarrow X$ be a surjective endomorphism on a projective variety with $\delta_f > 1$. Fix $d > 0$.*

- (1) *Let Y be an f -invariant subvariety of $X_{\overline{\mathbb{K}}}$. Then, is $X(d) \cap (f^{-s}(Y) \setminus f^{-(s-1)}(Y))$ empty for $s \gg 0$?*
- (2) *Is there a positive integer N such that $Y(d)$ is empty for any f -periodic subvariety $Y \subseteq X_{\overline{\mathbb{K}}}$ of small dynamical degree whose period is larger than N ?*

- (3) *Is the number of maximal f -invariant subvarieties of small dynamical degree in $X_{\overline{K}}$ finite?*

Remark 8.5. (1) and (2) are arithmetic questions but (3) is a purely geometric question. Note that (3) is true for abelian varieties (cf. Theorem 5.6) and projective surfaces (cf. Theorem 4.1 and [MZ19b, Section 5]).

REFERENCES

- [Ame03] E. Amerik, On endomorphisms of projective bundles, *Manuscripta Math.* **111** (2003), no. 1, 17–28.
- [AM69] M. Atiyah and I. Macdonald, *Introduction to commutative algebra*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont, 1969.
- [BGRS17] J. P. Bell, D. Ghioca, Z. Reichstein and M. Santriano, On the Medvedev–Scanlon conjecture for minimal threefolds of non-negative Kodaira dimension, *New York J. Math.* **23** (2017), 1185–1203.
- [Birk] G. Birkhoff, Linear transformations with invariant cones, *Amer. Math. Monthly* **74** (1967), 274–276.
- [CS93] G. S. Call and J. H. Silverman, Canonical heights on varieties with morphisms, *Compositio Math.* **89** (1993), no. 2, 163–205.
- [CS16] G. S. Call and J. H. Silverman, Degeneration of dynamical degrees in families of maps, *arXiv:1609.02119*
- [CMZ17] P. Cascini, S. Meng and D.-Q. Zhang, Polarized endomorphisms of normal projective threefolds in arbitrary characteristic, *Math. Ann.*, <https://doi.org/10.1007/s00208-019-01877-6>
- [Deb01] O. Debarre, *Higher-dimensional algebraic geometry*, Universitext, Springer-Verlag, 2001.
- [Fak03] N. Fakhruddin, Questions on self-maps of algebraic varieties, *J. Ramanujan Math. Soc.*, **18**(2):109–122, 2003.
- [FAG05] B. Fantechi, L. Göttsche, L. Illusie, S. L. Kleiman, N. Nitsure and A. Vistoli, *Fundamental algebraic geometry. Grothendieck’s FGA explained*. Mathematical Surveys and Monographs, 123. American Mathematical Society, Providence, RI.
- [Fuj02] Y. Fujimoto, Endomorphisms of smooth projective 3-folds with nonnegative Kodaira dimension, *Publ. RIMS, Kyoto Univ.* **38** (2002), 33–92.
- [FN07] Y. Fujimoto and N. Nakayama, Endomorphisms of smooth projective 3-folds with nonnegative Kodaira dimension, II, *J. Math. Kyoto. Univ.* **47** (2007), no. 1, 79–114.
- [HS00] M. Hindry and J. H. Silverman, *Diophantine Geometry: An Introduction*, Springer-Verlag, New York, 2000.
- [Kaw08] S. Kawaguchi, Projective surface automorphisms of positive topological entropy from an arithmetic viewpoint, *Amer. J. Math.* **130** (2008), no. 1, 159–186.
- [KS14] S. Kawaguchi and J. H. Silverman, Examples of dynamical degree equals arithmetic degree, *Michigan Math. J.* **63** (2014), no. 1, 41–63.
- [KS16a] S. Kawaguchi and J. H. Silverman, On the dynamical and arithmetic degrees of rational self-maps of algebraic varieties, *J. Reine Angew. Math.* **713** (2016), 21–48.

- [KS16b] S. Kawaguchi and J. H. Silverman, Dynamical canonical heights for Jordan blocks, arithmetic degrees of orbits, and nef canonical heights on abelian varieties, *Trans. Amer. Math. Soc.* **368** (2016), no. 7, 5009–5035.
- [KM98] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Univ. Press, 1998.
- [Lan83] S. Lang, *Fundamentals of Diophantine Geometry*, Springer-Verlag, New York, 1983.
- [LS18] J. Lesieutre and M. Santriano, Canonical heights on hyper-Kähler varieties and the Kawaguchi–Silverman Conjecture, *Int. Math. Res. Not.* (to appear), arXiv:**1802.07388**
- [Mat16] Y. Matsuzawa, On upper bounds of arithmetic degrees, arXiv:**1606.00598**
- [Mat19] Y. Matsuzawa, Kawaguchi–Silverman Conjecture for endomorphisms on several classes of varieties, arXiv:**1902.06072**
- [MS18] Y. Matsuzawa and K. Sano, Arithmetic and dynamical degrees of self-morphisms of semi-abelian varieties, arXiv:**1802.03173**
- [MSS18] Y. Matsuzawa, K. Sano and T. Shibata, Arithmetic degrees and dynamical degrees of endomorphisms on surfaces, *Algebra Number Theory* **12** (2018), no. 7, 1635–1657.
- [MY19] Y. Matsuzawa and S. Yoshikawa, Kawaguchi–Silverman Conjecture for endomorphisms on rationally connected varieties admitting an int-amplified endomorphism, arXiv:**1908.11537**
- [Men17] S. Meng, Building blocks of amplified endomorphisms of normal projective varieties, *Math. Z.* (to appear), arXiv:**1712.08995**
- [MZ18] S. Meng and D.-Q. Zhang, Building blocks of polarized endomorphisms of normal projective varieties, *Adv. Math.* **325** (2018), 243–273.
- [MZ19] S. Meng and D.-Q. Zhang, Characterizations of toric varieties via polarized endomorphisms, *Math. Z.* **292** (2019), no. 3-4, 1223–1231.
- [MZ19a] S. Meng and D.-Q. Zhang, Semi-group structure of all endomorphisms of a projective variety admitting a polarized endomorphism, *Math. Res. Lett.* (to appear), arXiv:**1806.05828**
- [MZ19b] S. Meng and D.-Q. Zhang, Kawaguchi–Silverman Conjecture for surjective endomorphisms, arXiv:**1908.01605**
- [Mer96] L. Merel, Bornes pour la torsion des courbes elliptiques sur les corps de nombres, *Invent. Math.* **124** (1996), no. 1-3, 437–449.
- [MS94] P. Morton and J. H. Silverman, Rational periodic points of rational functions, *Internat. Math. Res. Notices* 1994, no. 2, 97–110.
- [Mum74] D. Mumford, *Abelian varieties*, With appendices by C. P. Ramanujam and Yuri Manin, Corrected reprint of the second (1974) edition.
- [Nak08] N. Nakayama, On complex normal projective surfaces admitting non-isomorphic surjective endomorphisms, preprint September 2008.
- [NZ09] N. Nakayama and D.-Q. Zhang, Building blocks of étale endomorphisms of complex projective manifolds, *Proc. Lond. Math. Soc.* (3) **99** (2009), no. 3, 725–756.
- [NZ10] N. Nakayama and D.-Q. Zhang, Polarized endomorphisms of complex normal varieties, *Math. Ann.* **346** (2010), no. 4, 991–1018.
- [San16] K. Sano, Dynamical degree and arithmetic degree of endomorphisms on product varieties, arXiv:**1604.04174**
- [San17] K. Sano, The canonical heights for Jordan blocks of small eigenvalues, preperiodic points, and the arithmetic degrees, arXiv:**1712.07533**

- [Shi19] T. Shibata, Ample canonical heights for endomorphisms on projective varieties, *J. Math. Soc. Japan* **71** (2019), no. 2, 599–634.
- [Sil17] J. H. Silverman, Arithmetic and dynamical degrees on abelian varieties, *J. Théor. Nombres Bordeaux* **29** (2017), no. 1, 151–167.
- [Tru15] T. T. Truong, (Relative) dynamical degrees of rational maps over an algebraic closed field, *arXiv:1501.01523*
- [Uen] K. Ueno, *Classification theory of algebraic varieties and compact complex spaces*, Lecture Notes in Mathematics, Vol. **439**, Springer-Verlag, Berlin, 1975.
- [War90] J. Wahl, A characteristic number for links of surface singularities, *J. Amer. Math. Soc.* **3** (1990), no. 3, 625–637.

DEPARTMENT OF MATHEMATICS, BOX 1917, BROWN UNIVERSITY, RHODE ISLAND 02912, USA
E-mail address: matsuzawa@math.brown.edu

KOREAN INSTITUTE FOR ADVANCED STUDY, HOEGIRO 87, SEOUL 02455, KOREA
E-mail address: ms@u.nus.edu, shengmeng@kias.re.kr

NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119076, REPUBLIC OF SINGAPORE
E-mail address: mattash@nus.edu.sg

NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119076, REPUBLIC OF SINGAPORE
E-mail address: matzdz@nus.edu.sg