

Residual finiteness for central pushouts

Alexandru Chirvasitu

Abstract

We prove that pushouts $A *_C B$ of residually finite-dimensional (RFD) C^* -algebras over central subalgebras are always residually finite-dimensional provided the fibers A_p and B_p , $p \in \text{spec } C$ are RFD, recovering and generalizing results by Korchagin and Courtney-Shulman. This then allows us to prove that certain central pushouts of amenable groups have RFD group C^* -algebras. Along the way, we discuss the problem of when, given a central group embedding $H \leq G$, the resulting C^* -algebra morphism is a continuous field: this is always the case for amenable G but not in general.

Key words: C^* -algebra, amenable group, pushout, residually finite, residually finite-dimensional, Fell topology

MSC 2010: 46L09; 20E26; 22D10; 18A30

Introduction

Residual finiteness properties have elicited considerable interest, both in the operator algebra literature and in group theory. On the operator-algebraic side one typically considers *residually finite-dimensional* (henceforth RFD) C^* -algebras, i.e. those whose elements are separated by representations on finite-dimensional Hilbert spaces. The group-theoretic analogue is the concept of a *residually finite* (or RF) group, i.e. one whose finite-index normal subgroups intersect trivially (i.e. having “enough” finite quotients).

The literature on RFD C^* -algebras is rather substantial, as is that on RF groups; so much so, in fact, that it would be impossible to do it justice. For samplings (the best a short note such as this one can do) we direct the reader to, say, [15, 2, 11, 3, 18, 20, 21, 14, 25] (for RFD C^* -algebras) and [5, 4, 6] or [22, §6.5], [24, Chapters 6, 14, 15], [13, Chapter 2] (for RF groups), and references therein.

We are concerned here with the types of “permanence” properties for residual finiteness as studied, say, in [11, 3, 20, 21, 14, 25, 5], to the effect that various types of pushouts (also known as amalgamated free products) of RFD or RF objects are again such. Specifically, the main result of [20] is that pushouts of separable commutative C^* -algebras are RFD. More generally, the main theorem of [14] proves this for central pushouts $A *_C B$ with A and B separable and *strongly* RFD, i.e. such that all of their quotients are RFD.

The present note is motivated in part by the desire to recover these results without the separability and strong RFD-ness assumptions.

The preliminary Section 1 recalls some background and sets conventions.

In the short Section 2 we prove Theorem 2.1, stating that central pushouts of RFD C^* -algebras with RFD fibers are RFD.

Section 3 is devoted to the problem of when (or whether), given a central group embedding $H \leq G$, the resulting embedding $C^*(H) \rightarrow C^*(G)$ is a continuous field over $\text{spec } C^*(H)$ in the

sense of Definition 1.1 below. This holds for amenable groups (Theorem 3.2), but not, for instance, for property-(T) groups (Example 3.4).

Section 4 centers around the variant of [25, Theorem 6.9] obtained in Theorem 4.1. The latter proves that (the full group C^* -algebra of) $G_1 *_H G_2$ is RFD provided G_i are amenable residually finite (RF) and $H \leq G_i$ is a common central subgroup such that the quotients G_i/H are RF. The former result, on the other hand, assumes that $G_1 *_H G_2$ itself is RF.

Although Theorem 4.1 is formally stronger for that reason, we nevertheless show in Theorem 4.6 that its hypotheses imply the residual finiteness of the pushout $G_1 *_H G_2$. [25, Theorem 6.9] and Theorem 4.1 are thus equivalent, albeit non-obviously.

Acknowledgements

This work was partially supported by NSF grants DMS-1801011 and DMS-2001128.

I am grateful for much valuable input from Tatiana Shulman, Manuel L. Reyes and Amaury Freslon, as well as the anonymous referee's insightful suggestions.

1 Preliminaries

1.1 Fields of C^* -algebras

All C^* -algebras and pushouts are assumed unital. Pushouts over \mathbb{C} are undecorated, i.e. $A * B$ denotes what in the literature (e.g. [15]) is sometimes referred to as $A *_\mathbb{C} B$. Given a C^* morphism $C \rightarrow A$ with $C = C(X)$ commutative, we denote by A_p , the *fiber* of A at the point p of the spectrum X of C :

$$A_p := A / \langle \ker p \rangle,$$

where $p \in X$ is regarded as a character $p : C(X) \rightarrow \mathbb{C}$ and angled brackets denote the ideal generated by the respective set. Similarly, for $a \in A$ we denote by a_p its image through the surjection $A \rightarrow A_p$.

We will refer to a central C^* morphism $C \rightarrow A$ as a C -algebra A (e.g. [19, Definition 1.5] or [9, Définition 2.6]). Following standard practice (see [9, Définition 3.1] for instance), we have

Definition 1.1 A C -algebra A constitutes a *continuous field of C^* -algebras* if for every A the norm function

$$\text{spec } C \ni p \mapsto \|a\|_{A_p}$$

is continuous. ♦

The function is known to always be *upper* semicontinuous, for example by [23, Proposition 1.2].

1.2 Group filtrations

Following [5, §2.2], we need the following notion applicable to a residually finite group G .

Definition 1.2 A *filtration* on G is a family of finite-index normal subgroups $N_\alpha \trianglelefteq G$ with trivial intersection.

Let $H \leq G$ be a subgroup. A filtration $\{N_\alpha\}$ of G is an *H -filtration* provided

$$\bigcap_{\alpha} HN_\alpha = H. \quad \diamond$$

2 Pushouts over central subalgebras

The main result of this section is

Theorem 2.1 *Let A and B be two C^* -algebras, $C \subseteq A, B$ central C^* -embeddings, and assume all corresponding fibers A_p and B_p are RFD for*

$$p \in X := \text{spec } C.$$

*Then, $M := A *_C B$ is RFD.*

Proof Because C is central in A and B it is central in M . Note the isomorphism

$$M_p \cong A_p *_C B_p \cong A_p * B_p.$$

For $m \in M$ we have

$$\|m\| = \sup_{p \in X} \|m_p\|_{M_p}. \quad (2-1)$$

by [9, Proposition 2.8] and hence we can approximate the norm of m arbitrarily well with the norms of its images through representations of $M_p \cong A_p * B_p$ as p ranges over X . By the RFD-ness assumption on A_p and B_p , their coproduct $A_p * B_p$ is again RFD by [15, Theorem 3.2]. This finishes the proof. \blacksquare

Remark 2.2 Note that in fact, in the proof above one does not need the precise norm estimate (2-1): all that is needed is that every $0 \neq m \in M$ be non-zero in some quotient $M \rightarrow M_p, p \in X$. \blacklozenge

Remark 2.3 The conclusion of Theorem 2.1 cannot hold without the RFD-fiber assumption, as shown in [25, Proposition 8.3]. That result relies on the construction in [1] of an amenable RF group Γ with a central subgroup $N < \Gamma$ such that Γ/N is not RF, and in that context proves that the pushout $C^*(\Gamma) *_C C^*(\Gamma)$ is not RFD.

We can see directly, in this case, that the fiber

$$C^*(\Gamma)_p \cong C^*(\Gamma/N)$$

at the point $p \in \text{spec } C^*(N)$ corresponding to the trivial morphism $N \rightarrow \{1\}$ is not RFD. Indeed, since Γ/N is finitely generated (finitely presented, even) by [1], the RFD-ness of $C^*(\Gamma/N)$ would entail the residual finiteness of Γ/N (e.g. by [25, Proposition 2.4]), yielding a contradiction. \blacklozenge

3 Fibers over dense sets

It will be useful, in working with group C^* -algebras, to allow the points $p \in X$ from Theorem 2.1 to range only over a dense subset $Y \subseteq X := \text{spec } C$ rather than the entire spectrum. We cannot do this in full generality, as (2-1) does not hold for the supremum over only a dense subset $Y \subseteq X$:

Example 3.1 Let $I = [0, 1]$, $M = C_b(I)$ (all bounded functions) and $C = C(I) \subset M$ (continuous functions), both equipped with the supremum norm. The indicator function m of $\{0\} \subset I$ is non-zero, even though its image in every fiber

$$M_p, p \in Y := (0, 1] \subset I$$

vanishes. For that reason, we do *not* have

$$\|m\| = \sup_{p \in Y} \|m_p\|_{M_p}. \quad \blacklozenge$$

What went wrong in the above example is that M was not a continuous field over C in the sense of Definition 1.1. One might hope that the pathology in Example 3.1 is at least avoidable for group C^* -algebras; this is not the case, as Example 3.4 below shows. Nevertheless, for amenable groups one can do better (see also [10, Theorem 3.7] for a broader discussion of the continuity of pushout fields).

Theorem 3.2 *Let G_i , $i = 1, 2$ be two amenable groups and $H \leq G_i$ a common central subgroup. Then,*

$$C^*(G_1 *_H G_2) \cong C^*(G_1) *_{C^*(H)} C^*(G_2)$$

is a continuous field of C^ -algebras over $\widehat{H} = \text{spec } C^*(H)$ in the sense of Definition 1.1.*

Proof Since, as noted in §1.1, for $C := C^*(H)$ the map

$$C^*(G_1) *_C C^*(G_2) =: M \ni m \mapsto \|m_p\|_{M_p}$$

is always upper semicontinuous, it remains to prove that for every dense subset

$$Y \subseteq \widehat{H}$$

of the spectrum of and every $m \in M$ we have

$$\|m\| = \sup_{p \in Y} \|m_p\|_{M_p}.$$

First, fix an arbitrary unitary representation ρ_2 of G_2 where H acts by scalars, with central character $p_0 \in \widehat{H}$. Let

$$Y \ni p_\lambda \rightarrow p_0$$

be a net of characters from Y converging to p_0 and set

$$q_\lambda := p_\lambda p_0^{-1} \rightarrow 1 \in \widehat{H}$$

be the corresponding ‘‘error’’.

Because

$$G_i/H, \quad i = 1, 2$$

are amenable homogeneous spaces of G_i respectively in the sense of [8, §2], the trivial representation $\mathbf{1}_{G_2}$ is weakly contained in the induced representation $\text{Ind}_H^{G_2} \mathbf{1}_H$. Since $q_\lambda \rightarrow \mathbf{1}_H$ and induction is continuous [7, Theorem F.3.5], we have

$$\text{Ind}_H^{G_2} q_\lambda \rightarrow \text{Ind}_H^{G_2} \mathbf{1}_H \rightarrow \mathbf{1}_{G_2}$$

in the Fell topology, and hence

$$\rho_2 \otimes \text{Ind}_H^{G_2} q_\lambda \rightarrow \rho_2 \otimes \mathbf{1}_{G_2} \cong \rho_2. \quad (3-1)$$

Because H acts by p_0 in ρ_2 , it acts via $p_\lambda = q_\lambda p_0$ in the left hand side of (3-1). In short, we can Fell-approximate ρ_2 with unitary representations where H acts by characters from Y .

Now fix $m \in M$. According to [9, Proposition 2.8], the norm of m is achieved in some unitary representation ρ of $G := G_1 *_H G_2$ where H acts by scalars, with central character $p_0 \in \widehat{H}$. Working only with non-degenerate (indeed, unital) representations isomorphic to their own \aleph_0 -multiples, [16, Lemma 2.4] implies that it will suffice to approximate ρ arbitrarily well in the Fell topology [7, Appendix F.2] with unitary representations where H operates with central characters belonging to the dense subset $Y \subseteq \widehat{H}$.

In turn, in order to achieve the above it is enough to approximate the restrictions $\rho|_{G_i}$ with representations where H acts via elements of Y . This, however, is what the first part of the proof does. ■

Over the course of the proof of [Theorem 3.2](#) we have obtained

Proposition 3.3 *Let $H \leq G$ be a central subgroup of an amenable group. Then, for every dense subset*

$$Y \subseteq \widehat{H} = \text{spec } C^*(H)$$

the canonical map

$$C^*(G) \rightarrow \prod_{p \in Y} C^*(G)_p \tag{3-2}$$

is one-to-one. ■

In other words, a variant of [[9](#), Proposition 2.8] with only a *dense* subset of the spectrum rather than the entirety of it.

To prepare the ground for [Example 3.4](#), note that by [[7](#), Theorem F.4.4] (3-2) is an embedding if and only if the G -representations where H acts by characters in Y form a dense set in the Fell topology. This cannot possibly happen if

- G has the Kazhdan property (T) (e.g. [[7](#), §1.1]), and hence the trivial representation is isolated in the unitary dual;
- the dense subset $Y \subset \widehat{H}$ does not contain the trivial element.

To construct such examples all we need is a property-(T) group G with an *infinite* central subgroup H , whereupon we can simply take

$$Y = \widehat{H} \setminus \{1\}.$$

Example 3.4 Let G be the universal cover $\widetilde{Sp}_4(\mathbb{R})$ of the 4×4 real symplectic group. It is shown in [[17](#), Theorem 6.8] that G has property (T) (indeed, even the stronger property (T*); note that the authors of that paper denote Sp_4 by Sp_2).

The fundamental group of $Sp_4(\mathbb{R})$ is \mathbb{Z} , so we can simply take that copy of \mathbb{Z} as the infinite central subgroup $H < G$.

Finally, if a *discrete* group is desired then one can simply take the preimage through

$$\widetilde{Sp}_4(\mathbb{R}) \rightarrow Sp_4(\mathbb{R})$$

of any lattice in the latter (it will again have property (T) by [[7](#), Theorem 1.7.1]). See also [[7](#), §1.7] for a discussion of property (T) permanence under passage to universal covering groups. ◆

4 Central pushouts of RF groups

The present section attempts to prove a slightly more general version of [[25](#), Theorem 6.9]. The difference is that we only assume that G_i are individually RF rather than assuming that $G_1 *_H G_2$ is.

Theorem 4.1 *Let G_i , $i = 1, 2$ be two amenable groups and $H \leq G_i$ a common central subgroup such that*

- *each G_i is RF;*
- *each G_i/H is RF.*

Then, $G_1 *_H G_2$ is RFD.

Proof We will obtain the result as an application of Theorem 3.2, with

$$A = C^*(G_1), \quad B = C^*(G_2) \text{ and } C = C^*(H).$$

According to that result, what we have to argue is that the fibers A_p and B_p are RFD for a dense set of points p in the spectrum

$$\widehat{H} = \text{Pontryagin dual of } H = \text{spec } C^*(H).$$

This is precisely what Corollary 4.5 does, finishing the proof. ■

Lemma 4.2 *Let $H \leq G$ be a central inclusion such that G and G/H are both RF. Then, the characters $p : H \rightarrow \mathbb{S}^1$ whose kernels*

$$N = \ker p$$

give rise to RF quotients G/N form a dense subset of the Pontryagin dual \widehat{H} .

Proof Since G is RF the normal finite-index subgroups $G_\alpha \trianglelefteq G$ have trivial intersection. Additionally, since G/H is RF, G has an H filtration in the sense of Definition 1.2 (e.g. [25, Proposition 3.3]) and hence

$$\bigcap_{\alpha} HG_{\alpha} = H. \tag{4-1}$$

Since

$$H_{\alpha} := G_{\alpha} \cap H \leq H$$

have trivial intersection, the union of the duals

$$\widehat{H/H_{\alpha}} \subseteq \widehat{H}$$

is a dense subgroup. We claim that any $p \in \widehat{H/H_{\alpha}}$ will meet the requirements in the statement; proving this will achieve the desired conclusion, so it is the task we turn to next.

Fix such a character $p : H \rightarrow \mathbb{S}^1$, factoring through some

$$H \rightarrow H/H_{\alpha_0},$$

and let $N \leq H$ be its kernel. Then, by its very construction, N will contain $H_{\alpha_0} = H \cap G_{\alpha_0}$. This means that

$$NG_{\alpha_0} \cap H \subseteq N$$

(which is then an equality), and hence the intersection

$$\bigcap_{\alpha} NG_{\alpha} \subseteq H$$

(where the latter inclusion uses (4-1)) cannot possibly contain N *strictly*. In conclusion we have

$$\bigcap_{\alpha} NG_{\alpha} = N,$$

meaning that the filtration $\{G_{\alpha}\}$ is compatible with N (i.e. an N -filtration) and hence G/N is RF. ■

Remark 4.3 The RF requirements in Theorem 4.1 are both crucial:

On the one hand, [25, §8] recalls the example given by Abels in [1] of a central inclusion $\mathbb{Z} < G$ into an RF amenable group such that G/\mathbb{Z} is not RF.

On the other hand, [12] contains an example (attributed there to C. Kanta Gupta) of a non-residually finite amenable group G whose quotient by an order-two normal subgroup $K \trianglelefteq G$ is residually finite. Centrality is easy to arrange, since the centralizer of K in G will have finite index in the latter. \blacklozenge

Although Lemma 4.2 ensures that the quotients G/N for

$$N = \ker(p : H \rightarrow \mathbb{S}^1)$$

are RF, we have yet to prove that the resulting fiber algebras $C^*(G)/\langle h - p(h), h \in H \rangle$ are RFD.

Lemma 4.4 *Let $H \leq G$ be a central subgroup, $p \in \widehat{H}$ a character of finite order, and $N = \ker p$ its kernel in H . If G/N is RFD then so is the fiber C^* -algebra*

$$C^*(G)_p := C^*(G)/\langle h - p(h), h \in H \rangle \tag{4-2}$$

corresponding to p is RFD.

Proof Note that $C^*(G)_p$ is precisely the fiber of $C^*(G/N)$ at the character \bar{p} induced by p on the quotient (finite cyclic) group H/N : indeed, denoting images of elements $g \in G$ in G/N by \bar{g} , we have

$$C^*(G)_p = C^*(G)/\langle h - p(h), h \in H \rangle = C^*(G/N)/\langle \bar{h} - \bar{p}(\bar{h}), \bar{h} \in H/N \rangle = C^*(G/N)_{\bar{p}}$$

because the kernel of $C^*(G) \rightarrow C^*(G/N)$ is generated by $h - 1$, $h \in N$, which are already contained in the ideal we are modding out in (4-2).

For that reason, we may as well assume that

- N is trivial, and hence
- H is finite cyclic;
- G is RFD.

But now note that $C^*(G)_p$ is a fiber of the RFD C^* -algebra over the *finite-dimensional* central subalgebra $C^*(H) \leq C^*(G)$. In general, a C^* -algebra A will break up as a product of the fibers A_p over a finite-dimensional central C^* -subalgebra $C \leq A$, by simply cutting A with the minimal projections in C .

In particular, under our assumptions the fiber $C^*(G)_p$ is a Cartesian factor of $C^*(G)$, and hence the RFD-ness of the latter entails that of the former. \blacksquare

Corollary 4.5 *Under the hypotheses of Lemma 4.2 the characters $p : H \rightarrow \mathbb{S}^1$ for which the fiber (4-2) is RFD form a dense subset of \widehat{H} .*

Proof Indeed, the characters p in the proof of Lemma 4.2 are of finite order, and hence Lemma 4.4 applies. \blacksquare

4.1 Recovering residual finiteness for a pushout

Recall that [25, Theorem 6.9] assumes $G_1 *_C G_2$ is RF, whereas Theorem 4.1 only requires that G_i , $i = 1, 2$ be RF individually (along with G_i/H). We argue here that that distinction is only apparent:

Theorem 4.6 *Let $H \leq G_i$, $i = 1, 2$ be a common central subgroup such that G_i and G_i/H are all RF. Then, $G_1 *_H G_2$ is RF.*

Proof Note first that the case $G_1 = G_2$ is clear: indeed, the assumptions that G and G/H are RF then show that G has an H -filtration. The two copies of that filtration in the two copies of G are then (H, H, id) -compatible in the sense of [5, §2.2], and hence $G *_H G$ is RF by [5, Proposition 2].

It thus remains to reduce the problem to the case $G_1 = G_2$. To do this, consider the *tensor product*

$$G := G_1 \otimes_H G_2$$

defined by identifying the two copies of H in $G_1 \times G_2$; in other words, $G_1 \otimes_H G_2$ is $G_1 *_H G_2$ modulo the relations making the elements of

$$G_1 \text{ and } G_2 \leq G_1 *_H G_2$$

commute.

We then have

$$G/H \cong (G_1/H) \times (G_2/H),$$

which is thus RF by assumption. On the other hand, if we show that $G *_H G$ itself is RF then so is

$$G_1 *_H G_2 \leq G *_H G$$

(where the inclusions $G_i \leq G = G_1 \otimes_H G_2$ are the obvious ones).

To summarize, we have thus far

- observed that the conclusion holds when $G_1 = G_2$ (equal to a common group G , say);
- reduced the problem to its instance for $G = G_1 \otimes_H G_2$, modulo
- the desired hypothesis that that G is RF.

In conclusion, all that remains to be proven is that under our hypotheses $G := G_1 \otimes_H G_2$ is indeed RF; we relegate this to Lemma 4.7. ■

Lemma 4.7 *Let $H \leq G_i$, $i = 1, 2$ be a common central subgroup such that G_i and G_i/H are all RF. Then, $g := G_1 \otimes_H G_2$ is RF.*

Proof If $\{G_{i,\alpha}\}$ are H -filtrations of G_i respectively for $i = 1, 2$ then the images G_α of

$$G_{1,\alpha} \times G_{2,\alpha} \leq G_1 \times G_2$$

through the surjection

$$G_1 \times G_2 \rightarrow G = G_1 \otimes_H G_2$$

identifying the two copies of H will form an H -filtration for G . ■

References

- [1] Herbert Abels. An example of a finitely presented solvable group. In *Homological group theory (Proc. Sympos., Durham, 1977)*, volume 36 of *London Math. Soc. Lecture Note Ser.*, pages 205–211. Cambridge Univ. Press, Cambridge-New York, 1979.
- [2] R. J. Archbold. On residually finite-dimensional C^* -algebras. *Proc. Amer. Math. Soc.*, 123(9):2935–2937, 1995.
- [3] Scott Armstrong, Ken Dykema, Ruy Exel, and Hanfeng Li. On embeddings of full amalgamated free product C^* -algebras. *Proc. Amer. Math. Soc.*, 132(7):2019–2030, 2004.
- [4] Benjamin Baumslag and Marvin Tretkoff. Residually finite HNN extensions. *Comm. Algebra*, 6(2):179–194, 1978.
- [5] Gilbert Baumslag. On the residual finiteness of generalised free products of nilpotent groups. *Trans. Amer. Math. Soc.*, 106:193–209, 1963.
- [6] Gilbert Baumslag. Finitely generated cyclic extensions of free groups are residually finite. *Bull. Austral. Math. Soc.*, 5:87–94, 1971.
- [7] Bachir Bekka, Pierre de la Harpe, and Alain Valette. *Kazhdan’s property (T)*, volume 11 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2008.
- [8] Mohammed E. B. Bekka. Amenable unitary representations of locally compact groups. *Invent. Math.*, 100(2):383–401, 1990.
- [9] Étienne Blanchard. Déformations de C^* -algèbres de Hopf. *Bull. Soc. Math. France*, 124(1):141–215, 1996.
- [10] Etienne Blanchard. Amalgamated free products of C^* -bundles. *Proc. Edinb. Math. Soc. (2)*, 52(1):23–36, 2009.
- [11] Nathaniel P. Brown and Kenneth J. Dykema. Popa algebras in free group factors. *J. Reine Angew. Math.*, 573:157–180, 2004.
- [12] Robert Campbell. http://www.math.umbc.edu/~campbell/CombGpThy/RF_Thesis/2_RF_Results.html. Accessed: 2020-01-21.
- [13] Tullio Ceccherini-Silberstein and Michel Coornaert. *Cellular automata and groups*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2010.
- [14] Kristin Courtney and Tatiana Shulman. Free products with amalgamation over central C^* -subalgebras. *Proc. Amer. Math. Soc.*, 148(2):765–776, 2020.
- [15] Ruy Exel and Terry A. Loring. Finite-dimensional representations of free product C^* -algebras. *Internat. J. Math.*, 3(4):469–476, 1992.
- [16] J. M. G. Fell. Weak containment and induced representations of groups. *Canadian J. Math.*, 14:237–268, 1962.
- [17] Uffe Haagerup, Søren Knudby, and Tim de Laat. A complete characterization of connected Lie groups with the approximation property. *Ann. Sci. Éc. Norm. Supér. (4)*, 49(4):927–946, 2016.

- [18] Don Hadwin. A lifting characterization of RFD C^* -algebras. *Math. Scand.*, 115(1):85–95, 2014.
- [19] G. G. Kasparov. Equivariant KK -theory and the Novikov conjecture. *Invent. Math.*, 91(1):147–201, 1988.
- [20] Anton Korchagin. Amalgamated free products of commutative C^* -algebras are residually finite-dimensional. *J. Operator Theory*, 71(2):507–515, 2014.
- [21] Qihui Li and Junhao Shen. A note on unital full amalgamated free products of RFD C^* -algebras. *Illinois J. Math.*, 56(2):647–659, 2012.
- [22] Wilhelm Magnus, Abraham Karrass, and Donald Solitar. *Combinatorial group theory*. Dover Publications, Inc., Mineola, NY, second edition, 2004. Presentations of groups in terms of generators and relations.
- [23] Marc A. Rieffel. Continuous fields of C^* -algebras coming from group cocycles and actions. *Math. Ann.*, 283(4):631–643, 1989.
- [24] Derek J. S. Robinson. *A course in the theory of groups*, volume 80 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1996.
- [25] Tatiana Shulman. Central amalgamation of groups and the RFD property, 2019. arXiv:2001.00052.

DEPARTMENT OF MATHEMATICS, UNIVERSITY AT BUFFALO, BUFFALO, NY 14260-2900, USA
E-mail address: `achirvas@buffalo.edu`