

# A STRATIFICATION RESULT FOR AN EXPONENTIAL SUM MODULO $p^2$

KOSTADINKA LAPKOVA AND STANLEY YAO XIAO

ABSTRACT. In this note we consider algebraic exponential sums over the values of homogeneous nonsingular polynomials  $F(x_1, \dots, x_n) \in \mathbb{Z}[x_1, \dots, x_n]$  in the quotient ring  $\mathbb{Z}/p^2\mathbb{Z}$ . We provide an estimate of this exponential sum and a corresponding stratification of the space  $\mathbb{A}_{\mathbb{F}_p}^n$ , which in particular illustrates a general stratification theorem of Fouvry and Katz.

## 1. INTRODUCTION

Let  $F(\mathbf{x}) \in \mathbb{Z}[x_1, \dots, x_n]$  be a polynomial of degree  $d \geq 1$  and let  $V_F$  be the hypersurface defined by  $F(\mathbf{x}) = 0$ . Denote by  $K$  a finite field or a quotient ring of size  $q$ . We are interested in the exponential sums

$$(1.1) \quad \mathcal{S}_F(\mathbf{h}; K) = \sum_{\mathbf{x} \in V_F(K)} \exp\left(\frac{h_1x_1 + \dots + h_nx_n}{q}\right),$$

where we use the notation  $\exp(x) := e^{2\pi ix}$ . In this note we address the estimate of  $\mathcal{S}_F(\mathbf{h}, \mathbb{Z}/p^2\mathbb{Z})$  for large primes  $p$ .

A fundamental work in the estimation of general algebraic exponential sums is the paper of Fouvry and Katz [3]. When we drop a multiplicative character factor from the sums considered in [3], and take not a general nontrivial additive character but only  $\exp(\text{mod } q)$ , we arrive at sums similar to  $\mathcal{S}_F(\mathbf{h}; \mathbb{F}_p)$ , thus our result would be a special case of the sums considered in [3], but with the modulus being squares of primes rather than primes. An analogous algebraic exponential sum was treated by Fouvry and our main theorem resembles [2, Proposition 1.0] in the special case when the subscheme considered is a smooth hypersurface.

One significant aspect of the results of Fouvry and Fouvry-Katz [2, 3] is that they establish the existence of a decreasing filtration of subschemes (stratification)  $\mathbb{A}_{\mathbb{Z}}^n \supset \mathcal{G}_1 \supset \mathcal{G}_2 \supset \dots \supset \mathcal{G}_n$  of codimension  $\geq j$ , such that whenever  $\mathbf{h} \notin \mathcal{G}_j(\mathbb{F}_p)$  we have

$$(1.2) \quad \mathcal{S}_F(\mathbf{h}, \mathbb{F}_p) \ll p^{(n-1)/2+(j-1)/2}.$$

Thus the deviation from the bound  $p^{(n-1)/2}$ , which amounts to square-root cancellation and is the best that can be expected in general, is tamed by the fact that the dimensions of the subschemes  $\mathcal{G}_j$  for  $j \geq 1$  are small. However, the construction of this stratification is ineffective and inaccessible for non-specialists. Furthermore, the verifying of certain criteria that allow to apply the theorems of Fouvry-Katz [3] constitutes a separate problem, see for example [1]. Also see [14] for a certain generalization

---

*Date:* October 10, 2021.

of [3]. A further overview of the topic is given in the excellent review by P. Michel [11].

On the other hand, Stepanov [12] focuses not on the stratification but rather constructs a subset of the good locus  $\bar{\mathcal{G}}_1$  in which the optimal bound  $\mathcal{S}_F(\mathbf{h}, \mathbb{F}_q) \ll q^{(n-1)/2+\epsilon}$  holds for any finite field and not only for fields of prime order. Heath-Brown [4] treats the special case  $q = p^2$  and  $F \equiv \mathbf{1}$ .

We show that in the specific case considered in (1.1) the expected estimate holds and we give an explicitly constructed stratification.

For a given vector  $\mathbf{h} \in \mathbb{Z}^n$  consider the auxiliary affine variety  $W_{F,\mathbf{h}}$  given by the equations defined over  $\mathbb{F}_p$

$$(1.3) \quad \begin{aligned} F(\mathbf{y}) &= 0 \\ h_i \frac{\partial F}{\partial x_j}(\mathbf{y}) &= h_j \frac{\partial F}{\partial x_i}(\mathbf{y}), \quad 1 \leq i, j \leq n, i \neq j. \end{aligned}$$

Note that this is the singular locus of the variety  $\{F(\mathbf{y}) = 0, \mathbf{h} \cdot \mathbf{y} = 0\}$ . Now put  $\mathcal{G}_{F,j}$  for the set given by

$$(1.4) \quad \mathcal{G}_{F,j} = \{\mathbf{h} \in \mathbb{F}_p^n : \dim(W_{F,\mathbf{h}}) \geq j\}.$$

We have the following theorem, which can be viewed as an extension to Theorem 1.2 of Fouvry-Katz [3] to  $V(\mathbb{Z}/p^2\mathbb{Z})$  in a restricted setting.

**Theorem 1.1.** *Let  $F$  be a non-singular homogeneous polynomial in  $\mathbb{Z}[x_1, \dots, x_n]$ . Let  $V_F$  be the hypersurface defined by  $F$ . Further, suppose that  $F$  is non-singular modulo  $p$ . For  $\mathbf{h} \in \mathbb{Z}^n$  let  $S(\mathbf{h}; p^2)$  be the exponential sum*

$$(1.5) \quad S(\mathbf{h}; p^2) = \sum_{\mathbf{x} \in V_F(\mathbb{Z}/p^2\mathbb{Z})} \exp\left(\frac{h_1 x_1 + \dots + h_n x_n}{p^2}\right).$$

*Then the sets defined in (1.4) are varieties satisfying  $\mathcal{G}_{F,0} = \mathbb{A}_{\mathbb{F}_p}^n \supseteq \mathcal{G}_{F,1} \supseteq \dots \supseteq \mathcal{G}_{F,n}$ , where the codimension of each  $\mathcal{G}_{F,j}$  is  $\geq j$ , and we have the estimate*

$$S(\mathbf{h}; p^2) = O_F(p^{n+j-2})$$

*whenever  $\mathbf{h} \notin \mathcal{G}_{F,j}(\mathbb{F}_p)$ .*

The interest in this theorem is that unlike in the situation considered by Fouvry and Katz, a priori we have relatively little access to tools from algebraic geometry. This is because the base ring we consider is  $\mathbb{Z}/p^2\mathbb{Z}$  which is not a field, so many of the basic facts one takes for granted from algebraic geometry do not necessarily hold. Nevertheless, we are able to reduce to a case where we once again consider things over a field.

When  $n = 2$  and  $F$  is non-singular of degree  $d \geq 2$ , we have  $\dim W_{F,\mathbf{h}} = 1$  if and only if the equation  $h_2 \frac{\partial F}{\partial x_1}(\mathbf{y}) = h_1 \frac{\partial F}{\partial x_2}(\mathbf{y})$  is redundant. If  $h_1, h_2$  are not both zero then this implies that  $F$  shares a component with a curve of strictly smaller degree, which violates the assumption that  $F$  is non-singular. Therefore the only way to

have  $\dim W_{F, \mathbf{h}} = 1$  is for  $h_1 = h_2 = 0$  and so any non-singular  $F \in \mathbb{Z}[x_1, x_2]$  we have square-root cancellation, and obtain the bound

$$S(\mathbf{h}; p^2) \ll p^{n-1}.$$

We will soon see in the proof that this essentially follows from the Weil bound resolving the Riemann Hypothesis for curves, where for any (higher) dimension of  $V_F$  we apply the theorem of Lang-Weil.

In many sieving problems (see for example [7]) one applies a result like that of Fouvry-Katz [3] to use congruence conditions modulo somewhat large primes to show paucity of contributions, which results in sums of the shape

$$\sum_{x_1 < p < x_2} \left( \frac{B^n}{p} + 1 \right).$$

Notice that the sum over the lead terms  $1/p$  diverges, so in such a case the size of  $x_1$  is not relevant. On the other hand, if we apply our Theorem instead, then one would end up with a sum over  $1/p^2$  which is then convergent, so the sum would then be sensitive to the size of  $x_1$ . This is crucial if one wishes to obtain asymptotic formulae. Similar arguments could lead to some of the potential applications of our theorem.

## 2. PROOF OF THEOREM 1.1

Write  $\mathbf{x} = \mathbf{y} + p\mathbf{z}$ . Then we have

$$(2.1) \quad F(\mathbf{x}) = F(\mathbf{y}) + p \left( \sum_{i=1}^n z_i \frac{\partial F}{\partial x_i}(\mathbf{y}) \right) + p^2 H_F(\mathbf{y}; \mathbf{z}) \equiv F(\mathbf{y}) + p\mathbf{z} \cdot \nabla F(\mathbf{y}) \pmod{p^2}.$$

Now restrict  $\mathbf{y} \in \mathbb{F}_p$ . Then  $\mathbf{y} + p\mathbf{z} \in V(\mathbb{Z}/p^2\mathbb{Z})$  if and only if  $\mathbf{y} \in V(\mathbb{F}_p)$  and

$$\mathbf{z} \cdot \nabla F(\mathbf{y}) \equiv -F(\mathbf{y})/p \pmod{p}.$$

Then (1.5) becomes

$$\sum_{\mathbf{y} \in V(\mathbb{F}_p)} \exp \left( \frac{h_1 y_1 + \cdots + h_n y_n}{p^2} \right) \sum_{\substack{\mathbf{z} \pmod{p} \\ \mathbf{z} \cdot \nabla F(\mathbf{y}) \equiv -F(\mathbf{y})/p \pmod{p}}} \exp \left( \frac{h_1 z_1 + \cdots + h_n z_n}{p} \right).$$

We examine the inner sum in detail. Let us introduce the notation

$$T_p(\mathbf{a}, \mathbf{b}, c) := \sum_{\substack{\mathbf{z} \pmod{p} \\ \mathbf{b} \cdot \mathbf{z} \equiv c \pmod{p}}} \exp \left( \frac{\mathbf{a} \cdot \mathbf{z}}{p} \right).$$

We claim that the exponential sum  $T_p(\mathbf{a}, \mathbf{b}, c) = 0$  every time there exists a pair  $(i, j)$  with  $1 \leq i, j \leq n$  and  $i \neq j$ , such that  $a_i b_j \not\equiv a_j b_i \pmod{p}$ . Indeed, if the latter holds, we can assume that  $b_i \neq 0$ , since  $b_i = b_j = 0$  contradicts the assumption. We can assume that  $i = n$ . We write

$$z_n \equiv -b_n^{-1} (b_1 z_1 + \cdots + b_{n-1} z_{n-1} - c) \pmod{p}$$

and then  $T_p(\mathbf{a}, \mathbf{b}, c)$  equals

$$\sum_{z_1, \dots, z_{n-1} \pmod{p}} \exp\left(\frac{a_1 z_1 + \dots + a_{n-1} z_{n-1} - a_n b_n^{-1} (b_1 z_1 + \dots + b_{n-1} z_{n-1} - c)}{p}\right) = \\ \sum_{z_1, \dots, z_{n-1} \pmod{p}} \exp\left(\frac{\sum_{1 \leq j \leq n-1} (a_j - a_n b_n^{-1} b_j) z_j + a_n b_n^{-1} c}{p}\right).$$

As we assumed that for some  $j \in [1, n-1]$  we have  $a_j - a_n b_n^{-1} b_j \not\equiv 0 \pmod{p}$ , it follows that  $T_p(\mathbf{a}, \mathbf{b}, c) = 0$ . On the other hand, if  $a_i b_j \equiv a_j b_i \pmod{p}$  for every  $1 \leq i, j \leq n$  and  $i \neq j$ , we have trivially  $|T_p(\mathbf{a}, \mathbf{b}, c)| = p^{n-1}$ .

Now we conclude that (1.5) is equal to

$$\sum_{\mathbf{y} \in V(\mathbb{F}_p)} \exp\left(\frac{\mathbf{h} \cdot \mathbf{y}}{p^2}\right) T_p(\mathbf{h}, \nabla F(\mathbf{y}), -F(\mathbf{y})/p)$$

and this is bounded from above by

$$p^{n-1} \# \left\{ \mathbf{y} \in \mathbb{F}_p^n : F(\mathbf{y}) = 0, \text{rank} \begin{pmatrix} \mathbf{h} \\ \nabla F(\mathbf{y}) \end{pmatrix} < 2 \right\} = p^{n-1} \sum_{\mathbf{y} \in W_{F, \mathbf{h}}(\mathbb{F}_p)} 1.$$

If  $\mathbf{h} \notin \mathcal{G}_{F, j}(\mathbb{F}_p)$ , then

$$\dim(W_{F, \mathbf{h}}) \leq j - 1.$$

By the Lang-Weil theorem, it follows that

$$\sum_{\mathbf{y} \in W_{F, \mathbf{h}}(\mathbb{F}_p)} 1 \ll_{d, n} p^{j-1}$$

and the estimate of the exponential sum  $S(\mathbf{h}, p^2)$  follows. It remains to deal with the dimensions of the sets  $\mathcal{G}_{F, j}$ .

The following lemma is reminiscent of Heath-Brown's Lemma 2 in [5], proved using the method of dimension counting via incidence geometry. In [5] he constructs a stratification similar to ours, but for exponential sums modulo  $p$ . The lemma will help us estimate the dimensions of  $\mathcal{G}_{F, j}$ , which turn out to be affine varieties.

**Lemma 2.1.** *Let  $F \in K[x_1, \dots, x_n]$  be a non-singular homogenous polynomial of degree  $d$  in  $n$  variables defined over a field  $K$ . For a given  $\mathbf{z} \in K^n$  let*

$$V_{\mathbf{z}} = \text{sing}(X \cap H_{\mathbf{z}}),$$

where  $X = \{F(\mathbf{x}) = 0\} \subset \mathbb{P}_K^{n-1}$  and  $H_{\mathbf{z}} = \{\mathbf{z} \cdot \mathbf{x} = 0\} \subset \mathbb{P}_K^{n-1}$ . Let

$$T_s = \{z = [\mathbf{z}] \in \mathbb{P}_K^{n-1} : \dim(V_{\mathbf{z}}) \geq s\}, \quad s \geq -1.$$

Then  $T_s$  is a projective variety of dimension at most  $n - 2 - s$ .

*Proof.* Consider the variety

$$I = \left\{ (z, \mathbf{x}) \in \mathbb{P}_K^{n-1} \times X : \text{rank} \begin{pmatrix} \mathbf{z} \\ \nabla F(\mathbf{x}) \end{pmatrix} < 2 \right\}.$$

The projection  $\pi_2 : I \rightarrow X$  is onto, since  $[\nabla F(\mathbf{x})]$  is in the fibre for every  $\mathbf{x} \in X$ . Since  $X$  is non-singular we have  $\nabla F(\mathbf{x}) \neq 0$  for every  $\mathbf{x} \in X$ , therefore we have the condition

$$\text{rank} \begin{pmatrix} \mathbf{z} \\ \nabla F(\mathbf{x}) \end{pmatrix} = 1,$$

i.e. each minor in the matrix equals 0. It is easy to see that this implies that any other  $z \in \mathbb{P}^{n-1}$  in the same fibre is such that  $\mathbf{z}$  is proportional to  $\nabla F(\mathbf{x})$ , i.e.  $[z] = [\nabla F(\mathbf{x})]$ . This means that the fibre consists of a single point. Hence  $I$  is irreducible and by the theorem on the dimension of fibres we get

$$\dim(I) = \dim(\pi_2^{-1}(\mathbf{x})) + \dim(X) = 0 + n - 2 = n - 2.$$

Let  $U$  be an irreducible component of  $T_s$  and consider the projection

$$\pi_1 : I \rightarrow U.$$

Then

$$n - 2 = \dim(I) \geq s + \dim(U),$$

which yields  $\dim(T_s) \leq n - 2 - s$ . □

We will also need the following estimate of the dimensions of  $\mathcal{G}_{F,j}$ .

**Lemma 2.2.** *For any homogenous and non-singular polynomial  $F$  and  $j \geq 0$  we have*

$$\dim(\mathcal{G}_{F,j}(\mathbb{A}_{\mathbb{F}_p}^n)) \leq n - j.$$

*Proof.* When we regard  $X$  and  $H_{\mathbf{z}}$  over  $\mathbb{A}_{\mathbb{F}_p}^n$  as affine varieties, and respectively the affine singular locus of their intersection  $\text{sing}(X \cap H_{\mathbf{z}})(\mathbb{A}_{\mathbb{F}_p}^n) = V_{\mathbf{z}}(\mathbb{A}_{\mathbb{F}_p}^n)$ , we have  $\dim(V_{\mathbf{z}}(\mathbb{A}_{\mathbb{F}_p}^n)) = \dim(V_{\mathbf{z}}) + 1$ . Thus if  $[\mathbf{z}] \in T_s$  for  $s \geq -1$ , we have  $\dim(V_{\mathbf{z}}) \geq s$  and respectively  $\dim(W_{F,\mathbf{z}}(\mathbb{A}_{\mathbb{F}_p}^n)) \geq s + 1$ , so any of the corresponding  $\mathbf{z} \in \mathbb{F}_p^n$  satisfies  $\mathbf{z} \in \mathcal{G}_{F,s+1}$ . As  $\dim(T_s) \leq n - 2 - s$  and  $T_s$  is exactly  $\mathcal{G}_{F,s+1}(\mathbb{P}_{\mathbb{F}_p}^{n-1})$ , we have  $\dim(\mathcal{G}_{F,s+1}) \leq (n - 2 - s) + 1 = n - (s + 1)$ . We conclude that for  $j \geq 0$  we have  $\dim(\mathcal{G}_{F,j}(\mathbb{A}_{\mathbb{F}_p}^n)) \leq n - j$ . □

This completes the proof of Theorem 1.1.

**Acknowledgements.** We are grateful to Prof. Tim Browning for helping with Lemma 2.1. K. Lapkova is supported by a Hertha Firnberg grant [T846-N35] of the Austrian Science Fund (FWF).

## REFERENCES

- [1] L. Devin, *On the congruence class modulo prime numbers of the number of rational points of a variety*, Int. Math. Res. Not. (14) (2018), 4450–4479
- [2] E. Fouvry, *Consequences of a result of N. Katz and G. Laumon concerning trigonometric sums*, Israel J. Math. 120 (2000), part A, 81–96
- [3] E. Fouvry, N. Katz, *A general stratification theorem for exponential sums, and applications*, J. reine angew. Math **540** (2001), 115–166.
- [4] D. R. Heath-Brown, *A multiple exponential sum to modulus  $p^2$* , Canad. Math. Bull. (4) **28** (1985), 394–396.
- [5] D. R. Heath-Brown, *The density of rational points on nonsingular hypersurfaces*, Proc. Indian Acad. Sci. Math. Sci. (1) **104** (1994), 1329.

- [6] C. Hooley, *On the power free values of polynomials*, *Mathematika* **14** (1967), 21-26.
- [7] C. Hooley, *On the power-free values of polynomials in two variables*, *Analytic Number Theory: Essays in Honour of Klaus Roth* (2009).
- [8] C. Hooley, *On the power-free values of polynomials in two variables II*, *Journal of Number Theory* **129** (2009), 1443-1455.
- [9] K. Lapkova, S. Y. Xiao, *Density of power-free values of polynomials*, *Mathematika* (4) **65** (2019), 1038-1050
- [10] K. Lapkova, S. Y. Xiao, *Density of power-free values of polynomials II*, Preprint
- [11] P. Michel, review of "A general stratification theorem for exponential sums, and applications" (in *J. reine angew. Math* **540** (2001), 115-166, by E. Fouvry, N. Katz) , *Mathematical Reviews/MathSciNet* MR1868601
- [12] S. A. Stepanov, *Rational trigonometric sums on "algebraic varieties"*, *Mat. Zametki* (2) **39**(1986), 161–174, 301
- [13] S. Y. Xiao, *Square-free values of decomposable forms*, *Canadian J. of Math* **70** (2018), 1390-1415.
- [14] J. Xu, *Stratification for Multiplicative Character Sums*, *Int. Math. Res. Not. IMRN*, 05 2018.rny096.

INSTITUTE OF ANALYSIS AND NUMBER THEORY, GRAZ UNIVERSITY OF TECHNOLOGY, KOPERNIKUS-GASSE 24/II, 8010 GRAZ, AUSTRIA

*E-mail address:* lapkova@math.tugraz.at

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, BAHEN CENTRE, 40 ST. GEORGE ST., ROOM 6290, TORONTO, ONTARIO, CANADA M5S 2E4

*E-mail address:* stanley.xiao@math.toronto.edu