

# Super congruences concerning binomial coefficients and Apéry-like numbers

Zhi-Hong Sun

School of Mathematics and Statistics

Huaian, Jiangsu 223300, P.R. China

Email: zhsun@hytc.edu.cn

Homepage: <http://maths.hytc.edu.cn/szh1.htm>

## Abstract

Let  $p$  be a prime with  $p > 3$ , and let  $a, b$  be two rational  $p$ -integers. In this paper we present general congruences for  $\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{p}{k+b} \pmod{p^2}$ . For  $n = 0, 1, 2, \dots$  let  $D_n$  and  $b_n$  be Domb and Almkvist-Zudilin numbers, respectively. We also establish congruences for

$$\sum_{n=0}^{p-1} \frac{D_n}{16^n}, \quad \sum_{n=0}^{p-1} \frac{D_n}{4^n}, \quad \sum_{n=0}^{p-1} \frac{b_n}{(-3)^n}, \quad \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \pmod{p^2}$$

in terms of certain binary quadratic forms.

MSC: Primary 11A07, Secondary 05A19, 11B65, 11E25

Keywords: Congruence; binomial coefficients; Domb numbers; Almkvist-Zudilin numbers; binary quadratic form

## 1. Introduction

Let  $p > 3$  be a prime. In 2003, based on his work concerning hypergeometric functions and Calabi-Yau manifolds, Rodriguez-Villegas [RV] conjectured the following congruences:

$$(1.1) \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k} \equiv \left(\frac{-1}{p}\right) \pmod{p^2}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \equiv \left(\frac{-3}{p}\right) \pmod{p^2},$$

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k} \equiv \left(\frac{-2}{p}\right) \pmod{p^2}, \quad \sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k} \equiv \left(\frac{-1}{p}\right) \pmod{p^2},$$

where  $\left(\frac{a}{p}\right)$  is the Legendre symbol. These congruences were later confirmed by Mortenson [M1-M2] via the Gross-Koblitz formula. It is easily seen that (see [S7])

$$(1.2) \quad \left(\frac{-\frac{1}{2}}{k}\right)^2 = \frac{\binom{2k}{k}^2}{16^k}, \quad \left(\frac{-\frac{1}{3}}{k}\right) \left(\frac{-\frac{2}{3}}{k}\right) = \frac{\binom{2k}{k} \binom{3k}{k}}{27^k},$$

$$\left(\frac{-\frac{1}{4}}{k}\right) \left(\frac{-\frac{3}{4}}{k}\right) = \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k}, \quad \left(\frac{-\frac{1}{6}}{k}\right) \left(\frac{-\frac{5}{6}}{k}\right) = \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k}.$$

Let  $\mathbb{Z}$  be the set of integers. For a prime  $p$  let  $\mathbb{Z}_p$  be the set of rational numbers whose denominator is not divisible by  $p$ . In [S7] the author generalized (1.1) by proving that for  $a \in \mathbb{Z}_p$ ,

$$(1.3) \quad \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \equiv (-1)^{\langle a \rangle_p} \pmod{p^2},$$

where  $\langle a \rangle_p \in \{0, 1, \dots, p-1\}$  is given by  $a \equiv \langle a \rangle_p \pmod{p}$ . In [S9] the author established the congruence

$$(1.4) \quad \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{2a+1}{2k+1} \equiv 1 + 2 \frac{a - \langle a \rangle_p}{p} \pmod{p^2}.$$

Let  $p$  be an odd prime and  $a, b \in \mathbb{Z}_p$  with  $ab \not\equiv 0 \pmod{p}$ . In Section 2 we show that for  $\langle b \rangle_p \leq \langle a \rangle_p$  and  $\langle b \rangle_p \leq p-1 - \langle a \rangle_p$ ,

$$(1.5) \quad \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{1}{k+b} \equiv \frac{p(s+t+1)(s-t)}{b^2(s+1) \binom{\langle a \rangle_p}{\langle b \rangle_p} \binom{p-1-\langle a \rangle_p}{\langle b \rangle_p}} \pmod{p^2},$$

where  $t = (a - \langle a \rangle_p)/p$  and  $s = (b - \langle b \rangle_p)/p \not\equiv -1 \pmod{p}$ . In Section 3 we present a congruence for  $\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{p}{k-a} \pmod{p^2}$  under the condition  $1 \leq \langle a \rangle_p < \frac{p}{2}$ . In Section 4, using a combinatorial identity we obtain a congruence for  $\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{p}{16^k(k+b)} \pmod{p^2}$  under the condition  $\langle b \rangle_p > \frac{p}{2}$ . In particular, we prove that

$$\begin{aligned} \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{p}{16^k(6k+1)} &\equiv 4x^2 - 2p \pmod{p^2} \quad \text{for } p = x^2 + 3y^2 \equiv 1 \pmod{3}, \\ (2 - (-1)^{\frac{p-1}{2}}) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{p}{16^k(8k+1)} &\equiv 4x^2 - 2p \pmod{p^2} \quad \text{for } p = x^2 + 2y^2 \equiv 1, 3 \pmod{8}, \\ (3 \binom{p}{3} - 2) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{p}{16^k(12k+1)} &\equiv 4x^2 - 2p \pmod{p^2} \quad \text{for } p = x^2 + 4y^2 \equiv 1 \pmod{4}, \end{aligned}$$

where  $x$  and  $y$  are integers. We mention that in [S11, Theorem 2.8 and (6)] the author proved that

$$\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{p}{16^k(4k+1)} \equiv 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3} \quad \text{for prime } p = x^2 + 4y^2 \equiv 1 \pmod{4}.$$

In Section 5 we use congruences in Sections 2-4 to prove super congruences for three Apéry-like numbers. Let  $\{D_n\}$ ,  $\{b_n\}$  and  $\{W_n\}$  be Apéry-like numbers given by

$$(1.6) \quad D_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k} \binom{2n-2k}{n-k},$$

$$(1.7) \quad b_n = \sum_{k=0}^{\lfloor n/3 \rfloor} \binom{2k}{k} \binom{3k}{k} \binom{n}{3k} \binom{n+k}{k} (-3)^{n-3k},$$

$$(1.8) \quad W_n = \sum_{k=0}^{\lfloor n/3 \rfloor} \binom{2k}{k} \binom{3k}{k} \binom{n}{3k} (-3)^{n-3k},$$

where  $\lfloor x \rfloor$  is the greatest integer not exceeding  $x$ . The numbers  $D_n$  ( $n = 0, 1, 2, \dots$ ) are called Domb numbers, and  $b_n$  ( $n = 0, 1, 2, \dots$ ) are called Almkvist-Zudilin numbers.

For  $D_n, b_n, W_n$  See A002895, A125143 and A291898 in Sloane's database "The On-Line Encyclopedia of Integer Sequences", and related papers [AT], [CZ], [S8] and [S12].

For positive integers  $a, b$  and  $n$ , if  $n = ax^2 + by^2$  for some integers  $x$  and  $y$ , we briefly write that  $n = ax^2 + by^2$ . Let  $p$  be a prime with  $p > 3$ . In [Su2], the author's brother Z.W. Sun posed the following conjecture:

$$(1.9) \quad \sum_{n=0}^{p-1} \frac{D_n}{16^n} \equiv \sum_{n=0}^{p-1} \frac{D_n}{4^n} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } 3 \mid p-1 \text{ and so } p = x^2 + 3y^2, \\ 0 \pmod{p^2} & \text{if } 3 \mid p-2. \end{cases}$$

In [S12, Conjecture 4.15] the author conjectured that

$$\sum_{n=0}^{p-1} \frac{D_n}{4^n} \equiv \sum_{n=0}^{p-1} \frac{D_n}{16^n} \equiv 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3} \quad \text{for } p = x^2 + 3y^2 \equiv 1 \pmod{3},$$

and

$$\sum_{n=0}^{p-1} \frac{D_n}{4^n} \equiv -2 \sum_{n=0}^{p-1} \frac{D_n}{16^n} \equiv \frac{p^2}{2} \left( \frac{p-1}{2} \right)^{-2} \pmod{p^3} \quad \text{for } p \equiv 2 \pmod{3}.$$

In [S12, Conjecture 4.16] the author also conjectured that

$$\sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \equiv \begin{cases} 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3} & \text{if } 12 \mid p-1 \text{ and } p = x^2 + 9y^2, \\ 2p - 2x^2 + \frac{p^2}{2x^2} \pmod{p^3} & \text{if } 12 \mid p-5 \text{ and } 2p = x^2 + 9y^2 \end{cases}$$

and

$$\sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv -15 \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \equiv \begin{cases} -\frac{5}{3} p^2 \binom{[p/3]}{[p/12]}^{-2} \pmod{p^3} & \text{if } 12 \mid p-7, \\ \frac{5}{6} p^2 \binom{[p/3]}{[p/12]}^{-2} \pmod{p^3} & \text{if } 12 \mid p-11. \end{cases}$$

In Section 5 we prove (1.9) and

$$(1.10) \quad \sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \equiv \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{4k+1} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } 12 \mid p-1 \text{ and so } p = x^2 + 9y^2, \\ 2p - 2x^2 \pmod{p^2} & \text{if } 12 \mid p-5 \text{ and so } 2p = x^2 + 9y^2, \\ 0 \pmod{p^2} & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

We also determine  $\sum_{n=0}^{p-1} \frac{W_n}{(-3)^n}$  modulo  $p^2$ . In particular, we show that for  $p \equiv 1 \pmod{3}$  with  $4p = L^2 + 27M^2$  and  $L \equiv 1 \pmod{3}$ ,  $\sum_{n=0}^{p-1} \frac{W_n}{(-3)^n} \equiv -L + \frac{p}{L} \pmod{p^2}$ . This was conjectured by the author's brother Z.W. Sun in [Su1].

Throughout this paper, let  $q_p(a) = (a^{p-1} - 1)/p$  for given odd prime  $p$  and  $a \in \mathbb{Z}$ , and let  $H_0 = 0$  and  $H_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n}$  ( $n \geq 1$ ).

## 2. Congruences for $\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{1}{k+b} \pmod{p^2}$ .

We begin with basic congruences for harmonic numbers  $H_n$ . For odd prime  $p$  and  $k \in \{1, 2, \dots, p-1\}$ , it is clear that

$$(2.1) \quad H_{p-1-k} = \sum_{i=1}^{p-1} \frac{1}{i} - \sum_{i=1}^k \frac{1}{p-i} \equiv 0 + \sum_{i=1}^k \frac{1}{i} = H_k \pmod{p}.$$

For  $k = 1, 2, \dots, \frac{p-1}{2}$  we have

$$H_{\frac{p-1}{2}-k} = H_{\frac{p-1}{2}} - \sum_{i=1}^k \frac{1}{\frac{p+1}{2}-i} \equiv H_{\frac{p-1}{2}} + 2 \sum_{i=1}^k \frac{1}{2i-1} = H_{\frac{p-1}{2}} + 2(H_{2k} - \frac{1}{2}H_k) \pmod{p}.$$

It is well known that (see [L])

$$(2.2) \quad H_{\frac{p-1}{2}} \equiv -2q_p(2) \quad \text{and} \quad H_{[\frac{p}{3}]} \equiv -\frac{3}{2}q_p(3) \pmod{p} \text{ for } p > 3.$$

Thus

$$(2.3) \quad 2H_{2k} \equiv 2q_p(2) + H_k + H_{\frac{p-1}{2}-k} \pmod{p} \quad \text{for } k = 1, 2, \dots, \frac{p-1}{2}.$$

From (2.2) and (2.3) we deduce the known congruences (see [L]):

$$(2.4) \quad H_{[\frac{p}{4}]} \equiv -3q_p(2) \pmod{p} \quad \text{and} \quad H_{[\frac{p}{6}]} \equiv -2q_p(2) - \frac{3}{2}q_p(3) \pmod{p} \text{ for } p > 3.$$

Let  $p$  be an odd prime,  $a, m \in \mathbb{Z}_p$ ,  $k \in \{1, 2, \dots, p-1\}$  and  $a-i \not\equiv 0 \pmod{p^2}$  for  $i = 0, 1, \dots, k-1$ . Then clearly

$$\begin{aligned} \binom{a+mp}{k} &= \frac{(mp+a)(mp+a-1)\cdots(mp+a-k+1)}{k!} \\ &\equiv \frac{1}{k!} \left( a(a-1)\cdots(a-k+1) + mp \sum_{i=0}^{k-1} \frac{a(a-1)\cdots(a-k+1)}{a-i} \right) \pmod{p^2}. \end{aligned}$$

That is,

$$(2.5) \quad \binom{a+mp}{k} \equiv \binom{a}{k} \left( 1 + \sum_{i=0}^{k-1} \frac{mp}{a-i} \right) \pmod{p^2}.$$

By [S9,(2.2)], for  $n = 0, 1, 2, \dots$ ,

$$\sum_{k=0}^n \binom{a}{k} \binom{-1-a}{k} \frac{ab+1}{bk+1} - \sum_{k=0}^n \binom{a-1}{k} \binom{-a}{k} \frac{ab-1}{bk+1} = 2 \binom{a-1}{n} \binom{-a-1}{n}.$$

Replacing  $b$  with  $\frac{1}{b}$  yields

$$(2.6) \quad \sum_{k=0}^n \binom{a}{k} \binom{-1-a}{k} \frac{a+b}{k+b} - \sum_{k=0}^n \binom{a-1}{k} \binom{-a}{k} \frac{a-b}{k+b} = 2 \binom{a-1}{n} \binom{-a-1}{n}.$$

By [S9, Lemma 2.2], for odd prime  $p$  and  $a \in \mathbb{Z}_p$  with  $a \not\equiv 0 \pmod{p}$ ,

$$(2.7) \quad \binom{a-1}{p-1} \binom{-a-1}{p-1} \equiv \frac{(a - \langle a \rangle_p)(p + a - \langle a \rangle_p)}{a^2} \pmod{p^3}.$$

Now taking  $b = a$  in (2.6) and applying (2.7) yields

$$(2.8) \quad \sum_{k=0}^{p-1} \frac{\binom{a}{k} \binom{-1-a}{k}}{k+a} = \frac{1}{a} \binom{a-1}{p-1} \binom{-a-1}{p-1} \equiv \frac{(a - \langle a \rangle_p)(p + a - \langle a \rangle_p)}{a^3} \pmod{p^3}.$$

**Lemma 2.1.** *Let  $p$  be an odd prime,  $a, b \in \mathbb{Z}_p$ ,  $a \not\equiv 0 \pmod{p}$ ,  $b + \langle -b \rangle_p \not\equiv 0 \pmod{p^2}$  and*

$$S(a, b) = \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{p}{k+b}.$$

*Suppose that  $m \in \{0, 1, \dots, \langle a \rangle_p - 1\}$  and  $\langle a + b \rangle_p \notin \{0, 1, \dots, m\}$ . Then*

$$S(a, b) \equiv \frac{\binom{a-b}{m+1}}{\binom{a+b}{m+1}} S(a - m - 1, b) \pmod{p^3}.$$

*Proof.* Since  $b + \langle -b \rangle_p \not\equiv 0 \pmod{p^2}$  we see that  $b + k \not\equiv 0 \pmod{p^2}$  and so  $\frac{p}{k+b} \in \mathbb{Z}_p$  for  $k = 0, 1, \dots, p-1$ . Hence  $S(a, b) \in \mathbb{Z}_p$ . For  $k = 0, 1, \dots, \langle a \rangle_p - 1$  we have  $a - k \not\equiv 0 \pmod{p}$ . From (2.6) and (2.7),

$$\begin{aligned} & (a - k + b)S(a - k, b) - (a - k - b)S(a - k - 1, b) \\ &= 2p \binom{a - k - 1}{p-1} \binom{-(a - k) - 1}{p-1} \equiv 0 \pmod{p^3}. \end{aligned}$$

Thus, for  $k = 0, 1, \dots, m$ ,

$$S(a - k, b) \equiv \frac{a - b - k}{a + b - k} S(a - k - 1, b) \pmod{p^3}.$$

Therefore,

$$\begin{aligned} S(a, b) &\equiv \frac{a-b}{a+b} S(a-1, b) \equiv \frac{a-b}{a+b} \cdot \frac{a-b-1}{a+b-1} S(a-2, b) \\ &\equiv \dots \equiv \prod_{k=0}^m \frac{a-b-k}{a+b-k} \cdot S(a-m-1, b) \pmod{p^3}. \end{aligned}$$

To see the result, we note that

$$\prod_{k=0}^m \frac{a-b-k}{a+b-k} = \frac{\binom{a-b}{m+1}}{\binom{a+b}{m+1}}.$$

**Lemma 2.2.** *Let  $p$  be an odd prime,  $b, t \in \mathbb{Z}_p$ ,  $bt \not\equiv 0 \pmod{p}$  and  $s = (b - \langle b \rangle_p)/p \not\equiv -1 \pmod{p}$ . Then*

$$\sum_{k=0}^{p-1} \binom{pt}{k} \binom{-1-pt}{k} \frac{p}{k+b} \equiv \left(1 + \frac{t}{s+1}\right) \frac{p}{b} \left(1 - \frac{pt}{b}\right) \pmod{p^3}.$$

Proof. It is well known that  $\sum_{k=1}^{p-1} \frac{1}{k} \equiv 0 \pmod{p}$ . Thus,

$$\begin{aligned}
& \sum_{k=1}^{p-1} \binom{pt}{k} \binom{-1-pt}{k} \frac{p}{k+b} \\
&= \sum_{k=1}^{p-1} (-1)^k \frac{pt(pt+k)(p^2t^2-1^2)\cdots(p^2t^2-(k-1)^2)}{k!^2} \cdot \frac{p}{k+b} \\
&\equiv -\sum_{k=1}^{p-1} \frac{pt(pt+k)}{k^2} \cdot \frac{p}{k+b} = -p^2t^2 \sum_{k=1}^{p-1} \frac{p}{k^2(k+b)} - \frac{pt}{b} \sum_{k=1}^{p-1} \left( \frac{p}{k} - \frac{p}{k+b} \right) \\
&\equiv -\frac{p^2t^2}{(p-\langle b \rangle_p)^2(p-\langle b \rangle_p+b)/p} + \frac{pt}{b} \left( \frac{1}{(p-\langle b \rangle_p+b)/p} + \sum_{\substack{k=1 \\ k \neq p-\langle b \rangle_p}}^{p-1} \frac{p}{k+b} \right) \\
&\equiv -\frac{p^2t^2}{b^2(s+1)} + \frac{pt}{b} \left( \frac{1}{s+1} - \frac{p}{b} \right) \pmod{p^3},
\end{aligned}$$

which yields the result.

**Theorem 2.1.** *Let  $p$  be an odd prime,  $a, b \in \mathbb{Z}_p$ ,  $ab \not\equiv 0 \pmod{p}$  and  $\langle b \rangle_p \leq p-1-\langle a \rangle_p$ . Assume  $t = (a - \langle a \rangle_p)/p$  and  $s = (b - \langle b \rangle_p)/p \not\equiv -1 \pmod{p}$ . For  $\langle b \rangle_p \leq \langle a \rangle_p$  we have*

$$\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{1}{k+b} \equiv \frac{p(s+t+1)(s-t)}{b^2(s+1) \binom{\langle a \rangle_p}{\langle b \rangle_p} \binom{p-1-\langle a \rangle_p}{\langle b \rangle_p}} \pmod{p^2}.$$

For  $\langle b \rangle_p > \langle a \rangle_p$  we have

$$\begin{aligned}
\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{1}{k+b} &\equiv \frac{s+1+t}{b(s+1)} \cdot \frac{\binom{\langle b \rangle_p-1}{\langle a \rangle_p}}{\binom{p-1-\langle b \rangle_p}{\langle a \rangle_p}} \left( 1 + p \frac{s+1}{b} + p(2s+1)H_{\langle b \rangle_p-1} \right. \\
&\quad \left. - p(s-t)H_{\langle b \rangle_p-\langle a \rangle_p-1} - p(s+t+1)H_{\langle a \rangle_p+\langle b \rangle_p} \right) \pmod{p^2}.
\end{aligned}$$

Proof. For  $k \in \{0, 1, \dots, p-1\}$  clearly  $k+b \not\equiv 0 \pmod{p}$  for  $k \neq p-\langle b \rangle_p$ . For  $k = p-\langle b \rangle_p$  we have  $k+b \not\equiv 0 \pmod{p^2}$ ,  $\binom{a}{k} = \binom{a}{p-\langle b \rangle_p} \equiv \binom{\langle a \rangle_p}{p-\langle b \rangle_p} = 0 \pmod{p}$  and so  $\binom{a}{k} \frac{1}{k+b} \in \mathbb{Z}_p$ . Therefore,  $\binom{a}{k} \binom{-1-a}{k} \frac{1}{k+b} \in \mathbb{Z}_p$  for  $k = 0, 1, \dots, p-1$ . Now taking  $m = \langle a \rangle_p - 1$  in Lemma 2.1 and then applying Lemma 2.2 gives

$$(2.9) \quad \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{1}{k+b} \equiv \frac{\binom{a-b}{\langle a \rangle_p}}{\binom{a+b}{\langle a \rangle_p}} \left( 1 + \frac{t}{s+1} \right) \frac{1}{b} \left( 1 - \frac{pt}{b} \right) \pmod{p^2}.$$

We first assume  $\langle b \rangle_p \leq \langle a \rangle_p$ . Since  $a-b \equiv \langle a \rangle_p - \langle b \rangle_p \pmod{p}$ , we see that

$$\begin{aligned}
\frac{\binom{a-b}{\langle a \rangle_p}}{\binom{a+b}{\langle a \rangle_p}} &= (a-b - (\langle a \rangle_p - \langle b \rangle_p)) \frac{(a-b)(a-b-1)\cdots(a-b - (\langle a \rangle_p - \langle b \rangle_p - 1))}{\langle a \rangle_p!} \\
&\quad \times \frac{(a-b - (\langle a \rangle_p - \langle b \rangle_p + 1))\cdots(a-b - \langle a \rangle_p + 1)}{\binom{a+b}{\langle a \rangle_p}}
\end{aligned}$$

$$\begin{aligned}
&\equiv p(t-s) \frac{(\langle a \rangle_p - \langle b \rangle_p)!}{\langle a \rangle_p!} \cdot \frac{(-1)(-2) \cdots (-\langle b \rangle_p - 1)}{\binom{\langle a \rangle_p + \langle b \rangle_p}{\langle a \rangle_p}} \\
&= \frac{p(t-s)}{-\langle b \rangle_p} \cdot \frac{(\langle a \rangle_p - \langle b \rangle_p)! (-1)^{\langle b \rangle_p} \langle b \rangle_p!}{\langle a \rangle_p! (-1)^{\langle b \rangle_p} \binom{-1-\langle a \rangle_p}{\langle b \rangle_p}} \equiv \frac{p(s-t)}{b \binom{\langle a \rangle_p}{\langle b \rangle_p} \binom{p-1-\langle a \rangle_p}{\langle b \rangle_p}} \pmod{p^2}.
\end{aligned}$$

This together with (2.9) yields the result in the case  $\langle b \rangle_p \leq \langle a \rangle_p$ .

For  $\langle b \rangle_p > \langle a \rangle_p$ , using (2.5) we see that

$$\begin{aligned}
\frac{\binom{a-b}{\langle a \rangle_p}}{\binom{a+b}{\langle a \rangle_p}} &= \frac{\binom{b-1-a+\langle a \rangle_p}{\langle a \rangle_p}}{\binom{-b-1-a+\langle a \rangle_p}{\langle a \rangle_p}} = \frac{\binom{\langle b \rangle_p - 1 + p(s-t)}{\langle a \rangle_p}}{\binom{p-1-\langle b \rangle_p - p(s+t+1)}{\langle a \rangle_p}} \\
&\equiv \frac{\binom{\langle b \rangle_p - 1}{\langle a \rangle_p} (1 + p(s-t) \sum_{i=0}^{\langle a \rangle_p - 1} \frac{1}{\langle b \rangle_p - 1 - i})}{\binom{p-1-\langle b \rangle_p}{\langle a \rangle_p} (1 - p(s+t+1) \sum_{i=0}^{\langle a \rangle_p - 1} \frac{1}{p-1-\langle b \rangle_p - i})} \\
&\equiv \frac{\binom{\langle b \rangle_p - 1}{\langle a \rangle_p} (1 + p(s-t)(H_{\langle b \rangle_p - 1} - H_{\langle b \rangle_p - \langle a \rangle_p - 1})}{\binom{p-1-\langle b \rangle_p}{\langle a \rangle_p} (1 + p(s+t+1)(H_{\langle a \rangle_p + \langle b \rangle_p} - H_{\langle b \rangle_p})} \\
&\equiv \frac{\binom{\langle b \rangle_p - 1}{\langle a \rangle_p}}{\binom{p-1-\langle b \rangle_p}{\langle a \rangle_p}} (1 + p(s-t)(H_{\langle b \rangle_p - 1} - H_{\langle b \rangle_p - \langle a \rangle_p - 1}) \\
&\quad \times (1 - p(s+t+1)(H_{\langle a \rangle_p + \langle b \rangle_p} - H_{\langle b \rangle_p})) \\
&\equiv \frac{\binom{\langle b \rangle_p - 1}{\langle a \rangle_p}}{\binom{p-1-\langle b \rangle_p}{\langle a \rangle_p}} \left( 1 + p \frac{s+t+1}{b} + p(2s+1)H_{\langle b \rangle_p - 1} \right. \\
&\quad \left. - p(s-t)H_{\langle b \rangle_p - \langle a \rangle_p - 1} - p(s+t+1)H_{\langle a \rangle_p + \langle b \rangle_p} \right) \pmod{p^2}.
\end{aligned}$$

This together with (2.9) yields the remaining part.

**Corollary 2.1.** *Let  $p$  be a prime with  $p > 3$ ,  $a \in \mathbb{Z}_p$ ,  $1 \leq \langle a \rangle_p \leq \frac{p-3}{2}$  and  $t = (a - \langle a \rangle_p)/p \not\equiv -1 \pmod{p}$ . Then*

$$\begin{aligned}
&\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{1}{k+a+1} \\
&\equiv \frac{2t+1}{(a+1)(t+1) \binom{p-2-\langle a \rangle_p}{\langle a \rangle_p}} \left( 1 + p \frac{t+1}{a+1} - p(2t+1)(H_{2\langle a \rangle_p + 1} - H_{\langle a \rangle_p}) \right) \pmod{p^2}.
\end{aligned}$$

*Proof.* Set  $b = a + 1$ . Then  $\langle b \rangle_p = \langle a \rangle_p + 1$  and so  $\langle a \rangle_p < \langle b \rangle_p \leq p - 1 - \langle a \rangle_p$ . Now putting  $b = a + 1$  in Theorem 2.1 and noting that  $s = t$  yields the result.

**Theorem 2.2.** *Let  $p > 3$  be a prime. Suppose  $b \in \mathbb{Z}_p$ ,  $\langle b \rangle_p \neq 0$  and  $s = (b - \langle b \rangle_p)/p \not\equiv -1 \pmod{p}$ . Then*

$$(2.10) \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k (k+b)} \equiv \frac{(s+1/2)^2 p}{b^2 (s+1) \binom{(p-1)/2}{\langle b \rangle_p}^2} \pmod{p^2} \quad \text{for } \langle b \rangle_p < \frac{p}{2},$$

$$(2.11) \quad \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^2}{16^k(k+b)} \equiv \frac{b - \langle b \rangle_p}{b^2 \binom{(p-1)/2}{\langle b \rangle_p}} \pmod{p^2} \quad \text{for } \langle b \rangle_p < \frac{p}{2},$$

$$(2.12) \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k(k+b)} \equiv (-1)^{\langle b \rangle_p} \frac{(s+1/3)(s+2/3)p}{b^2(s+1) \binom{2\langle b \rangle_p}{\langle b \rangle_p} \binom{[p/3]+\langle b \rangle_p}{[p/3]-\langle b \rangle_p}} \pmod{p^2} \quad \text{for } \langle b \rangle_p < \frac{p}{3},$$

$$(2.13) \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k(k+b)} \equiv (-1)^{\langle b \rangle_p} \frac{(s+1/4)(s+3/4)p}{b^2(s+1) \binom{2\langle b \rangle_p}{\langle b \rangle_p} \binom{[p/4]+\langle b \rangle_p}{[p/4]-\langle b \rangle_p}} \pmod{p^2} \quad \text{for } \langle b \rangle_p < \frac{p}{4},$$

$$(2.14) \quad \sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k(k+b)} \equiv (-1)^{\langle b \rangle_p} \frac{(s+1/6)(s+5/6)p}{b^2(s+1) \binom{2\langle b \rangle_p}{\langle b \rangle_p} \binom{[p/6]+\langle b \rangle_p}{[p/6]-\langle b \rangle_p}} \pmod{p^2} \quad \text{for } \langle b \rangle_p < \frac{p}{6}.$$

Proof. Set  $a = -\frac{1}{2}$ . Then  $\langle a \rangle_p = \frac{p-1}{2}$  and  $t = (a - \langle a \rangle_p)/p = -1/2$ . For  $\langle b \rangle_p < \frac{p}{2}$  we have  $\langle b \rangle_p \leq p-1 - \langle a \rangle_p = \langle a \rangle_p$ . Since  $\binom{-1/2}{k}^2 = \binom{2k}{k} 16^{-k}$ , taking  $a = -\frac{1}{2}$  in Theorem 2.1 yields (2.10). Clearly,

$$\begin{aligned} \frac{1}{p} \binom{2(p - \langle b \rangle_p)}{p - \langle b \rangle_p} &= \frac{(2p - 2\langle b \rangle_p)(2p - (2\langle b \rangle_p - 1)) \cdots (p+1) \cdot (p-1) \cdots (p - (\langle b \rangle_p - 1))}{(p - \langle b \rangle_p)!} \\ &\equiv \frac{(p - 2\langle b \rangle_p)! (-1)^{\langle b \rangle_p - 1} (\langle b \rangle_p - 1)!}{(p - \langle b \rangle_p)!} \equiv \frac{2(-1)^{\langle b \rangle_p} \langle b \rangle_p!}{(p - \langle b \rangle_p) \cdots (p - 2\langle b \rangle_p)} \\ &\equiv \frac{2}{-b \binom{2\langle b \rangle_p}{\langle b \rangle_p}} \equiv \frac{2}{-b(-4)^{\langle b \rangle_p} \binom{(p-1)/2}{\langle b \rangle_p}} \pmod{p}. \end{aligned}$$

Since  $p \mid \binom{2k}{k}$  for  $\frac{p}{2} < k < p$ , using (2.10) we see that

$$\begin{aligned} \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^2}{16^k(k+b)} &\equiv \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k(k+b)} - \frac{\binom{2(p-\langle b \rangle_p)}{p-\langle b \rangle_p}^2}{16^{p-\langle b \rangle_p}(p-\langle b \rangle_p+b)} \\ &\equiv \frac{p(s+1/2)^2}{b^2(s+1) \binom{(p-1)/2}{\langle b \rangle_p}^2} - \frac{1}{16^{p-\langle b \rangle_p}(s+1)} \cdot \frac{4p}{16^{\langle b \rangle_p} b^2 \binom{(p-1)/2}{\langle b \rangle_p}^2} \\ &\equiv \frac{ps}{b^2 \binom{(p-1)/2}{\langle b \rangle_p}^2} = \frac{b - \langle b \rangle_p}{b^2 \binom{(p-1)/2}{\langle b \rangle_p}^2} \pmod{p^2}. \end{aligned}$$

This proves (2.11).

Now we consider (2.12)-(2.14). We first note that

$$\begin{aligned} \binom{\langle a \rangle_p}{\langle b \rangle_p} \binom{p-1-\langle a \rangle_p}{\langle b \rangle_p} &\equiv \binom{\langle a \rangle_p}{\langle b \rangle_p} \binom{-1-\langle a \rangle_p}{\langle b \rangle_p} = (-1)^{\langle b \rangle_p} \binom{\langle a \rangle_p}{\langle b \rangle_p} \binom{\langle a \rangle_p + \langle b \rangle_p}{\langle a \rangle_p} \\ &= (-1)^{\langle b \rangle_p} \binom{2\langle b \rangle_p}{\langle b \rangle_p} \binom{\langle a \rangle_p + \langle b \rangle_p}{\langle a \rangle_p - \langle b \rangle_p} \pmod{p}. \end{aligned}$$

Set  $a = -\frac{1}{3}$  or  $-\frac{2}{3}$  according as  $p \equiv 1 \pmod{3}$  or  $p \equiv 2 \pmod{3}$ . Then  $\langle a \rangle_p = [\frac{p}{3}]$  and  $t = (a - \langle a \rangle_p)/p = -\frac{1}{3}$ . Recall that  $\binom{-1/3}{k} \binom{-2/3}{k} = \binom{2k}{k} \binom{3k}{k} 27^{-k}$ . From Theorem 2.1 and the above we obtain (2.12).



Next consider (2.13). Set  $a = -\frac{1}{4}$  or  $-\frac{3}{4}$  according as  $p \equiv 1 \pmod{4}$  or  $p \equiv 3 \pmod{4}$ . Then  $\langle a \rangle_p = \left[\frac{p}{4}\right]$  and  $t = (a - \langle a \rangle_p)/p = -\frac{1}{4}$ . Recall that  $\binom{-1/4}{k} \binom{-3/4}{k} = \binom{2k}{k} \binom{4k}{2k} 64^{-k}$ , (2.13) follows from Theorem 2.1 (with  $a = -\frac{1}{4}$ ).

Finally consider (2.14). Set  $a = -\frac{1}{6}$  or  $-\frac{5}{6}$  according as  $p \equiv 1 \pmod{6}$  or  $p \equiv 5 \pmod{6}$ . Then  $\langle a \rangle_p = \left[\frac{p}{6}\right]$  and  $t = (a - \langle a \rangle_p)/p = -\frac{1}{6}$ . Since  $\binom{-1/6}{k} \binom{-5/6}{k} = \binom{3k}{k} \binom{6k}{3k} 432^{-k}$ , applying Theorem 2.1 yields (2.14). The proof is now complete.

Let  $p$  be a prime with  $p > 3$ . As examples, from (2.10), (2.11) and the fact  $\binom{(p-1)/2}{k} \equiv \binom{-1/2}{k} = \binom{2k}{k} 4^{-k} \pmod{p}$  we deduce that

$$\begin{aligned} - \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^2}{16^k(4k+3)} &\equiv 3 \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k(4k+3)} \equiv p \left(\frac{p-1}{4}\right)^{-2} \pmod{p^2} \text{ for } p \equiv 1 \pmod{4}, \\ 8 \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k(3k+1)} &\equiv 5 \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k(6k+1)} \equiv p \left(\frac{p-1}{6}\right)^{-2} \pmod{p^2} \text{ for } p \equiv 5 \pmod{6}, \\ 45 \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k(8k+1)} &\equiv 7 \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k(8k+3)} \equiv -p \left(\frac{[p/4]}{[p/8]}\right)^{-2} \pmod{p^2} \text{ for } p \equiv 5 \pmod{8}, \\ \frac{14}{9} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k(8k+1)} &\equiv 10 \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k(8k+3)} \equiv p \left(\frac{[p/4]}{[p/8]}\right)^{-2} \pmod{p^2} \text{ for } p \equiv 7 \pmod{8}. \end{aligned}$$

**Theorem 2.3.** *Let  $p$  be a prime with  $p > 3$ . Then*

$$(2.15) \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k(3k+2)} \equiv \frac{1}{2 \binom{(2(p-1)/3)}{(p-1)/3}} \pmod{p^2} \text{ for } p \equiv 1 \pmod{3},$$

$$(2.16) \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k(3k+1)} \equiv \frac{1+2p}{2 \binom{(2(p-2)/3)}{(p-2)/3}} \pmod{p^2} \text{ for } p \equiv 2 \pmod{3},$$

$$(2.17) \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k(4k+3)} \equiv (-1)^{\frac{p-1}{4}} \frac{2^{p-1} + 1}{6 \binom{(p-1)/2}{(p-1)/4}} \pmod{p^2} \text{ for } p \equiv 1 \pmod{4},$$

$$(2.18) \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k(4k+1)} \equiv (-1)^{\frac{p+1}{4}} \frac{2^{p-1} + 2p + 1}{6 \binom{(p-3)/2}{(p-3)/4}} \pmod{p^2} \text{ for } p \equiv 3 \pmod{4},$$

$$(2.19) \quad \sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k(6k+5)} \equiv \frac{1 - \frac{4}{3}(2^{p-1} - 1)}{5 \binom{(5(p-1)/6)}{(p-1)/6}} \pmod{p^2} \text{ for } p \equiv 1 \pmod{6},$$

$$(2.20) \quad \sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k(6k+1)} \equiv \frac{4}{5} \cdot \frac{1 + 5p - \frac{4}{3}(2^{p-1} - 1)}{\binom{(5p-7)/6}{(p-5)/6}} \pmod{p^2} \text{ for } p \equiv 5 \pmod{6}.$$

Proof. For  $a = -\frac{1}{3}$  and  $p \equiv 1 \pmod{3}$  we see that  $\langle a \rangle_p = \frac{p-1}{3}$ ,  $\frac{a - \langle a \rangle_p}{p} = -\frac{1}{3}$  and

$$H_{2\langle a \rangle_p + 1} - H_{\langle a \rangle_p} = H_{\frac{2p+1}{3}} - H_{\frac{p-1}{3}} \equiv H_{p-1-\frac{2p+1}{3}} - H_{\frac{p-1}{3}} = H_{\frac{p-4}{3}} - H_{\frac{p-1}{3}} = -\frac{3}{p-1} \equiv 3 \pmod{p}.$$

Thus, from Corollary 2.1 and (1.2),

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k (3k+2)} &= \frac{1}{3} \sum_{k=0}^{p-1} \binom{-\frac{1}{3}}{k} \binom{-\frac{2}{3}}{k} \frac{1}{k+2/3} \\ &\equiv \frac{2t+1}{3(a+1)(t+1)^{\binom{p-2-(p-1)/3}{(p-1)/3}}} \equiv \frac{1}{2^{\binom{2(p-1)/3}{(p-1)/3}}} \pmod{p^2}, \end{aligned}$$

For  $a = -\frac{2}{3}$  and  $p \equiv 2 \pmod{3}$  we see that  $\langle a \rangle_p = \frac{p-2}{3}$ ,  $\frac{a-\langle a \rangle_p}{p} = -\frac{1}{3}$  and

$$H_{2\langle a \rangle_p+1} - H_{\langle a \rangle_p} \equiv H_{p-2-2\langle a \rangle_p} - H_{\langle a \rangle_p} = H_{\frac{p-2}{3}} - H_{\frac{p-2}{3}} = 0 \pmod{p}.$$

Thus, from Corollary 2.1 and (1.2),

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k (3k+1)} &= \frac{1}{3} \sum_{k=0}^{p-1} \binom{-\frac{1}{3}}{k} \binom{-\frac{2}{3}}{k} \frac{1}{k+1/3} \\ &\equiv \frac{2t+1}{3(a+1)(t+1)^{\binom{p-2-(p-2)/3}{(p-2)/3}}} \left(1 + p \frac{t+1}{a+1}\right) = \frac{1+2p}{2^{\binom{2(p-2)/3}{(p-2)/3}}} \pmod{p^2}. \end{aligned}$$

For  $a = -\frac{1}{4}$  and  $p \equiv 1 \pmod{4}$  we see that  $\langle a \rangle_p = \frac{p-1}{4}$  and  $\frac{a-\langle a \rangle_p}{p} = -\frac{1}{4}$ . It is well known that  $H_{\frac{p-1}{2}} \equiv -2q_p(2) \pmod{p}$  and  $H_{\frac{p-1}{4}} \equiv -3q_p(2) \pmod{p}$ . Thus,

$$H_{2\langle a \rangle_p+1} - H_{\langle a \rangle_p} = \frac{1}{(p+1)/2} + H_{\frac{p-1}{2}} - H_{\frac{p-1}{4}} \equiv 2 - 2q_p(2) + 3q_p(2) = 2 + q_p(2) \pmod{p}.$$

By [S3, Lemma 2.5],

$$(-1)^{\frac{p-1}{4}} \binom{3(p-1)/4}{(p-1)/4} \equiv \left(3 - 2(-4)^{\frac{p-1}{4}}\right) \binom{(p-1)/2}{(p-1)/4} \equiv (1 - pq_p(2)) \binom{(p-1)/2}{(p-1)/4} \pmod{p^2}.$$

Therefore, taking  $a = -\frac{1}{4}$  in Corollary 2.1 and applying the above and (1.2) gives

$$\begin{aligned} &\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k (4k+3)} \\ &= \frac{1}{4} \sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k} \binom{-\frac{3}{4}}{k} \frac{1}{k+3/4} \equiv \frac{1}{4} \cdot \frac{\frac{1}{2}}{\frac{3}{4} \cdot \frac{3}{4} \binom{p-2-(p-1)/4}{(p-1)/4}} \left(1 + p - \frac{p}{2} (H_{\frac{p-1}{2}+1} - H_{\frac{p-1}{4}})\right) \\ &\equiv \frac{1}{3 \binom{3(p-1)/4}{(p-1)/4}} \left(1 - \frac{p}{2} q_p(2)\right) \equiv \frac{1}{3} (-1)^{\frac{p-1}{4}} \frac{1}{(1 - pq_p(2)) \binom{(p-1)/2}{(p-1)/4}} \left(1 - \frac{p}{2} q_p(2)\right) \\ &\equiv \frac{1}{3} (-1)^{\frac{p-1}{4}} \frac{1}{\binom{(p-1)/2}{(p-1)/4}} (1 + pq_p(2)) \left(1 - \frac{p}{2} q_p(2)\right) \equiv \frac{1}{3} (-1)^{\frac{p-1}{4}} \frac{1 + pq_p(2)/2}{\binom{(p-1)/2}{(p-1)/4}} \pmod{p^2}. \end{aligned}$$

For  $a = -\frac{3}{4}$  and  $p \equiv 3 \pmod{4}$  we see that  $\langle a \rangle_p = \frac{p-3}{4}$  and  $\frac{a-\langle a \rangle_p}{p} = -\frac{1}{4}$ . By [S3, Lemma 2.5],

$$(-1)^{\frac{p-3}{4}} \frac{\binom{(p-1)/2+(p-3)/4}{(p-3)/4}}{\binom{(p-1)/2}{(p-3)/4}} \equiv 3 - 2(-4)^{\frac{p-3}{4}} \frac{\binom{(p-1)/2}{(p-3)/4}}{\binom{(p-3)/2}{(p-3)/4}} = 3 - 2(-4)^{\frac{p-3}{4}} \frac{2p-2}{p+1}$$

$$\begin{aligned}
&\equiv 3 - 2 \cdot (-1)^{\frac{p+1}{4}} 2^{\frac{p-1}{2}} (1 - 2p) \\
&\equiv 3 - 2(1 + pq_p(2)/2) + 4p = 1 + 4p - pq_p(2) \pmod{p^2}.
\end{aligned}$$

Hence

$$\begin{aligned}
(-1)^{\frac{p-3}{4}} \binom{\frac{p-1}{2} + \frac{p-3}{4}}{\frac{p-3}{4}} &\equiv (1 + 4p - pq_p(2)) \binom{\frac{p-1}{2}}{\frac{p-3}{4}} \\
&= (1 + 4p - pq_p(2)) \frac{2p-2}{p+1} \binom{(p-3)/2}{(p-3)/4} \equiv -2(1 + 2p - pq_p(2)) \binom{(p-3)/2}{(p-3)/4}
\end{aligned}$$

Also,

$$H_{2\langle a \rangle_p + 1} - H_{\langle a \rangle_p} = H_{\frac{p-1}{2}} - H_{\frac{p-3}{4}} \equiv -2q_p(2) + 3q_p(2) = q_p(2) \pmod{p}.$$

Now, taking  $a = -\frac{3}{4}$  in Corollary 2.1 and then applying the above gives

$$\begin{aligned}
\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k (4k+1)} &= \frac{1}{4} \sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k} \binom{-\frac{3}{4}}{k} \frac{1}{k+1/4} \equiv \frac{2}{3} \cdot \frac{1 + 3p - pq_p(2)/2}{\binom{\frac{p-1}{2} + \frac{p-3}{4}}{\frac{p-3}{4}}} \\
&\equiv \frac{2}{3} (-1)^{\frac{p+1}{4}} \frac{1 + 3p - \frac{1}{2}pq_p(2)}{2(1 + 2p - pq_p(2)) \binom{\frac{p-3}{2}}{\frac{p-3}{4}}} \equiv \frac{1}{3} (-1)^{\frac{p+1}{4}} \frac{1 + p + \frac{1}{2}pq_p(2)}{\binom{\frac{p-3}{2}}{\frac{p-3}{4}}} \pmod{p^2}.
\end{aligned}$$

For  $a = -\frac{1}{6}$  and  $p \equiv 1 \pmod{6}$  we have  $\langle a \rangle_p = \frac{p-1}{6}$ ,  $\frac{a - \langle a \rangle_p}{p} = -\frac{1}{6}$ . Applying (2.2) and (2.4),

$$H_{2\langle a \rangle_p + 1} - H_{\langle a \rangle_p} = \frac{1}{(p+2)/3} + H_{\frac{p-1}{3}} - H_{\frac{p-1}{6}} \equiv \frac{3}{2} + 2q_p(2) \pmod{p}.$$

Now taking  $a = -\frac{1}{6}$  in Corollary 2.1 yields

$$\begin{aligned}
\sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{64^k (6k+5)} &= \frac{1}{6} \sum_{k=0}^{p-1} \binom{-\frac{1}{6}}{k} \binom{-\frac{5}{6}}{k} \frac{1}{k+5/6} \\
&\equiv \frac{\frac{1}{6} \cdot \frac{2}{3}}{\binom{5}{6} 2^{\binom{p-2-(p-1)/6}{(p-1)/6}}} \left(1 + p - \frac{2}{3}p \left(\frac{3}{2} + 2q_p(2)\right)\right) \equiv \frac{1 - \frac{4}{3}(2^{p-1} - 1)}{5 \binom{5(p-1)/6}{(p-1)/6}} \pmod{p^2}.
\end{aligned}$$

For  $a = -\frac{5}{6}$  and  $p \equiv 5 \pmod{6}$  we have  $\langle a \rangle_p = \frac{p-5}{6}$  and  $\frac{a - \langle a \rangle_p}{p} = -\frac{1}{6}$ . Also,  $H_{2\langle a \rangle_p + 1} - H_{\langle a \rangle_p} = H_{\lfloor \frac{p}{3} \rfloor} - H_{\lfloor \frac{p}{6} \rfloor} \equiv 2q_p(2) \pmod{p}$ . Now taking  $a = -\frac{5}{6}$  in Corollary 2.1 and applying the above yields (2.20). The proof is now complete.

**Theorem 2.4.** *Let  $p$  be a prime with  $p > 3$ . Then*

$$\sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k (3k+1)} \equiv \begin{cases} \frac{2^{p-1} + 2}{3 \cdot 2^{\frac{p-1}{3}}} \pmod{p^2} & \text{if } p \equiv 1 \pmod{6}, \\ 2^{\frac{p+1}{3}} (2^{p-3} - 1) \pmod{p^2} & \text{if } p \equiv 5 \pmod{6}. \end{cases}$$

*Proof.* We first assume  $p \equiv 1 \pmod{6}$ . Set  $a = -\frac{1}{6}$  and  $b = \frac{1}{3}$ . Then  $\langle a \rangle_p = \frac{p-1}{6}$ ,  $\langle b \rangle_p = \frac{2p+1}{3}$ ,  $t = (a - \langle a \rangle_p)/p = -\frac{1}{6}$  and  $s = (b - \langle b \rangle_p)/p = -\frac{2}{3}$ . Since  $\langle a \rangle_p < \langle b \rangle_p <$

$p-1-\langle a \rangle_p$ , from Theorem 2.1 and (2.1)-(2.4),

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k (3k+1)} &\equiv \frac{\binom{(2p-2)/3}{(p-1)/6}}{2 \binom{(p-4)/3}{(p-1)/6}} \left( 1 + p - \frac{p}{3} H_{\frac{2p-2}{3}} + \frac{1}{2} p H_{\frac{p-1}{2}} - \frac{p}{6} H_{\frac{5p+1}{6}} \right) \\ &\equiv \frac{\binom{(2p-2)/3}{(p-1)/6}}{2 \binom{(p-4)/3}{(p-1)/6}} \left( 1 + p - \frac{p}{3} H_{\frac{p-1}{3}} + \frac{1}{2} p H_{\frac{p-1}{2}} - \frac{p}{6} \left( H_{\frac{p-1}{6}} - \frac{1}{(p-1)/6} \right) \right) \\ &\equiv \frac{\binom{(2p-2)/3}{(p-1)/6}}{\binom{(p-1)/3}{(p-1)/6}} \left( 1 - \frac{2}{3} p q_p(2) + \frac{3}{4} p q_p(3) \right) \pmod{p^2} \end{aligned}$$

By [S3, Lemmas 2.4 and 2.5], for  $k = 1, 2, \dots, \frac{p-1}{2}$ ,

$$\begin{aligned} \binom{\frac{p-1}{2} + k}{k} &\equiv (-1)^k \binom{\frac{p-1}{2}}{k} \left( 1 + 2p \sum_{i=1}^k \frac{1}{2i-1} \right) \\ &\equiv \binom{2k}{k} 4^{-k} \left( 1 - p \sum_{i=1}^k \frac{1}{2i-1} \right) \left( 1 + 2p \sum_{i=1}^k \frac{1}{2i-1} \right) \\ &\equiv \binom{2k}{k} 4^{-k} \left( 1 + p \sum_{i=1}^k \frac{1}{2i-1} \right) \pmod{p^2}. \end{aligned}$$

That is,

$$(2.21) \quad \binom{\frac{p-1}{2} + k}{k} \equiv \binom{2k}{k} 4^{-k} \left( 1 + p \left( H_{2k} - \frac{1}{2} H_k \right) \right) \pmod{p^2}.$$

Thus,

$$\frac{\binom{(2p-2)/3}{(p-1)/6}}{\binom{(p-1)/3}{(p-1)/6}} \equiv 4^{-\frac{p-1}{6}} \left( 1 + p \left( H_{\frac{p-1}{3}} - \frac{1}{2} H_{\frac{p-1}{6}} \right) \right) \equiv 2^{-\frac{p-1}{3}} \left( 1 + p q_p(2) - \frac{3}{4} p q_p(3) \right) \pmod{p^2}.$$

Therefore,

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k (3k+1)} &\equiv \left( 1 - \frac{2}{3} p q_p(2) + \frac{3}{4} p q_p(3) \right) 2^{-\frac{p-1}{3}} \left( 1 + p q_p(2) - \frac{3}{4} p q_p(3) \right) \\ &\equiv 2^{-\frac{p-1}{3}} \left( 1 + \frac{1}{3} p q_p(2) \right) = \frac{2^{p-1} + 2}{3 \cdot 2^{\frac{p-1}{3}}} \pmod{p^2}. \end{aligned}$$

Now assume  $p \equiv 5 \pmod{6}$ . Set  $a = -\frac{5}{6}$  and  $b = \frac{1}{3}$ . Then  $\langle a \rangle_p = \frac{p-5}{6}$ ,  $\langle b \rangle_p = \frac{p+1}{3}$ ,  $t = (a - \langle a \rangle_p)/p = -\frac{1}{6}$  and  $s = (b - \langle b \rangle_p)/p = -\frac{1}{3}$ . Since  $\langle a \rangle_p < \langle b \rangle_p < p-1 - \langle a \rangle_p$ , from Theorem 2.1 and (2.1)-(2.4),

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k (3k+1)} &\equiv \frac{3}{4} \cdot \frac{\binom{(p-2)/3}{(p-5)/6}}{\binom{(2p-4)/3}{(p-5)/6}} \left( 1 + 2p + \frac{p}{3} H_{\frac{p-2}{3}} + \frac{p}{6} H_{\frac{p+1}{6}} - \frac{p}{2} H_{\frac{p-1}{2}} \right) \end{aligned}$$

$$\begin{aligned}
&\equiv \frac{3}{4} \cdot \frac{\binom{(p-2)/3}{(p-5)/6}}{\binom{(2p-4)/3}{(p-5)/6}} \left( 1 + 2p + \frac{p}{3} \left( -\frac{3}{2}q_p(3) \right) + \frac{p}{6} \left( \frac{1}{(p+1)/6} - 2q_p(2) - \frac{3}{2}q_p(3) \right) - \frac{p}{2}(-2q_p(2)) \right) \\
&\equiv \frac{3}{4} \cdot \frac{2(p-2)}{p+1} \cdot \frac{\binom{(p-5)/3}{(p-5)/6}}{\binom{(2p-4)/3}{(p-5)/6}} \left( 1 + 3p + \frac{2}{3}pq_p(2) - \frac{3}{4}pq_p(3) \right) \pmod{p^2}.
\end{aligned}$$

By (2.21),

$$\begin{aligned}
\binom{\frac{2p-4}{3}}{\frac{p-5}{6}} &\equiv \binom{\frac{p-5}{3}}{\frac{p-5}{6}} 4^{-\frac{p-5}{6}} \left( 1 + p \left( H_{\frac{p-5}{3}} - \frac{1}{2}H_{\frac{p-5}{6}} \right) \right) \\
&\equiv \binom{\frac{p-5}{3}}{\frac{p-5}{6}} 4^{-\frac{p-5}{6}} \left( 1 + p \left( -\frac{3}{2}q_p(3) - \frac{1}{(p-2)/3} - \frac{1}{2} \left( -2q_p(2) - \frac{3}{2}q_p(3) \right) \right) \right) \\
&\equiv \binom{\frac{p-5}{3}}{\frac{p-5}{6}} 2^{-\frac{p-5}{3}} \left( 1 + p \left( \frac{3}{2} + q_p(2) - \frac{3}{4}q_p(3) \right) \right) \pmod{p^2}.
\end{aligned}$$

Hence

$$\begin{aligned}
&\sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k (3k+1)} \\
&\equiv \frac{3}{2} \cdot \frac{(p-2)(1-p)}{1-p^2} \cdot \frac{2^{(p-5)/3}}{1+p(\frac{3}{2} + q_p(2) - \frac{3}{4}q_p(3))} \left( 1 + p \left( 3 + \frac{2}{3}q_p(2) - \frac{3}{4}q_p(3) \right) \right) \\
&\equiv \frac{3}{2} (-2 + 3p) 2^{\frac{p-5}{3}} \left( 1 - p \left( \frac{3}{2} + q_p(2) - \frac{3}{4}q_p(3) \right) \right) \left( 1 + p \left( 3 + \frac{2}{3}q_p(2) - \frac{3}{4}q_p(3) \right) \right) \\
&\equiv \frac{3}{2} \cdot 2^{\frac{p-5}{3}} \left( -2 + \frac{2}{3}pq_p(2) \right) = 2^{\frac{p+1}{3}} (2^{p-3} - 1) \pmod{p^2}.
\end{aligned}$$

This completes the proof.

**Conjecture 2.1.** *Let  $p$  be a prime with  $p \equiv 1 \pmod{4}$ . Then*

$$\sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k (4k+1)} \equiv \begin{cases} (-1)^y \pmod{p^2} & \text{if } p = x^2 + 9y^2 \equiv 1 \pmod{12}, \\ (-3)^{\frac{p-1}{4}} \pmod{p} & \text{if } p \equiv 5 \pmod{12}. \end{cases}$$

**Conjecture 2.2.** *Let  $p$  be a prime with  $p > 3$ . Then*

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k (3k+1)} \equiv \begin{cases} 1 \pmod{p^2} & \text{if } p \equiv 1 \pmod{3}, \\ -\frac{5}{4} \pmod{p^2} & \text{if } p \equiv 2 \pmod{3} \end{cases}$$

and

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k (3k+2)} \equiv \begin{cases} \frac{1}{2} \pmod{p^2} & \text{if } p \equiv 1 \pmod{3}, \\ -\frac{2}{5} \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

**Conjecture 2.3.** *Let  $p$  be a prime with  $p \equiv 1 \pmod{6}$  and so  $p = x^2 + 3y^2$ . Then*

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k} \cdot \frac{p}{6k+1} \equiv (-1)^{\frac{p-1}{6}} \left( 4x^2 - 2p - \frac{p^2}{4x^2} \right) \pmod{p^3}.$$

### 3. The congruence for $\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{p}{k-a} \pmod{p^2}$

Let  $p$  be an odd prime. Replacing  $a$  with  $-1-a$  in Corollary 2.1 we obtain a congruence for  $\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{1}{k-a} \pmod{p^2}$  under the condition  $\frac{p}{2} < \langle a \rangle_p < p-1$ . Now we present the congruence for  $\sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{p}{k-a} \pmod{p^2}$  under the condition  $1 \leq \langle a \rangle_p < \frac{p}{2}$ . We note that for  $k = \langle a \rangle_p$ ,  $\binom{a}{k} \binom{-1-a}{k} \frac{1}{k-a} \notin \mathbb{Z}_p$ , but  $\binom{a}{k} \binom{-1-a}{k} \frac{p}{k-a} \in \mathbb{Z}_p$  provided that  $a - \langle a \rangle_p \not\equiv 0 \pmod{p^2}$ .

**Theorem 3.1.** *Let  $p$  be an odd prime,  $a \in \mathbb{Z}_p$ ,  $1 \leq \langle a \rangle_p < \frac{p}{2}$  and  $t = \frac{a - \langle a \rangle_p}{p} \not\equiv 0 \pmod{p}$ . Then*

$$\begin{aligned} \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{p}{k-a} &\equiv \frac{2t+1}{t} (-1)^{\langle a \rangle_p - 1} \binom{2\langle a \rangle_p}{\langle a \rangle_p} + 2 \binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} \\ &\equiv (-1)^{\langle a \rangle_p - 1} \binom{2\langle a \rangle_p}{\langle a \rangle_p} \left( \frac{1}{t} + 2p(H_{2\langle a \rangle_p} - H_{\langle a \rangle_p}) \right) \pmod{p^2}. \end{aligned}$$

Proof. Set  $S(a) = \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{p}{k-a}$ . From [G, (1.41)],

$$(3.1) \quad \sum_{k=0}^n \binom{n}{k} (-1)^k \frac{1}{k+x} = \frac{1}{x \binom{x+n}{n}} = \frac{(-1)^n}{(x+n) \binom{-x}{n}}.$$

Thus,  $\sum_{r=0}^k \binom{k}{r} (-1)^r \frac{p}{r-a} = \frac{(-1)^k p}{(k-a) \binom{a}{k}}$ . From [S7, Theorem 2.4] and the above,

$$\begin{aligned} &(-1)^{\langle a \rangle_p} S(a) \\ &\equiv \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{(-1)^k p}{(k-a) \binom{a}{k}} = \sum_{k=0}^{p-1} (-1)^k \binom{-1-a}{k} \frac{p}{k-a} \\ &= \sum_{\substack{k=0 \\ k \neq \langle a \rangle_p}}^{p-1} (-1)^k \binom{-1-a}{k} \frac{p}{k-a} + (-1)^{\langle a \rangle_p} \binom{-1-a}{\langle a \rangle_p} \frac{1}{(\langle a \rangle_p - a)/p} \\ &\equiv \sum_{\substack{k=0 \\ k \neq \langle a \rangle_p}}^{p-1-\langle a \rangle_p} (-1)^k \binom{p-1-\langle a \rangle_p}{k} \frac{p}{k-a} + (-1)^{\langle a \rangle_p} \binom{-1-a}{\langle a \rangle_p} \frac{1}{(\langle a \rangle_p - a)/p} \\ &= \sum_{k=0}^{p-1-\langle a \rangle_p} (-1)^k \binom{p-1-\langle a \rangle_p}{k} \frac{p}{k-a} + (-1)^{\langle a \rangle_p} \binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} \frac{1}{t} - (-1)^{\langle a \rangle_p} \binom{-1-a}{\langle a \rangle_p} \frac{1}{t} \\ &= \frac{(-1)^{p-1-\langle a \rangle_p} p}{(p-1-\langle a \rangle_p - a) \binom{a}{p-1-\langle a \rangle_p}} + \frac{(-1)^{\langle a \rangle_p}}{t} \left( \binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} - \binom{-1-a}{\langle a \rangle_p} \right) \pmod{p^2}. \end{aligned}$$

Applying (2.5) we see that

$$\begin{aligned} S(a) &\equiv -\frac{p}{(a + \langle a \rangle_p - p + 1) \binom{a}{p-1-\langle a \rangle_p}} + \frac{1}{t} \left( \binom{-1-a+p(t+1)}{\langle a \rangle_p} - \binom{-1-a}{\langle a \rangle_p} \right) \\ &\equiv -\frac{p}{(a + \langle a \rangle_p - p + 1) \binom{a}{p-1-\langle a \rangle_p}} + \frac{1}{t} \binom{-1-a}{\langle a \rangle_p} \sum_{i=0}^{\langle a \rangle_p - 1} \frac{p(t+1)}{-1-a-i} \end{aligned}$$

$$= -\frac{(p-1-\langle a \rangle_p)! \cdot p}{a(a-1) \cdots (a-(p-1-\langle a \rangle_p))} - \frac{t+1}{t} \binom{-1-a}{\langle a \rangle_p} \sum_{i=0}^{\langle a \rangle_p - 1} \frac{p}{1+a+i} \pmod{p^2}.$$

Since  $\langle a \rangle_p < \frac{p}{2}$ ,

$$\begin{aligned} & \frac{(p-1-\langle a \rangle_p)! \cdot p}{a(a-1) \cdots (a-(p-1-\langle a \rangle_p))} \\ &= \frac{(p-1-\langle a \rangle_p)! \cdot p}{(pt+\langle a \rangle_p)(pt+\langle a \rangle_p-1) \cdots (pt+1)pt(pt-1) \cdots (pt-(p-1-2\langle a \rangle_p))} \\ &\equiv \frac{(p-1-\langle a \rangle_p)!}{t \cdot \langle a \rangle_p! (1+pt \sum_{i=1}^{\langle a \rangle_p} \frac{1}{i}) \cdot (p-1-2\langle a \rangle_p)! (1-pt \sum_{i=1}^{p-1-2\langle a \rangle_p} \frac{1}{i})} \\ &\equiv \frac{1}{t} \binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} \left(1-pt \sum_{i=1}^{\langle a \rangle_p} \frac{1}{i}\right) \left(1+pt \sum_{i=1}^{p-1-2\langle a \rangle_p} \frac{1}{i}\right) \\ &\equiv \frac{1}{t} \binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} \left(1-ptH_{\langle a \rangle_p} + ptH_{p-1-2\langle a \rangle_p}\right) \pmod{p^2} \end{aligned}$$

and

$$\begin{aligned} & \binom{-1-a}{\langle a \rangle_p} \sum_{i=0}^{\langle a \rangle_p - 1} \frac{p}{1+a+i} \\ &\equiv \binom{-1-a}{\langle a \rangle_p} p \sum_{i=0}^{\langle a \rangle_p - 1} \frac{1}{\langle a \rangle_p + 1 + i} \equiv p \binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} (H_{2\langle a \rangle_p} - H_{\langle a \rangle_p}) \pmod{p^2}. \end{aligned}$$

Recall that  $H_{p-1-k} \equiv H_k \pmod{p}$ . From the above we deduce that

$$\begin{aligned} & S(a) \\ &\equiv -\frac{1}{t} \binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} (1+pt(H_{2\langle a \rangle_p} - H_{\langle a \rangle_p})) - \frac{p(t+1)}{t} \binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} (H_{2\langle a \rangle_p} - H_{\langle a \rangle_p}) \\ &= -\frac{1}{t} \binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} - p \frac{2t+1}{t} \binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} (H_{2\langle a \rangle_p} - H_{\langle a \rangle_p}) \pmod{p^2}. \end{aligned}$$

By (2.5),

$$\binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} = (-1)^{\langle a \rangle_p} \binom{2\langle a \rangle_p - p}{\langle a \rangle_p} \equiv (-1)^{\langle a \rangle_p} \binom{2\langle a \rangle_p}{\langle a \rangle_p} (1-p(H_{2\langle a \rangle_p} - H_{\langle a \rangle_p})) \pmod{p^2}.$$

Hence

$$\begin{aligned} & S(a) \\ &\equiv -\frac{1}{t} (-1)^{\langle a \rangle_p} \binom{2\langle a \rangle_p}{\langle a \rangle_p} (1-p(H_{2\langle a \rangle_p} - H_{\langle a \rangle_p})) - p \frac{2t+1}{t} (-1)^{\langle a \rangle_p} \binom{2\langle a \rangle_p}{\langle a \rangle_p} (H_{2\langle a \rangle_p} - H_{\langle a \rangle_p}) \\ &= -\frac{1}{t} (-1)^{\langle a \rangle_p} \binom{2\langle a \rangle_p}{\langle a \rangle_p} - 2p (-1)^{\langle a \rangle_p} \binom{2\langle a \rangle_p}{\langle a \rangle_p} (H_{2\langle a \rangle_p} - H_{\langle a \rangle_p}) \\ &\equiv \frac{1}{t} (-1)^{\langle a \rangle_p - 1} \binom{2\langle a \rangle_p}{\langle a \rangle_p} + 2 \left( \binom{p-1-\langle a \rangle_p}{\langle a \rangle_p} - (-1)^{\langle a \rangle_p} \binom{2\langle a \rangle_p}{\langle a \rangle_p} \right) \pmod{p^2}. \end{aligned}$$

This completes the proof.

**Theorem 3.2.** *Let  $p$  be a prime with  $p \equiv 1 \pmod{4}$ . Then*

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k} \cdot \frac{p}{4k+1} \equiv (-1)^{\frac{p-1}{4}} \frac{3-2^{p-1}}{2} \binom{\frac{p-1}{2}}{\frac{p-1}{4}} \pmod{p^2}.$$

Proof. By (2.2) and (2.4),  $H_{\frac{p-1}{2}} \equiv -2q_p(2) \pmod{p}$  and  $H_{[\frac{p}{4}]} \equiv -3q_p(2) \pmod{p}$ . Taking  $a = -\frac{1}{4}$  in Theorem 3.1 and noting that  $\langle a \rangle_p = \frac{p-1}{4}$  and  $t = -\frac{1}{4}$  gives

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k} \cdot \frac{p}{4k+1} &= \frac{1}{4} \sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k} \binom{-\frac{3}{4}}{k} \frac{p}{k - (-\frac{1}{4})} \\ &\equiv (-1)^{\frac{p-1}{4}} \binom{\frac{p-1}{4}}{\frac{p-1}{4}} \left(1 - \frac{p}{2} (H_{\frac{p-1}{2}} - H_{\frac{p-1}{4}})\right) \equiv (-1)^{\frac{p-1}{4}} \binom{\frac{p-1}{2}}{\frac{p-1}{4}} \left(1 - \frac{p}{2} q_p(2)\right) \pmod{p^2} \end{aligned}$$

This proves the theorem.

**Theorem 3.3.** *Let  $p$  be a prime with  $p \equiv 1 \pmod{3}$ . Then*

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{3k+1} \equiv \binom{2(p-1)/3}{(p-1)/3} \pmod{p^2}.$$

Proof. Taking  $a = -\frac{1}{3}$  in Theorem 3.1 and noting that  $\langle a \rangle_p = \frac{p-1}{3}$ ,  $t = -\frac{1}{3}$  and  $\binom{-1/3}{k} \binom{-2/3}{k} = \binom{2k}{k} \binom{3k}{k} 27^{-k}$  yields the result.

**Theorem 3.4.** *Let  $p$  be a prime with  $p \equiv 1 \pmod{6}$ . Then*

$$\sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k} \cdot \frac{p}{6k+1} \equiv \frac{5-2^p}{3} (-1)^{\frac{p-1}{6}} \binom{(p-1)/3}{(p-1)/6} \pmod{p^2}.$$

Proof. By (2.2) and (2.4),  $H_{[\frac{p}{3}]} \equiv -\frac{3}{2}q_p(3) \pmod{p}$  and  $H_{[\frac{p}{6}]} \equiv -2q_p(2) - \frac{3}{2}q_p(3) \pmod{p}$ . Thus, putting  $a = -\frac{1}{6}$  in Theorem 3.1 and noting that  $\langle a \rangle_p = \frac{p-1}{6}$  and  $t = -\frac{1}{6}$  gives

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k} \cdot \frac{p}{6k+1} &= \frac{1}{6} \sum_{k=0}^{p-1} \binom{-\frac{1}{6}}{k} \binom{-\frac{5}{6}}{k} \frac{p}{k - (-\frac{1}{6})} \\ &\equiv (-1)^{\frac{p-1}{6}} \binom{(p-1)/3}{(p-1)/6} \left(1 - \frac{p}{3} (H_{[\frac{p}{3}]} - H_{[\frac{p}{6}]})\right) \\ &\equiv (-1)^{\frac{p-1}{6}} \binom{(p-1)/3}{(p-1)/6} \left(1 - \frac{2}{3} p q_p(2)\right) \pmod{p^2}. \end{aligned}$$

This proves the theorem.



## 4. Congruences for $\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{p}{16^k(k+b)} \pmod{p^2}$

Let  $p$  be an odd prime and  $b \in \mathbb{Z}_p$ . From (2.10) we have a congruence for  $\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k(k+b)}$   $\pmod{p^2}$  under the condition  $1 \leq \langle b \rangle_p \leq \frac{p-3}{2}$ . Now we present a congruence for  $\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{p}{16^k(k+b)} \pmod{p^2}$  under the condition  $\langle b \rangle_p > \frac{p}{2}$ .

**Theorem 4.1.** *Let  $p$  be an odd prime,  $b \in \mathbb{Z}_p$ ,  $\langle b \rangle_p > \frac{p}{2}$  and  $s = (b - \langle b \rangle_p)/p \not\equiv -1 \pmod{p}$ . Then*

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k} \cdot \frac{p}{k+b} &\equiv \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^2}{16^k} \cdot \frac{p}{k+b} \\ &\equiv \frac{\binom{2(\langle b \rangle_p - \frac{p+1}{2})}{\langle b \rangle_p - \frac{p+1}{2}}^2}{(s+1)16^{\langle b \rangle_p - \frac{p+1}{2}}} (1 + p(2s+1)(H_{p-\langle b \rangle_p} - H_{\langle b \rangle_p - \frac{p+1}{2}})) \\ &\equiv \frac{1}{s+1} \left( \frac{p-1}{p - \langle b \rangle_p} \right)^2 \left( 1 + p(2q_p(2) + (2s+2)H_{p-\langle b \rangle_p} - (2s+1)H_{\langle b \rangle_p - \frac{p+1}{2}}) \right) \\ &\equiv \frac{\binom{2(p-\langle b \rangle_p)}{p-\langle b \rangle_p}^2}{(s+1)16^{p-\langle b \rangle_p}} \left( 1 + p(2s+2)(H_{p-\langle b \rangle_p} - H_{\langle b \rangle_p - \frac{p+1}{2}}) \right) \pmod{p^2}. \end{aligned}$$

Proof. From [G, (3.100)] we know that

$$(4.1) \quad \sum_{k=0}^n \binom{2k}{k} (-1)^k \binom{n+k}{2k} \frac{1}{k+b} = (-1)^n \frac{(b-1)(b-2)\cdots(b-n)}{b(b+1)\cdots(b+n)}.$$

From [S3, Lemma 2.2],

$$\binom{\frac{p-1}{2} + k}{2k} \equiv \frac{\binom{2k}{k}}{(-16)^k} \left( 1 - p^2 \sum_{i=1}^k \frac{1}{(2i-1)^2} \right) \pmod{p^4} \quad \text{for } k = 1, 2, \dots, \frac{p-1}{2}.$$

Thus,

$$(4.2) \quad \frac{\binom{2k}{k}}{(-16)^k} \equiv \left( 1 + p^2 \sum_{i=1}^k \frac{1}{(2i-1)^2} \right) \binom{\frac{p-1}{2} + k}{2k} \pmod{p^4} \quad \text{for } k = 1, 2, \dots, \frac{p-1}{2}.$$

Appealing to (4.1) and (4.2),

$$\begin{aligned} &\sum_{k=0}^{\frac{p-1}{2}} \frac{\binom{2k}{k}^2}{16^k} \cdot \frac{p}{k+b} \\ &\equiv \sum_{k=0}^{\frac{p-1}{2}} \binom{2k}{k} (-1)^k \binom{\frac{p-1}{2} + k}{2k} \left( 1 + p^2 \sum_{i=1}^k \frac{1}{(2i-1)^2} \right) \frac{p}{k+b} \\ &\equiv \sum_{k=0}^{\frac{p-1}{2}} \binom{2k}{k} (-1)^k \binom{\frac{p-1}{2} + k}{2k} \frac{p}{k+b} \end{aligned}$$

$$\begin{aligned}
& + p^2 \binom{2(p - \langle b \rangle_p)}{p - \langle b \rangle_p} (-1)^{p - \langle b \rangle_p} \binom{\frac{p-1}{2} + p - \langle b \rangle_p}{2(p - \langle b \rangle_p)} \left( \sum_{i=1}^{p - \langle b \rangle_p} \frac{1}{(2i-1)^2} \right) \frac{p}{p - \langle b \rangle_p + b} \\
& \equiv (-1)^{\frac{p-1}{2}} \frac{(b-1)(b-2) \cdots (b - \frac{p-1}{2})}{b(b+1) \cdots (b+p - \langle b \rangle_p - 1) \frac{b+p - \langle b \rangle_p}{p} (b+p - \langle b \rangle_p + 1) \cdots (b + \frac{p-1}{2})} \\
& + \frac{(-1)^{p - \langle b \rangle_p}}{s+1} \binom{2(p - \langle b \rangle_p)}{p - \langle b \rangle_p} \left( \frac{\binom{2(p - \langle b \rangle_p)}{p - \langle b \rangle_p}}{(-16)^{p - \langle b \rangle_p}} - \binom{\frac{p-1}{2} + p - \langle b \rangle_p}{2(p - \langle b \rangle_p)} \right) \pmod{p^3}.
\end{aligned}$$

That is,

$$\begin{aligned}
(4.3) \quad & \sum_{k=0}^{\frac{p-1}{2}} \frac{\binom{2k}{k}^2}{16^k} \cdot \frac{p}{k+b} \equiv \frac{(-1)^{p - \langle b \rangle_p} \binom{b-1}{\frac{p-1}{2}} \binom{\frac{p-1}{2}}{\langle b \rangle_p - \frac{p+1}{2}}}{(s+1) \binom{-1+p+b - \langle b \rangle_p}{p - \langle b \rangle_p} \binom{-1-(p+b - \langle b \rangle_p)}{\langle b \rangle_p - \frac{p+1}{2}}} \\
& + \frac{(-1)^{p - \langle b \rangle_p}}{s+1} \binom{2(p - \langle b \rangle_p)}{p - \langle b \rangle_p} \left( \frac{\binom{2(p - \langle b \rangle_p)}{p - \langle b \rangle_p}}{(-16)^{p - \langle b \rangle_p}} - \binom{\frac{p-1}{2} + p - \langle b \rangle_p}{2(p - \langle b \rangle_p)} \right) \pmod{p^3}.
\end{aligned}$$

From (4.3), (2.5) and (4.2),

$$\begin{aligned}
& \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^2}{16^k} \cdot \frac{p}{k+b} \\
& \equiv \frac{(-1)^{p - \langle b \rangle_p} \binom{b-1}{\frac{p-1}{2}} \binom{\frac{p-1}{2}}{\langle b \rangle_p - \frac{p+1}{2}}}{(s+1) \binom{-1+p+b - \langle b \rangle_p}{p - \langle b \rangle_p} \binom{-1-(p+b - \langle b \rangle_p)}{\langle b \rangle_p - \frac{p+1}{2}}} = \frac{(-1)^{p - \langle b \rangle_p} \binom{\langle b \rangle_p - 1 + ps}{\frac{p-1}{2}} \binom{\frac{p-1}{2}}{\langle b \rangle_p - \frac{p+1}{2}}}{(s+1) \binom{-1+p(s+1)}{p - \langle b \rangle_p} \binom{-1-p(s+1)}{\langle b \rangle_p - \frac{p+1}{2}}} \\
& \equiv \frac{(-1)^{p - \langle b \rangle_p}}{s+1} \cdot \frac{\binom{\langle b \rangle_p - 1}{\frac{p-1}{2}} (1 + ps \sum_{i=1}^{\frac{p-1}{2}} \frac{1}{\langle b \rangle_p - i}) \binom{\frac{p-1}{2}}{\langle b \rangle_p - \frac{p+1}{2}}}{\binom{-1}{p - \langle b \rangle_p} (1 + p(s+1) \sum_{i=1}^{p - \langle b \rangle_p} \frac{1}{-i}) \binom{-1}{\langle b \rangle_p - \frac{p+1}{2}} (1 - p(s+1) \sum_{i=1}^{\langle b \rangle_p - \frac{p+1}{2}} \frac{1}{-i})} \\
& = \frac{1}{s+1} (-1)^{\langle b \rangle_p - \frac{p+1}{2}} \frac{\binom{\langle b \rangle_p - 1}{\frac{p-1}{2}} (1 + ps(H_{\langle b \rangle_p - 1} - H_{\langle b \rangle_p - \frac{p+1}{2}})) \binom{\frac{p-1}{2}}{\langle b \rangle_p - \frac{p+1}{2}}}{(1 - p(s+1)H_{p - \langle b \rangle_p}) (1 + p(s+1)H_{\langle b \rangle_p - \frac{p+1}{2}})} \\
& \equiv \frac{1}{s+1} (-1)^{\langle b \rangle_p - \frac{p+1}{2}} \binom{\langle b \rangle_p - 1}{\frac{p-1}{2}} \binom{\frac{p-1}{2}}{\langle b \rangle_p - \frac{p+1}{2}} (1 + ps(H_{p - \langle b \rangle_p} - H_{\langle b \rangle_p - \frac{p+1}{2}})) \\
& \quad \times (1 + p(s+1)H_{p - \langle b \rangle_p}) (1 - p(s+1)H_{\langle b \rangle_p - \frac{p+1}{2}}) \\
& \equiv (-1)^{\langle b \rangle_p - \frac{p+1}{2}} \frac{\binom{\frac{p-1}{2} + \langle b \rangle_p - \frac{p+1}{2}}{\langle b \rangle_p - \frac{p+1}{2}} \binom{\frac{p-1}{2}}{\langle b \rangle_p - \frac{p+1}{2}}}{s+1} (1 + p(2s+1)(H_{p - \langle b \rangle_p} - H_{\langle b \rangle_p - \frac{p+1}{2}})) \\
& = (-1)^{\langle b \rangle_p - \frac{p+1}{2}} \frac{\binom{2(\langle b \rangle_p - \frac{p+1}{2})}{\langle b \rangle_p - \frac{p+1}{2}} \binom{\frac{p-1}{2} + \langle b \rangle_p - \frac{p+1}{2}}{2(\langle b \rangle_p - \frac{p+1}{2})}}{s+1} (1 + p(2s+1)(H_{p - \langle b \rangle_p} - H_{\langle b \rangle_p - \frac{p+1}{2}})) \\
& \equiv \frac{\binom{2(\langle b \rangle_p - \frac{p+1}{2})}{\langle b \rangle_p - \frac{p+1}{2}}^2}{(s+1)16^{\langle b \rangle_p - \frac{p+1}{2}}} (1 + p(2s+1)(H_{p - \langle b \rangle_p} - H_{\langle b \rangle_p - \frac{p+1}{2}})) \pmod{p^2}.
\end{aligned}$$

By [S3, Lemma 2.4], for  $k = 1, 2, \dots, \frac{p-1}{2}$ ,

$$(4.4) \quad \binom{\frac{p-1}{2}}{k} \equiv \frac{\binom{2k}{k}}{(-4)^k} \left(1 - p \sum_{i=1}^k \frac{1}{2i-1}\right) \equiv \frac{\binom{2k}{k}}{(-4)^k} \left(1 - p(H_{2k} - \frac{1}{2}H_k)\right) \pmod{p^2}.$$

Thus,

$$(4.5) \quad \left(\binom{\frac{p-1}{2}}{k}\right)^2 \equiv \frac{\binom{2k}{k}^2}{16^k} \left(1 - p(H_{2k} - \frac{1}{2}H_k)\right)^2 \equiv \frac{\binom{2k}{k}^2}{16^k} (1 - 2pH_{2k} + pH_k) \pmod{p^2}$$

and so

$$(4.6) \quad \frac{\binom{2k}{k}^2}{16^k} \equiv \left(\binom{\frac{p-1}{2}}{k}\right)^2 (1 + 2pH_{2k} - pH_k) \pmod{p^2}.$$

Now, from the above and (2.3) we deduce that

$$\begin{aligned} & \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k}^2}{16^k} \cdot \frac{p}{k+b} \\ & \equiv \frac{1}{s+1} \left(\binom{\frac{p-1}{2}}{\langle b \rangle_p - \frac{p+1}{2}}\right)^2 (1 + 2pH_{2\langle b \rangle_p - p-1} - pH_{\langle b \rangle_p - \frac{p+1}{2}}) \\ & \quad \times (1 + p(2s+1)(H_{p-\langle b \rangle_p} - H_{\langle b \rangle_p - \frac{p+1}{2}})) \\ & \equiv \frac{1}{s+1} \left(\binom{\frac{p-1}{2}}{p - \langle b \rangle_p}\right)^2 (1 + p(2H_{2(p-\langle b \rangle_p)} + (2s+1)H_{p-\langle b \rangle_p} - (2s+2)H_{\langle b \rangle_p - \frac{p+1}{2}})) \\ & \equiv \frac{1}{s+1} \left(\binom{\frac{p-1}{2}}{p - \langle b \rangle_p}\right)^2 (1 + p(2q_p(2) + (2s+2)H_{p-\langle b \rangle_p} - (2s+1)H_{\langle b \rangle_p - \frac{p+1}{2}})) \\ & \equiv \frac{1}{(s+1)16^{p-\langle b \rangle_p}} \left(\frac{2(p-\langle b \rangle_p)}{p-\langle b \rangle_p}\right)^2 (1 - 2pH_{2p-2\langle b \rangle_p} + pH_{p-\langle b \rangle_p}) \\ & \quad \times (1 + p(2H_{2(p-\langle b \rangle_p)} + (2s+1)H_{p-\langle b \rangle_p} - (2s+2)H_{\langle b \rangle_p - \frac{p+1}{2}})) \\ & \equiv \frac{\binom{2(p-\langle b \rangle_p)}{p-\langle b \rangle_p}^2}{(s+1)16^{p-\langle b \rangle_p}} (1 + p(2s+2)(H_{p-\langle b \rangle_p} - H_{\langle b \rangle_p - \frac{p+1}{2}})) \pmod{p^2}. \end{aligned}$$

To complete the proof, we note that  $p \mid \binom{2k}{k}$  for  $\frac{p}{2} < k < p$ .

**Corollary 4.1.** *Let  $p$  be an odd prime,  $b \in \mathbb{Z}_p$ ,  $\langle b \rangle_p > \frac{p}{2}$  and  $s = (b - \langle b \rangle_p)/p \not\equiv -1, -\frac{1}{2} \pmod{p}$ . Then*

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k} \cdot \frac{p}{k + \frac{1}{2} - b} \equiv -\frac{2s+2}{2s+1} \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k} \cdot \frac{p}{k+b} \pmod{p^2}.$$

*Proof.* Set  $b' = \frac{1}{2} - b$  and  $s' = (b' - \langle b' \rangle_p)/p$ . Then  $\langle b' \rangle_p = p + \frac{p+1}{2} - \langle b \rangle_p > \frac{p}{2}$  and  $s' = -\frac{3}{2} - s$ . By Theorem 4.1,

$$(s'+1) \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k} \cdot \frac{p}{k+b'}$$

$$\begin{aligned}
&\equiv \binom{\frac{p-1}{2}}{p-\langle b' \rangle_p}^2 \left( 1 + p(2q_p(2) + (2s' + 2)H_{p-\langle b' \rangle_p} - (2s + 1)H_{\langle b' \rangle_p - \frac{p+1}{2}}) \right) \\
&= \binom{\frac{p-1}{2}}{p-\langle b \rangle_p}^2 \left( 1 + p(2q_p(2) + (2s + 2)H_{p-\langle b \rangle_p} - (2s + 1)H_{\langle b \rangle_p - \frac{p+1}{2}}) \right) \\
&\equiv (s + 1) \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k} \cdot \frac{p}{k+b} \pmod{p^2}.
\end{aligned}$$

This yields the result.

**Theorem 4.2.** *Let  $p$  be a prime with  $p \equiv 1 \pmod{3}$  and so  $p = x^2 + 3y^2$ . Then*

$$\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{3k+1} \equiv \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{6k+1} \equiv 4x^2 - 2p \pmod{p^2}.$$

Proof. Putting  $b = \frac{1}{3}$  in Theorem 4.1 and noting that  $\langle b \rangle_p = \frac{2p+1}{3}$  and  $s = -\frac{2}{3}$  gives

$$\begin{aligned}
&\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{3k+1} \\
&\equiv \left( \frac{\frac{p-1}{2}}{\frac{p-1}{3}} \right)^2 \left( 1 + p(2q_p(2) + \frac{2}{3}H_{\frac{p-1}{3}} + \frac{1}{3}H_{\frac{p-1}{6}}) \right) \\
&\equiv \left( \frac{\frac{p-1}{2}}{\frac{p-1}{3}} \right)^2 \left( 1 + p \left( 2q_p(2) + \frac{2}{3} \left( -\frac{3}{2}q_p(3) \right) + \frac{1}{3} \left( -2q_p(2) - \frac{3}{2}q_p(3) \right) \right) \right) \\
&= \left( \frac{\frac{p-1}{2}}{\frac{p-1}{6}} \right)^2 \left( 1 + p \left( \frac{4}{3}q_p(2) - \frac{3}{2}q_p(3) \right) \right) \pmod{p^2}.
\end{aligned}$$

By [Y] or [BEW, Theorem 9.4.4],

$$\left( \frac{\frac{p-1}{2}}{\frac{p-1}{6}} \right) \equiv \left( 2x - \frac{p}{2x} \right) \left( -1 + \frac{2}{3}pq_p(2) - \frac{3}{4}pq_p(3) \right) \pmod{p^2}.$$

Thus,

$$\left( \frac{\frac{p-1}{2}}{\frac{p-1}{6}} \right)^2 \equiv (4x^2 - 2p) \left( 1 - \frac{4}{3}pq_p(2) + \frac{3}{2}pq_p(3) \right) \pmod{p^2}.$$

Hence

$$\begin{aligned}
&\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{3k+1} \\
&\equiv (4x^2 - 2p) \left( 1 - p \left( \frac{4}{3}q_p(2) - \frac{3}{2}q_p(3) \right) \right) \left( 1 + p \left( \frac{4}{3}q_p(2) - \frac{3}{2}q_p(3) \right) \right) \equiv 4x^2 - 2p \pmod{p^2}.
\end{aligned}$$

Taking  $b = \frac{1}{3}$  in Corollary 4.1 gives

$$\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{6k+1} \equiv \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{3k+1} \pmod{p^2}.$$

Thus, the theorem is proved.

**Theorem 4.3.** *Let  $p$  be a prime with  $p > 3$ ,  $p \equiv 1, 3 \pmod{8}$  and so  $p = x^2 + 2y^2$ . Then*

$$\begin{aligned} (2 - (-1)^{\frac{p-1}{2}}) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{8k+1} &\equiv (2 + (-1)^{\frac{p-1}{2}}) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{8k+3} \\ &\equiv 4x^2 - 2p \pmod{p^2}. \end{aligned}$$

Proof. Since  $\frac{3}{8} = \frac{1}{2} - \frac{1}{8}$ , from Corollary 4.1 we only need to prove the congruence for  $\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{8k+1} \pmod{p^2}$ . Set  $b = \frac{1}{8}$ . For  $p \equiv 1 \pmod{8}$ ,  $\langle b \rangle_p = \frac{7p+1}{8} > \frac{p}{2}$ ,  $p - \langle b \rangle_p = \frac{p-1}{8}$ ,  $\langle b \rangle_p - \frac{p+1}{2} = \frac{3(p-1)}{8}$  and  $s = \frac{b - \langle b \rangle_p}{p} = -\frac{7}{8}$ . By Theorem 4.1,

$$\frac{1}{8} \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{k+1/8} \equiv \left( \frac{p-1}{[p/8]} \right)^2 \left( 1 + p \left( 2q_p(2) + \frac{3}{4} H_{[\frac{3p}{8}]} + \frac{1}{4} H_{[\frac{p}{8}]} \right) \right) \pmod{p^2}.$$

For  $p \equiv 3 \pmod{8}$ ,  $\langle b \rangle_p = \frac{5p+1}{8} > \frac{p}{2}$ ,  $p - \langle b \rangle_p = [\frac{3p}{8}]$ ,  $\langle b \rangle_p - \frac{p+1}{2} = [\frac{p}{8}]$  and  $s = \frac{b - \langle b \rangle_p}{p} = -\frac{5}{8}$ . By Theorem 4.1,

$$\frac{3}{8} \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{k+1/8} \equiv \left( \frac{p-1}{[p/8]} \right)^2 \left( 1 + p \left( 2q_p(2) + \frac{3}{4} H_{[\frac{3p}{8}]} + \frac{1}{4} H_{[\frac{p}{8}]} \right) \right) \pmod{p^2}.$$

By (2.3),  $H_{[\frac{p}{8}]} + H_{[\frac{3p}{8}]} \equiv 2H_{[\frac{p}{4}]} - 2q_p(2) \equiv -8q_p(2) \pmod{p}$ . Hence, for  $p \equiv 1, 3 \pmod{8}$ ,

$$\begin{aligned} (2 - (-1)^{\frac{p-1}{2}}) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{8k+1} &\equiv \left( \frac{p-1}{[\frac{p}{8}]} \right)^2 \left( 1 + p \left( 2q_p(2) + \frac{3}{4} (-8q_p(2)) - \frac{2}{4} H_{[\frac{p}{8}]} \right) \right) \\ &= \left( \frac{p-1}{[\frac{p}{8}]} \right)^2 \left( 1 - p \left( 4q_p(2) + \frac{1}{2} H_{[\frac{p}{8}]} \right) \right) \pmod{p^2}. \end{aligned}$$

On the other hand, taking  $n = [\frac{p}{8}]$  and  $t = -\frac{1}{4}$  in [S7, Theorem 3.2(ii)] gives

$$\begin{aligned} \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{128^k} &= \sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k} \binom{-\frac{3}{4}}{k} \frac{1}{2^k} \equiv \left( \frac{p-1}{[p/8]} \right) \left( 1 + p \left( \frac{3}{4} H_{[\frac{p}{4}]} - \frac{1}{4} H_{[\frac{p}{8}]} + \frac{1}{4} q_p(2) \right) \right) \\ &\equiv \left( \frac{p-1}{[p/8]} \right) \left( 1 - p \left( 2q_p(2) + \frac{1}{4} H_{[\frac{p}{8}]} \right) \right) \pmod{p^2}. \end{aligned}$$

From [S5, Theorem 4.3],

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{128^k} \equiv (-1)^{[\frac{p}{8}] + \frac{p-1}{2}} \left( 2x - \frac{p}{2x} \right) \pmod{p^2},$$

where  $x \equiv 1 \pmod{4}$ . Thus,

$$4x^2 - 2p \equiv \left( 2x - \frac{p}{2x} \right)^2 \equiv \left( \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{128^k} \right)^2 \equiv \left( \frac{p-1}{[p/8]} \right)^2 \left( 1 - p \left( 2q_p(2) + \frac{1}{4} H_{[\frac{p}{8}]} \right) \right)^2$$

$$\begin{aligned}
&\equiv \left(\frac{p-1}{[p/8]}\right)^2 \left(1 - 2p\left(2q_p(2) + \frac{1}{4}H_{[\frac{p}{8}]}\right)\right) \\
&\equiv (2 - (-1)^{\frac{p-1}{2}}) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{8k+1} \pmod{p^2}.
\end{aligned}$$

This proves the theorem.

**Theorem 4.4.** *Let  $p$  be a prime,  $p \equiv 1 \pmod{4}$  and so  $p = x^2 + y^2$  with  $2 \nmid x$ . Then*

$$\begin{aligned}
\left(3\binom{p}{3} - 2\right) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{12k+1} &\equiv \left(3\binom{p}{3} + 2\right) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{12k+5} \\
&\equiv 4x^2 - 2p \pmod{p^2}.
\end{aligned}$$

Proof. Since  $\frac{5}{12} = \frac{1}{2} - \frac{1}{12}$ , from Corollary 4.1 we only need to prove the congruence for  $\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{12k+1} \pmod{p^2}$ . Set  $b = \frac{1}{12}$ . For  $p \equiv 1 \pmod{12}$ ,  $\langle b \rangle_p = \frac{11p+1}{12} > \frac{p}{2}$ ,  $p - \langle b \rangle_p = \frac{p-1}{12}$ ,  $\langle b \rangle_p - \frac{p+1}{2} = \frac{5(p-1)}{12}$  and  $s = \frac{b - \langle b \rangle_p}{p} = -\frac{11}{12}$ . By Theorem 4.1,

$$\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{12k+1} \equiv \left(\frac{p-1}{[\frac{p}{12}]}\right)^2 \left(1 + p\left(2q_p(2) + \frac{1}{6}H_{[\frac{p}{12}]} + \frac{5}{6}H_{[\frac{5p}{12}]}\right)\right) \pmod{p^2}.$$

For  $p \equiv 5 \pmod{12}$ ,  $\langle b \rangle_p = \frac{7p+1}{12} > \frac{p}{2}$ ,  $p - \langle b \rangle_p = \frac{5p-1}{12}$ ,  $\langle b \rangle_p - \frac{p+1}{2} = \frac{p-5}{12}$  and  $s = \frac{b - \langle b \rangle_p}{p} = -\frac{7}{12}$ . By Theorem 4.1,

$$\left(1 - \frac{7}{12}\right) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{k+1/12} \equiv \left(\frac{p-1}{[\frac{p}{12}]}\right)^2 \left(1 + p\left(2q_p(2) + \frac{1}{6}H_{[\frac{p}{12}]} + \frac{5}{6}H_{[\frac{5p}{12}]}\right)\right) \pmod{p^2}.$$

Since  $[\frac{p}{12}] + [\frac{5p}{12}] = \frac{p-1}{2}$ , from (2.3) and (2.4),

$$(4.7) \quad H_{[\frac{5p}{12}]} + H_{[\frac{p}{12}]} \equiv 2H_{[\frac{p}{6}]} - 2q_p(2) \equiv -6q_p(2) - 3q_p(3) \pmod{p}.$$

Thus,

$$\begin{aligned}
2q_p(2) + \frac{1}{6}H_{[\frac{p}{12}]} + \frac{5}{6}H_{[\frac{5p}{12}]} &\equiv 2q_p(2) - \frac{4}{6}H_{[\frac{p}{12}]} + \frac{5}{6}(-6q_p(2) - 3q_p(3)) \\
&= -3q_p(2) - \frac{5}{2}q_p(3) - \frac{2}{3}H_{[\frac{p}{12}]} \pmod{p^2}.
\end{aligned}$$

On the other hand, putting  $n = [\frac{p}{12}]$  and  $t = -\frac{1}{6}$  in [S7, Theorem 3.1(ii)] we deduce that

$$\sum_{k=0}^{p-1} \frac{\binom{3k}{k}\binom{6k}{3k}}{864^k} = \sum_{k=0}^{p-1} \binom{-\frac{1}{6}}{k} \binom{-\frac{5}{6}}{k} \frac{1}{2^k} \equiv \left(\frac{p-1}{[\frac{p}{12}]}\right) \left(1 + p\left(\frac{5}{6}H_{[\frac{p}{6}]} - \frac{1}{3}H_{[\frac{p}{12}]} + \frac{1}{6}q_p(2)\right)\right) \pmod{p^2}.$$

By [S6, Theorem 3.2],  $\sum_{k=0}^{p-1} \frac{\binom{3k}{k}\binom{6k}{3k}}{864^k} \equiv 2c - \frac{p}{2c} \pmod{p^2}$ , where

$$(4.8) \quad c = \begin{cases} x & \text{if } p \equiv 1 \pmod{12} \text{ and } 3 \nmid x, \\ -x & \text{if } p \equiv 1 \pmod{12} \text{ and } 3 \mid x, \\ y & \text{if } p \equiv 5 \pmod{12} \text{ and } y \equiv x \pmod{3} \end{cases}$$

and  $x \equiv 1 \pmod{4}$ . By (2.4),  $H_{[\frac{p}{6}]} \equiv -2q_p(2) - \frac{3}{2}q_p(3) \pmod{p}$ . Thus,

$$(4.9) \quad 2c - \frac{p}{2c} \equiv \left( \frac{\frac{p-1}{2}}{[\frac{p}{12}]} \right) \left( 1 - p \left( \frac{3}{2}q_p(2) + \frac{5}{4}q_p(3) + \frac{1}{3}H_{[\frac{p}{12}]} \right) \right) \pmod{p^2}$$

and so

$$\begin{aligned} 4c^2 - 2p &\equiv \left( \frac{\frac{p-1}{2}}{[\frac{p}{12}]} \right)^2 \left( 1 - p \left( 3q_p(2) + \frac{5}{2}q_p(3) + \frac{2}{3}H_{[\frac{p}{12}]} \right) \right) \pmod{p^2} \\ &\equiv \begin{cases} \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{12k+1} \pmod{p^2} & \text{if } p \equiv 1 \pmod{12}, \\ 5 \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{12k+1} \pmod{p^2} & \text{if } p \equiv 5 \pmod{12}. \end{cases} \end{aligned}$$

To complete the proof we note that  $4c^2 - 2p = 4y^2 - 2p \equiv -(4x^2 - 2p) \pmod{p^2}$  for  $p \equiv 5 \pmod{12}$ .

Based on calculations with Maple, we pose the following conjectures.

**Conjecture 4.1.** *Let  $p$  be a prime such that  $p \equiv 1 \pmod{6}$  and so  $p = x^2 + 3y^2$ .*

*Then*

$$\sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{3k+1} \equiv \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{6k+1} \equiv 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3}.$$

**Conjecture 4.2.** *Let  $p$  be a prime with  $p > 3$ ,  $p \equiv 1, 3 \pmod{8}$  and so  $p = x^2 + 2y^2$ .*

*Then*

$$\begin{aligned} (2 - (-1)^{\frac{p-1}{2}}) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{8k+1} &\equiv (2 + (-1)^{\frac{p-1}{2}}) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{8k+3} \\ &\equiv 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3}. \end{aligned}$$

**Conjecture 4.3.** *Let  $p$  be a prime,  $p \equiv 1 \pmod{4}$  and so  $p = x^2 + y^2$  with  $2 \nmid x$ .*

*Then*

$$\begin{aligned} \left( 3 \binom{p}{3} - 2 \right) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{12k+1} &\equiv \left( 3 \binom{p}{3} + 2 \right) \sum_{k=0}^{p-1} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{12k+5} \\ &\equiv 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3}. \end{aligned}$$

## 5. Congruences for Domb and Almkvist-Zudilin numbers

This section is devoted to proving some conjectures on congruences for Domb and Almkvist-Zudilin numbers.

**Theorem 5.1.** *Let  $p$  be a prime with  $p > 3$ . Then*

$$\sum_{n=0}^{p-1} \frac{D_n}{16^n} \equiv \sum_{n=0}^{p-1} \frac{D_n}{4^n} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p \equiv 1 \pmod{3} \text{ and so } p = x^2 + 3y^2, \\ 0 \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

Proof. From [S8] we know that

$$(5.1) \quad D_n = (-1)^n \sum_{k=0}^n \binom{2k}{k}^2 \binom{3k}{k} \binom{n+2k}{3k} (-16)^{n-k}.$$

From [G, (1.52)],

$$(5.2) \quad \sum_{n=r}^m \binom{n}{r} = \binom{m+1}{r+1}.$$

Thus,

$$\begin{aligned} \sum_{n=0}^{p-1} \frac{D_n}{16^n} &= \sum_{n=0}^{p-1} \sum_{k=0}^n \binom{2k}{k}^2 \binom{3k}{k} \binom{n+2k}{3k} \frac{1}{(-16)^k} \\ &= \sum_{k=0}^{p-1} \binom{2k}{k}^2 \binom{3k}{k} \frac{1}{(-16)^k} \sum_{n=k}^{p-1} \binom{n+2k}{3k} = \sum_{k=0}^{p-1} \binom{2k}{k}^2 \binom{3k}{k} \frac{1}{(-16)^k} \binom{p+2k}{3k+1}. \end{aligned}$$

If  $\frac{2p}{3} < k < p$ , then  $\binom{2k}{k}^2 \binom{3k}{k} = \frac{(2k)!(3k)!}{k!^5} \equiv 0 \pmod{p^3}$ . If  $\frac{p}{2} < k < \frac{2p}{3}$ , then  $3k+2 \leq 2p < p+2k$  and so

$$\binom{p+2k}{3k+1} = \binom{p+2k}{p-1-k} = \frac{(p+2k)(p+2k-1)\cdots(3k+2)}{(p-1-k)!} \equiv 0 \pmod{p}.$$

Hence  $p^3 \mid \binom{2k}{k}^2 \binom{p+2k}{3k+1}$  and therefore

$$(5.3) \quad \begin{aligned} \sum_{n=0}^{p-1} \frac{D_n}{16^n} &= \sum_{k=0}^{p-1} \binom{2k}{k}^2 \binom{3k}{k} \frac{1}{(-16)^k} \binom{p+2k}{3k+1} \\ &\equiv \sum_{k=0}^{(p-1)/2} \binom{2k}{k}^2 \binom{3k}{k} \frac{1}{(-16)^k} \binom{p+2k}{3k+1} \pmod{p^3}. \end{aligned}$$

For  $1 \leq k < \frac{p}{2}$  with  $k \neq \frac{p-1}{3}$ ,

$$\begin{aligned} \binom{3k}{k} \binom{p+2k}{3k+1} &= \frac{(p+2k)\cdots(p+1)p(p-1)\cdots(p-k)}{k!(2k)!(3k+1)} \\ &= \frac{p}{3k+1} \cdot \frac{(p^2-1^2)\cdots(p^2-k^2)(p+k+1)\cdots(p+2k)}{k!(2k)!} \\ &\equiv \frac{p}{3k+1} \cdot \frac{(-1^2)\cdots(-k^2)(k+1)\cdots 2k}{k!(2k)!} \left(1 + p\left(\frac{1}{k+1} + \cdots + \frac{1}{2k}\right)\right) \\ &= \frac{(-1)^k}{3k+1} (p + p^2(H_{2k} - H_k)) \pmod{p^3}. \end{aligned}$$



For  $k = \frac{p-1}{3}$  we see that

$$\begin{aligned}
\binom{3k}{k} \binom{p+2k}{3k+1} &= \binom{p-1}{\frac{2(p-1)}{3}} \binom{p+\frac{2(p-1)}{3}}{p} \\
&= \frac{(p-1)(p-2)\cdots(p-\frac{2(p-1)}{3})}{\frac{2(p-1)}{3}!} \cdot \frac{(p+\frac{2(p-1)}{3})\cdots(p+1)p(p-1)\cdots(\frac{2(p-1)}{3}+1)}{p!} \\
&= \frac{(p^2-1^2)(p^2-2^2)\cdots(p^2-(\frac{2(p-1)}{3})^2)}{\frac{2(p-1)}{3}!^2} \equiv 1 - p^2 \sum_{i=1}^{2(p-1)/3} \frac{1}{i^2} \pmod{p^4}.
\end{aligned}$$

Let  $\{U_n\}$  be given by

$$(5.4) \quad U_0 = 1, \quad U_n = -2 \sum_{k=1}^{\lfloor n/2 \rfloor} \binom{n}{2k} U_{n-2k} \quad (n \geq 1).$$

Using [S1, Theorem 5.1(a)] and [S4, Theorem 3.3],

$$\sum_{i=1}^{2(p-1)/3} \frac{1}{i^2} \equiv \sum_{i=1}^{2(p-1)/3} \frac{1}{(p-i)^2} = \sum_{x=\frac{p-1}{3}+1}^{p-1} \frac{1}{x^2} = \sum_{x=1}^{p-1} \frac{1}{x^2} - \sum_{x=1}^{(p-1)/3} \frac{1}{x^2} \equiv 0 - 3U_{p-3} \pmod{p}.$$

From [S4, Theorem 3.2(iii)],

$$\sum_{i=\frac{p-1}{3}+1}^{2(p-1)/3} \frac{1}{i} = \sum_{i=1}^{2(p-1)/3} \frac{1}{i} - \sum_{i=1}^{(p-1)/3} \frac{1}{i} = \sum_{i=1}^{2(p-1)/3} \frac{(-1)^{i-1}}{i} \equiv 3pU_{p-3} \pmod{p^2}.$$

Hence for  $p \equiv 1 \pmod{3}$ ,

$$\binom{p-1}{\frac{p-1}{3}} \binom{p+\frac{2(p-1)}{3}}{p} \equiv 1 - p^2 \sum_{i=1}^{2(p-1)/3} \frac{1}{i^2} \equiv 1 + 3p^2 U_{p-3} \equiv 1 + p(H_{\frac{2(p-1)}{3}} - H_{\frac{p-1}{3}}) \pmod{p^3}.$$

Therefore, from (5.3) and the above congruence for  $\binom{3k}{k} \binom{p+2k}{3k+1} \pmod{p^3}$  we derive that

$$(5.5) \quad \sum_{n=0}^{p-1} \frac{D_n}{16^n} \equiv \sum_{k=0}^{(p-1)/2} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{3k+1} (1 + p(H_{2k} - H_k)) \pmod{p^3}.$$

Note that  $H_{\frac{2(p-1)}{3}} = H_{p-1-\frac{p-1}{3}} \equiv H_{\frac{p-1}{3}} \pmod{p}$ . We get  $\sum_{n=0}^{p-1} \frac{D_n}{16^n} \equiv \sum_{k=0}^{(p-1)/2} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{3k+1} \pmod{p^2}$ . Since

$$\binom{p-1}{\frac{p-1}{3}} \binom{p+\frac{p-1}{3}}{p} = \frac{(p^2-1^2)(p^2-2^2)\cdots(p^2-(\frac{p-1}{3})^2)}{(\frac{p-1}{3}!)^2} \equiv 1 \pmod{p^2},$$

from [S10, p.137] we know that  $\sum_{n=0}^{p-1} \frac{D_n}{4^n} \equiv \sum_{k=0}^{(p-1)/2} \binom{2k}{k}^2 \frac{1}{16^k} \cdot \frac{p}{3k+1} \pmod{p^2}$ . Now applying (2.11) and Theorem 4.2 yields the result.

**Remark 5.1** Suppose that  $p$  is a prime with  $p > 3$ . In [S10] the author proved that  $\sum_{n=0}^{p-1} \frac{D_n}{4^n} \equiv 0 \pmod{p^2}$  for  $p \equiv 2 \pmod{3}$ .

Let  $\{b_n\}$  be Almkvist-Zudilin numbers. From [S8, (5.1) and (5.2)] we know that

$$(5.6) \quad \begin{aligned} b_n &= \sum_{k=0}^{\lfloor n/3 \rfloor} \binom{2k}{k}^2 \binom{4k}{2k} \binom{n+k}{4k} (-3)^{n-3k} \\ &= \sum_{k=0}^n \binom{2k}{k}^2 \binom{4k}{2k} \binom{n+3k}{4k} (-27)^{n-k}. \end{aligned}$$

**Lemma 5.1.** *Suppose that  $p$  is a prime with  $p > 3$ . Then*

$$\sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \equiv \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{4k+1} \pmod{p^2}.$$

Proof. Using (5.6) and (5.2),

$$\begin{aligned} \sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} &= \sum_{n=0}^{p-1} \sum_{k=0}^{\lfloor n/3 \rfloor} \binom{2k}{k}^2 \binom{4k}{2k} \binom{n+k}{4k} \frac{1}{(-27)^k} \\ &= \sum_{k=0}^{\lfloor p/3 \rfloor} \binom{2k}{k}^2 \binom{4k}{2k} \frac{1}{(-27)^k} \sum_{n=3k}^{p-1} \binom{n+k}{4k} \\ &= \sum_{k=0}^{\lfloor p/3 \rfloor} \binom{2k}{k}^2 \binom{4k}{2k} \frac{1}{(-27)^k} \binom{p+k}{4k+1} \\ &= \sum_{k=0}^{\lfloor p/3 \rfloor} \frac{(p+k)(p+k-1) \cdots (p+1)p(p-1) \cdots (p-3k)}{(4k+1) \cdot (-27)^k \cdot k!^4} \\ &= \sum_{k=0}^{\lfloor p/3 \rfloor} \frac{p}{4k+1} \cdot \frac{(p^2-1^2)(p^2-2^2) \cdots (p^2-k^2) \cdot (p-(k+1)) \cdots (p-3k)}{k!^4 \cdot (-27)^k}. \end{aligned}$$

Thus,

$$\begin{aligned} &\sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \\ &\equiv \sum_{k=0}^{\lfloor p/3 \rfloor} \frac{p}{(4k+1)(-27)^k \cdot k!^4} \cdot (-1^2)(-2^2) \cdots (-k^2) \left(1 - p^2 \sum_{i=1}^k \frac{1}{i^2}\right) \\ &\quad \times (-(k+1)) \cdots (-3k) \left(1 - p \sum_{i=k+1}^{3k} \frac{1}{i} + p^2 \sum_{k+1 \leq i < j \leq 3k} \frac{1}{ij}\right) \\ &= \sum_{k=0}^{\lfloor p/3 \rfloor} \frac{p}{4k+1} \cdot \frac{(3k)!}{k!^3 \cdot 27^k} \left(1 - p^2 \sum_{i=1}^k \frac{1}{i^2}\right) \left(1 - p \sum_{i=k+1}^{3k} \frac{1}{i} + \frac{p^2}{2} \left( \left( \sum_{i=k+1}^{3k} \frac{1}{i} \right)^2 - \sum_{i=k+1}^{3k} \frac{1}{i^2} \right)\right) \\ &\equiv \sum_{k=0}^{\lfloor p/3 \rfloor} \frac{p}{4k+1} \cdot \frac{(3k)!}{k!^3 \cdot 27^k} \left(1 - p \sum_{i=k+1}^{3k} \frac{1}{i} + \frac{p^2}{2} \left( \left( \sum_{i=k+1}^{3k} \frac{1}{i} \right)^2 - \sum_{i=1}^{3k} \frac{1}{i^2} - \sum_{i=1}^k \frac{1}{i^2} \right)\right) \pmod{p^3}. \end{aligned}$$

For  $k = \frac{p-1}{4}$  we see that  $\sum_{i=k+1}^{3k} \frac{1}{i} = H_{p-1-\frac{p-1}{4}} - H_{\frac{p-1}{4}} \equiv 0 \pmod{p}$  and

$$\sum_{i=1}^{3(p-1)/4} \frac{1}{i^2} + \sum_{i=1}^{(p-1)/4} \frac{1}{i^2} \equiv \sum_{k=(p-1)/4+1}^{p-1} \frac{1}{k^2} + \sum_{k=1}^{(p-1)/4} \frac{1}{k^2} = \sum_{k=1}^{p-1} \frac{1}{k^2} \equiv 0 \pmod{p}.$$

Thus,

$$(5.7) \quad \sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv \sum_{k=0}^{[p/3]} \frac{p}{4k+1} \cdot \frac{(3k)!}{k!^3 \cdot 27^k} \left(1 - p \sum_{i=k+1}^{3k} \frac{1}{i}\right) \pmod{p^3}$$

and so  $\sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv \sum_{k=0}^{[p/3]} \frac{p}{4k+1} \cdot \frac{(3k)!}{k!^3 \cdot 27^k} \pmod{p^2}$ . Similarly,

$$\begin{aligned} \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} &= \sum_{n=0}^{p-1} \sum_{k=0}^n \binom{2k}{k}^2 \binom{4k}{2k} \binom{n+3k}{4k} \frac{1}{(-27)^k} \\ &= \sum_{k=0}^{p-1} \binom{2k}{k}^2 \binom{4k}{2k} \frac{1}{(-27)^k} \sum_{n=k}^{p-1} \binom{n+3k}{4k} = \sum_{k=0}^{p-1} \binom{2k}{k}^2 \binom{4k}{2k} \frac{1}{(-27)^k} \binom{p+3k}{4k+1} \\ &= p + \sum_{k=1}^{p-1} \frac{(p+3k)(p+3k-1) \cdots (p+1)p(p-1) \cdots (p-k)}{(4k+1) \cdot (-27)^k \cdot k!^4} \\ &= p + \sum_{k=1}^{p-1} \frac{p}{4k+1} \cdot \frac{(p^2-1^2)(p^2-2^2) \cdots (p^2-k^2) \cdot (p+(k+1)) \cdots (p+3k)}{k!^4 \cdot (-27)^k}. \end{aligned}$$

For  $\frac{p}{3} < k < p$  we see that  $\frac{p}{4k+1} \cdot \frac{(3k)!}{k!^3 \cdot 27^k} \equiv 0 \pmod{p^2}$ . Recall that  $\sum_{i=k+1}^{3k} \frac{1}{i} \equiv 0 \pmod{p}$  for  $k = \frac{p-1}{4}$ . From the above we then get

$$\begin{aligned} \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} &\equiv p + \sum_{k=1}^{p-1} \frac{p}{(4k+1)(-27)^k \cdot k!^4} (-1^2)(-2^2) \cdots (-k^2)(k+1) \cdots 3k \left(1 + \sum_{i=k+1}^{3k} \frac{p}{i}\right) \\ &= \sum_{k=0}^{p-1} \frac{p}{4k+1} \cdot \frac{(3k)!}{k!^3 \cdot 27^k} \left(1 + \sum_{i=k+1}^{3k} \frac{p}{i}\right) \equiv \sum_{k=0}^{[p/3]} \frac{p}{4k+1} \cdot \frac{(3k)!}{k!^3 \cdot 27^k} \left(1 + \sum_{i=k+1}^{3k} \frac{p}{i}\right) \\ &\equiv \sum_{k=0}^{[p/3]} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{4k+1} \pmod{p^2}. \end{aligned}$$

For  $\frac{2p}{3} < k < p$  we have  $p^2 \mid (3k)!$ , for  $\frac{p}{3} < k < \frac{2p}{3}$  we have  $p \nmid (4k+1)$  and  $p \mid (3k)!$ . Thus,

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{4k+1} \equiv \sum_{k=0}^{[p/3]} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{4k+1} \pmod{p^2}.$$

Combining the above proves the lemma.

**Theorem 5.2.** *Suppose that  $p$  is a prime with  $p > 3$  and  $p \equiv 3 \pmod{4}$ . Then*

$$\sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \equiv 0 \pmod{p^2}.$$

Proof. From Lemma 5.1 and (2.12) (with  $b = \frac{1}{4}$ ),

$$\sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \equiv \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{4k+1} \equiv 0 \pmod{p^2}.$$

**Theorem 5.3.** *Let  $p$  be a prime with  $p \equiv 1 \pmod{4}$ . Then*

$$\sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \equiv \begin{cases} 4x^2 - 2p \pmod{p^2} & \text{if } p \equiv 1 \pmod{12} \text{ and so } p = x^2 + 9y^2, \\ 2p - 2x^2 \pmod{p^2} & \text{if } p \equiv 5 \pmod{12} \text{ and so } 2p = x^2 + 9y^2. \end{cases}$$

Proof. Set  $S(a) = \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{p}{4k+1}$ . By (1.2),  $S(-\frac{1}{3}) = \sum_{k=0}^{p-1} \frac{(3k)!}{k!^3 \cdot 27^k} \cdot \frac{p}{4k+1}$ . Applying Lemma 5.1,

$$(5.8) \quad \sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \equiv S\left(-\frac{1}{3}\right) = S\left(-\frac{2}{3}\right) \pmod{p^2}.$$

For  $p \equiv 1 \pmod{12}$ , taking  $a = -\frac{1}{3}$ ,  $b = \frac{1}{4}$  and  $m = \frac{p-1}{12} - 1$  in Lemma 2.1 yields

$$(5.9) \quad S\left(-\frac{1}{3}\right) \equiv \frac{\binom{-7/12}{(p-1)/12}}{\binom{-1/12}{(p-1)/12}} S\left(-\frac{1}{4} - \frac{p}{12}\right) \pmod{p^3}.$$

For  $p \equiv 5 \pmod{12}$ , taking  $a = -\frac{2}{3}$ ,  $b = \frac{1}{4}$  and  $m = \frac{p-5}{12} - 1$  in Lemma 2.1 yields

$$(5.10) \quad S\left(-\frac{2}{3}\right) \equiv \frac{\binom{-11/12}{(p-5)/12}}{\binom{-5/12}{(p-5)/12}} S\left(-\frac{1}{4} - \frac{p}{12}\right) \pmod{p^3}.$$

Note that  $\sum_{k=1}^{(p-1)/2} \frac{1}{2k-1} = H_{p-1} - \frac{1}{2}H_{\frac{p-1}{2}} \equiv q_p(2) \pmod{p}$ . By [S2, Theorem 3.1(ii)],

$$(5.11) \quad \sum_{i=0}^{(p-1)/4-1} \frac{1}{4i+1} = \sum_{\substack{k=1 \\ k \equiv p \pmod{4}}}^{p-1} \frac{1}{k} \equiv \frac{3}{4}q_p(2) \pmod{p}.$$

Thus,

$$(5.12) \quad \sum_{i=0}^{\frac{p-1}{4}-1} \frac{1}{4i+3} = \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{2k-1} - \sum_{i=0}^{\frac{p-1}{4}-1} \frac{1}{4i+1} \equiv q_p(2) - \frac{3}{4}q_p(2) = \frac{1}{4}q_p(2) \pmod{p}.$$

Using (2.5), (5.11) and (5.12),

$$\begin{aligned} S\left(-\frac{1}{4} - \frac{p}{12}\right) &= \sum_{k=0}^{p-1} \binom{-\frac{1}{4} - \frac{p}{12}}{k} \binom{-\frac{3}{4} + \frac{p}{12}}{k} \frac{p}{4k+1} \\ &\equiv \sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k} \binom{-\frac{3}{4}}{k} \frac{p}{4k+1} \left(1 + \sum_{i=0}^{k-1} \frac{-\frac{p}{12}}{-\frac{1}{4} - i}\right) \left(1 + \sum_{i=0}^{k-1} \frac{\frac{p}{12}}{-\frac{3}{4} - i}\right) \end{aligned}$$

$$\begin{aligned}
&= \sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k} \binom{-\frac{3}{4}}{k} \frac{p}{4k+1} \left(1 + \frac{1}{3} \sum_{i=0}^{k-1} \frac{p}{4i+1}\right) \left(1 - \frac{1}{3} \sum_{i=0}^{k-1} \frac{p}{4i+3}\right) \\
&\equiv \sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k} \binom{-\frac{3}{4}}{k} \frac{p}{4k+1} \left(1 + \frac{1}{3} \sum_{i=0}^{k-1} \left(\frac{p}{4i+1} - \frac{p}{4i+3}\right)\right) \\
&\equiv \sum_{k=0}^{p-1} \binom{-\frac{1}{4}}{k} \binom{-\frac{3}{4}}{k} \frac{p}{4k+1} + \frac{p}{3} \binom{-\frac{1}{4}}{\frac{p-1}{4}} \binom{-\frac{3}{4}}{\frac{p-1}{4}} \sum_{i=0}^{(p-1)/4-1} \left(\frac{1}{4i+1} - \frac{1}{4i+3}\right) \\
&\equiv S\left(-\frac{1}{4}\right) + \frac{p}{3} \cdot \frac{\binom{\frac{p-1}{2}}{\frac{p-1}{4}} \binom{p-1}{\frac{p-1}{2}}}{64 \frac{p-1}{4}} \left(\frac{3}{4} q_p(2) - \frac{1}{4} q_p(2)\right) \\
&\equiv S\left(-\frac{1}{4}\right) + \frac{p}{6} \binom{\frac{p-1}{2}}{\frac{p-1}{4}} (-1)^{\frac{p-1}{4}} q_p(2) \pmod{p^2}.
\end{aligned}$$

Now applying Theorem 3.2 gives

$$S\left(-\frac{1}{4} - \frac{p}{12}\right) \equiv S\left(-\frac{1}{4}\right) + \frac{p}{6} \binom{\frac{p-1}{2}}{\frac{p-1}{4}} (-1)^{\frac{p-1}{4}} q_p(2) \equiv (-1)^{\frac{p-1}{4}} \binom{\frac{p-1}{2}}{\frac{p-1}{4}} \left(1 - \frac{1}{3} p q_p(2)\right) \pmod{p^2}.$$

Suppose  $p = A^2 + B^2$  with  $A \equiv 1 \pmod{4}$ . From [CDE] or [BEW],

$$(5.13) \quad \binom{(p-1)/2}{(p-1)/4} \equiv \left(1 + \frac{1}{2} p q_p(2)\right) \left(2A - \frac{p}{2A}\right) \pmod{p^2}.$$

Thus,

$$\begin{aligned}
S\left(-\frac{1}{4} - \frac{p}{12}\right) &\equiv (-1)^{\frac{p-1}{4}} \left(1 + \frac{1}{2} p q_p(2)\right) \left(2A - \frac{p}{2A}\right) \left(1 - \frac{1}{3} p q_p(2)\right) \\
&\equiv (-1)^{\frac{p-1}{4}} \left(1 + \frac{1}{6} p q_p(2)\right) \left(2A - \frac{p}{2A}\right) \pmod{p^2}.
\end{aligned}$$

Combining this with (5.8)-(5.10) yields

$$(5.14) \quad \begin{cases} \sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \\ \equiv \begin{cases} \frac{\binom{-7/12}{(p-1)/12}}{\binom{-1/12}{(p-1)/12}} (-1)^{\frac{p-1}{4}} \left(1 + \frac{1}{6} p q_p(2)\right) \left(2A - \frac{p}{2A}\right) \pmod{p^2} & \text{if } p \equiv 1 \pmod{12}, \\ \frac{\binom{-11/12}{(p-5)/12}}{\binom{-5/12}{(p-5)/12}} (-1)^{\frac{p-1}{4}} \left(1 + \frac{1}{6} p q_p(2)\right) \left(2A - \frac{p}{2A}\right) \pmod{p^2} & \text{if } p \equiv 5 \pmod{12}. \end{cases} \end{cases}$$

For  $p \equiv 1 \pmod{12}$  suppose  $p = x^2 + 9y^2$ . Using (2.5) we see that

$$\frac{\binom{-\frac{7}{12}}{\frac{p-1}{12}}}{\binom{-\frac{1}{12}}{\frac{p-1}{12}}} = \frac{\binom{\frac{7(p-1)}{12} - \frac{7p}{12}}{\frac{p-1}{12}}}{\binom{\frac{p-1}{12} - \frac{p}{12}}{\frac{p-1}{12}}} \equiv \frac{\binom{\frac{7(p-1)}{12}}{\frac{p-1}{12}} \left(1 - \frac{7}{12} p \sum_{i=0}^{\frac{p-1}{12}-1} \frac{1}{\frac{7(p-1)}{12} - i}\right)}{\binom{\frac{p-1}{12}}{\frac{p-1}{12}} \left(1 - \frac{p}{12} \sum_{i=0}^{\frac{p-1}{12}-1} \frac{1}{\frac{p-1}{12} - i}\right)}$$

$$\begin{aligned}
&= \left( \frac{7(p-1)}{12} \right) \frac{1 - \frac{7}{12}p(H_{\frac{7(p-1)}{12}} - H_{\frac{p-1}{2}})}{1 - \frac{p}{12}H_{\frac{p-1}{12}}} \\
&\equiv \left( \frac{7(p-1)}{12} \right) \left( 1 - \frac{7}{12}p(H_{\frac{7(p-1)}{12}} - H_{\frac{p-1}{2}}) \right) \left( 1 + \frac{p}{12}H_{\frac{p-1}{12}} \right) \\
&\equiv \left( \frac{7(p-1)}{12} \right) \left( 1 - \frac{7}{12}p(H_{\frac{7(p-1)}{12}} + 2q_p(2)) + \frac{p}{12}H_{\frac{p-1}{12}} \right) \pmod{p^2}.
\end{aligned}$$

By (4.7),  $H_{\frac{7(p-1)}{12}} \equiv H_{\frac{5(p-1)}{12}} \equiv -6q_p(2) - 3q_p(3) - H_{\frac{p-1}{12}} \pmod{p}$ . By [S3, Lemma 2.5],

$$\begin{aligned}
\left( \frac{7(p-1)}{12} \right) &\equiv (-1)^{\frac{p-1}{12}} \left( \frac{p-1}{12} \right) \left( 1 + 2p \left( H_{\frac{p-1}{6}} - \frac{1}{2}H_{\frac{p-1}{12}} \right) \right) \\
&\equiv (-1)^{\frac{p-1}{12}} \left( \frac{p-1}{12} \right) \left( 1 - p(4q_p(2) + 3q_p(3) + H_{\frac{p-1}{12}}) \right) \pmod{p^2}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
&(-1)^{\frac{p-1}{4}} \left( 1 + \frac{1}{6}pq_p(2) \right) \left( \frac{-7}{12} \right) \left( \frac{-1}{12} \right)^{-1} \\
&\equiv \left( 1 + \frac{1}{6}pq_p(2) \right) \left( \frac{p-1}{12} \right) \left( 1 - p(4q_p(2) + 3q_p(3) + H_{\frac{p-1}{12}}) \right) \\
&\quad \times \left( 1 - \frac{7}{12}p(-4q_p(2) - 3q_p(3) - H_{\frac{p-1}{12}}) + \frac{p}{12}H_{\frac{p-1}{12}} \right) \\
&\equiv \left( \frac{p-1}{12} \right) \left( 1 - p \left( \frac{3}{2}q_p(2) + \frac{5}{4}q_p(3) + \frac{1}{3}H_{\frac{p-1}{12}} \right) \right) \pmod{p^2}.
\end{aligned}$$

This together with (5.14) and (4.9) yields

$$\begin{aligned}
\sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} &\equiv \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \equiv \left( \frac{p-1}{12} \right) \left( 1 - p \left( \frac{3}{2}q_p(2) + \frac{5}{4}q_p(3) + \frac{1}{3}H_{\frac{p-1}{12}} \right) \right) \left( 2A - \frac{p}{2A} \right) \\
&\equiv \left( 2c - \frac{p}{2c} \right) \left( 2A - \frac{p}{2A} \right) \equiv \begin{cases} 4A^2 - 2p = 4x^2 - 2p \pmod{p^2} & \text{if } 3 \nmid A, \\ -(4A^2 - 2p) \equiv 4x^2 - 2p \pmod{p^2} & \text{if } 3 \mid A, \end{cases}
\end{aligned}$$

where  $c = A$  or  $-A$  according as  $3 \nmid A$  or  $3 \mid A$ .

Now assume  $p \equiv 5 \pmod{12}$  and  $p = A^2 + B^2$  with  $A \equiv 1 \pmod{4}$  and  $B \equiv A \pmod{3}$ . Appealing to (2.5),

$$\begin{aligned}
\frac{\left( \frac{-11}{12} \right)}{\left( \frac{p-5}{12} \right)} &= \frac{\left( \frac{7p-11}{12} - \frac{7p}{12} \right)}{\left( \frac{p-5}{12} - \frac{p}{12} \right)} \equiv \frac{\left( \frac{7p-11}{12} \right) \left( 1 - \frac{7}{12}p \sum_{i=0}^{\frac{p-5}{12}-1} \frac{1}{\frac{7p-11}{12}-i} \right)}{\left( \frac{p-5}{12} \right) \left( 1 - \frac{p}{12} \sum_{i=0}^{\frac{p-5}{12}-1} \frac{1}{\frac{p-5}{12}-i} \right)} \\
&= \left( \frac{7p-11}{12} \right) \frac{1 - \frac{7}{12}p(H_{\frac{7p-11}{12}} - H_{\frac{p-1}{2}})}{1 - \frac{p}{12}H_{\frac{p-5}{12}}} \\
&\equiv \left( \frac{7p-11}{12} \right) \left( 1 - \frac{7}{12}p(H_{\frac{7p-11}{12}} - H_{\frac{p-1}{2}}) \right) \left( 1 + \frac{p}{12}H_{\frac{p-5}{12}} \right)
\end{aligned}$$

$$\equiv \binom{\frac{7p-11}{12}}{\frac{p-5}{12}} \left(1 - \frac{7}{12}p(H_{\frac{7p-11}{12}} + 2q_p(2)) + \frac{p}{12}H_{\frac{p-5}{12}}\right) \pmod{p^2}.$$

By (4.7),

$$H_{\frac{7p-11}{12}} = H_{p-1-\frac{5p-1}{12}} \equiv H_{\frac{5p-1}{12}} \equiv -6q_p(2) - 3q_p(3) - H_{\frac{p-5}{12}}.$$

By [S3, Lemma 2.5],

$$\begin{aligned} \binom{\frac{7p-11}{12}}{\frac{p-5}{12}} &\equiv (-1)^{\frac{p-5}{12}} \binom{\frac{p-1}{2}}{\frac{p-5}{12}} \left(1 + 2p\left(H_{\frac{p-5}{6}} - \frac{1}{2}H_{\frac{p-5}{12}}\right)\right) \\ &\equiv (-1)^{\frac{p-5}{12}} \binom{\frac{p-1}{2}}{\frac{p-5}{12}} \left(1 - p\left(4q_p(2) + 3q_p(3) + H_{\frac{p-5}{12}}\right)\right) \pmod{p^2}. \end{aligned}$$

Hence

$$\begin{aligned} \frac{\binom{-\frac{11}{12}}{\frac{p-5}{12}}}{\binom{-\frac{5}{12}}{\frac{p-5}{12}}} &\equiv (-1)^{\frac{p-5}{12}} \binom{\frac{p-1}{2}}{\frac{p-5}{12}} \left(1 - p\left(4q_p(2) + 3q_p(3) + H_{\frac{p-5}{12}}\right)\right) \\ &\quad \times \left(1 - \frac{7}{12}p\left(-6q_p(2) - 3q_p(3) - H_{\frac{p-5}{12}} + 2q_p(2)\right) + \frac{p}{12}H_{\frac{p-5}{12}}\right) \\ &\equiv (-1)^{\frac{p-5}{12}} \binom{\frac{p-1}{2}}{\frac{p-5}{12}} \left(1 - p\left(\frac{5}{3}q_p(2) + \frac{5}{4}q_p(3) + \frac{1}{3}H_{\frac{p-5}{12}}\right)\right) \pmod{p^2} \end{aligned}$$

and so

$$\begin{aligned} &\binom{\frac{7p-11}{12}}{\frac{p-5}{12}} (-1)^{\frac{p-1}{4}} \left(1 + \frac{1}{6}pq_p(2)\right) \\ &\equiv -\binom{\frac{p-1}{2}}{\frac{p-5}{12}} \left(1 + \frac{1}{6}pq_p(2)\right) \left(1 - p\left(\frac{5}{3}q_p(2) + \frac{5}{4}q_p(3) + \frac{1}{3}H_{\frac{p-5}{12}}\right)\right) \\ &\equiv -\binom{\frac{p-1}{2}}{\frac{p-5}{12}} \left(1 - p\left(\frac{3}{2}q_p(2) + \frac{5}{4}q_p(3) + \frac{1}{3}H_{\frac{p-5}{12}}\right)\right) \equiv -\left(2B - \frac{p}{2B}\right) \pmod{p^2} \end{aligned}$$

by (4.9). This together with (5.14) yields

$$\sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \equiv \sum_{n=0}^{p-1} \frac{b_n}{(-27)^n} \equiv -\left(2A - \frac{p}{2A}\right) \left(2B - \frac{p}{2B}\right) \equiv -4AB \pmod{p^2}.$$

Suppose  $2p = x^2 + 9y^2$ . Then clearly  $\left(\frac{x+3y}{2}\right)^2 + \left(\frac{x-3y}{2}\right)^2 = p$  and  $\frac{x+3y}{2} \equiv \frac{x-3y}{2} \pmod{3}$ .

Thus,

$$2p - 2x^2 = 9y^2 - x^2 = -4 \cdot \frac{x+3y}{2} \cdot \frac{x-3y}{2} = -4AB.$$

Thus the theorem is proved.

**Corollary 5.1.** *Let  $p$  be a prime of the form  $12k + 5$ . Then  $p = A^2 + B^2$  with  $A \equiv 1 \pmod{4}$  and  $B \equiv A \pmod{3}$ , and  $2p = x^2 + 9y^2$ . We have*

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-12288)^k} \equiv -4AB \equiv -2x^2 \pmod{p}.$$

Proof. Taking  $u = -1/3$  in [S8, Theorem 5.1] gives  $\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2 \binom{4k}{2k}}{(-12288)^k} \equiv \sum_{n=0}^{p-1} \frac{b_n}{(-3)^n} \pmod{p}$ . Thus the result follows from Theorem 5.3 and its proof.

**Theorem 5.4.** *Let  $p > 3$  be a prime. Then*

$$\begin{aligned} \sum_{n=0}^{p-1} \frac{W_n}{(-3)^n} &\equiv \sum_{k=0}^{\lfloor p/3 \rfloor} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{3k+1} \\ &\equiv \begin{cases} -L + \frac{p}{L} \pmod{p^2} & \text{if } 3 \mid p-1 \text{ and so } 4p = L^2 + 27M^2 \text{ with } 3 \mid L-1, \\ -\frac{p}{3} \left( \frac{p-2}{3}! \right)^3 \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases} \end{aligned}$$

Proof. Using (1.8) and (5.2),

$$\begin{aligned} \sum_{n=0}^{p-1} \frac{W_n}{(-3)^n} &= \sum_{n=0}^{p-1} \sum_{k=0}^{\lfloor n/3 \rfloor} \binom{2k}{k} \binom{3k}{k} \binom{n}{3k} \frac{1}{(-27)^k} \\ &= \sum_{k=0}^{\lfloor p/3 \rfloor} \binom{2k}{k} \binom{3k}{k} \frac{1}{(-27)^k} \sum_{n=3k}^{p-1} \binom{n}{3k} = \sum_{k=0}^{\lfloor p/3 \rfloor} \binom{2k}{k} \binom{3k}{k} \frac{1}{(-27)^k} \binom{p}{3k+1} \\ &= \sum_{k=0}^{\lfloor p/3 \rfloor} \binom{2k}{k} \binom{3k}{k} \frac{1}{(-27)^k} \cdot \frac{p}{3k+1} \cdot \frac{(p-1)(p-2)\cdots(p-3k)}{(3k)!} \\ &\equiv \sum_{k=0}^{\lfloor p/3 \rfloor} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{3k+1} \pmod{p^2}. \end{aligned}$$

For  $\frac{p}{3} < k < p$  with  $k \neq \frac{2p-1}{3}$  we have  $p \nmid 3k+1$  and  $\binom{2k}{k} \binom{3k}{k} = \frac{(3k)!}{k!^3} \equiv 0 \pmod{p}$ . Thus,

$$\begin{aligned} &\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{3k+1} - \sum_{k=0}^{\lfloor p/3 \rfloor} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{3k+1} \\ &\equiv \begin{cases} 0 \pmod{p^2} & \text{if } p \equiv 1 \pmod{3}, \\ \frac{(2p-1)!}{\left(\frac{2p-1}{3}\right)!^3} \cdot \frac{1}{2 \cdot 27^{(2p-1)/3}} \pmod{p^2} & \text{if } p \equiv 2 \pmod{3}. \end{cases} \end{aligned}$$

For  $p \equiv 1 \pmod{3}$  it is well known that  $4p = L^2 + 27M^2$  with  $L, M \in \mathbb{Z}$  and  $L \equiv 1 \pmod{3}$ . By [BEW, Theorem 9.4.4],  $\binom{\frac{2(p-1)}{3}}{\frac{p-1}{3}} \equiv -L + \frac{p}{L} \pmod{p^2}$ . Hence applying the above and Theorem 3.3,

$$\sum_{n=0}^{p-1} \frac{W_n}{(-3)^n} \equiv \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{3k+1} \equiv \binom{\frac{2(p-1)}{3}}{\frac{p-1}{3}} \equiv -L + \frac{p}{L} \pmod{p^2}.$$

For  $p \equiv 2 \pmod{3}$ ,

$$\frac{(2p-1)!}{\left(\frac{2p-1}{3}\right)!^3} \cdot \frac{1}{2 \cdot 27^{(2p-1)/3}} = \frac{p(p^2-1^2)(p^2-2^2)\cdots(p^2-(p-1)^2) \cdot \left(\frac{p-2}{3}\right)!^3}{\left(\frac{2p-1}{3}\right)!^3 \left(\frac{p-2}{3}\right)!^3 \cdot 2 \cdot 3^{2p-1}}$$



$$\equiv p \cdot \left(\frac{p-2}{3}\right)!^3 \frac{\binom{p-1}{(p-2)/3}^3}{(p-1)! \cdot 2 \cdot 3} \equiv \frac{p}{6} \left(\frac{p-2}{3}\right)!^3 \pmod{p^2}$$

and

$$\left(\frac{\frac{2(p-2)}{3}}{\frac{p-2}{3}}\right)^{-1} = \frac{\left(\frac{p-2}{3}\right)!^2 (p-1)(p-2) \cdots (p - \frac{p+1}{3})}{(p-1)!} \equiv -\frac{1}{3} \left(\frac{p-2}{3}\right)!^3 \pmod{p}.$$

From (2.16) and the above,

$$\begin{aligned} \sum_{n=0}^{p-1} \frac{W_n}{(-3)^n} &\equiv \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{3k+1} - \frac{p}{6} \left(\frac{p-2}{3}\right)!^3 \equiv \frac{p}{2} \left(\frac{\frac{2(p-2)}{3}}{\frac{p-2}{3}}\right)^{-1} - \frac{p}{6} \left(\frac{p-2}{3}\right)!^3 \\ &\equiv -\frac{p}{6} \left(\frac{p-2}{3}\right)!^3 - \frac{p}{6} \left(\frac{p-2}{3}\right)!^3 = -\frac{p}{3} \left(\frac{p-2}{3}\right)!^3 \pmod{p^2}. \end{aligned}$$

Thus, the theorem is proved.

To conclude the paper, we pose two challenging conjectures.

**Conjecture 5.1.** *Let  $p > 3$  be a prime. Then*

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \cdot \frac{p}{4k+1} \equiv \begin{cases} 4x^2 - 2p - \frac{p^2}{4x^2} \pmod{p^3} & \text{if } 12 \mid p-1 \text{ and so } p = x^2 + 9y^2, \\ 2p - 2x^2 + \frac{p^2}{2x^2} \pmod{p^3} & \text{if } 12 \mid p-5 \text{ and so } 2p = x^2 + 9y^2, \\ -\frac{5}{27} p^2 \binom{[p/3]}{[p/12]}^{-2} \pmod{p^3} & \text{if } 12 \mid p-7, \\ \frac{5}{54} p^2 \binom{[p/3]}{[p/12]}^{-2} \pmod{p^3} & \text{if } 12 \mid p-11. \end{cases}$$

**Conjecture 5.2.** *Suppose that  $p$  is a prime of the form  $4k+1$ . Then*

$$(-1)^{[\frac{p}{12}]} \binom{[\frac{p}{3}]}{[\frac{p}{12}]} \equiv \begin{cases} 2x \pmod{p} & \text{if } p \equiv 1 \pmod{12} \text{ and so } p = x^2 + 9y^2 \text{ with } 3 \mid x-1, \\ x \pmod{p} & \text{if } p \equiv 5 \pmod{12} \text{ and so } 2p = x^2 + 9y^2 \text{ with } 3 \mid x-1. \end{cases}$$

## References

- [AT] T. Amdeberhan and R. Tauraso, *Supercongruences for the Almkvist-Zudilin numbers*, Acta Arith. **173**(2016), 255-268.
- [BEW] B.C. Berndt, R.J. Evans and K.S. Williams, *Gauss and Jacobi Sums*, Wiley, New York, 1998.
- [CZ] H.H. Chan and W. Zudilin, *New representations for Apéry-like sequences*, Mathematika **56**(2010), 107-117.

- [CDE] S. Chowla, B. Dwork and R. J. Evans, *On the mod  $p^2$  determination of  $\binom{(p-1)/2}{(p-1)/4}$* , J. Number Theory **24**( 1986), 188-196.
- [G] H.W. Gould, *Combinatorial Identities, A Standardized Set of Tables Listing 500 Binomial Coefficient Summations*, West Virginia University, Morgantown, WV, 1972.
- [L] E. Lehmer, *On congruences involving Bernoulli numbers and the quotients of Fermat and Wilson*, Ann. Math. **39**(1938), 350-360.
- [M1] E. Mortenson, *A supercongruence conjecture of Rodriguez-Villegas for a certain truncated hypergeometric function*, J. Number Theory **99**(2003), 139-147.
- [M2] E. Mortenson, *Supercongruences between truncated  ${}_2F_1$  hypergeometric functions and their Gaussian analogs*, Trans. Amer. Math. Soc. **355**(2003), 987-1007.
- [RV] F. Rodriguez-Villegas, *Hypergeometric families of Calabi-Yau manifolds*, in: Noriko Yui, James D. Lewis (Eds.), *Calabi-Yau Varieties and Mirror Symmetry*, Toronto, ON, 2001, in: Fields Inst. Commun., vol. 38, Amer. Math. Soc., Providence, RI, 2003, pp.223-231.
- [S1] Z.H. Sun, *Congruences concerning Bernoulli numbers and Bernoulli polynomials*, Discrete Appl. Math. **105**(2000), 193-223.
- [S2] Z.H. Sun, *Congruences involving Bernoulli and Euler numbers*, J. Number Theory **128**(2008), 280-312.
- [S3] Z. H. Sun, *Congruences concerning Legendre polynomials*, Proc. Amer. Math. Soc. **139**(2011), 1915-1929.
- [S4] Z.H. Sun, *Identities and congruences for a new sequence*, Int. J. Number Theory **8**(2012), 207-225.
- [S5] Z. H. Sun, *Congruences concerning Legendre polynomials II*, J. Number Theory **133**(2013), 1950-1976.
- [S6] Z. H. Sun, *Legendre polynomials and supercongruences*, Acta Arith. **159**(2013), 169-200.
- [S7] Z. H. Sun, *Generalized Legendre polynomials and related supercongruences*, J. Number Theory **143**(2014), 293-319.
- [S8] Z. H. Sun, *Congruences for Domb and Almkvist-Zudilin numbers*, Integral Transforms Spec. Funct. **26**(2015), 642-659.
- [S9] Z. H. Sun, *Supercongruences involving Bernoulli polynomials*, Int. J. Number Theory **12**(2016), 1259-1271.
- [S10] Z.H. Sun, *Congruences for sums involving Franel numbers*, Int. J. Number Theory **14**(2018), 123-142.
- [S11] Z.H. Sun, *Super congruences for two Apéry-like sequences*, J. Differ. Equ. Appl. **24**(2018), 1685-1713.

- [S12] Z.H. Sun, *Congruences involving binomial coefficients and Apéry-like numbers*, Publ. Math. Debrecen, to appear.
- [Su1] Z.W. Sun, *Connections between  $p = x^2 + 3y^2$  and Franel numbers*, J. Number Theory **133**(2013), 2914-2928.
- [Su2] Z.W. Sun, *Conjectures and results on  $x^2 \pmod{p^2}$  with  $4p = x^2 + dy^2$* , in: Number Theory and Related Area (eds., Y. Ouyang, C. Xing, F. Xu and P. Zhang), Higher Education Press & International Press, Beijing and Boston, 2013, pp.149-197.
- [Y] K.M. Yeung, *On congruences for binomial coefficients*, J. Number Theory **33**(1989), 1-17.