

A CONSTRUCTION OF THE FREE SKEW-BRACE

JUAN ORZA

ABSTRACT. The aim of this paper is to give a simple construction of the free skew-brace over a set X .

INTRODUCTION

The concept of skew-brace was introduced by Guarnieri and Vendramin [1] in order to study non-involutive set theoretical solutions of de braid equation using method of non-commutative ring theory. For the original definition we refer to this paper. In [3] the author ask for a construction of the skew-brace over a set X . The free skew-brace over X exist by general result of universal algebra, but the constructions of free objects valid in the context of universal algebra are very involved. So, in this note we present a simple construction of the free skew-brace using the following result established in [2]:

Proposition. A group (A, \circ) , endowed with a pair of left actions $b \mapsto a \cdot b$ y $b \mapsto a : b$ on A (as a set), is a skew-brace if and only if the following equality holds:

$$(a \cdot b) \circ a = (b : a) \circ b \quad \text{for all } a, b \in A. \quad (1)$$

1. A CONSTRUCTION OF THE FREE SKEW-BRACE OVER X

To begin with, we recursively define a set Y in the following way: first we set

$$Y_1 := X \sqcup X^*,$$

where $X^* := \{x^* : x \in X\}$ is a disjoint copy of the elements of X . Next we define the sets

$$\begin{aligned} X_n^{\cdot} &:= \{a \cdot b \mid a, b \in Y_{n-1}, b \neq a^* \cdot c \text{ for any } c \in Y_{n-1}\} \\ X_n^{:} &:= \{a : b \mid a, b \in Y_{n-1}, b \neq a^* : c \text{ for any } c \in Y_{n-1}\} \\ Y_n &:= Y_{n-1} \sqcup (X_n^{\cdot} \sqcup X_n^{:}) \sqcup (X_n^{\cdot} \sqcup X_n^{:})^* \end{aligned}$$

where $a \cdot b$ and $a : b$ are formal elements. Then we set $Y := \bigcup_{n \geq 1} Y_n$ and we let G denotes the free group $F(Y)$ over Y (with underlying set the collection of reduced words, with the element x^* playing the roll of the inverse of x). In the sequel, we are going to denote the multiplication of G by juxtaposition $a \circ b = ab$.

The next step is to define maps \cdot and $:$ from $Y \times Y$ to Y by

$$\begin{aligned} a \cdot b &= \begin{cases} a \cdot b & \text{if } b \neq a^* \cdot c, \\ c & \text{if } b = a^* \cdot c, \end{cases} \\ a : b &= \begin{cases} a : b & \text{if } b \neq a^* : c, \\ c & \text{if } b = a^* : c. \end{cases} \end{aligned}$$

We want to extend both \cdot and $:$ to left actions of G on G in such away that $(a \cdot b)a = (b : a)b$ for all $a, b \in G$, which happen if and only if

$$a \cdot b = (b : a)ba^{-1} \quad \text{for all } a, b \in G, \quad (2)$$

or, equivalently,

$$b : a = (a \cdot b)ab^{-1} \quad \text{for all } a, b \in G. \quad (3)$$

Taking $b = 1$ in this equalities we see that necessarily $a \cdot 1 = a : 1 = 1$ for all $a \in G$. Let $a, b, c \in G$. From (3) it follows that

$$\begin{aligned} & (ab) : c = a : (b : c) \\ \Leftrightarrow & (c \cdot (ab))c(ab)^{-1} = a : ((c \cdot b)cb^{-1}) \\ \Leftrightarrow & (c \cdot (ab))cb^{-1}a^{-1} = (((c \cdot b)cb^{-1}) \cdot a)(c \cdot b)cb^{-1}a^{-1} \\ \Leftrightarrow & c \cdot (ab) = ((b : c) \cdot a)(c \cdot b). \end{aligned}$$

Similarly, from (2) it follows that:

$$(ab) \cdot c = a \cdot (b \cdot c) \Leftrightarrow c : (ab) = ((b \cdot c) : a)(c \cdot b).$$

This motivates to extend both \cdot and $:$ to maps from $Y \times G$ to G , by defining

$$x \cdot (gy) := ((y : x) \cdot g)(x \cdot y), \quad (4)$$

$$x : (gy) := ((y \cdot x) : g)(x : y), \quad (5)$$

for each $x, y \in X$ and each reduced word gy , and then to extend these definitions to left actions of G on G , by defining

$$\begin{aligned} (x_1x_2 \dots x_n) \cdot g &= x_1 \cdot (x_2 \cdot (\dots (x_n \cdot g) \dots)), \\ (x_1x_2 \dots x_n) : g &= x_1 : (x_2 : (\dots (x_n : g) \dots)), \\ 1 \cdot g &= 1 : g = g, \end{aligned}$$

for all $x_i \in Y$ and $g \in G$.

Now we check that

$$x \cdot (ab) = ((b : x) \cdot a)(x \cdot b) \quad \text{for all } x \in Y \text{ and } a, b \in G.$$

We proceed by induction on the length of b . This is clear for $b = 1$. Assume that it is true for $b \in G$, and let us see that it is true for by with $y \in Y$:

$$\begin{aligned} x \cdot (a(by)) &= ((y : x) \cdot (ab))(x \cdot y) \\ &= ((b : (y : x)) \cdot a)((y : x) \cdot b)(x \cdot y) \\ &= (((by) : x) \cdot a)(x \cdot (by)) \end{aligned}$$

In a similar way it is proved that $x : (ab) = ((b \cdot x) : a)(x : b)$.

Lemma 1.1. *For all $a, b, c \in G$, the identities*

$$a \cdot (bc) = ((c : a) \cdot b)(a \cdot c) \quad \text{and} \quad a : (bc) = ((c \cdot a) : b)(a : c) \quad (6)$$

are fulfilled.

Proof. We prove the first equality and left the second one, which is similar, to the reader. Clearly the first equality in (6) is true if $a = 1$ or $c = 1$. Take $c = x \in Y$ and suppose that the first equality in (6) holds for $a \in G$. Then,

$$\begin{aligned} (ya) \cdot (bx) &= y \cdot (a \cdot (bx)) \\ &= y \cdot (((x : a) \cdot b)(a \cdot x)) \\ &= (((a \cdot x) : y) \cdot ((x : a) \cdot b))(y \cdot (a \cdot x)) \\ &= (((a \cdot x) : y)(x : a) \cdot b)((ya) \cdot x) \\ &= ((x : (ya)) \cdot b)((ya) \cdot x) \end{aligned}$$

Let $a, b, c \in G$, $z \in X$, and suppose that the first equality in (6) holds $c \in G$. We have

$$\begin{aligned} a \cdot (b(cz)) &= ((z : a) \cdot (bc))(a \cdot z) \\ &= ((c : (z : a)) \cdot b)((z : a) \cdot c)(a \cdot z) \\ &= ((cz : a) \cdot b)(a \cdot (cz)). \end{aligned}$$

which proves that the first equality in (6) also holds for cz . \square

It is not true that $(a \cdot b)a = (b : a)b$ for all $a, b \in G$. So, in order to finish our task we need to take a quotient G/\sim of G by a suitable congruence relation.

Proposition 1.2. *Let A be a group endowed with a pair of left actions $b \mapsto a \cdot b$ $y \mapsto a : b$ on A (as a set) such that*

$$a \cdot (bc) = ((c : a) \cdot b)(a \cdot c) \quad \text{and} \quad a : (bc) = ((c \cdot a) : b)(a : c) \quad \text{for all } a, b, c \in A,$$

and let X be a set of generators of A as a monoid. If $(x \cdot y)x = (y : x)y$ for all $x, y \in X$, then $(a \cdot b)a = (b : a)b$ for all $a, b \in A$.

Proof. Suppose that $(x \cdot b)x = (b : x)b$ for some $b \in G$ and all $x, y \in X$. Then, for all $x, y \in X$,

$$\begin{aligned} (x \cdot (by))x &= ((y : x) \cdot b)(x \cdot y)x \\ &= (b : (y : x))by \\ &= ((by) : x)by. \end{aligned}$$

So, $(x \cdot b)x = (b : x)b$ for all $x \in X$ and $b \in G$. Let $a \in G$. Suppose that $(a \cdot b)a = (b : a)b$ for all $b \in A$. Then

$$\begin{aligned} ((ax) \cdot b)ax &= (a \cdot (x \cdot b))ax \\ &= ((x \cdot b) : a)(x \cdot b)x \\ &= ((x \cdot b) : a)(b : x)b \\ &= (b : (ax))b, \end{aligned}$$

which finishes the proof. \square

Let \sim be the minimal equivalence relation in A such that:

- (1) $(x \cdot y)x \sim (y : x)y$ for all $x, y \in X$, and
- (2) If $a \sim b$ and $c \sim d$, then

$$ac \sim bd, \tag{7}$$

$$a \cdot c \sim b \cdot d, \tag{8}$$

$$a : c \sim b : d. \tag{9}$$

By Proposition 1.2 and the proposition in the introduction, G/\sim is a skew brace.

Theorem 1.3. *The group G/\sim , endowed with the binary operations \cdot and $:$ induced by the corresponding operations in G , is the free skew brace over X .*

Proof. Let A be a skew brace and let $f : X \rightarrow A$ be a map. Next, we extend f to a group morphism $\tilde{f} : G \rightarrow A$ such that

$$\tilde{f}(x \cdot y) = f(x) \cdot f(y) \quad \text{and} \quad \tilde{f}(x : y) = f(x) : f(y) \quad \text{for all } x, y \in Y.$$

For this, first we extend f to Y as follows:

- We extend f to Y_1 by defining $\tilde{f}(x^*) := f(x)^{-1}$ for each $x \in X$;
- Assuming that we have defined \tilde{f} on Y_{n-1} , we extend \tilde{f} to Y_n by defining $\tilde{f}(x \cdot y) := f(x) \cdot f(y)$ and $\tilde{f}(x : y) := f(x) : f(y)$, for all $x, y \in Y_{n-1}$.

Then, we extend f to G by defining

$$\tilde{f}(x_1 x_2 \dots x_n) := \tilde{f}(x_1) \tilde{f}(x_2) \cdots \tilde{f}(x_n) \quad \text{for } x_i \in Y.$$

Since equality (1) is satisfied in A , the map \tilde{f} induce a skew brace morphism $\varphi : G/\sim \rightarrow A$, unique such that $f = \varphi \circ i$, where $i : X \rightarrow G/\sim$ is the canonical map. \square

REFERENCES

- [1] L. Guarnieri and L. Vendramin, *Skew braces and the Yang-Baxter equation*, Math. Comput. **86** (2017), no. 307, 2519–2534 (English).
- [2] W. Rump, *A covering theory for non-involutive set-theoretic solutions to the Yang-Baxter equation*, J. Algebra **520** (2019), 136–170 (English).
- [3] L. Vendramin, *Problems on skew left braces*, Adv. Group Theory Appl. **7** (2019), 15–37 (English).

DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS EXACTAS Y NATURALES-UBA, PABELLÓN 1-CIUDAD UNIVERSITARIA, INTENDENTE GUIRALDES 2160 (C1428EGA) BUENOS AIRES, ARGENTINA.