

A Linear Algebra Approach for Detecting Binomiality of Steady State Ideals of Reversible Chemical Reaction Networks

Hamid Rahkooy¹, Ovidiu Radulescu², and Thomas Sturm^{1,3}

¹ CNRS, Inria, and the University of Lorraine, Nancy, France
hamid.rahkooy@inria.fr

² LPHI CNRS UMR 5235, and University of Montpellier, Montpellier, France
ovidiu.radulescu@umontpellier.fr



³ MPI-INF and Saarland University, Saarbrücken, Germany
thomas.sturm@loria.fr

Abstract. Motivated by problems from Chemical Reaction Network Theory, we investigate whether steady state ideals of reversible reaction networks are generated by binomials. We take an algebraic approach considering, besides concentrations of species, also rate constants as indeterminates. This leads us to the concept of unconditional binomiality, meaning binomiality for all values of the rate constants. This concept is different from conditional binomiality that applies when rate constant values or relations among rate constants are given. We start by representing the generators of a steady state ideal as sums of binomials, which yields a corresponding coefficient matrix. On these grounds we propose an efficient algorithm for detecting unconditional binomiality. That algorithm uses exclusively elementary column and row operations on the coefficient matrix. We prove asymptotic worst case upper bounds on the time complexity of our algorithm. Furthermore, we experimentally compare its performance with other existing methods.

Keywords: Binomial Ideals, Linear Algebra, Reversible Chemical Reaction Networks

1 Introduction

A *chemical reaction* is a transformation between two sets of chemical objects called chemical *complexes*. The objects that form a chemical complex are chemical *species*. In other words, complexes are formal sums of chemical species representing the left hand and the right hand sides of chemical reactions. A *chemical reaction network* is a set of chemical reactions. For example



is a chemical reaction network with two *reversible* reactions.

A *kinetics* of a chemical reaction network is an assignment of a rate function, depending on the concentrations of chemical species at the left hand side, to each reaction in the network. A kinetics for a chemical reaction network is called *mass-action* if for each reaction in the chemical reaction network, the rate function is a monomial in the concentrations of the chemical species with exponents given by the numbers of molecules of the species consumed in the reaction, multiplied by a constant called rate constant. Reactions are classified as zero-order, first-order, etc. according to the order of the monomial giving the rate. For reversible reactions, the net reaction rate is a binomial, the difference between the forward and backward rates. In the example above k_{12} , k_{21} , k_{23} , k_{32} are the rate constants. In this article we generally assume mass-action kinetics. We furthermore assume that reactions are reversible, unless explicitly specified otherwise.

The change in the concentration of each species over time in a reaction can be described via a system of autonomous ordinary differential equations. For instance, consider the chemical reaction network above and let x_1, x_2, x_3, x_4, x_5 be the indeterminates representing the concentrations of the species CO_2 , H_2 , CO , H_2O and C , respectively. The corresponding differential equations are

$$\dot{x}_1 = p_1, \quad p_1 = -k_{12}x_1x_2 + k_{21}x_3x_4 - k_{34}x_3^2 - k_{43}x_1x_5, \quad (1)$$

$$\dot{x}_2 = p_2, \quad p_2 = -k_{12}x_1x_2 + k_{21}x_3x_4, \quad (2)$$

$$\dot{x}_3 = p_3, \quad p_3 = k_{12}x_1x_2 - k_{21}x_3x_4 + -2k_{34}x_3^2 + 2k_{43}x_1x_5, \quad (3)$$

$$\dot{x}_4 = p_4, \quad p_4 = k_{12}x_1x_2 - k_{21}x_3x_4, \quad (4)$$

$$\dot{x}_5 = p_5, \quad p_5 = -k_{12}x_1x_2 + k_{21}x_3x_4 + k_{34}x_3^2 - k_{43}x_1x_5. \quad (5)$$

Each zero of the polynomials p_1, p_2, p_3, p_4, p_5 gives a concentration of species in which the system is in equilibrium. The zeros of p_1, p_2, p_3, p_4, p_5 are called the steady states of the chemical reaction network. Accordingly, the ideal $\langle p_1, p_2, p_3, p_4, p_5 \rangle \subseteq \mathbb{Q}[k_{12}, k_{21}, k_{34}, k_{43}, x_1, x_2, x_3, x_4, x_5]$ is called the *steady state ideal* of the chemical reaction network. We consider the coefficient field \mathbb{Q} because of computability issue. Otherwise, theoretically, our results hold for any coefficient field. The solutions of these polynomials can be in \mathbb{R} or in \mathbb{C} .

For a thorough introduction to chemical reaction network theory, we refer to Feinberg's Book [17] and his lecture notes [16]. We follow the notation of Feinberg's book in this article.

An ideal is called binomial if it is generated by a set of binomials. In this article we investigate whether the steady state ideal of a given chemical reaction network is binomial. We are interested in efficient algorithms for testing binomiality. Consider the steady state ideal

$$I = \langle p_1, p_2, p_3, p_4, p_5 \rangle \subseteq \mathbb{Q}[k_{12}, k_{21}, k_{34}, k_{43}, x_1, x_2, x_3, x_4, x_5], \quad (6)$$

given by Equations (1)–(5). Reducing p_1, p_3 and p_4 with respect to p_2 and p_5 , we have

$$I = \langle -k_{12}x_1x_2 + k_{21}x_3x_4, -k_{34}x_3^2 + k_{43}x_1x_5 \rangle, \quad (7)$$

which shows that the ideal I is binomial. In this article, we work over the ring $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$ and investigate binomiality over this ring.

Note that in the literature there exist also slightly different notions of binomiality. Eisenbud and Sturmfels in [12] call an ideal binomial if it is generated by polynomials with at most two terms. Following this definition, some authors, e.g., Dickenstein et al. in [32] have considered the steady state ideal as an ideal in the ring $\mathbb{Q}(k_{ij})[x_1, \dots, x_n]$ and studied the binomiality of these ideals in $\mathbb{R}[x_1, \dots, x_n]$ after specialising k_{ij} with positive real values. In order to distinguish between the two notions, we call *unconditionally binomial* a steady state ideal that is binomial in $\mathbb{Q}[k, x]$ (the notion used in this paper) and *conditionally binomial* a steady state ideal that is binomial in $\mathbb{Q}(k)[x]$, i.e. for specified parameters k (the notion used in [32]).

The notions of binomial ideals and toric varieties have roots in thermodynamics, dating back to Boltzmann. Binomiality corresponds to detailed balance, which for reaction networks means that at thermodynamic equilibrium the forward and backward rates should be equal for all reactions. Detailed balance is a very important concept in thermodynamics, for instance it has been used by Einstein in his Nobel prize winning theory of the photoelectric effect [11], by Wegscheider in his thermodynamic theory of chemical reaction networks [36] and by Onsager for deriving his famous reciprocity relations [33]. Because detailed balance implies time reversal symmetry, systems with detailed balance can not produce directed movement and can only dissipate heat. This is important in applications, for instance in molecular biology, where molecular motors can not function with detailed balance. Although most interesting molecular devices function without detailed balance and binomiality, some of their subsystems can satisfy these conditions. The interest of studying binomiality relies in the simplicity of the analysis of such subsystems. For instance, important properties such as multistationarity and stability are easier to establish for binomial systems. Toricity, also known as complex, or cyclic, or semi-detailed balance is also known since Boltzmann that has used it as a sufficient condition for deriving his famous H-theorem [1]. Binomiality implies toricity, but the converse is not true: in order to have binomiality, a toric system must obey constraints on the rates constants, such as the well known Wegscheider-Kolmogorov condition asking for the equality of the products of forward and backward rates constants in cycles of reversible reactions. In this paper we focus on the situation when detailed balance is satisfied without conditions on the rate constants.

Detecting binomiality of an ideal, particularly of a steady state ideal, is a difficult problem, both from a theoretical and a practical point of view. The problem is typically solved by computing a Gröbner basis, which is EXPSPACE-complete [29]. Recent linear algebra approaches for solving the problem in a different setting than our problem construct large matrices which also points at the difficulty of the problem [30,6].

There is quite comprehensive literature on chemical reaction network theory. An excellent reference to this topic is [17,16]. As mathematical concepts, binomiality and toricity have been widely studied and their properties have

been investigated by various authors, e.g., Fulton [18], Sturmfels [35], Eisenbud et al. [12]. Binomiality and toricity show up quite often in chemical reaction networks. Binomiality in the case of *detailed balancing* of reversible chemical reactions has been studied by Gorban et al. [20,21] and Grigoriev and Weber [25]. Feinberg [15] and Horn and Jackson [26] have studied toric dynamical systems. Gatermann et al. studied *deformed toricity* in [19]. Craciun, et al. have considered the toricity problem over the real numbers in [7] and have presented several interesting results in this regard, among them, they have shown that *complex balanced systems* are the same as toric dynamical systems, although *toric steady states* are different from that. It has been shown in [9,10] that the binomial structure will imply much simpler criteria for multistationarity. These results give strong motivation for one to study algorithms for detecting binomial networks. Especially, in [9], the authors defined *linearly binomial network* and they proposed sufficient conditions for a network to be linearly binomial. The proof is constructive even though it has not been presented as an algorithm. Their method is also quite straightforward and can handle more general networks in many applications.

Dickenstein et al. have presented sufficient linear algebra conditions with inequalities for binomiality of the steady state ideals in [30]. Their idea has been developed in [31], where the concept of MESSI reactions has been introduced. Conradi and Kahle have proved in [6] that for homogenous ideals (i.e. for chemical reaction networks without zero-order reactions), the sufficient condition of Dickenstein et al. is necessary as well and also introduced an algorithm for testing binomiality of homogenous ideals. As many biochemical networks are not homogeneous, the algorithm requires heuristics in such cases. The algorithm has been implemented in Maple and Macaulay II in [28,27] and experiments have been carried out on several biological models. Grigoriev et al. in [23] have considered the toricity of steady state ideals from a geometric point of view. Introducing shifted toricity, they presented algorithms, complexity bounds as well as experimental results for testing toricity using two important tools from symbolic computation, quantifier elimination [8,22,37] and Gröbner bases [4,5,13,14]. Recently, first order logic test for toricity have been introduced [34].

The main idea of this article is to consider the generators of the steady state ideal as sums of the binomials associated to the reactions rather than the monomials associated to the complexes. This is feasible for a reversible chemical reaction network. Following the above observation and assigning a binomial to each reaction, one can write the generators of the steady state ideal as sums of those binomials with integer coefficients.

As our main result, we have proved that a reversible chemical reaction network is unconditionally binomial if and only if it is “linearly” binomial (i.e., there exist linear combinations of the generators such that these combinations are binomials). More precisely, having represented of the generators of the steady state ideal as sum of binomials, one can test the binomiality exclusively using elementary row and column operations on the coefficient matrix of these binomials.

This can be done by computing the reduced row echelon form of the coefficient matrix, which yields an efficient method for testing binomiality.

Our main contributions in this article are the following.

1. We introduce a new representation of the generators of the steady state ideal of a reversible chemical reaction as a sum of certain binomials rather than monomials.
2. Using that representation, we assign a matrix with entries in \mathbb{Z} to a reversible chemical reaction network, such that the binomiality of the steady state ideal can be tested by computing the reduced row echelon form of this matrix.
3. We prove a worst-case upper bound on the time complexity of our binomiality test. We experimentally compare our test with the existing binomiality tests in the literature, which demonstrates the applicability of our method.

Our representation of the steady state ideal as a sum of certain binomials, as well as the matrices associated to them are further original ideas presented in this paper. While typically complex-species matrices are used for testing binomiality, we use reaction-species matrices for this purpose.


The plan of the article is as follows. Section 1 gives an introduction to the necessary concepts of chemical reaction network theory, reviews the literature and presents the idea of this work. Section 2 includes the main definitions and results. In this section we show our representation of the generators of the steady state ideal of a reversible chemical reaction network and present our algorithm for testing binomiality. In Section 3, we discuss the complexity of our method. We furthermore compare our algorithm with other existing algorithms in the literature via experiments. In Section 4 we summarise our results and draw some conclusions.

2 Testing Binomiality

In this section, we present our main result based on which we present an algorithm for testing unconditional binomiality of reversible chemical reaction networks. In Subsection 2.1 we introduce a representation for the generators of the steady state ideal of a chemical reaction network as sum of binomials. We show that this representation is unique for reversible reaction networks, considering rate constants as indeterminates. In Subsection 2.2, we define a matrix associated to a chemical reaction network which is essentially the species–reaction matrix, rather than the stoichiometric matrix which is the species–complex matrix. Having considered constant rates as indeterminates, the uniqueness of our matrix for reversible reactions comes from the uniqueness of representing of the generators of the steady state ideal as sum of binomials.

2.1 Sum of Binomial Representation

Consider the following reversible reaction between two complexes C_1 and C_2 .



Let m_i , $i = 1, 2$, be the product of the concentrations of the species in C_i with the stoichiometric coefficients as the powers. We call m_i the monomial associated to C_i . Also let x_1 be the concentration of a species that is in C_1 with the stoichiometric coefficient α_1 and is not in C_2 . The differential equation describing the kinetics of this species is

$$\dot{x}_1 = -\alpha_1 k_{12} m_1 + \alpha_1 k_{21} m_2. \quad (8)$$

For a species in C_2 with stoichiometric coefficient α_2 which is not in C_1 with the concentration x_2 , the differential equation will be


$$\dot{x}_2 = \alpha_2 k_{12} m_1 - \alpha_2 k_{21} m_2. \quad (9)$$

For a species with concentration x_3 that appears in both C_1 and C_2 , the differential equation will be $\dot{x}_3 = c(k_{12}m_1 - k_{21}m_2)$, where $c \in \mathbb{Z}$ is the difference between the corresponding stoichiometric coefficients in C_2 and C_1 . Set $b_{12} := -k_{12}m_1 + k_{21}m_2$ and $b_{21} := k_{12}m_1 - k_{21}m_2$. The steady state ideal of the above chemical reaction network is $\langle \alpha_1 b_{12}, \alpha_2 b_{21} \rangle$, which is equal to $\langle b_{12} \rangle$, since $b_{12} = -b_{21}$.

For a reversible reaction network with more than one reaction, one can associate a binomial of the form $b_{ij} := k_{ij}m_i - k_{ji}m_j$ to each reaction. Then the polynomials generating the steady state ideal can be written as sums of b_{ij} with integer coefficients. We make this more precise in the following definition.

Definition 1. Let \mathcal{C} be a reversible chemical reaction network with the complexes C_1, \dots, C_s , let k_{ij} , $1 \leq i \neq j \leq s$, be the rate constant of the reaction from C_i to C_j , and let x_1, \dots, x_n be the concentrations of the species in the chemical reaction network. We call a monomial m_i the monomial associated to C_i if m_i is the product of the concentrations of those species that appear in C_i with the stoichiometric coefficients of the species as the powers. If there is a reaction between C_i and C_j , then $b_{ij} := -k_{ij}m_i + k_{ji}m_j$ is called the binomial associated to the reaction from C_i to C_j , otherwise $b_{ij} := 0$.

Example 1. Recall the following chemical reaction network from Section 1:



Following the notation in Section 1, let x_1, x_2, x_3, x_4, x_5 be the concentrations of CO_2 , H_2 , CO , H_2O and C , respectively. The monomials associated to the complexes $CO_2 + H_2$, $CO + H_2O$, $2CO$ and $CO_2 + C$ are x_1x_2 , x_3x_4 , x_3^2 and x_1x_5 , respectively. The binomials associated to the two reactions in this network are $b_{12} = -k_{12}x_1x_2 + k_{21}x_3x_4$ and $b_{34} = -k_{34}x_3^2 + k_{43}x_1x_5$. As there is no reaction between the first and third complexes we have $b_{13} = b_{31} = 0$. Similarly, $b_{23} = b_{32} = 0$, $b_{14} = b_{41} = 0$ and $b_{24} = b_{42} = 0$. Also, by definition, $b_{21} = -b_{12}$,

$b_{34} = -b_{43}$, etc.. Using the binomials associated to the reactions, one can write the polynomials generating the steady state ideal as

$$p_1 = b_{12} - b_{34} \tag{10}$$

$$p_2 = b_{12} \tag{11}$$

$$p_3 = -b_{12} + 2b_{34} \tag{12}$$

$$p_2 = -b_{12} \tag{13}$$

$$p_2 = -b_{34}. \tag{14}$$

Hence, the steady state ideal can be written as

$$\langle p_1, p_2, p_3, p_4, p_5 \rangle = \langle b_{12}, b_{34} \rangle. \tag{15}$$


As Example 1 and the definition of the binomials b_{ij} in Definition 1 suggests one can write the generators of the steady state ideal of every reversible chemical reaction networks as sums of b_{ij} with integer coefficients, i.e., assuming that \mathcal{R} is the set of reactions in the chemical reaction network

$$\dot{x}_k = p_k = \sum_{C_i \rightarrow C_i \in \mathcal{R}} c_{ij}^{(k)} b_{ij}, \tag{16}$$

for $k = 1 \dots n$ and $c_{ij}^{(k)} \in \mathbb{Z}$.

For clarification, we may remind the reader that in this article we assume working over $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$. This is the case, in particular, for Definition 1 and the discussion afterwards. In [32], the authors specialise k_{ij} with positive real values, in which case, the steady state ideal may or may not be binomial over $\mathbb{R}[x_1, \dots, x_n]$. Similarly, specialising k_{ij} in Equation 16 can result in writing p_k as sum of different binomials. In other words, if k_{ij} specialised, the representation of p_k as sum of binomials in 16 is not necessarily unique. This is illustrated in the following example.

Example 2. [32, Example 2.3] Let $C_1 = 2A$, $C_2 = 2B$ and $C_3 = A + B$. Consider the reversible chemical reaction network given by the following reactions:



Assuming x_1 and x_2 to be the concentrations of A and B , respectively, by Definition 1,

$$b_{12} = -k_{12}x_1^2 + k_{21}x_2^2 \tag{17}$$

$$b_{13} = -k_{13}x_1^2 + k_{31}x_1x_2 \tag{18}$$

$$b_{23} = k_{23}x_2^2 - k_{32}x_1x_2. \tag{19}$$

It can be checked that the generators of the steady state ideal can be written as

$$p_1 = 2b_{12} + b_{13} + b_{23} \quad (20)$$

$$p_2 = -2b_{12} - b_{13} - b_{23}. \quad (21)$$

If $k_{31} = k_{32}$ then $k_{31}x_1x_2 = k_{32}x_1x_2$, hence $k_{31}x_1x_2$ will occur in b_{13} and b_{23} with opposite signs which will be cancelled out in $b_{13} + b_{23}$, resulting in writing p_1 as sum of b_{12} and $-k_{13}x_1^2 + k_{23}x_2^2$. This is another way of writing p_1 as sum of binomials. Because binomiality relies here on the condition $k_{31} = k_{32}$, this is an example of conditional binomiality.

If we consider the rate constants k_{ij} as indeterminates, i.e., if we consider the steady state ideal as an ideal over the ring $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$, then the representation in Equation (16) as sum of binomials b_{ij} will be unique. More precisely, we have the following.

Lemma 1. *Given a reversible chemical reaction network with the notation of Definition 1, if k_{ij} are indeterminates then the generators of the steady state ideal can be uniquely written as sum of the binomials presented in Equation 16.*

Proof. Assuming that k_{ij} , $1 \leq i, j \leq s$ are indeterminates, they will be algebraically independent over $\mathbb{Q}[x_1, \dots, x_n]$. Therefore for monomials m_t and $m_{t'}$ in $\mathbb{Q}[x_1, \dots, x_n]$ associated to two distinct complexes and for all $1 \leq i, j, i', j' \leq s$, $k_{ij}m_t$ and $k_{i'j'}m_{t'}$ will be distinct monomials in $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$. Hence binomials b_{ij} associated to the reversible reactions are not only pairwise distinct, but also their monomials are pairwise distinct in $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$. This implies that the generators of the steady state ideal have unique representations in $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$ as sum of b_{ij} with integer coefficients.

Having a unique representation as in Equation 16 enables us to represent our binomial coefficient matrix, defined later, which is the base of our efficient algorithm for testing unconditional binomiality of reversible chemical reaction networks.

Considering rate constants k_{ij} as indeterminates, if a steady state ideal is unconditionally binomial, i.e., binomial in the ring $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$, then its elimination ideal is binomial in the ring $\mathbb{Q}[x_1, \dots, x_n]$. Indeed, the elimination of a binomial ideal is a binomial ideal. This can be seen from elimination property of Gröbner bases. Authors of [12] have studied binomial ideals and their properties intensively. In particular Corollary 1.3 in the latter article state the binomiality of the elimination ideal of a binomial ideal. We remind the reader that the definition of binomiality in this article is different from [12]. In the latter, binomial ideals have binomial and monomial generators, however in the current article, we only consider binomial generators. Restricting the definition of binomial ideal to the ideals with only binomial generators, most of the result in [12] still holds, in particular the one about the elimination of binomial ideals. Therefore, if the steady state ideal of a chemical reaction network is binomial in $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$, then its elimination $I \cap \mathbb{Q}[x_1, \dots, x_n]$ is also a binomial ideal.

Geometrically, the above discussion can be explained via projection of the corresponding varieties. Given a chemical reaction network, assume that reaction rates k_{ij} are indeterminates and let the number of k_{ij} be t . Let V denote the steady state variety, i.e., the variety of the steady state ideal. V is a Zariski closed subset of \mathbb{K}^{t+n} , where \mathbb{K} is an appropriate field (e.g., \mathbb{C}). If V is a coset of a subgroup of the multiplicative group $(\mathbb{K}^*)^{t+n}$, then the projection of V into the space generated by $x_1 \dots, x_n$, i.e., $V \cap (\mathbb{K}^*)^n$ is also a coset. In particular, the projection of a group is a group. Since the variety of a binomial ideal is a coset [23,24], the projection of the variety of a binomial ideal is the variety of a binomial ideal. As special cases, the projection of a toric variety, a shifted toric variety and a binomial variety (defined in [23,24]) is a toric, a shifted toric and a binomial variety, respectively. For a detailed study of toricity of steady state varieties, we refer to [23].

Remark 1. – We may mention that in [7], the authors have studied *toric dynamical systems*, where they have considered working over $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$ and presented several interesting results. In particular, Theorem 7 in that article states that a chemical reaction network is toric if and only if the rate constants lie in the variety of a certain ideal in $\mathbb{Q}[k_{ij}]$, called the *moduli ideal*.
– Toric dynamical systems are known as *complex balancing* mass action systems [7].

2.2 The Algorithm

Definition 2. Let \mathcal{C} be a reversible chemical reaction network as in Definition 1 and assume that the generators of its steady state ideal are written as the linear combination of the binomials associated to its reactions as in Equation 16, i.e.,

$$p_k = \sum_{C_i \rightarrow C_i \in \mathcal{R}}^s c_{ij}^{(k)} b_{ij} \quad \text{for } k = 1, \dots, n.$$

We define the binomial coefficient matrix of \mathcal{C} to be the matrix whose rows are labeled by p_1, \dots, p_n and whose columns are labeled by non-zero b_{ij} and the entry in row p_k and column b_{ij} is $c_{ij}^{(k)} \in \mathbb{Z}$.

By the definition, the binomial coefficient matrix of a reversible chemical reaction network is the coefficient matrix of the binomials that occur in the representation of the generators of the steady state ideal as sum of binomials. As we consider k_{ij} indeterminates, the representation of the generators of the steady state ideal of a given complex is unique, which implies that the binomial coefficient matrix of a given complex is unique too.

Example 3. Consider the chemical reaction network in Example 1, with generators of the steady state ideal as follows.

$$p_1 = b_{12} - b_{34} \quad (22)$$

$$p_2 = b_{12} \quad (23)$$

$$p_3 = -b_{12} + 2b_{34} \quad (24)$$


$$p_4 = -b_{12} \quad (25)$$

$$p_5 = -b_{34}. \quad (26)$$

The binomial coefficient matrix of this chemical reaction network is

$$M = \begin{matrix} & & b_{12} & b_{34} \\ & p_1 & \begin{pmatrix} 1 & -1 \\ 1 & 0 \\ -1 & 2 \\ -1 & 0 \\ 0 & -1 \end{pmatrix} \\ & p_2 & \\ & p_3 & \\ & p_4 & \\ & p_5 & \end{matrix}. \quad (27)$$

Another simple example is the reaction



with the binomial associated to it as $b_{12} := -k_{12}x_1^4 + k_{21}x_1x_2$, where x_1 is the concentration of A and x_2 is the concentration of B . The steady state ideal is generated by $\{3b, -b\}$, and the binomial coefficient matrix for this network is $\begin{pmatrix} 3 \\ -1 \end{pmatrix}$.

One can test binomiality of the steady state ideal of a reversible reaction network using its binomial coefficient matrix.

Theorem 1. *The steady state ideal of a reversible chemical reaction network is unconditionally binomial, i.e., binomial in $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$, if and only if the reduced row echelon form of its binomial coefficient matrix has at most one non-zero entry at each row.*

Proof. Let $G = \{p_1, \dots, p_n\} \subseteq \mathbb{Q}[k_{ij}, x_1, \dots, x_n]$ be a generating set for the steady state ideal of a given reversible chemical reaction network \mathcal{C} , and let $\{b_{ij} \mid 1 \leq i \neq j \leq s\}$ be the ordered set of non-zero binomials associated to the reactions. Fix a term order on the monomials in $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$.

First we prove that if the reduced row echelon form of the binomial coefficient matrix has at most one non-zero entry at each row, then the steady state ideal is binomial. The proof of this side of the proposition comes from the definition of reduced row echelon form. In fact, the reduced row echelon form of the binomial coefficient matrix of \mathcal{C} can be computed by row reduction in that matrix, which is equivalent to the reduction of the generators of the steady state ideal with respect to each other. Therefore, computing the reduced row echelon form of the binomial coefficient matrix and multiplying it with the vector of binomials b_{ij} , one can obtain another basis for the steady state ideal. Having this, if the

reduced row echelon form has at most one non-zero entry at each row, then the new basis for the steady state ideal will only include b_{ij} . Therefore the steady state ideal will be binomial.

Now we prove the “only if” part of the proposition, that is, if the steady state ideal of \mathcal{C} is binomial, then the reduced row echelon form of the binomial coefficient matrix has at most one non-zero entry at each row. We claim that for each pair of polynomials $p_t, p_m \in G$, p_t is reducible with respect to p_m if and only if there exists a binomial b_{ij} that occurs in both p_t and p_m and includes their leading terms. The “only if” part of the claim is obvious. To prove the “if” part of the claim, let p_m be reducible with respect to p_t . Then the leading term of p_m divides the leading term of p_t . Since the leading terms are multiples of k_{ij} and these are disjoint indeterminates, this is only possible if both of the leading terms are equal. If the leading terms are equal, then b_{ij} in which the leading terms occur, must itself occur in both p_t and p_m . Therefore p_t and p_m share a binomial associated to a reaction, which is in contradiction with our assumption.

From the above claim and the definition of the reduced row echelon form one can see that p_1, \dots, p_n are pairwise irreducible if and only if the binomial coefficient matrix of \mathcal{C} is in reduced row echelon form.

Now we prove that p_1, \dots, p_n are pairwise irreducible if and only if they form a Gröbner basis in which polynomials are pairwise irreducible. Note that this does not necessarily imply that G is a reduced Gröbner basis, as p_i are not necessarily monic. Assume that p_1, \dots, p_n are pairwise irreducible. We prove that the greatest common divisor of each pair of the leading terms of the p_1, \dots, p_n is 1. By contradiction, assume that there exists a monomial not equal to 1 which divides the leading terms of both p_t, p_m , for $1 \leq t, m \leq n$. Then there exists a variable x_l such that x_l divides the leading terms of p_t and p_m . Since each leading term is the monomial associated to a complex, the species with concentration x_l occurs in two complexes with associated monomials as the leading terms of p_t and p_m . Then both p_t and p_m have as their summand the binomials that are associated to the reactions including those complexes. As for each complex there exists at least one binomial associated, both p_m and p_t have as a summand one common binomial b_{ij} . However, we had already proved that this implies that p_t and p_m are not pairwise irreducible, which is a contradiction to the assumption that the greatest common divisor of the leading terms of p_t and p_m is not 1. Now by Buchberger’s first criterion if the greatest common divisor of the leading terms of each pair of polynomials in G is 1 then G is a Gröbner basis. The other side of this claim is obvious.

From what we have proved until now, we can conclude that the binomial coefficient matrix of \mathcal{C} is in reduced row echelon form if and only if G is a Gröbner basis with pairwise irreducible elements. On the other hand, by a result of Eisenbud and Sturmfels [12], the steady state ideal of \mathcal{C} is binomial if and only if every Gröbner basis of it includes binomials. Therefore we conclude that the steady state ideal is binomial if and only if the reduced row echelon form of the binomial coefficient matrix has at most one non-zero entry in each row.

Otherwise, one can consider computing \tilde{M} as a preprocessing step and run another method, e.g., Gröbner bases, quantifier elimination as in [23], or the method in Dickenstein, et al [30].

3 Complexity & Comparisons

Proposition 1. *Let r be the number of reactions and n be the number of species of a reversible chemical reaction network \mathcal{C} . The asymptotic worst case time complexity of testing unconditional binomiality of the steady state ideal of \mathcal{C} via Algorithm 1 can be bounded by $\mathcal{O}(\max(r, n)^\omega)$ where $\omega \approx 2.3737$, which is also the complexity of matrix multiplication.*

Proof. The operations in steps 1–4 and 7–11 are at most linear in terms of r and n . Since M is a matrix of size $n \times r$, where $r = |b_{ij}|$, and B is a vector of size r , computing reduced row echelon form in step 5 and also the matrix multiplication in step 6 will cost at most $\mathcal{O}(\max(r, n)^\omega)$. Therefore the total number of operations in the algorithm can be bounded by $\mathcal{O}(\max(r, n)^\omega)$.

In [23, Section 4] it has been shown that there exists an exponential asymptotic worst case upper bound on the time complexity of testing toricity. An immediate consequence of that result is that the time complexity of testing binomiality can be bounded by the same exponential function. Following the arguments in [23, Section 4], one can show that there exists an algorithm for testing binomiality over $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$ and $\mathbb{Q}[x_1, \dots, x_n]$ simultaneously, with an exponential upper bound for the worst case time complexity.

As mentioned earlier in Section 2, the reduced Gröbner basis of a binomial ideal, with respect to every term order, includes only binomials. This directly can be seen from running Buchberger’s algorithm and that S-polynomials and their reductions by binomials are binomial. Eisenbud et. al’ article [12], with a slightly different definition of binomial ideals, investigates many properties of binomial ideals using the latter fact. Following this fact, a typical method for testing binomiality is via computing a reduced Gröbner basis of a steady state ideal $I \subseteq \mathbb{Q}[k_{ij}, x_1, \dots, x_n]$. The drawback of computing Gröbner bases is that this is EXPSPACE-complete [29]. So our algorithm is asymptotically considerably more efficient than Gröbner basis computation.

Example 5 (Models from the BioModels Repository⁴).

- There are twenty non-reversible biomodels in which Gröbner basis computations done in [23] for testing conditional binomiality do not terminate in a six-hour time limit, however our algorithm terminates in less than three seconds. Also there are six cases in which Gröbner basis computations terminate in less than six hours, but are at least 1000 times slower than our algorithm. Finally there are ten models in which Gröbner basis is at least 500 times slower than our computations.

⁴ <https://www.ebi.ac.uk/biomodels/>

- There are sixty nine biomodels that are not considered for computation in [23] because of the of the unclear numeric value of their rate constants. Our computations on almost all of those cases terminated in less than a second.
- (Reversible models from the BioModels Repository) Biomodels 491 and 492 are both reversible. Biomodel 491 has 52 species and 86 reactions. The binomial coefficient matrix of this biomodel has size 52×86 and has ± 1 entries. A reduced row echelon form computation in Maple reveals in 0.344 seconds that it is unconditionally binomial, while a Gröbner basis computation takes more than 12 seconds to check its conditional binomiality. BioModel 492 has also 52 species, and includes 88 reactions. The binomial coefficient matrix has entries ± 1 and is of size 52×88 . This biomodel is also unconditionally binomial. It takes 0.25 seconds for Maple to check its unconditional binomiality via Algorithm 1 in Maple, while a Gröbner basis computation takes near 18 seconds, as one can see in the computations in [23, Table 3], which show the group structure of the steady state varieties of the models.

Dickenstein et al. in [30] have proposed a method for testing toricity of a chemical reaction network. The definitions and purpose of that work are slightly different from our article, hence comparisons between those two methods should be treated with caution. While we focus on unconditional binomiality of the steady state ideals of reversible reaction networks, , i.e., binomiality in $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$, with the aim of efficiency of the computations, the authors of the above article are interested in conditional binomiality with algebraic dependencies between k_{ij} such that the elimination ideal is binomial. Having mentioned that, our method leads to the computation of reduced row echelon form of a matrix of size $n \times r$ with integer entries which is polynomial time, while Theorem 3.3. in [30] requires constructing a matrix of size $n \times s$ with entries from $\mathbb{Z}[k_{ij}]$ and finding a particular partition of its kernel.

Considering Example 2.3 in [32], our algorithm constructs the matrix M and its reduced row echelon form \tilde{M} :

$$M = \begin{pmatrix} 1 & 1 & -2 \\ -1 & -1 & 2 \end{pmatrix}, \quad \tilde{M} = \begin{pmatrix} 1 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}, \quad (29)$$

and we see that the steady state ideal is not unconditionally binomial over $\mathbb{Q}[k_{ij}, x_1, \dots, x_n]$. The method in [32] constructs

$$\begin{pmatrix} -2k_{12} - k_{13} & 2k_{21} + k_{23} & k_{31} - k_{32} \\ 2k_{12} + k_{13} & -2k_{21} - k_{23} & -k_{31} + k_{32} \end{pmatrix}, \quad (30)$$

and finds an appropriate partition, which shows that the steady state ideal is binomial in $\mathbb{Q}[x_1, \dots, x_n]$ if and only if $k_{31} = k_{32}$. As a larger example, consider the chemical reaction network given in Example 3.13 in [32] and assume that it is a reversible chemical reaction network. Our method constructs a matrix with entries ± 1 of size 9×8 and computes its reduced row echelon form (in this case reduced row echelon form, as entries are ± 1). The method described in [32] leads to a 9×10 matrix with entries as linear polynomials in $\mathbb{Z}[k_{ij}]$ and computes a particular partition of the kernel of the matrix.

For homogeneous ideals, Conradi and Kahle have shown in [6] that the sufficient condition for conditional binomiality in [32] is necessary, too. Their Algorithm 3.3 tests conditional binomiality of a homogeneous ideal, which can be generalised by homogenising. The algorithm computes a basis for the ideal degree by degree and performs reductions with respect to the computed basis elements at each degree step. Since our algorithm is intended for steady state ideals of reversible chemical reaction networks, which are not necessarily homogeneous, our following comparison with the Conradi–Kahle algorithm bears a risk of being biased by homogenisation. We discuss the execution of both algorithms on Example 3.15 in [32]. This chemical reaction network does not satisfy the sufficient condition presented in [32, Theorem 3.3]. Testing this condition leads to the construction of a 9×13 matrix with entries in $\mathbb{Z}[k_{ij}]$, followed by further computations, including finding a particular partition of its kernel. Theorem 3.19 in [32] is a generalisation of Theorem 3.3 there, which can test conditional binomiality of this example by adding further rows and columns to the matrix. Conradi and Kahle also treat this example with their algorithm. This requires the construction of a coefficient matrix of size 9×13 with entries in $\mathbb{Z}[k_{ij}]$ and certain row reductions. If we add reactions so that the reaction network becomes reversible, our algorithm will construct a matrix of size 9×9 with entries ± 1 and compute its reduced row echelon form to test unconditional binomiality in $\mathbb{Q}[k_{ij}, x_1, \dots, x_9]$.

4 Conclusions

Binomiality of steady state ideals is an interesting problem in chemical reaction network theory. It has a rich history and literature and is still an active research area. For instance, recently MESSI systems have been introduced [31] following the authors’ work on binomiality of a system. Finding binomiality and toricity is computationally hard from both a theoretical and a practical point of view. It typically involves computations of Gröbner bases, which is EXPSPACE-complete.

In a recent work [23] we investigated toricity of steady state varieties and gave efficient algorithms. In particular, we experimentally investigated toricity of biological models systematically via quantifier elimination. Besides that, we presented exponential theoretical bounds on the toricity problem. The current article, restricting to reversible reaction networks, aims at an efficient linear algebra approach to the problem of unconditional binomiality, which can be considered as a special case of the toricity problem.

In that course, considering rate constants as indeterminates, we assign a unique binomial to each reaction and construct the coefficient matrix with respect to these binomials. Our algorithm proposed here computes a reduced row echelon form of this matrix in order to detect unconditional binomiality. The algorithm is quite efficient, as it constructs comparatively small matrices whose entries are integers. It is a polynomial time algorithm in terms of the number of species and reactions. While other existing methods for testing conditional bino-

miality have different settings and purposes than our algorithm, for the common cases, our algorithm has advantages in terms of efficiency.

Acknowledgments

This work has been supported by the bilateral project ANR-17-CE40-0036 and DFG-391322026 SYMBIONT [2,3].

References

1. Ludwig Boltzmann. *Lectures on Gas Theory*. University of California Press, Berkeley and Los Angeles, CA, 1964.
2. François Boulier, François Fages, Ovidiu Radulescu, Satya Samal, Andreas Schuppert, Werner Seiler, Thomas Sturm, Sebastian Walcher, and Andreas Weber. The SYMBIONT project: Symbolic methods for biological networks. *ACM Communications in Computer Algebra*, 52(3):67–70, 2018.
3. François Boulier, François Fages, Ovidiu Radulescu, Satya Samal, Andreas Schuppert, Werner Seiler, Thomas Sturm, Sebastian Walcher, and Andreas Weber. The SYMBIONT project: Symbolic methods for biological networks. *F1000Research*, 7(1341), 2018.
4. Bruno Buchberger. *Ein Algorithmus zum Auffinden der Basiselemente des Restklassenringes nach einem nulldimensionalen Polynomideal*. Doctoral dissertation, Mathematical Institute, University of Innsbruck, Austria, 1965.
5. Bruno Buchberger. Ein Algorithmisches Kriterium für die Lösbarkeit eines algebraischen Gleichungssystems. *Aequationes Mathematicae*, 3:374–383, 1970.
6. Carsten Conradi and Thomas Kahle. Detecting binomiality. *Advances in Applied Mathematics*, 71:52–67, 2015.
7. Gheorghe Craciun, Alicia Dickenstein, Anne Shiu, and Bernd Sturmfels. Toric dynamical systems. *J. Symb. Comput.*, 44(11):1551–1565, November 2009.
8. James H. Davenport and Joos Heintz. Real quantifier elimination is doubly exponential. *J. Symb. Comput.*, 5(1–2):29–35, 1988.
9. Alicia Dickenstein, Mercedes Pérez Millán, Anne Shiu and Xiaoxian Tang. Multistationarity in Structured Reaction Networks. *Bulletin of Mathematical Biology*. 81: 1527–1581, 2019.
10. AmirHossein Sadeghimanesh and Elisenda Feliu. The multistationarity structure of networks with intermediates and a binomial core network. *Bulletin of Mathematical Biology*. 81: 2428–2462, 2019.
11. Albert Einstein. Strahlungs-emission und -absorption nach der Quantentheorie. *Verh. Dtsch. Phys. Ges.*, 18:318–323, 1916.
12. David Eisenbud and Bernd Sturmfels. Binomial ideals. *Duke Math. J.*, 84(1):1–45, 1996.
13. Jean-Charles Faugère. A new efficient algorithm for computing Gröbner bases (F4). *J. Pure Appl. Algebra*, 139(1):61–88, 1999.
14. Jean-Charles Faugère. A new efficient algorithm for computing gröbner bases without reduction to zero (F5). In *Proc. ISSAC 2002*, pages 75–83. ACM, New York, NY, 2002.
15. Martin Feinberg. Complex balancing in general kinetic systems. *Arch. Ration. Mech. An.*, 49(3):187–194, 1972.

16. Martin Feinberg. Lectures on chemical reaction networks, 1979.
17. Martin Feinberg. *Foundations of Chemical Reaction Network Theory*, volume 202 of *Applied Mathematical Sciences*. Springer, 2019.
18. William Fulton. *Introduction to Toric Varieties*, volume 131 of *Annals of Mathematics Studies*. Princeton University Press, 1993.
19. Karin Gatermann. Counting stable solutions of sparse polynomial systems in chemistry. In *Symbolic Computation: Solving Equations in Algebra, Geometry, and Engineering*, volume 286 of *Contemporary Mathematics*, pages 53–69. AMS, Providence, RI, 2001.
20. A. N. Gorban and V. N. Kolokoltsov. Generalized mass action law and thermodynamics of nonlinear Markov processes. *Math. Model. Nat. Phenom.*, 10(5):16–46, 2015.
21. A. N. Gorban and G. S. Yablonsky. Three waves of chemical dynamics. *Math. Model. Nat. Phenom.*, 10(5):1–5, 2015.
22. D. Yu. Grigoriev. Complexity of deciding Tarski algebra. *J. Symb. Comput.*, 5(1–2):65–108, 1988.
23. Dima Grigoriev, Alexandru Iosif, Hamid Rahkooy, Thomas Sturm, and Andreas Weber. Efficiently and effectively recognizing toricity of steady state varieties. *CoRR*, abs/1910.04100, 2019.
24. Dima Grigoriev and Pierre D. Milman. Nash resolution for binomial varieties as Euclidean division. A priori termination bound, polynomial complexity in essential dimension 2. *Adv. Math.*, 231(6):3389–3428, 2012.
25. Dima Grigoriev and Andreas Weber. Complexity of solving systems with few independent monomials and applications to mass-action kinetics. In *Proc. CASC 2012*, volume 7442 of *LNCS*, pages 143–154. Springer, 2012.
26. F. Horn and R. Jackson. General mass action kinetics. *Arch. Ration. Mech. An.*, 47(2):81–116, 1972.
27. Alexandru Iosif and Hamid Rahkooy. Analysis of the Conradi–Kahle algorithm for detecting binomiality on biological models. *CoRR*, abs/1912.06896, 2019.
28. Alexandru Iosif and Hamid Rahkooy. MapleBinomials, a Maple package for testing binomiality of ideals. <http://doi.org/10.5281/zenodo.3564428>, 2019.
29. Ernst W. Mayr and Albert R. Meyer. The complexity of the word problems for commutative semigroups and polynomial ideals. *Adv. Math.*, 46(3):305–329, 1982.
30. Mercedes Pérez Millán, Alicia Dickenstein, Anne Shiu, and Carsten Conradi. Chemical reaction systems with toric steady states. *Bulletin of Mathematical Biology*, 74(5):1027–1065, 2012.
31. Mercedes Pérez Millán and Alicia Dickenstein. The structure of MESSI biological systems. *SIAM J. Appl. Dyn. Syst.*, 17(2):1650–1682, 2018.
32. Mercedes Pérez Millán, Alicia Dickenstein, Anne Shiu, and Carsten Conradi. Chemical reaction systems with toric steady states. *Bull. Math. Biol.*, 74(5):1027–1065, 2012.
33. Lars Onsager. Reciprocal relations in irreversible processes. I. *Phys. Rev.*, 37(4):405, 1931.
34. Hamid Rahkooy and Thomas Sturm. First-order tests for toricity. *CoRR*, abs/2002.03586, 2020.
35. Bernd Sturmfels. *Gröbner Bases and Convex Polytopes*, volume 8 of *University Lecture Series*. AMS, Providence, RI, 1996.
36. Rudolf Wegscheider. Über simultane Gleichgewichte und die Beziehungen zwischen Thermodynamik und Reaktionskinetik homogener Systeme. *Monatsh. Chem. Verw. Tl.*, 22(8):849–906, 1901.

37. Volker Weispfenning. The complexity of linear problems in fields. *J. Symb. Comput.*, 5(1-2):3-27, 1988.