

# A NOTE ON STABLE COMMUTATOR LENGTH IN BRAIDED PTOLEMY-THOMPSON GROUPS

SHUHEI MARUYAMA

ABSTRACT. In this note, we show that the sets of all stable commutator lengths in the braided Ptolemy-Thompson groups are equal to non-negative rational numbers.

## 1. INTRODUCTION

Let  $G$  be a group and  $[G, G]$  its commutator subgroup. For an element  $g \in [G, G]$ , its commutator length  $\text{cl}(g)$  is defined by  $\text{cl}(g) = \min\{l \mid g = [a_1, b_1] \cdots [a_l, b_l]\}$ . For  $g \in G$  with  $g^k \in [G, G]$  for some positive integer  $k$ , its stable commutator length  $\text{scl}(g)$  is defined by  $\text{scl}(g) = \lim_{n \rightarrow \infty} \text{cl}(g^{kn})/kn$ . For  $g \in G$  with  $g^k \notin [G, G]$  for any  $k$ , we put  $\text{scl}(g) = \infty$ . We say that the group  $G$  has a gap in stable commutator length if there is a positive constant  $c$  such that either  $\text{scl}(g) \geq c$  or  $\text{scl}(g) = 0$  holds for any  $g \in G$ .

The values of stable commutator length in many classes of groups has been studied. For example, all stable commutator lengths are rational in the central extension of the Thompson group  $T$  corresponding to the Euler class (Ghys-Sergiescu[17]), free groups (Calegari[5]), free products of some classes of groups (Walker[22], Calegari[6], Chen[8]), amalgams of free abelian groups (Susse[21]), and certain graphs of groups (Clay-Forester-Louwsma[12], Chen[10]). It is known that there is a finitely presented group which has an irrational stable commutator length (Zhuang[23]). The gap in many classes of groups has also been studied. For example, the above central extension of the Thompson group has no gap, and the following groups have a gap; free groups (Duncan-Howie[13]), word-hyperbolic groups (Calegari-Fujiwara[7]), finite index subgroups of the mapping class group of a closed orientable surface (Bestvina-Bromberg-Fujiwara[3]), right-angled Artin groups (Heuer[18]), and the fundamental group of closed oriented connected 3-manifolds (Chen-Heuer[11]).

In this note, we show that all stable commutator lengths in the braided Ptolemy-Thompson groups  $T^*$  and  $T^\sharp$  (see Section 2.3) are rational and these groups have no gap. More precisely, we show the following theorem.

**Theorem 1.1.** The sets  $\text{scl}(T^*)$  and  $\text{scl}(T^\sharp)$  of all stable commutator lengths in the braided Ptolemy-Thompson groups are equal to the non-negative rational numbers  $\mathbb{Q}_{\geq 0}$ .

## 2. PRELIMINARIES

**2.1. quasi-morphisms and Bavard's duality theorem.** Let  $G$  be a group. A map  $\phi : G \rightarrow \mathbb{R}$  is a quasi-morphism if

$$D(\phi) = \sup_{g,h \in G} |\phi(gh) - \phi(g) - \phi(h)|$$

is finite. This  $D(\phi)$  is called the defect of  $\phi$ . A quasi-morphism is homogeneous if the condition  $\phi(g^n) = n\phi(g)$  holds for any  $g \in G$  and  $n \in \mathbb{Z}$ . Let  $Q(G)$  denote the  $\mathbb{R}$ -vector space consisting of all homogeneous quasi-morphisms on  $G$ . The key ingredient to calculate the stable commutator length is the following Bavard's duality theorem.

**Theorem 2.1.** [2] Let  $G$  be a group and  $g \in [G, G]$ , then

$$\text{scl}(g) = \sup_{\phi \in Q(G)} \frac{|\phi(g)|}{2D(\phi)}.$$

This theorem allows us to calculate the stable commutator length in groups with few homogeneous quasi-morphisms. For example, any homogeneous quasi-morphism on the infinite braid group  $B_\infty$  is equal to the abelianization homomorphism  $B_\infty \rightarrow \mathbb{Z}$  up to constant multiple (see Kotschick[19]). Thus, we have  $\text{scl}(g) = 0$  for  $g \in [B_\infty, B_\infty]$ . The central extension  $\tilde{T}$  of the Thompson group  $T$  corresponding to the Euler class is also a group with few quasi-morphisms. It is known that any homogeneous quasi-morphism on  $\tilde{T}$  is equal to the rotation number up to constant multiple. Ghys-Sergiescu[17] showed that the set of all rotation numbers on  $\tilde{T}$  coincides with the rational numbers  $\mathbb{Q}$ . Since  $\tilde{T} = [\tilde{T}, \tilde{T}]$  and the defect of rotation number is equal to 1, we have  $\text{scl}(\tilde{T}) = \mathbb{Q}_{\geq 0}$ .

**2.2. Central extensions of the Thompson group  $T$ .** Let  $n$  be an integer and  $0 \rightarrow \mathbb{Z} \rightarrow T_n \xrightarrow{p} T \rightarrow 1$  denote the central  $\mathbb{Z}$ -extension of  $T$  corresponding to  $n$  times the Euler class  $e \in H^2(T; \mathbb{Z})$ . Let  $\chi$  be a bounded cocycle representing the Euler class  $e \in H^2(T; \mathbb{Z})$ . Then the extension  $T_n$  can be obtained as a group  $T \times \mathbb{Z}$  with the product

$$(s, i)(t, j) = (st, i + j + n\chi(s, t)).$$

Since the Thompson group  $T$  is uniformly perfect and the Euler class  $e$  is bounded, there is a unique homogeneous quasi-morphism  $\phi_n \in Q(T_n)$  satisfying  $-\delta\phi_n = np^*e_b \in H_b^2(T_n; \mathbb{R})$  (see Barge-Ghys[1]). Here  $e_b \in H_b^2(T; \mathbb{R})$  is the bounded Euler class. The homogeneous quasi-morphism  $\phi_n$  is obtained as the homogenization of the quasi-morphism  $T_n \rightarrow \mathbb{R}; (t, j) \mapsto j$ . More explicitly, the quasi-morphism  $\phi_n$  is written as

$$\phi_n(t, j) = j + n \lim_{k \rightarrow \infty} \frac{a(t, k)}{k},$$

where  $a(t, k) = \chi(t, t) + \chi(t^2, t) + \cdots + \chi(t^{k-1}, t) \in \mathbb{Z}$ . Since  $T_1$  is equal to  $\tilde{T}$ , the homogeneous quasi-morphism  $\phi_1 : T_1 \rightarrow \mathbb{R}$  coincides with the rotation number on  $\tilde{T}$ . It is known that the defect of the rotation number on  $\tilde{T}$  is equal to 1 (see, for example, Calegari[5]), thus we have  $D(\phi_1) = 1$ .

**Lemma 2.2.** Let  $n$  be a non-zero integer, then  $D(\phi_n) = n$ .

*Proof.* Note that

$$\begin{aligned} 1 = D(\phi_1) &= \sup_{(s,i),(t,j)} |\phi_1(st, i + j + \chi(s, t)) - \phi_1(s, i) - \phi_1(t, j)| \\ &= \sup_{s,t} \left| \chi(s, t) + \lim_{k \rightarrow \infty} \frac{a(st, k) - a(s, k) - a(t, k)}{k} \right|. \end{aligned}$$

Thus we have

$$\begin{aligned} D(\phi_n) &= \sup_{(s,i),(t,j)} |\phi_n(st, i + j + \chi(s, t)) - \phi_n(s, i) - \phi_n(t, j)| \\ &= \sup_{s,t} \left| n\chi(s, t) + n \lim_{k \rightarrow \infty} \frac{a(st, k) - a(s, k) - a(t, k)}{k} \right| \\ &= nD(\phi_1) = n. \end{aligned}$$

□

**Lemma 2.3.** Let  $n$  be a non-zero integer, then  $\phi_n(T_n) = \mathbb{Q}$ .

*Proof.* Ghys-Sergiescu[17] showed that  $\phi_1(T_1)$  is equal to the rational numbers  $\mathbb{Q}$ . Thus, for any  $(t, j) \in T_1$ , the number  $\lim_{k \rightarrow \infty} (a(t, k))/k = \phi_1(t, j) - j$  is rational and therefore  $\phi_n(T_n) \subset \mathbb{Q}$  follows. Take  $q \in \mathbb{Q}$ . Then, for any  $q \in \mathbb{Q}$ , there is an element  $(t, j) \in T_1$  such that

$$\frac{q}{n} = \phi_1(t, j) = j + \lim_{k \rightarrow \infty} \frac{a(t, k)}{k}.$$

Thus we have  $q = \phi_n(t, nj)$ .

□

Since  $\dim Q(T_n) = 1$  and  $T_n = [T_n, T_n]$ , we have  $\text{scl}(T_n) = \mathbb{Q}_{\geq 0}$  for any non-zero integer  $n$ .

**2.3. The braided Ptolemy-Thompson groups.** In [14], Funar and Kapoudjian introduced two extensions  $T^*$  and  $T^\sharp$  of Thompson group  $T$  by the infinite braid group  $B_\infty$  and showed that they are finitely presented. Moreover, it is shown that their abelianization  $H_1(T^*)$  and  $H_1(T^\sharp)$  are isomorphic to  $\mathbb{Z}/12\mathbb{Z}$  and  $\mathbb{Z}/6\mathbb{Z}$  respectively, and therefore  $T^*$  and  $T^\sharp$  are not isomorphic. Let  $\diamond$  denote either  $*$  or  $\sharp$ . Since the abelianization homomorphism  $B_\infty \rightarrow \mathbb{Z}$  is  $T^\diamond$ -conjugation invariant, we have central  $\mathbb{Z}$ -extensions  $T_{\text{ab}}^\diamond$  of  $T$ , that is, we have the diagram

$$(2.1) \quad \begin{array}{ccccccc} 1 & \longrightarrow & B_\infty & \longrightarrow & T^\diamond & \longrightarrow & T \longrightarrow 1 \\ & & \downarrow & & \downarrow \rho & & \parallel \\ 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & T_{\text{ab}}^\diamond & \xrightarrow{p} & T \longrightarrow 1. \end{array}$$

The corresponding cohomology classes  $e(T_{\text{ab}}^*)$  and  $e(T_{\text{ab}}^\sharp)$  in  $H^2(T; \mathbb{Z})$  are equal to  $12e$  and  $21e$  respectively, where  $e \in H^2(T; \mathbb{Z})$  is the Euler class (see Funar-Sergiescu[16] and Funar-Kapoudjian-Sergiescu[15]). Thus  $T_{\text{ab}}^*$  is isomorphic to  $T_{12}$  and  $T_{\text{ab}}^\sharp$  is isomorphic to  $T_{21}$ .

### 3. THE STABLE COMMUTATOR LENGTH IN THE BRAIDED PTOLEMY-THOMPSON GROUPS

In this section, we show that the stable commutator lengths in  $T^\diamond$  are rational and all positive rationals are realized as the stable commutator length.

At first, we show the following proposition, which is essentially proved in Shtern[20].

**Proposition 3.1.** For an exact sequence  $1 \rightarrow K \rightarrow G \xrightarrow{p} H \rightarrow 1$ , we have an exact sequence

$$0 \rightarrow Q(H) \rightarrow Q(G) \rightarrow Q(K)^G,$$

where  $Q(K)^G$  is the vector space consisting of all  $G$ -conjugation invariant homogeneous quasi-morphisms on  $K$ .

*Proof.* Shtern[20] showed that, if  $\phi \in Q(G)$  is equal to 0 on  $K$ , there exists a homogeneous quasi-morphism  $\psi \in Q(H)$  such that  $\phi = p^*\psi$ . This implies that the above sequence is exact at  $Q(G)$ .  $\square$

Let us consider the diagram(2.1). There are homogeneous quasi-morphisms  $\phi_* = \phi_{12} \in Q(T_{12}) = Q(T_{ab}^*)$  and  $\phi_\# = \phi_{21} \in Q(T_{21}) = Q(T_{ab}^\#)$ . By Proposition 3.1 and the exact sequence  $1 \rightarrow [B_\infty, B_\infty] \rightarrow T^\diamond \xrightarrow{p} T_{ab}^\diamond \rightarrow 1$ , we have an injection  $\rho^* : Q(T_{ab}^\diamond) \rightarrow Q(T^\diamond)$ . Thus we have a non-trivial homogeneous quasi-morphism  $\psi_\diamond = \rho^*\phi_\diamond \in Q(T^\diamond)$  satisfying  $[\delta\psi_\diamond] = (p \circ \rho)^*e_b \in H_b^2(T^\diamond; \mathbb{R})$ .

**Lemma 3.2.** The dimension of  $Q(T^\diamond)$  is equal to 1.

*Proof.* Since there is a non-trivial element  $\psi \in Q(T^\diamond)$ , it is enough to show that the inequality  $\dim Q(T^\diamond) \leq 1$  holds. For an exact sequence  $1 \rightarrow B_\infty \rightarrow T^\diamond \rightarrow T \rightarrow 1$ , we have an exact sequence

$$0 \rightarrow Q(T) \rightarrow Q(T^\diamond) \rightarrow Q(B_\infty)^{T^\diamond}$$

by Proposition 3.1. Since the Thompson group  $T$  is uniformly perfect, we have  $Q(T) = 0$ . Thus the map  $Q(T^\diamond) \rightarrow Q(B_\infty)^{T^\diamond}$  is injective. The  $\mathbb{R}$ -vector space  $Q(B_\infty)$  is spanned by the abelianization homomorphism  $B_\infty \rightarrow \mathbb{Z}$  (see Kotschick[19]). Moreover, since the abelianization homomorphism is  $T^\diamond$ -conjugation invariant, we have  $Q(B_\infty) = Q(B_\infty)^{T^\diamond}$ . Thus we have  $\dim Q(T^\diamond) \leq \dim Q(B_\infty)^{T^\diamond} = \dim Q(B_\infty) = 1$ .  $\square$

*Proof of Theorem 1.1.* Note that, by Lemma 2.2 and the surjectivity of  $\rho : T^\diamond \rightarrow T_{ab}^\diamond$ , the defect  $D(\psi_*) = D(\rho^*\phi_*)$  is equal to  $D(\phi_*) = 12$  and  $D(\psi_\#) = D(\rho^*\phi_\#)$  is equal to  $D(\phi_\#) = 21$ . Since  $H_1(T^*) = \mathbb{Z}/12\mathbb{Z}$ , the power  $g^{12}$  is in  $[T^*, T^*]$  for any  $g \in T^*$ . Thus, by Lemma 3.2 and the Bavard's duality theorem, we have

$$\text{scl}(g) = \frac{\text{scl}(g^{12})}{12} = \frac{|\psi_*(g^{12})|}{288} = \frac{|\phi_*(\rho(g))|}{24}$$

for any  $g \in [T^*, T^*]$ . Thus, by Lemma 2.3, we have  $\text{scl}(T^*) = \mathbb{Q}_{\geq 0}$ . We can prove  $\text{scl}(T^\#) = \mathbb{Q}_{\geq 0}$  in the same way.  $\square$

## REFERENCES

- [1] J. Barge and É. Ghys: Cocycles d'Euler et de Maslov, *Math. Ann.* **294** (1992), 235–265.
- [2] C. Bavard: Longueur stable des commutateurs, *Enseign. Math. (2)* **37** (1991), 109–150.
- [3] M. Bestvina, K. Bromberg and K. Fujiwara: Stable commutator length on mapping class groups, *Ann. Inst. Fourier* **66** (2016), 871–898.
- [4] D. Calegari: scl, *MSJ Memoirs* **20**, Mathematical Society of Japan (2009).
- [5] D. Calegari: Stable commutator length is rational in free groups, *J. Amer. Math. Soc.* **22** (2009), 941–961.
- [6] D. Calegari: scl, sails, and surgery, *J. Topol.* **4** (2011), 305–326.
- [7] D. Calegari and K. Fujiwara: Stable commutator length in word-hyperbolic groups, *Groups Geom. Dyn.* **4** (2010), 59–90.
- [8] L. Chen: Scl in free products, *Algebr. Geom. Topol.* **18** (2018), 3279–3313.
- [9] L. Chen: Spectral gap of scl in free products, *Proc. Amer. Math. Soc.* **146** (2018), 3143–3151.
- [10] L. Chen: Scl in graphs of groups, arXiv:1904.08360.
- [11] L. Chen and N. Heuer: Spectral gap of scl in graphs of groups and 3-manifolds, arXiv:1910.14146.
- [12] M. Clay, M. Forester, and J. Louwsma: Stable commutator length in Baumslag-Solitar groups and quasimorphisms for tree actions, *Trans. Amer. Math. Soc.* **368** (2016), 4751–4785.
- [13] A. J. Duncan and J. Howie: The genus problem for one-relator products of locally indicable groups, *Math. Z.* **208** (1991), 225–237.
- [14] L. Funar and C. Kapoudjian: The braided Ptolemy-Thompson group is finitely presented, *Geom. Topol.* **12** (2008), 475–530.
- [15] L. Funar, C. Kapoudjian, and V. Sergiescu: Asymptotically rigid mapping class groups and Thompson's groups, *Handbook of Teichmüller theory*, IRMA Lect. Math. Theor. Phys. **17** (2012), 595–664.
- [16] L. Funar and V. Sergiescu: Central extensions of the Ptolemy-Thompson group and quantized Teichmüller theory, *J. Topol.* **3** (2010), 29–62.
- [17] É. Ghys and V. Sergiescu: Sur un groupe remarquable de difféomorphismes du cercle, *Comment. Math. Helv.* **62** (1987), 185–239.
- [18] N. Heuer: Gaps in SCL for amalgamated free products and RAAGs, *Geom. Funct. Anal.* **29** (2019), 198–237.
- [19] D. Kotschick: Stable length in stable groups, *Groups of diffeomorphisms*, *Adv. Stud. Pure Math.* **52** (2008), 401–413.
- [20] A. Shtern: Quasisymmetry. I, *Russian J. Math. Phys.* **2** (1994), 353–382.
- [21] T. Susse: Stable commutator length in amalgamated free products, *J. Topol. Anal.* **7** (2015), 693–717.
- [22] A. Walker: Stable commutator length in free products of cyclic groups, *Exp. Math.* **22** (2013), 282–298.
- [23] D. Zhuang: Irrational stable commutator length in finitely presented groups, *J. Mod. Dyn.* **2** (2008), 499–507.

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, JAPAN  
*E-mail address:* m17037h@math.nagoya-u.ac.jp