

Laws of large numbers for weighted sums of independent random variables: a game of mass

Luca Avena¹, Conrado da Costa²

March 18, 2022

Abstract

We consider weighted sums of independent random variables regulated by an increment sequence and provide operative conditions that ensure strong law of large numbers for such sums to hold in both the centred and non-centred case. In the literature, the existing criteria for the strong law are either implicit or based on restrictions on the increment sequence. In our set up we allow for arbitrary sequence of increments, possibly random, provided the random variables regulated by such increments satisfy some mild concentration conditions. In the non-centred case, convergence can be translated into the behavior of a deterministic sequence and it becomes a game of mass provided the expectation of the random variables is a function of the increments. We show how different limiting scenarios can emerge by identifying several classes of increments, for which concrete examples will be offered.

MSC 2020: 60F15, 60G50.

Keywords: Law of large numbers, weighted sums of independent random variables, Toeplitz matrices.

1 Setup, literature and overview

Let $\mathbb{X} := \{X_k, k \in \mathbb{N}\}$ be a sequence of independent real valued random variables with finite mean and $\mathbb{A} = (a_{n,k} \in \mathbb{R}_+; n, k \in \mathbb{N})$ a *Toeplitz summation matrix*, i.e., \mathbb{A} satisfies

$$\lim_n a_{n,k} = 0, \quad (1.1)$$

$$\lim_n \sum_k a_{n,k} = 1, \quad (1.2)$$

$$\sup_n \sum_k |a_{n,k}| < \infty. \quad (1.3)$$

In this set up, one seeks conditions on \mathbb{X} and \mathbb{A} to ensure convergence in probability or almost sure convergence for the sequence $\{S_n, n \in \mathbb{N}\}$, where

$$S_n := \sum_k a_{n,k} X_k.$$

¹Mathematical Institute, Leiden University, P.O. Box 9512, 2300 RA Leiden, The Netherlands

²Mathematical Sciences & Computer Science Building, Upper Mountjoy Campus, Stockton Road, Durham University, DH1 3LE, Durham, UK.

This type of questions, known in the literature as weak/strong Law of Large Numbers (LLN) have been investigated since the birth of probability theory, see [4], and has been extensively studied in the XX century, for different summation methods, e.g. Voronoi sums, Beurling moving averages see [6, 5], see also [13, 10] and references therein for a classical and a more recent account. The quest for operative conditions that apply to a wide range of (\mathbb{X}, \mathbb{A}) and ensure weak/strong convergence of S_n has been the the subject of [9, 11, 14, 12].

When the elements of \mathbb{X} are i.i.d. mean zero random variables, the *weak* LLN is equivalent to $\lim_n \max_k a_{n,k} = 0$, see [11, Theorem 1]. In [11, Theorem 2], the following sufficient conditions for the *strong* LLN are given:

$$\mathbb{E}[X_1^{1+\frac{1}{\gamma}}] < \infty \quad \text{and} \quad \limsup_n n^\gamma \max_k a_{n,k} < \infty, \quad \text{for some } \gamma > 0. \quad (1.4)$$

For (mean-zero) independent but not identically distributed variables, similar sufficient conditions have been examined in [9, 14, 12]. In particular, in analogy with the two conditions in (1.4), these references require that the variables X_k 's are stochastically dominated by a random variable X_* satisfying a moment condition, and that the associated coefficients $a_{n,k}$ decay sufficiently fast.

Unlike the above references, in this paper we impose concentration conditions on \mathbb{X} and obtain sufficient conditions for the weak/strong LLN when $\limsup_n \max_k a_{k,n} > 0$. Here, as in [9], we consider a family of weights, which we will refer to as *masses*, $\mathbf{m} := \{m_k \in \mathbb{R}_+, k \in \mathbb{N}\}$. We assume that the mass sequence \mathbf{m} is such that

$$\sum_{k \in \mathbb{N}} m_k = \infty. \quad (1.5)$$

Set $M_n := \sum_{k=1}^n m_k$ and

$$a_{n,k} := \begin{cases} \frac{m_k}{M_n} & \text{if } k \leq n, \\ 0 & \text{otherwise.} \end{cases} \quad (1.6)$$

Conditions (1.2) and (1.3) hold true by definition. Also, because (1.5) implies $\lim_n M_n = \infty$ it follows that (1.1) is in force and therefore \mathbb{A} is a Toeplitz summation matrix. We notice in particular that if the sum in (1.5) is finite, then no LLN can be expected. In fact, if the random variables are not all constant, the limit random variable will have finite yet strictly positive variance, what precludes convergence to a constant. To describe our results we depart from the set up of [9] and consider $X_k = X_k(m)$ to be a one parameter family of random variables.

Our first contribution: The main goal of this paper is to provide (near) optimal operative conditions on \mathbb{X} to ensure that for *any* sequence of positive masses $\mathbf{m} \in \mathbb{R}_+^{\mathbb{N}}$,

$$S_n = S_n(\mathbf{m}) := \sum_{k=1}^n \frac{m_k}{M_n} X_k(m_k) \quad (1.7)$$

converges to zero as $n \rightarrow \infty$ both in a weak and in a strong sense, going beyond the fast coefficient decay assumptions made in the existing literature. Due to the nature of the coefficients in (1.6) we will refer to the sum in (1.7) as *incremental sum*.

Starting motivation and applications in random media: Our original motivation to look at this type of incremental sums came from the analysis pursued in [3, 1, 2] of the asymptotic speed, and related large deviations, of a Random Walk (RW) in a dynamic random

media, referred to as Cooling Random Environment (CRE). This model is obtained as a perturbation of another process, the well-known RWRE, by adding independence through resetting. RWCRE, denoted by $(\bar{Z}_n)_{n \in \mathbb{N}_0}$, is a patchwork of independent RWRE displacements over different time intervals. More precisely, the classical RWRE consists of a random walk $(Z_n)_{n \in \mathbb{N}_0}$ on \mathbb{Z} with random transition kernel given by a random environment sampled from some law μ at time zero. To build the dynamic transition kernel of RWCRE we fix a partition of time into disjoint intervals $\mathbb{N}_0 = \bigcup_{k \in \mathbb{N}} I_k$ then, we sample a sequence of environments from a given law μ and assign them to the time intervals I_k . To obtain the sum in (1.7) we let $|I_k| = m_k$ and consider $S_n = \bar{Z}_{M_n}/M_n$. In this case, S_n represents the empirical speed of RWCRE at time M_n and $X_k(m_k)$ represents the displacement of an independent RWRE Z_{m_k} after m_k time steps.

This type of time-perturbation of RWRE by resetting in reality gives rise to a slightly more general sum than the incremental one in (1.7). Hence we will prove statements for the above incremental sum but also for the more general one, referred to as *gradual sum*, as defined in (2.1)–(2.2) below. It is worth saying that this patchwork construction can be naturally performed to perturb (in time or even in space) other models in random media, for example, to describe polymagnets [15] based on juxtaposition of independent Curie-Weiss spin-systems [8] of relative sizes m_k 's.

A first partial analysis of the asymptotic speed for RWCRE started in [3, Thm. 1.5] and in [1, Thm. 1.12]. The new results here offer a full characterization of when existence of the asymptotic averages in (1.7) holds and moreover, when moving from a centered zero-mean setup, which values can this limit attain.

Second contribution: Our second goal is thus to explore in the general non-centred case structural conditions on the masses \mathbf{m} that ensure convergence of the weighted sums. We will in particular identify different classes of masses for which the resulting limit exists and can be characterized, what we will refer to as *the game of mass*, for which many concrete examples will be presented.

Structure of the paper: In Section 2 we state the general LLNs for centred random variables: Theorems 2.1 and 2.2, respectively, for the weak and the strong laws for the incremental sum; Theorem 2.3 for the more general gradual sum. Section 3 is devoted to the game of mass where we study concrete convergence criteria for non-centred variables. A discussion on the nature of the hypotheses, illustrated by counterexamples, is presented in Section 4. Lastly, Section 5 contains the proofs of the main theorems. Appendix A covers a technical lemma adapted from [9] and used in the proof of Section 5.2.

2 LLNs for mean-zero variables

In what follows all random variables are defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and \mathbb{E} denotes expectation with respect to \mathbb{P} . Let $\mathbb{X} = \{X_k(m), m \in \mathbb{R}_+, k \in \mathbb{N}\}$ be a family of integrable random variables that are independent in k . Our first statement is the the weak LLN for mean-zero independent random variables.

Theorem 2.1 (Weak LLN). *Assume that \mathbb{X} satisfies the following conditions:*

C (Centering)

$$\forall m \in \mathbb{R}_+, k \in \mathbb{N}; \quad \mathbb{E}[X_k(m)] = 0.$$

W1 (Concentration)

$$\lim_{m \rightarrow \infty} \sup_k \mathbb{P}(|X_k(m)| > \varepsilon) = 0, \quad \forall \varepsilon > 0.$$

W2 (Uniform Integrability)

$$\lim_{A \rightarrow \infty} \sup_{k,m} \mathbb{E}[|X_k(m)| \mathbb{1}_{|X_k(m)| > A}] = 0.$$

Then, for any sequence $\mathbf{m} \in \mathbb{R}_+^{\mathbb{N}}$ that satisfies (1.5),

$$\lim_{n \rightarrow \infty} \mathbb{P}(|S_n| > \varepsilon) = 0, \quad \forall \varepsilon > 0.$$

In this centred case, as captured in the next theorem, to obtain a strong LLN for $\{S_n, n \in \mathbb{N}\}$ we impose further conditions on \mathbb{X} . In particular the concentration condition will be strengthened by requiring a mild polynomial decay and the uniform integrability by a uniform domination.

Theorem 2.2 (Strong LLN). *Assume that \mathbb{X} satisfies (C) and*

S1 (Polynomial decay) *There is a $\delta > 0$ such that for all $\varepsilon > 0$ there is a $C = C(\varepsilon)$ for which*

$$\sup_k \mathbb{P}(|X_k(m)| > \varepsilon) < \frac{C}{m^\delta}.$$

S2 (Stochastic domination and moment control) *There is a random variable X_* and $\gamma > 0$ such that $\mathbb{E}(|X_*|^{2+\gamma}) < \infty$ and for all $x \in \mathbb{R}$*

$$\sup_{k,m} \mathbb{P}(X_k(m) > x) \leq \mathbb{P}(X_* > x).$$

Then for any sequence $\mathbf{m} \in \mathbb{R}_+^{\mathbb{N}}$ that satisfies (1.5),

$$\mathbb{P}\left(\lim_{n \rightarrow \infty} S_n = 0\right) = 1.$$

As anticipated, motivated by the random walk in [3, 1], we next focus on a more general sum by considering a time parameter t that runs on the positive real line partitioned into intervals $I_k = [M_{k-1}, M_k)$ of size m_k : $[0, \infty) = \cup_k I_k$. As $t \rightarrow \infty$ the increments determined by the partition are gradually completed as captured in definition (2.2) below. For $\mathbf{m} \in \mathbb{R}_+^{\mathbb{N}}$, let

$$\ell_t = \ell_t(\mathbf{m}) := \inf\{\ell \in \mathbb{N} : M_\ell \geq t\}, \quad (2.1)$$

and set $\bar{t} := t - M_{\ell_t-1}$. We define the *gradual sum* by

$$\mathcal{S}_t = \mathcal{S}_t(\mathbf{m}) := \sum_{k=1}^{\ell_t-1} \frac{m_k}{t} X_k(m_k) + \frac{\bar{t}}{t} X_{\ell_t}(\bar{t}). \quad (2.2)$$

The next theorem, is the extension of Theorem 2.2 to treat the gradual sum \mathcal{S}_t .

Theorem 2.3 (Generalized strong LLN). *Assume that \mathbb{X} satisfies (C), (S1), (S2), and further:*

S3 (Slow relative increment growth) for every $\varepsilon > 0$ there is a $\beta > 1$, $C_\varepsilon > 0$ which for every $t, r > 0$

$$\sup_k \mathbb{P} \left(\sup_{s \leq m} |(r+s)X_k(r+s) - rX_k(r)| \geq t\varepsilon \right) \leq \frac{C_\varepsilon m^\beta}{t^\beta}.$$

Then for any sequence $\mathbf{m} \in \mathbb{R}_+^{\mathbb{N}}$ that satisfies (1.5)

$$\mathbb{P} \left(\lim_{t \rightarrow \infty} \mathcal{S}_t = 0 \right) = 1.$$

Remark 2.1. (Continuity assumption for the gradual sum) Assumption (S3) controls the oscillations between the times M_n . If the sequence $sX_k(s)$ was a martingale, Doob's L^p inequality would yield (S3). Also, if the increments

$$(r+s)X_k(r+s) - rX_k(r)$$

were bounded by $f(s)$ then this condition would also follow. Condition (S3) reveals that the one parameter families $\{X_k(m), m \in \mathbb{R}\}$ we consider here possess some dependence structure or satisfy some increment domination.

If the random variables \mathbb{X} are not centred, the convergence of \mathcal{S}_t will correspond to the convergence of $\mathbb{E}[\mathcal{S}_t]$. This is the content of the next result.

Corollary 2.1 (Non-centred strong LLN). Assume that \mathbb{X} satisfies (S1)–(S3) and that the sequence $\mathbf{m} \in \mathbb{R}_+^{\mathbb{N}}$ satisfies (1.5). Then, for any increasing sequence $\{t_k\}_{k \in \mathbb{N}}$ with $\lim_k t_k = \infty$ and $\lim_k \mathbb{E}[\mathcal{S}_{t_k}] =: v$,

$$\mathbb{P} \left(\lim_k \mathcal{S}_{t_k} = v \right) = 1.$$

We remark that it is not sufficient to examine the convergence of the sequence $t_k = M_k$, as the boundary term in the gradual sum may not be negligible as, for instance, in the case where

$$\mathbb{E}[X_k(m)] = \frac{2^k - m}{\max\{2^k, m\}}.$$

Indeed, for $m_k = 2^k$ $\mathbb{E}[X_k(m_k)] = 0$ and so for $t_k = M_k$, $\lim_k \mathbb{E}(\mathcal{S}_{t_k}) = 0$ but for $t_k = M_{k-1} + 2^{k-1}$, $\lim_k \mathbb{E}(\mathcal{S}_{t_k}) = 1/4$.

Interestingly, if $\mathbb{E}[X_k(m)] = v_m$ depends only of m , one can relate the convergence of \mathcal{S}_t to the structure of \mathbf{m} . This is what we call *the game of mass* and explore next.

3 The game of mass: operative conditions

As a consequence of Corollary 2.1, it is natural to seek for conditions on (\mathbb{X}, \mathbf{m}) that guarantee convergence of the full sequence \mathcal{S}_t . In this section, we assume that the expectation of $X_k(m)$ depends only on m and not on k , that is:

$$\mathbb{E}[X_k(m)] = v_m \quad \forall k \in \mathbb{N}, \tag{3.1}$$

and that

$$m \mapsto v_m \quad \text{is a bounded continuous function in } \bar{\mathbb{R}}_+, \tag{3.2}$$

where $\bar{\mathbb{R}}_+ = [0, \infty] := \mathbb{R}_+ \cup \{\infty\}$ be the compact metric space with the metric $d(x, y) := \arctan(x) - \arctan(y)$.

We will classify the mass-sequences \mathbf{m} into two classes: *regular* and *non-regular*. The notion of regularity will be captured by the existence of the weak limit of the empirical measure associated to a given mass sequence. We start Section 3.1 with the definition of regular masses and show that, contrary to the non-regular ones, the LLN always holds true. In Section 3.1.1, we consider other notions of regularities and examine how they relate to the convergence of the empirical measures. Section 3.2 is devoted to examples of bounded masses and their relation to the previously defined notions. In Section 3.3, we identify the regular regime of mass sequences that diverge. Finally, in Section 3.4 we investigate what can be said when the mass-sequence \mathbf{m} is *random*. The game of mass is summarized in Fig. 1.




Figure 1: Summary of the game of mass (\mathbf{m}, \mathbb{X}) . The above rectangle offers a visual classification of the possible different masses. The region in gray corresponds to masses for which the LLN is valid, that is, S_t converges. The vertical line divides the masses between regular (left) and irregular (right) ones according to definition 3.1. The horizontal line separates the mass sequences between bounded (down) and unbounded ones for which $\limsup m_k = \infty$ (up). Among the unbounded masses, those divergent in a Cèsaro sense, and in particular those divergent in a classical sense, are always regular. The black and red dotted boxes correspond to those masses for which the related frequencies are asymptotically stable, respectively, in a weak and in a L^1 sense. The roman numbers in each of the different subclasses correspond to the labels of the different illustrative examples from Sections 3.2-3.3-3.4.

3.1 Regular mass sequences

Let $\bar{\mathcal{P}}$ be the space of Borel measures on $\bar{\mathbb{R}}_+$. Recall (2.1), for a given weight sequence $\mathbf{m} \in \mathbb{R}_+^{\mathbb{N}}$, define $\{\mu_t(\cdot) = \mu_t^{(\mathbf{m})}(\cdot), t \geq 0\}$ to be the sequence of *empirical mass measures* on $\bar{\mathbb{R}}_+$, where

$\mu_t(\cdot)$ is given by

$$\mu_t(\cdot) := \frac{\bar{t}}{t} \delta_{\bar{t}}(\cdot) + \sum_{k=1}^{\ell_t-1} \frac{m_k}{t} \delta_{m_k}(\cdot). \quad (3.3)$$

Given a measure $\lambda \in \bar{\mathcal{P}}$ and a measurable function $f : \bar{\mathbb{R}}_+ \rightarrow \mathbb{R}$, we denote by $\langle \lambda, f \rangle = \int f(m) d\lambda(m)$ the integral of f with respect to λ . Furthermore, we say that a sequence $\{\lambda_t; t \geq 0\}$ of probability measures on $\bar{\mathbb{R}}_+$ converges *weakly* to a probability measure λ_* on $\bar{\mathbb{R}}_+$, denoted by $\lambda_t \xrightarrow{w} \lambda_*$ when

$$\lim_t \langle \lambda_t, f \rangle = \langle \lambda_*, f \rangle, \quad \text{for every } f \in C_b(\bar{\mathbb{R}}_+),$$

where $C_b(\bar{\mathbb{R}}_+)$ denotes the space of continuous functions on $\bar{\mathbb{R}}_+$. Note that this definition allows for $\lambda_*(\{\infty\}) := 1 - \lambda_*(\mathbb{R}_+)$ to be strictly positive.

Definition 3.1 (Regular mass sequence). We say that \mathbf{m} is a regular mass sequence when there is a probability measure $\mu_* \in \bar{\mathcal{P}}$ such that

$$\mu_t \xrightarrow{w} \mu_*. \quad (3.4)$$

The following proposition determines the limit of \mathcal{S}_t for regular mass sequences.

Proposition 3.1 (Limit characterization for regular sequences).

Assume \mathbb{X} satisfies (S1)–(S3), and also that it satisfies (3.1) and (3.2). Then, for any mass sequence \mathbf{m} and any $t \geq 0$:

$$\mathbb{E}[\mathcal{S}_t] = \int v_m d\mu_t(m). \quad (3.5)$$

In particular, if \mathbf{m} is regular and (3.4) holds true, then

$$\mathbb{P}\left(\lim_t \mathcal{S}_t = \int v_m d\mu_*(m)\right) = 1. \quad (3.6)$$

Proof. To prove (3.5) note first that by (3.2) we can define $f \in C_b(\bar{\mathbb{R}}_+)$ by $f(\infty) := \lim_m v_\infty$ and $f(m) = v_m$. Now note that

$$\begin{aligned} \mathbb{E}[\mathcal{S}_t] &= \mathbb{E}\left[\frac{\bar{t}}{t} X_{\ell_t}(\bar{t}) + \sum_{k=1}^{\ell_t-1} \frac{m_k}{t} X_k(m_k)\right] \\ &= v_{\bar{t}} \frac{\bar{t}}{t} + \sum_{k=1}^{\ell_t-1} v_{m_k} \frac{m_k}{t} = \int v_m d\mu_t(m) = \langle \mu_t, f \rangle. \end{aligned}$$

As a consequence, if \mathbf{m} is regular there is $\mu_* \in \bar{\mathcal{P}}$ for which (3.4) holds. Therefore, by (3.4) $\langle \mu_t, f \rangle \rightarrow \langle \mu_*, f \rangle$ and (3.6) follows from Corollary 2.1. \square

Remark 3.1. When \mathbf{m} is not regular, almost sure convergence is not prevented, in fact, if $v_m = 0$ for all m , then by Theorem 2.3, \mathcal{S}_t converges almost surely to 0. On the other hand, Examples XI, XIII presented in Section 3.3.1 below show that almost sure convergence may not hold for irregular masses.

3.1.1 Regularity and stability of empirical frequency

There are other notions of regularity beyond the one captured in Definition 3.1. For example, instead of the empirical measure in (3.3), we may examine the *empirical mass frequency* $\{F_t = F_t^{(\mathbf{m})}, t \geq 0\}$, where $F_t \in \bar{\mathcal{P}}$ is given by

$$F_t := \frac{\delta_{\bar{t}}}{\ell_t} + \sum_{k=1}^{\ell_t-1} \frac{\delta_{m_k}}{\ell_t}.$$

The reason to consider other notions of regularity is that it contributes to a finer control of the operative conditions for convergence for L^1 bounded sequences of increments, see also example XIV.

We note that, for any $t \geq 0$ and any arbitrary function f , the following relation between μ_t and F_t is in force:

$$\int f(m) d\mu_t(m) = \frac{\ell_t}{t} \int m f(m) dF_t(m). \quad (3.7)$$

In particular, if we take $f(m) = v_m$ and $f(m) \equiv 1$, respectively, we have that

$$\mathbb{E}[\mathcal{S}_t] = \frac{\ell_t}{t} \int v_m m dF_t(m),$$

and

$$\frac{t}{\ell_t} = \int m dF_t(m). \quad (3.8)$$

The relation in (3.7) may suggest to consider weak convergence of F_t as a natural alternative notion of regularity. However, as shown in the Proposition 3.1 below, these two notions are not equivalent. We find more convenient to adopt the notion in Definition 3.1 for the following two reasons. First, there are masses for which both μ_t and F_t converge weakly to some μ_* and F_* , respectively, but the limit of \mathcal{S}_t is determined by μ_* and not by F_* , see Examples II, IX, and VII below. Second, among the unbounded masses, those divergent in a Cèsaro sense will always be regular while the corresponding F_t is not guaranteed to admit a limit, see Examples VIII and X.

Yet, it is interesting to look at the LLN from the perspective of masses with “well-behaved” frequencies. In particular, the next proposition clarifies how the relation between μ_t and F_t expressed in (3.7) behaves in the limit. In particular it shows how to relate the behavior of the empirical frequencies and empirical masses under different modes of convergence, which we next define.

As is standard, see [7, Thm. 4.6.3, p. 245] we write $F_t \xrightarrow{L^1} F_*$, if there exists a measure $F_*(\cdot)$ on \mathbb{R}_+ for which

$$F_t \xrightarrow{w} F_* \quad \text{and} \quad \int m dF_t(m) \rightarrow \int m dF_*(m) < \infty. \quad (3.9)$$

In a somewhat dual manner, we write $\mu_t \xrightarrow{w^+} \mu_*$, if

$$\mu_t \xrightarrow{w} \mu_* \quad \text{and} \quad \int \frac{1}{m} d\mu_t(m) \rightarrow \int \frac{1}{m} d\mu_*(m) < \infty. \quad (3.10)$$

Proposition 3.2 (Regularity and stable frequencies). *Assume \mathbb{X} satisfies (3.1) and (3.2), consider a mass sequence \mathbf{m} and assume that for $\ell_t = \ell_t(\mathbf{m})$, the limit $A := \lim_{t \rightarrow \infty} \frac{\ell_t}{t} \in \bar{\mathbb{R}}_+$ exists. Then:*

a $F_t \xrightarrow{L^1} F_* \neq \delta_0 \Rightarrow \mu_t \xrightarrow{w} \mu_*$ with $\langle \mu_*, f \rangle := A \int m f(m) dF_*(m)$,

b $\mu_t \xrightarrow{w^+} \mu_* \neq \delta_\infty \Rightarrow F_t \xrightarrow{w} F_*$ with $\langle F_*, f \rangle := \frac{1}{A} \int \frac{1}{m} f(m) d\mu_*(m)$.

Furthermore if both cases above $A \in (0, \infty)$.

Proof. For item **a**, by (3.8), (3.9) and the assumption that $F_t(\cdot)$ converges in L^1 to $F_* \neq \delta_0$, we have that

$$\frac{t}{\ell_t} = \int m dF_t(m) \rightarrow A = \int m dF_*(m) \in (0, \infty).$$

The above relation, (3.7), and (3.9) imply that for any $f \in C_b(\bar{\mathbb{R}}_+)$

$$\mu_t(f) = \frac{\ell_t}{t} \int m f(m) dF_t(m) \rightarrow A \int m f(m) dF_*(m).$$

We now turn to the proof of item **b**. From (3.7), the convergence of $\langle \mu_t, f \rangle$ applied to the function $f(m) = 1/m$ and the assumption that $\mu_* \neq \delta_\infty$ we have that

$$\frac{\ell_t}{t} = \int \frac{1}{m} d\mu_t(m) \rightarrow \int \frac{1}{m} d\mu_*(m) = A \in (0, \infty). \quad (3.11)$$

Therefore, for any $f \in C_b(\bar{\mathbb{R}}_+)$, by (3.7), (3.10), and (3.11) we conclude that

$$\langle F_t, f \rangle = \frac{t}{\ell_t} \int \frac{1}{m} f(m) d\mu_t(m) \rightarrow \frac{1}{A} \int \frac{1}{m} f(m) d\mu_*(m).$$

□

Proposition 3.2 explains part of the different relations depicted in Fig. 1 among the dotted boxes corresponding to masses for which F_t convergences weakly and in L^1 . In what follows, with the help of examples, we explore more how these notions of weak and L^1 convergence for F_t relate to the regularity of μ_t . The examples are organized in the following sections, and in particular they clarify Figure 1. First, in Section 3.2, we examine bounded masses. Second, in Section 3.3.1, we examine Cèsaro divergent masses. Third, in Section 3.3.2, we consider unbounded masses that are not Cèsaro divergent. To conclude, in Section 3.4, we examine i.i.d. random masses.

3.2 Examples of bounded masses

If the sequence of \mathbf{m} is bounded then weak convergence of F_t implies L^1 convergence of F_t . We also remark that when the sequence is regular the limit of \mathcal{S}_t exists and is given by $v = \int v_m d\mu_*(m)$. The following examples show how regular masses relate with weak convergent frequencies.

I (Regular + L^1 -lim F_t). When $\sup_k m_k < \infty$, L^1 convergence follows from weak convergence of the empirical frequency plus uniform integrability. If $F_*(m) \neq \delta_0$, then the formula for the limit of \mathcal{S}_t is given in terms of F_* . Indeed, by item **a** of Proposition 3.2 we conclude that

$$\mathbb{P} \left(\lim_{t \rightarrow \infty} \mathcal{S}_t = v \right) = 1 \quad \text{where} \quad v = A \int v_m m dF_*(m), \quad A := \lim_t \frac{\ell_t}{t}.$$

II (Regular + L^1 -lim F_t). This example shows that if the limit $F_* = \delta_0$ then μ_* may not be given by the expression in item (a) of Proposition 3.2.

Consider the triangular array $\{a_{i,j}, i, j \in \mathbb{N}, j \leq i\}$ defined by $a_{i,1} := 1$ and for $1 < j \leq i$, $a_{i,j} := 2^{-i}$, represented below

$$\begin{aligned} &1, \\ &1, 2^{-1}, \\ &1, 4^{-1}, 4^{-1}, \\ &1, 8^{-1}, 8^{-1}, 8^{-1} \\ &\dots \end{aligned}$$

For the sequence of increment take m_k to be the k -th term of this array, more precisely let $i(k)$ be such that

$$\frac{(i(k)-1)i(k)}{2} \leq k \leq \frac{i(k)(i(k)+1)}{2}$$

and $j(k) := k - \frac{(i(k)-1)i(k)}{2}$. Let $m_k := a_{i(k),j(k)}$. In this example, $F_t \xrightarrow{L^1} \delta_0$ while $\mu_t \xrightarrow{w} \delta_1$. This shows that the L^1 limit of F_t is not sufficient to describe the limit of \mathcal{S}_t , which is given by $v_1 = \int v_m d\mu_*(m)$.

III (Regular + \nexists lim F_t). Let us now move to an example of bounded regular mass sequence such that the limit of F_t does not exist. Consider the sequence \mathbf{m} defined as follows:

- (i) Set $m_1 = 1$,
- (ii) while $F_{M(k)}(\{1\}) > 1/4$ set $m_k = 2^{-k}$ else, go to (iii),
- (iii) while $F_{M(k)}(\{1\}) < 3/4$ set $m_k = 1$ else, go to (ii).

In this case, $\mu_t \xrightarrow{w} \delta_1$ and F_t does not converge.

Note that if \mathbf{m} is not regular, then depending on $\{v_m, m \in \mathbb{R}_+\}$, $\mathbb{E}[\mathcal{S}_t]$ may or may not converge. If there are $K, L \in \mathbb{R}_+$ such that $v_K < v_L$, as in the example below, it is simple to construct a sequence \mathbf{m} for which $\mathbb{E}[\mathcal{S}_t]$ does not converge.

IV (Irregular + \nexists lim F_t). Let \mathbf{m} be the sequence composed of A_i increments of size K followed by B_i increments of size L where the sequences $(A_i)_i, (B_i)_i$ will be determined later. More formally, let $(A_i, B_i)_i$ be given, define $\tau_0 := 0$ $\tau_n := \tau_{n-1} + A_n + B_n$ and set

$$m_k = \begin{cases} K & \text{if } k \in (\tau_n, \tau_n + A_{n+1}] \text{ for some } n \geq 0, \\ L & \text{if } k \in (\tau_n + A_{n+1}, \tau_{n+1}] \text{ for some } n \geq 0. \end{cases} \quad (3.12)$$

Choose $(A_i, B_i; i \in \mathbb{N})$ such that for all $n \in \mathbb{N}$, $A_n < A_{n+1}$, $B_n < B_{n+1}$ and

$$\frac{L(B_1 + \dots + B_n)}{K(A_1 + \dots + A_{n+1})} \leq \frac{1}{n} \quad \text{and} \quad \frac{K(A_1 + \dots + A_{n+1})}{L(B_1 + \dots + B_{n+1})} \leq \frac{1}{n}.$$

If $v_K < v_L$ then $\mathbb{E}[\mathcal{S}_t]$ does not converge as

$$\limsup_t \mathbb{E}[\mathcal{S}_t] = v_L \neq v_K = \liminf_t \mathbb{E}[\mathcal{S}_t].$$

V (Irregular + L^1 -lim F_t). If we combine the sequence defined in Example II with the one defined in Example IV we can construct an irregular sequence for which $F_t \xrightarrow{w} F_*$. More precisely, let m'_k be the sequence defined in Example IV and consider a triangular array $a_{i,j}$ defined by $a_{i,1} := m'_i$ and for $1 < j \leq i$, set $a_{i,j} := 2^{-i}$. To conclude, set $m_k := a_{i(k),j(k)}$ with $i(k), j(k)$ as defined in Example II. Note that this sequence is irregular even though $F_t \xrightarrow{L^1} \delta_0$.

As we look back to item (b) of Proposition 3.2 we see that w^+ convergence can not occur in any of the examples of bounded regular mass for which the empirical frequency does not converge. Indeed, all those example have a significant amount of increments of negligible mass, and as such, they modify the empirical frequency without affecting the limit of the mass sequence. We now move to the study of unbounded masses.

3.3 Unbounded masses

3.3.1 Divergent and Cèsaro's divergent masses

We say that a sequence of masses \mathbf{m} is Cèsaro's divergent when

$$\lim_n \frac{m_1 + \dots + m_n}{n} \rightarrow \infty \quad (3.13)$$

In this case one has that $\mu_t \xrightarrow{w} \delta_\infty$. Therefore the Cèsaro divergent sequences are always regular and by Proposition 3.1 it follows that

$$\mathbb{P}(\lim_t \mathcal{S}_t = v_\infty) = 1. \quad (3.14)$$

A particular case of Cèsaro divergence is given by the divergent masses as captured in the next example.

VI (Divergent mass + w -lim F_t). We say that a sequence of masses \mathbf{m} is divergent when

$$\lim_{k \rightarrow \infty} m_k = \infty, \quad (3.15)$$

in which case (3.13) holds true and hence (3.14). Also, as $\lim_t F_t([0, A]) = 0$ for any $A > 0$, it follows that $F_t \xrightarrow{w} \delta_\infty$.

The regime captured in (3.15) is treated in [1, Theorem 1.10] in the context of random walks in dynamic environment. Theorem 2.3 can actually be seen as a generalization of the latter. As briefly mentioned at the end of Section 4, the present proofs could actually cover even more general cases, if, for example, we relax the assumption in Equation (3.1). The following example shows that in the Cèsaro divergent regime, the sequence F_t may converge, but may not be able to capture the limit of \mathcal{S}_t .

VII (Cèsaro divergent mass + w -lim F_t). Consider the sequence \mathbf{m}

$$m_k := \begin{cases} 1 & \text{if } k \text{ is odd, and} \\ k & \text{if } k \text{ is even.} \end{cases}$$

Informally, half the increments are 1, and the other half diverges. More precisely,

$$F_t \xrightarrow{w} \frac{1}{2}\delta_1 + \frac{1}{2}\delta_\infty.$$

As such, one might be tempted to say that $\mathbb{E}[\mathcal{S}_t] \rightarrow \frac{1}{2}v_1 + \frac{1}{2}v_\infty$ as $t \rightarrow \infty$. This is not the case because one has to take into account the relative weights of the sequences. As it turns out, the mass of increments of size 1 for this particular sequence vanishes in the limit. Indeed, note that the sum of the first $2k$ increments, M_{2k} is

$$M_{2k} = \frac{k(k-1)}{2} + k = \frac{k^2 + k}{2}.$$

Now note that $\frac{k}{M_{2k}} \rightarrow 0$ and therefore

$$\mathbb{E}[\mathcal{S}_{M_{2k}}] = \frac{k}{M_{2k}}v_1 + \frac{1}{M_{2k}} \sum_{i=1}^k iv_i \rightarrow v_\infty.$$

Also in this example, if $v_1 \neq v_\infty$, then the weak limit of F_t does not determine the limit of \mathcal{S}_t , even if it is well defined.

As in the bounded case, see Example III, also Cèsaro divergent sequences may not have well behaved empirical frequencies, as shown in the next example.

VIII (Cèsaro divergent mass + $\not\# w$ -lim F_t). It is possible to construct a sequence \mathbf{m} that is regular but such that F_t does not converge weakly. Take an irregular sequence \mathbf{m}' such as the one defined in (3.12) and intercalate it with a huge increment so that it diverges in the Cèsaro sense. To be more concrete, for $k \in \mathbb{N}$ let $m_{2k-1} := m'_k$ and $m_{2k} := k \sum_{i=1}^{2k-1} m_i$.

3.3.2 Unbounded sequences that do not diverge in the Cèsaro sense

When dealing with unbounded masses that are not Cèsaro divergent, then the sequence is not necessarily regular and more subtle scenarios may occur, as the following examples illustrate. We start with an example of a regular sequence that allows an asymptotic positive mass of increments of finite size and positive mass at infinity.

IX (Regular $\liminf m_k < \infty + w$ -lim F_t). Let the first elements of $\mathbf{m} \in \mathbb{R}_+^{\mathbb{N}}$ be defined by $m_1 := 1, m_2 := 2, m_3 := 1$. If $m_k := j > 1$ then the next $j-1$ increments will be of size 1 after that $m_{k+j} := j+1$. The sequence of increment sizes can be arranged in a triangular array $\{a_{i,j}\}_{i,j \geq 1}$, where $m_k := a_{i(k),j(k)}$ with $i(k), j(k)$ as in Example II.

$$\begin{array}{c} 1, \\ 2, 1, \\ 3, 1, 1, \\ \dots \end{array}$$

In this case $F_t \xrightarrow{w} \delta_1$ but $\mu_t \xrightarrow{w} \frac{1}{2}\delta_1 + \frac{1}{2}\delta_\infty$ and so $\lim_t \mathbb{E}[\mathcal{S}_t] = \frac{1}{2}v_1 + \frac{1}{2}v_\infty$. We notice in particular that if $v_1 > v_\infty$, the above mass sequence is another example of a regular sequence for which the weak limit of F_t does not determine the limit of \mathcal{S}_t , even when it exists.

The next example shows a regular sequence with unbounded increments and for which the empirical frequency does not converge.

X (Regular + $\not\rightarrow$ w -lim F_t). Take \mathbf{m} as in Example III but replace the k -th increment of mass 1 by the k -th increment of the sequence defined in Example IX. For this example, we have that

$$\limsup_t F_t(\{1\}) - \liminf_t F_t(\{1\}) \geq 1/2.$$

The above condition precludes the weak convergence of F_t . Furthermore, since the total mass on increments strictly smaller than 1 is finite, $\mu_t \xrightarrow{w} \frac{1}{2}\delta_1 + \frac{1}{2}\delta_\infty$ and the sequence is regular.

XI (Irregular + w -lim F_t). Only weak convergence of the empirical measure F_t does not imply convergence of \mathcal{S}_t . Indeed, assume that $v_1 > v_\infty$, and let $(K_i)_i$ and $(N_i)_i$ be auxiliary sequences that we will determine later. The sequence \mathbf{m} alternates one increment of size K_i with N_i increments of size 1. More precisely, let $\tau(j) = j + \sum_{i=1}^j N_i$ and set

$$m_k = \sum_{j=1}^{\infty} K_j \mathbb{1}_{\tau(j)} + \mathbb{1}_{(\tau(j), \tau(j+1))}.$$

Now, choose (N_i, K_i) such that

$$\frac{N_1 + \dots + N_i}{K_i} \leq \frac{1}{i} \text{ and } \frac{K_1 + \dots + K_i}{N_{i+1}} \leq \frac{1}{i}.$$

Note that $F_t \xrightarrow{w} \delta_1$, but if $v_1 > v_\infty$

$$\limsup_t \mathbb{E}[\mathcal{S}_t] = v_1 > v_\infty = \liminf_t \mathbb{E}[\mathcal{S}_t].$$

XII (Irregular + L^1 -lim F_t). In this example we construct an unbounded irregular sequence for which F_t converges in L^1 . In particular from item (a) of Proposition 3.2 it follows that this limit must be δ_0 . Let $(A_i)_i$ be auxiliary sequences to be defined later. Informally, this constructed as a combination of Example V and Example XI, where we intercalate an irregular unbounded sequence with a large number of increments of small mass. Let \mathbf{m}' be the sequence defined in Example XI set $\tau(1) := 1$ and for $j > 1$ set $\tau(j) := \tau(j-1) + A_j$. Now let

$$m_k := \sum_j m'_j \mathbb{1}_{\tau(j)} + 2^{-k} \mathbb{1}_{(\tau(j-1), \tau(j))}.$$

Finally choose A_i such that

$$\frac{\sum_{k=1}^i m'_k + 1}{A_i} < \frac{1}{i}.$$

Since

$$\int m dF_t(m) \leq \frac{\sum_{k=1}^{\ell_t} m'_k + 1}{\ell_t} \rightarrow 0, \text{ as } t \rightarrow \infty,$$

it follows that $F_t \xrightarrow{L^1} \delta_0$. Furthermore, if $v_1 > v_\infty$

$$\limsup_t \mathbb{E}[\mathcal{S}_t] = v_1 > v_\infty = \liminf_t \mathbb{E}[\mathcal{S}_t].$$

XIII (irregular + $\not\rightarrow$ w -lim F_t). It is also possible to construct a sequence \mathbf{m} that is irregular but such that F_t does not converge weakly. Take the sequence defined in Example III and replace the k -th increment of size 1 by the k -th increment of the sequence defined in XI.

3.4 Random masses

In this section consider random mass sequences \mathbf{m} . More specifically, we let m_k be an i.i.d. sequence of random variables, independent of \mathbb{X} , each distributed according to a measure ν on \mathbb{R}_+ . There are two cases depending on whether ν has finite or infinite mean. For notational ease, we model $(m_k, k \in \mathbb{N})$ as i.i.d. random variables in the probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

XIV (Regular + (un) bounded + L^1 -lim F_t). Assume that $\nu \neq 0$, and that $\int m\nu(dm) < \infty$. Now, let the increments m_k to be sampled independently from ν . By the Glivenko-Cantelli Theorem [7, Theorem 2.4.9] it follows that almost surely $F_t([0, x])$ converges (uniformly in x) to $\nu([0, x])$. By the classical LLN for iid. random variables, almost surely, $\int m dF_t(m) \rightarrow \int m dF_*(m)$. Therefore the conditions of (3.9) are satisfied almost surely and so $\mathbb{P}(F_t \xrightarrow{L^1} F_*) = 1$. By item (a) of Proposition 3.2 it follows that $\mathbb{P}(\mu_t \xrightarrow{w} \nu) = 1$. Therefore, almost surely, the sequence m_k is regular and

$$\mathbb{P}\left(\lim_t \mathbb{E}[\mathcal{S}_t] = \int v_x d\nu(x)\right) = 1.$$

XV (Regular + Cèsaro + $\exists w$ -lim F_t). Now, assume that $\int m d\nu(m) = \infty$ and define the increments m_k to be sampled independently from ν . In this case

$$\mathbb{P}\left(\frac{m_1 + \dots + m_k}{k} \rightarrow \infty\right) = 1. \quad (3.16)$$

Then note that after k increments, the mass of increments of size smaller than $a > 0$, $\mu_t([0, a])$, is bounded by $\frac{ka}{m_1 + \dots + m_k}$ and therefore, by (3.16), for any $a > 0$, almost surely $\mu_t([0, a]) \rightarrow 0$. This implies that $\mathbb{P}(\mu_t \xrightarrow{w} \delta_\infty) = 1$ and therefore

$$\mathbb{P}\left(\lim_t \mathcal{S}_t = v_\infty\right) = 1.$$

4 Hypothesis and counterexamples

4.1 Weak LLN: necessity of (W1) and (W2).

Both conditions (W1) (W2) are necessary for the weak LLN. The necessity for condition (W1) is due to [9, Theorem 1]. We show below that condition (W2) is necessary by means of a counter-example.

Counter-example: Consider a sequence $\{U_k, k \in \mathbb{N}\}$ of i.i.d. uniform random variables on $(0, 1)$ and $X_k(m) := V_m(U_k)$, where

$$V_m(u) = \begin{cases} A_m & \text{if } u \in [0, g(m)/2), \\ -A_m & \text{if } u \in (g(m)/2, g(m)], \\ 0 & \text{else.} \end{cases}$$

with this definition, it follows that $\mathbb{P}(|X_k(m_k)| > 0) = g(m_k)$. Assume that g is a strictly decreasing continuous function such that $\lim_{m \rightarrow \infty} g(m) = 0$. Let $m_k := \inf\{m : g(m) > 1/k\}$. This implies that $m_k \rightarrow \infty$ as $k \rightarrow \infty$ and so (1.5) is satisfied. Furthermore by the

definition of $X_k(m_k)$, the assumptions (C) and (W1) in Theorem 2.1 are verified. Now choose $\{A_{m_k}, k \in \mathbb{N}\}$ to be such that

$$\frac{m_n}{M_{N(n)}} A_{m_n} > 1 + \sum_{k=1}^{n-1} A_{m_k},$$

where $N(n)$ is such that

$$\mathbb{P}(\exists n \leq j \leq N(n) : X_j(m_j) \neq 0) > \frac{1}{2}.$$

Such an $N(n)$ exists and is finite since by the second Borel-Cantelli Lemma and the continuity of probability measures:

$$1 = \mathbb{P}(\exists j \geq n : X_j(m_j) \neq 0) = \lim_{N \rightarrow \infty} \mathbb{P}(\exists n \leq j < N : X_j(m_j) \neq 0).$$

With this choice of A_{m_n} it follows that if there is a j , $i \leq j \leq N(i)$ for which $|X_j(m_j)| > 0$ then $|S_{N(i)}| > 1$. Therefore for any $i \in \mathbb{N}$,

$$\mathbb{P}(|S_{N(i)}| > 1) > \frac{1}{2}.$$

As $\mathbb{P}(S_n > 0 \mid |S_n| > 0) = \frac{1}{2}$, we conclude that the weak LLN does not hold.

4.2 Strong LLN: near optimality of (S1).

One could try to improve the condition in (S1) by requiring a decay smaller than polynomial, that is:

$$\mathbb{P}(|X_k(m)| > \varepsilon) < \frac{C_\varepsilon}{f(m)},$$

for some $f : \mathbb{R}_+ \rightarrow \mathbb{R}_+$. When we look for a scale that grows slower than any polynomial, $f(m) = \log(m)$ is a natural candidate. However, as illustrated next, this already allows for counterexamples.

Counter-example: Let $\{U_k, k \in \mathbb{N}\}$ be a sequence of i.i.d. uniform random variables on $(0, 1)$ and let $X_k(m) := g_m(U_k)$ where

$$g_m(x) := \begin{cases} 1 & \text{if } x \in (0, \frac{1}{2 \log_2 m}), \\ -1 & \text{if } x \in [\frac{1}{2 \log_2 m}, \frac{1}{\log_2 m}), \\ 0 & \text{else.} \end{cases}$$

Note that \mathbb{X} fulfills assumptions (C), (S2), (S3), and instead of (S1) it satisfies

$$\mathbb{P}(|X_k(m)| > \varepsilon) = \frac{1}{\log_2 m}.$$

Now take \mathbf{m} with $m_k = 4^k$. For such an \mathbf{m} we see that the incremental sum S_n does not satisfy the strong LLN. Indeed, as

$$\mathbb{P}(|X_k(m_k)| = 1) = \frac{1}{2k},$$

by the second Borel Cantelli lemma,

$$\mathbb{P}(|X_k(m_k)| = 1, \text{ i.o.}) = 1.$$

Therefore, by (1.7) it follows that there is an $\varepsilon > 0$ for which

$$\mathbb{P}(|S_n - S_{n-1}| > \varepsilon \text{ i.o.}) = 1,$$

which means that almost surely S_n does not converge.

In light of the above example, we see that the condition S1 is near to optimal. Indeed, to improve it, we would need to find $f(m)$ satisfying

$$\log^k(m) \ll f(m) \ll m^\delta \quad \forall k \in \mathbb{N}, \delta > 0.$$

4.3 Concluding remarks

- **Independence.** Our examples above and proofs below are based on the independence in k of $\{X_k(m), m \in \mathbb{R}_+, k \in \mathbb{N}\}$. However, for certain choices of well-behaved mass sequences \mathbf{m} , it seems possible to adapt our arguments and still obtain a strong LLN in presence of “weak enough dependence”, though the notion of “weak enough dependence” would very much depend on the weight sequence and this is why we did not pursue this line of investigation.
- **Relaxing condition (3.1).** In the game of mass described in Section 3, for simplicity, we have restricted our analysis to variables with expected value independent of k , as captured in assumption (3.1). We note that this is not really needed, as we might, for example, consider $X_k(m)$ ’s with expected value, say, v_m and $v'_m \neq v_m$ depending on the parity of k . Yet, the resulting analysis would branch into many different regimes depending on how exactly condition (3.1) is violated.
- **Fluctuations and large deviations.** It is natural to consider “higher order asymptotics”, such as large deviations or scaling limit characterizations, for the sums in (1.7) or (2.2). However, the analysis for this type of questions relies heavily on the specific distribution of the sequence of variables \mathbb{X} thus preventing a general self-contained treatment. Still, it is interesting to note that these other questions can give rise to many subtleties and anomalous behavior. This is well illustrated by the specific RWCRE model in random media introduced in [3] that motivated the present paper, we refer the interested reader to [2] for results on crossovers phenomena in related fluctuations, and to [1] for stability results of large deviations rate functions.

5 Proofs

5.1 Weak law of large numbers

In this section we prove Theorem 2.1 by implementing a truncation argument following the ideas from [9]. For each $K > 0$, let S_n^K represent the contribution to S_n coming from the increments larger than K , i.e.

$$S_n^K := \sum_{k=1}^n \frac{m_k}{M_n} X_k(m_k) \mathbb{1}_{m_k > K}.$$

Now note that due to (W1) and (W2) it follows that

$$\lim_{K \rightarrow \infty} \sup_{m > K} \sup_k \mathbb{E}[|X_k(m)|] = 0. \quad (5.1)$$

Indeed, for any $\varepsilon > 0$ and any $A > \varepsilon$

$$\mathbb{E}[|X_k(m)|] \leq \varepsilon + A\mathbb{P}(\varepsilon < |X_k(m)| \leq A) + \mathbb{E}[|X_k(m)| \mathbb{1}_{|X_k(m)| > A}].$$

the right hand side above can be bounded by 3ε using (W1) and (W2) and since $\varepsilon > 0$ is arbitrary, (5.1) follows. Now let $\bar{S}_n^K := S_n - S_n^K$ be the contribution to S_n coming from the increments smaller than K . By the triangle inequality and the union bound it follows that

$$\mathbb{P}(|S_n| > \varepsilon) \leq \mathbb{P}(|S_n^K| + |\bar{S}_n^K| > \varepsilon) \leq \mathbb{P}\left(|S_n^K| > \frac{\varepsilon}{2}\right) + \mathbb{P}\left(|\bar{S}_n^K| > \frac{\varepsilon}{2}\right).$$

As $\sum_{k=1}^n \frac{m_k}{M_n} \mathbb{1}_{m_k > K} \leq 1$, (5.1) and Markov's inequality imply

$$\limsup_{K \rightarrow \infty} \mathbb{P}(S_n^K > \varepsilon) = 0,$$

and therefore for any $K > 0$,

$$\limsup_{n \rightarrow \infty} \mathbb{P}(|S_n| > \varepsilon) \leq \limsup_{n \rightarrow \infty} \mathbb{P}\left(|\bar{S}_n^K| > \frac{\varepsilon}{2}\right).$$

It remains to prove that the right-hand side above goes to zero for arbitrary $\varepsilon > 0$. To do so, consider the truncated random variables

$$Y_k(m_k) := X_k(m_k) \mathbb{1}_{|X_k(m_k)| < \frac{M_n}{m_k}} \mathbb{1}_{m_k < K}$$

and notice that as $M_n \rightarrow \infty$,

$$\lim_n \max_{1 \leq k \leq n} \frac{m_k}{M_n} \mathbb{1}_{m_k < K} = 0. \quad (5.2)$$

Set $\bar{s}_n^K := \sum_{k=1}^n Y_n(m_k)$. We will first argue that this truncated sum \bar{s}_n^K approximates well \bar{S}_n^K , and then show that the variance of truncation vanishes. To perform these two steps we will need the following lemma, whose proof is postponed to the end of this section.

Lemma 5.1. *If (C), (W1), and (W2) of Theorem 2.1 hold true, then*

$$\lim_{n \rightarrow \infty} \max_{1 \leq k \leq n} \frac{M_n}{m_k} \mathbb{P}\left(|X_k(m)| \mathbb{1}_{m_k < K} \geq \frac{M_n}{m_k}\right) = 0, \quad (5.3)$$

and

$$\lim_{n \rightarrow \infty} \max_{1 \leq k \leq n} \frac{m_k}{M_n} \mathbb{E}[Y_k^2(m_k)] = 0. \quad (5.4)$$

By the union bound, the definition of $Y_k(m_k)$, using that $\sum_k \frac{m_k}{M_n} \leq 1$ we have that

$$\begin{aligned} \limsup_n \mathbb{P}(\bar{S}_n^K \neq \bar{s}_n^K) &\leq \limsup_n \sum_{k=1}^n \mathbb{P}(X_k(m_k) \mathbb{1}_{m_k < K} \neq Y_k(m_k)) \\ &= \limsup_n \sum_{k=1}^n \mathbb{P}\left(|X_k(m)| \mathbb{1}_{m_k < K} \geq \frac{M_n}{m_k}\right) \\ &\leq \limsup_n \max_{1 \leq k \leq n} \frac{M_n}{m_k} \mathbb{P}\left(|X_k(m_k)| \mathbb{1}_{m_k < K} \geq \frac{M_n}{m_k}\right) \sum_{k=1}^n \frac{m_k}{M_n} \\ &\leq \limsup_n \max_{1 \leq k \leq n} \frac{M_n}{m_k} \mathbb{P}\left(|X_k(m_k)| \mathbb{1}_{m_k < K} \geq \frac{M_n}{m_k}\right), \end{aligned}$$

the latter can be made arbitrary small via (5.3). Hence it suffices to consider \bar{s}_n^K instead of \bar{S}_n^K . We next control the mean and the variance of \bar{s}_n^K .

The mean. As $X_k(m_k)$ is uniformly integrable family of centred random variables, by (5.2) it follows that $\limsup_n \sup_k \mathbb{E}[|Y_k(m_k)|] = 0$, from which it follows that $\lim_n \mathbb{E}(\bar{s}_n^K) = 0$.

The Variance. Similarly, by independence and (5.4) we can estimate

$$\begin{aligned} \limsup_n \text{Var}(\bar{s}_n^K) &= \limsup_n \sum_{k=1}^n \frac{m_k^2}{M_n^2} \text{Var}(Y_k(m_k)) \\ &\leq \limsup_n \sum_{k=1}^n \frac{m_k}{M_n} \max_{1 \leq k \leq n} \frac{m_k}{M_n} \text{Var}(Y_k(m_k)) \\ &\leq \limsup_n \max_{1 \leq k \leq n} \frac{m_k}{M_n} \mathbb{E}[Y_k^2(m_k)] = 0. \end{aligned} \quad (5.5)$$

Finally, $\lim_n \mathbb{E}(\bar{s}_n^K) = 0$ together with (5.5) and Chebyshev's inequality yield

$$\limsup_n \mathbb{P}(\bar{s}_n^K \geq \varepsilon) \leq \limsup_n \frac{4}{\varepsilon^2} \text{Var}(\bar{s}_n^K) = 0.$$

□

Proof of Lemma 5.1 Let $\bar{T}_n := \inf_{1 \leq k \leq n} \frac{M_n}{m_k} \mathbb{1}_{m_k < K}$. Since $\lim_n \bar{T}_n = \infty$, equation (5.3) follows from (W2) as

$$\lim_n \frac{M_n}{m_k} P\left(|X_k(m)| \geq \frac{M_n}{m_k}\right) \leq \lim_n \mathbb{E}\left[|X_k(m)| \mathbb{1}_{|X_k(m)| \geq \bar{T}_n}\right] = 0.$$

To prove (5.4), let $F_{k,m}(a) := \mathbb{P}(|X_k(m)| < a)$ and note first that integration by parts yields

$$\begin{aligned} \int_0^T x^2 dF_{k,m}(x) &= T^2 \mathbb{P}(|X_k(m)| < T) - 2 \int_0^T x \mathbb{P}(|X_k(m)| < x) dx \\ &= T^2 [1 - \mathbb{P}(|X_k(m)| \geq T)] - 2 \int_0^T x [1 - \mathbb{P}(|X_k(m)| \geq x)] dx \\ &= -T^2 \mathbb{P}(|X_k(m)| \geq T) + 2 \int_0^T x \mathbb{P}(|X_k(m)| \geq x) dx. \end{aligned} \quad (5.6)$$

Observe further that

$$\int_0^{\frac{M_n}{m_k}} x^2 dF_{k,m}(x) = \mathbb{E}[Y_k^2(m_k)],$$

and that by the uniform integrability (W2)

$$\lim_{T \rightarrow \infty} \sup_{k,m} T \mathbb{P}(|X_k(m)| \geq T) = 0. \quad (5.7)$$

By (5.7) and (5.6), it follows that

$$\lim_{n \rightarrow \infty} \max_{1 \leq k \leq n} \frac{m_k}{M_n} \mathbb{E}[Y_k^2(m_k)] \leq \lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \sup_{k \leq n, m} \frac{1}{T} \int_0^T x^2 dF_{k,m}(x) = 0.$$

□

5.2 Strong law of large numbers for the incremental sum

As in [1], our proof here relies on an iterative scale decomposition into “small” and “big” increments. At each scale, the small contribution is defined as the truncated sum that, thanks to the stochastic domination assumption (S2), can be dealt with the techniques of [12]. What is left, classified as “big”, is again split (in the next scale) into a “small” and a “big”. At this level, the small one is controlled in the same way as before. The iteration proceeds until we reach a scale where the condition (S1) is sufficient to ensure convergence. The proof is organized as follows. We first iteratively decompose the sum S_n into a finite number of sums of relatively small increments and one sum of large increments. Next we show that the large increment sum converges to zero almost surely. Finally we prove that each of the small increments also converge to zero almost surely.

The recursive decomposition. We take δ from (S1) and γ from (S2) and fix $K = K(\delta, \gamma) \in \mathbb{N}$ such that

$$\delta K > 1, \quad \text{and} \quad \frac{K}{K-1} < 1 + \gamma$$

Now, we let $M^0 := \mathbb{N}$, and let

$$M^{0,s} := \{j \in M^0 : m_j \leq 1\} \tag{5.8}$$

and $M^1 := M^0 \setminus M^{0,s}$. For $i \geq 1$, given M^i , if M^i is finite for $j > i$ set $M^j, M^{j,s} = \emptyset$, if M^i is an infinite set, let $k^i : \mathbb{N} \rightarrow M^i$ be an increasing map with $k^i(\mathbb{N}) = M^i$. define the (i)-st small increments by

$$M^{i,s} := \{k_j^i \in \mathbb{N} : m_{k_j^i} < j^{i/K}\},$$

and define the next level (large) increments $M^{k+1} := M^i \setminus M^{i,s}$. Now, we let the cardinality of indices less than n be denoted by

$$J(i; n) := \#\{j \in M^i : j \leq n\} \quad \text{and} \quad J(i, s; n) := \#\{j \in M^{i,s} : j \leq n\}.$$

To ease the notation set $X_k := X_k(m_k)$, $a_{j,n}^i := \frac{m_{k_j^i}}{M_n}$, and $a_{j,n}^{i,s} = \frac{m_{k_j^{i,s}}}{M_n}$. Since $\mathbb{N} = \bigcup_{i=0}^K M^{i,s} \cup M^K$, we have that

$$\begin{aligned} S_n &= \sum_{i=1}^K \underbrace{\sum_{j=1}^{J(i,s;n)} a_{j,n}^{i,s} X_{k_j^{i,s}}}_{S_n^{i,s}} + \underbrace{\sum_{j=1}^{J(K;n)} a_{j,n}^K X_{k_j^K}}_{S_n^K} \\ &= \sum_{i=1}^K S_n^{i,s} + S_n^K. \end{aligned}$$

In what follows we show that

$$\mathbb{P}\left(\limsup_n S_n^K = 0\right) = 1, \tag{5.9}$$

$$\mathbb{P}\left(\limsup_n S_n^{i,s} = 0\right) = 1 \text{ for } i \in \{0, 1, \dots, K\}. \tag{5.10}$$

The large increments sum. To prove (5.9) it is enough to show that for any $\varepsilon > 0$

$$\mathbb{P}(\limsup_n S_n^K \leq \varepsilon) = 1. \quad (5.11)$$

By (S1), and the fact that $m_{k_j^K} \geq j^K$, it follows that

$$\mathbb{P}(X_{k_j^K} > \varepsilon) \leq \frac{C_\varepsilon}{(m_{k_j^K})^\delta} \leq \frac{C_\varepsilon}{j^{K\delta}}.$$

Since $K\delta > 1$, we conclude that $\sum_{j=1}^\infty \mathbb{P}(X_{k_j^K} > \varepsilon) < \infty$, and by the Borel Cantelli Lemma, we have that

$$\mathbb{P}\left(\limsup_j |X_{k_j^K}| \leq \varepsilon\right) = 1.$$

As $M_n \rightarrow \infty$ and $\sum_{j=1}^{J(K,n)} m_{k_j^K} \leq M_n$, we conclude that (5.11) holds.

The small increment sums. The proof of (5.10) will be split in two parts, first we prove it for $i \geq 1$ and then we treat the case $i = 0$. For notation ease, set for any $J \in \mathbb{N}$,

$$\tilde{m}_j := m_{k_j^{i,s}}, \quad \tilde{M}_J := \sum_{j=1}^J \tilde{m}_j, \quad \tilde{a}_{j,J} := \frac{\tilde{m}_j}{\tilde{M}_J}, \quad \text{and let } \tilde{S}_J = \sum_{j=1}^J \tilde{a}_{j,J} X_{k_j^{i,s}}.$$

Now note that for any n

$$S_n^{i,s} = \frac{\tilde{M}_{J(i,s;n)}}{M_n} \tilde{S}_{J(i,s;n)}.$$

As $\frac{\tilde{M}_{J(i,s;n)}}{M_n} \leq 1$, it follows that $\limsup_n |S_n^{i,s}| \leq \limsup_J |\tilde{S}_J|$. Therefore, it suffices to show that

$$\mathbb{P}\left(\lim_J \tilde{S}_J = 0\right) = 1. \quad (5.12)$$

Since for $i \geq 1$, $m_{k_j^{i,s}} \in [j^{(i-1)/K}, j^{i/K}]$, there are $C, c > 0$ for which

$$\tilde{M}_J \geq cJ^{1+(i-1)/K}, \quad \tilde{a}_{j,J} \leq \frac{C}{J^{\frac{K-1}{K}}}. \quad (5.13)$$

Now, as $\lim_J \tilde{a}_{j,J} = 0$, $\sum_j \tilde{a}_{j,J} = 1$, and conditions (5.13) and (S2) hold, one can apply Theorem 2 in [12] with $\nu = \frac{K-1}{K}$ to obtain (5.12) and therefore (5.10) for $i \geq 1$. To conclude the proof of Theorem 2.2 it remains to verify that $S_n^{0,s}$ converges to 0 almost surely. The proof is an adaptation of Theorem 4 in [9] and is postponed to Appendix A. \square

5.3 Strong law for the gradual sum

In this section we prove Theorem 2.3. Recall the decomposition of \mathcal{S}_t from (2.2). We note that \mathcal{S}_t is a convex combination of S_{ℓ_t} and the boundary term $X_{\ell_t}(t)$ with $\bar{t} = t - M_{\ell_t}$.

By the proof of (2.2), To prove Theorem 2.3, it remains to show that the boundary term vanishes, i.e.

$$\mathbb{P} \left(\lim_{t \rightarrow \bar{t}} \frac{\bar{t}}{t} X_{\ell_t}(\bar{t}) = 0 \right) = 1. \quad (5.14)$$

We divide the proof of (5.14) in two steps. First we show (5.14) for a properly defined small increments, $m_{k+1} < (1 + \alpha_k)M_k$. Then we show (5.14) for the complement set that we refer to as the set of large increments.

5.3.1 The small Increments

Let $V_n = \sup \left\{ \frac{s}{(M_n+s)} |X_k(s)| : s \in [0, m_k] \right\}$ and note that

$$\limsup_t \frac{\bar{t}}{t} X_{\ell_t}(\bar{t}) = \limsup_n V_n. \quad (5.15)$$

Thanks to condition (S3), we can control the oscillations V_n for small increments that satisfy a growth condition defined as follows. Fix $\beta > 1$ as in (S3) and let $\alpha_j = \frac{1}{j^a}$ with $a \in (1/\beta, 1)$. The first small increment is defined by

$$k'_1 := \inf \{ k \in \mathbb{N} : M_{k+1} < (1 + \alpha_1)M_k \},$$

and define recursively for j -th small increment by

$$k'_{j+1} := \inf \{ k \in \mathbb{N} : k > k'_j, \quad M_{k+1} < (1 + \alpha_{j+1})M_k \}.$$

If for some $j, k_j = \infty$ this implies there are only finitely many small increments and we do not need to worry about them in (5.15). If for all $j, k'_j < \infty$, we claim that almost surely

$$\limsup_j V_{k'_j} = 0.$$

Indeed, as $m_{k'_j+1} < \alpha_j M_{k'_j}$, by (S3), with $r = 0$, it follows that for any $\varepsilon > 0$

$$\mathbb{P}(V_{k'_j} > \varepsilon) \leq \mathbb{P} \left[\sup_{s \leq \alpha_j M_{k'_j}} s |X_{k'_j}(s)| > \varepsilon M_{k'_j} \right] \leq \alpha_j^\beta C_\varepsilon.$$

As $\sum_j \alpha_j^\beta < \infty$, by the Borel-Cantelli lemma we conclude that

$$\mathbb{P} \left(\limsup_j V_{k'_j} \leq \varepsilon \right) = 1, \quad (5.16)$$

and since $\varepsilon > 0$ is arbitrary, it follows that

$$\mathbb{P} \left(\lim_j V_{k'_j} = 0 \right) = 1. \quad (5.17)$$

□

5.3.2 The large increments

By (5.17) we can restrict our attention to $\{k_1^*, k_2^*, \dots\} = \mathbb{N} \setminus \{k'_1, k'_2, \dots\}$. Note that since $\alpha_j \leq 1$

$$(1 + \alpha_j) \geq C \exp(\alpha_j/2), \quad (5.18)$$

for some $C > 0$. Therefore, for some $c_a > 0$ the following growth condition holds for the terms in the sequence $\{M_{k_i^*}\}_i$

$$M_{k_i^*} \geq \prod_{j=1}^i (1 + \alpha_j) M_1 \geq C \exp\left(\sum_{j=1}^i \frac{\alpha_j}{2}\right) M_1 \geq \exp(c_a i^{1-a}) M_1. \quad (5.19)$$

The proof now proceeds in two steps, we first show that the boundary term $\frac{\bar{t}}{t} X_{\ell_t}(\bar{t})$ converges to zero along a subsequence $\{t_{i,j}, i, j \in \mathbb{N} \cup \{0\}\}$, what we call *pinning*, and then based on this result we show that the full sequence converges to zero as we bound its oscillations on the intervals $[t_{i,j}, t_{i,j+1}]$.

Pinning. For the boundary increments $k \in \{k_1^*, k_2^*, \dots\}$ consider the following pinning procedure. Let $k_0^* = i(k_0^*) = 0$ and define recursively for $n \in \mathbb{N}$

$$i(k_n^*) := \inf\{i > i(k_{n-1}^*) : \prod_{j=i(k_{n-1}^*)}^i (1 + \alpha_j) M_{k_n^*} > M_{k_{n-1}^*}\}.$$

We note that (5.18) and $\sum_j \alpha_j = \infty$ imply that $i(k_n^*) < \infty$ for all n . Now to define the pinning sequence let $t_{i,0} := k_i^*$ and for $j \in \{0, \dots, i(k_i^*) - i(k_{i-1}^*)\}$ set

$$t_{i,j} := \begin{cases} (1 + \alpha_{i(k_{i-1}^*)+j}) t_{i,j-1} & \text{if } j < i(k_i^*) - i(k_{i-1}^*) \\ M_{k_{i+1}^*} & \text{if } j = i(k_i^*) - i(k_{i-1}^*). \end{cases}$$

Now it follows from the definition of \bar{t} that

$$\bar{t}_{i,j} = t_{i,j} - M_{k_i^*} = M_{k_i^*} \left[\prod_{n=1}^j (1 + \alpha_{i(k_{i-1}^*)+n}) - 1 \right]. \quad (5.20)$$

By the polynomial decay in (S1) it follows that for any $\varepsilon > 0$, and $i, j > 0$

$$\mathbb{P} \left[\left| \frac{\bar{t}_{i,j}}{t_{i,j}} X_{k_i^*}(\bar{t}_{i,j}) \right| \geq \varepsilon \right] \leq \mathbb{P} \left[|X_{k_i^*}(\bar{t}_{i,j})| \geq \varepsilon \right] \leq \frac{C_\varepsilon}{(\bar{t}_{i,j})^\delta}.$$

By (5.20) and (5.19), the sum over $i, j > 0$ of the above probability is finite and therefore for any $\varepsilon > 0$

$$\mathbb{P} \left[\frac{\bar{t}_{i,j}}{t_{i,j}} |X_{k_i^*}(\bar{t}_{i,j})| \geq \varepsilon \text{ for infinitely many } (i, j) \right] = 0,$$

which implies that

$$\mathbb{P} \left[\limsup_{i,j} \frac{\bar{t}_{i,j}}{t_{i,j}} |X_{k_i^*}(\bar{t}_{i,j})| = 0 \right] = 1.$$

It remains to understand the behaviour of the boundary term in $[t_{i,j}, t_{i,j+1}]$.

Oscillations. Now we use (S3) to compute the oscillations between the pinned values of the boundary. Fix $\varepsilon > 0$ and consider the event Ω_{i_0} defined by

$$\Omega_{i_0} := \left\{ \sup_j \left| \frac{\bar{t}_{i,j}}{t_{i,j}} X_{k_i^*}(\bar{t}_{k_i^*,j}) \right| \leq \varepsilon, \quad \text{for } i > i_0 \right\}. \quad (5.21)$$

Therefore, on Ω_{i_0} for $t \in [t_{i,j}, t_{i,j+1}]$, and $j \geq 1$

$$\begin{aligned} & \left| \frac{\bar{t}}{t} X_{k_i^*}(\bar{t}) - \frac{\bar{t}_{i,j}}{t_{i,j}} X_{k_i^*}(\bar{t}_{i,j}) \right| \\ &= \left| \frac{1}{t} [\bar{t} X_{k_i^*}(\bar{t}) - \bar{t}_{i,j} X_{k_i^*}(\bar{t}_{i,j})] + \left(\frac{\bar{t}}{t} - \frac{\bar{t}_{i,j}}{t_{i,j}} \right) X_{k_i^*}(\bar{t}_{i,j}) \right| \\ &\leq \frac{1}{t} |\bar{t} X_{k_i^*}(\bar{t}) - \bar{t}_{i,j} X_{k_i^*}(\bar{t}_{i,j})| + (C_\alpha - 1)\varepsilon \end{aligned}$$

where

$$C_\alpha = \sup_{i,j} \sup_{t \in [t_{i,j}, t_{i,j+1}]} \frac{\bar{t}}{t_{i,j}} < \infty$$

Let $s := t - t_{i,j}$. By (S3) it follows that on Ω_{i_0}

$$\mathbb{P} \left[\sup_{s \leq t_{i,j+1} - t_{i,j}} \left| \frac{\bar{t}_{i,j} + s}{t_{i,j} + s} X_{k_i^*}(\bar{t}_{i,j} + s) - \frac{\bar{t}_{i,j}}{t_{i,j}} X_{k_i^*}(\bar{t}_{i,j}) \right| > C_\alpha \varepsilon \right] \leq \alpha_{i,j}^\beta C_\varepsilon.$$

As the sum of the above terms over $i, j \in \mathbb{N}$ is finite, by (5.21), it follows that

$$\mathbb{P} \left[\limsup_k V_k \leq C_\alpha \varepsilon \right] \geq \limsup_{i_0} \mathbb{P}(\Omega_{i_0}) = 1 \quad (5.22)$$

Since $\varepsilon > 0$ is arbitrary, from (5.15), (5.16) and (5.22) we conclude that (5.14) holds. \square

A Bounded increments

To deal with the case $i = 0$, in the case of small increments defined in (5.8), if $\lim_n \sum_{i=1}^n m_n^{i,0} < \infty$ it follows that $S_n^{i,0}$ converges to 0. For this reason assume without loss of generality that $m_k = m_k^{i,0}$ and that

$$\lim_n M_n = \lim_n \sum_{i=1}^n m_n^{i,0} \rightarrow \infty.$$

We next consider the truncated versions of X_k

$$Y_k := X_k \mathbb{1}_{\{|X_k| \leq \frac{M_k}{m_k}\}}.$$

The proof proceeds in two steps: first we show that

$$\mathbb{P}(Y_k \neq X_k \text{ i.o.}) = 0. \quad (A.1)$$

This implies that the limit of S_n equals the limit of $\bar{S}_n := \sum_{k=1}^n a_{n,k} Y_k$. The proof will be complete once we prove that

$$\mathbb{P} \left(\lim_n \bar{S}_n = 0 \right) = 1. \quad (A.2)$$

Proof of (A.1). Let $N(x) := \{k: \frac{M_k}{m_k} \leq x\}$, $F^*(a) := \mathbb{P}(|X^*| < a)$, and note that by the stochastic domination (S2)

$$\begin{aligned} \sum_k \mathbb{P}(Y_k \neq X_k) &\leq \sum_k \mathbb{P}\left(|X_k| \geq \frac{M_k}{m_k}\right) \leq \sum_j \mathbb{P}\left(|X^*| \geq \frac{M_k}{m_k}\right) \\ &\leq \sum_j \int_{x \geq \frac{M_k}{m_k}} dF^*(x) = \int N(x) dF^*(x) = \mathbb{E}[N(|X^*|)]. \end{aligned}$$

To obtain (A.1) it remains to prove that

$$\mathbb{E}[N(|X^*|)] < \infty.$$

This step follows from Lemma 2 of [9] which states that

$$\limsup \frac{N(x)}{x \log x} \leq 2. \quad (\text{A.3})$$

By (A.3), it follows that $N(x) \leq Cx^{1+\gamma}$ and by (S2), $\mathbb{E}[N(|X^*|)] < \infty$. \square

Proof of (A.2) As

$$\lim_n \mathbb{E}[\bar{S}_n] = 0,$$

to prove (A.2) it suffices to show that

$$\sum_k \frac{m_k^2}{M_k^2} \text{Var}(Y_k) < \infty. \quad (\text{A.4})$$

As $\frac{M_k}{m_k} \rightarrow \infty$ it follows that there is a C such that

$$\mathbb{E}[Y_k^2] \leq C \int_{|x| \leq \frac{M_k}{m_k}} x^2 dF^*(x).$$

Therefore, the sum in (A.4) can be bounded by

$$C \sum_k \frac{m_k^2}{M_k^2} \int_{|x| < \frac{M_k}{m_k}} x^2 dF^*(x) = C \int x^2 \sum_{k: \frac{M_k}{m_k} \geq |x|} \frac{m_k^2}{M_k^2} dF^*(x).$$

To complete the proof it remains to show that the right-hand side above is finite. This follows from the following claims whose proofs are given right after:

$$\sum_{k: \frac{M_k}{m_k} \geq |x|} \frac{m_k^2}{M_k^2} \leq 2 \int_{y \geq |x|} \frac{N(y)}{y^3} dy, \quad (\text{A.5})$$

and

$$\int x^2 \int_{y \geq |x|} \frac{N(y)}{y^3} dy dF^*(x) < \infty. \quad (\text{A.6})$$

\square

proof of (A.6) By (A.3) it follows that there are $C > 0$ and $\gamma \in (0, 1)$ such that $N(x) \leq Cx^{1+\gamma}$. Therefore

$$\begin{aligned} & \int x^2 \int_{y \geq |x|} \frac{N(y)}{y^3} dy dF^*(x) \leq \int x^2 \int_{y \geq |x|} \frac{Cy^{1+\gamma}}{y^3} dy dF^*(x) \\ & = \int x^2 \int_{y \geq |x|} \frac{C}{y^{2-\gamma}} dy dF^*(x) = \int x^2 \frac{C}{(1-\gamma)x^{1-\gamma}} dF^*(x) \\ & = \frac{C}{1-\gamma} \mathbb{E} \left[|X^*|^{1+\gamma} \right] < \infty. \end{aligned}$$

□

proof of (A.5) Observe that by the definition of N and integration by parts

$$\begin{aligned} \sum_{k: |x| < \frac{M_k}{m_k} \leq z} \frac{m_k^2}{M_k^2} &= \int_{|x| < y < z} \frac{dN(y)}{y^2} \\ &= \frac{N(z)}{z^2} - \frac{N(|x|)}{x^2} + 2 \int_{|x| < y < z} \frac{N(y)}{y^3} dy. \end{aligned}$$

Furthermore, since $N(z) \leq N(y)$ for $z \leq y$ and $\frac{1}{z^2} = 2 \int_z^\infty \frac{1}{y^3} dy$ it follows from (A.6) that

$$\frac{N(z)}{z^2} \leq 2 \int_z^\infty \frac{N(y)}{y^3} dy \rightarrow 0,$$

and so

$$\begin{aligned} \sum_{k: |x| < \frac{M_k}{m_k}} \frac{m_k^2}{M_k^2} &= \lim_z \sum_{k: |x| < \frac{M_k}{m_k} < z} \frac{m_k^2}{M_k^2} \int_{|x| < y < z} \frac{dN(y)}{y^2} \\ &\leq 2 \int_{|x| < y} \frac{N(y)}{y^3} dy. \end{aligned}$$

□

Acknowledgements

The research in this paper was partially supported through NWO Gravitation Grant NETWORKS-024.002.003. We are grateful to Evgeny Verbitsky for pointing the paper of Rohatgi [12] and related useful discussions.

References

- [1] Luca Avena, Yuki Chino, Conrado da Costa, and Frank den Hollander. Random walk in cooling random environment: ergodic limits and concentration inequalities. *Electron. J. Probab.*, 24:1–35, 2019.
- [2] Luca Avena, Yuki Chino, Conrado da Costa, and Frank den Hollander. Random walk in cooling random environment: recurrence versus transience and mixed fluctuations. *To appear in AIHP*, arXiv:1903.09200, 2019.

- [3] Luca Avena and Frank den Hollander. Random walks in cooling random environments. In Vlasdas Sidoravicius, editor, *Sojourns in Probability Theory and Statistical Physics - III*, pages 23–42, Singapore, 2019. Springer Singapore.
- [4] Jacob Bernoulli. *Ars Conjectandi, Opus Posthumum*. Basileae, Impensis Thurnisiorum, Fratrum Werke 3, 1713.
- [5] N. H. Bingham. *Riesz Means and Beurling Moving Averages*, chapter 8, pages 159–172. World Scientific Europe, 2019.
- [6] N.H. Bingham and Bujar Gashi. Voronoi means, moving averages, and power series. *Journal of Mathematical Analysis and Applications*, 449(1):682–696, 2017.
- [7] Rick Durrett. *Probability: Theory and Examples*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 5 edition, 2019.
- [8] Sacha Friedli and Yvan Velenik. *Statistical Mechanics of Lattice Systems: A Concrete Mathematical Introduction*. Cambridge University Press, 2017.
- [9] Benton Jamison, Steven Orey, and William Pruitt. Convergence of weighted averages of independent random variables. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 4(1):40–44, Mar 1965.
- [10] Oleg Klesov. *Limit Theorems for Multi-Indexed Sums of Random Variables*, volume 71 of *Probability Theory and Stochastic Modelling*. Springer, Berlin, Heidelberg, 2014.
- [11] William Pruitt. Summability of independent random variables. *J. Math. and Mech.*, 15(5):769–776, 1966.
- [12] Vijay Rohatgi. Convergence of weighted sums of independent random variables. *Mathematical Proceedings of the Cambridge Philosophical Society*, 69(2):305–307, 1971.
- [13] Pál Révész. *Laws of Large Numbers*. Academic Pres, 1 edition, 1967.
- [14] William Stout. Some results on the complete and almost sure convergence of linear combinations of independent random variables and martingale differences. *Ann. Math. Statist.*, 39(5):1549–1562, 10 1968.
- [15] Matthew Sullivan, Steven Pirotta, V. Chernenko, GH Wu, G Balasubramanium, Susan Hua, and Harsh Chopra. Magnetic mosaics in crystalline tiles: The novel concept of polymagnets (invited). *International Journal of Applied Electromagnetics and Mechanics*, 22:11–23, 10 2005.