

Butson-Hadamard matrices and Plotkin-optimal p^k -ary codes

Damla Acar¹, Bülent Saraç¹, and Oğuz Yayla²

¹ Hacettepe University, Department of Mathematics,
Beytepe, 06800, Ankara, Turkey

damla.acar@hacettepe.edu.tr, bsarac@hacettepe.edu.tr

² Middle East Technical University, Institute of Applied Mathematics,
06800, Ankara, Turkey
oguz@metu.edu.tr

Abstract. A Butson-Hadamard matrix H is a square matrix of dimension n whose entries are complex roots of unity such that $HH^* = nI$. In the first part of this work, some new results on generalized Gray map are studied. In the second part, codes obtained from Butson-Hadamard matrices and some bounds on the minimum distance of these codes are proved. In particular, we show that the code obtained from a Butson-Hadamard matrix meets the Plotkin bound under a non-homogeneous weight. We also give the parameters of some code families which are obtained from modified Butson-Hadamard matrices under a (non)homogeneous Gray map.

Keywords: Butson-Hadamard matrices, codes, generalized Gray map, Plotkin bound

2020 AMS Subject Classification: 05B20,94B60,94B65

1 Introduction

A *Hadamard matrix* of order n is an $n \times n$ matrix whose entries are $1, -1$ satisfying $HH^T = nI_n$. Here I_n is the $n \times n$ identity matrix. Hadamard [8] conjectured that such matrices exist for every n that is a multiple of 4. See [9,11,16] for more information about Hadamard matrices and their applications. Hadamard matrices can be generalized in many ways. Two of them are Butson-Hadamard (BH) matrices which are introduced by Butson in [3] and generalized Hadamard (GH) matrices by Drake in [6].

A *Butson-Hadamard matrix* H is an $n \times n$ matrix whose entries are complex roots of unity such that

$$HH^* = nI_n$$

where H^* is the Hermitian transpose of H . If all the entries of H are k -th roots of unity then we say that H is a $BH(n, k)$.

Another generalization of Hadamard matrices is *(group) generalized Hadamard matrices*. Let $G = \{g_1, g_2, \dots, g_m\}$ be a finite group and let H be an $n \times n$ matrix whose entries are elements of G . For convenience, we identify the group G with

the naturally embedded copy of G in the group ring $\mathbb{Z}[G] = \{\sum_{i=1}^m a_i g_i \mid a_i \in \mathbb{Z}\}$. A natural involution on $\mathbb{Z}[G]$, which we shall call conjugation, is defined by

$$\overline{\left(\sum a_i g_i\right)} := \sum a_i g_i^{-1}. \quad (1)$$

Then, for the matrices with entries in $\mathbb{Z}[G]$, define an adjoint, $M^* = [m_{ij}]^* := [\overline{m_{ji}}]$. That is the adjoint is performed by first taking the transpose of the matrix and then conjugating each entry. We call that an $n \times n$ matrix H whose entries are in G is a *(group) generalized Hadamard matrix* if

$$HH^* \equiv nI_n \pmod{\sum_{g \in G} g}. \quad (2)$$

For brevity, we say that such a matrix H is a $\text{GH}(n, G)$.

For example, the following is a $\text{GH}(5, C_5)$, where ζ is a generator of the cyclic group C_5 of order 5.

$$H = \begin{pmatrix} 1 & \zeta & \zeta^4 & \zeta^4 & \zeta \\ \zeta & 1 & \zeta & \zeta^4 & \zeta^4 \\ \zeta^4 & \zeta & 1 & \zeta & \zeta^4 \\ \zeta^4 & \zeta^4 & \zeta & 1 & \zeta \\ \zeta & \zeta^4 & \zeta^4 & \zeta & 1 \end{pmatrix}$$

Also we can say that a $\text{GH}(n, G)$ can exist only if n is a multiple of $|G|$. See also [12] for more information about generalized Hadamard matrices.

It is known that a BH (resp. GH) matrix can be transformed to an equivalent matrix consisting of 1s in the first row and column by dividing rows or columns, or by interchanging rows or columns. So, a BH (resp. GH) matrix is said to be *normalized* if the first row and column consist entirely of 1s.

In this paper we first study the parameters for code families where the rows of BH and modified BH matrices are assigned to be codewords. The minimum distance between the rows of normalized GH matrices has been studied widely, see [11, Sections 4.4 and 9.4]. In addition to minimum distance, the rank and kernel of the codes obtained from a GH matrix were recently studied in [4,5], and codes from a cocyclic GH matrix and their equivalence to combinatorial difference sets are studied in [1,2]. On the other hand, Greferath, McGuire and O'Sullivan [7] showed that the codes obtained from BH matrices meet the Plotkin bound under any homogeneous weight. Stepanov [18] also constructed codes obtained from modified BH matrices, which have parameters close to the Plotkin bound.

In this paper, we give a lower bound for the minimum distance of the codes obtained from a normalized BH matrix and an amenable BH matrix in Theorem 2 and 3, respectively. Then, we prove their minimum distance under a homogeneous Gray map, in Corollary 1. We also consider codes obtained from the image of the modified BH matrices under a non-homogeneous Gray map, and we prove their minimum distances in Theorem 4. We also show that the distances between

the rows are same, hence the code is equidistant. Moreover, we show that the code meets the Plotkin bound in Corollary 4.

The paper is organized as follows. We study the generalized Plotkin bound in Section 2 and generalized Gray map in Section 3. In Section 4 we give some results on the minimum distance of codes obtained from BH and modified BH matrices. The codes obtained from the image of BH matrices under the non-homogeneous Gray map are given in Section 5.

2 The generalized Plotkin bound

Hadamard codes have been widely studied in the literature, see for instance [11]. Greferath et al. in [7] considered BH matrices and determined the parameters of the code obtained from normalized BH matrices, where they considered homogeneous weights. Their result is given below, but we first give the definition of a homogeneous weight.

Definition 1. [7] *A real-valued function w on the finite ring R is called a homogeneous weight if $w(0) = 0$ and the following hold:*

- (i) $Rx = Ry$ implies $w(x) = w(y)$ for all $x, y \in R$.
- (ii) There exists a real number γ such that

$$\sum_{y \in Rx} w(y) = \gamma |Rx|, \forall x \in R \setminus \{0\},$$

where the number γ is called the average value of w on R .

For instance, the Lee weight w_L on \mathbb{Z}_4 defined by $w_L(0) = 0, w_L(1) = w_L(3) = 1$ and $w_L(2) = 2$ is a homogeneous weight with the average value $\gamma = 1$.

A q -ary (n, M, d) code is defined to be a nonempty subset of \mathbb{Z}_q^n of size M , where d is the minimum distance of the code determined by the distance function induced by a specified homogeneous weight on \mathbb{Z}_q^n . For an (n, M, d) code with $d > \gamma n$, the *generalized Plotkin bound* states that $M \leq d/(d - \gamma n)$, and it is called *Plotkin-optimal* when

$$M > d/(d - \gamma n) - 1, \tag{3}$$

where γ is the average value of the specified homogeneous weight.

Theorem 1. [7] *Let $H = [\zeta_h^{a_{ij}}]$ be a normalized Butson-Hadamard matrix of type $\text{BH}(n, h)$ for some primitive h -th root of unity ζ_h . Then the code in $(\mathbb{Z}_h)^{n-1}$ formed by taking the rows of $H' = [a_{ij}]$ and omitting the first coordinate is a code over \mathbb{Z}_h with parameters $(n-1, n, \gamma n)$ that meets the Plotkin bound.*

3 Generalized Gray maps

In this section we consider two kinds of generalized Gray maps. At first, we give the generalized Gray map G_1 , originated in [10] (see Definition 2 below). Note that G_1 gives rise to a homogeneous weight. As Theorem 1 states, codes obtained from rows of BH matrices satisfy the Plotkin bound under any homogeneous weight [7]. On the other hand, very little is known when the map is non-homogeneous. Hence, we consider a generalized Gray map G_2 which yields a non-homogeneous weight w_2 (see Definition 3). We also prove some lemmas on G_2 in which we give the minimum distance of BH codes under the weight w_2 defined by G_2 .

Definition 2. Let k be a positive integer, u be any element of \mathbb{Z}_{p^k} and $u = \sum_{i=1}^k u_i p^{i-1}$ its p -ary expansion for some $u_i \in \{0, 1, \dots, p-1\}$. The image of u by a Gray map G_1 is defined to be the Boolean function on $GF(p)^{k-1}$:

$$G_1(u) : (y_1, \dots, y_{k-1}) \rightarrow u_k + \sum_{i=1}^{k-1} u_i y_i.$$

Example 1. Suppose that $k = 3$ and $p = 2$. Then the Gray map for u is defined as follows:

$$G_1(u) : (y_1, y_2) \rightarrow u_3 + u_1 y_1 + u_2 y_2,$$

where $(y_1, y_2) \in \mathbb{Z}_2^2$, in lexicographical order, are taken as follows:

$$\begin{array}{c|c} y_2 & y_1 \\ \hline 0 & 0 \\ 0 & 1 \\ 1 & 0 \\ 1 & 1 \end{array}$$

If $u = 6$, then its binary representation is equal to $(u_3 u_2 u_1) = (110)$. Therefore if $(y_1, y_2) = (0, 0)$ then we get $G_1(6) = 1$. If we continue like this we can get Boolean function $G_1(u) = (1100)$.

Let w be the Hamming weight and w_1 be a weight on \mathbb{Z}_{p^k} defined by $w_1(u) = w(G_1(u))$. We note that $w_1(0) = 0$ and for any $u \neq 0$

$$w_1(u) := \begin{cases} p^{k-1} - p^{k-2} & \text{if } u \in \mathbb{Z}_{p^k} \setminus \{p^{k-1}, 2p^{k-1}, \dots, (p-1)p^{k-1}\} \\ p^{k-1} & \text{otherwise.} \end{cases}$$

Then one can easily deduce from Definition 1 that w_1 is a homogeneous weight. In what follows, we list some properties of G_1 .

- i. Let k be a positive integer and $u = \sum_{i=1}^k u_i 2^{i-1} \in \mathbb{Z}_{2^k}$ for some $u_i \in \{0, 1\}$. Then $G_1(u)$ is a homomorphism from \mathbb{Z}_2^{k-1} to \mathbb{Z}_2 .
- ii. Let $a, b \in \mathbb{Z}_{2^k}$ such that $a \neq b$. Then

$$d(G_1(a), G_1(b)) = \begin{cases} 2^{k-1} & \text{if } a - b = 2^{k-1} \\ 2^{k-2} & \text{otherwise.} \end{cases}$$

Here $d(G_1(a), G_1(b))$ is the Hamming distance between $G_1(a)$ and $G_1(b)$.

In this paper we also consider another Gray map which yields a non-homogeneous weight.

Definition 3. [20] Let $p > 2$ be a prime number and k be a positive integer. A Gray map, denoted G_2 , on \mathbb{Z}_{p^k} is defined as follows:

- i. If $u \leq p^{k-1}$, then $G_2(u) \in \mathbb{Z}_p^{p^{k-1}}$ has 1 in the first u location and 0's elsewhere.
- ii. If $u \geq p^{k-1} + 1$, then $G_2(u) = \bar{q} + G_2(r)$, where q and $r < p^{k-1}$ are positive integers such that $u = qp^{k-1} + r$ and $\bar{q} = (qq \cdots qq)$.

Let w_2 be the weight on \mathbb{Z}_{p^k} defined by $w_2(u) = w(G_2(u))$. Then we have

$$w_2(u) := \begin{cases} u & \text{if } u \leq p^{k-1} \\ p^{k-1} & \text{if } p^{k-1} \leq u \leq p^k - p^{k-1} \\ p^k - u & \text{if } p^k - p^{k-1} < u \leq p^k - 1. \end{cases}$$

As the following example shows, w_2 is not homogeneous in general.

Example 2. Let $p = 3$, $k = 2$ and $R = \mathbb{Z}_9$. Now if we take $x = 2$ and $y = 8$, then $Rx = Ry$; but since $G_2(2) = (110)$, $G_2(8) = (002)$, $w_2(x)$ and $w_2(y)$ are not equal. Therefore w_2 is not homogeneous by Definition 1.

Now we give four lemmas that we will use in the proof of Theorem 4.

Lemma 1. Let $m, n \in \mathbb{Z}_{p^k}$ with $m, n \neq 0$. If either $m, n < p^{k-1}$ and $m + n = p^{k-1}$ or $m > n > (p - 1)p^{k-1}$ and $m - n = (p - 1)p^{k-1}$, then the total number of 0's in $G_2(m)$ and $G_2(n)$ is p^{k-1} .

Proof. If $m, n < p^{k-1}$ and $m + n = p^{k-1}$ then the result is clear from the definition of G_2 . Let $m > n$ and $m - n = (p - 1)p^{k-1}$. Since $m > n$ and $m = n + (p - 1)p^{k-1}$ we see that $n < p^{k-1}$. Therefore, the number of 0's in $G_2(n)$ is equal to $p^{k-1} - n$. On the other hand, $G_2(m) = G_2((p - 1)p^{k-1} + n)$, and by definition, the number of 0's in $G_2(m)$ is n . This completes the proof. \square

Example 3. Let $p = 3$, $k = 3$. If we take $n = 8$ and $m = n + (p - 1)p^{k-1} = 26$ then we can easily find that $G_2(n) = G_2(8) = (111111110)$ and $G_2(m) = G_2(26) = (000000002)$. Therefore the total number of 0's in $G_2(m)$ and $G_2(n)$ is equal to $9 = p^{k-1}$.

Lemma 2. Let p be a prime number, $k \in \mathbb{Z}^+$, $i, j \in \mathbb{Z}_{p^k}$ and $|i - j| = n$. Then

$$d(G_2(i), G_2(j)) = \begin{cases} n & \text{if } n < p^{k-1} \\ p^{k-1} & \text{otherwise.} \end{cases}$$

Proof. First we consider $n < p^{k-1}$. Without loss of generality we take $i > j$. If $i \leq p^{k-1}$ then $G_2(i) = (111 \dots 11000 \dots 00)$ by the definition of G_2 , where it has the number 1 in the first i coordinates and the others are 0. We also have $j < p^{k-1}$ and $j = i - n$ as $j < i$. Therefore $G_2(j) = (111 \dots 11000 \dots 00)$ such that

the first $i - n$ coordinates are 1 and the others are 0. So $d(G_2(i), G_2(j)) = n$. Now if $i > p^{k-1}$ then we can write $i = qp^{k-1} + r$ for some $q, r \in \mathbb{Z}^+$ such that $r < p^{k-1}$. Then $G_2(i) = \bar{q} + G_2(r)$. Also $j = i - n$ and $n < p^{k-1}$. Now, if $n < r$ then we can say that $j = qp^{k-1} + r - n$ and $r - n < p^{k-1}$. Therefore $d(G_2(i), G_2(j)) = n$. If $n \geq r$ then $j = (q-1)p^{k-1} + p^{k-1} - (n-r)$. Hence $G_2(i) = (q+1 \quad q+1 \dots q+1 \underbrace{qq \dots qq}_{p^{k-1}-r})$ and $G_2(j) = (\underbrace{qq \dots qq}_{n-r} \quad q-1 \quad q-1 \dots q-1)$. Also

we can say that $p^{k-1} - r > n - r$ since $n > r, n < p^{k-1}$. Therefore there are $p^{k-1} - r - n + r$ same coordinates in $G_2(i)$ and $G_2(j)$. So $d(G_2(i), G_2(j)) = n$.

Now suppose that $j = qp^{k-1} + r, n = mp^{k-1} + r'$ and $r + r' = lp^{k-1} + r''$ for some $q, m, l, r, r', r'' \in \mathbb{Z}^+$ such that $r, r', r'' < p^{k-1}$. Then $i = (q+m+l)p^{k-1} + r''$ and since $i < p^k$ we must have $q + m + l < p$. It follows that $G(i) = \overline{q+m+l} + G_2(r'')$ and $G_2(j) = \bar{q} + G_2(r)$. Since $r \neq r''$, we obtain $d(G_2(i), G_2(j)) = p^{k-1}$. Also it is clear that if $j \leq p^{k-1}$, then $d(G_2(i), G_2(j)) = p^{k-1}$. \square

Lemma 3. *Let p be a prime number, $k \in \mathbb{Z}^+$, ζ be a primitive p^k -th root of unity and $a_i \in \mathbb{Z}^+$ for $i = 1, 2, \dots, n$. If $\zeta^{a_1} + \dots + \zeta^{a_n} = 0$, then the total number of zeros in $[G_2(a_1), G_2(a_2), \dots, G_2(a_n)]$ is equal to np^{k-2} .*

Proof. We know that a_i 's satisfying $\zeta^{a_1} + \zeta^{a_2} + \dots + \zeta^{a_n} = 0$ will only be of the form $a_i = ip^x + j$, where $x < k, j < p^x$ and $i = 1, 2, \dots, p^{k-x}$, see [13, Theorem 3.3]. Here we will give a proof only for the case $j = 0$, since we will give the case where j is nonzero in Corollary. 3 We know that obtaining 0 in the images $G_2(a_i)$ for $i = 1, 2, \dots, n$ is only possible for $1 \leq a_i < p^{k-1}$ or $(p-1)p^{k-1} < a_i \leq p^k$. So the number of 0 in $G_2(a_1), G_2(a_2), \dots, G_2(a_{p^{k-x}})$ are equal to $p^{k-1} - p^x, p^{k-1} - 2p^x, \dots, (p^{k-1-x} - 1)p^x$, respectively. Here there are $p^{k-1-x} - 1$ elements. On the other hand, the number of zero in $G_2(a_i)$ for $(p-1)p^{k-1} < a_i \leq p^k$ are $p^x, 2p^x, \dots, (p^{k-1-x} - 1)p^{k-1}, p^{k-1}$, in order. Therefore, the total number of zeros in $G_2(a_1), G_2(a_2), \dots, G_2(a_{p^{k-x}})$ is

$$(p^{k-1-x} - 1)p^{k-1} + p^{k-1} = p^{k-2}p^{k-x} = np^{k-2}.$$

\square

Example 4. Let $p = 3, k = 3$ and ζ be the primitive 27-th root of unity. Also we know that $\zeta^3 + \zeta^6 + \zeta^9 + \zeta^{12} + \zeta^{15} + \zeta^{18} + \zeta^{21} + \zeta^{24} + \zeta^{27} = 0$. Then $G_2(3) = (111000000), G_2(6) = (111111000), G_2(9) = (111111111), G_2(12) = (222111111), G_2(15) = (222222111), G_2(18) = (222222222), G_2(21) = (000111111), G_2(24) = (000001111), G_2(27) = (000000000)$. Total number of zeros in these images is equal to $np^{k-2} = 27$.

4 Butson-Hadamard codes

In this section we will study codes obtained via assigning rows of normalized BH matrices and their translates as codewords. Namely, let ζ be a primitive k -th root

of unity, $H = [\zeta^{a_{ij}}]$ be a normalized BH(n, k) matrix and $[a_{ij}]$ be the matrix with entries taken from the exponents of corresponding entries of H . Then codes are constructed from the rows of $[a_{ij}]$.

Codes from normalized generalized Hadamard matrices and their translates are widely studied, see [11, Chapter 4] and references therein. Besides, codes constructed from modified quaternary complex BH matrices are given in [17], which are close to the Plotkin bound. Another type modified BH matrices and the codes obtained from them are studied in [14].

Definition 4. Let n, k be positive integers and ζ be a primitive k -th root of unity, $H = [\zeta^{a_{ij}}]$ be a normalized BH(n, k) matrix. Suppose that N be the set of all k -th roots of unity. If R_i is the i -th row of H then, uR_i is called a translate of R_i for some $u \in N$.

We use the definitions given in [11, Definition 4.32] and we take the Hamming weight for the codes obtained from BH matrices in the following theorem.

Theorem 2. Let n, k be positive integers, ζ be a primitive k -th root of unity, $H = [\zeta^{a_{ij}}]$ be a normalized BH(n, k) matrix and N be the set of all k -th roots of unity. Let $H' = [a_{ij}]$ and $l = \min\{i \geq 2 : i|k\}$.

- i. A_k the k -ary $(n-1, n, d_A)$ code consisting of the rows of H' with the first column deleted. Here $d_A \geq n - \frac{n}{l}$.
- ii. B_k the k -ary $(n-1, nk, d_B)$ code consisting of the rows of exponent matrix of the translates uH , $u \in N$ with first column deleted. Here $d_B \geq n - \frac{n}{l} - 1$
- iii. C_k the k -ary (n, nk, d_C) code consisting of the rows of exponent matrix of the translates uH , $u \in N$. Here $d_C \geq n - \frac{n}{l}$
- iv. D_k , $n = k$, the k -ary $(n+1, n^2, d_D)$ code consisting of the rows of exponent matrix of $(uH)_c$ for all $u \in N$ and any fixed noninitial column c of H . Here $d_D \geq n - \frac{n}{l}$.

Proof.

- i. We first note that l is a prime number. If ζ is a primitive k -th root of unity then $\zeta^{\frac{k}{l}}$ is l -th root of unity. Then $(\zeta^{\frac{k}{l}}) + (\zeta^{\frac{k}{l}})^2 + \dots + (\zeta^{\frac{k}{l}})^l = 0$. This implies that we must have at least l elements in order to get vanishing sum. On the other hand, since the rows of BH matrices are orthogonal, the number of same elements in two rows of H must be smaller than $\frac{n}{l}$. That is $n - d_A \leq \frac{n}{l}$. So we have $d_A \geq n - \frac{n}{l}$.
- ii. Let $u = \zeta^m$ be an element of N . Then it is clear that the codes A_k and shifting of A_k with m have same minimum distance. Now let R_i and R_i^u be the i -th row of H and uH , respectively, with the first column deleted for $i = 1, 2, \dots, n$. So $d(R_i, R_j^u) = n - 1$ if $i = j$. On the other hand, as $R_j^u = uR_j$ we have $d(R_i, R_j^u) = n - 1 - \#\{u|u \in R_i^{-1}R_j^u\}$ for $i \neq j$. If we use the same argument in the proof of (i), we can say that $\#\{u|u \in R_i^{-1}R_j^u\} \leq \frac{n}{l}$. Therefore $d_B \geq n - \frac{n}{l} - 1$.

- iii. C_k has codewords of length n . Similar to the proof of (ii), we have $d_C \geq n - \frac{n}{7}$
 iv. D_k and C_k are only one column different from each other. Therefore $d_D \geq n - \frac{n}{7}$.

□

Example 5. Let ζ be a primitive 36-th root of unity and

$$H_1 = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & \zeta^{12} & \zeta^{24} & \zeta^{28} & \zeta^4 & \zeta^{16} & 1 & \zeta^{12} & \zeta^{24} & 1 & \zeta^{12} & \zeta^{24} \\ 1 & \zeta^{24} & \zeta^{12} & \zeta^{20} & \zeta^8 & \zeta^{32} & 1 & \zeta^{24} & \zeta^{12} & 1 & \zeta^{24} & \zeta^{12} \\ 1 & \zeta^{27} & 1 & 1 & 1 & 1 & \zeta^{18} & \zeta^9 & \zeta^{18} & \zeta^{18} & \zeta^{18} & \zeta^{18} \\ 1 & \zeta^3 & \zeta^{24} & \zeta^{28} & \zeta^4 & \zeta^{16} & \zeta^{18} & \zeta^{21} & \zeta^6 & \zeta^{18} & \zeta^{30} & \zeta^6 \\ 1 & \zeta^{15} & \zeta^{12} & \zeta^{20} & \zeta^8 & \zeta^{32} & \zeta^{18} & \zeta^{33} & \zeta^{30} & \zeta^{18} & \zeta^6 & \zeta^{30} \\ 1 & 1 & 1 & \zeta^{18} & \zeta^{18} & \zeta^{18} & \zeta^9 & 1 & 1 & \zeta^{27} & \zeta^{18} & \zeta^{18} \\ 1 & \zeta^{12} & \zeta^{24} & \zeta^{10} & \zeta^{22} & \zeta^{34} & \zeta^9 & \zeta^{12} & \zeta^{24} & \zeta^{27} & \zeta^{30} & \zeta^6 \\ 1 & \zeta^{24} & \zeta^{12} & \zeta^2 & \zeta^{26} & \zeta^{14} & \zeta^9 & \zeta^{24} & \zeta^{12} & \zeta^{27} & \zeta^6 & \zeta^{30} \\ 1 & \zeta^{27} & 1 & \zeta^{18} & \zeta^{18} & \zeta^{18} & \zeta^{27} & \zeta^9 & \zeta^{18} & \zeta^9 & 1 & 1 \\ 1 & \zeta^3 & \zeta^{24} & \zeta^{10} & \zeta^{22} & \zeta^{34} & \zeta^{27} & \zeta^{21} & \zeta^6 & \zeta^9 & \zeta^{12} & \zeta^{24} \\ 1 & \zeta^{15} & \zeta^{12} & \zeta^2 & \zeta^{26} & \zeta^{14} & \zeta^{27} & \zeta^{33} & \zeta^{30} & \zeta^9 & \zeta^{24} & \zeta^{12} \end{pmatrix}$$

be a BH(12, 36) matrix. Here the minimum distance between the rows of H_1 is equal to $d_A = 7$ by using SageMath [19], therefore $d_A \geq n - \frac{n}{7}$ is satisfied, where $n = 12$ and $l = \min\{i \geq 2 : i|36\} = 2$. Similarly, for a primitive 10-th root of unity, ζ , let

$$H_2 = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & \zeta^5 & \zeta^3 & \zeta^3 & \zeta^5 & \zeta^9 & \zeta^8 & \zeta^7 & \zeta \\ 1 & \zeta^4 & \zeta^5 & \zeta^7 & \zeta & \zeta^3 & \zeta^5 & \zeta^9 & \zeta^9 \\ 1 & \zeta^3 & \zeta^7 & \zeta^5 & \zeta & \zeta^8 & \zeta^9 & \zeta^3 & \zeta^5 \\ 1 & \zeta^9 & \zeta & \zeta^5 & \zeta^5 & \zeta^3 & \zeta^7 & \zeta^2 & \zeta^7 \\ 1 & \zeta^9 & \zeta^5 & \zeta & \zeta^3 & \zeta^5 & \zeta & \zeta^7 & \zeta^6 \\ 1 & \zeta & \zeta^7 & \zeta^9 & \zeta^6 & \zeta & \zeta^5 & \zeta^5 & \zeta^3 \\ 1 & \zeta^7 & \zeta^9 & \zeta^4 & \zeta^9 & \zeta^5 & \zeta^3 & \zeta^5 & \zeta \\ 1 & \zeta^5 & \zeta^2 & \zeta^9 & \zeta^7 & \zeta^7 & \zeta^3 & \zeta & \zeta^5 \end{pmatrix}$$

be a BH(9, 10) matrix. It can be verified that the minimum distance between the rows of H_2 is equal to $d_A = 7$ and satisfies $d_A \geq n - \frac{n}{7}$, where $n = 9$ and $l = 2$. Now consider B_k . In this case, we have A_k and its translates as codewords. If we calculate the minimum distance by SageMath, we get $d_B = 5$ and so $d_B \geq n(1 - \frac{1}{7}) - 1$. Again, by using SageMath, we can get that the minimum distance for C_k is $d_C = 6$, which satisfies $d_C \geq n - \frac{n}{7}$.

Let ζ be a primitive 6-th root of unity. Then

$$H_2 = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & \zeta & \zeta^2 & \zeta^3 & \zeta^4 & \zeta^5 \\ 1 & \zeta^2 & \zeta^4 & 1 & \zeta^2 & \zeta^4 \\ 1 & \zeta^3 & 1 & \zeta^3 & 1 & \zeta^3 \\ 1 & \zeta^4 & \zeta^2 & 1 & \zeta^4 & \zeta^2 \\ 1 & \zeta^5 & \zeta^4 & \zeta^3 & \zeta^2 & \zeta \end{pmatrix}$$

is a BH(6, 6) matrix. If we calculate the minimum distance for D_k by SageMath, we get $d_D = 3$, so $d_D \geq n(1 - \frac{1}{7})$ holds.

Remark 1. We note that it is easy to get a code with larger alphabet size without decreasing the minimum distance by combining two BH matrices under Chinese Remainder Theorem. Let h, k be distinct prime numbers, ζ, α and β be primitive h -th, k -th and hk -th roots of unities, respectively. Also $H_1 = [\zeta^{a_{ij}}]$ and $H_2 = [\alpha^{b_{ij}}]$ be normalized BH(n, h) and BH(n, k) matrices, respectively. Let d_1 and d_2 be the minimum distance between the rows of $[a_{ij}]$ and $[b_{ij}]$, respectively. Let $H = [\beta^{c_{ij}}]$ be the normalized matrix obtained from H_1 and H_2 such that $c_{ij} \equiv a_{ij} \pmod{h}$ and $c_{ij} \equiv b_{ij} \pmod{k}$. Then the minimum distance of rows of $[c_{ij}]$ satisfies $d \geq \max\{d_1, d_2\}$.

Proposition 1. *Let R be a Frobenius ring and let $A = [a_{ij}]$ be an $n \times n$ matrix over R . Suppose that for any $1 \leq k, l \leq n$ with $k \neq l$, the set*

$$R_k - R_l := \{a_{k1} - a_{l1}, a_{k2} - a_{l2}, \dots, a_{kn} - a_{ln}\}, \quad (4)$$

is a disjoint union of the cosets of right or left ideals of R . Here R_k and R_l are the k -th and l -th rows of A , respectively. Then for any generating character χ of R , the matrix $H = [\chi(a_{ij})]$ is a Butson-Hadamard matrix.

Proof. Suppose that for $1 \leq i, j \leq n$ with $i \neq j$, S_i and S_j be the distinct rows of H . Then $S_i = (\chi(a_{i1}), \chi(a_{i2}), \dots, \chi(a_{in}))$ and $S_j = (\chi(a_{j1}), \chi(a_{j2}), \dots, \chi(a_{jn}))$.

Therefore $S_i S_j^* = \sum_{k=1}^n \chi(a_{ik} - a_{jk})$. We know that the sum of the images under χ over a right (or left) ideal of R is equal to 0. Then for a fixed element $a \in R$ and a right (or left) ideal I of R , we have $\sum_{i \in I} \chi(a + i) = \sum_{i \in I} \chi(a)\chi(i) = \chi(a) \sum_{i \in I} \chi(i) = 0$; so we can say that $S_i S_j^* = \sum_{k=1}^n \chi(a_{ik} - a_{jk}) = 0$.

If $i = j$ then $\sum_{k=1}^n \chi(a_{ik} - a_{jk}) = \sum_{k=1}^n \chi(a_{ik} - a_{jk}) = n$. Therefore H is a Butson-Hadamard matrix. \square

Definition 5. *Let R be a Frobenius ring and $A = [a_{ij}]$ be an $n \times n$ matrix over R . Also suppose that A satisfies (4). Then for any generating character χ of R , the Butson-Hadamard matrix $H = [\chi(a_{ij})]$ is called an amenable Butson-Hadamard matrix.*

Example 6. Suppose that R is a finite Frobenius ring of characteristic t and χ is a generating character for R , $B : L \times M \rightarrow R$ be a nondegenerate bilinear pairing for the finite left R -module L and finite right R -module M . Then $H = [\chi(B(x, y))]_{L \times M}$ is a BH(t, m) Butson-Hadamard matrix, where $m = |L| = |M|$ by [15, Theorem 10]. All matrices created in this way are amenable Butson-Hadamard matrices since the elements in the row differences in $[B(x, y)]_{L \times M}$ form left ideals of R .

Bilinear pairings give us amenable Butson-Hadamard matrices, but not all amenable BH matrices have to be of this type. The matrix below is a counterexample.

Example 7. Let $R = \mathbb{Z}_6$ and A be a 6×6 matrix over R given as below.

$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 3 & 3 & 4 \\ 0 & 1 & 3 & 5 & 0 & 3 & 3 \\ 0 & 2 & 5 & 4 & 3 & 0 & 2 \\ 0 & 3 & 4 & 2 & 5 & 2 & 0 \\ 0 & 4 & 2 & 0 & 3 & 4 & 1 \\ 0 & 4 & 2 & 3 & 1 & 0 & 4 \end{pmatrix}$$

It is readily checked that all pairwise row differences of A are the unions of the cosets of $3\mathbb{Z}_6$ and $2\mathbb{Z}_6$. Thus, if $\chi : R \rightarrow \mathbb{C}, \chi(a) = \zeta^a$, where ζ is a primitive 6-th root of unity, then $H = [\chi(a_{ij})]$ is an amenable $BH(7, 6)$ matrix.

Example 8. All matrices of type $BH(n, p^k)$ are amenable BH matrices where p is a prime number and $k, n \in \mathbb{Z}^+$, because every row difference in the additive representation of a $BH(n, p^k)$ forms up a union of cosets of ideals $p^i \mathbb{Z}_{p^k}$ for some $1 \leq i \leq k - 1$, see [13, Theorem 3.3].

Remark 2. The authors of this article do not know the existence of a Butson-Hadamard matrix which is not amenable.

In Theorem 2 we have given lower bounds for the minimum distances of codes defined via rows of BH matrices. But in the following theorem we have obtained equality for these minimum distances by taking BH matrices as amenable.

Now suppose that w is a homogeneous weight, $l = \max\{w(a) : a \in \mathbb{Z}_k\}$, $l' = \min\{w(a) : a \in \mathbb{Z}_k \setminus \{0\}\}$ and γ is the average value of w .

Theorem 3. *Let n and k be positive integers, $H = [\zeta^{a_{ij}}]$ be a normalized amenable $BH(n, k)$ matrix, $H' = [a_{ij}]$ and N be the set of all k -th roots of unity. Also suppose that A_k, B_k, C_k and D_k are defined as in Theorem 2. Then $d_{A_k} = \gamma n, d_{B_k} = \gamma n - l, d_{C_k} = \gamma n$ and $d_{D_k} \in \{\gamma n, \gamma n + l'\}$.*

Proof. We know that $d_H = \gamma n$ by Theorem 1. Now consider B_k . Let R_i, R_j be the i -th and j -th rows of H , respectively. Also suppose that R'_j is the j -th row of $\zeta^u H$ such that $1 \leq i, j \leq n$ and $i \neq j, u \in \mathbb{Z}_t$. So $R'_j = \zeta^u R_j$. If we use the same argument in the proof of 1 we can say that $f_{r+u} = |\{k \in \{1, \dots, n\} : \log_\zeta(R_{ik} R_{jk}^{-1}) = r + u\}|/n = f_r$ for all $r, u \in \mathbb{Z}_t$. Here $\log_\zeta(R_{ik} R_{jk}^{-1})$ is the k -th coordinate of $R_i R_j^{-1}$. Therefore the proof of the Theorem 1 is also provided here as well. So $\sum_{k=1}^{k=n} (w(R_{ik} R_{jk}^{-1})) = \gamma n$. But here since the first coordinates of R_i and $R'_j = u R_j$ will always be different if we delete the first coordinate, the minimum distance decrease by l . After all this we get $d_{B_k} = \gamma n - l$.

The C_k code differs from B_k only in the first coordinate. Therefore, from the results we obtained above we get $d_{C_k} = \gamma n$.

Finally we consider D_k code. Since D_k is obtained by adding a column to C_k we can say that $d_{D_k} \in \{\gamma n, \gamma n + l'\}$. \square

As noted Example 8, all BH matrices in the following corollary are amenable.

Corollary 1. *Let $p > 2$ be a prime number, $k \in \mathbb{Z}^+$ such that $k \geq 2$ and ζ be a primitive p^k -th root of unity, $H = [\zeta^{a_{ij}}]$ be an $n \times n$ normalized Butson-Hadamard matrix and N be the set of all p^k -th roots of unity. Also let $H' = [a_{ij}]$ and define A_k, B_k, C_k, D_k codes as in Theorem 2. Then, $d_{G_1(A_k)} = n(p-1)p^{k-2}$, $d_{G_1(B_k)} = n(p-1)p^{k-2} - p^{k-1}$, $d_{G_1(C_k)} = n(p-1)p^{k-2}$, $d_{G_1(D_k)} \in \{p^{2k-2}(p-1), p^{k-2}(p^{k+1} - p^k + p - 1)\}$.*

Proof.

$d_{G_1(A_k)}$ is a consequence of Theorem 1. We need to find the γ average value of w . Now since G_1 is balanced except for the elements of $L = \{p^{k-1}, 2p^{k-1}, \dots, p^k\}$. Namely for every $1 \leq \alpha \leq p^k$ such that $\alpha \notin L$ there are equal number of $0, 1, \dots, p-1$ in $G_1(\alpha)$. For any $\beta \in L \setminus \{0\}$, $w(\beta) = p^{k-1}$. Therefore $l = p^{k-1} - p^{k-2}$, $l' = p^{k-1}$ and

$$\gamma = \frac{(p-1)p^{k-1} + (p^k - p)(p^{k-1} - p^{k-2})}{p^k} = p^{k-1} - p^{k-2} = (p-1)p^{k-2}.$$

So the result is obtained when the γ is written instead of the Theorem 3. \square

Now we will give an example of Theorem 3.

Example 9. Let $p = 3, k = 2$ and ζ be the primitive 9-th root of unity. Then

$$H = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & \zeta & \zeta^2 & \zeta^3 & \zeta^4 & \zeta^5 & \zeta^6 & \zeta^7 & \zeta^8 \\ 1 & \zeta^2 & \zeta^4 & \zeta^6 & \zeta^8 & \zeta & \zeta^3 & \zeta^5 & \zeta^7 \\ 1 & \zeta^3 & \zeta^6 & 1 & \zeta^3 & \zeta^6 & 1 & \zeta^3 & \zeta^6 \\ 1 & \zeta^4 & \zeta^8 & \zeta^3 & \zeta^7 & \zeta^2 & \zeta^6 & \zeta & \zeta^5 \\ 1 & \zeta^5 & \zeta & \zeta^6 & \zeta^2 & \zeta^7 & \zeta^3 & \zeta^8 & \zeta^4 \\ 1 & \zeta^6 & \zeta^3 & 1 & \zeta^6 & \zeta^3 & 1 & \zeta^6 & \zeta^3 \\ 1 & \zeta^7 & \zeta^5 & \zeta^3 & \zeta & \zeta^8 & \zeta^6 & \zeta^4 & \zeta^2 \\ 1 & \zeta^8 & \zeta^7 & \zeta^6 & \zeta^5 & \zeta^4 & \zeta^3 & \zeta^2 & \zeta \end{pmatrix}$$

is an amenable $BH(9, 9)$ matrix. Then

$$G_1(H') = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 2 & 0 & 2 & 1 & 1 & 1 & 1 & 1 & 2 & 0 & 1 & 0 & 2 & 2 & 2 & 2 & 0 & 1 & 2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 2 & 1 & 1 & 2 & 0 & 2 & 2 & 2 & 2 & 1 & 0 & 0 & 1 & 2 & 1 & 1 & 1 & 1 & 0 & 2 & 2 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 & 2 & 2 & 2 & 0 & 0 & 0 & 1 & 1 & 1 & 2 & 2 & 2 & 0 & 0 & 0 & 1 & 1 & 1 & 2 & 2 & 2 \\ 0 & 0 & 0 & 1 & 2 & 0 & 2 & 1 & 0 & 1 & 1 & 1 & 2 & 0 & 1 & 0 & 2 & 1 & 2 & 2 & 2 & 0 & 1 & 2 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 0 & 2 & 0 & 1 & 2 & 2 & 2 & 2 & 0 & 2 & 1 & 2 & 0 & 1 & 1 & 1 & 1 & 2 & 1 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 2 & 2 & 2 & 1 & 1 & 1 & 0 & 0 & 0 & 2 & 2 & 2 & 1 & 1 & 1 & 0 & 0 & 0 & 2 & 2 & 2 & 1 & 1 & 1 \\ 0 & 0 & 0 & 2 & 0 & 1 & 1 & 0 & 2 & 1 & 1 & 1 & 0 & 1 & 2 & 2 & 1 & 0 & 2 & 2 & 2 & 1 & 2 & 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 2 & 1 & 0 & 2 & 0 & 1 & 2 & 2 & 2 & 1 & 0 & 2 & 1 & 2 & 0 & 1 & 1 & 1 & 0 & 2 & 1 & 0 & 1 & 2 \end{pmatrix}.$$

If we calculate the average weight we get $\gamma = 2$. By using SageMath [19], we obtain that $d_{G_1(A_k)} = 18 = \gamma n$, $d_{G_1(B_k)} = 15 = \gamma n - l$, $d_{G_1(C_k)} = 18 = \gamma n$. Here $l = \max\{w(a) : a \in \mathbb{Z}_k \setminus \{0\}\} = 3$. Let c be the 4-th column of H' . Then we get $d_{G_1(D_k)} = 18$. These minimum distances comply with the results in Theorem 3.

If w is a homogeneous weight with the average value γ over a Frobenius ring R , then $\sum_{r \in y+I} w(r) = \gamma|I|$ for every nonzero ideal I of R , see [7, Proposition 2.6], a property which plays a crucial role in proving that some linear codes produced from a bimodule over R by using a non-degenerated bilinear form meet the Plotkin bound [7, Theorem 4.3]. Note that this important property of a homogeneous weight is shared also by the non-homogeneous weight w_2 induced by the Gray map G_2 with $\gamma = (p-1)p^{k-2}$, see Corollary 2 below. In the following definition we formalize such a weight.

Definition 6. A weight w on the finite ring R is called quasi-homogeneous if $w(0) = 0$ and there exists a real number γ such that

$$\sum_{r \in a+I} w(r) = \gamma|I|,$$

for each nonzero ideal I of R . The number γ is called the average value of w .

Proposition 2. Let k be a positive integer and γ be a positive real number. Define a mapping $w : \mathbb{Z}_{p^k} \rightarrow \mathbb{R}$ by

$$w(u) = \begin{cases} \frac{\gamma}{p^{k-2}(p-1)}u, & \text{if } u_{k-1} = 0 \\ \frac{\gamma p}{p-1}, & \text{if } 0 < u_{k-1} \leq p-2 \\ \frac{\gamma p^2}{p-1} - \frac{\gamma}{p^{k-2}(p-1)}u, & \text{if } u_{k-1} = p-1 \end{cases}$$

for all $u \in \mathbb{Z}_{p^k}$ with the p -ary expansion $u = u_0 + u_1p + \cdots + u_{k-1}p^{k-1}$. Then w is a quasi-homogeneous weight on \mathbb{Z}_{p^k} with the average value γ .

Proof. Fix $0 \leq s \leq k-1$. We show that for any $a \in \mathbb{Z}_{p^k}$,

$$\sum_{u \in a + \langle p^s \rangle} w(u) = \gamma p^{k-s},$$

which completes the proof. Note that it is enough to assume $0 \leq a \leq p^s - 1$. Note also that for any $u \in a + \langle p^s \rangle$, there exists a unique $n \in \mathbb{Z}_{p^k}$ such that $0 \leq n \leq p^{k-s} - 1$ and $u = a + np^s$. Write $a = a_0 + a_1p + \cdots + a_{s-1}p^{s-1}$ and $n = n_0 + n_1p + \cdots + n_{k-s-1}p^{k-s-1}$, where $0 \leq a_i, n_j \leq p-1$ for each $i = 0, \dots, s-1$ and $j = 0, \dots, k-s-1$. Hence, a typical element u of $a + \langle p^s \rangle$ can be written uniquely as

$$u = a_0 + a_1p + \cdots + a_{s-1}p^{s-1} + n_0p^s + n_1p^{s+1} + \cdots + n_{k-s-2}p^{k-2} + n_{k-s-1}p^{k-1},$$

where the a_i 's and n_j 's all lie in $\{0, \dots, p-1\}$. We separate the sum of $w(u)$'s, where u ranges over $a + \langle p^s \rangle$, into three sums as follows:

(I) The sum of the $w(u)$'s for $u \in a + \langle p^s \rangle$ with $n_{k-s-1} = 0$, i.e.,

$$\frac{\gamma}{p^{k-2}(p-1)} \left[p^{k-s-1}a + p^s \frac{p^{k-s-1}(p^{k-s-1}-1)}{2} \right];$$

(II) the sum of the $w(u)$'s for $u \in a + \langle p^s \rangle$ with $0 \leq n_{k-s-1} \leq p-2$, i.e.,

$$\frac{\gamma p(p-2)p^{k-s-1}}{p-1};$$

(III) the sum of the $w(u)$'s for $u \in a + \langle p^s \rangle$ with $n_{k-s-1} = p-1$, i.e.,

$$\frac{\gamma p^2 p^{k-s-1}}{p-1} - \frac{\gamma}{p^{k-2}(p-1)} \left[p^{k-s-1}a + \frac{p^s p^{k-s-1}(p^{k-s-1}-1)}{2} + p^{k-s-1}(p-1)p^{k-1} \right].$$

It is now easy to see that the numbers in (I)–(III) sum up to γp^{k-s} . \square

Corollary 2. *The weight w_2 on \mathbb{Z}_{p^k} , where $p > 2$ is a prime number and $k \in \mathbb{Z}^+$, is a quasi-homogeneous weight with the average value $\gamma = p^{k-2}(p-1)$.*

Proof. Notice that w_2 coincides with the weight defined in Proposition 2 with $\gamma = p^{k-2}(p-1)$. \square

Remark 3. In view of the proof of Proposition 1, we see that both weights w_1 and w_2 share the same average value $\gamma = (p-1)p^{k-2}$. In fact, w_1 is the only homogeneous weight on \mathbb{Z}_{p^k} with the average value $\gamma = (p-1)p^{k-2}$, see Theorem 2.2 in [7]. It follows that our consideration of quasi-homogeneous weights enables us to work with more useful weights with the same common average value and flourishes our repository of weights in this manner.

Corollary 3. *Let $p > 2$ be a prime number, k be a positive integer, and $i \in \{1, 2, \dots, k-1\}$. Given any $j \in \mathbb{Z}_{p^k}$, the total number of 0's in the elements of the set $\{G_2(u) : u \in j + \langle p^i \rangle\}$ is independent of j and equals p^{2k-i-2} .*

Proof. By Corollary 2, the total number of 0's in the elements of the set $\{G_2(u) : u \in j + \langle p^i \rangle\}$ is

$$\begin{aligned} p^{k-1}p^{k-i} - \sum_{u \in j + \langle p^i \rangle} w_2(u) &= p^{2k-i-1} - p^{k-2}(p-1)p^{k-i} \\ &= p^{2k-i-2} \end{aligned}$$

\square

Example 10. Choose $p = 5$, $k = 2$, $i = 1$ and $j = 3$. The total number of 0's in $G_2(5) = (11111)$, $G_2(10) = (22222)$, $G_2(15) = (33333)$, $G_2(20) = (44444)$, $G_2(25) = (00000)$ is $p^{2k-i-2} = 5$. On the other hand, the total number of 0's in $G_2(8) = (22211)$, $G_2(13) = (33322)$, $G_2(18) = (44433)$, $G_2(23) = (00044)$, $G_2(3) = (11100)$ is $p^{2k-i-2} = 5$. Hence, the total number of 0's in both sets is the same.

Proposition 3. *Let $p > 2$ be a prime number, $k \in \mathbb{Z}^+$ such that $k \geq 2$ and ζ be a primitive p^k -th root of unity, $H = [\zeta^{a_{ij}}]$ be an $n \times n$ normalized Butson-Hadamard matrix, $H' = [a_{ij}]$ and w' be a quasi-homogeneous weight with the average value γ . Then the minimum distance of the code consisting the rows of H' with the first column deleted is equal to γn .*

Proof. Since H is a Butson-Hadamard matrix we know that there exists $m \in \mathbb{Z}^+$ such that $n = mp$. We will the proof by induction on m .

For $m = 1$ we have a $BH(p, p)$ matrix. Let R_i, R_j be the i -th and j -th rows of H' , respectively. Hence the elements of $R_i - R_j$ consists of the elements of $p^{k-1}\mathbb{Z}_{p^k}$ or the cosets of this ideal. More clearly $R_i - R_j = \{p^{k-1}, 2p^{k-1}, \dots, p^k\}$ or $\{p^{k-1} + i, 2p^{k-1} + i, \dots, p^k + i\}$ for all $1 \leq i \leq p^{k-1} - 1$. So for $I = p^{k-1}\mathbb{Z}_{p^k}$ we can say that $\sum_{r \in I+y} w'(r) = \gamma p = \gamma n$. Now let $m > 1$ be given and suppose

that the theorem is true for all $m = l$ with $l \in \mathbb{Z}^+$. Let us show the theorem is true for $m = l + 1$. Suppose that $m = l + 1$. We know that $d = \gamma n$ for $n = l$. If we take $n = l + 1$ this differs from the case $n = l$ only p elements. The p elements again only come from the $p^{k-1}\mathbb{Z}_{p^k}$ ideal or the cosets of this ideal. Therefore $d = \gamma lp + \gamma p = \gamma p(l + 1) = \gamma n$. This ends the proof. \square

Theorem 4. *Let $p > 2$ be a prime number, $k \in \mathbb{Z}^+$ such that $k \geq 2$ and ζ be a primitive p^k -th root of unity, $H = [\zeta^{a_{ij}}]$ be an $n \times n$ normalized Butson-Hadamard matrix and N be the set of all p^k -th roots of unity. Also let $H' = [a_{ij}]$ and define A_k, B_k, C_k, D_k codes as in Theorem 2. Then,*

$$d_{G_2(A_k)} = n(p-1)p^{k-2}, \quad d_{G_2(B_k)} = n-1, \quad d_{G_2(C_k)} = n, \quad d_{G_2(D_k)} \in \{n, n+1\}.$$

Proof. We can say that $d_{G_2(A_k)} = \gamma n$ by Proposition 3 and Corollary 2 we get $d_{G_2(A_k)} = n(p-1)p^{k-2}$.

Now consider $G_2(B_k)$. Let $R_i = [a_{i2} \dots a_{in}]$ and $R'_i = [a'_{i2} \dots a'_{in}]$ be the i -th row of the exponent matrix of H and ζH with deleted first column, respectively. Then $|a_{ij} - a'_{ij}| = 1$, so we get $d(G_2(a_{ij}), G_2(a'_{ij})) = 1$ for $j = 2, 3, \dots, n$ by Lemma 2. Therefore the distance between $G_2(R_i)$ and $G_2(R'_i)$ is equal to $n-1$. Now suppose that R'_j be the j -th row of the exponent matrix of uH for an arbitrary $u \in N$. By Corollary 3 and Lemma 3, we can say

$$d(G_2(R_i), G_2(R'_j)) = (n-1)p^{k-1} - np^{k-2} = p^{k-2}(n(p-1) - 1)$$

for $i \neq j$. Since $n-1 < p^{k-2}(n(p-1) - 1)$, the minimum distance of $G_2(B_k)$ is $d_{G_2(B_k)} = n-1$.

We next consider $G_2(C_k)$. Its codewords are constructed from the rows of the exponent matrix of uH for $u \in N$. Similar to (ii), by considering i -th rows R_i and R'_i of H and ζH respectively, we have $d(G_2(R_i), G_2(R'_i)) = n$. And so, $d_{G_2(C_k)} = n$.

Finally the code $G_2(D_k)$ is the Gray image of the rows of uH concatenated with a non-initial column c of H . So, by using (iii), we can say that $d_{G_2(D_k)}$ is either n or $n+1$. \square

Now we can directly obtain a Plotkin optimal code form G_2 image of A_k in the following corollary.

References

1. Armario, J.A., Bailera, I., Borges, J., Rifà, J.: Quasi-Hadamard full propelinear codes. *Mathematics in Computer Science* **12**(4), 419–428 (2018)
2. Armario, J.A., Bailera, I., Egan, R.: Generalized Hadamard full propelinear codes. arXiv preprint arXiv:1906.06220 (2019)
3. Butson, A.: Generalized Hadamard matrices. *Proceedings of the American Mathematical Society* **13**(6), 894–898 (1962)
4. Dougherty, S.T., Rifà, J., Villanueva, M.: Ranks and kernels of codes from generalized Hadamard matrices. *IEEE Transactions on Information Theory* **62**(2), 687–694 (2015)
5. Dougherty, S.T., Rifà, J., Villanueva, M.: Rank and kernel of \mathbb{F}_p -additive generalised Hadamard codes. arXiv preprint arXiv:2001.11609 (2020)
6. Drake, A.D.: Partial λ -geometries and generalized Hadamard matrices over groups. *Can. J. Math.* **XXXI**, No. 3, 617–627 (1979)
7. Greferath, M., McGuire, G., O’Sullivan, M.E.: On Plotkin-optimal codes over finite Frobenius rings. *Journal of Algebra and its Applications* **5**(06), 799–815 (2006)
8. Hadamard, J.: Résolution d’une question relative aux déterminants. *Bull. des sciences math.* **2**, 240–246 (1893)
9. Hedayat, A., Wallis, W.D.: Hadamard matrices and their applications. *The Annals of Statistics* **6**(6), 1184–1238 (1978)
10. Heng, Z., Yue, Q.: Generalized Gray map and a class of p -ary nonlinear codes. *Finite Fields and Their Applications* **36**, 81–97 (2015)
11. Horadam, K.J.: *Hadamard Matrices and Their Applications*. Princeton University Press (2007)
12. Jungnickel, D.: On difference matrices, resolvable transversal designs and generalized Hadamard matrices. *Mathematische Zeitschrift* **167**(1), 49–60 (1979)
13. Lam, T.Y., Leung, K.H.: On vanishing sums of roots of unity. *Journal of algebra* **224**(1), 91–109 (2000)
14. Lee, M., Jiang, X., Hwang, G., Pokhrel, S.S.: Error correcting codes from modified butson-hadamard matrices. In: 2006 International Conference on Systems and Networks Communications, pp. 65–65. IEEE Computer Society (2006)
15. McGuire, G., Ward, H.N.: Cocyclic hadamard matrices from forms over finite Frobenius rings. *Linear algebra and its applications* **430**(7), 1730–1738 (2009)
16. Seberry, J.: *Orthogonal designs*. Springer, Cham (2017)
17. Stepanov, S.A.: A new class of quaternary codes. *Problems of Information Transmission* **42**(1), 1–9 (2006)
18. Stepanov, S.A.: Nonlinear q -ary codes with large code distance. *Problems of Information Transmission* **53**(3), 242–250 (2017)
19. The Sage Developers: SageMath, the Sage Mathematics Software System (Version 8.2.0) (2020). <https://www.sagemath.org>
20. Yildiz, B., Ozger, Z.O.: Generalization of the Lee weight to \mathbb{Z}_{p^k} . *TWMS Journal of Applied and Engineering Mathematics* **2**(2), 145 (2012)