

Interacting Bose gas across a narrow Feshbach resonance

Fan Yang and Ran Qi*

Department of Physics, Renmin University of China, Beijing, 100872, P. R. China

We investigate the ground state properties of a weakly interacting Bose gas across a narrow Feshbach resonance. Strong deviation from a wide Feshbach resonance were discovered in various thermodynamic quantities. The energy per particle was found to saturate as the density increases which leads to significant reduction of the inverse compressibility. We also show that these effects can be observed in the density profiles in harmonic traps.

I. INTRODUCTION

The studies of weakly interacting Bose gas have a very long history ever since the pioneer works in early 20th century. A cornerstone progress has been made by Lee, Huang and Yang in their famous paper [1]. In this work, a low density expansion was obtained for the equation of state at zero temperature:

$$\frac{E}{N} = \frac{2\pi\hbar^2 a_s}{m} n \left(1 + \frac{128}{15\sqrt{\pi}} \sqrt{na_s^3} \right), \quad (1)$$

where E/N is the energy per particle of the ground state, n is total density of the Bose gas and a_s is the s-wave scattering length [1]. Based on Eq. (1), all thermal dynamic quantities can be derived at zero temperature. For example, the chemical potential and compressibility are given as:

$$\mu = \left(\frac{\partial E}{\partial N} \right)_V = \frac{4\pi\hbar^2 a_s}{m} n \left(1 + \frac{32}{3\sqrt{\pi}} \sqrt{na_s^3} \right), \quad (2)$$

$$\kappa^{-1} = \left(\frac{\partial \mu}{\partial n} \right)_N = \frac{4\pi\hbar^2 a_s}{m} \left(1 + \frac{16}{\sqrt{\pi}} \sqrt{na_s^3} \right). \quad (3)$$

Although these results was published more than 60 years ago, its experimental verification is only made possible very recently, thanks to the realization and high precision measurement of Bose Einstein condensation in ultracold quantum gases [2–4].

However, one should note that the validity of expansion (1) does not only require the low density condition $n^{1/3}a_s \ll 1$, but also relies on the fact that the two-body scattering process is fully determined by a single parameter a_s . For quantum gases close to a Feshbach resonance, while the second condition is usually full filled for a wide Feshbach resonance, it may break down for a very narrow resonance. As shown in several previous works, for a Feshbach resonance the two-body scattering amplitude is generally determined by an energy dependent scattering length $a_s(E) = a_{bg} \left(1 + \frac{\alpha_r^2}{g_r(E-\nu_r)} \right)$ where E is the total energy of the two particle under collision in the center of mass frame [5–8]. Across a wide resonance, the energy dependence in $a_s(E)$ can be safely neglected and

one has $a_s(E) \simeq a_s(0)$. In contrast, for a narrow resonance $a_s(E)$ has very sensitive dependence on E and thus the full functional form must be taken into account [6–8].

In particular, for a bose gas across a narrow Feshbach resonance, one can qualitatively estimate the effect of energy dependent scattering length as following. Since the chemical potential μ is the lowest energy to excite a single particle out of the condensate, the typical two-body scattering energy can be estimate as $E \sim 2\mu$. As a result, the effective interacting strength should be determined by $a_s(2\mu)$ instead of the zero energy scattering length $a_s(0)$. In this paper, we consider a particular case $\nu_r > 0, a_{bg} > 0$ and a schematic plot of $a_s(E)$ is shown in Fig. 1. In this case, the function $a_s(E)$ decreases from $a_s(0) > 0$ to 0 as energy E increases from 0 to a critical value E_c . At fixed magnetic field if one increases the density, the chemical potential μ will increase according to Eq. (2) such that $a_s(2\mu)$ decrease which tends to prevent μ from further increasing. As a result, we expect that μ may saturate to $E_c/2$ as the density increases which leads to a nearly zero inverse compressibility. In the following, we will show that above qualitative analysis is correct based on a meanfield plus Gaussian fluctuation calculation. We will provide a modified low density expansion for the equation of state which can be seen as a generalization of Eq. (1) that applies to bose gases across either wide or narrow Feshbach resonances. The effect of energy dependent scattering length on varies thermodynamic properties will be investigated systematically.




FIG. 1. energy-dependent scattering length

Our paper is organized as follows. In Sec.II, we introduce our model for a spinless bose gas across narrow Feshbach resonance and adopt the path integral method approach to obtain the thermodynamic partition func-

* qiran@ruc.edu.cn

tion [10]. In Sec.III we present mean-field results on thermodynamic properties in detail and show the effect of energy dependent scattering length on both uniform and trapped systems. In Sec.IV, we calculate the Gaussian fluctuation correction to the results obtained in Sec.III and show that including Gaussian fluctuation does not qualitatively change the meanfield results. In Sec.V, we summarize our main findings and conclude the paper.

II. MODEL AND PATH INTEGRAL APPROACH

For boson atoms interacting across a magnetic feshbach resonance, we adopt the widely used two-channel model:

$$\hat{H} = \hat{H}_a + \hat{H}_m + \hat{H}_c, \quad (4)$$

$$\hat{H}_a = \int d\mathbf{x} \left[\psi^\dagger(\mathbf{x}) \left(-\frac{\nabla^2}{2m} + V_a(\mathbf{x}) \right) \psi(\mathbf{x}) + \frac{g}{2} \psi^\dagger(\mathbf{x}) \psi^\dagger(\mathbf{x}) \psi(\mathbf{x}) \psi(\mathbf{x}) \right], \quad (5)$$

$$\hat{H}_b = \int d\mathbf{x} b^\dagger(\mathbf{x}) \left(-\frac{\nabla^2}{2M} + \nu_b + V_b(\mathbf{x}) \right) b(\mathbf{x}), \quad (6)$$

$$\hat{H}_c = \int d\mathbf{x} \left[\frac{\alpha}{\sqrt{2}} \psi^\dagger(\mathbf{x}) \psi^\dagger(\mathbf{x}) b(\mathbf{x}) + h.c. \right], \quad (7)$$

where ψ^\dagger and b^\dagger are creation field operators for atom and molecule, respectively. V_a and V_b are extra potential of atom and molecule. The molecule detuning ν_b , the inter-channel coupling α and the interaction parameter in open channel g are bare quantities which need to be renormalized as follows ($\hbar = 1$):

$$\begin{aligned} \nu_b &= \nu_r - [1 - Z(\Lambda)] \alpha_r^2 / g_r \\ \alpha &= Z(\Lambda) \alpha_r \\ g &= Z(\Lambda) g_r \end{aligned} \quad (8)$$

where $Z(\Lambda) = (1 - g_r \Lambda)^{-1}$ and $\Lambda = \frac{1}{V} \sum_{\mathbf{k}}' (2\varepsilon_{\mathbf{k}})^{-1}$. The renormalized quantities g_r , α_r and ν_r are related to a_s as

$$\frac{4\pi a_s(E)}{m} = \left[\left(g + \frac{\alpha^2}{E - \nu_b} \right)^{-1} + \Lambda \right]^{-1} = g_r + \frac{\alpha_r^2}{E - \nu_r} \quad (9)$$

The partition function of Hamiltonian (4) at arbitrary temperature can be written into the following imaginary time path integral form

$$Z = \int D[\phi^*] D[\phi] D[\beta^*] D[\beta] \exp(-S), \quad (10)$$

where

$$S = \int dx \left[\phi^*(x) \frac{\partial}{\partial \tau} \phi(x) + \beta^*(x) \frac{\partial}{\partial \tau} \beta(x) + h_a(x) + h_b(x) + h_c(x) \right], \quad (11)$$

$$h_a(x) = \phi^*(x) \left[-\frac{\nabla^2}{2m} + V_a(\mathbf{x}) - \mu \right] \phi(x) + \frac{g}{2} \phi^*(x) \phi^*(x) \phi(x) \phi(x), \quad (12)$$

$$h_b(x) = \beta^*(x) \left[-\frac{\nabla^2}{4m} + V_b(\mathbf{x}) - 2\mu \right] \beta(x),$$

$$h_c(x) = \frac{\alpha}{\sqrt{2}} \phi^*(x) \phi^*(x) \beta(x) + h.c.,$$

and we have defined $x \equiv (\mathbf{x}, \tau)$ and $\int dx \equiv \int d\mathbf{x} \int_0^\beta d\tau$ with $\beta = (k_B T)^{-1}$.

The path integral (10) can not be performed exactly due to the interaction terms. However, for a weakly interacting bose gas with very low density, it is a good approximation to expand the action S in Eq. (11) around its saddle point solution [10]:

$$\phi(x) = \phi_0(\mathbf{x}) + \phi'(x), \quad \beta(x) = \beta_0(\mathbf{x}) + \beta'(x) \quad (13)$$

where $\phi_0(\mathbf{x})$ and $\beta_0(\mathbf{x})$ are saddle point solutions which minimizes the action S , $\phi'(x)$ and $\beta'(x)$ are fluctuation field of atom and molecule.

Later in Sec.III, we will take mean-field approximation by neglecting all the fluctuation terms. The approximate action is then given by

$$\begin{aligned} S^{(0)} = & \beta \int d\mathbf{x} \left\{ \phi_0^*(\mathbf{x}) \left[-\frac{\nabla^2}{2m} + V_a(\mathbf{x}) - \mu \right] \phi_0(\mathbf{x}) \right. \\ & + \frac{g}{2} |\phi_0(\mathbf{x})|^4 + \beta_0^*(\mathbf{x}) \left[-\frac{\nabla^2}{4m} + V_b(\mathbf{x}) - 2\mu \right] \beta_0(\mathbf{x}) \\ & \left. + \frac{\alpha}{\sqrt{2}} \phi_0^*(\mathbf{x})^2 \beta_0(\mathbf{x}) + \frac{\alpha^*}{\sqrt{2}} \beta_0^*(\mathbf{x}) (\phi_0(\mathbf{x}))^2 \right\} \end{aligned} \quad (14)$$

Then in Sec.IV we will include the contribution from Gaussian fluctuation around the saddle point. For a uniform system, the saddle point solution is uniform, i.e. $\phi_0(\mathbf{x}) \equiv \phi_0$, $\beta_0(\mathbf{x}) \equiv \beta_0$, and we obtain the quadratic action as

$$S \approx S^{(0)} + S^{(2)} \quad (15)$$

$$\begin{aligned} S^{(2)} = & \int dx \phi'^*(x) \left[\frac{\partial}{\partial \tau} - \frac{\nabla^2}{2m} - \mu + 2g|\phi_0|^2 \right] \phi'(x) \\ & + \frac{1}{2} \int dx \left\{ \phi'^*(x)^2 \left[g\phi_0^2 + \frac{\alpha}{\sqrt{2}} \beta_0 \right] + c.c. \right\} \\ & + \int dx \beta'^*(x) \left[\frac{\partial}{\partial \tau} - \frac{\nabla^2}{4m} - 2\mu + \nu_b \right] \beta'(x) \\ & + \int dx \left[2 \frac{\alpha}{\sqrt{2}} \phi'^*(x) \phi_0^* \beta'(x) + c.c. \right] \end{aligned} \quad (16)$$

To perform the path integral for this quadratic action, it is more convenient to first transform the action into momentum-frequency space,

$$\begin{aligned}\phi'(x) &= \frac{1}{\sqrt{\beta V}} \sum_{k,n} \phi'_{\mathbf{k},n} \exp[i(\mathbf{k} \cdot \mathbf{x} - \omega_n \tau)] \\ \beta'(x) &= \frac{1}{\sqrt{\beta V}} \sum_{k,n} \beta'_{\mathbf{k},n} \exp[i(\mathbf{k} \cdot \mathbf{x} - \omega_n \tau)]\end{aligned}\quad (17)$$

where ω_n is bosonic Matsubara frequency, \mathbf{k} is free wave vector. And the action can be written more compactly as a matrix multiplication in Nambu space

$$\begin{aligned}S^{(2)} &= -\frac{\beta}{2} \sum_{\mathbf{k} \neq 0} (\epsilon_{\mathbf{k}}^b + \epsilon_{\mathbf{k}}^a + 2g|\phi_0|^2) \\ &\quad - \frac{1}{2} \sum_{\mathbf{k} \neq 0, n} \Phi_{\mathbf{k}}^\dagger \mathbf{G}^{-1}(\mathbf{k}, i\omega_n) \Phi_{\mathbf{k}}\end{aligned}\quad (18)$$

where $\epsilon_{\mathbf{k}}^b = \frac{k^2}{4m} + \nu_b - 2\mu$, $\epsilon_{\mathbf{k}}^a = \frac{k^2}{2m} - \mu$. We write as $k \equiv (\mathbf{k}, i\omega_n)$ and

$$\Phi_{\mathbf{k}}^\dagger = [\beta_{\mathbf{k}}'^*, \phi_{\mathbf{k}}'^*, \phi'_{-\mathbf{k}}, \beta'_{-\mathbf{k}}] \quad (19)$$

The first sum term in Eq.(18) comes from the order exchanging between the creation and annihilation field during the transformation into Nambu space [10]. And the 4×4 matrix \mathbf{G}^{-1} gives the interacting inverse Green's function

$$\mathbf{G}^{-1} = \mathbf{G}_0^{-1} - \Sigma \quad (20)$$

$$\mathbf{G}_0^{-1} = \begin{bmatrix} G_{b0}^{-1}(k) & & & \\ & G_{a0}^{-1}(k) & & \\ & & G_{a0}^{-1}(-k) & \\ & & & G_{b0}^{-1}(-k) \end{bmatrix} \quad (21)$$

$$\Sigma = \begin{bmatrix} 0 & \tilde{\alpha}^* & & \\ \tilde{\alpha} & 2g|\phi_0|^2 & \tilde{g} & \\ & \tilde{g}^* & 2g|\phi_0|^2 & \tilde{\alpha}^* \\ & & \tilde{\alpha} & 0 \end{bmatrix} \quad (22)$$

where $G_{a0}(k) = (i\omega_n - \epsilon_{\mathbf{k}}^a)^{-1}$, $G_{b0}(k) = (i\omega_n - \epsilon_{\mathbf{k}}^b)^{-1}$, $\tilde{\alpha} = \sqrt{2}\alpha\phi_0^*$ and $\tilde{g} = g\phi_0^2 + \sqrt{2}\alpha\beta_0$

It is known that the quasi-particle excitation corresponds to the poles of the Green's function. Here we obtain two branches of excitations

$$\begin{aligned}\omega_{\mathbf{k}}^\pm &= \sqrt{(B \pm \sqrt{B^2 - 4C})} / 2 \\ B &= (\epsilon_{\mathbf{k}}^a + 2g|\phi_0|^2)^2 + (\epsilon_{\mathbf{k}}^b)^2 + 2|\tilde{\alpha}|^2 - \mu^2 \\ C &= [(\epsilon_{\mathbf{k}}^a + 2g|\phi_0|^2)\epsilon_{\mathbf{k}}^b - |\tilde{\alpha}|^2]^2 - \mu^2 (\epsilon_{\mathbf{k}}^b)^2\end{aligned}\quad (23)$$

which also means the Green's function can be diagonalized into [9]

$$-\mathbf{U}^T \mathbf{G}^{-1}(\mathbf{k}, i\omega_n) \mathbf{U} = \begin{bmatrix} -i\omega_n + \omega_{\mathbf{k}}^+ & & & \\ & -i\omega_n + \omega_{\mathbf{k}}^- & & \\ & & i\omega_n + \omega_{\mathbf{k}}^- & \\ & & & i\omega_n + \omega_{\mathbf{k}}^+ \end{bmatrix}. \quad (24)$$

Finally, we obtain the diagonalized Gaussian action

$$\begin{aligned}S &= S^{(0)} + \frac{\beta}{2} \sum_{\mathbf{k}} (\omega_{\mathbf{k}}^+ + \omega_{\mathbf{k}}^-) \\ &\quad - \frac{\beta}{2} \sum_{\mathbf{k}} (\epsilon_{\mathbf{k}}^a + 2g|\phi_0|^2 + \epsilon_{\mathbf{k}}^b) \\ &\quad + \sum_{\mathbf{k}} (-i\omega_n + \omega_{\mathbf{k}}^+) \beta_{\mathbf{k},n}^* \beta_{\mathbf{k},n} \\ &\quad + \sum_{\mathbf{k}} (-i\omega_n + \omega_{\mathbf{k}}^-) \phi_{\mathbf{k},n}^* \phi_{\mathbf{k},n}\end{aligned}\quad (25)$$

where the sum of $\omega_{\mathbf{k}}^+$ and $\omega_{\mathbf{k}}^-$ comes from the exchanging of the field again [10].

In the following sections we calculate the thermodynamic potential Ω and density n with action given by Eq.(14) and Eq.(25) through following thermodynamic relations

$$\frac{\Omega}{V} = -\frac{1}{\beta V} \ln(Z) \quad (26)$$

$$n = -\frac{\partial(\Omega/V)}{\partial\mu} \quad (27)$$

III. MEAN-FIELD CALCULATION

In this section we neglect the contribution of Gaussian fluctuation, and the action is governed by Eq.(14). After minimizing the action, we obtain the two-channel G-P equations

$$\begin{aligned}\left[-\frac{\nabla^2}{2m} + V_a + g_r|\phi_0(\mathbf{x})|^2 - \mu\right] \phi_0(\mathbf{x}) + \sqrt{2}\alpha_r\phi_0^*(\mathbf{x})\beta_0(\mathbf{x}) &= 0 \\ \left[-\frac{\nabla^2}{4m} + \nu_r + V_b - 2\mu\right] \beta_0(\mathbf{x}) + \frac{\alpha_r^*}{\sqrt{2}}\phi_0(\mathbf{x})^2 &= 0\end{aligned}\quad (28)$$

The renormalization should be taken to the same order, so that in mean-field level we should use the renormalized parameters directly in Eq.(28) [10].

Uniform system. We first consider a uniform system, and thus the solution of Eq. (28) for ground state are constants ϕ_0 and β_0 . Assuming all the atoms and molecules condense on the zero momentum state in mean-field approximation, the atomic density n_a and molecular den-

sity n_b are given as

$$\begin{aligned} n_a &= |\phi_0|^2 = \frac{\mu}{g(2\mu)} \\ n_b &= |\beta_0|^2 = \frac{1}{2} \left[\frac{\mu}{(2\mu - \nu_r)g(2\mu)} \right]^2 \cdot |\alpha_r|^2 \end{aligned} \quad (29)$$

where $g(2\mu) = 4\pi\hbar^2 a_s(2\mu)/m$. One can see that n_a and n_b diverge when effective interaction strength $g(2\mu)$ tends to zero. Since $g(2\mu) = 0$ at $\mu_c = \nu_r - |\alpha|^2/g_r$, the chemical potential will saturate to μ_c as the density increases.




FIG. 2. (a):The chemical potential as a function of total density n . (b):The proportion of molecules as a function of total density n . $\tilde{\Delta} = |\alpha_r|^2/g_r\nu_r = 1/100$

Substituting Eq.(29) into mean-field action(14), we can easily obtain the energy per particle

$$\frac{E}{N} = \frac{1}{n^0} \frac{F_L}{V} + \mu = \frac{\mu}{2} \left(1 + \frac{2n_b}{n^0} \right) \quad (30)$$

where F_L and n^0 are thermodynamic potential and total density given by Eq.(26) and Eq.(27) with $S = S^{(0)}$.

We find that as the density increases, the energy per particle saturates to μ_c while in Eq. (1) it increases monotonically. The difference can be attributed to the energy dependence of scattering length. As the average scattering energy increase to the zero point of a_s , the system turns from a finite repulsive interacting gas into a nearly noninteracting gas, and the atoms in open channel are in favour of converting into the closed channel molecules of which the typical energy is the detuning. This energy dependent effect can also change the behavior of compressibility κ . Here, as the density increases, κ will diverge at $g(2\mu) = 0$ which means the gas becomes more and more soft.

$$\kappa = \left(\frac{\partial n}{\partial \mu} \right)_N = \frac{1}{\mu} \left(n_a + 4n_b + 4n_b \frac{\nu_r - \frac{\alpha_r^2}{g_r}}{\nu_r - \frac{\alpha_r^2}{g_r} - 2\mu} \right) \quad (31)$$

Density profile in harmonic trap. Now we consider the system in a harmonic trap with $V_b(r)/2 = V_a(r) = \frac{1}{2}m\omega^2 r^2$. For a large particle number in the trap, the kinetic energy is much smaller than other energy so that can be neglected [11]. The Thomas-Fermi distribution is




FIG. 3. (a):The energy per particle as a function of total density n . (b):The inverse compressibility κ^{-1} as a function of total density n . $\tilde{\Delta} = |\alpha_r|^2/g_r\nu_r = 1/100$. The Blue line and yellow line are narrow resonance and broad resonance respectively.

given by Eq.(29) while μ replaced with $\mu - V(r)$ and the result is shown in Fig.4(a).

We know for interacting Bose gas across a broad Feshbach resonance, the distribution in the harmonic trap is approximately governed by the balance between interaction energy and the trap potential, and Thomas-Fermi approximation gives the inverted parabola distribution [11]. Here, the distribution behaves as broad resonance Bose gas at the edge of the trap while more particles can be accommodated in the center where the local chemical potential is larger such that the energy dependence in a_s is more significant. A sharp peak appears if the effective local chemical potential $\mu - V(r) = (\nu_r - |\alpha_r|^2/g_r)2$. This divergency can also be interpreted by the calculation of the compressibility.

IV. GAUSSIAN FLUCTUATION CALCULATION

In this section, we take the correction of Gaussian fluctuation into account, and the action is governed by Eq.(25). The field integral of the action gives the thermodynamic potential

$$\begin{aligned} \frac{\Omega}{V} &= \frac{\Omega_0}{V} + \frac{1}{V\beta} \sum_{\mathbf{k}} \ln \left(1 - e^{-\beta\omega_{\mathbf{k}}^-} \right) \\ &+ \frac{1}{V\beta} \sum_{\mathbf{k}} \ln \left(1 - e^{-\beta\omega_{\mathbf{k}}^+} \right) \end{aligned} \quad (32)$$




FIG. 4. Distribution $n a_s^3(E = 0)$ of a Bose gas in a harmonic trap with fixed particle number $N = 2 \times 10^5$. The resonance width $\tilde{\Delta} = |\alpha_r|^2/g_r \nu_r = 1/1000$ and $a_o = \sqrt{m\omega/\hbar} = 60a_s(E = 0)$. **(a)**:Distribution in mean-field approximation for gas across narrow Feshbach resonance(blue line) and broad Feshbach resonance(green line). **(b)**:Distribution of gas across broad Feshbach resonance with(blue line) and without(green line) Gaussian fluctuation. **(c)**:Distribution of gas across narrow Feshbach resonance with(blue line) and without(green line) Gaussian fluctuation.

The ground state thermodynamic potential Ω_0 at zero temperature is given by

$$\frac{\Omega_0}{V} = \frac{F_L}{V} + \frac{F_{fluc}}{V} \quad (33)$$

$$\frac{F_L}{V} = -\frac{\mu^2}{2} \frac{1}{g_r + \frac{|\alpha_r|^2}{2\mu - \nu_r}} \quad (34)$$

$$\begin{aligned} \frac{F_{fluc}}{V} &= \frac{1}{2V} \sum_{\mathbf{k}} \left[(\omega_{\mathbf{k}}^- + \omega_{\mathbf{k}}^+) \right. \\ &\quad \left. - (\epsilon_{\mathbf{k}}^a + 2g_r |\phi_0|^2 + \epsilon_{\mathbf{k}}^b) + \frac{\mu^2}{2\epsilon_{\mathbf{k}}} \right] \\ &= \frac{1}{4\pi^2} (2m)^{3/2} \mu^{5/2} F(\tilde{\mu}, \tilde{\Delta}) \end{aligned} \quad (35)$$

where $\tilde{\mu} = \frac{\mu}{\nu_r}$, $\tilde{\Delta} = \frac{\Delta}{g_r}$ and $\Delta = \frac{|\alpha_r|^2}{g_r}$. F_L is the mean-field thermodynamic potential that has been renormalized with complete renormalization relation(8), and F_{fluc} is the correction by Gaussian fluctuation in which the parameters can be approximately replaced by the renormalized ones. The dimensionless function F is defined as

$$F(\tilde{\mu}, \tilde{\Delta}) = \int_0^\infty \left[\frac{1}{2} + x^2 g(x, \tilde{\mu}, \tilde{\Delta}) \right] dx \quad (36)$$

Here we defined another dimensionless function,

$$\begin{aligned} g(x, \tilde{\mu}, \tilde{\Delta}) &= \sqrt{\omega_a^2 + \omega_b^2 + 4\xi - 1 + 2\sqrt{(\omega_a \omega_b - 2\xi)^2 - \omega_b^2}} \\ &\quad - \omega_a - \omega_b \end{aligned} \quad (37)$$

with

$$\begin{aligned} \omega_a &= x^2 + 1 - 2 \frac{\tilde{\Delta}}{2\tilde{\mu} - 1 + \tilde{\Delta}} \\ \omega_b &= \frac{x^2}{2} + \frac{1}{\tilde{\mu}} - 2 \\ \xi &= \frac{\tilde{\Delta}}{\tilde{\mu}} \frac{2\tilde{\mu} - 1}{2\tilde{\mu} - 1 + \tilde{\Delta}} \end{aligned} \quad (38)$$

If chemical potential μ is very small where the scattering is far from resonance or the resonance width Δ is very small while μ is less than μ_{\max} , the system has little energy dependence and the fluctuation correction is close to the broad resonance case with $F(0,0) \rightarrow \frac{8\sqrt{2}}{15}$. As Δ or μ increases, $F(\tilde{\mu}, \tilde{\Delta})$ increases monotonically.

Below, we analyze the asymptotic behavior of $F(\tilde{\mu}, \tilde{\Delta})$ in three different limit.

In the limit $\tilde{\Delta} \ll 1$ which called narrow Resonance Limit, we have

$$F(\tilde{\mu}, \tilde{\Delta}) = \frac{8\sqrt{2}}{15} + \frac{G(\tilde{\mu})}{1 - 2\tilde{\mu}} \tilde{\Delta} + O(\tilde{\Delta}^2) \quad (39)$$

$$G(\tilde{\mu}) = \int_0^\infty \eta(x, \tilde{\mu}) dx \quad (40)$$

where

$$\begin{aligned} \eta(x, \tilde{\mu}) &= \frac{1}{\sqrt{(x^2 + 1)^2 - 1 + (x^2 + 1)}} \\ &\quad \frac{2x^2}{\sqrt{(x^2 + 1)^2 - 1 + \frac{x^2}{2} - 2 + \frac{1}{\tilde{\mu}}}} \\ &\quad \left(1 + \frac{x^2}{2\sqrt{(x^2 + 1)^2 - 1}} \right) \end{aligned} \quad (41)$$

If $\tilde{\mu}$ also approaches 0 in this case, then we have

$$G(\tilde{\mu}) \frac{\sqrt{6}\pi}{4} \sqrt{\tilde{\mu}} + O(\tilde{\mu}) \quad (42)$$

In the limit $\tilde{\mu} \ll 1$ while $\tilde{\Delta}$ remains finite, which called wide resonance limit, we have

$$F(\tilde{\mu}, \tilde{\Delta}) = \frac{8\sqrt{2}}{15} + \frac{\sqrt{6}\pi}{4} \frac{\tilde{\Delta}}{1 - \tilde{\Delta}} \sqrt{\tilde{\mu}} + O(\tilde{\mu}) \quad (43)$$

In the limit $\tilde{\mu} \rightarrow (1 - \tilde{\Delta})/2 - 0^+$ such that $g(2\mu) \rightarrow 0^+$

which called super soft limit, we have

$$F(\bar{\mu}, \bar{\Delta}) = \frac{\pi\sqrt{\bar{\Delta}}}{4} \frac{1}{\sqrt{z}} + \frac{1 - (21 - 8\sqrt{6})\bar{\Delta}}{8(1 - \bar{\Delta})\sqrt{\bar{\Delta}}} \pi\sqrt{z} + O(z^{3/2}) \quad (44)$$

where $z = \frac{1-\bar{\Delta}}{2} - \bar{\mu}$

We investigate this Gaussian fluctuation contribution to energy per particle.

$$\frac{E}{N} = \frac{1}{n} \left[\frac{\Omega^0}{V} + \frac{1}{4\pi^2} (2m)^{\frac{3}{2}} \mu^{\frac{5}{2}} F(\tilde{\mu}, \tilde{\Delta}) \right] + \mu \quad (45)$$

$$n = n^0 - \frac{(2\mu)^{\frac{3}{2}}}{4\pi^2} \left[\tilde{\mu} F'(\tilde{\mu}, \tilde{\Delta}) + \frac{5}{2} F(\tilde{\mu}, \tilde{\Delta}) \right] \quad (46)$$

We find the Gaussian fluctuation correction increases the energy as density increases, however it will not change the mean-field result qualitatively. The numerical results are shown in Fig.(6).

Then we evaluate the density distribution in harmonic trap by Eq.(46) and substituting μ by $\mu - V(r)$. The distribution compared with mean-field result with fixed particle number is shown in Fig.4(c). One can see that the density at trap center is further increased when including the Gaussian fluctuation.

V. CONCLUSION

We studied the ground state properties of weakly interacting bose gas close to a Feshbach resonance. With the help of a path integral approach, we established a low density expansion for the equation of state of this system. We have shown that for very narrow Feshbach resonance, the energy dependence in scattering length leads to a vanishingly small inverse compressibility. When the bose gas is trapped in a harmonic potential, this effect leads to a super soft core in the trap center thus significantly increase the density in this region. Such phenomena should be able to be observed in current cold atom experiments.




FIG. 5. $F(\tilde{\mu}, \tilde{\Delta}) - F(0, 0)$ in three different limits. **(a)**: $F(\tilde{\mu}, \tilde{\Delta}) - F(0, 0)$ as a function of $\tilde{\mu}$ with $\tilde{\Delta} = 1/20$. **(b)**: $F(\tilde{\mu}, \tilde{\Delta}) - F(0, 0)$ as a function of $\tilde{\Delta}$ with $\tilde{\mu} = 1/200$. **(c)**: $F(\tilde{\mu}, \tilde{\Delta}) - F(0, 0)$ as a function of $\tilde{\mu}$ with $\tilde{\Delta} = 1/100$. Blue lines are exact result given by Eq.(36), while green dashed lines in **(a)**-**(c)** are asymptotic result in limits given by Eq.(39), Eq.(43) and Eq.(44)




FIG. 6. Energy per particle in mean-field approximation (blue dashed line) and with Gaussian fluctuation correction (red line). Parameter in pictures are resonance width $\tilde{\Delta} = \frac{|\alpha|^2}{g_r \nu_r} = \{1/100, 1/20, 1/5\}$ from top to bottom, and background scattering length $a_{bg}^2 \cdot \nu_r = \{1/64, 1/16, 1/4\}$ from left to right.

SUPPLEMENTAL MATERIAL

In this supplemental material, we provide the details on how to obtain transformation matrix \mathbf{U} appeared in Eq (24) in the main text. Considering symmetry, the transformation matrix \mathbf{U} can be written as

$$\begin{pmatrix} \eta_2 & u_2 & v_2 & \xi_2 \\ \eta_1 & u_1 & v_1 & \xi_1 \\ \xi_1 & v_1 & u_1 & \eta_1 \\ \xi_2 & v_2 & u_2 & \eta_2 \end{pmatrix} \quad (47)$$

where for convenience, we set matrix element as real numbers. The constraint that the quasiparticles after transformation are still Bosons requires

$$\begin{pmatrix} \eta_2 & u_2 & v_2 & \xi_2 \\ \eta_1 & u_1 & v_1 & \xi_1 \\ \xi_1 & v_1 & u_1 & \eta_1 \\ \xi_2 & v_2 & u_2 & \eta_2 \end{pmatrix}^T \cdot \begin{pmatrix} -1 & & & \\ & -1 & & \\ & & 1 & \\ & & & 1 \end{pmatrix} \cdot \begin{pmatrix} \eta_2 & u_2 & v_2 & \xi_2 \\ \eta_1 & u_1 & v_1 & \xi_1 \\ \xi_1 & v_1 & u_1 & \eta_1 \\ \xi_2 & v_2 & u_2 & \eta_2 \end{pmatrix} = \begin{pmatrix} -1 & & & \\ & -1 & & \\ & & 1 & \\ & & & 1 \end{pmatrix} \quad (48)$$

and that the inverse Green's function is diagonalized requires

$$\begin{pmatrix} \eta_2 & u_2 & v_2 & \xi_2 \\ \eta_1 & u_1 & v_1 & \xi_1 \\ \xi_1 & v_1 & u_1 & \eta_1 \\ \xi_2 & v_2 & u_2 & \eta_2 \end{pmatrix}^T \cdot \begin{pmatrix} \epsilon_b & \alpha & & \\ \alpha & \epsilon_a & g & \\ & g & \epsilon_a & \alpha \\ & & \alpha & \epsilon_b \end{pmatrix} \cdot \begin{pmatrix} \eta_2 & u_2 & v_2 & \xi_2 \\ \eta_1 & u_1 & v_1 & \xi_1 \\ \xi_1 & v_1 & u_1 & \eta_1 \\ \xi_2 & v_2 & u_2 & \eta_2 \end{pmatrix} = \begin{pmatrix} \omega^+ & & & \\ & \omega^- & & \\ & & \omega^- & \\ & & & \omega^+ \end{pmatrix} \quad (49)$$

where

$$\begin{aligned} \epsilon_a &\rightarrow \frac{k^2}{2m} + 2g\phi_0^2 - \mu \\ \epsilon_b &\rightarrow \frac{k^2}{m} + \nu_b - 2\mu \\ g &\rightarrow g\phi_0^2 + \sqrt{2}\alpha\beta_0 \\ \alpha &\rightarrow \sqrt{2}\alpha\phi_0 \end{aligned} \quad (50)$$

Eq.(48) gives four equations

$$\begin{aligned} u_1^2 + u_2^2 - v_1^2 - v_2^2 &= 1 \\ \eta_1^2 + \eta_2^2 - \xi_1^2 - \xi_2^2 &= 1 \\ -u_1\eta_1 - u_2\eta_2 + v_1\xi_1 + v_2\xi_2 &= 0 \\ -v_1\eta_1 - v_2\eta_2 + u_1\xi_1 + u_2\xi_2 &= 0 \end{aligned} \quad (51)$$

Eq.(49) gives six equations

$$\begin{aligned} 2\eta_1\xi_1 \epsilon_a + 2\eta_2\xi_2 \epsilon_b + (\eta_1^2 + \xi_1^2) g + 2(\eta_2\xi_1 + \eta_1\xi_2) \alpha &= 0 \\ 2u_1v_1 \epsilon_a + 2u_2v_2 \epsilon_b + (u_1^2 + v_1^2) g + 2(u_2v_1 + u_1v_2) \alpha &= 0 \\ (\eta_1u_1 + \xi_1v_1) \epsilon_a + (\eta_2u_2 + \xi_2v_2) \epsilon_b + (\xi_1u_1 + \eta_1v_1) g + (\eta_1u_2 + \eta_2u_1 + \xi_1v_2 + \xi_2v_1) \alpha &= 0 \\ (\xi_1u_1 + \eta_1v_1) \epsilon_a + (\xi_2u_2 + \eta_2v_2) \epsilon_b + (\eta_1u_1 + \xi_1v_1) g + (\xi_1u_2 + \xi_2u_1 + \eta_1v_2 + \eta_2v_1) \alpha &= 0 \\ (\eta_1^2 + \xi_1^2) \epsilon_a + (\eta_2^2 + \xi_2^2) \epsilon_b + 2\eta_1\xi_1g + (2\eta_1\eta_2 + 2\xi_1\xi_2) \alpha &= \omega^+ \\ (u_1^2 + v_1^2) \epsilon_a + (u_2^2 + v_2^2) \epsilon_b + 2u_1v_1g + (2u_1u_2 + 2v_1v_2) \alpha &= \omega^- \end{aligned} \quad (52)$$

Using Eq.(51) and first four equations in Eq.(52), we can obtain $u_1, v_1, \eta_1, \xi_1, u_2, v_2, \eta_2, \xi_2$, and substituting them into last two equations in Eq.(52) we can obtain ω^+, ω^- . First we take substitutions as follows,

$$\begin{aligned} u_1 &= (x_1 + w_1)/2 & v_1 &= (x_1 - w_1)/2 \\ \eta_1 &= (y_1 + z_1)/2 & \xi_1 &= (y_1 - z_1)/2 \\ u_2 &= (x_2 + w_2)/2 & v_2 &= (x_2 - w_2)/2 \\ \eta_2 &= (y_2 + z_2)/2 & \xi_2 &= (y_2 - z_2)/2 \end{aligned} \quad (53)$$

Then the equations can be simplified into

$$\begin{aligned}
x_1 w_1 + x_2 w_2 &= 1 \\
y_1 z_1 + y_2 z_2 &= 1 \\
w_1 y_1 + w_2 y_2 &= 0 \\
x_1 z_1 + x_2 z_2 &= 0 \\
(\epsilon_a + g)x_1^2 - (\epsilon_a - g)w_1^2 + \epsilon_b x_2^2 - \epsilon_b w_2^2 + 2\alpha(x_1 x_2 - w_1 w_2) &= 0 \\
(\epsilon_a + g)y_1^2 - (\epsilon_a - g)z_1^2 + \epsilon_b y_2^2 - \epsilon_b z_2^2 + 2\alpha(y_1 y_2 - z_1 z_2) &= 0 \\
(\epsilon_a - g)w_1 z_1 + \epsilon_b w_2 z_2 + \alpha(w_2 z_1 + w_1 z_2) &= 0 \\
(\epsilon_a + g)x_1 y_1 + \epsilon_b x_2 y_2 + \alpha(x_2 y_1 + x_1 y_2) &= 0
\end{aligned} \tag{54}$$

taking $x_2 = s x_1$, $w_2 = t w_1$ and $y_1 = s y_2$, $z_1 = t z_2$, we obtain

$$\begin{aligned}
x_1 &= \left(\frac{1}{1 + s_1 t_1} \sqrt{\frac{\epsilon_a - g + \epsilon_b t_1^2 + 2\alpha t_1}{\epsilon_a + g + \epsilon_b s_1^2 + 2\alpha s_1}} \right)^{1/2} \\
w_1 &= \left(\frac{1}{1 + s_1 t_1} \sqrt{\frac{\epsilon_a + g + \epsilon_b s_1^2 + 2\alpha s_1}{\epsilon_a - g + \epsilon_b t_1^2 + 2\alpha t_1}} \right)^{1/2} \\
x_2 &= \left(\frac{s}{1 + s_1 t_1} \sqrt{\frac{\epsilon_a - g + \epsilon_b t_1^2 + 2\alpha t_1}{\epsilon_a + g + \epsilon_b s_1^2 + 2\alpha s_1}} \right)^{1/2} \\
w_2 &= \left(\frac{t}{1 + s_1 t_1} \sqrt{\frac{\epsilon_a + g + \epsilon_b s_1^2 + 2\alpha s_1}{\epsilon_a - g + \epsilon_b t_1^2 + 2\alpha t_1}} \right)^{1/2}
\end{aligned} \tag{55}$$

$$\begin{aligned}
y_2 &= \left(\frac{1}{1 + s_2 t_2} \sqrt{\frac{(\epsilon - g) t_2^2 + \epsilon_b + 2\alpha t_2}{(\epsilon + g) s_2^2 + \epsilon_b + 2\alpha s_2}} \right)^{1/2} \\
z_2 &= \left(\frac{1}{1 + s_2 t_2} \sqrt{\frac{(\epsilon + g) s_2^2 + \epsilon_b + 2\alpha s_2}{(\epsilon - g) t_2^2 + \epsilon_b + 2\alpha t_2}} \right)^{1/2} \\
y_1 &= \left(\frac{s_2}{1 + s_2 t_2} \sqrt{\frac{(\epsilon - g) t_2^2 + \epsilon_b + 2\alpha t_2}{(\epsilon + g) s_2^2 + \epsilon_b + 2\alpha s_2}} \right)^{1/2} \\
z_1 &= \left(\frac{t_2}{1 + s_2 t_2} \sqrt{\frac{(\epsilon + g) s_2^2 + \epsilon_b + 2\alpha s_2}{(\epsilon - g) t_2^2 + \epsilon_b + 2\alpha t_2}} \right)^{1/2}
\end{aligned} \tag{56}$$

There are two sets solutions of s_1, t_1 and s_2, t_2 respectively, however, considering that in the limits $\alpha \rightarrow 0$ $u_2, v_2, \xi_2, \eta_1, \xi_1$ should tends to 0 and thus s_1, t_1, s_2, t_2 should also tends to 0, we can determine the right solutions as follows

$$\begin{aligned}
s_1 &\rightarrow \frac{g^2 - \epsilon_a^2 + \epsilon_b^2 - \sqrt{(g^2 - \epsilon_a^2 + \epsilon_b^2)^2 - 4\alpha^2(g - \epsilon_a - \epsilon_b)(g + \epsilon_a + \epsilon_b)}}{2\alpha(\epsilon_a + \epsilon_b - g)}, \\
t_1 &\rightarrow \frac{g^2 - \epsilon_a^2 + \epsilon_b^2 - \sqrt{(g^2 - \epsilon_a^2 + \epsilon_b^2)^2 - 4\alpha^2(g - \epsilon_a - \epsilon_b)(g + \epsilon_a + \epsilon_b)}}{2\alpha(g + \epsilon_a + \epsilon_b)}
\end{aligned} \tag{57}$$

$$\begin{aligned}
s_2 &\rightarrow \frac{-g^2 + \epsilon_a^2 - \epsilon_b^2 + \sqrt{(g^2 - \epsilon_a^2 + \epsilon_b^2)^2 - 4\alpha^2(g - \epsilon_a - \epsilon_b)(g + \epsilon_a + \epsilon_b)}}{2\alpha(g + \epsilon_a + \epsilon_b)}, \\
t_2 &\rightarrow \frac{-g^2 + \epsilon_a^2 - \epsilon_b^2 + \sqrt{(g^2 - \epsilon_a^2 + \epsilon_b^2)^2 - 4\alpha^2(g - \epsilon_a - \epsilon_b)(g + \epsilon_a + \epsilon_b)}}{2\alpha(\epsilon_a + \epsilon_b - g)}
\end{aligned} \tag{58}$$

Substituting the solutions of x, w, y, z (55)(56) into Eq.(53) we can finally obtain the transformation matrix.

The quantum depletion on \mathbf{k} momentum states is thus given by

$$\begin{aligned}\langle G|\hat{a}_{\mathbf{k}}^\dagger\hat{a}_{\mathbf{k}}|G\rangle &= \langle G|\left(\xi_1^2\hat{\beta}_{-\mathbf{k}}\hat{\beta}_{-\mathbf{k}}^\dagger + v_1^2\hat{\alpha}_{-\mathbf{k}}\hat{\alpha}_{-\mathbf{k}}^\dagger\right)|G\rangle \\ &= \xi_1^2 + v_1^2\end{aligned}\tag{59}$$

$$\begin{aligned}\langle G|\hat{b}_{\mathbf{k}}^\dagger\hat{b}_{\mathbf{k}}|G\rangle &= \langle G|\left(\xi_2^2\hat{\beta}_{-\mathbf{k}}\hat{\beta}_{-\mathbf{k}}^\dagger + v_2^2\hat{\alpha}_{-\mathbf{k}}\hat{\alpha}_{-\mathbf{k}}^\dagger\right)|G\rangle \\ &= \xi_2^2 + v_2^2\end{aligned}\tag{60}$$

where $|G\rangle$ is the ground state of quasiparticle at zero temperature.

-
- | | |
|--|---|
| <p>[1] T. D. Lee, K. Huang, and C. N. Yang, Phys. Rev. 106, 1135 (1957).</p> <p>[2] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman and E. A. Cornell, Science 269, 198(1995).</p> <p>[3] K. B. Davis, M. O. Mewes, M. R. Andrews, N. J. Druten, D. S. Durfee, D. M. Kurn and W. Ketterle, Phys. Rev. Lett. 75, 3969 (1995).</p> <p>[4] N. Navon, S. Piatecki, K. Günter, B. Rem, T. C. Nguyen, F. Chevy, W. Krauth, and C. Salomon, Phys. Rev. Lett. 107, 135301 (2011).</p> <p>[5] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, Rev. Mod. Phys. 82, 1225 (2010).</p> | <p>[6] T.-L. Ho, X. Cui, and W. Li, Phys. Rev. Lett. 108, 250401 (2012).</p> <p>[7] Ran Qi and Hui Zhai Phys. Rev. A 85, 041603(R) (2012).</p> <p>[8] Christian Trefzger and Yvan Castin, Phys. Rev. A 85, 053612 (2012).</p> <p>[9] Details on how to diagonalize the \mathbf{G} matrix can be found in supplementary material.</p> <p>[10] H. T. C. Stoof, Dennis B. M. Dickerscheid, and K. Gubbels, <i>Ultracold Quantum Fields</i>, (The Netherlands, Springer, 2009).</p> <p>[11] M. Edward and K. Burnett, Phys. Rev. A 51, 1382 (1995). G. Baym and C. J. Pethick, Phys. Rev. Lett. 76, 6 (1996).</p> |
|--|---|