

The Courtade-Kumar Most Informative Boolean Function Conjecture and a Symmetrized Li-Médard Conjecture are Equivalent

Leighton Pate Barnes and Ayfer Özgür
Stanford University, Stanford, CA 94305
Email: {lpb, aozgur}@stanford.edu

Abstract—We consider the Courtade-Kumar most informative Boolean function conjecture for balanced functions, as well as a conjecture by Li and Médard that dictatorship functions also maximize the L^α norm of $T_p f$ for $1 \leq \alpha \leq 2$ where T_p is the noise operator and f is a balanced Boolean function. By using a result due to Laguerre from the 1880's, we are able to bound how many times an L^α -norm related quantity can cross zero as a function of α , and show that these two conjectures are essentially equivalent.

I. INTRODUCTION

In 2014, Courtade and Kumar published [1] in which they introduced a conjecture about which Boolean function maximizes mutual information when applied to a noisy input. More concretely, they considered the following scenario. Suppose X is uniformly distributed on the n -dimensional Hamming cube $\{0, 1\}^n$. Let Y be a noisy copy of X which is the output of a memoryless binary symmetric channel with crossover probability $0 < p < \frac{1}{2}$ when X is the input. The optimization problem of interest is to find which function $f : \{0, 1\}^n \rightarrow \{0, 1\}$ maximizes the mutual information $I(f(Y); X)$ (or equivalently $I(f(X); Y)$).

Courtade and Kumar conjectured that

$$I(f(Y); X) \leq 1 - h(p) \quad (1)$$

where $h(p) = -p \log p - (1 - p) \log(1 - p)$ is the binary entropy function. This mutual information is achieved when

$$f(Y) = f(Y_1, \dots, Y_n) = Y_i$$

for any $i = 1, \dots, n$. Such functions are called *dictatorship* functions. In other words, Courtade and Kumar conjecture that there is no more informative Boolean function than simply taking one of the coordinate values.

In recent years there has been substantial interest in solving this problem, as well as some partial results. Chandar and Tchamkerten show the bound

$$I(f(Y); X) \leq (1 - 2p)^2 \quad (2)$$

in [2]. Ordentlich, Shayevitz and Weinstein prove the bound

$$I(f(Y); X) \leq \frac{\log(e)}{2} (1 - 2p)^2 + 9 \left(1 - \frac{\log(e)}{2}\right) (1 - 2p)^4$$

for $\frac{1}{2} \left(1 - \frac{1}{\sqrt{3}}\right) \leq p \leq \frac{1}{2}$ in [3], which is tighter than (2) for $\frac{1}{3} \leq p \leq \frac{1}{2}$. Using this improved bound, as well as a strengthened version of “Mrs. Gerber’s Lemma”, Samorodnitsky was able to prove the conjecture is true for all $p \geq \frac{1}{2} - \delta$ where $\delta > 0$ is some absolute constant in [4].

Along with these partial results, there have also been some related conjectures made in [5], [6]. In these works it is conjectured that dictatorship functions maximize other functionals of f , in such a way that the conjectures are stronger than the Courtade-Kumar conjecture – meaning that if the conjectures hold then (1) must also hold. Of particular interest to us is the conjecture from Li and Médard in [5] that focuses on *balanced* Boolean functions f that have fixed mean

$$\frac{1}{2^n} \sum_{y \in \{0, 1\}^n} f(y) = \frac{1}{2}.$$

They conjecture that among all such balanced functions f , the L^α -norm of the function

$$T_p f(x) = \mathbb{P}(f(Y) = 1 | X = x) \quad (3)$$

is maximized by dictatorship functions when $1 \leq \alpha \leq 2$. Li and Médard show that if their conjecture is true, then (1) must also be true for all balanced functions f . Our main contribution in this paper is to show that a slightly modified converse statement also holds. In particular, if we “symmetrize” the Li-Médard conjecture by including both $T_p f(x)$ and $1 - T_p f(x)$ in the calculation of the L^α -norm, then the Courtade-Kumar conjecture for balanced functions f also implies this symmetrized conjecture. In this sense, the two conjectures are equivalent.

In order to show this equivalence, we study the quantity

$$N_\alpha(f) = \sum_{x \in \{0, 1\}^n} (T_p f(x))^\alpha \quad (4)$$

or its symmetrized version

$$N_\alpha^{\text{sym}}(f) = \sum_{x \in \{0, 1\}^n} (T_p f(x))^\alpha + (1 - T_p f(x))^\alpha \quad (5)$$

for any $\alpha \in \mathbb{R}$. Note that these quantities are not technically norms of $T_p f$ except in the case of (4) when $\alpha = 1$. However,

(4) can be thought of as the α -power of the L^α -norm when $\alpha \geq 1$. Letting f_0 be a dictatorship function, we define

$$g_f(\alpha) = N_\alpha(f) - N_\alpha(f_0) \quad (6)$$

and the corresponding

$$g_f^{\text{sym}}(\alpha) = N_\alpha^{\text{sym}}(f) - N_\alpha^{\text{sym}}(f_0). \quad (7)$$

By using a result due to Laguerre from the 1880's [7], we will show that both $g_f(\alpha)$ and $g_f^{\text{sym}}(\alpha)$ can have at most four zeros (unless, of course, they are identically zero). This allows us to relate the validity of the Courtade-Kumar conjecture, which can be thought of as a local property of $g_f^{\text{sym}}(\alpha)$ around $\alpha = 1$, to the global properties of $g_f^{\text{sym}}(\alpha)$ such as where the zeros are located. We believe this is a powerful insight that may shed further light on the validity of these sorts of conjectures.

II. MAIN RESULTS

In this section we state precisely the relevant conjectures and how they are equivalent. In order to make sense of the first conjecture below, we first need to analyze the mutual information quantity of interest $I(f(Y); X)$. For balanced f we can write this quantity as follows:

$$\begin{aligned} I(f(Y); X) &= H(f(Y)) - H(f(Y)|X) \\ &= 1 - \frac{1}{2^n} \sum_{x \in \{0,1\}^n} h(\mathbb{P}(f(Y) = 1|X = x)) \\ &= 1 + \frac{1}{2^n} \sum_{x \in \{0,1\}^n} \left(T_p f(x) \log T_p f(x) \right. \\ &\quad \left. + (1 - T_p f(x)) \log(1 - T_p f(x)) \right). \quad (8) \end{aligned}$$

Looking at (8), it makes sense to talk about an ‘‘unsymmetrized’’ Courtade-Kumar conjecture where we consider which balanced Boolean function maximizes only the part $\sum_{x \in \{0,1\}^n} T_p f(x) \log T_p f(x)$. Recall that we say a function $f : \{0,1\}^n \rightarrow \{0,1\}$ is balanced if $\mathbb{E}[f(Y)] = \frac{1}{2}$ and that f_0 represents a dictatorship function.

Conjecture 1 (unsymmetrized Courtade-Kumar). *For any balanced f ,*

$$\sum_{x \in \{0,1\}^n} T_p f(x) \log T_p f(x) \leq \sum_{x \in \{0,1\}^n} T_p f_0(x) \log T_p f_0(x).$$

Conjecture 2 (Courtade-Kumar). *For any balanced f ,*

$$I(f(Y); X) \leq 1 - h(p).$$

Conjecture 3 (Li-Médard). *For any balanced f ,*

$$N_\alpha(f) \leq N_\alpha(f_0)$$

for $1 \leq \alpha \leq 2$.

Conjecture 4 (symmetrized Li-Médard). *For any balanced f ,*

$$N_\alpha^{\text{sym}}(f) \leq N_\alpha^{\text{sym}}(f_0)$$

for $1 \leq \alpha \leq 2$.

We establish equivalences between these conjectures in our main results below.

Theorem 1. *Conjecture 1 is true if and only if Conjecture 3 is true.*

Theorem 2. *Conjecture 2 is true if and only if Conjecture 4 is true.*

Another relationship between these conjectures is that Conjecture 1 would imply Conjecture 2, and similarly Conjecture 3 would imply Conjecture 4. This is because if f is balanced then $1 - f$ is also balanced, and $T_p(1 - f) = 1 - T_p f$. Furthermore, $1 - f_0$ is equivalent to a dictatorship function and will achieve the same value for any of the relevant functionals. We therefore have the following relationships between the various conjectures.

$$\begin{array}{ccc} \text{Conjecture 1} & \iff & \text{Conjecture 3} \\ \downarrow & & \downarrow \\ \text{Conjecture 2} & \iff & \text{Conjecture 4} \end{array}$$

It is worth pointing out that Conjecture 2 was proven in [4] in the high-noise case, i.e. for all $\frac{1}{2} - \delta < p < \frac{1}{2}$ where $\delta > 0$ is an absolute constant. Therefore our result immediately implies that Conjecture 4 is also true in this high-noise case. It is also known that Conjecture 2 holds in the low-noise case (see [1], [5]) when $0 < p < \delta_n$ and δ_n depends on n . In the same way Conjecture 4 must also hold in this dimensionally-dependent low-noise case.

III. PROOF OF THEOREMS 1 AND 2

In this section we develop the machinery needed to prove Theorems 1 and 2, as well as finish their proofs. In both cases it is straightforward to show that the Li-Médard conjectures imply their corresponding Courtade-Kumar conjectures, but the other direction is more involved. We start with the easy direction.

A. *Conjecture 3 \implies Conjecture 1*

Taking the derivative of (4) with respect to α and evaluating at $\alpha = 1$ gives

$$\left. \frac{\partial}{\partial \alpha} N_\alpha(f) \right|_{\alpha=1} = \sum_{x \in \{0,1\}^n} T_p f(x) \log T_p f(x). \quad (9)$$

Note that (9) matches the quantity from the inequality in Conjecture 1. Writing out the derivative in (9) as a difference quotient,

$$\left. \frac{\partial}{\partial \alpha} N_\alpha(f) \right|_{\alpha=1} = \lim_{\epsilon \rightarrow 0} \frac{N_{1+\epsilon}(f) - N_1(f)}{\epsilon}.$$

If Conjecture 3 is true, then $N_{1+\epsilon}(f) \leq N_{1+\epsilon}(f_0)$ for any $0 \leq \epsilon \leq 1$. Furthermore, by the law of total probability,

$$\begin{aligned} N_1(f) &= \sum_{x \in \{0,1\}^n} \mathbb{P}(f(Y) = 1 | X = x) \\ &= 2^n \mathbb{P}(f(Y) = 1) \\ &= 2^{n-1} \\ &= N_1(f_0). \end{aligned}$$

Therefore

$$\left. \frac{\partial}{\partial \alpha} N_\alpha(f) \right|_{\alpha=1} \leq \left. \frac{\partial}{\partial \alpha} N_\alpha(f_0) \right|_{\alpha=1}$$

and (9) implies Conjecture 1 must be true.

B. Conjecture 4 \implies Conjecture 2

This implication is very similar to that of Section III-A above. Differentiating (5) with respect to α and evaluating at $\alpha = 1$ gives

$$\begin{aligned} \left. \frac{\partial}{\partial \alpha} N_\alpha^{\text{sym}}(f) \right|_{\alpha=1} &= \sum_{x \in \{0,1\}^n} T_p f(x) \log T_p f(x) \\ &\quad + (1 - T_p f(x)) \log(1 - T_p f(x)) \\ &= - \sum_{x \in \{0,1\}^n} h(\mathbb{P}(f(Y) = 1 | X = x)). \end{aligned} \quad (10)$$

Conjecture 4 implies that $N_{1+\epsilon}^{\text{sym}}(f) \leq N_{1+\epsilon}^{\text{sym}}(f_0)$ for $0 \leq \epsilon \leq 1$ and $N_1^{\text{sym}}(f) = 2^n = N_1^{\text{sym}}(f_0)$. Thus

$$\left. \frac{\partial}{\partial \alpha} N_\alpha^{\text{sym}}(f) \right|_{\alpha=1} \leq \left. \frac{\partial}{\partial \alpha} N_\alpha^{\text{sym}}(f_0) \right|_{\alpha=1}$$

and Conjecture 2 holds.

C. Conjecture 2 \implies Conjecture 4

We approach this implication by contrapositive, and show that if there exists an $1 \leq \alpha \leq 2$ such that $N_\alpha^{\text{sym}}(f) > N_\alpha^{\text{sym}}(f_0)$, then $I(f(Y); X) > 1 - h(p)$. We will need the following two lemmas.

We say that a function $g : \mathbb{R} \rightarrow \mathbb{R}$ has a zero at α if $g(\alpha) = 0$, and that it has an m th order zero (or a zero with multiplicity m) at α if both $g(\alpha) = 0$ and the derivatives $g^{(j)}(\alpha) = 0$ for all $j = 1, \dots, m-1$.

Lemma 1 (Laguerre 1883 [7]). *Suppose*

$$g(\alpha) = \sum_{i=1}^N A_i e^{c_i \alpha}$$

with

$$c_1 < c_2 < \dots < c_N$$

and $A_i \neq 0$. Then g has at most as many zeros (counting multiplicities) as the number of sign changes in the sequence A_1, A_2, \dots, A_N .

Proof. The proof is by induction on the number of sign changes. For the base case note that if each value in the

sequence A_1, \dots, A_N has the same sign, then either $g(\alpha) > 0$ or $g(\alpha) < 0$ for all $\alpha \in \mathbb{R}$.

For the inductive step, suppose that the sequence A_1, \dots, A_N changes sign k times as we move from A_1 to A_N . Suppose that one of the sign changes occurs between indices l and $l+1$, i.e., that $\text{sign}(A_l) \neq \text{sign}(A_{l+1})$. Letting $c_l < b < c_{l+1}$, we can rewrite $g(\alpha)$ as

$$g(\alpha) = e^{b\alpha} \sum_{i=1}^N A_i e^{(c_i-b)\alpha}. \quad (11)$$

We isolate the second factor from (11) above and define

$$h(\alpha) = \sum_{i=1}^N A_i e^{(c_i-b)\alpha}. \quad (12)$$

Since $e^{b\alpha} > 0$, the general Leibniz rule for derivatives of products implies that both g and h will have zeros in the same locations and with the same multiplicities. We will therefore focus on counting the zeros of $h(\alpha)$. Suppose that h has zeros at $\alpha_1, \dots, \alpha_r$ with corresponding multiplicities m_1, \dots, m_r and let $M = \sum_{j=1}^r m_r$.

The derivative $h'(\alpha)$ will have $\sum_{j=1}^r \max\{m_j - 1, 0\}$ zeros at the points $\alpha_1, \dots, \alpha_r$, and by Rolle's theorem [8] at least $r-1$ zeros between these points. So in total $h'(\alpha)$ will have at least $M-1$ zeros. Differentiating (12) with respect to α ,

$$\frac{\partial}{\partial \alpha} \sum_{i=1}^N A_i e^{(c_i-b)\alpha} = \sum_{i=1}^N (c_i - b) A_i e^{(c_i-b)\alpha}. \quad (13)$$

Display (13) fits exactly the form of the function required by the Lemma – it is a sum of ordered exponentials with $k-1$ sign changes in the coefficients. Hence, by the inductive assumption, the function $h'(\alpha)$ can have at most $k-1$ zeros. Putting this all together we have $M-1 \leq k-1$ and $M \leq k$ as desired. \square

Lemma 2. *For any balanced f , $N_2(f) \leq N_2(f_0)$ and $N_2^{\text{sym}}(f) \leq N_2^{\text{sym}}(f_0)$ with equality only if f is a dictatorship function.*

Lemma 2 follows immediately from taking the Fourier transform of $T_p f$ and using the Parseval/Rayleigh/Plancherel Theorem. See also [2], [5]. A proof is included in the appendix for completeness

In order to apply Lemma 1, we will consider the function $g_f^{\text{sym}}(\alpha)$ from (7). This function takes the required form since we can write it as

$$\begin{aligned} g_f^{\text{sym}}(\alpha) &= \sum_{x \in \{0,1\}^n} e^{\alpha \log T_p f(x)} + e^{\alpha \log(1-T_p f(x))} \\ &\quad - e^{\alpha \log T_p f_0(x)} - e^{\alpha \log(1-T_p f_0(x))}. \end{aligned} \quad (14)$$

It is clear that $T_p f_0(x) = p$ for 2^{n-1} different values of x , and $T_p f_0(x) = 1-p$ for the remaining 2^{n-1} values of x . We will assume without loss of generality that $T_p f(x_0) < p$ or

$T_p f(x_0) > 1 - p$ for some x_0 because if this is not the case we would necessarily have

$$(T_p f(x))^\alpha + (1 - T_p f(x))^\alpha \leq (T_p f_0(x))^\alpha + (1 - T_p f_0(x))^\alpha$$

for all x and $\alpha \geq 1$ and there would be nothing to prove. Furthermore, we must have at least one x such that $p < T_p f(x) < 1 - p$ otherwise this would contradict Lemma 2. The ordered sequence of the c_i (which take the form $\log(T_p f(x))$, $\log(1 - T_p f(x))$, $\log(T_p f_0(x))$, and $\log(1 - T_p f_0(x))$) will therefore give rise to a sequence of signs that looks like $+ - + - +$ in the coefficients A_i . There are therefore exactly four sign changes in the sequence A_i , and applying Lemma 1, $g_f^{\text{sym}}(\alpha)$ can have at most four zeros where $N_\alpha^{\text{sym}}(f_0) = N_\alpha^{\text{sym}}(f)$.

Two of the zeros of $g_f^{\text{sym}}(\alpha)$ must occur at $\alpha = 0$ and $\alpha = 1$. At $\alpha = 2$, Lemma 2 implies that $g_f^{\text{sym}}(2) < 0$. However, as α gets large, $N_\alpha^{\text{sym}}(f_0) < N_\alpha^{\text{sym}}(f)$ and $g_f^{\text{sym}}(\alpha) > 0$ since $T_p(x_0) > 1 - p$ or $1 - T_p(x_0) > 1 - p$. So by the continuity of $N_\alpha^{\text{sym}}(f)$ with respect to α we must have another zero of $g_f^{\text{sym}}(\alpha)$ with $\alpha > 2$. In a similar way, $N_\alpha^{\text{sym}}(f_0) < N_\alpha^{\text{sym}}(f)$ as α approaches $-\infty$.

With all of these restrictions on $g_f^{\text{sym}}(\alpha)$ in mind, we return to showing that Conjecture 2 implies Conjecture 4 by contrapositive. If there exists an $1 \leq \alpha \leq 2$ such that $N_\alpha^{\text{sym}}(f) > N_\alpha^{\text{sym}}(f_0)$, then the fourth and final zero of $g_f^{\text{sym}}(\alpha)$ must occur for some $1 < \alpha < 2$ (since $g_f^{\text{sym}}(2) < 0$ and the function is continuous). In this case, the function $g_f^{\text{sym}}(\alpha)$ must take the form depicted in Figure 1, and in particular it must be positive for $1 < \alpha < 1 + \epsilon$ for some $\epsilon > 0$. By looking at the difference quotient,

$$\left. \frac{\partial}{\partial \alpha} N_\alpha^{\text{sym}}(f) \right|_{\alpha=1} \geq \left. \frac{\partial}{\partial \alpha} N_\alpha^{\text{sym}}(f_0) \right|_{\alpha=1}.$$

We can rule out the possibility that

$$\left. \frac{\partial}{\partial \alpha} N_\alpha^{\text{sym}}(f) \right|_{\alpha=1} = \left. \frac{\partial}{\partial \alpha} N_\alpha^{\text{sym}}(f_0) \right|_{\alpha=1}$$

because it would imply $g_f^{\text{sym}}(\alpha)$ had a second-order zero at $\alpha = 1$ which would contradict Lemma 1. Thus

$$\left. \frac{\partial}{\partial \alpha} N_\alpha^{\text{sym}}(f) \right|_{\alpha=1} > \left. \frac{\partial}{\partial \alpha} N_\alpha^{\text{sym}}(f_0) \right|_{\alpha=1}$$

and because of (10),

$$I(f(Y); X) > 1 - h(p).$$

D. Conjecture 1 \implies Conjecture 3

The proof of this implication is roughly the same as that of Section III-C. The main difference is that there may exist an x_0 with $T_p f(x_0) < p$ or $T_p f(x_0) > 1 - p$, and since we are not including the corresponding $1 - T_p f(x_0)$ term, there might not be any x_1 with $T_p f(x_1) > 1 - p$ or $T_p f(x_1) < p$, respectively. This opens up the possibility that there could be only three sign changes in the coefficients of $g_f(\alpha)$. If there are four sign changes, then the result follows just as in Section III-C. Let us therefore assume that there are three sign changes.




Fig. 1. The $g_f^{\text{sym}}(\alpha)$ curve as a function of α from the proof in Section III-C. The position of the four unique zeros relative to each other are marked, and it is clear where the curve must be positive and negative. Note that this is not an actual $g_f^{\text{sym}}(\alpha)$ curve corresponding to a specific Boolean function f – if that were true it would be a counterexample to the Courtade-Kumar conjecture, none of which are known.

There are two trivial zeros of $g_f(\alpha)$ at $\alpha = 0$ and $\alpha = 1$, and if there exists an $1 \leq \alpha \leq 2$ such that $N_\alpha(f) > N_\alpha(f_0)$, then by Lemma 2 and the continuity of $g_f(\alpha)$ the third zero must occur for some $1 < \alpha < 2$. In this case, the curve $g_f(\alpha)$ must be positive between $\alpha = 1$ and the third zero, and

$$\left. \frac{\partial}{\partial \alpha} N_\alpha(f) \right|_{\alpha=1} > \left. \frac{\partial}{\partial \alpha} N_\alpha(f_0) \right|_{\alpha=1}$$

just like in Section III-C. Therefore,

$$\sum_{x \in \{0,1\}^n} T_p f(x) \log T_p f(x) > \sum_{x \in \{0,1\}^n} T_p f_0(x) \log T_p f_0(x)$$

and the proof by contrapositive is complete.

IV. AN EXAMPLE $g_f^{\text{sym}}(\alpha)$

In order to see what these $g_f^{\text{sym}}(\alpha)$ curves can look like, consider the following example. Let $n = 3$ and

$$f(y) = \begin{cases} 1 & , \text{ if } y_1 + y_2 + y_3 \geq 2 \\ 0 & , \text{ otherwise} \end{cases}$$

i.e., the majority function. For this f we have

$$\begin{aligned} g_f^{\text{sym}}(\alpha) &= 2 \left((1-p)^3 + 3p(1-p)^2 \right)^\alpha \\ &\quad + 6 \left((1-p)^3 + (1-p)^2 p + 2(1-p)p^2 \right)^\alpha \\ &\quad + 6 \left(2(1-p)^2 p + (1-p)p^2 + p^3 \right)^\alpha \\ &\quad + 2 \left(p^3 + 3(1-p)p^2 \right)^\alpha \\ &\quad - 8p^\alpha - 8(1-p)^\alpha. \end{aligned}$$

The theory from Lemma 1 proves that this function will have at most four zeros. Trivial zeros occur at $\alpha = 0$ and $\alpha = 1$, and there must be an additional one for $\alpha > 2$. In Figure 2 we show this curve for three different values of p , showing three different possible behaviors.




Fig. 2. The $g_f^{\text{sym}}(\alpha)$ curve as a function of α for the $n = 3$ majority function with $p = .21$ (top), $p = .068$ (middle), and $p = .017$ (bottom). There is a zero at large α in the middle and bottom curves that is not visible in these plots. The fourth zero for the top curve occurs at $\alpha < 0$. In the middle curve, the “fourth zero” is accounted for by a second order zero at $\alpha = 0$. In the bottom curve, the fourth zero occurs at $0 < \alpha < 1$. None of the curves has a zero at $1 < \alpha < 2$, as this would be equivalent to f being a counterexample to the Courtade-Kumar conjecture.

V. APPENDIX

In this appendix, we briefly introduce the Fourier analysis of Boolean functions in order to prove Lemma 2. For a full treatment we recommend [9]. A Boolean function $f : \{0, 1\}^n \rightarrow \{0, 1\}$ has Fourier expansion

$$f(y) = \sum_{v \in \{0,1\}^n} \hat{f}(v) W_v(y)$$

where the set of Boolean functions $\{W_v\}_{v \in \{0,1\}^n}$ defined by

$$W_v(y) = \frac{1}{\sqrt{2^n}} (-1)^{\sum_{i=1}^n v_i y_i}$$

forms an orthonormal basis.

One way to think about the probability $\mathbb{P}(f(Y) = 1 | X = x)$, and the origin of the notation from (3), is that it is the function of x obtained via the noise operator T_p applied to f . The function $T_p f$ can be expressed as

$$T_p f(x) = \sum_{y \in \{0,1\}^n} p^{d(x,y)} (1-p)^{n-d(x,y)} f(y)$$

where $d(\cdot, \cdot)$ is the Hamming distance, so the operator T_p can also be thought of as convolution with the kernel $\varphi_p(x) = p^{d(0,x)} (1-p)^{n-d(0,x)}$. Thus using the Convolution Theorem,

$$T_p f(x) = \sum_{v \in \{0,1\}^n} (1-2p)^{d(0,v)} \hat{f}(v) W_v(x)$$

where $\lambda_v = (1-2p)^{d(0,v)}$ for $v \in \{0,1\}^n$ are the eigenvalues of the T_p operator. Using Parseval’s theorem,

$$\begin{aligned} N_2(f) &= \sum_{x \in \{0,1\}^n} (T_p f(x))^2 \\ &= \sum_{v \in \{0,1\}^n} (1-2p)^{2d(0,v)} (\hat{f}(v))^2 \\ &= \frac{2^n}{4} + \sum_{v \neq 0} (1-2p)^{2d(0,v)} (\hat{f}(v))^2 \end{aligned} \quad (15)$$

where the last equality (15) follows from f being balanced. Since

$$\sum_{y \in \{0,1\}^n} (f(y))^2 = \frac{2^n}{2} = \sum_{v \in \{0,1\}^n} (\hat{f}(v))^2$$

for all balanced f , it is clear that in order to maximize (15) the remaining (non-DC-component) energy in f should all be concentrated in the Fourier coefficients with $d(0,v) = 1$, i.e., the “weight” one Fourier coefficients. The only balanced Boolean functions with all of their energy concentrated in the weight zero and one Fourier coefficients are the dictatorship functions, so we have

$$N_2(f) \leq N_2(f_0)$$

with equality only when f is a dictatorship function. Since f and $1-f$ are both dictatorship functions that maximize this quantity, we similarly have

$$N_2^{\text{sym}}(f) \leq N_2^{\text{sym}}(f_0).$$

REFERENCES

- [1] T. A. Courtade and G. R. Kumar, “Which boolean function maximize mutual information on noisy inputs?” *IEEE Transactions on Information Theory*, vol. 60, no. 8, 2014.
- [2] V. Chandar and A. Tchamkerten, “Most informative quantization functions,” *Proc. ITA Workshop, San Diego, CA, USA*, 2014.
- [3] O. Ordentlich, O. Shayevitz, and O. Weinstein, “An improved upper bound for the most informative boolean function conjecture,” *Proceedings of the International Symposium on Information Theory (ISIT)*, 2016.
- [4] A. Samorodnitsky, “On the entropy of a noisy function,” *IEEE Transactions on Information Theory*, vol. 62, no. 10, p. 5446–5464, 2016.
- [5] J. Li and M. Médard, “Boolean functions: noise stability, non-interactive correlation distillation, and mutual information.” [Online]. Available: <https://arxiv.org/pdf/1801.04462.pdf>
- [6] V. Anantharam, A. Bogdanov, A. Chakrabarti, T. Jayram, and C. Nair, “A conjecture regarding optimality of the dictator function under hellinger distance.” [Online]. Available: <http://chandra.ie.cuhk.edu.hk/pub/papers/HC/hel-conj.pdf>
- [7] E. N. Laguerre, “Sur la théorie des équations numériques,” *Journal de Mathématiques pures et appliquées*, 1883. [Online]. Available: <http://sepwww.stanford.edu/oldsep/stew/laguerre.pdf>
- [8] A. Mattuck, *Introduction to Analysis*. Prentice Hall, 1999.
- [9] R. O’Donnell, *Analysis of Boolean Functions*. Cambridge University Press, 2014.

ACKNOWLEDGEMENTS

This work was supported in part by NSF award CCF-1704624 and by the Center for Science of Information (CSoI), an NSF Science and Technology Center, under grant agreement CCF-0939370.