

A NOTE ON PAIR DEPENDENT LINEAR STATISTICS WITH SLOWLY GROWING VARIANCE.

ANDER AGUIRRE AND ALEXANDER SOSHNIKOV

ABSTRACT. We prove Gaussian fluctuation for pair counting statistics of the form $\sum_{1 \leq i \neq j \leq N} f(\theta_i - \theta_j)$ for the Circular Unitary Ensemble (CUE) of random matrices in the case of a slowly growing variance in the limit of large N .

1. INTRODUCTION

The study of random matrix theory goes back to the principal component data analysis of J. Wishart in 1920s-1930s and revolutionary ideas of E. Wigner in quantum physics in 1950s that linked statistical properties of the energy levels of heavy-nuclei atoms with spectral properties of Hermitian random matrices with independent components.

In 1960s, F. Dyson introduced three archetypal types of matrix ensembles: Circular Orthogonal Ensemble (COE), Circular Unitary Ensemble (CUE), and Circular Symplectic Ensemble (CSE), see e.g. [7]-[10]. The probability density of the eigenvalues $\{e^{i\theta_j}\}_{j=1}^N$, $0 \leq \theta_1, \dots, \theta_N < 2\pi$, is given by

$$(1.1) \quad p_N(\bar{\theta}) = \frac{1}{Z_N(\beta)} \prod_{1 \leq j < k \leq N} |e^{i\theta_j} - e^{i\theta_k}|^\beta,$$

where $\beta = 1, 2, \text{ and } 4$ correspond to COE, CUE, and CSE, correspondingly. For arbitrary $\beta > 0$, a (sparse) random matrix model with eigenvalues distribution following (1.1) was introduced in [15]. The ensemble (1.1) for arbitrary $\beta > 0$ is known as the Circular Beta Ensemble ($C\beta U$).

The Circular Unitary Ensemble ($\beta = 2$) corresponds to the joint distribution of the eigenvalues of an $N \times N$ random unitary matrix U distributed according to the Haar measure. In particular, the partition function is given by

$$(1.2) \quad Z_N(2) = (2\pi)^N \times N!$$

In [1], pair counting statistics of the form

$$(1.3) \quad S_N(f) = \sum_{1 \leq i \neq j \leq N} f(L_N(\theta_i - \theta_j)),$$

were studied for $C\beta E$ (1.1) for $1 \leq L_N \leq N$ under certain smoothness assumptions on f . The research in [1] was motivated by a classical result of Montgomery on pair correlation of zeros of the Riemann zeta function [18]-[19]. Assuming the Riemann

2010 *Mathematics Subject Classification.* Primary: 60F05.

Key words and phrases. Random Matrices, Circular Unitary Ensembles, Pair Counting Statistics, Central Limit Theorem.

Hypothesis, Montgomery studied the distribution of the “non-trivial” zeros on the critical line $1/2 + i\mathbb{R}$. Rescaling zeros $\{1/2 \pm \gamma_n\}$,

$$\tilde{\gamma}_n = \frac{\gamma_n}{2\pi} \log(\gamma_n),$$

Montgomery considered the statistic $\sum_{0 < \tilde{\gamma}_j \neq \tilde{\gamma}_k < T} f(\tilde{\gamma}_j - \tilde{\gamma}_k)$ for large T and sufficiently fast decaying f with $\text{supp } \mathcal{F}(f) \subset [-\pi, \pi]$, where $\mathcal{F}(f)$ denotes the Fourier transform of f . The results of [18]-[19] imply that the two-point correlations of the (rescaled) critical zeros coincide in the limit with the local two point correlations of the eigenvalues of a CUE random matrix.

The results of [1] deal with the limiting behavior of (1.3) in three different regimes, namely macroscopic ($L_N = 1$), mesoscopic ($1 \ll L_N \ll N$) and microscopic ($L_N = N$). In the unscaled $L_N = 1$ case it was shown that

$$(1.4) \quad S_N(f) = \sum_{1 \leq i \neq j \leq N} f(\theta_i - \theta_j),$$

has a non-Gaussian fluctuation in the limit $N \rightarrow \infty$ provided f is a sufficiently smooth function on the unit circle.

Namely, let f be a real even integrable function on the unit circle. Denote the Fourier coefficients of f as

$$(1.5) \quad \hat{f}(k) = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-ikx} dx.$$

Let us assume that $f' \in L^2(\mathbb{T})$ for $\beta = 2$, $\sum_{k \in \mathbb{Z}} |\hat{f}(k)| |k| < \infty$ for $\beta < 2$, $\sum_{k \in \mathbb{Z}} |\hat{f}(k)| \times |k| \log(|k| + 1) < \infty$ for $\beta = 4$, and $\sum_{k \in \mathbb{Z}} |\hat{f}(k)| |k|^2 < \infty$ for $\beta \in (2, 4) \cup (4, \infty)$. Then we have the following convergence in distribution as $N \rightarrow \infty$:

$$S_N(f) - \mathbb{E}S_N(f) \xrightarrow{\mathcal{D}} \frac{4}{\beta} \sum_{k=1}^{\infty} \hat{f}(k) k (\varphi_k - 1),$$

where φ_m are i.i.d. exponential random variables with $\mathbb{E}(\varphi_m) = 1$. For $\beta = 2$ the result was proven under the optimal condition $\sum_{k \in \mathbb{Z}} |\hat{f}(k)|^2 |k|^2 < \infty$.

The goal of this paper is to study the fluctuation of the pair counting statistic (1.4) when $\text{Var}(S_N(f))$ slowly grows with N to infinity.

Definition 1.1. A positive sequence V_N is said to be slowly varying in sense of Karamata ([4]) if

$$(1.6) \quad \lim_{n \rightarrow \infty} \frac{V_{\lfloor \lambda N \rfloor}}{V_N} = 1, \quad \forall \lambda > 0,$$

where $\lfloor m \rfloor$ denotes the integer part of m .

The following notation will be used throughout the paper:

$$(1.7) \quad V_N = \sum_{k=-N}^{k=N} |f(\hat{k})|^2 |k|^2.$$

Theorem 1.2. *Let $f \in L^2(\mathbb{T})$ be a real even function such that $V_N = \sum_{k=-N}^{k=N} |f(\hat{k})|^2 |k|^2$, $N = 1, 2, \dots$, is a slowly varying sequence that diverges to infinity as $N \rightarrow \infty$. Then we have the following convergence in distribution*

$$\frac{S_N(f) - \mathbb{E}S_N(f)}{\sqrt{2 \sum_{-N}^N |\hat{f}(k)|^2 |k|^2}} \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1)$$

Linear statistics of the eigenvalues of random matrices $\sum_{j=1}^N f(\lambda_j)$ have been studied extensively in the literature. Johansson ([11]) proved for (1.1) for arbitrary $\beta > 0$ and sufficiently smooth real-valued f that

$$\frac{\sum_{j=1}^N f(\theta_j) - N\hat{f}(0)}{\sqrt{\frac{2}{\beta} \sum_{-\infty}^{\infty} |\hat{f}(k)|^2 |k|}}$$

converges in distribution to a standard Gaussian random variable. In particular, for $\beta = 2$ he proved the result under the optimal conditions on f , namely

$$\sum_{-\infty}^{\infty} |\hat{f}(k)|^2 |k| < \infty.$$

If the variance of the linear statistic goes to infinity with N , Diaconis and Evans [5] proved the CLT in the case $\beta = 2$ provided the sequence $\{\sum_{-N}^N |\hat{f}(m)|^2 |m|\}_{n \in \mathbb{N}}$ is slowly varying.

For the results on the linear eigenvalue statistics in the mesoscopic regime $\sum_{j=1}^N f(L_N \theta_j)$, $1 \ll L_N \ll N$, we refer the reader to [21], [3], [16], and references therein. For additional results on the spectral properties of C β U we refer the reader to [6], [12], [2], [22], [14], [17], [20], [23], [24], and references therein.

The proof of the main result of the paper (Theorem 1.2) is given in the next section. Throughout the paper, the notation $a_N = O(b_N)$ means that the ratio a_N/b_N is bounded from above in absolute value. The notation $a_N = o(b_N)$ means that $a_N/b_N \rightarrow 0$ as $N \rightarrow \infty$. Occasionally, for non-negative quantities, in this case we will also use the notation $a_N \ll b_N$.

Research has been partially supported by the Simons Foundation Collaboration Grant for Mathematicians #312391.

2. PROOF OF THEOREM 1.2

The section is devoted to the proof of Theorem 1.2. We start by recalling the formula for the variance of $S_N(f)$ from Proposition 4.1 of [1]:

$$(2.1) \quad \begin{aligned} \text{Var}(S_N(f)) = & 4 \sum_{1 \leq s \leq N-1} s^2 |\hat{f}(s)|^2 + 4(N^2 - N) \sum_{N \leq s} |\hat{f}(s)|^2 \\ & - 4 \sum_{\substack{1 \leq s, t \\ 1 \leq |s-t| \leq N-1 \\ N \leq \max(s, t)}} (N - |s-t|) \hat{f}(s) \hat{f}(t) - 4 \sum_{\substack{1 \leq s, t \leq N-1 \\ N+1 \leq s+t}} ((s+t) - N) \hat{f}(s) \hat{f}(t). \end{aligned}$$

Our first goal is to show that the last two (off-diagonal) terms in the variance expression (2.1) are much smaller than $V_N = \sum_{s=-N}^N s^2 |\hat{f}(s)|^2$ for large N provided (1.6) is satisfied.

Lemma 2.1. *Let V_N from (1.7) be a slowly varying sequence diverging to infinity as $N \rightarrow \infty$. Then, as $N \rightarrow \infty$, we have*

(i)

$$\sum_{\substack{1 \leq s, t \leq N \\ s+t \geq N+1}} s |\hat{f}(s)| \cdot |\hat{f}(t)| = o(V_N),$$

(ii)

$$(N+1) \sum_{\substack{s-t \leq N \\ s \geq N+1 \\ 1 \leq t \leq N}} |\hat{f}(s)| \cdot |\hat{f}(t)| = o(V_N),$$

(iii)

$$N \sum_{\substack{|s-t| \leq N-1 \\ s, t \geq N}} |\hat{f}(s)| \cdot |\hat{f}(t)| = o(V_N).$$

The proof of the lemma is somewhat similar to the proof of Lemma 4.4. in [1]. For the convenience of the reader, we give the full details of the proof below.

Proof of Lemma 2.1

Proof of (i).

Let $x_s = s |\hat{f}(s)|$ for $1 \leq s \leq N$ and $X_N = \{x_s\}_{s=1}^N$. Define a vector $Y_N := X_N \times \mathbf{1}_{(s > N/2)}$, so that the first $\lfloor N/2 \rfloor$ coordinates of Y_N are zero and the rest coincide with the corresponding coordinates of X_N . We note that

$$(2.2) \quad 2 \|X_N\|_2^2 = V_N \text{ and } \|Y_N\|_2^2 = o(V_N),$$

where $\|X\|_2$ denotes the Euclidean norm of a vector $X \in \mathbb{R}^N$. The last bound follows from the condition (1.6) on the slow growth of V_N .

We now write the off diagonal variance term in (i) as a bilinear form:

$$(2.3) \quad \begin{aligned} \sum_{\substack{1 \leq s, t \leq N \\ s+t \geq N+1}} s |\hat{f}(s)| \cdot |\hat{f}(t)| &= \sum_{t=1}^N x_t \cdot \left(\frac{1}{t} \sum_{s=N-t+1}^N x_s \right) \\ &= \sum_{t=1}^N x_t \cdot \left(\frac{1}{t} \sum_{s=1}^t (U_N X_N)_s \right) = \langle X_N, A_N X_N \rangle, \end{aligned}$$

with $A_N = B_N U_N$, where U_N is a unitary permutation matrix given by $(U_N)_{s,t} = \mathbf{1}_{(t=N-s+1)}$ and B_N is a lower triangular matrix given by $(B_N)_{s,t} = (1/s) \mathbf{1}_{(t \leq s)}$. The

matrix A_N is given by:

$$A_N = \begin{pmatrix} 0 & 0 & 0 & 0 & \dots & 1 \\ 0 & 0 & \ddots & \ddots & \frac{1}{2} & \frac{1}{2} \\ 0 & \ddots & \ddots & \frac{1}{3} & \frac{1}{3} & \frac{1}{3} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \frac{1}{N} & \frac{1}{N} & \frac{1}{N} & \frac{1}{N} & \dots & \frac{1}{N} \end{pmatrix}$$

Taking into account that the upper-right $\lfloor N/2 \rfloor \times \lfloor N/2 \rfloor$ block of A_N is zero, we can bound $\langle X_N, A_N X_N \rangle \leq \langle X_N, A_N Y_N \rangle + \langle Y_N, A_N X_N \rangle$. It was shown in [1] that $\|A_N\|_{op} \leq 3$, where $\|A\|_{op}$ denotes the operator norm. This implies that the expression in (2.3) is bounded from above by $3\|X_N\|_2\|Y_N\|_2 = o(V_N)$ by (2.2). This completes the proof of Lemma 2.1(i).

To prove part (ii), let B_N be defined as in the proof of part (i). Similarly, let $x_s = s|\hat{f}(s)|$, $1 \leq s \leq 2N$, and $X_N = \{x_s\}_{s=1}^{2N}$. Now, X_N is a $2N$ -dimensional vector such that $\|X_N\|_2^2$ grows slowly in N . Define $Y_N := X_N \mathbf{1}_{(s > N)}$, so that the first N coordinates of Y_N are zero and the rest coincide with the corresponding coordinates of X_N . Observe that

$$\begin{aligned} N \sum_{\substack{s-t \leq N \\ s \geq N+1 \\ 1 \leq t \leq N}} |\hat{f}(s)| \cdot |\hat{f}(t)| &\leq \sum_{t=1}^N x_t \left(\frac{1}{t} \sum_{s=N+1}^{N+t} x_s \right) \\ &= \langle C_N X_N, D_N X_N \rangle, \end{aligned}$$

where

$$C_N = \begin{pmatrix} I_N & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad D_N = \begin{pmatrix} 0 & B_N \\ 0 & 0 \end{pmatrix}$$

We note that $\|D_N\|_{op} \leq 3$ and $\|C_N\|_{op} = 1$. Once again:

$$\langle C_N X_N, D_N X_N \rangle = \langle C_N X_N, D_N Y_N \rangle \leq 3\|X_N\|_2\|Y_N\|_2 = o(V_N).$$

The proof of (ii) is completed.

Proof of (iii). We start by estimating the l.h.s. of (iii) from above by

$$(2.4) \quad 2N \sum_{\substack{t-N+1 \leq s \leq N+t-1 \\ t \geq s \geq N}} |\hat{f}(s)| \cdot |\hat{f}(t)|.$$

As before, let $x_s = s|\hat{f}(s)|$ for $s \geq 1$. Define $X = \{x_s\}_{s=1}^{\infty}$ and $X^{(j)} = X \mathbf{1}_{(jN \leq s < (j+1)N)}$, $j = 0, 1, 2, \dots$. We can bound (2.4) from above by

$$(2.5) \quad 2 \sum_{t=N}^{\infty} x_t \left(\frac{1}{t} \sum_{s=t-N+1}^t x_s \right) = 2 \sum_{j=1}^{\infty} \sum_{t=jN}^{(j+1)N-1} x_t \left(\frac{1}{t} \sum_{s=t-N+1}^t x_s \right),$$

One can write the second sum at the r.h.s. of (2.5) as

$$(2.6) \quad \sum_{t=jN}^{(j+1)N-1} x_t \left(\frac{1}{t} \sum_{s=t-N+1}^t x_s \right) = \langle X^{(j)}, R_{N,j}(X^{(j-1)} + X^{(j)}) \rangle,$$

where $R_{N,j}$ is a bounded linear operator such that

$$(2.7) \quad (R_{N,j})_{t,s} = \frac{1}{t} \mathbf{1}_{(t-N+1 \leq s \leq t)} \mathbf{1}_{(jN \leq t < (j+1)N)}.$$

The operator norm of $R_{N,j}$ is bounded from above by its Hilbert-Schmidt norm

$$(2.8) \quad \|R_{N,j}\|_{op} \leq \|R_{N,j}\|_2 = \sqrt{N \sum_{t=jN}^{(j+1)N-1} \frac{1}{t^2}} \leq \sqrt{\frac{N^2}{j^2 N^2}} = \frac{1}{j}.$$

Now, by the Cauchy-Schwarz inequality, the r.h.s. of (2.6) can be bounded from above by

$$(2.9) \quad \langle X^{(j)}, R_{N,j}(X^{(j-1)} + X^{(j)}) \rangle \leq \|X^{(j)}\|_2 \|R_{N,j}\|_{op} (\|X^{(j-1)}\|_2 + \|X^{(j)}\|_2)$$

$$(2.10) \quad \leq \frac{1}{j} \|X^{(j)}\|_2^2 + \frac{1}{j} \|X^{(j)}\|_2 \|X^{(j-1)}\|_2.$$

Summing up the r.h.s. of the last inequality with respect to $j \geq 1$ gives $o(V_N)$. Indeed, $2\|X^{(0)}\|_2^2 = V_N$ and summation by parts gives

$$(2.11) \quad \sum_{j=1}^{\infty} \frac{1}{j} \|X^{(j)}\|_2^2 \leq \sum_{j=1}^{\infty} \frac{1}{j^2} (V_{jN} - V_N).$$

The condition (1.6) on the slow growth of V_N (1.6) implies that the r.h.s. of (2.11) is $o(V_N)$. To sum the second term in (2.10), we write

$$(2.12) \quad \sum_{j=1}^{\infty} \frac{1}{j} \|X^{(j)}\|_2 \|X^{(j-1)}\|_2 = \|X^{(1)}\|_2 \|X^{(0)}\|_2 + \sum_{j=2}^{\infty} \frac{1}{j} \|X^{(j)}\|_2 \|X^{(j-1)}\|_2.$$

The first term on the r.h.s. of (2.12) is $\sqrt{(V_{2N} - V_N)V_N} = o(V_N)$. To deal with the second term, we use the Cauchy-Schwarz inequality and proceed as in (2.11). This completes the proof of Lemma 2.1.

Proof of Theorem 1.2

We now proceed to finish the proof of Theorem 1.2. We recall that

$$(2.13) \quad S_N(f) = \sum_{1 \leq i \neq j \leq N} f(\theta_i - \theta_j) = 2 \sum_{k=1}^{\infty} \hat{f}(k) \left| \sum_{j=1}^N \exp(ik\theta_j) \right|^2 + \hat{f}(0)N^2 - Nf(0).$$

Denote by $t_{N,k}$ the trace of the k -th power of a CUE matrix, i.e.

$$(2.14) \quad t_{N,k} := \sum_{j=1}^N e^{ik\theta_j}, \quad k = 0, \pm 1, \pm 2, \dots$$

Then

$$(2.15) \quad \frac{S_N(f) - \mathbb{E}S_N(f)}{\sqrt{V_N}} = \frac{2}{\sqrt{V_N}} \sum_{k=1}^{\infty} \hat{f}(k)(|t_{N,k}|^2 - \mathbb{E}|t_{N,k}|^2) = \frac{2}{\sqrt{V_N}} \sum_{k=1}^{\lfloor N/M_N \rfloor} \hat{f}(k)(|t_{N,k}|^2 - \mathbb{E}|t_{N,k}|^2) + \frac{2}{\sqrt{V_N}} \sum_{\lfloor N/M_N \rfloor + 1}^{\infty} \hat{f}(k)(|t_{N,k}|^2 - \mathbb{E}|t_{N,k}|^2).$$

Here $\{M_N\}_{N=1}^{\infty}$ is positive integer-valued sequence sufficiently slowly growing to infinity as $N \rightarrow \infty$ in such a way that

$$(2.16) \quad \lim_{N \rightarrow \infty} \frac{V_{\lfloor NM_N \rfloor}}{V_N} = \lim_{N \rightarrow \infty} \frac{V_N}{V_{\lfloor N/M_N \rfloor}} = 1.$$

The existence of such a sequence follows from (1.6).

It follows from Lemma 2.1 that the second moment of the second term on the r.h.s. of (2.15) is going to zero as $N \rightarrow \infty$. We formulate this result in the next lemma.

Lemma 2.2.

$$(2.17) \quad \text{Var} \left(\sum_{\lfloor N/M_N \rfloor + 1}^{\infty} \hat{f}(k)|t_{N,k}|^2 \right) = o(V_N).$$

Proof of Lemma 2.2

It follows from (2.1) that

$$(2.18) \quad \text{Var} \left(\sum_{\lfloor N/M_N \rfloor + 1}^{\infty} \hat{f}(k)|t_{N,k}|^2 \right) = 4 \sum_{N/M_N < s \leq N-1} s^2 |\hat{f}(s)|^2 + 4(N^2 - N) \sum_{N \leq s} |\hat{f}(s)|^2 - 4 \sum_{\substack{N/M_N \leq s, t \\ 1 \leq |s-t| \leq N-1 \\ N \leq \max(s, t)}} (N - |s-t|) \hat{f}(s) \hat{f}(t) - 4 \sum_{\substack{N/M_N \leq s, t \leq N-1 \\ N+1 \leq s+t}} ((s+t) - N) \hat{f}(s) \hat{f}(t).$$

The first term on the r.h.s. of (2.18) is equal to $2(V_N - V_{\lfloor N/M_N \rfloor})$ and is $o(V_N)$ by (2.16). The last two terms on the r.h.s. of (2.18) are $o(V_N)$ by Lemma 2.1. Finally, the second term on the r.h.s. of (2.18) is bounded from above by

$$(2.19) \quad 4N^2 \sum_{N \leq s} |\hat{f}(s)|^2 \leq 2 \sum_{j=1}^{\infty} \frac{1}{j^2} (V_{(j+1)N} - V_{jN}) = o(V_N),$$

where the last estimate follows from (1.6). This completes the proof of Lemma 2.2.

To finish the proof of the theorem, we have to show that the first term on the r.h.s. of (2.15) converges in distribution to a standard Gaussian random variable. To do this, we first show that the first $\lfloor M_N/2 \rfloor$ moments of

$$(2.20) \quad \frac{2}{\sqrt{V_N}} \sum_{k=1}^{\lfloor N/M_N \rfloor} \hat{f}(k)(|t_{N,k}|^2 - \mathbb{E}|t_{N,k}|^2)$$

coincide with the first $\lfloor M_N/2 \rfloor$ moments of

$$(2.21) \quad \frac{2}{\sqrt{V_N}} \sum_{k=1}^{\lfloor N/M_N \rfloor} \hat{f}(k) k (\varphi_k - 1),$$

where φ_k , $k \geq 1$, are i.i.d. exponential random variables.

Lemma 2.3. *Let m be a positive integer such that $1 \leq m < \frac{M_N}{2}$. Then*

$$(2.22) \quad \mathbb{E} \left(\frac{2}{\sqrt{V_N}} \sum_{k=1}^{\lfloor N/M_N \rfloor} \hat{f}(k) (|t_{N,k}|^2 - \mathbb{E}|t_{N,k}|^2) \right)^m = \mathbb{E} \left(\frac{2}{\sqrt{V_N}} \sum_{k=1}^{\lfloor N/M_N \rfloor} \hat{f}(k) k (\varphi_k - 1) \right)^m.$$

Proof of Lemma 2.3 The formula (2.22) follows from the identity

$$(2.23) \quad \mathbb{E} \prod_{i=1}^l |t_{N,k_i}|^2 = \mathbb{E} \prod_{i=1}^l k_i \varphi_{k_i},$$

provided

$$(2.24) \quad 2 \sum_{i=1}^l |k_i| \leq N, \quad 0 < k_1, \dots, k_l.$$

(2.24) was established in [6] and [21] (see also [13]) where it was shown that a large number of the joint moments/cumulants of

$$k^{-1/2} t_{N,k} = k^{-1/2} \text{Tr}(U^k), \quad k \geq 1,$$

coincide with the corresponding joint moments/cumulants of a sequence of i.i.d. standard complex Gaussian random variables. Namely, if we denote by $\kappa(t_{N,k_1}, \dots, t_{N,k_n})$ the joint cumulant of $\{t_{N,k_j}, 1 \leq j \leq n\}$ then

$$(2.25) \quad \kappa(t_{N,k_1}, \dots, t_{N,k_n}) = 0$$

if at one of the following two conditions is satisfied:

- (i) $n \geq 1$, and $\sum_{j=1}^n k_j \neq 0$
- (ii) $n > 2$, $\sum_{j=1}^n k_j = 0$, and $\sum_{j=1}^n |k_j| \leq N$.

In addition, $\kappa(t_{N,k}, t_{N,-k}) = \min(|k|, N)$. We refer the reader to Lemma 5.2 of [1] for the details. Taking into account that the absolute value squared of a standard complex Gaussian random variable is distributed according to the exponential law we obtain (2.23-2.24). This finishes the proof of Lemma 2.3. Now, the proof of Theorem 1.2 immediately follows from the following standard lemma.

Lemma 2.4. *For any $t \in \mathbb{R}$*

$$(2.26) \quad \mathbb{E} \exp \left(\frac{t}{\sqrt{\sum_{k=1}^N a_k^2}} \sum_{k=1}^N a_k (\varphi_k - 1) \right) \rightarrow \exp(t^2/2)$$

as $N \rightarrow \infty$, provided

$$(2.27) \quad \sum_{k=1}^{\infty} a_k^2 = \infty \quad \text{and} \quad \max_{1 \leq k \leq N} |a_k| = o\left(\sqrt{\sum_{k=1}^N a_k^2}\right).$$

Proof of Lemma 2.4 The result of Lemma 2.4 follows from standard direct computations using independence of φ_k 's.

Setting $a_k = \hat{f}(k)k$, $k \geq 1$ and applying Lemma 2.4, we obtain the convergence of the exponential moment (and, hence, all moments) of $\sqrt{\frac{2}{V_N}} \sum_{k=1}^{\lfloor N/M_N \rfloor} \hat{f}(k)k(\varphi_k - 1)$ to that of a standard real Gaussian random variable. Thus, both

$$\sqrt{\frac{2}{V_N}} \sum_{k=1}^{\lfloor N/M_N \rfloor} \hat{f}(k)k(\varphi_k - 1)$$

and

$$\sqrt{\frac{2}{V_N}} \sum_{k=1}^{\lfloor N/M_N \rfloor} \hat{f}(k)(|t_{N,k}|^2 - \mathbb{E}|t_{N,k}|^2)$$

converge in distribution to a standard real Gaussian random variable. Since the second term in (2.15) goes to 0 in L^2 we conclude that Theorem 1.2 is proven.

REFERENCES

- [1] Aguirre A., Soshnikov A. and Sumpter J. *Pair Dependent Linear Statistics for $C\beta E$* [arXiv:1912.07110](https://arxiv.org/abs/1912.07110) [math.PR], Dec 2019
- [2] Baker, T.H., Forrester, P. J., *Finite-N Fluctuation Formulas for Random Matrices*. J. Stat. Phys., **88**, (1997), 1371–1386.
- [3] Bekerman, F., Lodhia, A., *Mesoscopic Central Limit Theorem for general β -ensembles*. Ann. Inst. H. Poincaré Prob. Stat. **54**, (2018), 1917–1938.
- [4] Bingham, N.H., Goldie, C.M., Teugels, J.L., *Regular Variation*. Cambridge University Press, 1987.
- [5] Diaconis, P., Evans, S.N., *Linear functionals of eigenvalues of random matrices*. Trans. Amer. Math. Soc. **353**, (2001), 2615–2633.
- [6] Diaconis, P., Shahshahani, M. *On eigenvalues of random matrices*. J. Appl. Probab., **31A**, (1994), 49–62.
- [7] Dyson, F.J. *Statistical theory of the energy levels of complex systems. I* J. Math. Phys. **3**, (1962), 140–156.
- [8] Dyson, F.J. *Statistical theory of the energy levels of complex systems. I* J. Math. Phys. **3**, (1962), 166–175.
- [9] Dyson, F.J. *Statistical theory of the energy levels of complex systems. I* J. Math. Phys. **3**, (1962), 1191–1198.
- [10] Dyson, F.J. *Correlations between the eigenvalues of a random matrix*. Comm. Math. Phys., **19(3)**, (1970) 235–250.

- [11] Johansson, K. *On Szego's Asymptotic Formula for Toeplitz Determinants and Generalizations*. Duke Math. J. **91** (1988), 151–204.
- [12] Johansson, K. *On Random Matrices from the Compact Classical Groups*. Ann. Math (2) **145**, (1997), 519–545.
- [13] Jiang, T., Matsumoto, S. *Moments of Traces of Circular β -ensembles*. Ann. Probab. **43**, Number 6 (2015), 3279–3336
- [14] Hughes, C.P., Keating, J.P., O'Connell, N. *On the characteristic polynomial of a random unitary matrix*. Comm. Math. Phys., **220(2)**, (2001), 429–451.
- [15] Killip, R., Nenciu, I. *Matrix models for circular ensembles.*, Int. Math. Res. Not. **50**, (2004), 2665–2701.
- [16] Lambert, G. *Mesoscopic central limit theorem for the circular beta-ensembles and applications*. available at [arXiv:1902.06611](https://arxiv.org/abs/1902.06611) math.PR.
- [17] Meckes, E. S., Meckes, M. W. *Self-similarity in the circular unitary ensemble*. Discrete Anal. (2016), paper No.9, 14pp.
- [18] Montgomery, H.L. *On pair correlation of zeros of the zeta function*. Proc. Sympos. Pure Math., **24**, (1973), 181–193.
- [19] Montgomery, H.L. *Distribution of the zeros of the Riemann zeta function.*, Proc. Internat. Congr. Math., **1**, Vancouver, BC (1974), 379–381.
- [20] Paquette, E., Zeitouni, O. *The Maximum of the CUE Field*. IMRN, **16**, (2018), 5028–5119
- [21] Soshnikov, A. *Central Limit Theorem for local linear statistics in classical compact groups and related combinatorial identities*. Ann. Probab. **28**, (2000), 1353–1370
- [22] Soshnikov, A. *Level spacings distribution for large random matrices: Gaussian fluctuations*. Ann. Math (2) **148**, (1998), 573–617.
- [23] Webb, C., *Linear statistics of the circular ensemble, Stein's method, and circular Dyson Brownian motion.*, E. J. Probab., **20**, (2015), No. 104, 21pp,
- [24] Witte N.S., Forrester, P.J. *Moments of the Gaussian Beta Ensembles Ensembles and the large- N expansion of the densities*. J. Math. Phys., **55**, (2014),

UNIVERSITY OF CALIFORNIA AT DAVIS, DEPARTMENT OF MATHEMATICS, 1 SHIELDS AVENUE,
DAVIS CA 95616, UNITED STATES OF AMERICA

E-mail address: aaguirre@ucdavis.edu

UNIVERSITY OF CALIFORNIA AT DAVIS, DEPARTMENT OF MATHEMATICS, 1 SHIELDS AVENUE,
DAVIS CA 95616, UNITED STATES OF AMERICA

E-mail address: soshniko@math.ucdavis.edu