

Genuine Multipartite Nonlocality Is Intrinsic to Quantum Networks

Patricia Contreras-Tejada,¹ Carlos Palazuelos,^{2,1} and Julio I. de Vicente³

¹*Instituto de Ciencias Matemáticas, E-28049 Madrid, Spain*

²*Departamento de Análisis Matemático y Matemática Aplicada,
Universidad Complutense de Madrid, E-28040 Madrid, Spain*

³*Departamento de Matemáticas, Universidad Carlos III de Madrid, E-28911, Leganés (Madrid), Spain*

Quantum entanglement and nonlocality are inextricably linked. However, while entanglement is necessary for nonlocality, it is not always sufficient in the standard Bell scenario. We derive sufficient conditions for entanglement to give rise to genuine multipartite nonlocality in networks. We find that any network where the parties are connected by bipartite pure entangled states is genuine multipartite nonlocal, independently of the amount of entanglement in the shared states and of the topology of the network. As an application of this result, we also show that all pure genuine multipartite entangled states are genuine multipartite nonlocal in the sense that measurements can be found on finitely many copies of any genuine multipartite entangled state to yield a genuine multipartite nonlocal behaviour. Our results pave the way towards feasible manners of generating genuine multipartite nonlocality using any connected network.

Correlations between quantum particles may be much stronger than those between classical particles. Their applications are manifold: cryptography [1, 2], randomness extraction, amplification and certification [3], communication complexity reduction [4], etc., and the study of these *nonlocal* correlations has led to the growing field of device-independent quantum information processing [5–7] (see also Ref. [8]).

While bipartite nonlocality has been well researched in the past three decades, much less is known about the multipartite case. Still, correlations in quantum multi-component systems have gained increasing attention recently, with applications in multiparty cryptography [9], the understanding of condensed matter physics [10, 11], and the development of quantum networks [12–18], particularly for quantum computation [19–21] and correlating particles which never interacted [22, 23].

A necessary condition for nonlocality is quantum entanglement. Indeed, this is one reason why entangled states are useful for communication-related tasks. However, not all entangled states are nonlocal: some bipartite entangled states only yield local distributions [24, 25]. Still, for pure bipartite states, entanglement *is* sufficient for nonlocality, which is the content of Gisin’s theorem [26, 27], and multipartite entangled pure states are never fully local [28, 29]. Interestingly, distributing certain bipartite entangled states in certain multipartite networks yields nonlocality even if the involved states are individually local [12, 14, 30–33].

Multipartite nonlocality is in principle harder to generate than bipartite nonlocality. By exploring the relationship between entanglement and nonlocality in the multipartite regime, in this Letter we show that networks simplify the job considerably: distributing arbitrarily low node-to-node entanglement is sufficient to observe truly multipartite nonlocal effects involving all parties in the network independently of its geometry. Added to its practical consequences for applications, this fact points

to a deep property of quantum networks.

The multipartite setting has a richer structure than the bipartite one, as different forms of entanglement and nonlocality can be identified. Full separability (full locality) refers to systems that do not display any form of entanglement (locality) whatsoever. However, falsifying these models need not imply truly multipartite quantum correlations since spreading them among two parties is sufficient. Hence, a *genuine multipartite* notion which inextricably relates all parties together is more often considered. Here, a state is genuine multipartite entangled (GME) if it is not a tensor product of states of two subsets of parties, M and its complement \overline{M} , i.e. of the form $|\psi\rangle = |\psi_M\rangle \otimes |\psi_{\overline{M}}\rangle$, or a convex combination of such states $|\psi\rangle\langle\psi|$ across all bipartitions. Analogously, a probability distribution $\{P(\alpha_1\alpha_2\dots\alpha_n|\chi_1\chi_2\dots\chi_n)\}_{\alpha_1,\dots,\alpha_n,\chi_1,\dots,\chi_n}$ (with input χ_i and output α_i for party i) which is not of the form

$$\begin{aligned} &P(\alpha_1\alpha_2\dots\alpha_n|\chi_1\chi_2\dots\chi_n) \\ &= \sum_{M \subseteq [n]} \sum_{\lambda} q_M(\lambda) P_M(\{\alpha_i\}_{i \in M} | \{\chi_i\}_{i \in M}, \lambda) \\ &\quad \times P_{\overline{M}}(\{\alpha_i\}_{i \in \overline{M}} | \{\chi_i\}_{i \in \overline{M}}, \lambda), \end{aligned} \quad (1)$$

where $q_M(\lambda) \geq 0 \forall \lambda, M$, $\sum_{\lambda, M} q_M(\lambda) = 1$ and $[n] := \{1, \dots, n\}$, is genuine multipartite nonlocal (GMNL) [34–36], and a state is GMNL if measurements giving rise to a GMNL distribution exist. The original definition [34] leaves the distributions $P_M, P_{\overline{M}}$ in equation (1) unrestricted; however, this has been shown to lead to operational problems [37–42]. Hence, like most recent works on the topic, we assume these distributions are nonsignalling, which captures most physical situations better [43, 44]. This means the marginal distributions for any subset of parties are independent of the inputs outside this subset, which is guaranteed if this holds for the marginals corresponding to ignoring just one party

[45]:

$$\sum_{\alpha_j} P_M(\{\alpha_i\}_{i \in M, i \neq j}, \alpha_j | \{\chi_i\}_{i \in M, i \neq j}, \chi_j, \lambda) = \sum_{\alpha_j} P_M(\{\alpha_i\}_{i \in M, i \neq j}, \alpha_j | \{\chi_i\}_{i \in M, i \neq j}, \chi'_j, \lambda), \quad (2)$$

for all $\lambda, \chi_j \neq \chi'_j$ and all parties j , and similarly for $P_{\overline{M}}$.

In this Letter we show that the nonlocality arising from networks of bipartite pure entangled states is a generic property and manifests in its strongest form, GMNL. Specifically, we obtain that any connected network of bipartite pure entangled states is GMNL. It was already known that a star network of maximally entangled states is GMNL [12], but we provide a full, qualitative generalisation of this result by making it independent of both the amount of entanglement shared and the network topology. Thus, we show GMNL is an intrinsic property of networks of pure bipartite entangled states.

Further, there are known mixed GME states that are not GMNL [46, 47]—some are even fully local [48]. Still, it is not known whether Gisin’s theorem extends to the genuine multipartite regime. Recent results show that, for pure n -qubit symmetric states [49] and all pure 3-qubit states [50], GME implies GMNL (at the single-copy level) [51]. Our result above shows that all pure GME states that have a network structure are GMNL; interestingly, we further apply this property to establish a second result: all pure GME states are GMNL in the sense that measurements can be found on finitely many copies of any GME state to yield a GMNL behaviour. We thus tighten the relationship between multipartite entanglement and nonlocality.

Our construction exploits the fact that the set of non-GME states is not closed under tensor products, i.e. GME can be superactivated by taking tensor products of states that are unentangled across different bipartitions. Thus, GME can be achieved by distributing bipartite entangled states among different pairs of parties. To obtain our results, we extend the superactivation property [52–54] from the level of states to that of probability distributions, i.e. GMNL can be superactivated by taking Cartesian products of probability distributions that are local across different bipartitions. In fact, when considering copies of quantum states, we only consider local measurements performed on each copy separately, thus pointing at a stronger notion of superactivation to achieve GMNL.

Definitions and preliminaries We consider distributions arising from GME states, and ask whether they satisfy (1). The set of distributions of the form (1) is a polytope: indeed, the set of local distributions across each bipartition $M|\overline{M}$ is a polytope, and convex combinations preserve that structure. We call this n -partite

polytope \mathcal{B}_n . We call an inequality

$$\sum_{\substack{\alpha_i \chi_i \\ i \in [n]}} c_{\alpha_1 \dots \alpha_n \chi_1, \dots, \chi_n} P(\alpha_1 \dots \alpha_n | \chi_1 \dots \chi_n) \leq c_0 \quad (3)$$

which holds for all P of the form (1) a GMNL inequality.

We use results from Ref. [55] to lift inequalities to account for more parties, inputs and outputs. They consider the fully local polytope \mathcal{L} , which only includes distributions

$$P(\alpha\beta|\chi v) = \sum_{\lambda} q(\lambda) P_A(\alpha|\chi, \lambda) P_B(\beta|v, \lambda) \quad (4)$$

where each party may have different numbers of inputs and outputs (more parties may be considered by adding more distributions correlated only by λ). Polytope \mathcal{B}_n includes convex combinations of distributions that are local across different bipartitions $M|\overline{M}$ of the parties, but the lifting results in [55] still hold. Indeed, to check an inequality holds for a polytope, it is sufficient by convexity to check the extremal points. As all extremal points in \mathcal{B}_n are contained in some polytope \mathcal{L} , lifting results for \mathcal{L} can be straightforwardly extended to \mathcal{B}_n .

We also use the EPR2 decomposition [56] and its multipartite extension [57]: any distribution P can be expressed (nonuniquely) as

$$P(\alpha_1 \dots \alpha_n | \chi_1 \dots \chi_n) = \sum_{M \subseteq [n]} p_L^M P_L^M(\alpha_1 \dots \alpha_n | \chi_1 \dots \chi_n) + p_{NS} P_{NS}(\alpha_1 \dots \alpha_n | \chi_1 \dots \chi_n) \quad (5)$$

where $\sum_{M \subseteq [n]} p_L^M + p_{NS} = 1$, P_L^M is local across $M|\overline{M}$ (i.e. satisfies equation (4) with parties grouped as per $M|\overline{M}$), and P_{NS} is nonsignalling. P is GMNL if all such decompositions have $p_{NS} > 0$, and fully GMNL if all such decompositions have $p_{NS} = 1$. A state ρ is fully GMNL if, $\forall \varepsilon > 0$, there exist local measurements giving rise to some P such that any decomposition (3) has $p_{NS} > 1 - \varepsilon$. Bipartite distributions and states may be nonlocal or fully nonlocal [58] analogously.

GMNL from bipartite entanglement Our first result shows that any connected network of pure bipartite entanglement (see Figure 1) is GMNL.

Theorem 1. *Any connected network of bipartite pure entangled states is GMNL.*

We now outline the proof for a tripartite network where A_1 is entangled to each of A_2 and A_3 , and leave the general case to [59]. Since it turns out sufficient to measure individually on each party’s different particles (see Figure 1 for the n -partite structure), the shared distribution $P(a_1^1 a_1^2, a_2^1, a_3^1 | x_1^1 x_1^2, x_2^1, x_3^1)$ takes the form

$$P_1(a_1^1 a_2^1 | x_1^1 x_2^1) P_2(a_1^2 a_3^2 | x_1^2 x_3^2) \quad (6)$$




Figure 1. Connected network of bipartite entanglement. For each $i \in [n]$, party A_i has input x_i^k and output a_i^k on the particle at edge k . Particles connected by an edge are entangled.

where parties A_i, A_j are connected by edge k (we label vertices and edges independently), and $P_k(a_i^k a_j^k | x_i^k x_j^k)$ is the distribution arising from the state at edge k .

The proof considers three cases, depending on whether the shared states are maximally entangled. If none are, we devise inequalities to detect bipartite nonlocality at each edge of the network, and combine them to form a multipartite inequality. Then, we find measurements on the shared states to violate it. If both states are maximally entangled, existing results show the network is fully GMNL [12, 57]. Combining these two cases for a heterogeneous network completes the proof.

To prove the first case, we take bipartite inequalities between A_1 and each other party, lift them to three parties and combine them using Refs. [55, 60], to obtain the following GMNL inequality:

$$\begin{aligned}
 I_3 &= I^1 + I^2 + P(00, 0, 0|00, 0, 0) \\
 &\quad - \sum_{a_1^2=0,1} P(0a_1^2, 0, 0|00, 0, 0) \\
 &\quad - \sum_{a_1^1=0,1} P(a_1^1 0, 0, 0|00, 0, 0) \leq 0.
 \end{aligned} \tag{7}$$

Here,

$$\begin{aligned}
 I^1 &= \sum_{a_1^2=0,1} [P(0a_1^2, 0, 0|00, 0, 0) - P(0a_1^2, 1, 0|00, 1, 0) \\
 &\quad - P(1a_1^2, 0, 0|10, 0, 0) - P(0a_1^2, 0, 0|10, 1, 0)] \leq 0;
 \end{aligned} \tag{8}$$

$$\begin{aligned}
 I^2 &= \sum_{a_1^1=0,1} [P(a_1^1 0, 0, 0|00, 0, 0) - P(a_1^1 0, 0, 1|00, 0, 1) \\
 &\quad - P(a_1^1 1, 0, 0|01, 0, 0) - P(a_1^1 0, 0, 0|01, 0, 1)] \leq 0
 \end{aligned} \tag{9}$$

are liftings of

$$I = P(00|00) - P(01|01) - P(10|10) - P(00|11) \leq 0 \tag{10}$$

to three parties with A_1 having 4 inputs and 4 outputs. Inequality (10) is equivalent to the CHSH inequality [61] for nonsignalling distributions [60]. Thus, inequalities (8), (9) are satisfied by distributions that are local across $A_1|A_2$ and $A_1|A_3$ respectively. To see that equation (7) is a GMNL inequality it is sufficient to check it holds for distributions that are local across some bipartition. This is straightforwardly done by observing the cancellations that occur when I^1 or I^2 are ≤ 0 .

Since both states are less-than-maximally entangled, A_1 can satisfy Hardy's paradox [62, 63] with each other party, achieving

$$P_k(00|00) > 0 = P_k(01|01) = P_k(10|10) = P_k(00|11) \tag{11}$$

for both k (the proof for qubits in Refs. [62, 63] is extended to qudits by measuring on a two-dimensional subspace, see Ref. [59]). Then, each negative term in I^1 and I^2 is zero, as

$$\sum_{a_1^2=0,1} P(0a_1^2, 1, 0|00, 1, 0) = P_1(01|01) \sum_{a_1^1=0,1} P_2(a_1^2 0|00) \tag{12}$$

and similarly for the others. Hence, only

$$P(00, 0, 0|00, 0, 0) = P_1(00|00)P_2(00|00) > 0 \tag{13}$$

survives, violating the inequality.

If, instead, $A_1 A_2$ share a maximally entangled state, and $A_2 A_3$ share a less-than-maximally entangled state, then $A_1 A_3$ can measure so that P_2 satisfies Hardy's paradox; hence $\exists \varepsilon > 0$ such that its local component in any EPR2 decomposition satisfies

$$p_{L,2} \leq 1 - \varepsilon. \tag{14}$$

Since the maximally entangled state is fully nonlocal [64], for this ε , $A_1 A_2$ can measure such that any EPR2 decomposition of P_1 satisfies

$$p_{L,1} < \varepsilon. \tag{15}$$

Then, we assume for a contradiction that $P(a_1^1 a_1^2, a_2^1, a_3^2 | x_1^1 x_1^2, x_2^1, x_2^2)$ is not GMNL and decompose it in its bipartite splittings,

$$\begin{aligned}
 &P(a_1^1 a_1^2, a_2^1, a_3^2 | x_1^1 x_1^2, x_2^1, x_2^2) \\
 &= \sum_{\lambda} (p_L(\lambda) P_{A_1 A_2}(a_1^1 a_1^2, a_2^1 | x_1^1 x_1^2, x_2^1, \lambda) P_{A_3}(a_3^2 | x_2^2, \lambda) \\
 &\quad + q_L(\lambda) P_{A_1 A_3}(a_1^1 a_1^2, a_3^2 | x_1^1 x_1^2, x_2^2, \lambda) P_{A_2}(a_2^1 | x_2^1, \lambda) \\
 &\quad + r_L(\lambda) P_{A_1}(a_1^1 a_1^2 | x_1^1 x_1^2, \lambda) P_{A_2 A_3}(a_2^1, a_3^2 | x_2^1, x_2^2, \lambda))
 \end{aligned} \tag{16}$$




Figure 2. Element $i \in [n-1]$ of the star network of bipartite entanglement created from a GME state $|\Psi\rangle$. Parties $\{B_j\}_{j \in [n-1], j \neq i}$ have already measured $|\Psi\rangle$ and are left unentangled. Alice and party B_i share a pure bipartite entangled state. Alice has input x_i and output a_i while each party B_j , $j \in [n-1]$, has input y_j^i and output b_j^i .

where $\sum_{\lambda} [p_L(\lambda) + q_L(\lambda) + r_L(\lambda)] = 1$.

Summing equation (16) over a_1^2, a_3^2 and using equation (6), we get an EPR2 decomposition of P_1 with local components q_L, r_L . By equation (26), this entails $\sum_{\lambda} [q_L(\lambda) + r_L(\lambda)] < \varepsilon$, so

$$\sum_{\lambda} p_L(\lambda) > 1 - \varepsilon. \quad (17)$$

Summing, instead, equation (16) over a_1^1, a_2^1 , we obtain an EPR2 decomposition of P_2 whose only nonnegligible component, $\sum_{\lambda} p_L(\lambda)$, is local in $A_1|A_3$, contradicting equation (24). Therefore, P must be GMNL.

GMNL from GME By Theorem 1, a star network whose central node shares pure entanglement with all others is GMNL. We now ask whether all GME states are GMNL (i.e. the genuine multipartite extension of Gisin's theorem). We show $(n-1)$ copies of any pure GME n -partite state suffice to generate n -partite GMNL. We do this by generating a distribution from these copies that mimics the star network configuration.

Theorem 2. *Any GME state $|\Psi\rangle \in \mathcal{H}_1 \otimes \dots \otimes \mathcal{H}_n \cong (\mathbb{C}^d)^{\otimes n}$ is such that $|\Psi\rangle^{\otimes(n-1)}$ is GMNL.*

The full proof is given in [59], and we presently outline the tripartite case. Hence, we consider two copies of the state. For each copy, we derive measurements for Bob1 and Bob2 that leave Alice bipartitely entangled with Bob2 and Bob1 respectively. This yields a network as in equation (6) but postselected on the inputs and outputs of these measurements. We generalise Theorem 1 to show this network is also GMNL.

For $i, j = 1, 2$, on copy i , B_j 's measurements have input y_j^i and output b_j^i and Alice's measurement has input x_i and output a_i . We denote B_j 's inputs and outputs in terms of their digits as $v_j = y_j^1 y_j^2$ and $\beta_j = b_j^1 b_j^2$. Then, after measurement, the parties share a distribution

$$P(\alpha\beta_1\beta_2|\chi v_1 v_2) = P_1(a_1, b_1^1 b_2^1 | x_1, y_1^1 y_2^1) P_2(a_2, b_1^2 b_2^2 | x_2, y_1^2 y_2^2). \quad (18)$$

For each $i, j = 1, 2$, $i \neq j$, we assume B_j uses input 0_j^i and output 0_j^i to project the i th copy of $|\Psi\rangle$ onto $|\phi_i\rangle_{AB_i}$, as shown in Figure 2 for n parties. Then, Refs. [28, 29] and a continuity argument serve to show we only have two possibilities for each i : either there exists an input and output per party such that $|\phi_i\rangle_{AB_i}$ is less-than-maximally entangled, or there exists an input per party such that, for all outputs, $|\phi_i\rangle_{AB_i}$ is maximally entangled. In each case we generalise the proof in Theorem 1 to show $|\Psi\rangle^{\otimes 2}$ is GMNL.

If both $|\phi_i\rangle_{AB_i}$, $i = 1, 2$ are less-than-maximally entangled, we use the following expression, which is a GMNL inequality by the same reasoning as in Theorem 1:

$$I_3 = \sum_{i=1}^2 I^i + P(00, 00, 00|00, 00, 00) - \sum_{i=1}^2 \sum_{\substack{a_j, b_j^i, \\ b_j^j=0,1, \\ j \neq i}} P(0_i a_j, 0_i^i b_j^i, 0_j^i b_j^j | 0_i 0_j, 0_i^i 0_j^i, 0_j^i 0_j^j) \leq 0, \quad (19)$$

where

$$I^i = \sum_{\substack{a_j, b_j^i, b_j^j=0,1, \\ j \neq i}} \left[P(0_i a_j, 0_i^i b_j^i, 0_j^i b_j^j | 0_i 0_j, 0_i^i 0_j^i, 0_j^i 0_j^j) - P(0_i a_j, 1_i^i b_j^i, 0_j^i b_j^j | 0_i 0_j, 1_i^i 0_j^i, 0_j^i 0_j^j) - P(1_i a_j, 0_i^i b_j^i, 0_j^i b_j^j | 1_i 0_j, 0_i^i 0_j^i, 0_j^i 0_j^j) - P(0_i a_j, 0_i^i b_j^i, 0_j^i b_j^j | 1_i 0_j, 1_i^i 0_j^i, 0_j^i 0_j^j) \right]. \quad (20)$$

Evaluating the inequality on the distribution (39), we find again that all negative terms in each I^i can be sent to zero. For each i we get, for example,

$$\sum_{\substack{a_j, b_j^i, b_j^j \\ =0,1}} P(0_i a_j, 1_i^i b_j^i, 0_j^i b_j^j | 0_i 0_j, 1_i^i 0_j^i, 0_j^i 0_j^j) = P_i(0_i 1_i^i 0_j^i | 0_i 1_i^i 0_j^i) \quad (21)$$

as the sum over P_j is 1. But, conditioned on B_j 's input and output being 0_j^i , parties AB_i can measure so P_i satisfies Hardy's paradox, hence this term is zero, and similarly for the other two negative terms. This means all terms in I_3 are zero except $P(00, 00, 00|00, 00, 00) > 0$, violating the inequality. Therefore, $|\Psi\rangle^{\otimes 2}$ is GMNL.

If, for both $i = 1, 2$, there exists a local measurement for party B_j , $j \neq i$ such that, for *all* outputs, $|\Psi\rangle$ is projected onto a maximally entangled state $|\phi_i\rangle_{AB_i}$, then $|\Psi\rangle$ satisfies Theorem 2 in Ref. [57], so $|\Psi\rangle$ itself is GMNL. Therefore so is $|\Psi\rangle^{\otimes 2}$.

Finally, if $|\phi_1\rangle_{AB_1}$ is maximally entangled for all of B_2 's outputs, and $|\phi_2\rangle_{AB_2}$ is less-than-maximally entangled, using Refs. [28, 57] we deduce that the bipartite EPR2 components of $P_{1,2}$ across $A|B_{1,2}$ respectively are bounded like in Theorem 1. That is, $\exists \varepsilon > 0$ such that the local component of any EPR2 decomposition across $A|B_2$ satisfies

$$p_{L,2}^{A|B_2} \leq 1 - \varepsilon \quad (22)$$

and, given this ε , parties AB_1 can measure locally such that all bipartite EPR2 decompositions across $A|B_1$ have a local component

$$p_{L,1}^{A|B_1} < \varepsilon. \quad (23)$$

Then, we assume $P(\alpha\beta_1\beta_2|\chi v_1 v_2)$ is not GMNL and decompose it in local terms across different bipartitions, like in equation (16) in Theorem 1. Summing over a_2, b_j^2 , $j = 1, 2$ gives an EPR2 decomposition of P_1 whose local components can be bounded using equation (23). Summing over a_1, b_j^1 , $j = 1, 2$ instead gives an EPR2 decomposition of P_2 . But the bound on the local component of P_1 entails a bound on that of P_2 which contradicts equation (22), proving P is GMNL.

Conclusions We have shown that GMNL can be obtained by distributing arbitrary pure bipartite entanglement, which paves the way towards feasible generation of GMNL from any network. In fact, our results imply that, given a set of nodes, distributing entanglement in the form of a tree is sufficient to observe GMNL. In practical applications, the entanglement shared by the nodes would unavoidably degrade to mixed-state form. By continuity, the GMNL in the networks of pure bipartite entanglement considered here must be robust to some noise. Quantifying this tolerance is interesting for future work.

Further, we have shown that a tensor product of finitely many GME states is always GMNL. The question whether all single-copy pure GME states are GMNL remains open.

The assumption that the distributions $P_M, P_{\overline{M}}$ are nonsignalling in the GMNL definition is physically natural. Still, removing it raises the stakes to achieve nonlocality, and establishing analogous results with the stronger definition is an open question.

Very recently, Ref. [65] proposed the concept of “genuine network entanglement”, a stricter notion than GME which rules out states which are a tensor product of non-GME states. One might hope that states that are GME but not genuine network entangled might be detected device independently by not passing GMNL tests. However,

our results show this will not work. Any distribution of pure bipartite states, even with arbitrarily weak entanglement, always displays GMNL as long as all parties are connected. This further motivates searching for an analogous concept of genuine network nonlocality that may detect genuine network entanglement.

The authors thank Aleksander M. Kubicki and Elie Wolfe for enlightening discussions. This research was funded by the Spanish MINECO through Grant No. MTM2017-88385-P and by the Comunidad de Madrid through Grant No. QUITEMAD-CMS2018/TCS-4342. We also acknowledge funding from the Spanish MINECO, through the “Severo Ochoa Programme for Centres of Excellence in R&D” SEV-2015-0554 and from the Spanish National Research Council, through the “Ayuda extraordinaria a Centros de Excelencia Severo Ochoa” 20205CEX001 (PCT and CP) and the Spanish MINECO Grant No. MTM2017-84098-P (JIdV). PCT is grateful for the hospitality of Perimeter Institute where part of this work was carried out. Research at Perimeter Institute is supported in part by the Government of Canada through the Department of Innovation, Science and Economic Development Canada and by the Province of Ontario through the Ministry of Economic Development, Job Creation and Trade.

Supplemental material

We prove Theorems 1 and 2 in the main text. For the reader's convenience, we also restate some definitions and the results.

We say a probability distribution $\{P(\alpha_1\alpha_2\dots\alpha_n|\chi_1\chi_2\dots\chi_n)\}_{\alpha_1,\dots,\alpha_n,\chi_1,\dots,\chi_n}$ (with input χ_i and output α_i for party i) is genuine multipartite nonlocal (GMNL) if it cannot be written in the form

$$P(\alpha_1\alpha_2\dots\alpha_n|\chi_1\chi_2\dots\chi_n) = \sum_M \sum_\lambda q_M(\lambda) P_M(\{\alpha_i\}_{i \in M} | \{\chi_i\}_{i \in M}, \lambda) P_{\overline{M}}(\{\alpha_i\}_{i \in \overline{M}} | \{\chi_i\}_{i \in \overline{M}}, \lambda), \quad (1)$$

where here and in the rest of the paper we take $\emptyset \neq M \subsetneq [n]$, and we ignore duplicate bipartitions (e.g. by assuming M always contains party A_1), $q_M(\lambda) \geq 0$ for each λ, M , $\sum_{\lambda, M} q_M(\lambda) = 1$ and the distributions $P_M, P_{\overline{M}}$ on each bipartition are nonsignalling. We will use the notation $[n] := \{1, \dots, n\}$ throughout. A state is GMNL if there exist measurements which give rise to a distribution that cannot be written as (1).

Given an inequality

$$\sum_{\substack{\alpha_i \chi_i \\ i \in [n]}} c_{\alpha_1 \dots \alpha_n \chi_1, \dots, \chi_n} P(\alpha_1 \dots \alpha_n | \chi_1 \dots \chi_n) \leq c_0 \quad (2)$$

which holds for all distributions P of the form (1), we refer to it as a GMNL inequality. The set of points $P(\alpha_1 \dots \alpha_n | \chi_1 \dots \chi_n)$ which satisfy it with equality is a *face* of the polytope \mathcal{B}_n of n -partite distributions (1). Inequalities are said to *support* faces of the polytope. Faces $F \neq \mathcal{B}_n$ of maximal dimension are *facets*, and inequalities which support facets are called *facet inequalities*.

The multipartite EPR2 decomposition [56, 57] of a probability distribution P is

$$\begin{aligned} P(\alpha_1 \dots \alpha_n | \chi_1 \dots \chi_n) \\ = \sum_M p_L^M P_L^M(\alpha_1 \dots \alpha_n | \chi_1 \dots \chi_n) + p_{NS} P_{NS}(\alpha_1 \dots \alpha_n | \chi_1 \dots \chi_n), \end{aligned} \quad (3)$$

where $p_L^M \geq 0$ for every M , $p_{NS} \geq 0$ and

$$\sum_M p_L^M + p_{NS} = 1, \quad (4)$$

P_L^M is local across the bipartition $M|\overline{M}$, and P_{NS} is nonsignalling. We are interested in decompositions which maximise the local EPR2 components, in order to deduce properties about the distributions. For a distribution P , we define

$$EPR2(P) = \max \left\{ \sum_M p_L^M : P = \sum_M p_L^M P_L^M + p_{NS} P_{NS}, \sum_M p_L^M + p_{NS} = 1 \right\} \quad (5)$$

and, for a state ρ , we define (with a slight abuse of notation)

$$EPR2(\rho) = \inf \left\{ EPR2(P) : P = \text{tr} \left(\bigotimes_{i=1}^n E_{\alpha_i | \chi_i}^i \rho \right) \right\}, \quad (6)$$

where the infimum is taken over local measurements $E_{\alpha_i | \chi_i}^i$ on each particle such that

$$E_{\alpha_i | \chi_i}^i \geq 0 \forall \alpha_i, \chi_i, \sum_{\alpha_i} E_{\alpha_i | \chi_i}^i = 1 \forall \chi_i, \forall i \in [n], \quad (7)$$

with any number of inputs and outputs. Then, a distribution P or a state ρ are GMNL if $EPR2(\cdot) < 1$, while they are fully GMNL if $EPR2(\cdot) = 0$. When considering bipartite distributions and states, the analogous property is termed full-nonlocality. Notice that the optimisation for probability distributions yields a maximum since the number of inputs and outputs is fixed. Instead, the optimisation for a state may involve measurements with an arbitrarily large number of inputs or outputs, as is the case for the maximally entangled state [64]. In this work, the number of inputs and outputs is always finite, and this will become relevant when bounding the EPR2 components of distributions arising from maximally entangled states in Theorems 1 and 2.

GMNL from bipartite entanglement

Theorem 1. *Any connected network of bipartite pure entangled states is GMNL.*

Proof. We consider the network as a connected graph where vertices are parties and edges are states. The graph is such that, at each vertex, there is one particle for every incident edge [66]. We label the edges as $k = 1, \dots, |E|$ (where $|E|$ is the number of edges of the graph) and the parties as A_1, \dots, A_n . Since it will be enough to consider individual measurements on each particle, we denote the input and output of party A_i at edge k as x_i^k, a_i^k respectively. We group the inputs and outputs of each party as $\chi_i = \{x_i^k\}_{k \in E_i}$, $\alpha_i = \{a_i^k\}_{k \in E_i}$ where E_i is the set of edges incident to vertex i . Then, the shared distribution is of the form

$$P(\alpha_1, \dots, \alpha_n | \chi_1, \dots, \chi_n) = \prod_{k=1}^{|E|} P_k(a_i^k a_j^k | x_i^k x_j^k), \quad (8)$$

where parties A_i, A_j are connected by edge k (notice that we label vertices and edges independently), and $P_k(a_i^k a_j^k | x_i^k x_j^k)$ is the distribution arising from the state at edge k . It will be sufficient to consider tree graphs, i.e. such that every pair of vertices (parties) is connected by exactly one path of edges. If the given graph is not a tree, any extra edges can be ignored.

Depending on the nature of the shared states, we consider three cases:

- (i) every shared state is less-than-maximally entangled;
- (ii) every shared state is maximally entangled;
- (iii) some shared states are maximally entangled, some are not.

Case (i): if all states are less-than-maximally entangled, we prove the result by deriving an inequality that detects GMNL and finding measurements on the shared states to violate it. To derive the inequality, we will find bipartite inequalities that can be violated by the state at each edge k , lift them to more inputs, outputs and parties using the techniques in Ref. [55] and combine them to obtain a GMNL inequality using tools in Ref. [60]. We will consider 2-input 2-output measurements on each particle. Thus, the global distribution will have $2^{|E_i|}$ inputs and outputs for each party A_i .

We start from the inequality

$$I = P(00|00) - P(01|01) - P(10|10) - P(00|11) \leq 0, \quad (9)$$

which is a facet inequality equivalent to the CHSH inequality [61] for nonsignalling distributions [60]. This inequality detects any bipartite nonlocality present in any bipartition that splits the parties connected by edge k [60]. To lift it to n parties, each with $2^{|E_i|}$ inputs and outputs (see Ref. [55]), we must set the inputs and outputs of the parties that are not connected by edge k to a fixed value (0, wlog). For the parties i that are connected by edge k , any extra inputs other than $x_i^k = 0_i^k, 1_i^k$ can be ignored. Outputs must be grouped, by summing over some of their digits, in order to get an effective 2-output distribution. It will be convenient to add over the output components a_i^k that do not correspond to edge k , varying only the digit $a_i^k = 0_i^k, 1_i^k$. Thus, we obtain the following n -partite inequality at each edge k :

$$I^k = \sum_{\vec{a}_i^k, \vec{a}_j^k} \left(P \left(0_i^k \vec{a}_i^k, 0_j^k \vec{a}_j^k, \vec{0}_{\bar{i}, \bar{j}} \mid 0_i^k 0_i^k, 0_j^k 0_j^k, \vec{0}_{\bar{i}, \bar{j}} \right) - P \left(0_i^k \vec{a}_i^k, 1_j^k \vec{a}_j^k, \vec{0}_{\bar{i}, \bar{j}} \mid 0_i^k 0_i^k, 1_j^k 0_j^k, \vec{0}_{\bar{i}, \bar{j}} \right) \right. \\ \left. - P \left(1_i^k \vec{a}_i^k, 0_j^k \vec{a}_j^k, \vec{0}_{\bar{i}, \bar{j}} \mid 1_i^k 0_i^k, 0_j^k 0_j^k, \vec{0}_{\bar{i}, \bar{j}} \right) - P \left(0_i^k \vec{a}_i^k, 0_j^k \vec{a}_j^k, \vec{0}_{\bar{i}, \bar{j}} \mid 1_i^k 0_i^k, 1_j^k 0_j^k, \vec{0}_{\bar{i}, \bar{j}} \right) \right) \leq 0, \quad (10)$$

where the sum is over each binary digit a_i^k, a_j^k of the outputs of parties i, j (which are connected by edge k), except digits a_i^k, a_j^k which are fixed to 0 or 1 in each term. The term $\vec{0}_{\bar{i}, \bar{j}}$ denotes input or output 0 for all components of all parties that are not i, j . Thus, each inequality I^k detects the bipartite nonlocality present in the distribution P across any bipartition that splits the parties connected by edge k . In the particular case of the distribution (8), it tells whether the component P_k is nonlocal.

Now, we can combine the inequalities I^k to form a GMNL inequality:

$$I_n = \sum_{k=1}^{|E|} I^k + P(\vec{0}, \vec{0} | \vec{0}, \vec{0}) - \sum_{k=1}^{|E|} \sum_{\vec{a}_i^k, \vec{a}_j^k} P \left(0_i^k \vec{a}_i^k, 0_j^k \vec{a}_j^k, \vec{0}_{\bar{i}, \bar{j}} \mid 0_i^k 0_i^k, 0_j^k 0_j^k, \vec{0}_{\bar{i}, \bar{j}} \right) \leq 0. \quad (11)$$

To show that this is indeed a GMNL inequality, we must show that it holds for any distribution P that is local across some bipartition. A bipartition of the network defines a cut of the graph. Because the graph is assumed connected, for every cut there exists an edge k_0 which crosses the cut. Therefore, if P is local across a bipartition which is crossed by edge k_0 , then by Ref. [60] we have

$$I^{k_0} \leq 0. \quad (12)$$

Hence,

$$I_n \leq \sum_{\substack{k=1 \\ k \neq k_0}}^{|E|} I^k + P(\vec{0}, \vec{0} | \vec{0}, \vec{0}) - \sum_{k=1}^{|E|} \sum_{\vec{a}_i^k, \vec{a}_j^k} P\left(0_i^k \vec{a}_i^k, 0_j^k \vec{a}_j^k, \vec{0}_{i,j} | 0_i^k 0_i^k, 0_j^k 0_j^k, \vec{0}_{i,j}\right). \quad (13)$$

For each $k \neq k_0$, the only nonnegative term gets subtracted in the final summation. The term $P(\vec{0}, \vec{0} | \vec{0}, \vec{0})$ then cancels out with the first term in the final summation for $k = k_0$, leaving only negative terms in the expression as required.

To complete the proof, we find local measurements for each party to violate inequality (11). Since all shared states are nonseparable and less-than-maximally entangled, the parties can choose local measurements on each particle such that all resulting distributions satisfy Hardy's paradox [62, 63]:

$$P_k(00|00) > 0 = P_k(01|01) = P_k(10|10) = P_k(00|11) \quad (14)$$

for each $k = 1, \dots, |E|$. This was proven for qubits in Refs. [62, 63], and we show the extension to any local dimension in Proposition 1 below. Because the distribution is of the form (8), each term in each inequality (10) simplifies significantly. For example, the second term gives

$$\begin{aligned} & \sum_{\vec{a}_i^k, \vec{a}_j^k} P\left(0_i^k \vec{a}_i^k, 1_j^k \vec{a}_j^k, \vec{0}_{i,j} | 0_i^k 0_i^k, 1_j^k 0_j^k, \vec{0}_{i,j}\right) \\ &= P_k(0_i^k 1_j^k | 0_i^k 1_j^k) \prod_{\ell} \sum_{a_i^{\ell}} P_{\ell}(a_i^{\ell} 0_j^{\ell} | 0_i^{\ell} 0_j^{\ell}) \prod_{\ell'} \sum_{a_j^{\ell'}} P_{\ell'}(0_i^{\ell'} a_j^{\ell'} | 0_i^{\ell'} 0_j^{\ell'}) \prod_m P_m(0_{i'}^m 0_{j'}^m | 0_{i'}^m 0_{j'}^m) \\ &= P_k(0_i^k 1_j^k | 0_i^k 1_j^k) p_k, \end{aligned} \quad (15)$$

where edges ℓ connect party i to party $j' \neq j$, edges ℓ' connect party j to party $i' \neq i$, and edges m connect parties i' and j' where $i', j' \neq i, j$. (Depending on the structure of the graph, there may be no edges ℓ , ℓ' or m for a given pair of parties i, j , but that does not affect the proof.)

The product of the terms P_{ℓ} , $P_{\ell'}$ and P_m will give a number p_k . This is similar for the third and fourth terms, which factorise to

$$\begin{aligned} & P_k(1_i^k 0_j^k | 1_i^k 0_j^k) p_k, \\ & P_k(0_i^k 0_j^k | 1_i^k 1_j^k) p_k \end{aligned} \quad (16)$$

respectively. The first term of each I^k cancels out with the last summation in I_n , and the only term that remains is

$$P(\vec{0}, \vec{0} | \vec{0}, \vec{0}) = \prod_{k=1}^{|E|} P_k(0_i^k 0_j^k | 0_i^k 0_j^k). \quad (17)$$

Since P_k satisfies Hardy's paradox for every k , then the components of each P_k appearing in equations (15), (16) are all zero, while the only surviving term, $P(\vec{0}, \vec{0} | \vec{0}, \vec{0})$, is strictly greater than zero. Thus, the inequality I_n is violated, showing that P is GMNL.

Case (ii): for every bipartition, there is an edge that crosses the corresponding cut, and each of these edges already contains a maximally entangled state. Therefore, the present network meets the requirements of Theorem 2 in [57], so the network is GMNL—in fact it is fully GMNL.

Case (iii): assume wlog that each edge $k = 1, \dots, K$ contains a less-than-maximally entangled state, while each edge $k = K + 1, \dots, |E|$ contains a maximally entangled state. Let

$$P = P_H P_+ \quad (18)$$

where

$$\begin{aligned}
P_H(\{a_i^k\}_{k \leq K, i \in [n]} | \{x_i^k\}_{k \leq K, i \in [n]}) &= \prod_{k=1}^K P_k(a_i^k a_j^k | x_i^k x_j^k), \\
P_+(\{a_i^k\}_{k > K, i \in [n]} | \{x_i^k\}_{k > K, i \in [n]}) &= \prod_{k=K+1}^{|E|} P_k(a_i^k a_j^k | x_i^k x_j^k)
\end{aligned} \tag{19}$$

where, on the right-hand side, parties i, j are connected by edge k . For $k = 1, \dots, K$, terms P_k satisfy Hardy's paradox (equation (14)), as they arise from the measurements performed in Case (i). For $k = K + 1, \dots, |E|$, the terms P_k arise from measurements on the maximally entangled state to be specified later. We now classify bipartitions depending on whether or not they are crossed by an edge $k \leq K$ or $k > K$: let $S_{\leq K}$ be the set of bipartitions $M | \overline{M}$ (indexed by M) which are crossed by an edge $k \leq K$, and $T_{\leq K}$ be its complement, i.e. the set of bipartitions which are *not* crossed by an edge $k \leq K$. Similarly, $S_{> K}$ (respectively, $T_{> K}$) is the set of bipartitions which are (not) crossed by an edge $k > K$.

Let I_H^k be an inequality detecting nonlocality on edge k , for the distribution P_H . That is, I_H^k is as in equation (10) but where the sum over \vec{a}_i^k, \vec{a}_j^k concerns only the components of parties A_i, A_j that belong only to edges $k' \leq K, k' \neq k$. Then, consider the following functional acting on distributions of the form of P_H :

$$I_H = \sum_{k=1}^K I_H^k + P(\vec{0}, \vec{0} | \vec{0}, \vec{0}) - \sum_{k=1}^K \sum_{\vec{a}_i^k, \vec{a}_j^k} P\left(0_i^k \vec{a}_i^k, 0_j^k \vec{a}_j^k, \vec{0}_{\bar{i}, \bar{j}} \mid 0_i^k 0_i^k, 0_j^k 0_j^k, \vec{0}_{\bar{i}, \bar{j}}\right). \tag{20}$$

Again, the summation in the last term concerns only components that belong to edges $k' \leq K, k' \neq k$. We claim that the functional I_H is nonpositive for any distribution P that is local across a bipartition of type $S_{\leq K}$, i.e. one that is crossed by an edge $k_0 \leq K$. The reasoning is similar to that in Case (i): if P is local across a bipartition crossed by an edge $k_0 \leq K$, then $I_H^{k_0} \leq 0$ will be satisfied, and so

$$I_H \leq \sum_{\substack{k=1 \\ k \neq k_0}}^K I_H^k + P(\vec{0}, \vec{0} | \vec{0}, \vec{0}) - \sum_{k=1}^K \sum_{\vec{a}_i^k, \vec{a}_j^k} P\left(0_i^k \vec{a}_i^k, 0_j^k \vec{a}_j^k, \vec{0}_{\bar{i}, \bar{j}} \mid 0_i^k 0_i^k, 0_j^k 0_j^k, \vec{0}_{\bar{i}, \bar{j}}\right). \tag{21}$$

Now, for each $k \neq k_0$, the only nonnegative term gets subtracted in the final summation. The term $P(\vec{0}, \vec{0} | \vec{0}, \vec{0})$ then cancels out with the first term in the final summation for $k = k_0$, leaving only negative terms in the expression as required.

We now show that, for $P = P_H$, we have $I_H > 0$. Indeed, the terms in I_H^k simplify in a similar manner to Case (i). Then, since each P_k , $k \leq K$ satisfies Hardy's paradox, the second, third and fourth terms in each I_H^k are zero, the first cancels out with the last summation, and the only surviving term is

$$P(\vec{0}, \vec{0} | \vec{0}, \vec{0}) = \prod_{k=1}^K P_k(0_i^k 0_j^k | 0_i^k 0_j^k) > 0. \tag{22}$$

This means that there exists an $\varepsilon > 0$ such that, for any EPR2 decomposition of P_H ,

$$P_H = \sum_M p_{L,H}^M P_{L,H}^M + p_{NS,H} P_{NS,H}, \tag{23}$$

we have that the terms where $P_{L,H}^M$ is local across a bipartition such that $M \in S_{\leq K}$ satisfy

$$\sum_{M \in S_{\leq K}} p_{L,H}^M \leq 1 - \varepsilon. \tag{24}$$

Also, it can be deduced from Ref. [57] that, given the ε above, the parties can choose suitable measurements such that P_+ is fully nonlocal across all bipartitions $S_{> K}$. That is, any multipartite EPR2 decomposition of P_+ ,

$$P_+ = \sum_M p_{L,+}^M P_{L,+}^M + p_{NS,+} P_{NS,+}, \tag{25}$$

is such that the terms where $P_{L,+}^M$ is local across a bipartition such that $M \in S_{>K}$ satisfy

$$\sum_{M \in S_{>K}} p_{L,+}^M < \varepsilon. \quad (26)$$

To prove that the global distribution P is GMNL, as is our goal, we assume the converse, and we derive a contradiction from the nonlocality properties of P_H and P_+ . Assuming P is not GMNL, we can express the distribution as

$$P = \sum_{\lambda, M} p_L^M(\lambda) P_M(\{\alpha_i\}_{i \in M} | \{\chi_i\}_{i \in M}, \lambda) P_{\overline{M}}(\{\alpha_i\}_{i \in \overline{M}} | \{\chi_i\}_{i \in \overline{M}}, \lambda), \quad (27)$$

where $p_L^M(\lambda)$ are nonnegative numbers for every M, λ such that

$$\sum_{\lambda, M} p_L^M(\lambda) = 1, \quad (28)$$

for each $\alpha_i, \chi_i, i = 1, \dots, n$.

Then, summing over the output components a_i^k for all $k \leq K$ and all i , we get P_+ on the left-hand side, from equation (18). On the right-hand side, we get two types of terms (depending on the type of bipartition) that turn out to form an EPR2 decomposition of P_+ [67]. Indeed, the local terms are given by bipartitions such that $M \in S_{>K}$, while the nonlocal terms are given by bipartitions such that $M \in T_{>K}$ (since all terms are nonsignalling). By equation (26), the choice of measurements on the particles involved in P_+ ensures that

$$\sum_{\lambda, M \in S_{>K}} p_L^M(\lambda) < \varepsilon, \quad (29)$$

while

$$\sum_{\lambda, M \in T_{>K}} p_L^M(\lambda) > 1 - \varepsilon. \quad (30)$$

If, instead, we sum over the output components a_i^k for all $k > K$ and all i , we get P_H on the left-hand side, from equation (18). On the right-hand side, by similar reasoning we find an EPR2 decomposition of P_H . This time, $S_{\leq K}$ will give the local terms and $T_{\leq K}$ will give the nonlocal terms. By equation (24), we have

$$\sum_{\lambda, M \in S_{\leq K}} p_L^M(\lambda) \leq 1 - \varepsilon. \quad (31)$$

Now, since the graph is connected, if a bipartition is not crossed by an edge $k > K$, then it must be crossed by an edge $k \leq K$. That is, $T_{>K} \subseteq S_{\leq K}$. This means that equation (31) also holds if the sum is over $T_{>K}$, but this contradicts equation (30). Therefore, the distribution P must be GMNL. \square

In Theorem 1 we assumed that all less-than-maximally entangled states satisfy Hardy's paradox. This is shown for qubits in [63], and we now extend the proof to any dimension.

Proposition 1. *Let $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B \cong (\mathbb{C}^d)^{\otimes 2}$ be a nonseparable and less-than-maximally entangled pure state. Then, $|\psi\rangle$ satisfies Hardy's paradox.*

Proof. Let $|\psi\rangle$ be as in the statement of the Proposition. We present 2-input, 2-output measurements for $|\psi\rangle$ to generate a distribution which satisfies Hardy's paradox [62, 63] using tools from Ref. [60].

Consider the Schmidt decomposition

$$|\psi\rangle = \sum_{i=0}^{d-1} \lambda_i^{1/2} |ii\rangle \quad (32)$$

and assume the coefficients are ordered such that $0 \neq \lambda_0 \neq \lambda_1 \neq 0$, which is always possible if the state is nonseparable and less-than-maximally entangled. Wlog assume the Schmidt basis of the state is the canonical basis. Let $\alpha \in]0, \pi/2[$

and $\delta \in \mathbb{R}$ and consider the dual vectors

$$\begin{aligned}
\langle e_{0|0} | &= \cos \alpha \langle 0 | + e^{i\delta} \sin \alpha \langle 1 | \\
\langle e_{1|1} | &= \lambda_0 \cos \alpha \langle 0 | + \lambda_1 e^{i\delta} \sin \alpha \langle 1 | \\
\langle f_{0|0} | &= \lambda_1^{3/2} e^{i\delta} \sin \alpha \langle 0 | - \lambda_0^{3/2} \cos \alpha \langle 1 | \\
\langle f_{1|1} | &= \lambda_1^{1/2} e^{i\delta} \sin \alpha \langle 0 | - \lambda_0^{1/2} \cos \alpha \langle 1 |
\end{aligned} \tag{33}$$

(one can write the projectors in the Schmidt basis of the state instead of assuming the state decomposes into the canonical basis). Define the measurements $E_{a|x}$ for Alice, with input x and output a , and $F_{b|y}$ for Bob, with input y and output b , given by

$$\begin{aligned}
E_{0|0} &= |e_{0|0}\rangle\langle e_{0|0}| \\
E_{1|0} &\propto |e_{0|0}\rangle\langle e_{0|0}|^\perp \oplus \mathbb{1}_{2,\dots,d-1} \\
E_{0|1} &\propto |e_{1|1}\rangle\langle e_{1|1}|^\perp \\
E_{1|1} &\propto |e_{1|1}\rangle\langle e_{1|1}| \oplus \mathbb{1}_{2,\dots,d-1} \\
F_{0|0} &\propto |f_{0|0}\rangle\langle f_{0|0}| \\
F_{1|0} &\propto |f_{0|0}\rangle\langle f_{0|0}|^\perp \oplus \mathbb{1}_{2,\dots,d-1} \\
F_{0|1} &\propto |f_{1|1}\rangle\langle f_{1|1}|^\perp \oplus \mathbb{1}_{2,\dots,d-1} \\
F_{1|1} &\propto |f_{1|1}\rangle\langle f_{1|1}|
\end{aligned} \tag{34}$$

where $|e_{0|0}\rangle\langle e_{0|0}|^\perp$ denotes the density matrix corresponding to the vector orthogonal to $|e_{0|0}\rangle$ when restricted to the subspace spanned by $\{|0\rangle, |1\rangle\}$, and $\mathbb{1}_{2,\dots,d-1}$ is the identity operator on the subspace spanned by $\{|i\rangle\}_{i=2}^{d-1}$, for either Alice or Bob. Note that, since we are only interested in whether some probabilities are equal or different from zero, normalisation will not play a role.

We now show that the distribution given by

$$P(ab|xy) = \text{tr}(E_{a|x} \otimes F_{b|y} |\psi\rangle\langle\psi|) \tag{35}$$

satisfies Hardy's paradox. Indeed, because of the probabilities considered and the form of the measurements, only the terms in $i = 0, 1$ contribute to the probabilities that appear in Hardy's paradox, therefore

$$\begin{aligned}
P(01|01) &\propto \left| \sum_{i=0}^1 \lambda_i^{1/2} (\langle e_{0|0} | \otimes \langle f_{1|1} |) |ii\rangle \right|^2 = 0 \\
P(10|10) &\propto \left| \sum_{i=0}^1 \lambda_i^{1/2} (\langle e_{1|1} | \otimes \langle f_{0|0} |) |ii\rangle \right|^2 = 0 \\
P(00|11) &\propto \left| \sum_{i=0}^1 \lambda_i^{1/2} (\langle e_{0|1} | \otimes \langle f_{0|1} |) |ii\rangle \right|^2 = 0.
\end{aligned} \tag{36}$$

For $P(00|00)$, we find

$$\begin{aligned}
P(00|00) &\propto \left| \sum_{i=0}^1 \lambda_i^{1/2} (\langle e_{0|0} | \otimes \langle f_{0|0} |) |ii\rangle \right|^2 \\
&= \left| e^{i\delta} \sin \alpha \cos \alpha \lambda_0^{1/2} \lambda_1^{1/2} (\lambda_1 - \lambda_0) \right|^2,
\end{aligned} \tag{37}$$

which is strictly greater than zero when $\alpha \in]0, \pi/2[$ and $0 \neq \lambda_0 \neq \lambda_1 \neq 0$, like we assumed. This proves the claim. \square

GMNL from GME

We fix some notation that we will use in Theorem 2. The result considers a GME state $|\Psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_{B_1} \dots \otimes \mathcal{H}_{B_{n-1}} \cong (\mathbb{C}^d)^{\otimes n}$, $n-1$ copies of which are shared between n parties A, B_1, \dots, B_{n-1} . Each party measures locally on each particle, like in Theorem 1. We denote Alice's input and output, respectively, as $\chi \equiv x_1 \dots x_{n-1}$, $\alpha \equiv a_1 \dots a_{n-1}$ in terms of the digits x_i, a_i corresponding to each particle $i \in [n-1]$. We let the measurement made by party B_j on copy i have input y_j^i and output b_j^i , where $i, j = 1, \dots, n-1$, and for each j we denote $v_j = y_j^1 \dots y_j^{n-1}$ and $\beta_j = b_j^1 \dots b_j^{n-1}$ digit-wise. Then, after measurement, the parties share a distribution

$$\{P(\alpha\beta_1 \dots \beta_{n-1} | \chi v_1 \dots v_{n-1})\}_{\substack{\alpha, \beta_1 \dots \beta_{n-1} \\ \chi, v_1 \dots v_{n-1}}} \quad (38)$$

Because we are considering local measurements made on each particle, this distribution is of the form

$$P(\alpha\beta_1 \dots \beta_{n-1} | \chi v_1 \dots v_{n-1}) = \prod_{i=1}^{n-1} P_i(a_i b_1^i \dots b_{n-1}^i | x_i y_1^i \dots y_{n-1}^i), \quad (39)$$

where each P_i is the distribution arising from copy i of the state $|\Psi\rangle$. As advanced in the main text, each copy i of the state $|\Psi\rangle$ will give an edge of a star network connecting Alice and party B_i . Because of the structure of this particular network, we can simplify the notation with respect to Theorem 1 and identify the index of each party B_i with its corresponding edge i .

Theorem 2. *Any GME state $|\Psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_{B_1} \dots \otimes \mathcal{H}_{B_{n-1}} \cong (\mathbb{C}^d)^{\otimes n}$ is such that $|\Psi\rangle^{\otimes(n-1)}$ is GMNL.*

Proof. For each copy $i = 1, \dots, n-1$ of the state $|\Psi\rangle$, we will find measurements for parties $\{B_j\}_{j \neq i}$ that leave Alice and party B_i with a bipartite entangled state. This will yield a network in a similar configuration to Theorem 1 for a star network, but conditionalised on the inputs and outputs of these measurements. We will generalise the result of Theorem 1 as it applies to a star network to show that this network is also GMNL.

Let $i \in [n-1]$ and consider the i th copy of $|\Psi\rangle$. Suppose each party B_j , $j \neq i$, performs a local, projective measurement onto a basis $\{|b_j\rangle\}_{b_j=0}^{d-1}$. We pick the computational basis on each party's Hilbert space to be such that the measurement performed by the parties B_j , $j \neq i$, leave Alice and B_i in state $|\phi_{\vec{b}}\rangle_{AB_i}$, where $\vec{b} = b_1 \dots b_{i-1} b_{i+1} \dots b_{n-1}$ denotes the output obtained by the parties B_j , $j \neq i$ (we briefly omit the script i referring to the copy of the state, for readability). This means that we can write the state $|\Psi\rangle$ as

$$|\Psi\rangle = \sum_{\vec{b}} \lambda_{\vec{b}} |\phi_{\vec{b}}\rangle_{AB_i} |\vec{b}\rangle_{B_1 \dots B_{i-1} B_{i+1} \dots B_{n-1}} \quad (40)$$

Ref. [28], whose proof was completed in Ref. [29], showed that there always exist measurements (i.e. bases) $\{|b_j\rangle\}_{b_j=0}^{d-1}$ such that $|\phi_{\vec{b}}\rangle_{AB_i}$ is entangled for a certain output \vec{b} . We now show that this opens up only two possibilities for each i : either there exists an output such that $|\phi_{\vec{b}}\rangle_{AB_i}$ is less-than-maximally entangled, or for all outputs \vec{b} , $|\phi_{\vec{b}}\rangle_{AB_i}$ is maximally entangled. Indeed, the only option left to discard is one where, for some $\vec{b} = \vec{b}^*$, $|\phi_{\vec{b}^*}\rangle_{AB_i}$ is maximally entangled, and for some other $\vec{b} = \vec{b}^{**}$, $|\phi_{\vec{b}^{**}}\rangle_{AB_i}$ is separable. But it is easy to see, by using a continuity argument, that in this case the bases $\{|b_j\rangle\}_{b_j=0}^{d-1}$ can be modified so that there exists one output for which AB_i are projected onto a less-than-maximally entangled state: it suffices to consider one (normalised) element of the measurement basis to be $c_0 |b_j^*\rangle + c_1 |b_j^{**}\rangle$ for some values $c_0, c_1 \in \mathbb{C}$, for each j .

Therefore, we consider the following cases:

- (i) for all $i \in [n-1]$, there exists an input and output for each $B_j, j \neq i$ such that $|\phi_i\rangle_{AB_i}$ is less-than-maximally entangled;
- (ii) for all $i \in [n-1]$, there exists an input for each $B_j, j \neq i$ such that $|\phi_i\rangle_{AB_i}$ is maximally entangled for all outputs;
- (iii) there exist $i, k \in [n-1]$ such that $|\phi_i\rangle_{AB_i}$ is as in Case (ii) and $|\phi_k\rangle_{AB_k}$ is as in Case (i).

Case (i): let $i \in [n-1]$. Suppose parties $\{B_j\}_{j \neq i}$ perform the measurements explained above that leave Alice and B_i less-than-maximally entangled. Then, Alice and B_i can perform local measurements on the resulting state to satisfy Hardy's paradox. We will modify the inequality in Theorem 1 and show that these measurements on $|\Psi\rangle^{\otimes(n-1)}$ give a distribution which violates the inequality.

To modify the inequality in Theorem 1, we import the same strategy to lift inequality (9) to n parties, each with 2^{n-1} inputs and outputs. We want I^{AB_i} to detect bipartite nonlocality between Alice's i th particle and B_i 's i th particle, that is, nonlocality in $a_i b_i^i |x_i y_i^i$. Therefore, for each i we now need to fix all other inputs x_j, y_i^j, y_j^j and add over all other outputs $a_j, b_i^j, b_j^j, j \neq i$, so that

$$I^{AB_i} = \sum_{a_i, b_i^i, b_i^i=0,1} \left(P(0_i a_{\bar{i}}, 0_i^i b_{\bar{i}}^i, 0_i^i b_{\bar{i}}^i | 0_i 0_{\bar{i}}, 0_i^i 0_{\bar{i}}^i, 0_i^i 0_{\bar{i}}^i) - P(0_i a_{\bar{i}}, 1_i^i b_{\bar{i}}^i, 0_i^i b_{\bar{i}}^i | 0_i 0_{\bar{i}}, 1_i^i 0_{\bar{i}}^i, 0_i^i 0_{\bar{i}}^i) \right. \\ \left. - P(1_i a_{\bar{i}}, 0_i^i b_{\bar{i}}^i, 0_i^i b_{\bar{i}}^i | 1_i 0_{\bar{i}}, 0_i^i 0_{\bar{i}}^i, 0_i^i 0_{\bar{i}}^i) - P(0_i a_{\bar{i}}, 0_i^i b_{\bar{i}}^i, 0_i^i b_{\bar{i}}^i | 1_i 0_{\bar{i}}, 1_i^i 0_{\bar{i}}^i, 0_i^i 0_{\bar{i}}^i) \right), \quad (41)$$

where the outputs in the first term are denoted as follows: $0_i a_{\bar{i}}$ denotes output $\alpha = a_1 \dots 0_i \dots a_{n-1}$, $0_i^i b_{\bar{i}}^i$ denotes output $\beta_i = b_1^1 \dots 0_i^i \dots b_i^{n-1}$, and $0_i^i b_{\bar{i}}^i$ denotes output $\beta_j = b_j^1 \dots 0_j^j \dots b_j^{n-1}$ for all $j \neq i$. Inputs are denoted similarly, and the notation is similar for the other three terms. Then, the inequality

$$I_n = \sum_{i=1}^{n-1} I^{AB_i} + P(\vec{0}, \vec{0} | \vec{0}, \vec{0}) - \sum_{i=1}^{n-1} \sum_{a_{\bar{i}}, b_i^i, b_i^i=0,1} P(0_i a_{\bar{i}}, 0_i^i b_{\bar{i}}^i, 0_i^i b_{\bar{i}}^i | 0_i 0_{\bar{i}}, 0_i^i 0_{\bar{i}}^i, 0_i^i 0_{\bar{i}}^i) \leq 0 \quad (42)$$

is a GMNL inequality, by the same reasoning as in Theorem 1.

Evaluating the inequality on the distribution (39), we find again that each term simplifies. For each i we get, for example,

$$\sum_{a_{\bar{i}}, b_i^i, b_i^i=0,1} P(0_i a_{\bar{i}}, 1_i^i b_{\bar{i}}^i, 0_i^i b_{\bar{i}}^i | 0_i 0_{\bar{i}}, 1_i^i 0_{\bar{i}}^i, 0_i^i 0_{\bar{i}}^i) \\ = P_i(0_i 1_i^i 0_i^i | 0_i 1_i^i 0_i^i) \prod_{\substack{j=1 \\ j \neq i}}^{n-1} \sum_{\substack{a_j, b_k^j=0,1 \\ k \neq j}} P_j(a_j b_1^j \dots b_{j-1}^j b_{j+1}^j \dots b_{n-1}^j | 0_j 0_1^j \dots 0_{j-1}^j 0_{j+1}^j \dots 0_{n-1}^j) \\ = P_i(0_i 1_i^i 0_i^i | 0_i 1_i^i 0_i^i) \quad (43)$$

and, similarly,

$$\sum_{a_{\bar{i}}, b_i^i, b_i^i=0,1} P(1_i a_{\bar{i}}, 0_i^i b_{\bar{i}}^i, 0_i^i b_{\bar{i}}^i | 1_i 0_{\bar{i}}, 0_i^i 0_{\bar{i}}^i, 0_i^i 0_{\bar{i}}^i) = P_i(1_i 0_i^i 0_i^i | 1_i 0_i^i 0_i^i); \\ \sum_{a_{\bar{i}}, b_i^i, b_i^i=0,1} P(0_i a_{\bar{i}}, 0_i^i b_{\bar{i}}^i, 0_i^i b_{\bar{i}}^i | 1_i 0_{\bar{i}}, 1_i^i 0_{\bar{i}}^i, 0_i^i 0_{\bar{i}}^i) = P_i(0_i 0_i^i 0_i^i | 1_i 1_i^i 0_i^i). \quad (44)$$

Also,

$$P(\vec{0}, \vec{0} | \vec{0}, \vec{0}) = \prod_{i=1}^{n-1} P_i(0_i 0_i^i 0_i^i | 0_i 0_i^i 0_i^i). \quad (45)$$

Now each P_i in equation (39) arises from measurements by $\{B_j\}_{j \neq i}$ to create a less-than-maximally entangled state between Alice and B_i , who can then choose measurements to satisfy Hardy's paradox. Hence all terms are zero except $P(\vec{0}, \vec{0} | \vec{0}, \vec{0}) > 0$, and so the inequality is violated. Therefore, $|\Psi\rangle^{\otimes(n-1)}$ is GMNL.

Case (ii): we assumed that, for all $i \in [n-1]$, there exist local measurements on $|\Psi\rangle$ for parties $\{B_j\}_{j \neq i}$ that, for all outcomes, create a maximally entangled state $|\phi_i\rangle_{AB_i}$ shared between Alice and B_i . Since all bipartitions can be expressed as $A|B_i$ for some i , we find that $|\Psi\rangle$ meets the requirements of Theorem 2 in [57], and so $|\Psi\rangle$ is GMNL. That is, one copy of the shared state $|\Psi\rangle$ is already GMNL, and therefore so is $|\Psi\rangle^{\otimes(n-1)}$.

Case (iii): assume wlog that the state $|\phi_i\rangle_{AB_i}$ is less-than-maximally entangled for $i = 1, \dots, K$ and maximally entangled for $i = K+1, \dots, n-1$. We will show that $|\Psi\rangle^{\otimes(K+1)}$ is GMNL, which implies that $|\Psi\rangle^{\otimes(n-1)}$ is so too.

It will be useful to classify bipartitions $M|\overline{M}$ like in Theorem 1. We will always assume that Alice belongs to M in order not to duplicate the bipartitions. Let $S_{\leq K}$ be the set of bipartitions $M|\overline{M}$ (indexed by M) which are crossed by an edge $j \leq K$, i.e., where \overline{M} contains at least one index $j \in \{1, \dots, K\}$, and $T_{\leq K}$ be its complement, i.e. the set of bipartitions where \overline{M} contains only indices $j \in \{K+1, \dots, n-1\}$. Similarly, $S_{>K}$ (respectively, $T_{>K}$) is the set of bipartitions which are (not) crossed by an edge $j > K$. That is, in $S_{>K}$, there is some $j \in \{K+1, \dots, n-1\}$ which belongs to \overline{M} , while in $T_{>K}$, \overline{M} contains only indices $j \in \{1, \dots, K\}$.

For each $i = 1, \dots, K$, parties AB_i can perform measurements on their shared state $|\phi_i\rangle_{AB_i}$ which, together with the measurements of parties $\{B_j\}_{j \neq i}$ that projected $|\Psi\rangle$ onto $|\phi_i\rangle_{AB_i}$, give rise to a distribution

$$P_i(a_i b_1^i \dots b_{n-1}^i | x_i y_1^i \dots y_{n-1}^i) \quad (46)$$

which satisfies Hardy's paradox when post-selected on the inputs and outputs of parties $\{B_j\}_{j \neq i}$. Then, the distribution arising from the first K copies of $|\Psi\rangle$ is

$$P_H(\{a_i\}_{i \leq K} \{b_j^i\}_{i \leq K, j \in [n-1]} | \{x_i\}_{i \leq K} \{y_j^i\}_{i \leq K, j \in [n-1]}) = \prod_{i=1}^K P_i(a_i b_1^i \dots b_{n-1}^i | x_i y_1^i \dots y_{n-1}^i), \quad (47)$$

with P_i as in equation (46). This distribution is similar to that in Case (i) when post-selected on the inputs and outputs of parties $\{B_j\}_{j > K}$. More precisely, by the nonsignalling condition, we have

$$\begin{aligned} P_H(\{a_i\}_{i \leq K} \{b_j^i\}_{i \leq K, j \leq K} \{b_j^i = 0_j^i\}_{i \leq K, j > K} | \{x_i\}_{i \leq K} \{y_j^i\}_{i \leq K, j \leq K} \{y_j^i = 0_j^i\}_{i \leq K, j > K}) = \\ P_{AB_1 \dots B_K}(\{a_i\}_{i \leq K} \{b_j^i\}_{i \leq K, j \leq K} | \{x_i\}_{i \leq K} \{y_j^i\}_{i \leq K, j \leq K}, \{b_j^i = 0_j^i\}_{i \leq K, j > K}, \{y_j^i = 0_j^i\}_{i \leq K, j > K}) \\ \times P_{B_{K+1} \dots B_{n-1}}(\{b_j^i = 0_j^i\}_{i \leq K, j > K} | \{y_j^i = 0_j^i\}_{i \leq K, j > K}), \end{aligned} \quad (48)$$

where by Case (i) we know that $P_{AB_1 \dots B_K}$ is GMNL in its parties. Then, P_H must be $(K+1)$ -way nonlocal (i.e., GMNL when restricted to parties A, B_1, \dots, B_K). Indeed, if this were not the case, by equation (48) we could obtain a decomposition of the form (1) for $P_{AB_1 \dots B_K}$, which would contradict the fact that this distribution is GMNL.

Therefore, there exists an $\varepsilon > 0$ such that any EPR2 decomposition of P_H as

$$P_H = \sum_M p_{L,H}^M P_{L,H}^M + p_{NS,H} P_{NS,H} \quad (49)$$

we have that the terms where $P_{L,H}^M$ is local across a bipartition such that $M \in S_{\leq K}$ satisfy

$$\sum_{M \in S_{\leq K}} p_{L,H}^M \leq 1 - \varepsilon. \quad (50)$$

On the other hand, $|\Psi\rangle$ satisfies Theorem 1 in Ref. [57] for all bipartitions $A|B_i$ for $i = K+1, \dots, n-1$, hence it is fully nonlocal across all such bipartitions. This means that, for any $\delta_i > 0$, there exist local measurements on $|\Psi\rangle$ (which depend on i) that lead to a distribution

$$P_+(ab_1 \dots b_{n-1} | xy_1 \dots y_{n-1}) \quad (51)$$

such that any bipartite EPR2 decomposition across a bipartition $A|B_i$, for $i = K+1, \dots, n-1$,

$$P_+ = p_{L,+}^{A|B_i} P_{L,+}^{A|B_i} + (1 - p_{L,+}^{A|B_i}) P_{NS,+}^{A|B_i} \quad (52)$$

satisfies

$$p_{L,+}^{A|B_i} < \delta_i. \quad (53)$$

Thus, considering the possibility of implementing all the above measurements for each i leads to a distribution of the form (51) in which equation (53) holds for every $i = K+1, \dots, n-1$.

Therefore, given the ε above, the parties can choose suitable δ_i to bound the bipartitely local components and hence ensure that any multipartite EPR2 decomposition of P_+ ,

$$P_+ = \sum_M p_{L,+}^M P_{L,+}^M + p_{NS,+} P_{NS,+} \quad (54)$$

is such that the terms where $P_{L,+}^M$ is local across a bipartition such that $M \in S_{>K}$ satisfy

$$\sum_{M \in S_{>K}} p_{L,+}^M < \varepsilon. \quad (55)$$

Since we only need to consider $(K+1)$ copies of the state, we denote the inputs and outputs of Alice and each party B_j , $j \in [n-1]$ by $\chi = x_1 \dots x_{K+1}$, $v_j = y_j^1 \dots y_j^{K+1}$; $\alpha = a_1 \dots a_{K+1}$, $\beta_j = b_j^1 \dots b_j^{K+1}$ respectively. Then, the global distribution obtained from $|\Psi\rangle^{\otimes(K+1)}$ is

$$\begin{aligned} P(\alpha\beta_1 \dots \beta_{n-1} | \chi v_1 \dots v_{n-1}) = \\ P_H(\{a_i\}_{i \leq K} \{b_j^i\}_{i \leq K, j \in [n-1]} | \{x_i\}_{i \leq K} \{y_j^i\}_{i \leq K, j \in [n-1]}) \\ \times P_+(a_{K+1} b_1^{K+1} \dots b_{n-1}^{K+1} | x_{K+1} y_1^{K+1} \dots y_{n-1}^{K+1}), \end{aligned} \quad (56)$$

where P_H comes from equation (47) and the EPR2 components of P_H, P_+ are as per equations (50), (55).

We now follow a similar strategy to that in Theorem 1. To prove that the global distribution P is GMNL, as is our goal, we assume the converse, and we derive a contradiction from the nonlocality properties of P_H and P_+ . Assuming P is not GMNL, we can express the distribution as

$$P(\alpha\beta_1 \dots \beta_{n-1} | \chi v_1 \dots v_{n-1}) = \sum_{\lambda, M} p_L^M(\lambda) P_M(\alpha\{\beta_j\}_{j \in M} | \chi\{v_j\}_{j \in M}, \lambda) P_{\overline{M}}(\{\beta_j\}_{j \in \overline{M}} | \{v_j\}_{j \in \overline{M}}, \lambda), \quad (57)$$

where

$$\sum_{\lambda, M} p_L^M(\lambda) = 1, \quad (58)$$

for each $\alpha, \beta_j, \chi, v_j, j = 1, \dots, n-1$, where we recall that each $\beta_j = b_j^1 \dots b_j^{K+1}$ and similarly for v_j .

Now, if we sum equation (57) over a_i, b_j^i for $i = 1, \dots, K$ and $j = 1, \dots, n-1$ (that is, we sum over the i th digit, $i \leq K$, of Alice and all parties B_j), we obtain P_+ on the left-hand side, from equation (56). On the right-hand side, we obtain, for each M , [68]

$$\sum_{\lambda} p_L^M(\lambda) P_M(a_{K+1} \{b_j^{K+1}\}_{j \in M} | \chi\{v_j\}_{j \in M}, \lambda) P_{\overline{M}}(\{b_j^{K+1}\}_{j \in \overline{M}} | \{v_j\}_{j \in \overline{M}}, \lambda), \quad (59)$$

whose sum turns out to form an EPR2 decomposition of P_+ . Indeed, local terms are given by bipartitions such that $M \in S_{>K}$, as in these terms there is some digit b_j^{K+1} with $j > K$ appearing in $P_{\overline{M}}$, thus they are local across $A|B_j$ for some $j > K$. The nonlocal terms are given by bipartitions such that $M \in T_{>K}$ (since all terms are nonsignalling). Therefore, the choice of measurements which generated P_+ ensures (by equation (55)) that

$$\sum_{\lambda, M \in S_{>K}} p_L^M(\lambda) < \varepsilon \quad (60)$$

and hence

$$\sum_{\lambda, M \in T_{>K}} p_L^M(\lambda) > 1 - \varepsilon. \quad (61)$$

Going back now to equation (57), we sum over a_{K+1}, b_j^{K+1} for $j = 1, \dots, n-1$ (that is, we sum over the $(K+1)$ th digit of Alice and all parties B_j). Then, we obtain P_H on the left-hand side, from equation (56). On the right-hand side, we obtain for each M ,

$$\sum_{\lambda} p_L^M(\lambda) P_M(\{a_i\}_{i \leq K} \{b_j^i\}_{i \leq K, j \in M} | \chi\{v_j\}_{j \in M}, \lambda) P_{\overline{M}}(\{b_j^i\}_{i \leq K, j \in \overline{M}} | \{v_j\}_{j \in \overline{M}}, \lambda), \quad (62)$$

whose sum over M gives an EPR2 decomposition of P_H . This time, $S_{\leq K}$ will give the local terms, as $P_{\overline{M}}$ will contain at least some digit b_j^i for $j \leq K$, while $T_{\leq K}$ will give the nonlocal terms. By equation (50), our choice of ε implies that

$$\sum_{\lambda, M \in S_{\leq K}} p_L^M(\lambda) \leq 1 - \varepsilon. \quad (63)$$

Now, any bipartition in $T_{>K}$ is such that all $j \in \{K+1, \dots, n-1\}$ are in M . Hence, there must be some $j \leq K$ in \overline{M} , otherwise \overline{M} would be empty. Therefore, $P_{\overline{M}}$ always contains at least one digit b_j^j for some $j \leq K$, and so terms where $M \in T_{>K}$ are local across the bipartition $A|B_j$ for some $j \leq K$. That is, $T_{>K} \subseteq S_{\leq K}$.

This means that equation (63) also holds if the sum is over $T_{>K}$, but this is in contradiction with equation (61). \square

-
- [1] N. Gisin, G. Ribordy, W. Tittel, and H. Zbinden, Quantum Cryptography, *Rev. Mod. Phys.* **74**, 145 (2002), arXiv:quant-ph/0101098.
- [2] S. Pirandola, U. L. Andersen, L. Banchi, M. Berta, D. Bunandar, R. Colbeck, D. Englund, T. Gehring, C. Lupo, C. Ottaviani, J. Pereira, M. Razavi, J. S. Shaari, M. Tomamichel, V. C. Usenko, G. Vallone, P. Villoresi, and P. Wallden, *Advances in Quantum Cryptography*, *Adv. Opt. Photon.* **12**, 1012 (2020), arXiv:1906.01645.
- [3] A. Acín and L. Masanes, Certified randomness in quantum physics, *Nature* **540**, 213 (2016), arXiv:1708.00265.
- [4] H. Buhrman, R. Cleve, S. Massar, and R. de Wolf, Nonlocality and communication complexity, *Rev. Mod. Phys.* **82**, 665 (2010), arXiv:0907.3584 [quant-ph].
- [5] D. Mayers and A. Yao, Quantum Cryptography with Imperfect Apparatus, in *Proceedings of the 39th Annual Symposium on Foundations of Computer Science*, FOCS '98 (IEEE Computer Society, USA, 1998) p. 503, arXiv:quant-ph/9809039.
- [6] A. Acín, N. Brunner, N. Gisin, S. Massar, S. Pironio, and V. Scarani, Device-independent security of quantum cryptography against collective attacks, *Phys. Rev. Lett.* **98**, 230501 (2007), arXiv:quant-ph/0702152.
- [7] R. Colbeck, *Quantum And Relativistic Protocols For Secure Multi-Party Computation*, Ph.D. thesis (2011), arXiv:0911.3814.
- [8] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, Bell nonlocality, *Rev. Mod. Phys.* **86**, 419 (2014), arXiv:1303.2849.
- [9] L. Aolita, R. Gallego, A. Cabello, and A. Acín, Fully nonlocal, monogamous, and random genuinely multipartite quantum correlations, *Phys. Rev. Lett.* **108**, 100401 (2012), arXiv:1109.3163.
- [10] J. Tura, R. Augusiak, A. B. Sainz, T. Vértesi, M. Lewenstein, and A. Acín, Detecting nonlocality in many-body quantum states, *Science* **344**, 1256 (2014), arXiv:1306.6860 [quant-ph].
- [11] J. Tura, G. De las Cuevas, R. Augusiak, M. Lewenstein, A. Acín, and J. I. Cirac, Energy as a Detector of Nonlocality of Many-Body Spin Systems, *Phys. Rev. X* **7**, 021005 (2017), arXiv:1607.06090 [quant-ph].
- [12] D. Cavalcanti, M. L. Almeida, V. Scarani, and A. Acín, Quantum networks reveal quantum nonlocality, *Nat Commun* **2**, 184 (2011), arXiv:1010.0900.
- [13] N. Gisin, Q. Mei, A. Tavakoli, M. O. Renou, and N. Brunner, All entangled pure quantum states violate the bilocality inequality, *Phys. Rev. A* **96**, 020304(R) (2017), arXiv:1702.00333.
- [14] I. Šupić, P. Skrzypczyk, and D. Cavalcanti, Measurement-device-independent entanglement and randomness estimation in quantum networks, *Phys. Rev. A* **95**, 042340 (2017).
- [15] A. Tavakoli, M. O. Renou, N. Gisin, and N. Brunner, Correlations in star networks: From Bell inequalities to network inequalities, *New J. Phys.* **19**, 073003 (2017), arXiv:1702.03866.
- [16] M.-O. Renou, E. Bäumer, S. Boreiri, N. Brunner, N. Gisin, and S. Beigi, Genuine Quantum Nonlocality in the Triangle Network, *Phys. Rev. Lett.* **123**, 140401 (2019), arXiv:1905.04902 [quant-ph].
- [17] N. Gisin, J.-D. Bancal, Y. Cai, P. Remy, A. Tavakoli, E. Zambrini Cruzeiro, S. Popescu, and N. Brunner, Constraints on nonlocality in networks from no-signaling and independence, *Nature Communications* **11**, 2378 (2020).
- [18] T. Kriváchy, Y. Cai, D. Cavalcanti, A. Tavakoli, N. Gisin, and N. Brunner, A neural network oracle for quantum nonlocality problems in networks, *npj Quantum Information* **6**, 1 (2020).
- [19] J. I. Cirac, A. Ekert, S. F. Huelga, and C. Macchiavello, Distributed Quantum Computation over Noisy Channels, *Phys. Rev. A* **59**, 4249 (1999), arXiv:quant-ph/9803017.
- [20] M. Howard and J. Vala, Nonlocality as a Benchmark for Universal Quantum Computation in Ising Anyon Topological Quantum Computers, *Phys. Rev. A* **85**, 022304 (2012), arXiv:1112.1516.
- [21] M. Howard, J. J. Wallman, V. Veitch, and J. Emerson, Contextuality supplies the magic for quantum computation, *Nature* **510**, 351 (2014), arXiv:1401.4174.
- [22] C. Branciard, N. Gisin, and S. Pironio, Characterizing the Nonlocal Correlations Created via Entanglement Swapping, *Physical Review Letters* **104**, 10.1103/PhysRevLett.104.170401 (2010), arXiv:0911.1314 [quant-ph].
- [23] C. Branciard, D. Rosset, N. Gisin, and S. Pironio, Bilocal versus non-bilocal correlations in entanglement swapping experiments, *Phys. Rev. A* **85**, 032119 (2012), arXiv:1112.4502.
- [24] R. F. Werner, Quantum states with Einstein-Podolsky-Rosen correlations admitting a hidden-variable model, *Phys. Rev. A* **40**, 4277 (1989).
- [25] J. Barrett, Nonsequential positive-operator-valued measurements on entangled mixed states do not always violate a Bell inequality, *Physical Review A* **65**, 10.1103/PhysRevA.65.042302 (2002), arXiv:quant-ph/0107045.
- [26] N. Gisin, Bell's inequality holds for all non-product states, *Physics Letters A* **154**, 201 (1991).
- [27] N. Gisin and A. Peres, Maximal violation of Bell's inequality for arbitrarily large spin, *Physics Letters A* **162**, 15 (1992).
- [28] S. Popescu and D. Rohrlich, Generic quantum nonlocality, *Physics Letters A* **166**, 293 (1992).
- [29] M. Gachechiladze and O. Gühne, Completing the proof of "Generic quantum nonlocality", *Physics Letters A* **381**, 1281 (2017), arXiv:1607.02948 [quant-ph].

- [30] A. Sen(De), U. Sen, Č. Brukner, V. Bužek, and M. Żukowski, Entanglement swapping of noisy states: A kind of superadditivity in nonclassicality, *Phys. Rev. A* **72**, 042310 (2005), arXiv:quant-ph/0311194.
- [31] D. Cavalcanti, R. Rabelo, and V. Scarani, Nonlocality Tests Enhanced by a Third Observer, *Phys. Rev. Lett.* **108**, 040402 (2012), arXiv:1110.3656 [quant-ph].
- [32] M.-X. Luo, Nonlocality of all quantum networks, *Physical Review A* **98**, 10.1103/PhysRevA.98.042317 (2018), arXiv:1806.09758 [quant-ph].
- [33] M.-X. Luo, A nonlocal game for witnessing quantum networks, *npj Quantum Inf* **5**, 1 (2019), arXiv:1906.08398 [quant-ph].
- [34] G. Svetlichny, Distinguishing three-body from two-body nonseparability by a Bell-type inequality, *Phys. Rev. D* **35**, 3066 (1987).
- [35] M. Seevinck and G. Svetlichny, Bell-Type Inequalities for Partial Separability in N -Particle Systems and Quantum Mechanical Violations, *Physical Review Letters* **89**, 10.1103/PhysRevLett.89.060401 (2002), arXiv:quant-ph/0201046.
- [36] D. Collins, N. Gisin, S. Popescu, D. Roberts, and V. Scarani, Bell-Type Inequalities to Detect True n -Body Nonseparability, *Physical Review Letters* **88**, 10.1103/PhysRevLett.88.170405 (2002), arXiv:quant-ph/0201058.
- [37] F. Buscemi, All Entangled Quantum States Are Nonlocal, *Phys. Rev. Lett.* **108**, 200401 (2012), arXiv:1106.6095 [quant-ph].
- [38] R. Gallego, L. E. Würflinger, A. Acín, and M. Navascués, Operational Framework for Nonlocality, *Phys. Rev. Lett.* **109**, 070401 (2012), arXiv:1112.2647 [quant-ph].
- [39] J.-D. Bancal, J. Barrett, N. Gisin, and S. Pironio, Definitions of multipartite nonlocality, *Phys. Rev. A* **88**, 014102 (2013), arXiv:1112.2626 [quant-ph].
- [40] J. Geller and M. Piani, Quantifying non-classical and beyond-quantum correlations in the unified operator formalism, *J. Phys. A: Math. Theor.* **47**, 424030 (2014), arXiv:1401.8197.
- [41] J. I. de Vicente, On nonlocality as a resource theory and nonlocality measures, *J. Phys. A: Math. Theor.* **47**, 424017 (2014), arXiv:1401.6941 [quant-ph].
- [42] R. Gallego and L. Aolita, Nonlocality free wirings and the distinguishability between Bell boxes, *Phys. Rev. A* **95**, 032118 (2017), arXiv:1611.06932 [quant-ph].
- [43] D. Schmid, D. Rosset, and F. Buscemi, The type-independent resource theory of local operations and shared randomness, *Quantum* **4**, 262 (2020), arXiv:1909.04065.
- [44] E. Wolfe, D. Schmid, A. B. Sainz, R. Kunjwal, and R. W. Spekkens, Quantifying Bell: The Resource Theory of Nonclassicality of Common-Cause Boxes, *Quantum* **4**, 280 (2020).
- [45] E. Hänggi, *Device-Independent Quantum Key Distribution*, Ph.D. thesis, ETH Zürich (2010), arXiv:1012.3878.
- [46] R. Augusiak, M. Demianowicz, J. Tura, and A. Acín, Entanglement and nonlocality are inequivalent for any number of particles, *Phys. Rev. Lett.* **115**, 030404 (2015), arXiv:1407.3114.
- [47] R. Augusiak, M. Demianowicz, and J. Tura, Constructions of genuinely entangled multipartite states with applications to local hidden variables (LHV) and states (LHS) models, *Phys. Rev. A* **98**, 012321 (2018), arXiv:1803.00279.
- [48] J. Bowles, J. Francfort, M. Fillettaz, F. Hirsch, and N. Brunner, Genuinely multipartite entangled quantum states with fully local hidden variable models and hidden multipartite nonlocality, *Phys. Rev. Lett.* **116**, 130401 (2016), arXiv:1511.08401.
- [49] Q. Chen, S. Yu, C. Zhang, C. H. Lai, and C. H. Oh, Test of Genuine Multipartite Nonlocality without Inequalities, *Phys. Rev. Lett.* **112**, 140404 (2014), arXiv:1305.4472 [quant-ph].
- [50] S. Yu and C. H. Oh, Tripartite entangled pure states are tripartite nonlocal, arXiv Preprint (2013), arXiv:1306.5330.
- [51] Hidden GMNL for three parties beyond qubits can be shown if some form of preprocessing is allowed.
- [52] M. Navascués and T. Vértesi, Activation of Nonlocal Quantum Resources, *Physical Review Letters* **106**, 10.1103/PhysRevLett.106.060403 (2011), arXiv:1010.5191 [quant-ph].
- [53] C. Palazuelos, Super-activation of quantum non-locality, *Phys. Rev. Lett.* **109**, 190401 (2012), arXiv:1205.3118.
- [54] P. Caban, A. Molenda, and K. Trzcińska, Activation of the violation of the Svetlichny inequality, *Physical Review A* **92**, 10.1103/PhysRevA.92.032119 (2015).
- [55] S. Pironio, Lifting Bell inequalities, *Journal of Mathematical Physics* **46**, 062112 (2005), arXiv:quant-ph/0503179.
- [56] A. C. Elitzur, S. Popescu, and D. Rohrlich, Quantum nonlocality for each pair in an ensemble, *Physics Letters A* **162**, 25 (1992).
- [57] M. L. Almeida, D. Cavalcanti, V. Scarani, and A. Acín, Multipartite fully-nonlocal quantum states, *Phys. Rev. A* **81**, 052111 (2010), arXiv:0911.3559.
- [58] Not to be confused with “nonfully local”, which is the opposite of “fully local”. “fully nonlocal” is a particular case of “nonfully local”.
- [59] See Supplemental Material for full proofs of the results in the main text.
- [60] F. J. Curchod, M. L. Almeida, and A. Acín, A versatile construction of Bell inequalities for the multipartite scenario, *New J. Phys.* **21**, 023016 (2019), arXiv:1808.10688 [quant-ph].
- [61] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Proposed Experiment to Test Local Hidden-Variable Theories, *Phys. Rev. Lett.* **23**, 880 (1969).
- [62] L. Hardy, Quantum mechanics, local realistic theories, and Lorentz-invariant realistic theories, *Phys. Rev. Lett.* **68**, 2981 (1992).
- [63] L. Hardy, Nonlocality for two particles without inequalities for almost all entangled states, *Phys. Rev. Lett.* **71**, 1665 (1993).
- [64] J. Barrett, A. Kent, and S. Pironio, Maximally Nonlocal and Monogamous Quantum Correlations, *Physical Review Letters* **97**, 10.1103/PhysRevLett.97.170409 (2006), arXiv:quant-ph/0605182.

- [65] M. Navascues, E. Wolfe, D. Rosset, and A. Pozas-Kerstjens, Genuine Network Multipartite Entanglement, *Phys. Rev. Lett.* **125**, 240505 (2020), arXiv:2002.02773.
- [66] Throughout the proof we assume $k \geq 2$. If $k = 1$, there are only two parties sharing bipartite pure entangled states, so the network is nonlocal by Refs. [26, 27].
- [67] Note that, while each of the terms on the right-hand side may depend on the whole of each party's input χ_i , the left-hand side does not, because the distribution is of the form (18). That is, the resulting EPR2 decomposition of P_+ holds for any fixed value of the inputs $\{x_i^k\}_{k \leq K}$ on the left- and right-hand sides.
- [68] Note that, once more, the distribution obtained by summing over only some of the digits of a party's output still depends on the whole input as it may be signalling in the different digits of the party's input. However, as in Theorem 1, these extra inputs can be fixed to an arbitrary value as the left-hand side is independent of them.