

Universal subdiffusion in strongly tilted many-body systems

Pengfei Zhang¹

¹*Institute for Quantum Information and Matter and Walter Burke Institute for Theoretical Physics, California Institute of Technology, Pasadena, California 91125, USA*

(Dated: December 30, 2021)

The quantum dynamics away from equilibrium is of fundamental interest for interacting many-body systems. In this letter, we study tilted many-body systems using the effective Hamiltonian derived from the microscopic description. We first give general arguments for the density relaxation rate satisfying $1/\tau \propto k^4$ for a large class of systems, including the Fermi Hubbard model case as observed in the recent experiment [1]. Here k is the wave vector of the density wave. The main ingredients are the emergence of the reflection symmetry and dipole moment conservation to the leading non-trivial order of the large tilted strength. To support our analysis, we then construct a solvable model with large local Hilbert space dimension by coupling sites described by the Sachdev-Ye-Kitaev models, where the density response can be computed explicitly. The tilt strength and the temperature dependence of the subdiffusion constant are also discussed.

Introduction In recent years, much understanding of strongly interacting systems has been gained thanks to the quantum simulation with cold atom systems. By using the quantum gas microscope for fermions [2–7], the antiferromagnetic correlation [8–11] has been observed in the Fermi Hubbard model, which is suspected to be related to the understanding of the high-temperature superconductivity. Besides observables in thermal ensembles, non-equilibrium quantum dynamics can also be used as a probe of underlying physics. As an example, the spin [12] and charge [13] diffusion have been studied experimentally, which shows the bad-metal behavior. Except for these standard transport experiments, the precise control of the Fermi Hubbard model in optical lattice also paves a new way to study physics in some extreme conditions. In a recent work [1], by applying an additional laser beam, authors realize a strong linear potential, or tilt, in the 2D Fermi Hubbard model. This is an analogy of applying a strong electric field in solid-state systems. A crossover between a traditional diffusion in the weak tilt limit [14] and a new subdiffusion behavior in the strong tilt limit has been observed.

In this work, we find this subdiffusion behavior is universal for a large class of tilted systems. Using the large tilt strength expansion, we obtain the low-energy effective Hamiltonian which governs the dynamics of the strongly tilted system. The subdiffusion behavior is then found to be a direct consequence of the emergent reflection symmetry and the dipole moment conservation. We also analyze the scaling of the subdiffusion constant with respect to the tilt strength. The analysis is under the assumption of hydrodynamical behavior, which is motivated by the experiments. Note that this is different from the low-dimensional models [15–18] which show transitions between ergodic and nonergodic phases. Very recently, subdiffusion behavior is also observed in a specific circuit model that conserves the dipole momentum [19], where both symmetries are imposed from the beginning [40], and in more sophisticated fracton models [20, 21]. To further support our analysis, we construct a solvable model by adding strong tilted potential to the quadratically coupled Sachdev-Ye-Kitaev model (SYK) [22–25]. Without the tilted potential, the model is a strongly




FIG. 1: A schematic of the tilted models on a two-dimensional square lattice. J denotes the hopping strength and F is the slope of the linear potential. Balls with different colors indicate atoms with different internal states. We focus on the relaxation of the initial density wave $\rho(j_x, t) = \rho_0 + A(t) \cos(k j_x)$.

interacting non-Fermi liquid with diffusive charge transport at finite temperature [25], which provides an ideal platform for our study. We explicitly compute the density response in the leading order of large tilted potential which shows the expected subdiffusion behavior.

General Analysis In this work, we consider quantum many-body systems with additional tilted potential. To be concrete, we focus on the two-dimensional square lattice with nearest-neighbor hopping and on-site interactions, as illustrated in

Fig. 1. The Hamiltonian reads

$$H = - \sum_{\mathbf{j},a} \left(J_x c_{\mathbf{j}+\hat{x},a}^\dagger c_{\mathbf{j},a} + J_y c_{\mathbf{j}+\hat{y},a}^\dagger c_{\mathbf{j},a} + \text{H.C.} \right) + \sum_{\mathbf{j}} H^{\text{int}}(c_{\mathbf{j},a}^\dagger, c_{\mathbf{j},a}) - \sum_{\mathbf{j}} F j_x n_{\mathbf{j}}. \quad (1)$$

Here $c_{\mathbf{j},a}$ ($c_{\mathbf{j},a}^\dagger$) is the annihilation (creation) operator on site $\mathbf{j} = (j_x, j_y)$ with additional internal-state index a . We have set the lattice constant $a_{\text{Lattice}} = 1$ and assumed the interactions preserve the local particle number $n_{\mathbf{j}} \equiv \sum_a n_{\mathbf{j},a} = \sum_a c_{\mathbf{j},a}^\dagger c_{\mathbf{j},a}$. For the Fermi-Hubbard case [1], we have $a \in \{\uparrow, \downarrow\}$ and $H^{\text{int}}(c_{\mathbf{j},a}^\dagger, c_{\mathbf{j},a}) = U n_{\mathbf{j},\uparrow} n_{\mathbf{j},\downarrow}$. The last term in (1) presents a tilted potential in the x direction. Note that although in this letter we only give explicit examples in term of fermions, the general argument presented below also applies for interacting bosons or boson-fermion mixtures.

We are interested in charge diffusion dynamics $\rho(\mathbf{j}, t) = \langle \psi | n_{\mathbf{j}}(t) | \psi \rangle$ with some initial state $|\psi\rangle$. For initial configurations with density waves $\rho(j_x, t) = \rho_0 + A(t) \cos(k j_x)$, we expect its amplitude decays exponentially as $A(t) \propto e^{-t/\tau}$ for general interacting systems [41]. We would like to understand the relation between τ and the wave vector k of the density wave. To analyze this problem, we first consider applying a gauge transformation $\mathcal{U} = \exp(i \sum_{\mathbf{j}} F j_x n_{\mathbf{j}} t)$. The Hamiltonian becomes

$$\tilde{H}(t) = - \sum_{\mathbf{j},a} \left(J_x e^{-iFt} c_{\mathbf{j}+\hat{x},a}^\dagger c_{\mathbf{j},a} + J_y c_{\mathbf{j}+\hat{y},a}^\dagger c_{\mathbf{j},a} + \text{H.C.} \right) + \sum_{\mathbf{j}} H^{\text{int}}(c_{\mathbf{j},a}^\dagger, c_{\mathbf{j},a}). \quad (2)$$

$\tilde{H}(t)$ is time-periodic with frequency F . For F much larger than any other energy scales, one can make a high-frequency expansion to obtain a Floquet effective Hamiltonian, which governs the dynamics of the system at integer times of the driven period $T = 2\pi/F$. The Floquet effective Hamiltonian H_{eff} is defined as

$$U(T) = \mathcal{T} \exp\left(-i \int_0^T \tilde{H}(t) dt\right) = \exp(-i H_{\text{eff}} T). \quad (3)$$

Here \mathcal{T} is the real-time ordering operator. Defining $\tilde{H}(t) = \sum_{m=0,\pm 1} H_m e^{imFt}$ with

$$H_0 = -J_y \sum_{\mathbf{j},a} \left(c_{\mathbf{j}+\hat{y},a}^\dagger c_{\mathbf{j},a} + \text{H.C.} \right) + \sum_{\mathbf{j}} H^{\text{int}}(c_{\mathbf{j},a}^\dagger, c_{\mathbf{j},a}), \quad (4)$$

$$H_1 = -J_x \sum_{\mathbf{j},a} c_{\mathbf{j},a}^\dagger c_{\mathbf{j}+\hat{x},a} = H_{-1}^\dagger,$$

the effective Hamiltonian reads [26]:

$$H_{\text{eff}} \equiv H_0 + \Delta H = H_0 + \frac{[H_{-1}, [H_0, H_1]]}{F^2}. \quad (5)$$

Here we keep terms to the order of $(|H|/F)^2$, with $|H|$ being the typical energy scale of hopping or interaction terms.

Importantly, the term proportional to $1/F$ vanishes since $[H_1, H_{-1}] = 0$ and the leading order non-trivial correlation between sites with different x comes from the $1/F^2$ term. It is also straightforward to show that only the interaction term in H_0 contributes to ΔH . However, for the present discussion, the specific form of this term is not significant.

For strongly interacting systems, it is plausible to assume the evolution of (coarse-grained) density $\rho(\mathbf{x}, t)$ can be described by the hydrodynamical equation $\partial_t \rho(\mathbf{x}, t) + \nabla \cdot \mathbf{J}(\mathbf{x}, t) = 0$ in the long-time limit. The existence of the subdiffusion behavior (in x direction) in general interacting many-body systems with large tilted potential comes from two observations:

Firstly, the system has the translation symmetry and an emergent reflection symmetry \mathcal{P} , which generates $x \rightarrow -x$. This can be understood by realizing that the microscopic Hamiltonian (1) satisfies

$$\mathcal{P} H(F) \mathcal{P} = H(-F). \quad (6)$$

Consequently, we have an emergent reflection symmetry $\mathcal{P} H_{\text{eff}} \mathcal{P} = H_{\text{eff}}$ since the effective Hamiltonian only depends on F^2 . This implies that if take a gradient expansion of J_x in terms of ρ , it should only contain an odd number of derivatives

$$J_x(\mathbf{x}, t) = -D^{(1)} \partial_x \rho(\mathbf{x}, t) + D^{(3)} \partial_x^3 \rho(\mathbf{x}, t) + \dots \quad (7)$$

Here $D^{(1)}$ is the usual charge diffusion constant, while $D^{(3)}$ is the subdiffusion constant.

Secondly, the effective Hamiltonian conserves the dipole moment $\mathcal{D} = \sum_{\mathbf{j}} j_x n_{\mathbf{j}}$:

$$[\mathcal{D}, H_{\text{eff}}] = 0, \quad (8)$$

since $[\mathcal{D}, H_0] = 0$ and $[\mathcal{D}, H_{\pm 1}] = \mp H_{\pm 1}$. This emergence of dipole moment conservation can also be directly understood as taking the large F limit of (1). As a result, we expect local conservation of dipole moment

$$\partial_t d = x \partial_x J_x = \partial_x [x J_x - D^{(1)} \rho + D^{(3)} \partial_x^2 \rho] \equiv \partial_x J_x^d. \quad (9)$$

Here we consider only variations in the x direction. $d(x, t) = x \rho(x, t)$ is the local dipole moment with current

$$J_x^d = x J_x - D^{(1)} \rho + D^{(3)} \partial_x^2 \rho.$$

Since the Hamiltonian is translation invariant, we expect $J_x^d = 0$ for a homogeneous charge filling. As a result, we should have $D^{(1)} = 0$ and the hydrodynamical equation writes

$$\partial_t \rho(x, t) + D^{(3)} \partial_x^4 \rho = 0, \quad (10)$$

which leads to the relaxation rate $1/\tau \propto k^4$. Similar subdiffusion equation has also been found recently in fracton models [21] and specific circuit model with dipole moment conservation [19]. Our analysis here can be directly applied to the experimental case of the Fermi Hubbard model [1]. Since the analysis is based on the symmetry of the system and does not depend on the details of the local interaction term, the

The phase fluctuation can be introduced as

$$c_{j,a,\pm}(t) \rightarrow e^{-i\phi_{\pm}(x,t)} c_{j,a}(t).$$

Here $\phi_{\pm}(x)$ is a slow-varying phase factor. The non-trivial coupling terms between ϕ and fermions read

$$S_K^c = \sum_m \int dt m \left[\sum_{j,a} c_{j,a,m}^\dagger (\partial_t \phi_m) c_{j,a,m} + \sum_{j,abcd} \frac{J_x^2 U_{ab;cd}^j}{2F^2} \times (c_{j+\hat{x},a,m}^\dagger c_{j-\hat{x},b,m}^\dagger c_{j,c,m} c_{j,d,m} e^{i\partial_x^2 \phi_m} + \text{H.C.}) \right]. \quad (17)$$

Since we focus on the density response $\Pi_R(t, \mathbf{j}) = -i\theta(t)[n_j(t), n_0]$. We further introduce source term:

$$S_K^s = \sum_m \int dt m \left[\sum_j n_{j,m} \mathcal{J}_{j,m} \right]. \quad (18)$$

which leads to $\Pi_R = -\frac{i}{2} \frac{\partial^2 \log Z}{\partial \mathcal{J}_{cl} \partial \mathcal{J}_q}$ with $O_{cl/q} \equiv (O_+ \pm O_-)/2$. To calculate Π_R to the leading order of $1/N$, we keep the action of ϕ to the quadratic order [24, 25]. The action takes the form:

$$\begin{aligned} \frac{iS_{q,cl}^\phi[\phi, J]}{N} &= \text{diagram 1} + \text{diagram 2} + \text{diagram 3} \\ &= \sum_{\omega, k_x} \Lambda_1(\omega) (\partial_t \phi_{cl} + \mathcal{J}_{cl})(\omega, k_x) (\partial_t \phi_q + \mathcal{J}_q)(-\omega, -k_x) \\ &\quad + k_x^4 (\Lambda_2(\omega) - \Lambda_2(0)) \phi_{cl}(\omega, k_x) \phi_q(-\omega, -k_x). \end{aligned} \quad (19)$$

Here the wavy line represents the phase fluctuation ϕ or source \mathcal{J} . Focusing on the retarded part, we only maintain terms contain both the classical and the quantum components. The first term describes the response of a homogeneous source field. As a result in the low-energy limit $\Lambda_1(\omega \rightarrow 0) \rightarrow 2iK$ with KN being the compressibility [33]. The other coefficient writes:

$$\begin{aligned} \Lambda_2(t) &= -\frac{J_x^4}{2F^4} (G_R(t) \Sigma_K(-t) + G_A(-t) \Sigma_K(-t) \\ &\quad + \Sigma_R(t) G_K(-t) + \Sigma_A(-t) G_K(-t)). \end{aligned} \quad (20)$$

Here we have defined the retarded Green's function $G_{R/A}(t) = -i\theta(\pm t) \{c_{j,a}^\dagger(t), c_{j,a}(0)\}$ with $\Sigma_{R/A}(t)$ being corresponding self-energy. The Keldysh component reads $G_K(\omega) = (G_R(\omega) - G_A(\omega))(1 - 2n_F(\omega)) \equiv -2\pi i A_G(\omega)(1 - 2n_F(\omega))$ and similarly for Σ_K . For small ω , we expand $\Lambda_2(\omega) - \Lambda_2(0) \approx -2\omega KD^{(3)}$ with

$$KD^{(3)} = -\frac{\pi J_x^4}{F^4} \int dq_0 A_G(q_0) A_\Sigma(q_0) \partial_{q_0} n_F(q_0) \approx \frac{J_x^4}{2F^4}. \quad (21)$$

We have used the conformal Green's function (15). Using the effective action (19), the density response can be computed as:

$$\Pi_R(\omega, k_x) = \frac{KND^{(3)}k_x^4}{-i\omega + D^{(3)}k_x^4}. \quad (22)$$

The pole $-i\omega = -D^{(3)}k_x^4$ leads to the correct hydrodynamical behavior as in (10). We see the subdiffusion constant $D^{(3)}$ has a scaling of $1/F^4$ consistent with our previous analysis. Interestingly, the factor of U in $KD^{(3)}$ cancels out, which is distinct from the expectation for weakly interacting systems.

We finally comment on the different temperature dependence of the (sub)diffusion constant compared to the result with $F = 0$, as computed in [25]. Without the tilted potential, the diffusion constant is proportional to $1/T$, leading to a linear resistivity as in strange metals. This is because the correlation between different sites is from a quadratic term, which is relevant at the low energy limit with a scaling dimension $1/2$. However, here the correlation is from a term that is marginal, leading to a constant subdiffusion constant. More generally, we could consider generalizing the local interaction term to be the SYK_q model. While scaling of the diffusion constant with $F = 0$ changes, the subdiffusion constant for the strongly tilted potential remains a constant. In this sense, we expect a weaker temperature dependence for $D^{(3)}$ in strongly correlated systems.

Outlook In this work we have analyzed quantum many-body systems in strongly tilted potential. We find the charge transport show subdiffusion behavior due to the emergent reflection symmetry and dipole momentum conservation. We further determine the subdiffusion constant in a solvable coupled SYK model as an explicit example.

Several interesting questions can be asked along this direction. Firstly, we could consider generalizing the tilted potential to impose dynamical conservation of high-order moments. However, this would break the translation symmetry and make the analysis harder. Secondly, here we have considered the case where F is the largest energy scale of the system. We could ask what happens if we instead consider the near-resonant case with $F \approx U$ in the Hubbard model case [35–39]. It is also interesting to study the possible interplay between the spin diffusion and the charge diffusion in the strongly tilted systems. We postpone all these questions for future studies.

Acknowledgment We would like to thank Lei Pan for bringing our attention to the experiment [1].

-
- [1] E. Guardado-Sanchez, A. Morningstar, B. M. Spar, P. T. Brown, D. A. Huse, and W. S. Bakr, *Physical Review X* **10**, 011042 (2020).
 - [2] E. Haller, J. Hudson, A. Kelly, D. A. Cotta, B. Peaudecerf, G. D. Bruce, and S. Kuhr, *Nature Physics* **11**, 738 (2015).
 - [3] L. W. Cheuk, M. A. Nichols, M. Okan, T. Gersdorf, V. V. Ramasesh, W. S. Bakr, T. Lompe, and M. W. Zwierlein, *Physical review letters* **114**, 193001 (2015).
 - [4] G. J. Edge, R. Anderson, D. Jervis, D. C. McKay, R. Day, S. Trotzky, and J. H. Thywissen, *Physical Review A* **92**, 063406 (2015).
 - [5] A. Omran, M. Boll, T. A. Hilker, K. Kleinlein, G. Salomon, I. Bloch, and C. Gross, *Physical review letters* **115**, 263001 (2015).

- [6] D. Greif, M. F. Parsons, A. Mazurenko, C. S. Chiu, S. Blatt, F. Huber, G. Ji, and M. Greiner, *Science* **351**, 953 (2016).
- [7] L. W. Cheuk, M. A. Nichols, K. R. Lawrence, M. Okan, H. Zhang, and M. W. Zwierlein, *Physical review letters* **116**, 235301 (2016).
- [8] M. F. Parsons, A. Mazurenko, C. S. Chiu, G. Ji, D. Greif, and M. Greiner, *Science* **353**, 1253 (2016).
- [9] M. Boll, T. A. Hilker, G. Salomon, A. Omran, J. Nespolo, L. Pollet, I. Bloch, and C. Gross, *Science* **353**, 1257 (2016).
- [10] L. W. Cheuk, M. A. Nichols, K. R. Lawrence, M. Okan, H. Zhang, E. Khatami, N. Trivedi, T. Paiva, M. Rigol, and M. W. Zwierlein, *Science* **353**, 1260 (2016).
- [11] T. A. Hilker, G. Salomon, F. Grusdt, A. Omran, M. Boll, E. Demler, I. Bloch, and C. Gross, *Science* **357**, 484 (2017).
- [12] M. A. Nichols, L. W. Cheuk, M. Okan, T. R. Hartke, E. Mendez, T. Senthil, E. Khatami, H. Zhang, and M. W. Zwierlein, *Science* **363**, 383 (2019).
- [13] P. T. Brown, D. Mitra, E. Guardado-Sanchez, R. Nourafkan, A. Reymbaut, C.-D. Hébert, S. Bergeron, A.-M. Tremblay, J. Kokalj, D. A. Huse, *et al.*, *Science* **363**, 379 (2019).
- [14] S. Mandt, A. Rapp, and A. Rosch, *Physical review letters* **106**, 250602 (2011).
- [15] P. Sala, T. Rakovszky, R. Verresen, M. Knap, and F. Pollmann, *Physical Review X* **10**, 011047 (2020).
- [16] E. van Nieuwenburg, Y. Baum, and G. Refael, *Proceedings of the National Academy of Sciences* **116**, 9269 (2019).
- [17] M. Schulz, C. Hooley, R. Moessner, and F. Pollmann, *Physical review letters* **122**, 040606 (2019).
- [18] A. Morningstar, V. Khemani, and D. A. Huse, arXiv preprint arXiv:2004.00096 (2020).
- [19] J. Feldmeier, P. Sala, G. de Tomasi, F. Pollmann, and M. Knap, arXiv preprint arXiv:2004.00635 (2020).
- [20] M. Pretko, X. Chen, and Y. You, arXiv preprint arXiv:2001.01722 (2020).
- [21] A. Gromov, A. Lucas, and R. M. Nandkishore, arXiv preprint arXiv:2003.09429 (2020).
- [22] A. Kitaev, in *Talk given at the Fundamental Physics Prize Symposium*, Vol. 10 (2014).
- [23] J. Maldacena and D. Stanford, *Physical Review D* **94**, 106002 (2016).
- [24] P. Zhang, *Physical Review B* **96**, 205138 (2017).
- [25] X.-Y. Song, C.-M. Jian, and L. Balents, *Physical review letters* **119**, 216601 (2017).
- [26] A. Eckardt, *Reviews of Modern Physics* **89**, 011004 (2017).
- [27] A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, *Reviews of Modern Physics* **68**, 13 (1996).
- [28] S. Banerjee and E. Altman, *Physical Review B* **95**, 134302 (2017).
- [29] X. Chen, R. Fan, Y. Chen, H. Zhai, and P. Zhang, *Physical review letters* **119**, 207603 (2017).
- [30] S.-K. Jian and H. Yao, arXiv preprint arXiv:1703.02051 (2017).
- [31] Y. Gu, X.-L. Qi, and D. Stanford, *Journal of High Energy Physics* **2017**, 125 (2017).
- [32] R. A. Davison, W. Fu, A. Georges, Y. Gu, K. Jensen, and S. Sachdev, *Physical Review B* **95**, 155131 (2017).
- [33] Y. Gu, A. Kitaev, S. Sachdev, and G. Tarnopolsky, *Journal of High Energy Physics* **2020**, 1 (2020).
- [34] A. Kamenev, *Field Theory of Non-Equilibrium Systems* (Cambridge University Press, 2011).
- [35] M. Bukov, M. Kolodrubetz, and A. Polkovnikov, *Physical review letters* **116**, 125301 (2016).
- [36] A. Itin and M. Katsnelson, *Physical review letters* **115**, 075301 (2015).
- [37] F. Görg, M. Messer, K. Sandholzer, G. Jotzu, R. Desbuquois, and T. Esslinger, *Nature* **553**, 481 (2018).
- [38] W. Xu, W. Morong, H.-Y. Hui, V. W. Scarola, and B. DeMarco, *Physical Review A* **98**, 023623 (2018).
- [39] N. Sun, P. Zhang, and H. Zhai, *Physical Review A* **99**, 043629 (2019).
- [40] In this paper, authors also study the consequence of the higher-moment conservation.
- [41] Note that this is an important assumption, which excludes the possibility of non-ergodicity.
- [42] We have used the fact that the hopping term in H_0 commutes with H_1 .