

# Some variations of a “divergent” Ramanujan-type $q$ -supercongruence

Victor J. W. Guo

School of Mathematical Sciences, Huaiyin Normal University, Huai’an 223300, Jiangsu,  
People’s Republic of China  
jwguo@hytc.edu.cn

**Abstract.** Using the  $q$ -Wilf–Zeilberger method and a  $q$ -analogue of a “divergent” Ramanujan-type supercongruence, we give several  $q$ -supercongruences modulo the fourth power of a cyclotomic polynomial. One of them is a  $q$ -analogue of a supercongruence recently proved by Wang: for any prime  $p > 3$ ,

$$\sum_{k=0}^{p-1} (3k-1) \frac{(\frac{1}{2})_k (-\frac{1}{2})_k^2}{k!^3} 4^k \equiv p - 2p^3 \pmod{p^4},$$

where  $(a)_k = a(a+1)\cdots(a+k-1)$  is the Pochhammer symbol.

*Keywords:* central  $q$ -binomial coefficients; Wilf–Zeilberger method;  $q$ -WZ method;  $q$ -WZ pair; cyclotomic polynomials.

*2010 Mathematics Subject Classifications:* 11B65 (Primary) 05A10, 05A30 (Secondary)

## 1. Introduction

By making use of the Wilf–Zeilberger (abbr. WZ) method [25,26], Guillera and Zudilin [3] established the following supercongruence: for any odd prime  $p$ ,

$$\sum_{k=0}^{(p-1)/2} \frac{3k+1}{16^k} \binom{2k}{k}^3 \equiv p \pmod{p^3}, \quad (1.1)$$

We can also sum  $k$  in (1.1) up to  $p-1$ , since the  $p$ -adic order of  $(\frac{1}{2})_k/k!$  is 1 for  $k$  in the range  $(p+1)/2 \leq k \leq p-1$ . In the spirit of [27], the supercongruence (1.1) corresponds to a divergent Ramanujan-type series for  $1/\pi$ :

$$\sum_{k=0}^{\infty} \frac{3k+1}{16^k} \binom{2k}{k}^3 \stackrel{“=”}{=} \frac{-2i}{\pi} \quad (1.2)$$

(see [3, (47)]). Here the summation in (1.2) must be understood as the analytic continuation of the corresponding hypergeometric series.

Still using the WZ method and the divisibility result: for  $n > 1$ ,

$$2n \binom{2n}{n} \left| \sum_{k=0}^{n-1} (3k+1) \binom{2k}{k}^3 16^{n-k-1}, \right.$$

which was conjectured by Z.-W. Sun [20] and confirmed by Mao and Zhang [15], B.Y. Sun [19] proved the following result: for  $n > 1$ ,

$$2n \binom{2n}{n} \left| \sum_{k=0}^{n-1} \frac{6k^4}{2k-1} \binom{2k}{k}^3 16^{n-k-1}. \right. \quad (1.3)$$

Motivated by B.Y. Sun's work, we found the following supercongruence: for any prime  $p > 3$ ,

$$\sum_{k=0}^{p-1} \frac{6k^4}{16^k(2k-1)} \binom{2k}{k}^3 \equiv p + 2p^3 \pmod{p^4}. \quad (1.4)$$

We shall prove the supercongruence (1.4) by establishing its  $q$ -analogue. Recall that the  $q$ -shifted factorial is defined by  $(a; q)_0 = 1$  and  $(a; q)_n = (1-a)(1-aq) \cdots (1-aq^{n-1})$  for  $n \geq 1$ , and the  $q$ -integer is defined as  $[n] = [n]_q = (1-q^n)/(1-q)$  (see [1]). Moreover, the  $n$ -th cyclotomic polynomial  $\Phi_n(q)$  is given by

$$\Phi_n(q) := \prod_{\substack{1 \leq k \leq n \\ \gcd(k, n) = 1}} (q - \zeta^k),$$

where  $\zeta$  is an  $n$ -th primitive root of unity. Our  $q$ -analogue of (1.4) can be stated as follows.

**Theorem 1.1.** *Let  $n > 1$  be an odd integer. Then*

$$\begin{aligned} & \sum_{k=0}^{n-1} \frac{[3k][2k][k]^2}{[2k-1](-q; q)_k^4} \binom{2k}{k}^3 q^{-(k^2+3k)/2} \\ & \equiv [n]q^{-(n+1)/2} + (1+q)[n]^3 + \frac{(n^2-1)(1-q)^2}{24} [n]^3 q^{-(n+1)/2} \pmod{[n]\Phi_n(q)^3}, \end{aligned} \quad (1.5)$$

where  $\binom{2k}{k} = (q; q)_{2k}/(q; q)_k^2$  denotes the central  $q$ -binomial coefficient.

Here we say that two rational functions  $A(q)$  and  $B(q)$  are congruent modulo a polynomial  $P(q)$  if and only if  $P(q)$  divides the numerator of the reduced form of  $A(q) - B(q)$  in the polynomial ring  $\mathbb{Z}[q]$ .

It is easy to see that, letting  $n = p > 3$  be a prime and taking  $q \rightarrow 1$  in (1.5), we obtain the supercongruence (1.4). Furthermore, we can also deduce from (1.5) that, for any prime  $p > 3$  and  $r \geq 2$ ,

$$\sum_{k=0}^{p^r-1} \frac{6k^4}{16^k(2k-1)} \binom{2k}{k}^3 \equiv p^r \pmod{p^{r+3}}.$$

Recently, via the WZ method and the summation package `Sigma` [17], Wang [21] proved the following supercongruence: for any prime  $p > 3$ ,

$$\sum_{k=0}^{p-1} (3k-1) \frac{\left(\frac{1}{2}\right)_k \left(-\frac{1}{2}\right)_k^2}{k!^3} 4^k \equiv p - 2p^3 \pmod{p^4}, \quad (1.6)$$

where  $(a)_k = a(a+1)\cdots(a+k-1)$  is the Pochhammer symbol. This also extends a conjectural result of the author and Schlosser [11, Conjecture 6.2].

In this paper, we shall give a  $q$ -analogue of (1.6) as follows:

**Theorem 1.2.** *Let  $n > 1$  be an odd integer. Then*

$$\begin{aligned} & \sum_{k=0}^{n-1} [3k-1] \frac{(q; q^2)_k (q^{-1}; q^2)_k^2}{(q; q)_k^2 (q^2; q^2)_k} q^{(3k-k^2)/2} \\ & \equiv [n] q^{-(n+1)/2} - (1+q)[n]^3 + \frac{(n^2-1)(1-q)^2}{24} [n]^3 q^{-(n+1)/2} \pmod{[n]\Phi_n(q)^3}. \end{aligned} \quad (1.7)$$

As before, letting  $n = p > 3$  be a prime and taking  $q \rightarrow 1$  in (1.7), we are led to (1.4). Moreover, it follows from (1.5) that, for any prime  $p > 3$  and  $r \geq 2$ ,

$$\sum_{k=0}^{p^r-1} (3k-1) \frac{\left(\frac{1}{2}\right)_k \left(-\frac{1}{2}\right)_k^2}{k!^3} 4^k \equiv p^r \pmod{p^{r+3}}.$$

We shall prove Theorems 1.1 and 1.2 by making use of the  $q$ -WZ method [25, 26] and the following  $q$ -supercongruence: for odd  $n$ ,

$$\begin{aligned} & \sum_{k=0}^{n-1} [3k+1] \frac{(q; q^2)_k^3 q^{-\binom{k+1}{2}}}{(q; q)_k^2 (q^2; q^2)_k} \\ & \equiv q^{(1-n)/2} [n] + \frac{(n^2-1)(1-q)^2}{24} q^{(1-n)/2} [n]^3 \pmod{[n]\Phi_n(q)^3}. \end{aligned} \quad (1.8)$$

This  $q$ -supercongruence was originally conjectured in [7] and recently proved in [9] with the help of the “creative microscoping” method [12] and the Chinese remainder theorem. It is easy to see that (1.1) follows from (1.8) by taking  $n = p$  and  $q \rightarrow 1$ . The  $n = p^r$  being an odd prime power and  $q \rightarrow 1$  case of (1.8) was conjectured by Z.-W. Sun [20]. We point out that some other interesting  $q$ -supercongruences were given in [2, 4–11, 13, 14, 16, 18, 22–24, 28].

The paper is organized as follows. We first give two lemmas in the next section. The proofs of Theorems 1.1 and 1.2 will be given in Sections 3 and 4, respectively. Two more similar  $q$ -supercongruences are given in Section 5. Finally, we propose a related open problem in Section 6.

## 2. Two lemmas

In this section we give three simple  $q$ -congruences. The first one may be deemed a  $q$ -analogue of Fermat's little theorem  $2^{p-1} \equiv 1 \pmod{p}$  for any odd prime  $p$ . The third one is a generalization of a recent result of Wang and Ni [22, Lemma 2.2].

**Lemma 2.1.** *Let  $n > 1$  be an odd integer. Then*

$$(-q; q)_{n-1} \equiv 1 \pmod{\Phi_n(q)}. \quad (2.1)$$

*Proof.* It is well known that

$$\frac{x^n - 1}{x - 1} = \prod_{k=1}^{n-1} (x - \zeta^k), \quad (2.2)$$

where  $\zeta$  is an  $n$ -th primitive root of unity. Letting  $x = -1$  in (2.2), we get  $(-\zeta; \zeta)_{n-1} = 1$ , which is equivalent to (2.1).  $\square$

**Lemma 2.2.** *Let  $n > 1$  be an odd integer. Then*

$$\frac{(q; q^2)_n}{(1-q)(q; q)_{n-1}} \equiv [n] \pmod{[n]\Phi_n(q)}, \quad (2.3)$$

$$\frac{(q; q^2)_{n-1}}{(q; q)_{n-1}} \equiv -[n]q \pmod{[n]\Phi_n(q)}. \quad (2.4)$$

*Proof.* It is easy to see that

$$\frac{(q; q^2)_n}{(1-q)(q; q)_{n-1}} = [n] \begin{bmatrix} 2n \\ n \end{bmatrix} \frac{1}{(-q; q)_n}, \quad (2.5)$$

and

$$\begin{aligned} \begin{bmatrix} 2n \\ n \end{bmatrix} \frac{1}{(-q; q)_n} &= \frac{(q; q^2)_{(n-1)/2} (q^{n+2}; q^2)_{(n-1)/2}}{(q; q)_{n-1}} \\ &\equiv \frac{(q; q^2)_{(n-1)/2} (q^2; q^2)_{(n-1)/2}}{(q; q)_{n-1}} = 1 \pmod{\Phi_n(q)} \end{aligned} \quad (2.6)$$

in view of  $q^n \equiv 1 \pmod{\Phi_n(q)}$ . Since  $\begin{bmatrix} 2n \\ n \end{bmatrix}$  is a polynomial in  $q$  and  $[n]$  is relatively prime to  $(-q; q)_n$  for odd  $n$ , the  $q$ -congruence (2.3) immediately follows from (2.5) and (2.6).

Observing that

$$\begin{aligned} \frac{(q; q^2)_{n-1}}{(q; q)_{n-1}} &= \frac{1}{[2n-1]} \frac{(q; q^2)_n}{(1-q)(q; q)_{n-1}}, \\ [2n-1] &= (1 - q^{2n-1}) / (1 - q) \equiv -q^{-1} \pmod{\Phi_n(q)}, \end{aligned}$$

and  $[2n-1]$  and  $[n]$  are relatively prime polynomials in  $q$ , we deduce (2.4) from (2.3).  $\square$

### 3. Proof of Theorem 1.1

Define two functions  $F(n, k)$  and  $G(n, k)$  as follows:

$$F(n, k) = [3n + 2k + 1] \frac{(q; q^2)_n (q^{2k+1}; q^2)_n^2 q^{-\binom{n+1}{2} - (2n+1)k}}{(q; q)_n^2 (q^2; q^2)_n}, \quad (3.1)$$

$$G(n, k) = -\frac{(1 + q^{n+2k-1})(q; q^2)_n (q^{2k+1}; q^2)_{n-1}^2 q^{-\binom{n}{2} - (2n-1)k}}{(1-q)(q; q)_{n-1}^2 (q^2; q^2)_{n-1}}, \quad (3.2)$$

where we have assumed that  $1/(q^2; q^2)_n = 0$  for any negative integer  $n$ . It is easy to check that

$$F(n, k-1) - F(n, k) = G(n+1, k) - G(n, k). \quad (3.3)$$

That is, the functions  $F(n, k)$  and  $G(n, k)$  form a  $q$ -WZ pair.

We now let  $m > 1$  be an odd integer. Summing (3.3) over  $n$  from 0 to  $m-1$ , we obtain

$$\sum_{n=0}^{m-1} F(n, k-1) - \sum_{n=0}^{m-1} F(n, k) = G(m, k) - G(0, k) = G(m, k). \quad (3.4)$$

In light of (2.1) and (2.3), we have

$$\begin{aligned} G(m, 1) &= -\frac{(1 + q^{m+1})(q; q^2)_m (q^3; q^2)_{m-1}^2 q^{-\binom{m}{2} - 2m+1}}{(1-q)(q; q)_{m-1}^3 (-q; q)_{m-1}} \\ &= -\frac{(1 + q^{m+1})(q; q^2)_m^3 q^{-\binom{m}{2} - 2m+1}}{(1-q)^3 (q; q)_{m-1}^3 (-q; q)_{m-1}} \\ &\equiv -(1+q)q[m]^3 \pmod{[m]^3 \Phi_m(q)}, \end{aligned} \quad (3.5)$$

since  $q^m \equiv 1 \pmod{\Phi_m(q)}$ . Combining (3.4) and (3.5), we conclude that

$$\sum_{n=0}^{m-1} F(n, 1) \equiv \sum_{n=0}^{m-1} F(n, 0) + (1+q)q[m]^3 \pmod{[m]^3 \Phi_m(q)}. \quad (3.6)$$

It is easy to see that

$$\begin{aligned} \sum_{n=0}^{m-1} F(n, 1) &= \sum_{n=0}^{m-1} [3n+3] \frac{(q; q^2)_n (q^3; q^2)_n^2 q^{-\binom{n+1}{2} - (2n+1)}}{(q; q)_n^2 (q^2; q^2)_n} \\ &= \sum_{n=1}^m [3n] \frac{(q; q^2)_{n-1} (q^3; q^2)_{n-1}^2 q^{-\binom{n}{2} - (2n-1)}}{(q; q)_{n-1}^2 (q^2; q^2)_{n-1}} \\ &= \sum_{n=1}^m \frac{[3n][2n][n]^2}{[2n-1](-q; q)_n^4} \begin{bmatrix} 2n \\ n \end{bmatrix}^3 q^{-(n^2+3n)/2+1}. \end{aligned} \quad (3.7)$$

On the other hand, by (1.8) we have

$$\begin{aligned} \sum_{n=0}^{m-1} F(n, 0) &= \sum_{n=0}^{m-1} [3n+1] \frac{(q; q^2)_n^3 q^{-\binom{n+1}{2}}}{(q; q)_n^2 (q^2; q^2)_n} \\ &\equiv q^{(1-m)/2} [m] + \frac{(m^2-1)(1-q)^2}{24} q^{(1-m)/2} [m]^3 \pmod{[m] \Phi_m(q)^3}. \end{aligned} \quad (3.8)$$

Substituting (3.7) and (3.8) into (3.6), and noticing the  $m$ -th summand on the right-hand side of (3.7) is congruent to 0 modulo  $[m]^4$ , we are led to (1.5) with  $n \mapsto m$  differing only by a factor  $q$ .

*Remark.* Usually the basic hypergeometric functions satisfying the condition

$$F(n, k+1) - F(n, k) = G(n+1, k) - G(n, k)$$

are called a  $q$ -WZ pair. It is also reasonable to call the basic hypergeometric functions satisfying (3.3) a  $q$ -WZ pair (see Zudilin [27]). The  $q$ -WZ pair in the proof of Theorem 1.1 was found by the author [7] in his proof of a weaker form of (1.8) modulo  $[n] \Phi_n(q)^2$ . But the corresponding WZ pair (the limiting case  $q \rightarrow 1$ ) was first given by Guillera and Zudilin [3] in their proof of (1.1).

## 4. Proof of Theorem 1.2

Let the functions  $F(n, k)$  and  $G(n, k)$  be given by (3.1) and (3.2), respectively. Again, let  $m$  be an odd integer greater than 1. In view of (2.1), (2.3), (2.4) and  $q^m \equiv 1 \pmod{\Phi_m(q)}$ , we have

$$\begin{aligned} G(m, 0) &= -\frac{(1+q^{m-1})(q; q^2)_m (q; q^2)_{m-1}^2 q^{-\binom{m}{2}}}{(1-q)(q; q)_{m-1}^3 (-q; q)_{m-1}} \\ &\equiv -(1+q^{-1})q^2 [m]^3 \pmod{[m]^3 \Phi_m(q)}. \end{aligned} \quad (4.1)$$

It follows from (3.4) and (4.1) that

$$\sum_{n=0}^{m-1} F(n, -1) \equiv \sum_{n=0}^{m-1} F(n, 0) - (1+q)q [m]^3 \pmod{[m]^3 \Phi_m(q)}. \quad (4.2)$$

By the definition of (3.1), we have

$$\sum_{n=0}^{m-1} F(n, -1) = \sum_{n=0}^{m-1} [3n-1] \frac{(q; q^2)_n (q^{-1}; q^2)_n^2 q^{-\binom{n+1}{2} + (2n+1)}}{(q; q)_n^2 (q^2; q^2)_n}. \quad (4.3)$$

Finally, substituting (3.8) and (4.3) into (4.2), and dividing both sides by  $q$ , we arrive at (1.5) with  $n \mapsto m$ .

## 5. More similar $q$ -supercongruences

From (1.8) and (3.4) we can deduce more  $q$ -supercongruences besides (1.5) and (1.7). Here we give two such examples.

**Theorem 5.1.** *Let  $n > 3$  be an odd integer. Then, modulo  $[n]\Phi_n(q)^3$ ,*

$$\begin{aligned} & \sum_{k=0}^{n-1} [3k+5] \frac{(q; q^2)_k (q^5; q^2)_k^2}{(q; q)_k^2 (q^2; q^2)_k} q^{-(k^2+9k)/2} \\ & \equiv [n]q^{(5-n)/2} + (1+q)q^3[n]^3 + \frac{(1+q^3)q^4}{(1+q+q^2)^2} [n]^3 + \frac{(n^2-1)(1-q)^2}{24} [n]^3 q^{(5-n)/2}. \end{aligned} \quad (5.1)$$

*Proof.* Let  $m > 3$  be an odd integer. Then

$$\sum_{n=0}^{m-1} F(n, 2) = \sum_{k=0}^{m-1} [3k+5] \frac{(q; q^2)_k (q^5; q^2)_k^2}{(q; q)_k^2 (q^2; q^2)_k} q^{-(k^2+9k)/2-2}. \quad (5.2)$$

By (3.4), we get

$$\sum_{n=0}^{m-1} F(n, 2) = \sum_{n=0}^{m-1} F(n, 0) - G(m, 1) - G(m, 2). \quad (5.3)$$

Moreover, in view of (2.3), we have

$$\begin{aligned} G(m, 2) &= -\frac{(1+q^{m+3})(q; q^2)_m (q^5; q^2)_{m-1}^2 q^{-\binom{m}{2}-4m+2}}{(1-q)(q; q)_{m-1}^3 (-q; q)_{m-1}} \\ &= -\frac{(1-q^{2m+1})^2 (1+q^{m+3})(q; q^2)_m^3 q^{-\binom{m}{2}-4m+2}}{(1-q^3)^2 (1-q)^3 (q; q)_{m-1}^3 (-q; q)_{m-1}} \\ &\equiv -\frac{(1+q^3)q^2}{(1+q+q^2)^2} [m]^3 \pmod{[m]\Phi_m(q)^3}, \end{aligned} \quad (5.4)$$

Substituting (3.5), (3.8), (5.2) and (5.4) into (5.3), we arrive at (5.1) with  $n \mapsto m$  differing only by a factor  $q^{-2}$ .  $\square$

**Theorem 5.2.** *Let  $n > 3$  be an odd integer. Then, modulo  $[n]\Phi_n(q)^3$ ,*

$$\begin{aligned} & \sum_{k=0}^{n-1} [3k-3] \frac{(q; q^2)_k (q^{-3}; q^2)_k^2}{(q; q)_k^2 (q^2; q^2)_k} q^{(7k-k^2)/2} \\ & \equiv [n]q^{-(n+3)/2} - \frac{1+q}{q} [n]^3 - \frac{1+q^3}{(1+q+q^2)^2} [n]^3 + \frac{(n^2-1)(1-q)^2}{24} [n]^3 q^{-(n+3)/2}. \end{aligned} \quad (5.5)$$

*Proof.* The proof is similar to that of Theorem 5.1. This time we need to use

$$\sum_{n=0}^{m-1} F(n, -2) = \sum_{n=0}^{m-1} F(n, 0) + G(m, 0) + G(m, -1),$$

and

$$\begin{aligned} G(m, -1) &= -\frac{(1+q^{m-3})(q; q^2)_m (q^{-1}; q^2)_{m-1}^2 q^{-\binom{m}{2}+2m-1}}{(1-q)(q; q)_{m-1}^3 (-q; q)_{m-1}} \\ &\equiv -\frac{(1+q^3)q^2}{(1+q+q^2)^2} [m]^3 \pmod{[m]\Phi_m(q)^3}. \end{aligned}$$

□

## 6. An open problem

For Sun's divisibility result (1.3), we found that the following stronger version holds:

$$4n \binom{2n}{n} \left| \sum_{k=0}^{n-1} \frac{6k^4}{2k-1} \binom{2k}{k} \right|^3 16^{n-k-1}.$$

Furthermore, we believe that the following  $q$ -version should be true.

**Conjecture 6.1.** *Let  $n$  be a positive integer. Then*

$$\frac{1}{(1+q)^3 [2n+1] \begin{bmatrix} 2n \\ n \end{bmatrix}} \sum_{k=1}^n \frac{[3k][2k][k]^2 (-q; q)_n^4}{[2k-1](-q; q)_k^4} \begin{bmatrix} 2k \\ k \end{bmatrix}^3 q^{-(k^2+3k)/2}$$

*is a Laurent polynomial in  $q$ .*

**Acknowledgments.** This work was partially supported by the National Natural Science Foundation of China (grant 11771175).

## References

- [1] G. Gasper and M. Rahman, Basic hypergeometric series, Second Edition, Encyclopedia of Mathematics and Its Applications, Vol. 96, Cambridge University Press, Cambridge, 2004.
- [2] C.-Y. Gu and V.J.W. Guo,  $q$ -Analogues of two supercongruences of Z.-W. Sun, Czechoslovak Math. J., in press; <https://doi.org/10.21136/CMJ.2020.0516-18>
- [3] J. Guillera and W. Zudilin, "Divergent" Ramanujan-type supercongruences, Proc. Amer. Math. Soc. 140 (2012), 765–777.
- [4] V.J.W. Guo, Proof of a generalization of the (B.2) supercongruence of Van Hamme through a  $q$ -microscope, Adv. Appl. Math. 116 (2020), Art. 102016.



- [5] V.J.W. Guo,  $q$ -Analogues of Dwork-type supercongruences, *J. Math. Anal. Appl.* 487 (2020), Art. 124022.
- [6] V.J.W. Guo, A  $q$ -analogue of the (A.2) supercongruence of Van Hamme for primes  $p \equiv 1 \pmod{4}$ , *Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM*, in press; <https://doi.org/10.1007/s13398-020-00854-y>
- [7] V.J.W. Guo,  $q$ -Analogues of two “divergent” Ramanujan-type supercongruences, *Ramanujan J.*, in press, <https://doi.org/10.1007/s11139-019-00161-0>
- [8] V.J.W. Guo and J.-C. Liu,  $q$ -Analogues of two Ramanujan-type formulas for  $1/\pi$ , *J. Difference Equ. Appl.* 24 (2018), 1368–1373.
- [9] V.J.W. Guo,  $q$ -Supercongruences modulo the fourth power of a cyclotomic polynomial via creative microscoping, preprint, 2019, arXiv:1912.00765.
- [10] V.J.W. Guo and M.J. Schlosser, Proof of a basic hypergeometric supercongruence modulo the fifth power of a cyclotomic polynomial, *J. Difference Equ. Appl.* 25 (2019), 921–929.
- [11] V.J.W. Guo and M.J. Schlosser, Some  $q$ -supercongruences from transformation formulas for basic hypergeometric series, preprint, December 2018; arXiv:1812.06324.
- [12] V.J.W. Guo and W. Zudilin, A  $q$ -microscope for supercongruences, *Adv. Math.* 346 (2019), 329–358.
- [13] V.J.W. Guo and W. Zudilin, A common  $q$ -analogue of two supercongruences, *Results Math.* 75 (2020), Art. 46.
- [14] J.-C. Liu and F. Petrov, Congruences on sums of  $q$ -binomial coefficients, *Adv. Appl. Math.* 116 (2020), Art. 102003.
- [15] G.-S. Mao and T. Zhang, Proof of Sun’s conjectures on super congruences and the divisibility of certain binomial sums, *Ramanujan J.* 50 (2019), 1–11.
- [16] H.-X. Ni and H. Pan, Some symmetric  $q$ -congruences modulo the square of a cyclotomic polynomial, *J. Math. Anal. Appl.* 481 (2020), Art. 123372.
- [17] C. Schneider, Symbolic summation assists combinatorics, *Sémin. Lothar. Comb.* 56 (2007), B56b.
- [18] A. Straub, Supercongruences for polynomial analogs of the Apéry numbers, *Proc. Amer. Math. Soc.* 147 (2019), 1023–1036.
- [19] B.Y. Sun, Note on a new divisibility property of binomial sums, *Util. Math.* 107 (2018), 131–135.
- [20] Z.-W. Sun, Super congruences and Euler numbers, *Sci. China Math.* 54 (2011), 2509–2535.
- [21] C. Wang, Symbolic summation methods and hypergeometric supercongruences, *J. Math. Anal. Appl.* 488 (2020), Art. 124068.
- [22] C. Wang and H.-X. Ni, Some  $q$ -congruences arising from certain identities, preprint, March 2020; arXiv:2003.10883.
- [23] X. Wang and M. Yue, Some  $q$ -supercongruences from Watson’s  ${}_8\phi_7$  transformation formula, *Results Math.*, in press; <https://doi.org/10.1007/s00025-020-01195-3>
- [24] X. Wang and M. Yue, A  $q$ -analogue of the (A.2) supercongruence of Van Hamme for any prime  $p \equiv 3 \pmod{4}$ , *Int. J. Number Theory*, in press; <https://doi.org/10.1142/S1793042120500694>
- [25] H.S. Wilf, D. Zeilberger, An algorithmic proof theory for hypergeometric (ordinary and “ $q$ ”) multisum/integral identities, *Invent. Math.* 108 (1992), 575–633.

- [26] H.S. Wilf, D. Zeilberger, Rational function certification of multisum/integral/“ $q$ ” identities, *Bull. Amer. Math. Soc. (N.S.)*, 27 (1992), 148–153.
- [27] W. Zudilin, Ramanujan-type supercongruences, *J. Number Theory* 129 (2009), 1848–1857.
- [28] W. Zudilin, Congruences for  $q$ -binomial coefficients, *Ann. Combin.* 23 (2019), 1123–1135.