

DIOPHANTINE PROBLEMS RELATED TO THE OMEGA CONSTANT

WADIM ZUDILIN

To the memory of Jonathan Sondow (1943–2020)

The Omega constant

$$\Omega = 0.567143290409783872999968662210355549753815787186512508135131\dots$$

is the unique real solution of the transcendental equation $\Omega e^\Omega = 1$, which has several equivalent formulations including $-\log \Omega = \Omega$. It can be numerically computed via the iterates $x_{n+1} = (1 + x_n)/(1 + e^{x_n})$ starting, for example, from $x_0 = 1$. The constant shows up naturally in relation with Abel's polynomial interpolation [2, Section 3.5]: the function $f(x) = e^{\Omega x}$ satisfies $f^{(n)}(n) = 1$ for all $n = 0, 1, 2, \dots$, where $f^{(n)}(x)$ denotes the n th derivative of $f(x)$. The transcendence of Ω directly follows from the Lindemann–Weierstrass theorem (in fact, from its ‘Lindemann’ part!), however the quality of rational (and algebraic) approximations to Ω remains an open problem. I learned the following question from J. Sondow in 2008; it remains completely untouched because of lack of approaches.

Question 1. Is this true that the Omega constant is *not* a Liouville number? Give an estimate for the irrationality measure of Ω .

Applying the general lower bounds [3, Theorem 1] for $|e^\theta - \alpha| + |\theta - \beta|$, where $\theta \in \mathbb{C} \setminus \{0\}$ is fixed and α, β are varying algebraic numbers, with the choice $\theta = -\Omega$ and $\alpha = -\beta \in \mathbb{Q}$ one gets $|\Omega - p/q| > \exp(-C \log^2 |q|)$ for some absolute constant $C > 0$. Question 1 explicitly asks whether one can achieve the sharper bound $|\Omega - p/q| > \exp(-C \log |q|)$ for some $C > 0$.

The Omega constant is closely related to the Lambert W -function which is (a particular branch of) the functional inverse of ze^z ; in other words, $z = W(ze^z)$. The ‘standard’ branch of $W(x)$ is real-valued for real $x \geq -1/e$; for it we have $\Omega = W(1)$. The branch admits the series expansions

$$W(x)^k = kx^k \sum_{n=0}^{\infty} \frac{(n+k)^{n-1}}{n!} (-1)^n x^n, \quad |x| < \frac{1}{e},$$

for each $k = 1, 2, \dots$. Again, it follows from Lindemann's theorem that the values of $W(x)$ at non-zero rational (and algebraic) numbers are transcendental, and a natural question to ask is whether one can give some stronger estimates, than those implied by [3], for approximations of the values by rational (and algebraic) numbers.

Similar problems may be addressed for the values of $W(x)$ at non-zero algebraic points θ with $0 < |\theta|_p < p^{-1/(p-1)}$ in the p -adic settings. Mahler's result [1] from

1932 about the transcendence of $\exp(\theta)$ replaces Lindemann's theorem above. This strategy, however, gives one no way to assign a specific quantity from \mathbb{C}_p to $W(1)$, perhaps leaving the transcendental number $W(p) \in \mathbb{C}_p$ as an appropriate candidate for a p -adic Omega constant.

Possibly a more intrinsic p -adic version of the Lambert W -function comes from the formal series inverse $W_p(x)$ of $xE_p(x)$, where

$$E_p(x) = \exp\left(x + \frac{x^p}{p} + \frac{x^{p^2}}{p^2} + \cdots + \frac{x^{p^n}}{p^n} + \cdots\right) = \prod_{\substack{n=1 \\ (n,p)=1}}^{\infty} (1 - x^n)^{-\mu(n)/n} \in \mathbb{Z}_p[[x]]$$

stands for the Artin–Hasse exponential (and $\mu(n)$ denotes the Möbius function). Then $W_p(x) \in \mathbb{Z}_p[[x]]$ so that the values of the series are well defined inside the p -adic unit disk $|x|_p < 1$. Observe that

$$\prod_{n=1}^{\infty} (1 - x^n)^{-\mu(n)/n} = \exp(x)$$

corresponds to the product side of $E_p(x)$ with $p = 1$, and in this sense $W_1(x)$ is exactly the real-valued branch $W(x)$ above. This setup suggests $\Omega_p = W_p(p)$ as a legitimate p -adic Omega constant; alternatively, it can be defined as the limit of the iterates $x_{n+1} = (p + x_n)/(1 + E_p(x_n))$ with $x_0 = p$. In absence of Lindemann's theorem for the values of function $E_p(x)$ at algebraic points, already the *irrationality* of Ω_p looks rather challenging.

Acknowledgements. I thank Michel Waldschmidt for related comments and helpful feedback.

REFERENCES

- [1] K. MAHLER, Ein Beweis der Transcendenz der p -adischen Exponentialfunktion, *J. Reine Angew. Math.* **169** (1932), 61–66.
- [2] M. WALDSCHMIDT, Integer-valued functions, Hurwitz functions and related topics: a survey, *Preprint arXiv: 2002.01223 [math.NT]* (2020), 24 pp.
- [3] M. WALDSCHMIDT and YU. V. NESTERENKO, On the approximation of the values of exponential function and logarithm by algebraic numbers, in: Diophantine Approximations, Collection of papers dedicated to the memory of Prof. N. I. Feldman, *Mat. Zapiski* **2** (1996), 23–42 (Russian); English transl., *arXiv: math/0002047 [math.NT]* (2000), 17 pp.

DEPARTMENT OF MATHEMATICS, IMAPP, RADBOUD UNIVERSITY, PO Box 9010, 6500 GL NIJMEGEN, NETHERLANDS

E-mail address: w.zudilin@math.ru.nl