

ON SOME CONGRUENCES USING MULTIPLE HARMONIC SUMS OF LENGTH THREE AND FOUR

WALID KEHILA

ABSTRACT. In the present paper, we determine the sums $\sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}}$ and $\sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)} H_j^{(s_4)}}{j^{s_2}}$ modulo p and modulo p^2 in certain cases. This is done by using multiple harmonic sums of length three and four, as well as, many other results. In addition, We recover three congruences conjectured by Z.-W Sun and solved later by the author himself and R. Meštrović.

1. INTRODUCTION

Multiple Harmonic Sums (MHS) are defined by

$$H(s_1, \dots, s_k; n) := \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \frac{1}{j_1^{s_1} \dots j_k^{s_k}},$$

with the conventions $H(s_1, \dots, s_k; r) = 0$ for $r = 0, \dots, k-1$, and, $H(\emptyset; 0) = 1$. They satisfy the following recurrence relation [7]

$$H(s_1, \dots, s_k; n) = \sum_{k=1}^n \frac{1}{k^{s_k}} H(s_1, \dots, s_{k-1}; k-1).$$

In the case when $s_1 = \dots = s_k = s$, these sums are called the homogeneous multiple harmonic sums, and denoted

$$H(\underbrace{\{s\}^k}_{k \text{ times}}; n) := H(s, \dots, s; n) = \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq n} \frac{1}{(j_1 \dots j_k)^s}.$$

When $k = 1$, we find the sequence of generalized harmonic numbers, we may denote it

$$H_n^{(s)} = \sum_{k=1}^n \frac{1}{k^s},$$

note that the superscript is omitted in the case $s = 1$.

When $n = p-1$ we may simplify notations as follows

$$H(s_1, \dots, s_k) := \sum_{1 \leq j_1 < j_2 < \dots < j_k \leq p-1} \frac{1}{j_1^{s_1} \dots j_k^{s_k}} \quad \text{and} \quad H(s) := H_{p-1}^{(s)} = \sum_{k=1}^{p-1} \frac{1}{k^s}.$$

In [5] Sun has proposed some conjectures namely (Conjecture 1.1 and Conjecture 1.2); later, in [6] he proved (Conjecture 1.2). Meštrović [4], on the other hand, established the second part of Conjecture 1.1 using congruences of (MHS) of length three that can be found in [8].

In the present paper, we may unify the proof of these conjectures, as well as, proving many other congruences.

2010 *Mathematics Subject Classification.* 11A07, 11B68, 11B50, 11B83.

Key words and phrases. Harmonic numbers, Generalized harmonic numbers, Multiple harmonic sums.

Now, we give some results which will be using in the present paper.

Theorem 1.1. [8] *Let s and l be two positive integers. Let p be an odd prime such that $p \geq l + 2$ and $p - 1$ divides non of ls and $ks + 1$ for $k = 1, \dots, l$. Then*

$$H(\{s\}^l) \equiv \begin{cases} 0 & (\text{mod } p^2) \text{ for } ls - 1 \text{ even} \\ 0 & (\text{mod } p) \text{ for } ls - 1 \text{ odd.} \end{cases}$$

In particular, when $p \geq ls + 3$, the above is always true.

Theorem 1.2. [8] *Let s and k be two non-negative integers. Suppose also that $p \geq sk + 3$, then*

$$H(\{s\}^k) \equiv \begin{cases} (-1)^k \frac{s(sk+1)p^2}{2(sk+2)} B_{p-sk-2} & (\text{mod } p^3) \text{ for } ks \text{ odd} \\ (-1)^{k-1} \frac{sp}{sk+1} B_{p-sk-1} & (\text{mod } p^2) \text{ for } ks \text{ even,} \end{cases}$$

where $(B_n)_{n \in \mathbb{N}}$ is the sequence of Bernoulli numbers.

Theorem 1.3. [8] *Let s_1, s_2 be two positive integers and $p \geq 3$. Let $s_1 \equiv m, s_2 \equiv n \pmod{p-1}$ where $0 \leq m, n \leq p-2$. If $m, n \geq 1$ then*

$$H(s_1, s_2) \equiv \begin{cases} \frac{(-1)^n}{m+n} \binom{m+n}{m} B_{p-m-n} & (\text{mod } p) \text{ for } p \geq m+n \\ 0 & (\text{mod } p) \text{ for } p < m+n. \end{cases}$$

Furthermore, when $s_1 + s_2$ is even and $p > s_1 + s_2 + 1$

$$\begin{aligned} H(s_1, s_2) &\equiv p \left((-1)^{s_1} s_2 \binom{s_1 + s_2 + 1}{s_1} - (-1)^{s_1} s_1 \binom{s_1 + s_2 + 1}{s_2} - s_1 - s_2 \right) \\ &\quad \times \frac{B_{p-s_1-s_2-1}}{2(s_1 + s_2 + 1)} \pmod{p^2}. \end{aligned}$$

Theorem 1.4. [8][1] *Suppose that $w := s_1 + s_2 + s_3$ is odd, then for primes $p > w$ we have*

$$H(s_1, s_2, s_3) \equiv \left((-1)^{s_1} \binom{w}{s_1} - (-1)^{s_3} \binom{w}{s_3} \right) \frac{B_{p-w}}{2w} \pmod{p}.$$

In particular, when $s_1 = s_3$, and s_2 is odd, we have

$$H(s_1, s_2, s_1) \equiv 0 \pmod{p}.$$

2. CONGRUENCES USING MHS OF LENGTH THREE AND TWO

We start by establishing the following theorem which is motivated by a paper of Meštrović [4].

Theorem 2.1. *For all positive integers s_1, s_2, s_3 , we have the following*

$$\sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} = -H(s_1, s_2, s_3) + H(s_3, s_1 + s_2) + H(s_1 + s_2 + s_3) + H(s_3)H(s_1, s_2). \quad (2.1)$$

Equivalently, we have

$$\sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} = H(s_1, s_3, s_2) + H(s_3, s_1, s_2) + H(s_3, s_1 + s_2) + H(s_1 + s_3, s_2) + H(s_1, s_2 + s_3) + H(s_1 + s_2 + s_3). \quad (2.2)$$

Proof. For $j = 1, \dots, p-1$, we have

$$\frac{1}{(j+1)^{s_3}} + \dots + \frac{1}{(p-1)^{s_3}} = -H_j^{(s_3)} + H(s_3).$$

We obtain the following

$$\begin{aligned} H(s_1, s_2, s_3) &= \sum_{1 \leq i < j < k \leq p-1} \frac{1}{i^{s_1} j^{s_2} k^{s_3}} = \sum_{j=2}^{p-1} \frac{1}{j^{s_2}} \sum_{i=1}^{j-1} \frac{1}{i^{s_1}} \sum_{k=j+1}^{p-1} \frac{1}{k^{s_3}} \\ &= \sum_{j=2}^{p-1} \frac{1}{j^{s_2}} \left(1 + \frac{1}{2^{s_1}} + \dots + \frac{1}{(j-1)^{s_1}}\right) \left(\frac{1}{(j+1)^{s_3}} + \dots + \frac{1}{(p-1)^{s_3}}\right) \\ &= \sum_{j=2}^{p-1} \frac{1}{j^{s_2}} \left(1 + \frac{1}{2^{s_1}} + \dots + \frac{1}{(j-1)^{s_1}}\right) (-H_j^{(s_3)} + H(s_3)) \\ &= -\sum_{j=1}^{p-1} \frac{1}{j^{s_2}} \left(H_j^{(s_1)} - \frac{1}{j^{s_1}}\right) H_j^{(s_3)} + H(s_3) \sum_{j=1}^{p-1} \frac{1}{j^{s_2}} H_{j-1}^{(s_1)} \\ &= -\sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} + \sum_{j=1}^{p-1} \frac{1}{j^{s_1+s_2}} \left(H_{j-1}^{(s_3)} + \frac{1}{j^{s_3}}\right) + H(s_3) \sum_{j=1}^{p-1} \frac{1}{j^{s_2}} H_{j-1}^{(s_1)} \\ &= -\sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} + H(s_3, s_1 + s_2) + H(s_1 + s_2 + s_3) + H(s_3) H(s_1, s_2). \end{aligned}$$

Therefore

$$\sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} = -H(s_1, s_2, s_3) + H(s_3, s_1 + s_2) + H(s_1 + s_2 + s_3) + H(s_3) H(s_1, s_2).$$

Equation (2.2) follows immediately by the shuffle product relation

$$H(s_3) H(s_1, s_2) = H(s_3, s_1, s_2) + H(s_1, s_3, s_2) + H(s_1, s_2, s_3) + H(s_3 + s_1, s_2) + H(s_1, s_3 + s_2). \quad \square$$

Remark 1. One can easily observe that the previous proof does not depend on $p-1$, so for all positive integers n , we have

$$\sum_{j=1}^n \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} = -H(s_1, s_2, s_3; n) + H(s_3, s_1 + s_2; n) + H(s_1 + s_2 + s_3; n) + H(s_3; n) H(s_1, s_2; n).$$

and

$$\begin{aligned} \sum_{j=1}^n \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} &= H(s_1, s_3, s_2; n) + H(s_3, s_1, s_2; n) + H(s_3, s_1 + s_2; n) \\ &\quad + H(s_1 + s_3, s_2; n) + H(s_1, s_2 + s_3; n) + H_n^{(s_1+s_2+s_3)}. \end{aligned}$$

2.1. Congruences mod p . From Theorem 2.1 one can see that determining $\sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}}$ depends on determining MHS of length three and two, Using Theorems 1.4 and 1.3 we obtain the following result.

Theorem 2.2. *Suppose that w is odd then*

$$\sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} \equiv \left[\frac{(-1)^{s_1+1}}{2w} \binom{w}{s_1} + \frac{(-1)^{s_3} + 2(-1)^{s_1+s_2}}{2w} \binom{w}{s_3} \right] B_{p-w} \pmod{p}. \quad (2.3)$$

Proof. From the first part of Theorem 2.1, and in light of Theorems 1.4 and 1.3, we have modulo p

$$\begin{aligned} \sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} &\equiv - \left((-1)^{s_1} \binom{w}{s_1} - (-1)^{s_3} \binom{w}{s_3} \frac{B_{p-w}}{2w} \right) + \frac{(-1)^{s_1+s_2}}{w} \binom{w}{s_3} B_{p-w} \\ &\equiv \left[\frac{(-1)^{s_1+1}}{2w} \binom{w}{s_1} + \frac{(-1)^{s_3} + 2(-1)^{s_1+s_2}}{2w} \binom{w}{s_3} \right] B_{p-w}. \end{aligned}$$

□

Corollary 2.2.1. *Let $s \geq 1$ and r is odd, then for $p > 2s + r$, we have*

$$\sum_{j=1}^{p-1} \frac{(H_j^{(s)})^2}{j^r} \equiv (-1)^{s+r} \binom{2s+r}{s} \frac{B_{p-2s-r}}{2s+r} \pmod{p}. \quad (2.4)$$

For all $s \geq 1$ and any prime $p \geq 3$ such that $p-1$ does not divide $3s$, in particular when $p \geq 3s+3$, we have

$$\sum_{j=1}^{p-1} \frac{(H_j^{(s)})^2}{j^s} \equiv \binom{3s}{s} \frac{B_{p-3s}}{3s} \pmod{p}. \quad (2.5)$$

Moreover, when s is even, then

$$\sum_{j=1}^{p-1} \frac{(H_j^{(s)})^2}{j^s} \equiv 0 \pmod{p}. \quad (2.6)$$

Proof. For the first part, we apply Theorems 1.1 and 1.4.

For the second part, we apply Theorems 1.1 and 1.3; hence the last part yields immediately since Bernoulli numbers vanishes for odd values. This would solve the first part of Sun Conjecture see ([5] Conjecture 1.2) and ([6] Theorem 1.2). □

2.2. Some special cases. In this section, we shall prove some congruences modulo p that are not covered by Theorem 2.2.

Lemma 2.1. [2] Theorem 7.2, [8] page 94 and Proposition 3.8

For $p \geq 11$ we have

$$\frac{1}{3} H(2, 3, 1) \equiv -\frac{1}{2} H(3, 2, 1) \equiv -H(3, 1, 2) \equiv -\frac{1}{2} H(1, 4, 1) \equiv H(4, 1, 1) \equiv -\frac{1}{6} B_{p-3}^2 \pmod{p}. \quad (2.7)$$

For $p \geq 7$ we have

$$H(1, 2, 2) \equiv -\frac{3}{2} B_{p-5} \pmod{p}. \quad (2.8)$$

For $p \geq 17$ we have

$$H(5, 3, 4) \equiv H(4, 3, 5) \equiv 0 \pmod{p}. \quad (2.9)$$

Theorem 2.3. For $p \geq 11$ we have

$$2 \sum_{j=1}^{p-1} \frac{H_j^{(2)} H_j}{j^3} \equiv -3 \sum_{j=1}^{p-1} \frac{H_j^{(3)} H_j}{j^2} \equiv -6 \sum_{j=1}^{p-1} \frac{H_j^{(3)} H_j^{(2)}}{j} \equiv -3 \sum_{j=1}^{p-1} \frac{(H_j)^2}{j^4} \equiv 6 \sum_{j=1}^{p-1} \frac{H_j^{(4)} H_j}{j} \equiv B_{p-3}^2 \pmod{p}, \quad (2.10)$$

$$\sum_{j=1}^{p-1} \frac{H_j H_j^{(2)}}{j^2} \equiv -\frac{1}{2} B_{p-5} \pmod{p}. \quad (2.11)$$

For $p \geq 17$ we have

$$\sum_{j=1}^{p-1} \frac{H_j^{(5)} H_j^{(4)}}{j^3} \equiv 0 \pmod{p}. \quad (2.12)$$

Proof. We shall prove only the last one, since the others are proved in a similar manner.

From Theorems 2.1, 1.3 and the previous Lemma we find

$$\sum_{j=1}^{p-1} \frac{H_j^{(5)} H_j^{(4)}}{j^3} \equiv -H(5, 3, 4) + H(4, 8) \equiv 0 + \frac{165}{4} B_{p-12} = 0 \pmod{p}.$$

□

2.3. Congruences mod p^2 . In this section, we prove some congruences modulo p^2 in certain cases only, since determining (MHS) of length three seems to be a much more involved problem.

Lemma 2.2. [8] *Proposition 3.7*

For $p \geq 7$ we have

$$\frac{10}{9} H(1, 2, 1) \equiv \frac{5}{3} H(2, 1, 1) \equiv \frac{10}{11} H(1, 1, 2) \equiv p B_{p-5} \pmod{p^2}, \quad (2.13)$$

$$H(1, 3, 1) \equiv 0 \pmod{p^2}, \quad (2.14)$$

$$H(4, 1) \equiv -B_{p-5} \pmod{p^2}. \quad (2.15)$$

Corollary 2.3.1. For $p \geq 7$ we have

$$\sum_{j=1}^{p-1} \frac{H_j^2}{j^2} \equiv H(4) \equiv \frac{4}{5} p B_{p-5} \pmod{p^2}, \quad (2.16)$$

$$\sum_{j=1}^{p-1} \frac{H_j^{(2)} H_j}{j} \equiv -\frac{7}{10} p B_{p-5} \pmod{p^2}, \quad (2.17)$$

$$\sum_{j=1}^{p-1} \frac{H_j^2}{j^3} \equiv B_{p-5} \pmod{p^2}. \quad (2.18)$$

Proof. From Theorem 1.3 and the previous Lemma we obtain

$$H(1, 2, 1) \equiv \frac{9}{10} p B_{p-5} \equiv H(1, 3) \pmod{p^2}.$$

Substituting in (2.1) and using Theorem 1.2 we find

$$\sum_{j=1}^{p-1} \frac{H_j^2}{j^2} \equiv H(4) \equiv \frac{4}{5} p B_{p-5} \pmod{p^2}.$$

This establishes the second part of Sun Conjecture see ([5] conjecture 1.1) proved by Meštrović see [4].

From Theorems 1.3, 1.2, and the previous Lemma we get

$$\sum_{j=1}^{p-1} \frac{H_j^{(2)} H_j}{j} \equiv -\frac{3}{5} p B_{p-5} - \frac{9}{10} p B_{p-5} + \frac{4}{5} p B_{p-5} \equiv -\frac{7}{10} p B_{p-5} \pmod{p^2}.$$

From Equation (2.1) and the previous Lemma we obtain

$$\sum_{j=1}^{p-1} \frac{H_j^2}{j^3} \equiv H(1, 4) \pmod{p^2}.$$

Since

$$H(4, 1) \equiv -B_{p-5} \pmod{p^2},$$

we have from Theorem 3.2 in [8]

$$H(4, 1) + H(1, 4) \equiv -B_{p-5} + H(1, 4) \equiv -\frac{5}{6}pB_{p-6} = 0 \pmod{p^2}.$$

□

Corollary 2.3.2. *Suppose s is even, then for $p > 3s + 1$ we have*

$$\sum_{j=1}^{p-1} \frac{(H_j^{(s)})^2}{j^s} \equiv \left[\binom{3s+1}{s-1} + \frac{s}{2} \right] p \frac{B_{p-3s-1}}{3s+1} \pmod{p^2}. \quad (2.19)$$

Proof. Using the identity

$$\binom{n}{k+1} = \frac{n-k}{k+1} \binom{n}{k},$$

and from the first part of Theorem 2.1, we find modulo p^2

$$\begin{aligned} \sum_{j=1}^{p-1} \frac{(H_j^{(s)})^2}{j^s} &\equiv -H(s, s, s) + H(s, 2s) + H(3s) \\ &\equiv \left[-s + \frac{2s \binom{3s+1}{s} - s \binom{3s+1}{2s} - 3s}{2} + 3s \right] p \frac{B_{p-3s-1}}{3s+1} \\ &= \left[2 + \frac{\binom{3s+1}{s} - \frac{s}{s+1} \binom{3s+1}{s} - 3}{2} \right] sp \frac{B_{p-3s-1}}{3s+1} \\ &= \left[1 + \frac{\binom{3s+1}{s}}{s+1} \right] \frac{sp}{2} \frac{B_{p-3s-1}}{3s+1} \\ &= \left[\binom{3s+1}{s-1} + \frac{s}{2} \right] p \frac{B_{p-3s-1}}{3s+1}. \end{aligned}$$

Congruences (2.5) and (2.19) establishes Conjecture 1.2 (see [5] and [6] Theorem 1.2) with a slightly different approach. □

3. SOME SPECIAL CONGRUENCES USING MHS OF LENGTH FOUR

In this section, we prove some congruences using some results on (MHS) of length four.

Theorem 3.1. *The following equality holds true*

$$\begin{aligned} -\sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)} H_j^{(s_4)}}{j^{s_2}} &= H(s_4) \sum_{j=1}^{p-1} \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} + H(s_1, s_3, s_2, s_4) + H(s_3, s_1, s_2, s_4) \\ &\quad + H(s_3, s_1 + s_2, s_4) + H(s_1 + s_3, s_2, s_4) + H(s_1, s_2 + s_3, s_4) + H(s_1 + s_2 + s_3, s_4). \quad (3.1) \end{aligned}$$

Proof. From Remark 1 and Equation (2.2) we find

$$\begin{aligned}
 \sum_{l=1}^{p-1} \frac{1}{l^{s_4}} \sum_{j=1}^{l-1} \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} &= \sum_{l=1}^{p-1} \frac{1}{l^{s_4}} \left[H(s_1, s_3, s_2; l-1) + H(s_3, s_1, s_2; l-1) + H(s_3, s_1 + s_2; l-1) \right. \\
 &\quad \left. + H(s_1 + s_3, s_2; l-1) + H(s_1, s_2 + s_3; l-1) + H(s_1 + s_2 + s_3; l-1) \right] \\
 \sum_{j=1}^{p-1} \left(\sum_{l=j+1}^{p-1} \frac{1}{l^{s_4}} \right) \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} &= H(s_1, s_3, s_2, s_4) + H(s_3, s_1, s_2, s_4) + H(s_3, s_1 + s_2, s_4) + H(s_1 + s_3, s_2, s_4) \\
 &\quad + H(s_1, s_2 + s_3, s_4) + H(s_1 + s_2 + s_3, s_4) \\
 \sum_{j=1}^{p-1} (H(s_4) - H_j^{(s_4)}) \frac{H_j^{(s_1)} H_j^{(s_3)}}{j^{s_2}} &= H(s_1, s_3, s_2, s_4) + H(s_3, s_1, s_2, s_4) + H(s_3, s_1 + s_2, s_4) + H(s_1 + s_3, s_2, s_4) \\
 &\quad + H(s_1, s_2 + s_3, s_4) + H(s_1 + s_2 + s_3, s_4).
 \end{aligned}$$

Hence, 3.1 yields immediately. □

Corollary 3.1.1. *For $p \geq 7$ we have*

$$\sum_{j=1}^{p-1} \frac{H_j^3}{j} \equiv \frac{3}{2} p B_{p-5} \pmod{p^2}. \quad (3.2)$$

Proof. From the previous Theorem, we have modulo p^2

$$\begin{aligned}
 - \sum_{j=1}^{p-1} \frac{H_j^3}{j} &\equiv 2H(\{1\}^4) + 2H(1, 2, 1) + H(2, 1, 1) + H(3, 1) \\
 - \sum_{j=1}^{p-1} \frac{H_j^3}{j} &\equiv -\frac{2}{5} p B_{p-5} - \frac{9}{5} p B_{p-5} + \frac{3}{5} p B_{p-5} + \frac{1}{10} p B_{p-5} = -\frac{3}{2} p B_{p-5}.
 \end{aligned}$$

Note that another proof can be found in [4]. □

Lemma 3.1. [1] *Let a, b be non-negative integers and a prime $p > 2a + 2b + 3$. Then*

$$H(\{2\}^a, 3, \{2\}^b) \equiv \frac{(-1)^{a+b}(a-b)}{(a+1)(b+1)} \binom{2a+2b+2}{2a+1} B_{p-2a-2b-3} \pmod{p}, \quad (3.3)$$

And for primes such that $p > 2a + 2b + 1$, we have

$$H(\{2\}^a, 1, \{2\}^b) \equiv \frac{4(-1)^{a+b}(a-b)(1-4^{-a-b})}{(2a+1)(2b+1)} \binom{2a+2b}{2a} B_{p-2a-2b-1} \pmod{p}. \quad (3.4)$$

Corollary 3.1.2. *For $p \geq 11$ we have*

$$-\frac{1}{13} \sum_{j=1}^{p-1} \frac{(H_j^{(2)})^2 H_j^{(3)}}{j^2} \equiv \frac{3}{83} \sum_{j=1}^{p-1} \frac{(H_j^{(2)})^3}{j^3} \equiv B_{p-9} \pmod{p}, \quad (3.5)$$

$$-\frac{8}{21} \sum_{j=1}^{p-1} \frac{(H_j^{(2)})^2 H_j}{j^2} \equiv \frac{1}{3} \sum_{j=1}^{p-1} \frac{(H_j^{(2)})^3}{j} \equiv B_{p-7} \pmod{p}. \quad (3.6)$$

Proof. For the first congruence, we apply the previous Lemma, Theorems 3.1, 1.3 and 1.4 and obtain

$$\begin{aligned} \sum_{j=1}^{p-1} \frac{(H_j^{(2)})^2 H_j^{(3)}}{j^2} &= -2H(\{2\}^3, 3) - 2H(2, 4, 3) - H(4, 2, 3) - H(6, 3) \\ &\equiv \left[+12 - \frac{40}{3} - \frac{35}{3} - 0 \right] B_{p-9} = -13B_{p-9}, \end{aligned}$$

also

$$\begin{aligned} \sum_{j=1}^{p-1} \frac{(H_j^{(2)})^3}{j^3} &= -2H(\{2\}^2, 3, 2) - 2H(2, 5, 2) - H(5, 2, 2) - H(7, 2) \\ &\equiv \left[+\frac{56}{3} - 0 + 9 - 0 \right] B_{p-9} = \frac{83}{3} B_{p-9}. \end{aligned}$$

The second part of the corollary is proved in a quite similar manner. □

REFERENCES

- [1] KH. Hessami Pilehrood, T. Hessami Pilehrood, and R. Tauraso, *New properties of multiple harmonic sums modulo p and p-analogues of Leshchiner's series*, Trans. Amer. Math. Soc, Volume 366, Number 6, June 2014, Pages 3131–3159.
- [2] M.E. Hoffman, *Quasi-symmetric functions and mod p multiple harmonic sums*, [math.NT/0401319](https://arxiv.org/abs/math/0401319).
- [3] R. Meštrović, *An Extension of a Congruence by Tauraso*, ISRN Combinatorics, Volume 2013, Article ID 363724, 7 pages.
- [4] R. Meštrović, *Proof of a congruence for harmonic numbers conjectured by Z.-W.Sun*, Int. J. Number Theory Vol.8, No.4 (2012), 1081–1085.
- [5] Z.-W. Sun, *Arithmetic theory of harmonic numbers*. Proc. Am. Math. Soc. 140, 415–428 (2012).
- [6] Z.-W. Sun, L.-L. Zhao, *Arithmetic theory of harmonic numbers (II)*. Colloq. Math. 130, 67–78 (2013).
- [7] R. Tauraso, *New harmonic number identities with applications*, Sémin. Lothar. Combin. 63 (2010) Article B63g.
- [8] J. Zhao, *Wolstenholme type theorem for multiple harmonic sum*, Int. J. of Number Theory 4 (2008), 73–106.

* UNIVERSITY OF SCIENCE AND TECHNOLOGY
 HOUARI BOUMEDIENNE
 USTHB
 FACULTY OF MATHEMATICS, P.B. 32 EL ALIA, 16111
 ALGIERS, ALGERIA
E-mail address: wkehilausthb@gmail.com; wkehila@usthb.dz