

# FIRST-PASSAGE TIMES FOR RANDOM WALKS IN THE TRIANGULAR ARRAY SETTING

DENIS DENISOV, ALEXANDER SAKHANENKO, AND VITALI WACHTEL

ABSTRACT. In this paper we continue our study of exit times for random walks with independent but not necessarily identically distributed increments. Our paper “First-passage times for random walks with non-identically distributed increments” (2018) was devoted to the case when the random walk is constructed by a fixed sequence of independent random variables which satisfies the classical Lindeberg condition. Now we consider a more general situation when we have a triangular array of independent random variables. Our main assumption is that the entries of every row are uniformly bounded by a deterministic sequence, which tends to zero as the number of the row increases.

## 1. INTRODUCTION AND THE MAIN RESULT.

1.1. **Introduction.** Suppose that for each  $n = 1, 2, \dots$  we are given independent random variables  $X_{1,n}, \dots, X_{n,n}$  such that

$$\mathbf{E}X_{i,n} = 0 \quad \text{for all } i \leq n \quad \text{and} \quad \sum_{i=1}^n \mathbf{E}X_{i,n}^2 = 1. \quad (1)$$

For each  $n$  we consider a random walk

$$S_{k,n} := X_{1,n} + \dots + X_{k,n}, \quad k = 1, 2, \dots, n. \quad (2)$$

Let  $\{g_{k,n}\}_{k=1}^n$  be deterministic real numbers, and let

$$T_n := \inf\{k \geq 1 : S_{k,n} \leq g_{k,n}\} \quad (3)$$

be the first crossing over the moving boundary  $\{g_{k,n}\}$  by the random walk  $\{S_{k,n}\}$ . The main purpose of the present paper is to study the asymptotic behaviour, as  $n \rightarrow \infty$ , of the probability

$$\mathbf{P}(T_n > n) = \mathbf{P}\left(\min_{1 \leq k \leq n} (S_{k,n} - g_{k,n}) > 0\right). \quad (4)$$

We shall always assume that the boundary  $\{g_{k,n}\}$  is of a small magnitude, that is,

$$g_n^* := \max_{1 \leq k \leq n} |g_{k,n}| \rightarrow 0. \quad (5)$$

Here and in what follows, all unspecified limits are taken with respect to  $n \rightarrow \infty$ .

---

2020 *Mathematics Subject Classification.* Primary 60G50; Secondary 60G40, 60F17.

*Key words and phrases.* Random walk, triangular array, first-passage time, central limit theorem, moving boundary, transition phenomena.

A.S. and V.W. were supported by RFBR and DFG according to the research project № 20-51-12007.

Furthermore, to avoid trivialities, we shall assume that

$$\mathbf{P}(T_n > n) > 0 \quad \text{for all } n \geq 1. \quad (6)$$

An important particular case of the triangular array scheme is given by the following construction. Let  $X_1, X_2, \dots$  be independent random variables with finite variances such that

$$\mathbf{E}X_i = 0 \quad \text{for all } i \geq 1 \quad \text{and} \quad B_n^2 := \sum_{i=1}^n \mathbf{E}X_i^2 \rightarrow \infty. \quad (7)$$

For a real deterministic sequence  $\{g_1, g_2, \dots\}$  set

$$T := \inf\{k \geq 1 : S_k \leq g_k\}, \quad \text{where } S_k := X_1 + \dots + X_k. \quad (8)$$

Stopping time  $T$  is the first crossing over the moving boundary  $\{g_k\}$  by the random walk  $\{S_k\}$ . Clearly, (7) – (8) is a particular case of (1) – (3). Indeed to obtain (1) – (3) it is sufficient to set

$$X_{k,n} = \frac{X_k}{B_n}, \quad S_{k,n} = \frac{S_k}{B_n}, \quad g_{k,n} = \frac{g_k}{B_n}. \quad (9)$$

However, the triangular array scheme is much more general than (7) – (9).

If the classical Lindeberg condition holds for the sequence  $\{X_k\}$  and  $g_n = o(B_n)$  then, according to Theorem 1 in [2],

$$\mathbf{P}(T > n) \sim \sqrt{\frac{2}{\pi}} \frac{U(B_n^2)}{B_n}, \quad (10)$$

where  $U$  is a positive slowly varying function with the values

$$U(B_n^2) = \mathbf{E}[S_n - g_n; T > n], \quad n \geq 1.$$

The constant  $\sqrt{\frac{2}{\pi}}$  in front of the asymptotics has been inherited from the tail asymptotics of exit time of standard Brownian motion. Indeed, let  $W(t)$  be the standard Brownian motion and set

$$\tau_x^{bm} := \inf\{t > 0 : x + W(t) \leq 0\}, \quad x > 0.$$

Then,

$$\mathbf{P}(\tau_x^{bm} > t) = \mathbf{P}(|W(t)| \leq x) = \mathbf{P}\left(|W(1)| \leq \frac{x}{\sqrt{t}}\right) \sim \sqrt{\frac{2}{\pi}} \frac{x}{\sqrt{t}}, \quad \text{as } \frac{x}{\sqrt{t}} \rightarrow 0.$$

The continuity of paths of  $W(t)$  implies that  $x + W(\tau_x^{bm}) = 0$ . Combining this with the optional stopping theorem, we obtain

$$\begin{aligned} x &= \mathbf{E}[x + W(\tau_x^{bm} \wedge t)] = \mathbf{E}[x + W(t); \tau_x^{bm} > t] + \mathbf{E}[x + W(\tau_x^{bm}); \tau_x^{bm} \leq t] \\ &= \mathbf{E}[x + W(t); \tau_x^{bm} > t]. \end{aligned}$$

Therefore, for any fixed  $x > 0$ ,

$$\mathbf{P}(\tau_x^{bm} > t) \sim \sqrt{\frac{2}{\pi}} \frac{x}{\sqrt{t}} = \sqrt{\frac{2}{\pi}} \frac{\mathbf{E}[x + W(t); \tau_x^{bm} > t]}{\sqrt{t}}, \quad \text{as } t \rightarrow \infty.$$

Thus, the right hand sides here and in (10) are of the same type.

**1.2. Main result.** The purpose of the present note is to generalise the asymptotic relation (10) to the triangular array setting. More precisely, we are going to show that the following relation holds

$$\mathbf{P}(T_n > n) \sim \sqrt{\frac{2}{\pi}} E_n, \quad (11)$$

where

$$E_n := \mathbf{E}[S_{n,n} - g_{n,n}; T_n > n] = \mathbf{E}[-S_{T_n,n}; T_n \leq n] - g_{n,n} \mathbf{P}(T_n > n). \quad (12)$$

Unexpectedly for the authors, in contrast to the described above case of a single sequence, the Lindeberg condition is not sufficient for the validity of (11), see Example 6. Thus, one has to find a more restrictive condition for (11) to hold. In this paper we show that (11) holds under the following assumption: there exists a sequence  $r_n$  such that

$$\max_{1 \leq i \leq n} |X_{i,n}| \leq r_n \rightarrow 0. \quad (13)$$

It is clear that under this assumption the triangular array satisfies the Lindeberg condition and, hence, the Central Limit Theorem holds.

At first glance, (13) might look too restrictive. However we shall construct a triangular array, see Example 7, in which the assumption (13) becomes necessary for (11) to hold. Now we state our main result.

**Theorem 1.** *Assume that (5) and (13) are valid. Then there exists an absolute constant  $C_1$  such that*

$$\mathbf{P}(T_n > n) \geq \sqrt{\frac{2}{\pi}} E_n (1 - C_1 (r_n + g_n^*)^{2/3}). \quad (14)$$

*On the other hand, there exists an absolute constant  $C_2$  such that*

$$\mathbf{P}(T_n > n) \leq \sqrt{\frac{2}{\pi}} E_n (1 + C_2 (r_n + g_n^*)^{2/3}), \quad \text{if } r_n + g_n^* \leq 1/24. \quad (15)$$

*In addition, for  $m \leq n$ ,*

$$\mathbf{P}(T_n > m) \leq \frac{4E_n}{B_m^{(n)}} \quad (16)$$

*provided that*

$$B_m^{(n)} := \left( \sum_{k=1}^m \mathbf{E} X_{k,n}^2 \right)^{1/2} \geq 24(r_n + g_n^*).$$

**Corollary 2.** *Under conditions (5), (6) and (13) relation (11) takes place.*

Estimates (14) and (15) can be seen as an improved version of (11), with a rate of convergence. Moreover, the fact, that the dependence on  $r_n$  and  $g_n$  is expressed in a quite explicit way, is very important for our work [3] in progress, where we analyse unbounded random variables. In this paper we consider first-passage times of walks  $S_n = X_1 + X_2 + \dots + X_n$  for which the central limit theorem is valid but the Lindeberg condition may fail. We use Theorem 1 to analyse the behaviour of triangular arrays obtained from  $\{X_n\}$  by truncation.

**1.3. Triangular arrays of weighted random variables.** Theorem 1 and Corollary 2 can be used in studying first-passage times of weighted sums of independent random variables.

Suppose that we are given independent random variables  $X_1, X_2, \dots$  such that

$$\mathbf{E}X_i = 0 \quad \text{and} \quad \mathbf{P}(|X_i| \leq M_i) = 1 \quad \text{for all } i \geq 1, \quad (17)$$

where  $M_1, M_2, \dots$  are deterministic. For each  $n$  we consider a random walk

$$U_{k,n} := u_{1,n}X_1 + \dots + u_{k,n}X_k, \quad k = 1, 2, \dots, n, \quad (18)$$

and let

$$\tau_n := \inf\{k \geq 1 : U_{k,n} \leq G_{k,n}\} \quad (19)$$

be the first crossing over the moving boundary  $\{G_{k,n}\}$  by the random walk  $\{U_{k,n}\}$ . The main purpose of the present example is to study the asymptotic behaviour, as  $n \rightarrow \infty$ , of the probability

$$\mathbf{P}(\tau_n > n) = \mathbf{P}\left(\min_{1 \leq k \leq n} (U_{k,n} - G_{k,n}) > 0\right). \quad (20)$$

We suppose that  $\{u_{k,n}, G_{k,n}\}_{k=1}^n$  are deterministic real numbers such that

$$M := \sup_{k,n \geq 1} (|u_{k,n}|M_k + |G_{k,n}|) < \infty \quad (21)$$

and

$$\sigma_n^2 := \sum_{k=1}^n u_{k,n}^2 \mathbf{E}X_k^2 \rightarrow \infty. \quad (22)$$

Moreover, we assume that

$$u_{k,n} \rightarrow u_k \quad \text{and} \quad G_{n,k} \rightarrow g_k \quad \text{for every } k \geq 1. \quad (23)$$

**Corollary 3.** *Assume that the distribution functions of all  $X_k$  are continuous. Then, under assumptions (17), (21), (22) and (23),*

$$\sigma_n \mathbf{P}(\tau_n > n) \rightarrow \sqrt{\frac{2}{\pi}} \mathbf{E}[-U_\tau] \in [0, \infty), \quad (24)$$

where

$$U_k := u_1X_1 + \dots + u_kX_k \quad \text{and} \quad \tau := \inf\{k \geq 1 : U_k \leq g_k\}. \quad (25)$$

It follows from condition (23) that random walks  $\{U_{k,n}\}$  introduced in (18) may be considered as perturbations of the walk  $\{U_k\}$  defined in (25). Thus, we see from (24) that the influence of perturbations on the behavior of the probability  $\mathbf{P}(\tau_n > n)$  is concentrated in the  $\sigma_n$ .

**Example 4.** As an example we consider the following method of summation, which has been suggested by Gaposkin [4]. Let  $f : [0, 1] \mapsto \mathbb{R}^+$  be a non-degenerate continuous function. For random variables  $\{X_k\}$  define

$$U_k(n, f) := \sum_{j=1}^k f\left(\frac{j}{n}\right) X_j, \quad j = 1, 2, \dots, n.$$

This sequence can be seen as a stochastic integral of  $f$  with respect to the random walk  $S_k = X_1 + X_2 + \dots + X_k$  normalized by  $n$ .

We assume that the random variables  $\{X_k\}$  are independent and identically distributed. Furthermore, we assume that  $X_1$  satisfies (17) and that its distribution function is continuous. In this case

$$\sigma_n^2(f) := \frac{1}{n} \mathbf{E} X_1^2 \sum_{j=1}^n f^2\left(\frac{j}{n}\right) \rightarrow \sigma^2(f) := \mathbf{E} X_1^2 \int_0^1 f(t) dt > 0.$$

From Corollary 3 with  $u_{k,n} := f\left(\frac{j}{n}\right) \rightarrow f(0) =: u_k$ ,  $G_{k,n} \equiv 0$  and  $\sigma_n := \sqrt{n} \sigma_n(f)$  we immediately obtain

$$\sqrt{n} \mathbf{P}\left(\min_{k \leq n} U_k(n, f) > 0\right) \rightarrow \sqrt{\frac{2}{\pi}} \frac{f(0)}{\sigma(f)} \mathbf{P}[-S_\tau] \in [0, \infty), \quad (26)$$

where

$$S_k := X_1 + \cdots + X_k \quad \text{and} \quad \tau := \inf\{k \geq 1 : S_k \leq 0\}. \quad (27)$$

◇

Clearly, (26) gives one exact asymptotics only when  $f(0) > 0$ . The case  $f(0) = 0$  seems to be much more delicate. If  $f(0) = 0$  then one needs an information on the behaviour of  $f$  near zero. If, for example,  $f(t) = t^\alpha$  with some  $\alpha > 0$  then, according to Example 12 in [2],

$$\mathbf{P}\left(\min_{k \leq n} U_k(n, f) > 0\right) = \mathbf{P}\left(\min_{k \leq n} \sum_{j=1}^k j^\alpha X_j > 0\right) \sim \frac{Const}{n^{\alpha+1/2}}.$$

Now we give an example of application of our results to study of transition phenomena.

**Example 5.** Consider an autoregressive sequence

$$U_n(\gamma) = \gamma U_{n-1}(\gamma) + X_n, \quad n \geq 0, \quad n = 1, 2, \dots, \quad \text{where} \quad U_0(\gamma) = 0, \quad (28)$$

with a non-random  $\gamma = \gamma_n \in (0, 1)$  and with independent, identically distributed innovations  $X_1, X_2, \dots$ . As in the previous example, we assume that  $X_1$  satisfies (17) and that its distribution function is continuous. Consider the exit time

$$T(\gamma) := \inf\{n \geq 1 : U_n(\gamma) \leq 0\}.$$

We want to understand the behavior of the probability  $\mathbf{P}(T(\gamma) > n)$  in the case when  $\gamma = \gamma_n$  depends on  $n$  and

$$\gamma_n \in (0, 1) \quad \text{and} \quad \sup_n n(1 - \gamma_n) < \infty. \quad (29)$$

We now show that the autoregressive sequence  $U_n(\gamma)$  can be transformed to a random walk, which satisfies the conditions of Corollary 3. First, multiplying (28) by  $\gamma^{-n}$ , we get

$$U_n(\gamma) \gamma^{-n} = U_n(\gamma) \gamma^{-(n-1)} + X_n \gamma^{-n} = \sum_{k=1}^n \gamma^{-k} X_k, \quad n \geq 1.$$

Thus, for each  $n \geq 1$ ,

$$\{T(\gamma_n) > n\} = \left\{ \sum_{j=1}^k \gamma_n^{-j} X_j > 0 \quad \text{for all} \quad k \leq n \right\}. \quad (30)$$

Comparing (30) with (18) and (20), we see that we have a particular case of the model in Corollary 3 with  $u_{k,n} = \gamma_n^{-k}$  and  $G_{k,n} = 0$ . Clearly,  $u_{k,n} \rightarrow 1$  for every fixed  $k$ . Furthermore, we infer from (29) that

$$\gamma_n^{-n} = e^{-n \log \gamma_n} = e^{O(n|\gamma_n-1|)} = e^{O(1)}$$

and

$$\sigma_n^2(\gamma_n) := \frac{\gamma_n^{-2n} - 1}{1 - \gamma_n^2} = \gamma_n^{-2} + \gamma_n^{-4} + \dots + \gamma_n^{-2n} = ne^{O(1)}.$$

These relations imply that (23) and (21) are fulfilled. Applying Corollary 3, we arrive at

$$\sigma_n(\gamma_n) \mathbf{P}(T(\gamma_n) > n) \rightarrow \sqrt{\frac{2}{\pi \mathbf{E}X_1^2}} \mathbf{E}[-S_\tau] \in (0, \infty), \quad (31)$$

where  $\tau$  is defined in (27).  $\diamond$

**1.4. Discussion of the assumption (13).** Based on the validity of CLT and considerations in [2] one can expect that the Lindeberg condition will again be sufficient. However the following example shows that this is not the case and the situation is more complicated.

**Example 6.** Let  $X_2, X_3, \dots$  and  $Y_2, Y_3, \dots$  be mutually independent random variables such that

$$\mathbf{E}X_k = \mathbf{E}Y_k = 0, \quad \mathbf{E}X_k^2 = \mathbf{E}Y_k^2 = 1 \quad \text{and} \quad \mathbf{P}(|X_k| \leq M) = 1 \quad \text{for all } k \geq 2 \quad (32)$$

for some finite constant  $M$ . It is easy to see that the triangular array

$$X_{1,n} := \frac{Y_n}{\sqrt{n}}, \quad X_{k,n} := \frac{X_k}{\sqrt{n}}, \quad k = 2, 3, \dots, n; \quad n > 1 \quad (33)$$

satisfies the Lindeberg condition. Indeed,  $\sum_{i=1}^n \mathbf{E}X_{i,n}^2 = 1$  and for every  $\varepsilon > \frac{M}{\sqrt{n}}$  one has

$$\sum_{i=1}^n \mathbf{E}[X_{i,n}^2; |X_{i,n}| > \varepsilon] = \mathbf{E}[X_{1,n}^2; |X_{1,n}| > \varepsilon] \leq \mathbf{E}X_{1,n}^2 = \frac{\mathbf{E}Y_n^2}{n} = \frac{1}{n} \rightarrow 0 \quad (34)$$

due to the fact that  $|X_{k,n}| \leq \frac{M}{\sqrt{n}}$  for all  $k \geq 2$ .

We shall also assume that  $g_{k,n} \equiv 0$ . For each  $n > 1$  let random variables  $Y_n$  be defined as follows

$$Y_n := \begin{cases} N_n, & \text{with probability } p_n := \frac{1}{2N_n^2}, \\ 0, & \text{with probability } 1 - 2p_n, \\ -N_n, & \text{with probability } p_n, \end{cases} \quad (35)$$

where  $N_n \geq 1$ . Note that  $\mathbf{E}Y_n = 0$  and  $\mathbf{E}Y_n^2 = 1$ .

For every  $n > 1$  we set

$$U_n := X_2 + X_3 + \dots + X_n \quad \text{and} \quad \underline{U}_n := \min_{2 \leq i \leq n} U_i. \quad (36)$$

It is easy to see that

$$\{T_n > n\} = \{Y_n = N_n\} \cap \{\underline{U}_n > -N_n\}.$$

Noting now that  $\underline{U}_n \geq -(n-1)M$ , we infer that

$$\{T_n > n\} = \{Y_n = N_n\}, \quad \text{for any } N_n > (n-1)M. \quad (37)$$

In this case we have

$$\begin{aligned}
E_n &= \mathbf{E}[S_{n,n}; T_n > n] = \mathbf{E}\left[\frac{Y_n + U_n}{\sqrt{n}}; Y_n = N_n\right] \\
&= \mathbf{P}(Y_n = N_n)\mathbf{E}\left[\frac{N_n + U_n}{\sqrt{n}}\right] = \mathbf{P}(Y_n = n)\frac{N_n + \mathbf{E}U_n}{\sqrt{n}} \\
&= \mathbf{P}(Y_n = n)\frac{N_n}{\sqrt{n}}.
\end{aligned} \tag{38}$$

In particular, from (37) and (38) we conclude that

$$\mathbf{P}(T_n > n) = \mathbf{P}(Y_n = n) = \frac{E_n\sqrt{n}}{N_n} < \frac{E_n\sqrt{n}}{M(n-1)} = o(E_n)$$

provided that  $N_n > (n-1)M$ .

This example shows that (11) can not hold for all triangular arrays satisfying the Lindeberg condition.  $\diamond$

We now construct an array, for which the assumption (13) becomes necessary for the validity of (11).

**Example 7.** We consider again the model from the previous example and assume additionally that the variables  $X_2, X_3, \dots$  have the Rademacher distribution, that is,

$$\mathbf{P}(X_k = \pm 1) = \frac{1}{2}.$$

Finally, in order to have random walks on lattices, we shall assume that  $N_n$  is a natural number.

It is then clear that  $r_n := \frac{N_n}{\sqrt{n}}$  is the minimal deterministic number such that

$$\max_{k \leq n} |X_{k,n}| \leq r_n.$$

As in Example 6, we shall assume that  $g_{k,n} \equiv 0$ .

In order to calculate  $E_n$  we note that

$$\begin{aligned}
E_n &= \mathbf{E}[S_{n,n}; T_n > n] = \mathbf{P}(X_{1,n} = r_n) \mathbf{E}\left[r_n + \frac{U_n}{\sqrt{n}}; r_n + \frac{U_n}{\sqrt{n}} > 0\right] \\
&= \mathbf{P}(X_{1,n} = r_n) \frac{1}{\sqrt{n}} \mathbf{E}[N_n + U_n; N_n + \underline{U}_n > 0].
\end{aligned}$$

It is well known that for  $m \geq 1$  the sequence  $(N + U_m)1_{\{N + \underline{U}_m > 0\}}$  is a martingale with  $U_1 = \underline{U}_1 = 0$ . This implies that

$$\mathbf{E}[N + U_m; N + \underline{U}_m > 0] = N \quad \text{for all } m, N \geq 1.$$

Consequently,

$$E_n = p_n \frac{N_n}{\sqrt{n}} = p_n r_n. \tag{39}$$

Furthermore,

$$\mathbf{P}(T_n > n) = \mathbf{P}(X_{1,n} = r_n) \mathbf{P}\left(\frac{N_n}{\sqrt{n}} + \frac{U_n}{\sqrt{n}} > 0\right) = p_n \mathbf{P}(N_n + \underline{U}_n > 0).$$

Using the reflection principle for the symmetric simple random walk, one can show that

$$\mathbf{P}(N + \underline{U}_m > 0) = \mathbf{P}(-N < U_m \leq N) \quad \text{for all } m, N \geq 1. \tag{40}$$

Consequently,  $\mathbf{P}(T_n > n) = p_n \mathbf{P}(-N_n < U_n \leq N_n)$ . Combining this equality with (39), we obtain

$$\frac{\mathbf{P}(T_n > n)}{E_n} = \frac{1}{r_n} \mathbf{P}\left(-r_n < \frac{U_n}{\sqrt{n}} \leq r_n\right). \quad (41)$$

Using the central limit theorem, one obtains

$$\mathbf{P}\left(-r_n < \frac{U_n}{\sqrt{n}} \leq r_n\right) \sim \Psi(r_n), \quad (42)$$

where

$$\varphi(u) := \frac{1}{\sqrt{2\pi}} e^{-u^2/2} \quad \text{and} \quad \Psi(x) := 2 \int_0^{x^+} \varphi(u) du. \quad (43)$$

We will postpone the proof of (40) and (42) till the end of the paper. Assuming that (40) and (42) are true, as a result we have

$$\frac{\mathbf{P}(T_n > n)}{E_n} \sim \frac{\Psi(r_n)}{r_n}.$$

Noting now that  $\frac{\Psi(a)}{a} < 2\varphi(0) = \sqrt{\frac{2}{\pi}}$  for every  $a > 0$ , we conclude that the assumption  $r_n \rightarrow 0$  is necessary and sufficient for the validity of (11). More precisely,

- $\mathbf{P}(T_n > n) \sim \sqrt{\frac{2}{\pi}} E_n$  iff  $r_n \rightarrow 0$ ;
- $\mathbf{P}(T_n > n) \sim \frac{\Psi(a)}{a} E_n$  iff  $r_n \rightarrow a > 0$ ;
- $\mathbf{P}(T_n > n) = o(E_n)$  iff  $r_n \rightarrow \infty$ .

◇

## 2. PROOFS.

In this section we are going to obtain estimates, which are valid for each fixed  $n$ . For that reason we will sometimes omit the subscript  $n$  and introduce the following simplified notation:

$$T := T_n, \quad X_k := X_{k,n}, \quad S_k := S_{k,n}, \quad g_k := g_{k,n}, \quad 1 \leq k < n \quad (44)$$

and

$$\rho := r_n + g_n^*, \quad B_k^2 := \sum_{i=1}^k \mathbf{E}X_i^2, \quad B_{k,n}^2 := B_n^2 - B_k^2 = 1 - B_k^2, \quad 1 \leq k < n. \quad (45)$$

**2.1. Some estimates in the central limit theorem.** For every integer  $1 \leq k \leq n$  and every real  $y$  define

$$Z_k := S_k - g_k, \quad \widehat{Z}_k := Z_k \mathbf{1}\{T > k\} \quad \text{and} \quad Q_{k,n}(y) := \mathbf{P}\left(y + \min_{k \leq j \leq n} (Z_j - Z_k) > 0\right). \quad (46)$$

**Lemma 1.** *For all  $y \in \mathbb{R}$  and for all  $0 \leq k < n$  with  $B_{k,n} > 0$*

$$\left| Q_{k,n}(y) - \Psi\left(\frac{y}{B_{k,n}}\right) \right| \leq \frac{C_0 \rho}{B_{k,n}} \mathbf{1}\{y > 0\}, \quad (47)$$

where  $C_0$  is an absolute constant.



*Proof.* For non-random real  $y$  define

$$q_{k,n}(y) := \mathbf{P}\left(y + \min_{k \leq j \leq n} (S_j - S_k) > 0\right), \quad n > k \geq 1. \quad (48)$$

It follows from Corollary 1 in Arak [1] that there exists an absolute constant  $C_A$  such that

$$\left|q_{k,n}(y) - \Psi\left(\frac{y}{B_{k,n}}\right)\right| \leq \frac{C_A}{B_{k,n}} \max_{k < j \leq n} \frac{\mathbf{E}|X_j|^3}{\mathbf{E}X_j^2} \leq \frac{C_A r_n}{B_{k,n}}, \quad (49)$$

where maximum is taken over all  $j$  satisfying  $\mathbf{E}X_j^2 > 0$ . In the second step we have used the inequality  $\mathbf{E}|X_j|^3 \leq r_n \mathbf{E}X_j^2$  which follows from (13).

We have from (46) that  $|Z_k - S_k| = |g_k| \leq g_n^*$ . Hence, for  $Q_{k,n}$  and  $q_{k,n}$  defined in (46) and (48), we have

$$q_{k,n}(y_-) \leq Q_{k,n}(y) \leq q_{k,n}(y_+), \quad \text{where } y_{\pm} := y \pm 2g_n^*. \quad (50)$$

Then we obtain from (49) that

$$\left|q_{k,n}(y_{\pm}) - \Psi\left(\frac{y_{\pm}}{B_{k,n}}\right)\right| \leq \frac{C_A r_n}{B_{k,n}}. \quad (51)$$

On the other hand, it is easy to see from (43) that

$$\left|\Psi\left(\frac{y_{\pm}}{B_{k,n}}\right) - \Psi\left(\frac{y}{B_{k,n}}\right)\right| \leq \frac{2\varphi(0)|y_{\pm} - y|}{B_{k,n}} = \frac{4\varphi(0)g_n^*}{B_{k,n}}.$$

Applying this inequality together with (50) and (51) we immediately obtain (47) for  $y > 0$   $C_0 := C_A + 4\varphi(0)$ . For  $y \leq 0$  inequality (47) immediately follows since  $Q_{k,n}(y) = 0 = \Psi(y)$ .  $\square$

**Lemma 2.** *If  $1 \leq m \leq n$ , then*

$$\mathbf{E}S_m^+ \geq \frac{3}{8}B_m - r_n. \quad (52)$$

*Moreover, for all  $m$  satisfying  $B_m \geq 24(r_n + g_n^*)$  we have*

$$\mathbf{P}(T > m) \leq 3 \frac{\mathbf{E}\widehat{Z}_m}{B_m}. \quad (53)$$

*Proof.* We will use the following extension of the Berry-Esseen inequality due to Tyurin [5]:

$$\sup_{x \in \mathbb{R}} |\mathbf{P}(S_m > x) - \mathbf{P}(B_m \eta > x)| \leq 0.5606 \frac{\sum_{j=1}^m \mathbf{E}|X_j|^3}{B_m^3} \leq 0.5606 \frac{r_n}{B_m},$$

when  $B_m > 0$ . Here  $\eta$  is a random variable that follows the standard normal distribution. This inequality implies that, for every  $C > 0$ ,

$$\begin{aligned} \mathbf{E}S_m^+ &= \int_0^{\infty} \mathbf{P}(S_m > x) dx \geq \int_0^{CB_m} \mathbf{P}(S_m > x) dx \\ &\geq \int_0^{CB_m} \left( \mathbf{P}(B_m \eta > x) - 0.5606 \frac{r_n}{B_m} \right) dx = B_m \mathbf{E}(\eta^+ \wedge C) - 0.5606 C r_n. \end{aligned}$$

Further,

$$\begin{aligned}\mathbf{E}(\eta^+ \wedge C) &= \int_0^\infty (x \wedge C)\varphi(x)dx = \int_0^C x \frac{1}{2\pi} e^{-x^2/2} dx + C \int_C^\infty \varphi(x)dx \\ &= \varphi(0) - \varphi(C) + C \int_C^\infty \varphi(x)dx.\end{aligned}$$

Taking here  $C = 1/0.5606$  and using tables of the standard normal distribution we conclude that  $\mathbf{E}(\eta^+ \wedge C) > 0.375 > \frac{3}{8}$  and (52) holds.

Next, according to Lemma 25 in [2],

$$\mathbf{E}Z_m^+ \mathbf{P}(T > m) \leq \mathbf{E}\widehat{Z}_m, \quad 1 \leq m \leq n. \quad (54)$$

Therefore, it remains to derive a lower bound for  $\mathbf{E}Z_m^+$ . We first note that

$$S_m = Z_m + g_m \leq Z_m^+ + g_m^+ \leq Z_m^+ + g_n^*.$$

Hence,  $S_m^+ \leq Z_m^+ + g_n^*$  and, taking into account (52), we get

$$\mathbf{E}Z_m^+ \geq \mathbf{E}S_m^+ - g_n^* \geq \frac{3}{8}B_m - (r_n + g_n^*). \quad (55)$$

If  $m$  is such that  $\frac{B_m}{24} \geq r_n + g_n^*$ , then we infer from (54) and (55) that

$$\begin{aligned}\mathbf{E}\widehat{Z}_m &\geq \mathbf{E}Z_m^+ \mathbf{P}(T > m) \geq \left(\frac{3}{8}B_m - (r_n + g_n^*)\right) \mathbf{P}(T > n) \\ &\geq \left(\frac{3}{8} - \frac{1}{24}\right) B_m \mathbf{P}(T > m) = \frac{1}{3}B_m \mathbf{P}(T > m).\end{aligned}$$

Thus, (53) is proven.  $\square$

## 2.2. Estimates for expectations of $\widehat{Z}_k$ .

**Lemma 3.** *Let  $\alpha$  be a stopping time such that  $1 \leq \alpha \leq l \leq n$  with probability one. Then*

$$\mathbf{E}\widehat{Z}_\alpha - \mathbf{E}\widehat{Z}_l \leq 2g_n^* p(\alpha, l) \quad \text{with} \quad p(\alpha, l) := \mathbf{P}(\alpha < T, \alpha < l). \quad (56)$$

Moreover,

$$\mathbf{E}\widehat{Z}_\alpha - \mathbf{E}\widehat{Z}_l \geq \mathbf{E}[X_T; \alpha < T \leq l] - 2g_n^* p(\alpha, l) \geq -(2g_n^* + r_n)p(\alpha, l). \quad (57)$$

In addition, the equality in (12) takes place.

*Proof.* Define events

$$A_1 := \{\alpha < T \leq l\} \quad \text{and} \quad A_2 := \{\alpha < l < T\}.$$

Then, clearly,  $\{\alpha < T, \alpha < l\} = A_1 \cup A_2$ . Using Lemma 20 from [2], we obtain

$$\begin{aligned}\mathbf{E}\widehat{Z}_\alpha + \mathbf{E}[S_T; T \leq \alpha] &= -\mathbf{E}[g_\alpha; \alpha < T] \\ &= -\mathbf{E}[g_\alpha; A_2] - \mathbf{E}[g_l; \alpha = l < T] - \mathbf{E}[g_\alpha; A_1], \\ \mathbf{E}\widehat{Z}_l + \mathbf{E}[S_T; T \leq l] &= -\mathbf{E}[g_l; T > l] = -\mathbf{E}[g_l; A_2] - \mathbf{E}[g_l; \alpha = l < T].\end{aligned} \quad (58)$$

Thus,

$$\mathbf{E}\widehat{Z}_\alpha - \mathbf{E}\widehat{Z}_l = \mathbf{E}[S_T - g_\alpha; A_1] + \mathbf{E}[g_l - g_\alpha; A_2]. \quad (59)$$

Next, by the definition of  $T$ ,

$$g_T \geq S_T = S_{T-1} + X_T > g_{T-1} + X_T.$$

Hence,

$$\mathbf{E}[S_T - g_\alpha; A_1] \leq \mathbf{E}[g_T - g_\alpha; A_1] \leq 2g_n^* \mathbf{P}(A_1)$$

and

$$\begin{aligned} \mathbf{E}[S_T - g_\alpha; A_1] &\geq \mathbf{E}[g_{T-1} - g_\alpha + X_T; A_1] \\ &\geq \mathbf{E}[X_T; A_1] - 2g_n^* \mathbf{P}(A_1) \geq -(2g_n^* + r_n) \mathbf{P}(A_1). \end{aligned}$$

Furthermore,

$$|\mathbf{E}[g_n - g_\alpha; A_2]| \leq 2g_n^* \mathbf{P}(A_2).$$

Plugging these estimates into (59), we arrive at desired bounds.

The equality in (12) follows from (58) with  $l = n$ .  $\square$

For every  $h > 0$  define

$$\nu(h) := \inf\{k \geq 1 : S_k \geq g_k + h\} = \inf\{k \geq 1 : Z_k \geq h\}. \quad (60)$$

**Lemma 4.** *Suppose that  $m \leq n$  is such that the inequality (53) takes place,*

$$B_m \geq 24g_n^* \quad \text{and} \quad h \geq 6g_n^*. \quad (61)$$

Then

$$2\mathbf{E}\widehat{Z}_{\nu(h)\wedge m} \leq 3\mathbf{E}\widehat{Z}_m \leq 4\mathbf{E}\widehat{Z}_n = 4E_n, \quad \mathbf{P}(\widehat{Z}_{\nu(h)\wedge m} > 0) \leq \varkappa E_n, \quad (62)$$

$$2\varkappa g_n^* E_n \geq \mathbf{E}\widehat{Z}_{\nu(h)\wedge m} - E_n \geq \delta(h) - 2\varkappa g_n^* E_n, \quad (63)$$

where

$$0 \geq \delta(h) := \mathbf{E}[X_T; n \geq T > \nu(h) \wedge m] \geq -\varkappa r_n E_n \quad \text{and} \quad \varkappa := \frac{2}{h} + \frac{4}{B_m}. \quad (64)$$

In particular, (16) takes place.

*Proof.* First, we apply Lemma 3 with  $l = m$  and  $\alpha = \nu(h) \wedge m$ . For this choice of the stopping time one has

$$\begin{aligned} p(\nu(h) \wedge m, m) &= \mathbf{P}(\nu(h) \wedge m < T, \nu(h) \wedge m < m) \\ &\leq \mathbf{P}(\widehat{Z}_{\nu(h)\wedge m} \geq h) \leq \frac{\mathbf{E}\widehat{Z}_{\nu(h)\wedge m}}{h}. \end{aligned}$$

Plugging this bound into (56) and using the inequality  $h \geq 6g_n^*$ , we get

$$\mathbf{E}\widehat{Z}_{\nu(h)\wedge m} - \mathbf{E}\widehat{Z}_m \leq \frac{2g_n^*}{h} \mathbf{E}\widehat{Z}_{\nu(h)\wedge m} \leq \frac{\mathbf{E}\widehat{Z}_{\nu(h)\wedge m}}{3}$$

and hence

$$\frac{2}{3} \mathbf{E}\widehat{Z}_{\nu(h)\wedge m} \leq \mathbf{E}\widehat{Z}_m. \quad (65)$$

Next, we apply Lemma 3 with  $l = n$  and  $\alpha = m$ . In this case  $p(m, n) = \mathbf{P}(T > m)$  and we may use (53). Substituting these estimates into (56) and using (61), we obtain

$$\mathbf{E}\widehat{Z}_m - \mathbf{E}\widehat{Z}_n \leq 2g_n^* \mathbf{P}(T > m) \leq \frac{6g_n^*}{B_m} \mathbf{E}\widehat{Z}_m \leq \frac{1}{4} \mathbf{E}\widehat{Z}_m.$$

Therefore,

$$\frac{3}{4} \mathbf{E}\widehat{Z}_m \leq \mathbf{E}\widehat{Z}_n. \quad (66)$$

We conclude from (65) and (66) that the first relation in (62) takes place. In particular, from (53) and (66) we get that (16) holds under assumptions of Lemma 4.

At last, we are going to apply Lemma 3 with  $l = n > m$  and  $\alpha = \nu(h) \wedge m$ . For this choice of the stopping time one has

$$\begin{aligned} p(\nu(h) \wedge m, n) &= \mathbf{P}(T > \nu(h) \wedge m) = \mathbf{P}(\widehat{Z}_{\nu(h) \wedge m} > 0) \\ &\leq \mathbf{P}(\widehat{Z}_{\nu(h) \wedge m} \geq h) + \mathbf{P}(T > m) \\ &\leq \frac{\mathbf{E}\widehat{Z}_{\nu(h) \wedge m}}{h} + \frac{3\mathbf{E}\widehat{Z}_m}{B_m} \leq \frac{2E_n}{h} + \frac{4E_n}{B_m} = \varkappa E_n. \end{aligned} \quad (67)$$

Plugging this bound into (56) and (57), we immediately obtain (63). The second inequality in (62) also follows from (67); and using (13) together with (67) we find (64).

Thus, all assertions of Lemma 4 are proved.  $\square$

**2.3. Proof of Theorem 1.** According to the representation (36) in [2],

$$\begin{aligned} \mathbf{P}(T > n) &= \mathbf{E}[Q_{\nu(h) \wedge m, n}(Z_{\nu(h) \wedge m}); T > \nu(h) \wedge m] \\ &= \mathbf{E}Q_{\nu(h) \wedge m, n}(\widehat{Z}_{\nu(h) \wedge m}). \end{aligned} \quad (68)$$

**Lemma 5.** *Suppose that all assumptions of Lemma 4 are fulfilled and that  $B_{m,n} > 0$ . Then one has*

$$\begin{aligned} \left| \mathbf{P}(T > n) - \mathbf{E}\Psi\left(\frac{\widehat{Z}_{\nu(h) \wedge m}}{B_{\nu(h) \wedge m, n}}\right) \right| &\leq \frac{C_0\rho}{B_{m,n}} \mathbf{P}(\widehat{Z}_{\nu(h) \wedge m} > 0) \\ &\leq 2\varphi(0) \frac{1.3C_0\varkappa\rho E_n}{B_{m,n}}. \end{aligned} \quad (69)$$

In addition,

$$\mathbf{E}\Psi\left(\frac{\widehat{Z}_{\nu(h) \wedge m}}{B_{\nu(h) \wedge m, n}}\right) \leq \frac{2\varphi(0)E_n(1 + 2\varkappa g_n^*)}{B_{m,n}}, \quad (70)$$

$$\mathbf{E}\Psi\left(\frac{\widehat{Z}_{\nu(h) \wedge m}}{B_{\nu(h) \wedge m, n}}\right) \geq 2\varphi(0)E_n \left(1 - \frac{(r_n + h)^2}{6} - 2\varkappa g_n^* - \varkappa r_n\right). \quad (71)$$

*Proof.* Using (47) with  $y = \widehat{Z}_{\nu(h) \wedge m}$ , we obtain the first inequality in (69) as a consequence of (68). The second inequality in (69) follows from (62).

Next, it has been shown in [2, p. 3328] that

$$2\varphi(0)a \geq \Psi(a) \geq 2\varphi(0)a(1 - a^2/6) \quad \text{for all } a \geq 0. \quad (72)$$

Recall that  $0 \leq z := \widehat{Z}_{\nu(h) \wedge m} \leq r_n + h$  and  $B_n = 1$ . Hence, by (72),

$$\Psi\left(\frac{z}{B_{\nu(h) \wedge m, n}}\right) \leq \Psi\left(\frac{z}{B_{m,n}}\right) \leq \frac{2\varphi(0)z}{B_{m,n}}, \quad (73)$$

$$\Psi\left(\frac{z}{B_{\nu(h) \wedge m, n}}\right) \geq \Psi\left(\frac{z}{B_n}\right) \geq \frac{2\varphi(0)z}{B_n} \left(1 - \frac{z^2}{6B_n^2}\right) \geq 2\varphi(0)z \left(1 - \frac{(r_n + h)^2}{6}\right). \quad (74)$$

Taking mathematical expectations in (73) and (74) with  $z = \widehat{Z}_{\nu(h) \wedge m}$ , we obtain:

$$\frac{2\varphi(0)\mathbf{E}\widehat{Z}_{\nu(h) \wedge m}}{B_{m,n}} \geq \mathbf{E}\Psi\left(\frac{\widehat{Z}_{\nu(h) \wedge m}}{B_{\nu(h) \wedge m, n}}\right) \geq 2\varphi(0)\mathbf{E}\widehat{Z}_{\nu(h) \wedge m} \left(1 - \frac{(r_n + h)^2}{6}\right). \quad (75)$$

Now (70) and (71) follow from (75) together with (62) and (63).  $\square$

**Lemma 6.** *Assume that  $\rho \leq 1/64$ . Then inequalities (14) and (15) take place with some absolute constants  $C_1$  and  $C_2$ .*

*Proof.* Set

$$m := \min\{j \leq n : B_j \geq \frac{3}{2}\rho^{1/3}\} \quad \text{and} \quad h := \rho^{1/3}. \quad (76)$$

Noting that  $r_n \leq \rho \leq \rho^{1/3}/4^2$  we obtain

$$B_m^2 = B_{m-1}^2 + \mathbf{E}X_m^2 < \left(\frac{3}{2}\rho^{1/3}\right)^2 + r_n^2 \leq \frac{9}{4}\rho^{2/3} + \frac{1}{4^6} < \frac{1}{7}. \quad (77)$$

Consequently,  $B_{m,n}^2 = 1 - B_m^2$  and we have from (76) that

$$B_{m,n}^2 > \frac{6}{7}, \quad 24\rho \leq \frac{24}{4^2}\rho^{1/3} = \frac{3}{2}\rho^{1/3} \leq B_m, \quad 6g_n < \frac{6}{4^2}\rho^{1/3} < \rho^{1/3} = h. \quad (78)$$

Thus, all assumptions of Lemmas 4 and 5 are satisfied. Hence, Lemma 5 implies that

$$2\varphi(0)E_n(1 - \rho_1 - \rho_2 - 2\kappa\rho) \leq \mathbf{P}(T > n), \quad (79)$$

$$\mathbf{P}(T > n) \leq 2\varphi(0)E_n(1 + \rho_1)(1 + 2\kappa\rho)(1 + \rho_3), \quad (80)$$

where we used that  $2g_n^* + r_n \leq 2\rho$  and

$$\rho_1 := 1.3C_0\kappa\rho, \quad \rho_2 := \frac{(r_n + h)^2}{6}, \quad \rho_3 := \frac{1}{B_{m,n}} - 1. \quad (81)$$

Now from (64) and (76) with  $\rho^{1/3} \leq 1/4$  we have

$$\rho\kappa = \frac{2\rho}{h} + \frac{4\rho}{B_m} \leq 2\rho^{2/3} + \frac{4\rho^{2/3}}{3/2} < 4.7\rho^{2/3}, \quad r_n + h \leq \frac{1}{4^2}\rho^{1/3} + \rho^{1/3}.$$

Then, by (77),

$$\frac{1}{B_{m,n}} = \frac{B_{m,n}}{B_{m,n}^2} = \frac{\sqrt{1 - B_m^2}}{1 - B_m^2} \leq \frac{1 - B_m^2/2}{1 - B_m^2} = 1 + \frac{B_m^2}{2B_{m,n}^2} < 1 + 1.4\rho^{2/3}.$$

So, these calculations and (81) yield

$$\rho_1 < 5C_0\rho^{2/3}, \quad \rho_2 < 0.2\rho^{2/3}, \quad \rho_3 < 1.4\rho^{2/3}, \quad 2\kappa\rho < 9.4\rho^{2/3}. \quad (82)$$

Substituting (82) into (79) we obtain (14) with any  $C_1 \geq 5C_0 + 9.6$ . On the other hand from (82) and (80) we may obtain (15) with a constant  $C_2$  which may be calculated in the following way:

$$C_2 = \sup_{\rho^{1/3} \leq 1/4} [5C_0(1 + 2\kappa\rho)(1 + \rho_3) + 9.4(1 + \rho_3) + 1.4] < \infty.$$

$\square$

Thus, when  $\rho \leq 1/4^3$ , the both assertions of Theorem 1 immediately follow from Lemma 6. But if  $\rho > 1/4^3$  then (15) is valid with any  $C_1 \geq 4^2 = 16$  because in this case right-hand side in (15) is negative.

Let us turn to the upper bound (15). If  $\rho \leq \frac{1}{24}$  but  $\rho > \frac{1}{64}$  then (16) holds for  $m = n$ ; and as a result we have from (16) with any  $C_2 \geq 32/\varphi(0)$  that

$$\mathbf{P}(T_n > n) \leq 4E_n \leq 4^3 E_n \rho^{2/3} \leq 2\varphi(0)E_n(1 + C_2\rho^{2/3}) \quad \text{for} \quad \rho^{1/3} > 1/4.$$

So, we have proved all assertions of Theorem 1 in all cases.

**2.4. Proof of Corollary 3.** In order to apply Corollary 2 we introduce the following triangular array:

$$X_{j,n} := \frac{u_{j,n}X_j}{\sigma_n}, \quad g_{j,n} := \frac{G_{j,n}}{\sigma_n}, \quad 1 \leq j \leq n, \quad n \geq 1. \quad (83)$$

The assumptions in (21) and (22) imply that the array introduced in (83) satisfies (13) and (5). Thus,

$$\begin{aligned} \mathbf{P}(\tau_n > n) &= \mathbf{P}(T_n > n) \sim \sqrt{\frac{2}{\pi}} \mathbf{E}[S_{n,n} - g_{n,n}; T_n > n] \\ &= \sqrt{\frac{2}{\pi}} \left( \mathbf{E}[S_{n,n}; T_n > n] - g_{n,n} \mathbf{P}(T_n > n) \right). \end{aligned}$$

Here we also used (12). Since  $g_{n,n} \rightarrow 0$ , we conclude that

$$\mathbf{P}(\tau_n > n) \sim \sqrt{\frac{2}{\pi}} \mathbf{E}[S_{n,n}; T_n > n].$$

Noting that  $S_{n,n} = U_{n,n}/\sigma_n$ , we get

$$\mathbf{P}(\tau_n > n) \sim \sqrt{\frac{2}{\pi}} \frac{1}{\sigma_n} \mathbf{E}[U_{n,n}; \tau_n > n]. \quad (84)$$

By the optional stopping theorem,

$$\mathbf{E}[U_{n,n}; \tau_n > n] = -\mathbf{E}[U_{\tau_n,n}; \tau_n \leq n].$$

It follows from (23) that, for every fixed  $k \geq 1$ ,

$$U_{k,n} \rightarrow U_k \text{ a.s.} \quad (85)$$

and, taking into account the continuity of distribution functions,

$$\begin{aligned} \mathbf{P}(\tau_n > k) &= \mathbf{P}(U_{1,n} > G_{1,n}, U_{2,n} > G_{2,n}, \dots, U_{k,n} > G_{k,n}) \\ &\rightarrow \mathbf{P}(U_1 > g_1, U_2 > g_2, \dots, U_k > g_k) = \mathbf{P}(\tau > k). \end{aligned} \quad (86)$$

Obviously, (86) implies that

$$\mathbf{P}(\tau_n = k) \rightarrow \mathbf{P}(\tau = k) \quad \text{for every } k \geq 1. \quad (87)$$

Furthermore, it follows from the assumptions (17) and (21) that

$$|U_{\tau_n,n}| \leq M \quad \text{on the event } \{\tau_n \leq n\}. \quad (88)$$

Then, combining (85), (87) and (88), we conclude that

$$\mathbf{E}[U_{\tau_n,n}; \tau_n \leq k] = \sum_{j=1}^k \mathbf{E}[U_{j,n}; \tau_n = j] \rightarrow \sum_{j=1}^k \mathbf{E}[U_j; \tau = j] = \mathbf{E}[U_\tau; \tau \leq k]. \quad (89)$$

Note also that, by (88) and (86),

$$\limsup_{n \rightarrow \infty} |\mathbf{E}[U_{\tau_n,n}; k < \tau_n \leq n]| \leq M \limsup_{n \rightarrow \infty} \mathbf{P}(\tau_n > k).$$

Therefore,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \mathbf{E}[U_{\tau_n,n}; \tau_n \leq n] &\leq \limsup_{n \rightarrow \infty} \mathbf{E}[U_{\tau_n,n}; \tau_n \leq k] + \limsup_{n \rightarrow \infty} |\mathbf{E}[U_{\tau_n,n}; k < \tau_n \leq n]| \\ &= \mathbf{E}[U_\tau; \tau \leq k] + M \mathbf{P}(\tau > k) \end{aligned} \quad (90)$$

and

$$\begin{aligned} \liminf_{n \rightarrow \infty} \mathbf{E}[U_{\tau_n, n}; \tau_n \leq n] &\geq \liminf_{n \rightarrow \infty} \mathbf{E}[U_{\tau_n, n}; \tau_n \leq k] - \limsup_{n \rightarrow \infty} |\mathbf{E}[U_{\tau_n, n}; k < \tau_n \leq n]| \\ &= \mathbf{E}[U_\tau; \tau \leq k] - M\mathbf{P}(\tau > k). \end{aligned} \quad (91)$$

Letting  $k \rightarrow \infty$  in (90) and (91), and noting that  $\tau$  is almost surely finite, we infer that

$$\mathbf{E}[U_{\tau_n, n}; \tau_n \leq n] \rightarrow \mathbf{E}[U_\tau].$$

Consequently, by the optional stopping theorem,

$$\mathbf{E}[U_{\tau_n, n}; \tau_n > n] = -\mathbf{E}[U_{\tau_n, n}; \tau_n \leq n] \rightarrow \mathbf{E}[-U_\tau].$$

Plugging this into (84), we obtain the desired result.

### 2.5. Calculations related to Example 7.

**Lemma 7.** *For the simple symmetric random walk  $\{U_m\}$  one has*

$$\mathbf{P}(N + \underline{U}_m > 0) = \mathbf{P}(-N < U_m \leq N) \quad \text{for all } m, N \geq 1$$

and

$$\sup_{N \geq 1} \left| \frac{\mathbf{P}(-N < U_n \leq N)}{\Psi(N/\sqrt{n})} - 1 \right| \rightarrow 0.$$

*Proof.* By the reflection principle for symmetric simple random walks,

$$\mathbf{P}(N + U_m = k, N + \underline{U}_m \leq 0) = \mathbf{P}(U_m = N + k) \quad \text{for every } k \geq 1.$$

Thus, by the symmetry of the random walk  $U_m$ ,

$$\mathbf{P}(N + U_m > 0, N + \underline{U}_m \leq 0) = \mathbf{P}(U_m < -N) = \mathbf{P}(U_m > N).$$

Therefore,

$$\begin{aligned} \mathbf{P}(N + \underline{U}_m > 0) &= \mathbf{P}(N + U_m > 0) - \mathbf{P}(N + U_m > 0, N + \underline{U}_m \leq 0) \\ &= \mathbf{P}(U_m > -N) - \mathbf{P}(U_m > N) = \mathbf{P}(-N < U_m \leq N). \end{aligned}$$

We now prove the second statement. Recall that  $U_n$  is the sum of  $n - 1$  independent, Rademacher distributed random variables. By the central limit theorem,  $U_n/\sqrt{n} - 1$  converges to the standard normal distribution. Therefore,  $U_n/\sqrt{n}$  has the same limit. This means that

$$\varepsilon_n^2 := \sup_{x > 0} |\mathbf{P}(-x\sqrt{n} < U_n \leq x\sqrt{n}) - \Psi(x)| \rightarrow 0.$$

Taking into account that  $\Psi(x)$  increases, we conclude that, for every  $\delta > 0$ ,

$$\sup_{x \geq \delta} \left| \frac{\mathbf{P}(-x\sqrt{n} < U_n \leq x\sqrt{n})}{\Psi(x)} - 1 \right| \leq \frac{\varepsilon_n^2}{\Psi(\delta)}.$$

Choose here  $\delta = \varepsilon_n$ . Noting that  $\Psi(\varepsilon_n) \sim 2\varphi(0)\varepsilon_n$ , we obtain

$$\sup_{N \geq \varepsilon_n \sqrt{n}} \left| \frac{\mathbf{P}(-N < U_n \leq N)}{\Psi(N/\sqrt{n})} - 1 \right| \leq \frac{\varepsilon_n^2}{\Psi(\varepsilon_n)} \sim \frac{\varepsilon_n}{2\varphi(0)} \rightarrow 0.$$

It remains to consider the case  $N \leq \varepsilon_n \sqrt{n}$ . Here we shall use the local central limit theorem. Since  $U_n$  is 2-periodic,

$$\sup_{k: k \equiv n-1 \pmod{2}} |\sqrt{n-1} \mathbf{P}(U_n = k) - 2\varphi(k/\sqrt{n-1})| \rightarrow 0.$$

Noting that

$$\sup_{k \leq \varepsilon_n \sqrt{n}} |\varphi(k/\sqrt{n-1}) - \varphi(0)| \rightarrow 0,$$

we obtain

$$\sup_{N \leq \varepsilon_n \sqrt{n}} \left| \frac{\sqrt{n-1} \mathbf{P}(-N < U_n \leq N)}{2\varphi(0)m(n, N)} - 1 \right| \rightarrow 0,$$

where

$$m(n, N) = \#\{k \in (-N, N] : k \equiv n-1 \pmod{2}\}.$$

Since the interval  $(-N, N]$  contains  $N$  even and  $N$  odd lattice points,  $m(n, N) = N$  for all  $n, N \geq 1$ . Consequently,

$$\sup_{N \leq \varepsilon_n \sqrt{n}} \left| \frac{\sqrt{n-1} \mathbf{P}(-N < U_n \leq N)}{2\varphi(0)N} - 1 \right| \rightarrow 0,$$

It remains now to notice that

$$\Psi(N/\sqrt{n}) \sim \frac{2\varphi(0)N}{\sqrt{n}}$$

uniformly in  $N \leq \varepsilon_n \sqrt{n}$ . □

#### REFERENCES

- [1] Arak, T.V. On the distribution of the maximum of the successive partial sums of independent random variables. *Theory Probab. Appl.*, **19**, 245-266, 1975.
- [2] Denisov, D., Sakhanenko, A. and Wachtel, V. First-passage times for random walks with non-identically distributed increments. *Ann. Probab.* **46**(6): 3313-3350, 2018.
- [3] Denisov, D., Sakhanenko, A. and Wachtel, V. First-passage times for random walks without Lindeberg condition. work in progress.
- [4] Gaposhkin, V.F. The law of the iterated logarithm for Cesaro's and Abel's methods of summation. *Theory Probab. Appl.*, **10**:411-420, 1965.
- [5] Tyurin, I.S. Refinement of the upper bounds of the constants in Lyapunov's theorem. *Russian Math. Surveys* **65**(3):586-588, 2010.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANCHESTER, OXFORD ROAD, MANCHESTER M13 9PL, UK

*Email address:* `denis.denisov@manchester.ac.uk`

SOBOLEV INSTITUTE OF MATHEMATICS, 630090 NOVOSIBIRSK, RUSSIA

*Email address:* `aisakh@mail.ru`

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT AUGSBURG, 86135 AUGSBURG, GERMANY

*Email address:* `vitali.wachtel@math.uni-augsburg.de`