

# Fluctuations in Mean-Field Ising models

Nabarun Deb and Sumit Mukherjee \*

*Department of Statistics, Columbia University*  
*e-mail: nd2560@columbia.edu; sm3949@columbia.edu*

In this paper, we study the fluctuations of the average magnetization in an Ising model on an approximately  $d_N$  regular graph  $G_N$  on  $N$  vertices. In particular, if  $G_N$  is “well connected”, we show that whenever  $d_N \gg \sqrt{N}$ , the fluctuations are universal and same as that of the Curie-Weiss model in the entire Ferro-magnetic parameter regime. We give a counterexample to demonstrate that the condition  $d_N \gg \sqrt{N}$  is tight, in the sense that the limiting distribution changes if  $d_N \sim \sqrt{N}$  except in the high temperature regime. By refining our argument, we extend universality in the high temperature regime up to  $d_N \gg N^{1/3}$ . Our results conclude universal fluctuations of the average magnetization in Ising models on regular graphs, Erdős-Rényi graphs (directed and undirected), stochastic block models, and sparse regular graphons. In fact, our results apply to general matrices with non-negative entries, including Ising models on a Wigner matrix, and the block spin Ising model. As a by-product of our proof technique, we obtain Berry-Esseen bounds for these fluctuations, exponential concentration for the average of spins, and tight error bounds for the Mean-Field approximation of the partition function.

**MSC 2010 subject classifications:** Primary 82B20; secondary 82B26.

**Keywords and phrases:** Berry-Esseen bound, Ising model, Regular graphs, Mean-Field, Partition function.

## 1. Introduction

The Ising model is a discrete Markov random field which was initially introduced as a mathematical model of ferromagnetism in statistical physics, and has received extensive attention in Probability and Statistics (c.f. [1, 4, 5, 10, 14, 16, 17, 20, 21, 23, 24, 25, 26, 27, 29, 31, 32, 34, 35] and references therein). The model can be described by the following probability mass function in  $\sigma := (\sigma_1, \dots, \sigma_N) \in \{-1, 1\}^N$ :

$$\mathbb{P}(\sigma) := \frac{1}{Z_N(\beta, B)} \exp\left(\frac{\beta}{2} \sigma^\top A_N \sigma + B \sum_{i=1}^N \sigma_i\right). \quad (1.1)$$

Here  $A_N$  is a symmetric  $N \times N$  matrix with non-negative entries, and has zeroes on its diagonal, and  $\beta > 0$  and  $B \in \mathbb{R}$  are scalar parameters often referred to in the Statistical Physics literature as *inverse temperature* and *external magnetic field* respectively. The factor  $Z_N(\beta, B)$  is the normalizing constant/partition function of the model. The most common choice of the coupling matrix  $A_N$  is the adjacency matrix of graph  $G_N$  on  $N$  vertices, scaled by the average degree  $\bar{d}_N := \frac{1}{N} \sum_{i,j=1}^N G_N(i, j)$ . Here and throughout the rest of the paper, we use the notation  $G_N$  to denote both a graph and its adjacency matrix. A pivotal quantity of interest which has attracted extensive attention in the literature is the average sum of spins/magnetization density, defined by

$$\bar{\sigma} := \frac{\sum_{i=1}^N \sigma_i}{N}.$$

The fluctuations for  $\bar{\sigma}$  are mostly known for very few choices of the graph  $G_N$ , including the complete graph (see e.g., [14, 19, 21]), the directed Erdős-Rényi graph (see [26]), sparse Erdős-Rényi graphs (see [24]). In this paper, we focus on studying fluctuations of  $\bar{\sigma}$ , when  $A_N$  is the scaled adjacency matrix of an approximately regular graph  $G_N$ . The motivation for this work is the recent paper [4], where the authors show universal asymptotics of the partition function  $Z_N(\beta, B)$  on *any* sequence of approximately regular graphs with diverging average degree, which is governed by the Mean-Field prediction formula. In particular, it follows from [4, Theorem 2.1] that the Mean-Field prediction formula is asymptotically correct for any sequence of approximately  $d_N$  regular graphs  $G_N$  with  $d_N \rightarrow \infty$ . A natural follow up question is to what extent this universality extends to other properties of such “Mean-Field” Ising models. In this paper we try to address this question partially by studying universal behavior of the statistic  $\bar{\sigma}$ .

\*Research partially supported by NSF grant DMS-1712037

### 1.1. Main results

We begin with a definition which partitions the parameter set  $\{(\beta, B) : \beta > 0, B \in \mathbb{R}\}$  into different domains.

**Definition 1.1.** Let  $\Theta_{11} := \{(\beta, 0) : 0 < \beta < 1\}$ ,  $\Theta_{12} := \{(\beta, B) : \beta > 0, B \neq 0\}$ ,  $\Theta_2 := \{(\beta, 0) : \beta > 1\}$ ,  $\Theta_3 := (1, 0)$ . Finally, let  $\Theta_1 := \Theta_{11} \cup \Theta_{12}$ . We will refer to  $\Theta_1$  as the uniqueness regime,  $\Theta_2$  as the non uniqueness regime, and  $\Theta_3$  as the critical point. The names of the different regimes are by motivated by next lemma, the proof of which follows from simple calculus (see for e.g. [17, Page 10]).

**Lemma 1.1.** Consider the fixed point equation

$$\phi(x) = 0, \text{ where } \phi(x) := x - \tanh(\beta x + B). \quad (1.2)$$

- (a) If  $(\beta, B) \in \Theta_{11}$ , then (1.2) has a unique solution at  $t = 0$ , and  $\phi'(0) > 0$ .
- (b) If  $(\beta, B) \in \Theta_{12}$ , then (1.2) has a unique root  $t$  with the same sign as that of  $B$ , and  $\phi'(t) > 0$ .
- (c) If  $(\beta, B) \in \Theta_2$ , then (1.2) has two non zero roots  $\pm t$  of this equation, where  $t > 0$ , and  $\phi'(\pm t) > 0$ .
- (d) If  $(\beta, B) \in \Theta_3$ , then (1.2) has a unique solution at  $t = 0$ , and  $\phi'(0) = 0$ .

We will use  $t$  as defined in the above lemma throughout the paper, noting that  $t$  does depend on  $(\beta, B)$ . The following result summarizes the fluctuations of  $\bar{\sigma}$  in the Curie-Weiss model (see [21]), which is the Ising model on the complete graph.

**Lemma 1.2.** Suppose  $\sigma$  is a random vector from the Curie Weiss model  $\mathbb{P}^{CW}$  with p.m.f.

$$\mathbb{P}^{CW}(\sigma) = \frac{1}{Z_N^{CW}(\beta, B)} \exp\left(\frac{N\beta}{2}\bar{\sigma}^2 + B \sum_{i=1}^N \sigma_i\right). \quad (1.3)$$

Let  $Z_\tau \sim N(0, \tau)$  with  $\tau := \frac{1-t^2}{1-\beta(1-t^2)}$  for  $(\beta, B) \notin \Theta_3$ , and let  $W$  be a continuous random variable with density proportional to  $e^{-x^4/12}$ . Then the following holds:

$$\begin{aligned} \sqrt{N}(\bar{\sigma} - t) &\xrightarrow{d} Z_\tau && \text{if } (\beta, B) \in \Theta_1, \\ \sqrt{N}(\bar{\sigma} - M(\sigma)) &\xrightarrow{d} Z_\tau && \text{if } (\beta, B) \in \Theta_2, \\ N^{1/4}\bar{\sigma} &\xrightarrow{d} W && \text{if } (\beta, B) \in \Theta_3. \end{aligned}$$

Here  $M(\sigma)$  is a random variable which equals  $t$  if  $\bar{\sigma} \geq 0$ , and  $-t$  otherwise.

We will now explore to what extent the fluctuations of  $\bar{\sigma}$  are universal. We need the following notations to state our main results.

**Definition 1.2.** (i) Given two positive sequences  $x_N, y_N$ , we use the notation  $x_N \lesssim y_N$  to denote the existence of a finite constant  $C$  free of  $N$ , such that  $x_N \leq C y_N$ .

(ii) Given a symmetric matrix  $A_N$ , let  $R_i := \sum_{j=1}^N A_N(i, j)$  denote the row sums of  $A_N$ , and let  $(\lambda_1(A_N), \dots, \lambda_N(A_N))$  denote its eigenvalues arranged in decreasing order. Let  $\|A_N\|_F$  and  $\|A_N\|_{op}$  denote the Frobenius norm and operator norm of  $A_N$  respectively.

(iii) Given two real valued random variables  $X, Y$ , define the Kolmogorov-Smirnov distance between  $X$  and  $Y$  by

$$d_{KS}(X, Y) := \sup_{x \in \mathbb{R}} |\mathbb{P}(X \leq x) - P(Y \leq x)|.$$

**Theorem 1.1.** Suppose that  $(\beta, B) \in \Theta_1$ . Assume further that the sequence of matrices  $A_N$  satisfies the following two conditions:

$$\max_{1 \leq i \leq N} R_i \lesssim 1, \quad (1.4)$$

$$\lim_{N \rightarrow \infty} \lambda_1(A_N) = 1. \quad (1.5)$$

If  $\sigma$  is a random vector from the Ising model (1.1), then we have

$$d_{KS}\left(\sqrt{N}(\bar{\sigma} - t), Z_\tau\right) \lesssim \frac{1}{\sqrt{N}} \left( \|A_N\|_F^2 + \sum_{i=1}^N (R_i - 1)^2 + t \left| \sum_{i=1}^N (R_i - 1) \right| \right). \quad (1.6)$$

Note that Theorem 1.1 leaves out the parameter regime  $\Theta_2 \cup \Theta_3$ . The following example shows that such a universal behavior is not expected in this parameter regime, unless we assume some notion of connectivity for  $A_N$ .

**Example 1.1.** *With  $N$  even, let  $A_N$  be the adjacency matrix of two disjoint complete graphs  $K_{N/2}$ , scaled by  $N/2$ . Then the following holds:*

(a) *If  $(\beta, B) \in \Theta_2$ , then  $\bar{\sigma} \xrightarrow{d} \frac{1}{2}\delta_0 + \frac{1}{4}(\delta_t + \delta_{-t})$ .*

(b) *If  $(\beta, B) \in \Theta_3$ , then  $N^{1/4}\bar{\sigma} \xrightarrow{d} (W_1 + W_2)/2^{3/4}$ , where  $W_1, W_2$  are i.i.d. with the same distribution as that of  $W$ .*

The above example shows that if we want universal fluctuations in the regimes  $\Theta_2 \cup \Theta_3$ , the matrix  $A_N$  needs to be ‘‘well connected’’ in some asymptotic sense. If  $A_N$  is exactly the adjacency matrix of a  $d_N$  regular graph  $G_N$  scaled by  $d_N$ , then  $\lambda_1(A_N) = 1$ , and it is easy to check that the graph  $G_N$  is connected iff  $\lambda_2(A_N) < 1$ . Motivated by this, we propose the following asymptotic notion of well connectedness.

**Definition 1.3.** *We say a sequence of symmetric matrices  $\{A_N\}_{N \geq 1}$  with non-negative entries is well connected, if*

$$\limsup_{N \rightarrow \infty} \frac{\lambda_2(A_N)}{\lambda_1(A_N)} < 1. \quad (1.7)$$

We note that assumption (1.7) is somewhat weak in the sense that it does not imply connectivity in general. In particular this allows the existence of small disconnected subgraphs in  $G_N$ , as shown in the following example.

**Example 1.2.** *Let  $G_N$  denote a graph which is the disjoint union of a  $d_N$  regular graph  $G_{1, N_1}$  on  $N_1$  vertices, and an arbitrary graph  $G_{2, N_2}$  on  $N_2$  vertices, with  $N_1 + N_2 = N$  and  $N_2 = o(d_N)$ . Then the average degree of the whole graph  $G_N$  is  $\tilde{d}_N = d_N(1 + o(1))$ . It is easy to check that if  $G_{1, N_1}$  satisfies (1.7), then  $G_N$  satisfies (1.7), even though  $G_N$  is disconnected.*

Under the additional assumption of well connectedness, we show universal fluctuations in the non-uniqueness and critical regimes.

**Theorem 1.2.** *Suppose that  $(\beta, B) \in \Theta_2$ . Assume further that the sequence of matrices  $A_N$  satisfies (1.4), (1.5), and (1.7). If  $\sigma$  is a random vector from the Ising model (1.1), then we have*

$$d_{KS}\left(\sqrt{N}(\bar{\sigma} - M(\sigma)), Z_\tau\right) \lesssim \frac{1}{\sqrt{N}} \left( \|A_N\|_F^2 + \sum_{i=1}^N (R_i - 1)^2 + \left| \sum_{i=1}^N (R_i - 1) \right| \right). \quad (1.8)$$

where  $M(\sigma)$  is as in Lemma 1.2.

**Theorem 1.3.** *Suppose that  $(\beta, B) \in \Theta_3$ . If  $\sigma$  is a random vector from the Ising model (1.1) where  $A_N$  satisfies*

$$\limsup_{N \rightarrow \infty} N^{1/4} \max_{1 \leq i \leq N} |R_i - 1| \lesssim 1. \quad (1.9)$$

Then we have

$$d_{KS}\left(N^{1/4}\bar{\sigma}, W\right) \lesssim \frac{\varepsilon_N}{\sqrt{N}} + \frac{\varepsilon_N r_N}{N^{1/4}} + \frac{(\log N)^2}{N^{1/4}} \sqrt{\sum_{i=1}^N (R_i - 1)^2 + N^{-1/2} \left[ \sum_{i=1}^N (R_i - 1) \right]^2}, \quad (1.10)$$

where

$$r_N := \sqrt{(\log N)^3 \max_{1 \leq i \leq N} \sum_{j=1}^N A_N(i, j)^2 + \log N \max_{1 \leq i \leq N} |R_i - 1|},$$

$$\varepsilon_N := \|A_N\|_F^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1) \right]^2 + \frac{1}{N} \sum_{i=1}^N (R_i - 1)^2 + \log N,$$

and  $W$  is as in Lemma 1.2.

**Remark 1.1.** *Using these results, in section 1.2 we will show that for any sequence of well connected  $d_N$  regular graphs the fluctuation of  $\bar{\sigma}$  is universal in  $\Theta_1 \cup \Theta_2$  if  $d_N \gg \sqrt{N}$ , and in  $\Theta_3$  if  $d_N \gg \sqrt{N} \log N$ . We now give an example to show that the above conditions are actually necessary (up to log factor in the critical regime), which suggests that the convergence rates obtained in this paper are tight. The proof of this example will appear in an upcoming draft [36].*

**Example 1.3.** Let  $G_N$  denote the line graph of the complete graph  $K_n$ , so that  $N = \binom{n}{2} = \frac{n^2}{2}(1 + o(1))$ . This is a regular graph with degree  $d_N = 2(n-2) = 2\sqrt{2N}(1 + o(1))$ , and its top two eigenvalues are  $\lambda_1(G_N) = 2(n-2)$  and  $\lambda_2(G_N) = n-2$  (see [15, Lemma 2]). It follows that  $A_N = \frac{1}{d_N}G_N$  does satisfy (1.7), and

$$\lim_{N \rightarrow \infty} \frac{1}{\sqrt{N}} \|A_N\|_F^2 = \sqrt{N} \max_{1 \leq i \leq N} \sum_{j=1}^N A_N(i, j)^2 = \frac{1}{2\sqrt{2}} \neq 0.$$

In this case we have the following limiting distributions across different regimes:

$$\begin{aligned} \sqrt{N}(\bar{\sigma}_N - t) + \mu &\xrightarrow{w} Z_\tau && \text{if } (\beta, B) \in \Theta_1, \\ \sqrt{N}(\bar{\sigma}_N - M(\sigma)) + \text{sgn}(M(\sigma))\mu &\xrightarrow{w} Z_\tau && \text{if } (\beta, B) \in \Theta_2, \\ N^{1/4}\bar{\sigma}_N &\xrightarrow{w} \widetilde{W} && \text{if } (\beta, B) \in \Theta_3, \end{aligned}$$

where  $\mu := \frac{\beta t}{\sqrt{2}(1-\beta(1-t^2)) \cdot (2-\beta(1-t^2))} \geq 0$ , and  $\widetilde{W}$  has density proportional to  $\exp(-\frac{w^4}{12} - \frac{w^2}{\sqrt{2}})$ . Therefore, the fluctuations do not match that of the Curie-Weiss model unless  $(\beta, B) \in \Theta_{11}$ .

Note that in the above example,  $\bar{\sigma}$  has a different limit compared to the Curie-Weiss model in  $\Theta_{12} \cup \Theta_2 \cup \Theta_3$ , but continues to have universal fluctuations in the high parameter regime  $\Theta_{11}$ . We now state a modified theorem for the regime  $\Theta_{11}$ , which shows that in this regime we can do better.

**Theorem 1.4.** Suppose that  $(\beta, B) \in \Theta_{11}$ , and  $A_N$  satisfies

$$\lim_{N \rightarrow \infty} \max_{1 \leq i \leq N} R_i = 1. \quad (1.11)$$

If  $\sigma$  is a random vector from the Ising model (1.1), then setting  $\alpha_N := \max_{1 \leq i \leq N} \sum_{j=1}^N A_N(i, j)^2$  we have

$$d_{KS}(\sqrt{N}\bar{\sigma}, Z_\tau) \lesssim \frac{1}{\sqrt{N}} + \frac{\|A_N\|_F^2 \sqrt{\alpha_N \log N}}{\sqrt{N}} + \left[1 + \|A_N\|_F \alpha_N \log N\right] \sqrt{\frac{\sum_{i=1}^N (R_i - 1)^2}{N}}, \quad (1.12)$$

**Remark 1.2.** It follows from the above result that in the regime  $\Theta_{11}$ ,  $\bar{\sigma}$  has universal fluctuations on regular graphs of degree  $d_N \gg (N \log N)^{1/3}$ . We believe this is not tight, and universal fluctuations should hold on any sequence of regular graphs with  $d_N \rightarrow \infty$ . In [26] the authors prove such a result when  $G_N$  is a non symmetric Erdős-Rényi graph in the regime  $\Theta_{11}$  (details in example section below).

Note that we only expect similar behavior as the Curie Weiss model, if the underlying graphs are approximately regular and have large degree. Quantifying this philosophy, the bounds in each of the theorems have two terms, the first term controls the sparsity of the underlying graph/matrix, and the second term controls the extent of regularity of the graph/matrix. Recall example 1.3, which suggests that the term controlling the sparsity is optimal. In a similar spirit, the following example suggests that the term controlling the extent of regularity is also optimal.

**Example 1.4.** (a) Assume that  $\sqrt{N}$  is an integer, and let  $G_N$  be the disjoint union of two complete graphs of size  $N - \sqrt{N}$  and  $\sqrt{N}$  respectively. Let  $\bar{d}_N$  denote the average degree of  $G_N$  and  $A_N = (\bar{d}_N)^{-1}G_N$ . In this case

$$\lim_{N \rightarrow \infty} \frac{1}{\sqrt{N}} \sum_{i=1}^N (R_i - 1)^2 > 0,$$

but every other term in the RHS of (1.6) converges to 0. If  $\sigma$  is a random vector from the Ising model (1.1) with  $B \neq 0$ , then  $\sqrt{N}(\bar{\sigma} - t) \xrightarrow{w} \mu + Z_\tau$ , where  $\mu := \frac{\beta t(1-t^2)}{1-\beta(1-t^2)} + \tanh(B) - t \neq 0$ .

(b) With  $G_N = K_N$ , let  $A_N = \frac{1}{N-\sqrt{N}}G_N$ . In this case

$$\lim_{N \rightarrow \infty} \frac{1}{\sqrt{N}} \sum_{i=1}^N (R_i - 1) > 0,$$

but every other term in the RHS of (1.6) converges to 0. If  $\sigma$  is a random vector from the Ising model (1.1) with  $B \neq 0$ , then  $\sqrt{N}(\bar{\sigma} - t) \xrightarrow{w} \mu + Z_\tau$ , where  $\mu := \frac{\beta t(1-t^2)}{1-\beta(1-t^2)} \neq 0$ .

The main ingredient of our proof technique is comparing the Ising model on an approximately regular graph to that of an i.i.d. model/Curie-Weiss model. As a byproduct of this approach, we also obtain quantitative bounds for the following asymptotics of the log partition function via the Mean-Field prediction formula, defined via the following lower bound (c.f. [4]):

$$Z_N(\beta, B) \geq \sup_{\sigma \in \{-1, 1\}^N} \left\{ \frac{\beta}{2} \sigma^\top A_N \sigma + B \sum_{i=1}^N \sigma_i - \sum_{i=1}^N I(\sigma_i) \right\},$$

where  $I(x) := \frac{1+x}{2} \log \frac{1+x}{2} + \frac{1-x}{2} \log \frac{1-x}{2}$  is the binary entropy function. By choosing  $\sigma = t\mathbf{1}$  with  $t$  as defined in Lemma 1.2, we get the further lower bound

$$Z_N(\beta, B) \geq N \left\{ \frac{\beta t^2}{2} + Bt - I(t) \right\} + \frac{\beta t^2}{2} \sum_{i=1}^N (R_i - 1) =: M_N(\beta, B), \quad (1.13)$$

where  $t$  is as in Lemma 1.1. It follows from [4, Theorem 2.1] that  $\log Z_N(\beta, B) - M_N(\beta, B) = o(N)$ , as soon as  $\|A_N\|_F^2 + \sum_{i=1}^N (R_i - 1)^2 = o(N)$ . Our next result gives a bound to the approximation error of the partition function  $Z_N(\beta, B)$  by  $M_N(\beta, B)$ , which we henceforth refer to as the Mean-Field prediction in this paper.

**Theorem 1.5.** *Let  $A_N$  satisfy (1.4) and (1.5).*

(a) *If  $(\beta, B) \in \Theta_1$  then we have*

$$\log Z_N(\beta, B) - M_N(\beta, B) \lesssim \|A_N\|_F^2 + t^2 \sum_{i=1}^N (R_i - 1)^2.$$

(b) *If  $(\beta, B) \in \Theta_2$ , then the same conclusion as in part (a) holds under the extra assumption that  $A_N$  satisfies (1.7).*

(c) *If  $(\beta, B) \in \Theta_3$ , then under the extra assumption that  $A_N$  satisfies (1.7) we have*

$$\log Z_N(\beta, B) - M_N(\beta, B) \lesssim \|A_N\|_F^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1)^2 \right]^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1) \right]^2 + \log N.$$

**Remark 1.3.** *To see how the error bounds of the above theorem compare to existing error bounds for the Mean-Field prediction formula in the literature, let us take the example where  $A_N$  is the (scaled) adjacency matrix of a  $d_N$ -regular graph  $G_N$ . In this case, the above theorem gives the error bound  $O(N/d_N)$  for the Mean-Field prediction formula. This immediately improves the bounds from [4, Theorem 1.1] —  $o(N)$ , [25, Theorem 1.1] —  $O(N/d_N^{1/3})$ , [20, Example 3] —  $O(N/d_N^{1/2-o(1)})$  under strong expander type conditions not needed here) and [2, Corollary 2.9 and Example 2.10] —  $O(N/\sqrt{d_N})$ .*

For our next result, define an i.i.d. probability measure  $\mathbb{Q}$  on  $\{-1, 1\}^N$  by setting

$$\mathbb{Q}(\sigma_1, \dots, \sigma_N) := (\exp(-\beta t - B) + \exp(\beta t + B))^N \exp \left( (\beta t + B) \sum_{i=1}^N \tau_i \right). \quad (1.14)$$

Our next theorem shows that if an event is unlikely under the above i.i.d. measure/ the Curie Weiss model (depending on  $(\beta, B)$ ), then it is also unlikely under an Ising model on an approximately regular graph with large degree.

**Theorem 1.6.** *Let  $A_N$  satisfy (1.4) and (1.5). Also, let  $\mathcal{E}_N \subset \{-1, 1\}^N$  be arbitrary.*

(a) *If  $(\beta, B) \in \Theta_1$ , then we have*

$$\log \mathbb{P}(\mathcal{E}_N) \lesssim \log \mathbb{Q}(\mathcal{E}_N) + \|A_N\|_F^2 + t^2 \sum_{i=1}^N (R_i - 1)^2.$$

(b) *If  $(\beta, B) \in \Theta_2$ , then under the further assumption (1.7) we have*

$$\log \mathbb{P}(\mathcal{E}_N) \lesssim \log \mathbb{P}^{CW}(\mathcal{E}_N) + \|A_N\|_F^2 + \sum_{i=1}^N (R_i - 1)^2.$$

(c) If  $(\beta, B) \in \Theta_3$ , then under the further assumption (1.7) we have

$$\log \mathbb{P}(\mathcal{E}_N) \lesssim \log \mathbb{P}^{CW}(\mathcal{E}_N) + \|A_N\|_F^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1)^2 \right]^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1) \right]^2 + \log N.$$

As an application of the above theorem, we immediately get the following exponential concentration for  $\bar{\sigma}$ .

**Corollary 1.1.** *Suppose  $A_N$  satisfies (1.4), (1.5), and*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N (R_i - 1)^2 = 0, \quad \lim_{N \rightarrow \infty} \frac{1}{N} \|A_N\|_F^2 = 0.$$

- If  $(\beta, B) \in \Theta_1$ , then for every  $\delta > 0$  we have

$$\limsup_{n \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(|\bar{\sigma} - t| > \delta) < 0.$$

Same conclusion holds for  $(\beta, B) \in \Theta_3$ , under the extra assumption that  $A_N$  satisfies (1.7).

- If  $(\beta, B) \in \Theta_2$ , then under the extra assumption that  $A_N$  satisfies (1.7), for every  $\delta > 0$  we have

$$\limsup_{n \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(|\bar{\sigma} - M(\sigma)| > \delta) < 0.$$

Similar concentration results can be obtained for other higher order polynomials of  $\sigma$ , as studied in [1, 12, 23] and the references therein. However, these papers focus exclusively on the high temperature regime  $\Theta_{11}$ , whereas our result applies to all temperatures.

## 1.2. Examples

As mentioned before, the most common example of a coupling matrix  $A_N$  in model (1.1) is the scaled adjacency matrix  $\frac{1}{d_N} G_N$ , where  $G_N$  is the adjacency matrix of a simple labelled graph on  $N$  vertices with degree vector  $(d_1, \dots, d_N)$ , and  $\bar{d}_N := \frac{1}{N} \sum_{i=1}^N d_i$  is the average degree of  $G_N$ . The scaling discussed in the above definition ensures that the resulting Ising model has non-trivial phase transition properties (see e.g., [4, 31]). Below we consider some specific examples of graphs to illustrate our theorems.

- (a) **Regular graphs:** Let  $G_N$  be a  $d_N$  regular graph. Then  $\|A_N\|_F^2 = \frac{N}{d_N}$  and  $R_i = 1$ , and so applying Theorems 1.1, 1.2, 1.3 and 1.4 give

$$\begin{aligned} d_{KS}(\sqrt{N}(\bar{\sigma} - t), Z_\tau) &\lesssim \sqrt{\frac{N \log N}{d_N^3}} + \frac{1}{\sqrt{N}} && \text{if } (\beta, B) \in \Theta_{11}, \\ d_{KS}(\sqrt{N}(\bar{\sigma} - t), Z_\tau) &\lesssim \frac{\sqrt{N}}{d_N} && \text{if } (\beta, B) \in \Theta_{12}, \\ d_{KS}(\sqrt{N}(\bar{\sigma} - M(\sigma)), Z_\tau) &\lesssim \frac{\sqrt{N}}{d_N} && \text{if } (\beta, B) \in \Theta_2 \text{ and } G_N \text{ satisfies (1.7),} \\ d_{KS}(N^{1/4}\bar{\sigma}, W) &\lesssim \left(\frac{\sqrt{N} \log N}{d_N}\right)^{3/2} + \frac{\sqrt{N}}{d_N} + \frac{\log N}{\sqrt{N}} && \text{if } (\beta, B) \in \Theta_3 \text{ and } G_N \text{ satisfies (1.7).} \end{aligned}$$

In particular this means that  $\bar{\sigma}$  has the same fluctuations as that of the Curie Weiss model as soon as

$$\begin{aligned} d_N &\gg (N \log N)^{1/3} && \text{if } (\beta, B) \in \Theta_{11}, \\ d_N &\gg \sqrt{N} && \text{if } (\beta, B) \in \Theta_{12}, \\ d_N &\gg \sqrt{N} && \text{if } (\beta, B) \in \Theta_2 \text{ and (1.7) holds,} \\ d_N &\gg \sqrt{N} \log N && \text{if } (\beta, B) \in \Theta_3 \text{ and (1.7) holds.} \end{aligned} \tag{1.15}$$

Further, as already shown in Example 1.3, the requirement  $d_N \gg \sqrt{N}$  is sharp in the regimes  $\Theta_{12} \cup \Theta_2 \cup \Theta_3$ . Note that for the particular case of the Curie-Weiss model at criticality we get the convergence rate of  $N^{-1/2} \log N$ , which matches the rate obtained in [14] up to the log factor. In fact, it is easy to modify our argument in the special case of the Curie-Weiss model to get rid of the log factor. We observe that for the case of random  $d_N$  regular graphs, condition (1.7) holds with high probability, as  $\lambda_2(G_N) = O_p(\sqrt{d_N}) \ll d_N$  (see [11]), and so our results apply directly to random regular graphs if  $d_N$  satisfies (1.15). We stress that our results apply to regular bipartite graphs as well, and does not need the graph to be an expander as in [10].

- (b) **Erdős-Rényi graphs:** Suppose  $G_N \sim \mathcal{G}(N, p_N)$  be the symmetric Erdős Rényi random graph with  $0 < p_N \leq 1$ . Define  $A_N(i, j) := \frac{1}{(N-1)p_N} G_N(i, j)$ , and note that

$$\max_{1 \leq i \leq N} |R_i - 1| = O_p\left(\sqrt{\frac{\log N}{Np_N}}\right), \quad \left|\sum_{i=1}^N (R_i - 1)\right| = O_p\left(\frac{1}{\sqrt{p_N}}\right), \quad \sum_{i=1}^N (R_i - 1)^2 = O_p\left(\frac{1}{p_N}\right). \quad (1.16)$$

Since  $\lambda_2(G_N) = O_p(\sqrt{Np_N}) \ll Np_N$  ([22, Theorem 1.1]), (1.7) holds as well. Then our theorems conclude universal fluctuations for  $\bar{\sigma}$  as soon as

$$\begin{aligned} p_N &\gg (\log N)^{1/3} N^{-2/3} && \text{if } (\beta, B) \in \Theta_{11}, \\ p_N &\gg N^{-1/2} && \text{if } (\beta, B) \in \Theta_{12} \cup \Theta_2, \\ p_N &\gg (\log N)^4 N^{-1/2} && \text{if } (\beta, B) \in \Theta_3, \end{aligned} \quad (1.17)$$

both in the quenched and annealed setting. We note that our results also apply to the asymmetric Erdős-Rényi random graph  $\tilde{\mathcal{G}}(N, p_N)$ , under the same regime of  $p_N$  as in the symmetric case. This is because an Ising model on the asymmetric Erdős-Rényi graph is equivalent to an Ising model with the symmetric coupling matrix  $A_N(i, j) = \frac{\tilde{G}_N(i, j) + \tilde{G}_N(j, i)}{2(N-1)p_N}$ . The asymmetric case was studied recently in [26], where the authors derive fluctuations as soon as  $Np_N \rightarrow \infty$ , but only in the sub parameter regime  $\Theta_{11} \cup \Theta_3$ . The authors conjecture similar results for the symmetric case, which we are able to verify partially in this paper. Moreover, our theorems apply simultaneously to both the symmetric and the asymmetric cases with explicit convergence rates, thereby yielding fluctuations even in the hitherto unexplored regimes  $\Theta_{12}$  and  $\Theta_2$ .

- (c) **Balanced stochastic block model:** Suppose  $G_N$  is a stochastic block model with 2 communities of size  $N/2$  (assume  $N$  is even). Let the probability of an edge within the community be  $a_N$ , and across communities be  $b_N$ . This is the well known stochastic block model, which has received considerable attention in Probability, Statistics and Machine Learning (see [18, 27, 30] and references within). If we take  $A_N = \frac{2}{N(a_N + b_N)} G_N$ , universal asymptotics hold for  $\bar{\sigma}$  as soon as  $p_N := \frac{a_N + b_N}{2}$  satisfies (1.17), and  $\liminf_{N \rightarrow \infty} \frac{b_N}{a_N} > 0$  (needed to ensure (1.7)). Similar results hold when the number of communities is larger than 2.
- (d) **Sparse regular graphons:** Suppose that  $W$  be a symmetric measurable function from  $[0, 1]^2$  to  $[0, 1]$ , such that  $\int_{[0, 1]} W(x, y) dy = a > 0$  for all  $x \in [0, 1]$ , and  $\lambda_2(W) < a$ , where  $\{\lambda_i(W)\}_{i \geq 1}$  are the countable set of ordered eigenvalues. Also let  $(U_1, \dots, U_N) \stackrel{i.i.d.}{\sim} U(0, 1)$ . For  $\gamma \in (0, 1]$ , let

$$\{G_N(i, j)\}_{1 \leq i < j \leq N} \stackrel{i.i.d.}{\sim} \text{Bern}\left(\frac{W(U_i, U_j)}{N^\gamma}\right).$$

Such random graph models have been studied in the literature under the name  $W$  random graphons (c.f. [6, 7, 8, 9, 28]). In this case for the choice  $A_N = \frac{1}{Np_N} G_N$  with  $p_N = aN^{-\gamma}$ , universal fluctuation holds as soon as  $\gamma > 1/2$ . Indeed, writing

$$R_i - 1 = R_i - \frac{\sum_{j=1}^N W(U_i, U_j)}{aN} + \frac{\sum_{j=1}^N W(U_i, U_j)}{aN} - 1,$$

it is easy to check that (1.16) holds. Also with  $W_N$  denoting the  $N \times N$  matrix with  $W_N(i, j) = W(U_i, U_j)$ , using [3, Corollary 3.3] we have  $\|A_N - (aN)^{-1} W_N\|_{\text{op}} = O_p\left(\frac{\sqrt{N}}{Np_N}\right)$ . Since  $W_N$  converges in cut norm to  $W$ , it follows using [28, Section 11.6] we have

$$\lim_{N \rightarrow \infty} \lambda_2(A_N) = a^{-1} \lim_{N \rightarrow \infty} \frac{\lambda_2(W_N)}{N} = a^{-1} \lambda_2(W) < 1$$

and so  $A_N$  satisfies (1.7). By our results, universal fluctuations hold for  $\bar{\sigma}$  as soon as (1.16) holds.

- (e) **Block spin Ising model:** Suppose that  $N$  is even, and

$$\begin{aligned} A_N(i, j) &= a_N \text{ if } i, j \leq N/2 \text{ or } i, j > N/2, \\ &= b_N \text{ if } i \leq N/2, j > N/2, \text{ or } i > N/2, j \leq N/2. \end{aligned}$$

$A_N$  can be thought of as the expectation of a stochastic block Model with 2 communities. In the particular case  $a_N = \frac{\beta}{N}, b_N = \frac{\alpha}{N}$ , this model has been studied in [5, 29] under the name block spin

Ising model. Again in this case universal asymptotics holds for  $\bar{\sigma}$  as soon as  $d_N := \frac{N(a_N + b_N)}{2}$  satisfies (1.15), and  $\liminf_{N \rightarrow \infty} \frac{b_N}{a_N} > 0$ . This in particular matches the results obtained from [29, Theorems 1.2, 1.4] which studies the sub parameter regime  $\Theta_{11} \cup \Theta_3$ . Our results apply to the whole parameter regime of  $(\beta, B)$  and a wide regime of scalings of  $(a_N, b_N)$ , providing explicit convergence rates. Similar extension holds when the matrix  $A_N$  has more than 2 groups as well.

- (f) **Wigner matrices:** To demonstrate that our techniques apply to examples well beyond scaled adjacency matrices, let  $A_N$  be a Wigner matrix with its entries  $\{A_N(i, j), 1 \leq i < j \leq N\}$  i.i.d. from a distribution  $F$  scaled by  $N\mu$ , where  $F$  is a distribution on non-negative reals with finite exponential moment and mean  $\mu > 0$ . In this case we have

$$\max_{1 \leq i \leq N} |R_i - 1| = O_p\left(\sqrt{\frac{\log N}{N}}\right), \quad \left|\sum_{i=1}^N (R_i - 1)\right| = O_p(1), \quad \sum_{i=1}^N (R_i - 1)^2 = O_p(1).$$

Also [3, Corollary 3.5] shows that  $\|A_N - \frac{1}{N}\mathbf{1}\mathbf{1}^\top\|_{\text{op}} = N^{-1/2}$ , and so (1.7) holds. Thus our theorems apply giving universal fluctuations for  $\bar{\sigma}$ .

## 2. Proof of main results

We first state a lemma which will be needed in all parameter regimes.

**Lemma 2.1.** *Suppose  $\sigma \sim (1.1)$  for some  $A_N$  satisfying (1.4), and  $\beta > 0, B \in \mathbb{R}$ .*

- (a) *Setting  $m_i(\sigma) := \sum_{j=1}^N A_N(i, j)\sigma_j$  we have*

$$\mathbb{E}\left[\sum_{i=1}^N (\sigma_i - \tanh(\beta m_i(\sigma) + B)) \tanh(\beta m_i(\sigma) + B)\right]^2 \lesssim N.$$

- (b) *For any  $\mathbf{c} = (c_1, \dots, c_n) \in \mathbb{R}^n$  we have*

$$\log \mathbb{P}\left(\left|\sum_{i=1}^N c_i (\sigma_i - \tanh(\beta m_i(\sigma) + B))\right| > t\right) \lesssim -\frac{t^2}{\|\mathbf{c}\|_2^2}.$$

Here, part (a) follows by invoking [13, Lemma 3.2] and (b) can be obtained by making minor adjustments in the proof of [32, Lemma 1].

We now present an exponential moment control lemma in all parameter regimes, which is one of the main estimates of this paper, and is itself new and possibly of independent interest. The proof of this is deferred to Section 3.

**Lemma 2.2.** *Suppose  $\sigma \sim (1.1)$ , with  $A_N$  satisfying (1.4) and (1.5).*

- (a) *If  $(\beta, B) \in \Theta_1$ , then there exists a fixed positive number  $\delta > 0$  such that*

$$\log \mathbb{E}\left[\exp\left(\frac{\delta}{2} \sum_{i=1}^N (m_i(\sigma) - t)^2\right)\right] \lesssim \|A_N\|_F^2 + t^2 \sum_{i=1}^N (R_i - 1)^2. \quad (2.1)$$

- (b) *If  $(\beta, B) \in \Theta_2$ , then the conclusion of part (a) holds under the additional assumption that  $A_N$  satisfies (1.7).*

- (c) *If  $(\beta, B) \in \Theta_3$ , then under the additional assumption that  $A_N$  satisfies (1.7) there exists a fixed positive number  $\delta > 0$  such that*

$$\log \mathbb{E}\left[\exp\left(\frac{\delta}{2} \sum_{i=1}^N (m_i(\sigma) - \bar{\mathbf{m}}(\sigma))^2\right)\right] \lesssim \|A_N\|_F^2 + \frac{1}{N} \left[\sum_{i=1}^N (R_i - 1)^2\right]^2 + \frac{1}{N} \left[\sum_{i=1}^N (R_i - 1)\right]^2 + \log N, \quad (2.2)$$

where  $\bar{\mathbf{m}}(\sigma) := N^{-1} \sum_{i=1}^N m_i(\sigma)$ .



## 2.1. Proof of Theorem 1.1 and Theorem 1.2

Without loss of generality we may assume that the RHS of (1.6) and (1.8) are bounded by 1, because otherwise the bound is trivial. With  $\sigma$  an observation from the Ising model (1.1), form an exchangeable pair  $(\sigma, \sigma')$  as follows: Let  $I$  denote a randomly sampled index from  $\{1, 2, \dots, N\}$ . Given  $I = i$ , replace  $\sigma_i$  with an independent  $\pm 1$  valued random variable  $\sigma'_i$  with mean  $\tanh(\beta m_i(\sigma) + B) = \mathbb{E}[\sigma_i | (\sigma_j, j \neq i)]$ , and let  $\sigma' := (\sigma_1, \dots, \sigma_{i-1}, \sigma'_i, \sigma_{i+1}, \dots, \sigma_N)$ . Then  $(\sigma, \sigma')$  is an exchangeable pair. Extend the definition of  $M(\sigma)$  to all parameter regimes by setting  $M(\sigma) := t$  if  $(\beta, B) \in \Theta_1 \cup \Theta_3$ . Then, with  $T_N := \sqrt{N}(\bar{\sigma} - M(\sigma))$  and  $T'_N := \sqrt{N}(\bar{\sigma}' - M(\sigma'))$ , the pair  $(T_N, T'_N)$  is exchangeable as well. A direct computation gives

$$\mathbb{E}[T_N - T'_N | T_N] = \frac{1}{N^{3/2}} \sum_{i=1}^N \mathbb{E}[\sigma_i - \tanh(\beta m_i(\sigma) + B) | T_N] - \sqrt{N} \mathbb{E}[M(\sigma) - M(\sigma') | T_N], \quad (2.3)$$

where the second term in the RHS above can be expanded as

$$\begin{aligned} \sum_{i=1}^N \tanh(\beta m_i(\sigma) + B) &= N \tanh(\beta M(\sigma) + B) + \beta(1 - t^2) \sum_{i=1}^N (m_i(\sigma) - M(\sigma)) + \sum_{i=1}^N \xi_i (m_i - M(\sigma))^2 \\ &= NM(\sigma) + \beta(1 - t^2) \sum_{i=1}^N (\sigma_i - M(\sigma)) + \beta(1 - t^2) M(\sigma) \sum_{i=1}^N (R_i - 1) \\ &\quad + \beta(1 - t^2) \sum_{i=1}^N (R_i - 1) (\sigma_i - M(\sigma)) + \sum_{i=1}^N \xi_i (m_i(\sigma) - M(\sigma))^2 \end{aligned} \quad (2.4)$$

for random variables  $(\xi_i)_{1 \leq i \leq N}$  satisfying  $\max_{1 \leq i \leq N} |\xi_i| \lesssim 1$ , where the second line uses the identity  $M(\sigma) = \tanh(\beta M(\sigma) + B)$ . Setting  $h_i = \beta(1 - t^2)(R_i - 1)$  and plugging (2.4) into (2.3) we get

$$\begin{aligned} \mathbb{E}[T_N - T'_N | T_N] &= \underbrace{\frac{T_N}{N} (1 - \beta(1 - t^2))}_{g(T_N)} - \underbrace{\frac{1}{N\sqrt{N}} \sum_{i=1}^N \mathbb{E}[\xi_i (m_i(\sigma) - M(\sigma))^2 | T_N]}_{H_1(T_N)} - \underbrace{\frac{1}{N\sqrt{N}} \mathbb{E} \left[ \sum_{i=1}^N h_i (\sigma_i - M(\sigma)) | T_N \right]}_{H_2(T_N)} \\ &\quad - \underbrace{\sqrt{N} \mathbb{E}[M(\sigma) - M(\sigma') | T_N] - N^{-3/2} \beta(1 - t^2) M(\sigma) \sum_{i=1}^N (R_i - 1)}_{H_3(T_N)}. \end{aligned} \quad (2.5)$$

Set  $G(x) := \int_0^x g(y) dy = (1 - \beta(1 - t^2))x^2/2N$ ,  $c_0 := N/(1 - t^2)$  and  $c_1 := (2\pi\tau)^{-1/2}$ , and note the existence of positive constants  $c_2$  and  $c_3$  free of  $N$  such that assumptions (H1), (H2) and (H3) from [14, Page 2] are all satisfied. By [14, Theorem 1.2], we then have

$$\begin{aligned} d_{KS}(T_N, Z_\tau) &\lesssim \mathbb{E} \left| 1 - \frac{N}{2(1 - t^2)} \mathbb{E}[(T_N - T'_N)^2 | T_N] \right| \\ &\quad + \frac{N}{2(1 - t^2)} c_1 \mathbb{E}[|T_N - T'_N|^3] + \frac{N c_1 c_2}{1 - t^2} c_2 \mathbb{E} \left[ \sum_{a=1}^3 |H_a(T_N)| \right] + \frac{\mathbb{E}|T_N| + 1}{\sqrt{N}}. \end{aligned} \quad (2.6)$$

We will now estimate each term in the RHS above. First observe that

$$N \mathbb{E}[|T_N - T'_N|^3] \lesssim \frac{N}{N\sqrt{N}} \mathbb{E}[|\sigma_I - \sigma'_I|^3] + N\sqrt{N} \mathbb{E}[|M(\sigma) - M(\sigma')|^3] \lesssim \frac{1}{\sqrt{N}} \quad (2.7)$$

where the control on the second term for  $(\beta, B) \in \Theta_2$  follows on using part (b) of Theorem 1.6 with  $\mathcal{E}_N := \{\sum_{i=1}^N \sigma_i \in \{-1, 0, 1\}\}$  along with part (c) of Proposition 5.1 to note that

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(M(\sigma) \neq M(\sigma')) \leq \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(\mathcal{E}_N) < 0. \quad (2.8)$$

Proceeding to control  $\mathbb{E}|H_1(T_N)|$  we have

$$N\sqrt{N}|H_1(T_N)| = \left| \sum_{i=1}^N \mathbb{E}(\xi_i (m_i(\sigma) - M(\sigma))^2 | T_N) \right| \lesssim \mathbb{E} \left( \sum_{i=1}^N (m_i(\sigma) - M(\sigma))^2 | T_N \right), \quad (2.9)$$

and so

$$N\sqrt{N}\mathbb{E}|H_1(T_N)| \lesssim \mathbb{E} \sum_{i=1}^N \left( m_i(\boldsymbol{\sigma}) - M(\boldsymbol{\sigma}) \right)^2 \leq \eta_N \quad (2.10)$$

using Lemma 2.2, with  $\eta_N := \|A_N\|_F^2 + t^2 \sum_{i=1}^N (R_i - 1)^2$  denoting the RHS of (2.1). Next, we have

$$\begin{aligned} N\sqrt{N}|H_2(T_N)| &\leq \left| \mathbb{E} \left( \sum_{i=1}^N h_i \left( \sigma_i - \tanh(\beta m_i(\boldsymbol{\sigma}) + B) \right) \middle| T_N \right) \right| \\ &\quad + \left| \mathbb{E} \left( \sum_{i=1}^N h_i \left( \tanh(\beta m_i(\boldsymbol{\sigma}) + B) - \tanh(\beta M(\boldsymbol{\sigma}) + B) \right) \middle| T_N \right) \right| \end{aligned} \quad (2.11)$$

and so

$$\begin{aligned} N\sqrt{N}\mathbb{E}|H_2(T_N)| &\lesssim \sqrt{\sum_{i=1}^n h_i^2} + \sqrt{\sum_{i=1}^N h_i^2} \sqrt{\mathbb{E} \sum_{i=1}^N \left( m_i(\boldsymbol{\sigma}) - M(\boldsymbol{\sigma}) \right)^2} \\ &\lesssim \sqrt{\sum_{i=1}^N (R_i - 1)^2 (1 + \sqrt{\eta_N})} \lesssim \eta_N + \sum_{i=1}^N (R_i - 1)^2, \end{aligned} \quad (2.12)$$

where the penultimate line uses part (b) of Lemma 2.1, and the last line again uses (2.10). Also observe that,

$$N\sqrt{N}|H_3(T_N)| \lesssim N^2 \mathbb{E} \left( |M(\boldsymbol{\sigma}) - M(\boldsymbol{\sigma}')| T_N \right) + t \left| \sum_{i=1}^N (R_i - 1) \right|, \quad (2.13)$$

where the first term has an expectation which is exponentially small in  $N$  using (2.8). Finally we have

$$\left| \mathbb{E} \left[ 1 - \frac{N}{2(1-t^2)} (T_N - T'_N)^2 \middle| T_N \right] \right| \lesssim \mathbb{E} \left| \mathbb{E} \left[ 1 - \frac{(\sigma_I - \sigma'_I)^2}{2(1-t^2)} \middle| T_N \right] \right| + N^2 \mathbb{E}[|M(\boldsymbol{\sigma}) - M(\boldsymbol{\sigma}')|].$$

The second term on the RHS above is exponentially small, by (2.8). For the first term on the RHS, note that

$$\mathbb{E} \left[ 1 - (\sigma_I - \sigma'_I)^2 / 2(1-t^2) \middle| \boldsymbol{\sigma} \right] \lesssim N^{-1} \left| \sum_{i=1}^N (\sigma_i \tanh(\beta m_i(\boldsymbol{\sigma}) + B) - t^2) \right|.$$

As a result we have

$$\begin{aligned} &\mathbb{E} \left| \mathbb{E} \left[ 1 - (\sigma_I - \sigma'_I)^2 / 2(1-t^2) \middle| T_N \right] \right| \\ &\lesssim \mathbb{E} \left| \sum_{i=1}^N (\sigma_i - \tanh(\beta m_i(\boldsymbol{\sigma}) + B)) \tanh(\beta m_i(\boldsymbol{\sigma}) + B) \right| + \frac{1}{N} \sum_{i=1}^N \mathbb{E} (m_i(\boldsymbol{\sigma}) - M(\boldsymbol{\sigma}))^2 \\ &\lesssim \frac{1}{\sqrt{N}} + \frac{1}{N} \sum_{i=1}^N \mathbb{E} (m_i(\boldsymbol{\sigma}) - M(\boldsymbol{\sigma}))^2 \leq \frac{\eta_N}{N} + \frac{1}{\sqrt{N}}, \end{aligned} \quad (2.14)$$

where we have used (2.10), and part (a) of Lemma 2.1. We now claim that

$$\mathbb{E} T_N^2 \lesssim 1. \quad (2.15)$$

Given this claim, combining the estimates from (2.6), (2.7), (2.10), (2.12), (2.13), and (2.14) we get

$$d_{KS}(T_N, Z_\tau) \lesssim \frac{1}{\sqrt{N}} + \frac{\eta_N}{\sqrt{N}} + \frac{1}{\sqrt{N}} \sum_{i=1}^N (R_i - 1)^2 + \frac{t}{\sqrt{N}} \left| \sum_{i=1}^N (R_i - 1) \right|,$$

from which the desired bound follows on noting that  $\eta_N \gtrsim \|A_N\|_F^2 \gtrsim 1$ .

It thus suffices to prove (2.15). To this effect, using (2.5) we get:

$$\left| \mathbb{E}[T_N - T'_N | T_N] - \frac{T_N}{N}(1 - \beta(1 - t^2)) \right| \lesssim \sum_{a=1}^3 |H_a(T_N)|.$$

By multiplying both sides of the above display by  $|T_N|$  and taking expectation gives

$$\mathbb{E}[T_N^2] \lesssim \left| \mathbb{E}[N(T_N - T'_N)T_N] \right| + \mathbb{E} \left[ N|T_N| \left( \sum_{a=1}^3 |H_a(T_N)| \right) \right],$$

where we have used the fact that  $\beta(1 - t^2) \neq 1$ . By the exchangeability of  $T_N$  and  $T'_N$  we have

$$\mathbb{E}[N(T_N - T'_N)T_N] = \mathbb{E}[N(T'_N - T_N)T'_N] = \frac{1}{2} \mathbb{E}[N(T_N - T'_N)^2] \lesssim 1.$$

Also, from (2.9), (2.11) and (2.13) we have

$$N \sum_{a=1}^3 \mathbb{E}[H_a(T_N)]^2 \lesssim \frac{\eta_N + \sum_{i=1}^N (R_i - 1)^2}{\sqrt{N}} \lesssim 1,$$

where the last bound uses the fact that the RHS of (1.6) and (1.8) are bounded. Using Chebyshev's inequality then gives

$$\mathbb{E}[T_N^2] \lesssim 1 + \sqrt{\mathbb{E}[T_N^2]} \sqrt{\sum_{a=1}^3 \mathbb{E}(NH_a(T_N))^2} \lesssim 1 + \sqrt{\mathbb{E}[T_N^2]}$$

which implies  $\mathbb{E}[T_N^2] \lesssim 1$ . This verifies (2.15), and hence completes the proof of the theorem.

## 2.2. Proof of Theorem 1.4

For proving Theorem 1.4 we need the following lemma, whose proof we defer to section 4.

**Lemma 2.3.** *Assume that  $\sigma$  is an observation from (1.1) with  $(\beta, B) \in \Theta_{11}$ , and  $A_N$  satisfies (1.11). Setting  $\alpha_N = \max_{1 \leq i \leq N} \sum_{j=1}^N A_N(i, j)^2$  as in Theorem 1.4, the following conclusions hold:*

(a)  $\log \mathbb{P}(\max_{1 \leq i \leq N} |m_i(\sigma)| \geq \lambda \sqrt{\alpha_N \log N}) \lesssim -\lambda^2$ , for any  $\lambda > 0$ .

(b)  $\mathbb{E} \left[ \sum_{i=1}^N (R_i - 1) \sigma_i \right]^2 \lesssim \left( \sum_{i=1}^N (R_i - 1)^2 \right) \left[ 1 + \|A_N\|_F^2 \alpha_N^2 (\log N)^2 \right]$ .

*Proof.* Without loss of generality we can assume that the RHS of (1.12) is bounded as before. As in the proof of the previous theorems, it suffices to bound the RHS of (2.6), but with  $t = M(\sigma) = 0$  which implies  $H_3(T_N) = 0$ . To begin, use (2.14) to get

$$\mathbb{E} \left| \mathbb{E} \left[ 1 - \frac{N}{2} (T_N - T'_N)^2 | T_N \right] \right| \lesssim \frac{\|A_N\|_F^2}{N} + \frac{1}{\sqrt{N}}, \quad (2.16)$$

using (2.1), which allows us to replace  $\eta_N$  in the previous proof by  $\|A_N\|_F^2$ . Proceeding to bound  $\mathbb{E}|H_1(T_N)|$ , use the first equality of (2.9) along with Cauchy-Schwarz inequality to note that

$$N\sqrt{N} \mathbb{E}|H_1(T_N)| \lesssim \mathbb{E} \max_{1 \leq i \leq N} |m_i(\sigma)| \sum_{i=1}^N m_i(\sigma)^2 \leq \sqrt{\mathbb{E} \max_{1 \leq i \leq N} m_i(\sigma)^2} \sqrt{\mathbb{E} \left( \sum_{i=1}^N m_i(\sigma)^2 \right)^2} \lesssim \|A_N\|_F^2 \sqrt{\alpha_N \log N}, \quad (2.17)$$

where the last inequality uses part (a) of Lemma 2.3. Finally, for  $\mathbb{E}|H_2(T_N)|$  we have

$$N\sqrt{N} \mathbb{E}|H_2(T_N)| \leq \mathbb{E} \left| \sum_{i=1}^N (R_i - 1) \sigma_i \right| \lesssim \sqrt{\left( \sum_{i=1}^N (R_i - 1)^2 \right) \left[ 1 + \|A_N\|_F^2 \alpha_N \log N \right]},$$

where we use part (b) of Lemma 2.3. Plugging in the above bounds in (2.6), we have

$$d_{KS}(T_N, Z_\tau) \lesssim \frac{1 + \mathbb{E}(T_N^2)}{\sqrt{N}} + \frac{\|A_N\|_F^2 \sqrt{\alpha_N \log N}}{\sqrt{N}} + \left[1 + \|A_N\| \alpha_N \log N\right] \sqrt{\frac{(\sum_{i=1}^N (R_i - 1)^2)}{N}},$$

from which the claimed bound follows immediately, if we can verify (2.15). But the proof of this is the same as in the previous theorem, and so we are done.  $\square$

### 2.3. Proof of Theorem 1.3

For proving Theorem 1.3 we need the following lemma, whose proof we again defer to section 4.

**Lemma 2.4.** *Suppose  $\sigma \sim (1.1)$  with  $(\beta, B) \in \Theta_3$ , such that  $A_N$  satisfies (1.9) and (1.7). Suppose further that the RHS of (1.10) is bounded. Then the following conclusions hold:*

- (a)  $\log \mathbb{P} \left( \max_{1 \leq i \leq N} |m_i(\sigma) - \bar{m}(\sigma)| \geq \lambda \sqrt{\alpha_N (\log N)^3} + \log N \max_{1 \leq i \leq N} |R_i - 1| \right) \lesssim -\lambda^2$ , for any  $\lambda > 0$ .
- (b)  $\mathbb{E} \left[ \sum_{i=1}^N (R_i - 1) \sigma_i \right]^2 \lesssim \left( \sum_{i=1}^N (R_i - 1)^2 + N^{-1/2} \left[ \sum_{i=1}^N (R_i - 1) \right]^2 \right) (\log N)^4$ .
- (c)  $N^{3/2} \mathbb{E}(\bar{\sigma}^6) \lesssim 1$ .

*Proof.* With  $(\sigma, \sigma')$  the usual exchangeable pair, setting  $T_N := N^{1/4} \bar{\sigma}$  and  $T'_N := N^{1/4} \bar{\sigma}'$  we have

$$\mathbb{E}[T_N - T'_N | \sigma] = N^{-3/4} (\bar{\sigma} - \tanh(\bar{\sigma})) + N^{-3/4} (\tanh(\bar{\sigma}) - \tanh(\bar{m}(\sigma))) + N^{-7/4} \sum_{i=1}^N (\tanh(m_i(\sigma)) - \tanh(\bar{m}(\sigma))).$$

Using Taylor's expansion, this gives

$$\begin{aligned} & |\mathbb{E}[T_N - T'_N | \sigma] - N^{-3/4} (\bar{\sigma} - \tanh(\bar{\sigma}))| \\ & \lesssim N^{-3/4} |\bar{\sigma} - \bar{m}(\sigma)| + N^{-7/4} |\bar{\sigma}| \sum_{i=1}^N (m_i(\sigma) - \bar{m}(\sigma))^2 + N^{-7/4} \left| \sum_{i=1}^N (m_i(\sigma) - \bar{m}(\sigma))^3 \right|, \end{aligned} \quad (2.18)$$

and so we have  $\mathbb{E}[T_N - T'_N | T_N] = g(T_N) + H(T_N)$  where  $g(x) = N^{-3/2} x^3 / 3$ , and  $H(T_N)$  satisfies

$$\begin{aligned} \mathbb{E}[|H(T_N)|] & \lesssim N^{-2} \mathbb{E}[|T_N|^5] + N^{-3/4} \mathbb{E}[|\bar{\sigma} - \bar{m}(\sigma)|] + N^{-2} \mathbb{E} \left[ |T_N| \sum_{i=1}^N (m_i(\sigma) - \bar{m}(\sigma))^2 \right] \\ & \quad + N^{-7/4} \mathbb{E} \left[ \left| \sum_{i=1}^N (m_i(\sigma) - \bar{m}(\sigma))^3 \right| \right]. \end{aligned}$$

Invoking [14, Theorem 1.2] with  $G(x) := \int_0^x g(t) dt = N^{-3/2} x^4 / 12$  we have

$$d_{KS}(T_N, W) \lesssim \mathbb{E} \left| 1 - \frac{N^{3/2}}{2} \mathbb{E}[(T_N - T'_N)^2 | T_N] \right| + N^{3/2} \mathbb{E}[|H(T_N)|] + N^{-3/4} \mathbb{E}|T_N|^3. \quad (2.19)$$

By part (c) of Lemma 2.4 we have  $\mathbb{E}[|T_N|^5] \lesssim 1$ . Set  $\delta_N := \sum_{i=1}^N (R_i - 1)^2 + N^{-1/2} \left[ \sum_{i=1}^N (R_i - 1) \right]^2$ , and use part (b) of Lemma 2.4 and the Cauchy-Schwarz inequality to get

$$\mathbb{E}[|\bar{\sigma} - \bar{m}(\sigma)|] \lesssim \sqrt{\mathbb{E}(\bar{\sigma} - \bar{m}(\sigma))^2} \lesssim N^{-1} (\log N)^2 \sqrt{\delta_N}.$$

Similarly, by the Cauchy-Schwarz inequality and part (c) of Lemmas 2.2 along with part (a) of Lemma 2.4, we get:

$$\mathbb{E} \left[ |T_N| \sum_{i=1}^N (m_i(\sigma) - \bar{m}(\sigma))^2 \right] \leq \sqrt{\mathbb{E}(T_N)^2} \sqrt{\mathbb{E} \left( \sum_{i=1}^n (m_i(\sigma) - \bar{m}(\sigma))^2 \right)^2} \lesssim \varepsilon_N,$$

$$\mathbb{E} \left[ \sum_{i=1}^N |m_i(\boldsymbol{\sigma}) - \bar{m}(\boldsymbol{\sigma})|^3 \right] \leq \sqrt{\mathbb{E} \max_{1 \leq i \leq N} (m_i(\boldsymbol{\sigma}) - \bar{m}(\boldsymbol{\sigma}))^2} \sqrt{\mathbb{E} \left( \sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{m}(\boldsymbol{\sigma}))^2 \right)^2} \lesssim r_N \varepsilon_N,$$

where  $\varepsilon_N$  is as in the statement of Theorem 1.3. Combining the above observations, we get

$$N^{3/2} \mathbb{E}[|H(T_N)|] \lesssim N^{-1/2} + N^{-1/4} (\log N)^2 \sqrt{\delta_N} + N^{-1/4} r_N \varepsilon_N. \quad (2.20)$$

Finally, we have

$$\begin{aligned} \mathbb{E} \left| 1 - \frac{N^{3/2}}{2} \mathbb{E} [(T_N - T'_N)^2 | T_N] \right| &\lesssim \frac{1}{N} \mathbb{E} \left| \sum_{i=1}^N \sigma_i \tanh m_i(\boldsymbol{\sigma}) \right| \\ &\lesssim \frac{1}{N} \mathbb{E} \left| \sum_{i=1}^N (\sigma_i - \tanh m_i(\boldsymbol{\sigma})) \tanh m_i(\boldsymbol{\sigma}) \right| + \frac{1}{N} \mathbb{E} \sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{m}(\boldsymbol{\sigma}))^2 + \mathbb{E} \bar{\boldsymbol{\sigma}}^2 \\ &\lesssim \frac{1}{\sqrt{N}} + \frac{\varepsilon_N}{N} + \frac{1}{\sqrt{N}} \end{aligned} \quad (2.21)$$

where the last inequality follows from part (a) of Lemma 2.1, part (c) of Lemma 2.2, and part (c) of Lemma 2.4. Combing (2.20) and (2.21) along with (2.19) gives

$$d_{KS}(T_N, W) \lesssim \frac{1}{\sqrt{N}} + \frac{\sqrt{\delta_N} (\log N)^2}{N^{1/4}} + \frac{r_N \varepsilon_N}{N^{1/4}},$$

as desired.  $\square$

### 3. Proofs of Theorems 1.5, 1.6 and Lemma 2.2

We will need the following proposition which expresses the Curie-Weiss model as a mixture of i.i.d. random variables, first shown in [31, Lemma 3].

**Proposition 3.1.** *Let  $\boldsymbol{\sigma} \sim \mathbb{P}^{CW}$  be an observation from the Curie-Weiss model (1.3). Given  $\boldsymbol{\sigma}$ , let  $W_N$  be a Gaussian random variable with mean  $\bar{\boldsymbol{\sigma}}_N$  and variance  $(N\beta)^{-1}$ . Then the following conclusions hold:*

(a) *Given  $W_N$ , the random variables  $(\sigma_1, \sigma_2, \dots, \sigma_N)$  are i.i.d. with mean  $\widetilde{W}_N := \tanh(\beta W_N + B)$ .*

(b) *The marginal density of  $W_N$  is proportional to  $\exp(-Nf(w))$ , where  $f(w) = \frac{\beta w^2}{2} - \log \cosh(\beta w + B)$ .*

We state two more lemmas necessary for proving the results of this section, the proofs of which we defer to Section 5. The first lemma is a version of the Hanson-Wright inequality, which controls exponential moment of quadratic forms of binary random variables.

**Lemma 3.1.** *Suppose  $X_1, X_2, \dots, X_N, N \geq 1$  are i.i.d.  $\pm 1$  valued random variables such that  $\mathbb{E}[X_1] = \mu$ . Set  $s_\mu := 2\mu/(\log(1+\mu) - \log(1-\mu))$  with  $s_0$  being 1. Also assume that  $D_N$  is a  $N \times N$  symmetric matrix such that  $s_\mu \limsup_{N \rightarrow \infty} \lambda_1(D_N) < 1$ . Then, given any vector  $\mathbf{c}^\top := (c_1, c_2, \dots, c_N)$ , we get:*

$$\log \left\{ \mathbb{E}_0 \left[ \exp \left( \frac{1}{2} \sum_{i,j=1}^N D_N(i,j) \widetilde{X}_i \widetilde{X}_j + \sum_{i=1}^N c_i \widetilde{X}_i \right) \right] \right\} \lesssim \text{Tr}^+(D_N) + \|D_N\|_F^2 + \sum_{i=1}^N c_i^2$$

where  $\widetilde{X}_i = X_i - \mu$  for  $1 \leq i \leq N$ , and  $\text{Tr}^+(D_N) = \sum_{i=1}^N \max(D_N(i,i), 0)$ .

The second lemma gives a quantitative estimate which allows us to neglect the region where  $\widetilde{W}_N$  is not close to  $t$ .

**Lemma 3.2.** *Suppose (1.4), (1.5) and (1.7) holds, and further assume that  $\|A_N\|_F^2 = o(N)$ ,  $\sum_{i=1}^N (R_i - 1) = o(N)$ . Also, let  $V_N$  be any random variable such that  $V_N \leq cN$  for some fixed  $c > 0$ , and  $\varepsilon > 0$  be fixed. Setting  $\mathcal{A}_N := A_N - \mathbf{1}\mathbf{1}^\top/N$ , for any  $(\beta, B) \in \Theta_2 \cup \Theta_3$  there exists  $\delta = \delta(\varepsilon, c, \beta) > 0$  such that,*

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{E}^{CW} \left[ \exp \left( \delta V_N + \frac{\beta}{2} \boldsymbol{\sigma}^\top \mathcal{A}_N \boldsymbol{\sigma} \right) \mathbb{1}(|\widetilde{W}_N - M(\boldsymbol{\sigma})| \geq \varepsilon) \right] < 0. \quad (3.1)$$

*Proof of Theorem 1.5.* (a) To begin, note that

$$\frac{\beta}{2} \boldsymbol{\sigma}^\top A_N \boldsymbol{\sigma} + B \sum_{i=1}^N \sigma_i = \frac{\beta}{2} (\boldsymbol{\sigma} - t)^\top A_N (\boldsymbol{\sigma} - t) + \sum_{i=1}^N (\beta t R_i + B) \sigma_i - (\beta t^2/2) \mathbf{1}^\top A_N \mathbf{1},$$

which on using (1.13) gives

$$\frac{Z_N(\beta, B)}{\exp(M_N(\beta, B))} = \mathbb{E}^{\mathbb{Q}} \exp \left( \frac{\beta}{2} \sum_{i,j=1}^N (\sigma_i - t) A_N(i, j) (\sigma_j - t) + \beta t \sum_{i=1}^N (R_i - 1) (\sigma_i - t) \right) \quad (3.2)$$

where  $\mathbb{Q}$  is as defined in (1.14). In this case with  $D_N = \beta A_N$  we have

$$s_t \limsup_{N \rightarrow \infty} \lambda_1(D_N) = \beta s_t \limsup_{N \rightarrow \infty} \lambda_1(\mathcal{A}_N) \leq \beta s_t = \frac{\beta t}{\beta t + B} < 1,$$

for  $(\beta, B) \in \Theta_{12}$ . If  $(\beta, B) \in \Theta_{11}$ , then we have  $t = 0$ , and  $s_0 = 1$ , and so with  $D_N = \beta A_N$  as before, we have  $s_t \limsup_{N \rightarrow \infty} \lambda_1(D_N) = \beta < 1$ . Thus in both cases Lemma 3.1 is applicable with  $D_N = \beta A_N, c_i = \beta t(R_i - 1)$ , which using (3.2) gives

$$\mathbb{E}^{\mathbb{Q}} \exp \left( \frac{\beta}{2} \sum_{i,j=1}^N (\sigma_i - t) A_N(i, j) (\sigma_j - t) + \beta t \sum_{i=1}^N (R_i - 1) (\sigma_i - t) \right) \lesssim \|A_N\|_F^2 + t^2 \sum_{i=1}^N (R_i - 1)^2. \quad (3.3)$$

The conclusion of part (a) follows from this combined with (3.2).

(b) Define

$$Y_N := (\boldsymbol{\sigma} - \widetilde{W}_N)^\top A_N (\boldsymbol{\sigma} - \widetilde{W}_N) + 2\widetilde{W}_N \sum_{i=1}^N (R_i - 1) (\sigma_i - \widetilde{W}_N) + (\widetilde{W}_N^2 - t^2) \sum_{i=1}^N (R_i - 1), \quad (3.4)$$

and note that  $\boldsymbol{\sigma}^\top A_N \boldsymbol{\sigma} = Y_N + t^2 \sum_{i=1}^N (R_i - 1)$ . Using this, with  $J_{N,\epsilon} := \{|t| - \epsilon \leq |\widetilde{W}_N| \leq |t| + \epsilon\}$  for some  $\epsilon > 0$ , by a similar calculation as in part (a) we have:

$$\begin{aligned} \frac{Z_N(\beta, B)}{Z_N^{CW}(\beta, B)} &= \mathbb{E}^{CW} \exp \left( \frac{\beta}{2} \boldsymbol{\sigma}^\top A_N \boldsymbol{\sigma} \right) \\ &= \mathbb{E}^{CW} \left[ \exp \left( \frac{\beta}{2} \boldsymbol{\sigma}^\top A_N \boldsymbol{\sigma} \right) \mathbb{1}(J_{N,\epsilon}^c) \right] + \exp \left( \frac{\beta}{2} t^2 \sum_{i=1}^N (R_i - 1) \right) \mathbb{E}^{CW} \left( e^{\frac{\beta}{2} Y_N} \mathbb{1}(J_{N,\epsilon}) \right). \end{aligned} \quad (3.5)$$

The first term in the right hand side of (3.5) is  $o(1)$  by invoking Lemma 3.2 with  $\delta = 0$ . For the second term, by Proposition 3.1, the inner (conditional) expectation is taken with respect to i.i.d.  $\pm 1$  valued random variables with mean  $\widetilde{W}_N$ . In this regime  $\beta s_t = \beta t/\beta t = 1$ . But since  $\limsup_{N \rightarrow \infty} \lambda_1(\mathcal{A}_N) < 1$  by (1.7), on the set  $J_{N,\epsilon}$  we have

$$\limsup_{N \rightarrow \infty} s_{\widetilde{W}_N} \lambda_1(\beta \mathcal{A}_N) \leq \limsup_{N \rightarrow \infty} \sup_{\mu \in J_{N,\epsilon}} s_\mu \lambda_1(\beta \mathcal{A}_N) < 1$$

for  $\epsilon$  small enough. Therefore, Lemma 3.1 is applicable with  $D_N = \beta \mathcal{A}_N$  and  $c_i = 2\widetilde{W}_N(R_i - 1)$  to give

$$\log \mathbb{E}^{CW} \left( e^{\frac{\beta}{2} Y_N} \mathbb{1}(J_{N,\epsilon}) \mid \widetilde{W}_N \right) \leq C \left\{ \|A_N\|_F^2 + \sum_{i=1}^N (R_i - 1)^2 \right\} + \frac{\beta}{2} \left| (\widetilde{W}_N^2 - t^2) \sum_{i=1}^N (R_i - 1) \right|$$

for some  $C < \infty$ , which on taking another expectation gives

$$\begin{aligned} \log \mathbb{E}^{CW} \left( e^{\frac{\beta}{2} Y_N} \mathbb{1}(J_{N,\epsilon}) \right) &\leq C \left\{ \|A_N\|_F^2 + \sum_{i=1}^N (R_i - 1)^2 \right\} + \log \mathbb{E} e^{\frac{\beta}{2} \left| (\widetilde{W}_N^2 - t^2) \sum_{i=1}^N (R_i - 1) \right|} \\ &\lesssim \|A_N\|_F^2 + \sum_{i=1}^N (R_i - 1)^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1) \right]^2, \end{aligned} \quad (3.6)$$

where the last step uses part (b) of Proposition 5.1. This along with (3.5) gives

$$\log Z_N(\beta, B) - \log Z_N^{CW}(\beta, B) - \frac{\beta t^2}{2} \sum_{i=1}^N (R_i - 1) \lesssim \|A_N\|_F^2 + \sum_{i=1}^N (R_i - 1)^2,$$

from which the desired conclusion follows by another application of part (a) of Proposition 5.1 to note that  $\log Z_N^{CW}(\beta, B) - N \left[ \frac{\beta t}{2} + Bt - I(t) \right] \lesssim 1$ .

- (c) In this regime we have  $t = 0$ , and so  $s_t = s_0 = 1$ , and  $\beta s_0 = 1$ . As in the proof of part (b), the first term in the RHS of (3.5) is  $o(1)$  invoking Lemma 3.2 with  $\delta = 0$ . For handling the second term, invoking (1.7) gives

$$\limsup_{N \rightarrow \infty} s_{\widetilde{W}_N} \lambda_1(\mathcal{A}_N) \leq \limsup_{N \rightarrow \infty} \sup_{\mu \in J_{N,\varepsilon}} s_\mu \lambda_1(\mathcal{A}_N) < 1$$

for  $\varepsilon$  small enough. Also Lemma 3.1 with  $D_N = \mathcal{A}_N$  and  $c_i = 2\widetilde{W}_N(R_i - 1)$  gives

$$\log \mathbb{E}^{CW} \left( e^{\frac{\beta}{2} Y_N} \mathbb{1}(J_{N,\varepsilon}) | \widetilde{W}_N \right) \leq C \left\{ \|A_N\|_F^2 + \widetilde{W}_N^2 \sum_{i=1}^N (R_i - 1)^2 \right\} + \frac{\beta}{2} \widetilde{W}_N^2 \sum_{i=1}^N (R_i - 1)$$

for some  $C < \infty$  free of  $N$ . This, on taking another expectation along with (3.5) gives

$$\begin{aligned} \log \mathbb{E}^{CW} \left( e^{\frac{\beta}{2} Y_N} \mathbb{1}(J_{N,\varepsilon}) \right) &\leq C \|A_N\|_F^2 + \log \mathbb{E} \exp \left( C \widetilde{W}_N^2 \sum_{i=1}^N (R_i - 1)^2 + \frac{\beta}{2} \widetilde{W}_N^2 \sum_{i=1}^N (R_i - 1) \right) \\ &\lesssim \|A_N\|_F^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1) + \sum_{i=1}^N (R_i - 1)^2 \right]^2, \end{aligned} \quad (3.7)$$

where the last bound uses part (b) of Proposition 5.1. Combining (3.5) and (3.7) gives

$$\log Z_N(\beta, B) - \log Z_N^{CW}(\beta, B) \lesssim \|A_N\|_F^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1) \right]^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1)^2 \right]^2.$$

We incur an additional log factor in the final answer because  $\log Z_N^{CW}(\beta, B) - N \left[ \frac{\beta t}{2} + Bt - I(t) \right] \lesssim \log N$  by part (a) of Proposition 5.1.

□

*Proof of Theorem 1.6.* (a) Using a similar calculation as in (3.5), we get:

$$\mathbb{P}(\boldsymbol{\sigma} \in \mathcal{E}_N) = c(N) \mathbb{E}^{\mathbb{Q}} \left[ \exp \left( \frac{\beta}{2} \sum_{i,j=1}^N (\sigma_i - t) A_N(\sigma_j - t) + \beta t \sum_{i=1}^N (R_i - 1)(\sigma_i - t) \right) \mathbb{1}(\boldsymbol{\sigma} \in \mathcal{E}_N) \right] \quad (3.8)$$

where the deterministic sequence  $c(N)$  satisfies

$$c(N) = \frac{\exp(\beta t^2 (\mathbf{1}^\top A_N \mathbf{1} - N)) (\exp(\beta t + B) + \exp(-\beta t - B))^N}{Z_N(\beta, B) \exp((\beta t^2 / 2) \mathbf{1}^\top A_N \mathbf{1})} \leq 1,$$

on invoking the Mean-Field lower bound (1.13). Next, by using Hölder's inequality with exponent  $p$  (to be chosen later), the left hand side of (3.8) can be bounded above by,

$$\left\{ \mathbb{E}^{\mathbb{Q}} \exp \left( \frac{\beta(1+p)}{2} \sum_{i,j=1}^N (\sigma_i - t) A_N(\sigma_j - t) + \beta t(1+p) \sum_{i=1}^N (R_i - 1)(\sigma_i - t) \right) \right\}^{\frac{1}{1+p}} (\mathbb{Q}(\mathcal{E}_N))^{\frac{p}{1+p}}. \quad (3.9)$$

Using arguments similar to the derivation of (3.3) shows that for  $p$  small enough we have

$$\log \mathbb{E}^{\mathbb{Q}} \left[ \exp \left( \frac{\beta(1+p)}{2} \sum_{i,j} (\sigma_i - t) A_N(\sigma_j - t) + \beta t(1+p) \sum_{i=1}^N (R_i - 1)(\sigma_i - t) \right) \right] \lesssim \|A_N\|_F^2 + t^2 \sum_{i=1}^N (R_i - 1)^2.$$

Combining this along with (3.8) and (3.9) gives the desired conclusion follows.

- (b) With  $Y_N$  as in (3.4), using a similar calculation as in the derivation of (3.5) we can bound  $P(\boldsymbol{\sigma} \in \mathcal{E}_N)$  by

$$\begin{aligned} & \frac{Z_N^{CW}(\beta, B)}{Z_N(\beta, B)} \mathbb{E}^{CW} e^{\frac{\beta}{2} \boldsymbol{\sigma}^\top A_N \boldsymbol{\sigma}} \mathbb{1}(\boldsymbol{\sigma} \in \mathcal{E}_N) \\ & \leq \frac{Z_N^{CW}(\beta, B)}{Z_N(\beta, B)} \left[ \mathbb{E}^{CW} e^{\frac{\beta}{2} \boldsymbol{\sigma}^\top A_N \boldsymbol{\sigma}} \mathbb{1}(\boldsymbol{\sigma} \in J_{N,\varepsilon}^c) + e^{\frac{\beta t^2}{2} \sum_{i=1}^N (R_i - 1)} \mathbb{E}^{CW} e^{\frac{\beta}{2} Y_N} \mathbb{1}(\boldsymbol{\sigma} \in \mathcal{E}_N) \mathbb{1}(J_{N,\varepsilon}) \right]. \end{aligned} \quad (3.10)$$

For controlling the ratio of partition functions in the RHS of (3.10), use the Mean-Field approximation (1.13) to get a lower bound for  $\log Z_N(\beta, B)$ , whereas part (a) of Proposition 5.1 gives  $\log Z_N^{CW}(\beta, B) - N \left[ \frac{\beta t}{2} + Bt - I(t) \right] \lesssim 1$ . Combining these two observations, we get:

$$\log Z_N^{CW}(\beta, B) - \log Z_N(\beta, B) + \frac{\beta t^2}{2} \sum_{i=1}^N (R_i - 1) \lesssim 1. \quad (3.11)$$

Also, the first term inside the parenthesis in the RHS of (3.10) is exponentially small in  $N$ , by invoking Lemma 3.2 with  $\delta = 0$ . Proceeding to control the second term in the RHS of (3.10) we have

$$\mathbb{E}^{CW} e^{\frac{\beta}{2} Y_N} \mathbb{1}(\boldsymbol{\sigma} \in \mathcal{E}_N) \mathbb{1}(J_{N,\varepsilon}) \leq \left[ \mathbb{E}^{CW} e^{\frac{\beta(1+p)}{2} Y_N} \mathbb{1}(J_{N,\varepsilon}) \right]^{\frac{1}{1+p}} \left[ \mathbb{P}^{CW}(\boldsymbol{\sigma} \in \mathcal{E}_N) \right]^{\frac{p}{1+p}}. \quad (3.12)$$

where the last step uses Holder's inequality with  $p > 0$  to be chosen later. For controlling the first term inside the bracket in the RHS of (3.12), repeating the same argument as in the derivation of (3.6) with  $p > 0$  small enough, yields:

$$\log \mathbb{E}^{CW} (e^{\frac{\beta(1+p)}{2} Y_N} \mathbb{1}(J_{N,\varepsilon})) \lesssim \|A_N\|_F^2 + \sum_{i=1}^N (R_i - 1)^2. \quad (3.13)$$

Combining (3.10), (3.11), (3.12) and (3.13), the desired conclusion follows.

- (c) All steps of part (b) above go through verbatim, except the RHS of (3.11) gets replaced by  $\log N$  (by part (a) of Proposition 5.1), and (3.13) is replaced by (c.f. (3.7))

$$\log \mathbb{E}^{CW} e^{\frac{\beta(1+p)}{2} Y_N} \mathbb{1}(J_{N,\varepsilon}) \lesssim \|A_N\|_F^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1)^2 \right]^2 + \frac{1}{N} \left[ \sum_{i=1}^N \sum_{i=1}^N (R_i - 1) \right]^2. \quad (3.14)$$

Combining this with (3.10) and (3.14) gives the desired conclusion.  $\square$

*Proof of Lemma 2.2.* (a) Invoking Theorem 1.6 and changing  $\delta$  if necessary, it suffices to show the desired conclusion under  $\mathbb{Q}$ , where  $\mathbb{Q}$  is the i.i.d. measure defined in (1.14). A direct calculation shows that  $m_i(\boldsymbol{\sigma}) - t$  equals  $\sum_{j=1}^N A_N(i, j)(\sigma_j - t) + t(R_i - 1)$ , and so

$$\begin{aligned} \sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - t)^2 & \leq 2 \sum_{i=1}^N \left[ \sum_{j=1}^N A_N(i, j)(\sigma_j - t) \right]^2 + 2t^2 \sum_{i=1}^N (R_i - 1)^2 \\ & = 2 \sum_{i=1}^N \sum_{j=1}^N A_N^2(i, j)(\sigma_i - t)(\sigma_j - t) + 2t^2 \sum_{i=1}^N (R_i - 1)^2. \end{aligned} \quad (3.15)$$

It therefore suffices to control the exponential moment of the first term in the RHS of (3.15). Since  $\limsup_{N \rightarrow \infty} \lambda_1(A_N^2) \leq 1$ , for any  $\delta \in (0, 1/2)$ , using Lemma 3.1 with  $D_N = \delta A_N^2$  and  $c_i = 0$  we have

$$\log \mathbb{E}^{\mathbb{Q}} \exp(\delta(\boldsymbol{\sigma} - t)^\top A_N^2(\boldsymbol{\sigma} - t)) \lesssim \|A_N^2\|_F^2 = \sum_{i=1}^N \lambda_i^4 \lesssim \sum_{i=1}^N \lambda_i^2 = \|A_N\|_F^2.$$

This gives the desired conclusion.



- (c) By invoking Theorem 1.6, it suffices to show the desired conclusion under the Curie-Weiss model. Start by noting that  $\bar{m}(\boldsymbol{\sigma}) = \frac{1}{N} \sum_{i=1}^N R_i \sigma_i$ , and so

$$m_i(\boldsymbol{\sigma}) - \bar{m}(\boldsymbol{\sigma}) = \sum_{j=1}^N A_N(i, j) (\sigma_j - \widetilde{W}_N) + \frac{1}{N} \sum_{i=1}^N R_i (\sigma_i - \widetilde{W}_N) + \widetilde{W}_N (R_i - \bar{R}).$$

This shows that  $\sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{m}(\boldsymbol{\sigma}))^2$  is bounded by

$$\begin{aligned} & 3 \sum_{i=1}^N \left[ \sum_{j=1}^N A_N(i, j) (\sigma_j - \widetilde{W}_N) \right]^2 + \frac{3}{N} \left[ \sum_{i=1}^N R_i (\sigma_i - \widetilde{W}_N) \right]^2 + 3 \widetilde{W}_N^2 \sum_{i=1}^N (R_i - \bar{R})^2 \\ & \leq 3 \sum_{i, j=1}^N \left( A_N^2(i, j) + \frac{3}{N} R_i R_j \right) (\sigma_i - \widetilde{W}_N) (\sigma_j - \widetilde{W}_N) + 3 \widetilde{W}_N^2 \sum_{i=1}^N (R_i - 1)^2. \end{aligned} \quad (3.16)$$

Conditioning on  $\widetilde{W}_N$ , we now control the exponential moment of the first term in the RHS of the above display under the Curie-Weiss model. By Proposition 3.1, under the Curie Weiss model, given  $\widetilde{W}_N$ , the random vector  $(\sigma_1, \dots, \sigma_N)$  are i.i.d. with mean  $\widetilde{W}_N$ . Since

$$\limsup_{N \rightarrow \infty} \lambda_1 \left( A_N^2 + \frac{3}{N} \mathbf{R} \mathbf{R}^\top \right) \leq \limsup_{N \rightarrow \infty} \lambda_1(A_N^2) + \limsup_{N \rightarrow \infty} \frac{3}{N} \lambda_1(\mathbf{R} \mathbf{R}^\top) \lesssim 1,$$

on invoking Lemma 3.1 with  $c_i = 0$ ,  $D_N = \delta \left( A_N^2 + \frac{3}{N} \mathbf{R} \mathbf{R}^\top \right)$  for  $\delta$  small enough, and noting that  $\| \frac{1}{N} \mathbf{R} \mathbf{R}^\top \|_F^2 = \frac{1}{N^2} (\sum_{i=1}^N R_i^2)^2 \lesssim 1$  by (1.5), we get

$$\log \mathbb{E}^{CW} e^{\delta \sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{m}(\boldsymbol{\sigma}))^2} - 3 \delta \widetilde{W}_N^2 \sum_{i=1}^N (R_i - 1)^2 \lesssim \|A_N^2\|_F^2 + \text{tr}(A_N^2) \lesssim \|A_N\|_F^2,$$

from which the desired conclusion follows on noting that

$$\log \mathbb{E}^{CW} e^{3 \delta \widetilde{W}_N^2 \sum_{i=1}^N (R_i - 1)^2} \lesssim \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1)^2 \right]^2,$$

which follows from part (b) of Proposition 5.1.

- (b) To begin note that

$$\sum_{i=1}^N (m_i - M(\boldsymbol{\sigma}))^2 \lesssim \sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{m}(\boldsymbol{\sigma}))^2 + \frac{1}{N} \left[ \sum_{i=1}^N R_i (\sigma_i - \widetilde{W}_N) \right]^2 + (\widetilde{W}_N - M(\boldsymbol{\sigma}))^2 \left| \sum_{i=1}^N (R_i - 1) \right|. \quad (3.17)$$

By Hölder's inequality, it suffices to bound the exponential moments of the first three terms of the above display at some  $\delta > 0$ . Exponential moment of the third term in the RHS of (3.17) is bounded by part (b) of Proposition 5.1, as  $\sum_{i=1}^N |R_i - 1| = o(N)$ . Proceeding to bound the sum of the first two terms, use (3.16) to get

$$\sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{m}(\boldsymbol{\sigma}))^2 + \frac{1}{N} \left[ \sum_{i=1}^N R_i (\sigma_i - \widetilde{W}_N) \right]^2 \leq \sum_{i, j=1}^N \left( A_N^2(i, j) + \frac{4}{N} R_i R_j \right) (\sigma_i - \widetilde{W}_N) (\sigma_j - \widetilde{W}_N) + 3 \sum_{i=1}^N (R_i - 1)^2,$$

and so it suffices to bound

$$\log \mathbb{E}^{CW} \exp \left( \delta \sum_{i, j=1}^N \left( A_N^2(i, j) + \frac{4}{N} R_i R_j \right) (\sigma_i - \widetilde{W}_N) (\sigma_j - \widetilde{W}_N) \right)$$

for  $\delta$  small enough. But this follows on invoking Lemma 3.1 with  $D_N = \delta \left( A_N^2 + \frac{4}{N} \mathbf{R} \mathbf{R}^\top \right)$  and  $c_i = 0$  to get

$$\log \mathbb{E}^{CW} \exp \left( \delta \sum_{i, j=1}^N \left( A_N^2(i, j) + \frac{4}{N} R_i R_j \right) (\sigma_i - \widetilde{W}_N) (\sigma_j - \widetilde{W}_N) \right) \lesssim \|A_N^2\|_F^2 + \text{tr}(A_N^2) \lesssim \|A_N\|_F^2,$$

which completes the proof of part (b).  $\square$

#### 4. Proof of Lemmas 2.3 and 2.4

*Proof of Lemma 2.3.* (a) To begin, note that it suffices to prove the bound for  $\lambda$  large enough. To this effect, using part (b) of Lemma 2.1 we have the existence of a constant  $M$  free of  $N$ , such that for all  $\lambda > 0$  we have

$$\mathbb{P}\left(\left|m_i(\boldsymbol{\sigma}) - \sum_{j=1}^N A_N(i, j) \tanh(\beta m_j(\boldsymbol{\sigma}))\right| > \lambda \sqrt{\log N \sum_{j=1}^N A_N(i, j)^2}\right) \leq 2e^{-\frac{\lambda^2 \log N}{M}},$$

which on using a union bound with  $\alpha_N = \max_{1 \leq i \leq N} \sum_{j=1}^N A_N(i, j)^2$  (as in Theorem 1.4) gives

$$\mathbb{P}\left(\max_{1 \leq i \leq N} \left|m_i(\boldsymbol{\sigma}) - \sum_{j=1}^N A_N(i, j) \tanh(\beta m_j(\boldsymbol{\sigma}))\right| > \lambda \sqrt{\alpha_N \log N}\right) \leq 2Ne^{-\frac{\lambda^2 \log N}{M}}.$$

On the set  $\left\{\max_{1 \leq i \leq N} \left|m_i(\boldsymbol{\sigma}) - \sum_{j=1}^N A_N(i, j) \tanh(\beta m_j(\boldsymbol{\sigma}))\right| \leq \lambda \sqrt{\alpha_N \log N}\right\}$  using the bound  $|\tanh(x)| \leq |x|$  we have

$$\max_{1 \leq i \leq N} |m_i(\boldsymbol{\sigma})| \leq \sqrt{\alpha_N \log N} + \beta \max_{1 \leq i \leq N} R_i \max_{1 \leq i \leq N} |m_i(\boldsymbol{\sigma})|,$$

which on using (1.11) gives  $\max_{1 \leq i \leq N} |m_i(\boldsymbol{\sigma})| \lesssim \sqrt{\alpha_N \log N}$ . Thus there exists a constant  $c'$  such that

$$\mathbb{P}\left(\max_{1 \leq i \leq N} |m_i(\boldsymbol{\sigma})| > c' \lambda \sqrt{\alpha_N \log N}\right) \leq 2Ne^{-\frac{\lambda^2 \log N}{M}},$$

from which the desired conclusion follows for all  $\lambda$  large enough.

(b) More generally, we will show that for any vector  $\mathbf{c} \in \mathbb{R}^N$  we have

$$\mathbb{E}\left(\sum_{i=1}^N c_i \sigma_i\right)^2 \lesssim (\log N)^{3/2} \sum_{i=1}^N c_i^2. \quad (4.1)$$

To this effect, for every non-negative integer  $\ell$  set  $\mathbf{c}^{(\ell)} := \beta^\ell A_N^\ell \mathbf{c}$ , and  $x_\ell := \mathbb{E}[(\sum_i c_i^{(\ell)} \sigma_i)^2]$ , and note that  $\mathbf{c}^{(0)} = \mathbf{c}$ , and the LHS of (4.1) is just  $x_0$ . For any  $\ell \geq 0$  we can write

$$x_\ell = T_{1,\ell} + T_{2,\ell} + T_{3,\ell}, \quad (4.2)$$

where

$$T_{1,\ell} := \mathbb{E}\left[\left(\sum_{i=1}^N c_i^{(\ell)} (\sigma_i - \tanh(\beta m_i(\boldsymbol{\sigma})))\right)^2\right], \quad T_{2,\ell} := \mathbb{E}\left[\left(\sum_{i=1}^N c_i^{(\ell)} \tanh(\beta m_i(\boldsymbol{\sigma}))\right)^2\right]$$

$$T_{3,\ell} = 2\mathbb{E}\left[\left(\sum_{i \neq j} c_i^{(\ell)} c_j^{(\ell)} (\sigma_i - \tanh(\beta m_i(\boldsymbol{\sigma}))) \tanh(\beta m_j(\boldsymbol{\sigma}))\right)\right].$$

For controlling  $T_{3,\ell}$ , setting  $m_i^j(\boldsymbol{\sigma}) := \sum_{k=1, k \neq j}^N A_N(i, k) \sigma_k \sigma_j$  we have

$$|T_{3,\ell}| = 2 \left| \sum_{i \neq j} c_i^{(\ell)} c_j^{(\ell)} \mathbb{E}\left[(\sigma_i - \tanh(\beta m_i(\boldsymbol{\sigma}))) (\tanh(\beta m_j(\boldsymbol{\sigma})) - \tanh(\beta m_i^j(\boldsymbol{\sigma})))\right] \right|$$

$$\lesssim \sum_{i \neq j} |c_i^{(\ell)}| |c_j^{(\ell)}| A_N(i, j) \lesssim \|\mathbf{c}^{(\ell)}\|_2^2 \quad (4.3)$$

where we use  $\mathbb{E}[\sigma_i - \tanh(\beta m_i(\boldsymbol{\sigma})) | \sigma_j, j \neq i] = 0$  in the first line, and the bound  $|\tanh(\beta m_i(\boldsymbol{\sigma})) - \tanh(\beta m_i^j(\boldsymbol{\sigma}))| \lesssim A_N(i, j)$  in the second line.

Proceeding to bound  $T_{2,\ell}$ , use a Taylor's series expansion to get  $\tanh(\beta m_i(\boldsymbol{\sigma})) = \beta m_i(\boldsymbol{\sigma}) + \xi_i m_i(\boldsymbol{\sigma})^3$  for random variables  $\{\xi_i\}_{1 \leq i \leq N}$  uniformly bounded by 1 in absolute value. Consequently,

$$\begin{aligned} T_{2,\ell} - x_{\ell+1} &= \mathbb{E} \left[ \left( \sum_{i=1}^N c_i^{(\ell)} \{m_i(\boldsymbol{\sigma})\beta + \xi_i m_i(\boldsymbol{\sigma})^3\} \right)^2 \right] - \mathbb{E} \left[ \left( \sum_{i=1}^N c_i^{(\ell)} m_i(\boldsymbol{\sigma}) \right)^2 \right] \\ &\leq 2\sqrt{x_{\ell+1}} \|\mathbf{c}^{(\ell)}\|_2 \sqrt{\mathbb{E} \left[ \sum_i m_i(\boldsymbol{\sigma})^6 \right]} + \|\mathbf{c}^{(\ell)}\|_2^2 \mathbb{E} \left[ \sum_i m_i(\boldsymbol{\sigma})^6 \right]. \end{aligned} \quad (4.4)$$

Finally, using Cauchy-Schwarz inequality gives

$$\mathbb{E} \left( \sum_{i=1}^N m_i(\boldsymbol{\sigma})^6 \right) \leq \sqrt{\mathbb{E} \left( \sum_{i=1}^N m_i^2 \right)^2} \sqrt{\mathbb{E} \max_{1 \leq i \leq N} |m_i(\boldsymbol{\sigma})|^8} \leq C^2 \|A_N\|_F^2 \alpha_N^2 (\log N)^2 \quad (4.5)$$

for some  $C$  free of  $N$ , where the last inequality uses part (a) of this lemma and part (b) of Lemma 2.2. Noting that  $T_{1,\ell} \lesssim \|\mathbf{c}^{(\ell)}\|_2^2$  by part (b) of Lemma 2.1, combining (4.3), (4.4) and (4.5) along with (4.2) gives the existence of a constant  $D$  free of  $N, \ell$  such that

$$x_\ell \leq x_{\ell+1} + 2\sqrt{x_{\ell+1}} \|\mathbf{c}\|_2 \beta_N^\ell \delta_N + \|\mathbf{c}\|_2^2 \beta_N^{2\ell} \delta_N^2 + D \beta_N^{2\ell} \|\mathbf{c}\|_2^2, \quad (4.6)$$

where we have also used the bound  $\|\mathbf{c}^{(\ell)}\|_2 \leq \beta_N^\ell \|\mathbf{c}\|_2$  with  $\beta_N := \beta \|A_N\|_2$ , and we set  $\delta_N := \max(1, C \|A_N\|_F \alpha_N \log N)$ . Since  $\beta_N \rightarrow \beta < 1$ , for all  $N$  large we have  $\beta_N \leq \beta_0$  for some  $\beta_0 < 1$ . Given constants  $\beta_0 \in (0, 1), D > 0$ , there exists  $M$  large enough such that  $M > (\beta_0 \sqrt{M} + 1)^2 + D$ . With this  $M, \beta_0$  we claim that for all  $\ell$ , we have

$$x_\ell \leq M \|\mathbf{c}\|_2^2 \beta_0^{2\ell} \delta_N^2, \quad (4.7)$$

from which (4.1) is immediate on setting  $\ell = 0$ . For proving (4.7) we use backwards induction on  $\ell$ . Using Cauchy-Schwarz inequality gives

$$x_\ell \leq N \|\mathbf{c}^{(\ell)}\|_2^2 \leq N \beta_N^\ell \|\mathbf{c}\|_2^2,$$

and so (4.7) holds for all  $\ell$  large enough, as  $\beta_N < \beta_0$ . Assume that the result holds for  $x_{\ell+1}$  for some  $\ell$ , i.e.  $x_{\ell+1} \leq M \|\mathbf{c}\|_2^2 \beta_0^{2\ell+2} \delta_N^2$ . Using (4.6) gives

$$x_\ell \leq \|\mathbf{c}\|_2^2 \beta_0^{2\ell} \delta_N^2 \left( M \beta_0^2 + 2\sqrt{M} \beta_0 + 1 + D \right) \leq M \|\mathbf{c}\|_2^2 \beta_0^{2\ell} \delta_N^2,$$

where the last step uses the choice of  $M$ . This verifies the claim for  $\ell$ , and hence proves (4.7) by backward induction, for all  $\ell \geq 0$ .  $\square$

*Proof of Lemma 2.4.* (a) As in the proof of part (a) of Lemma 2.3, it suffices to prove the result for  $\lambda$  large. To this effect, define an  $N \times N$  matrix  $\tilde{A}_N$  by setting  $\tilde{A}_N(i, j) := A_N(i, j)/R_{\max}$  for  $i \neq j$  and  $\tilde{A}_N(i, i) := 1 - R_i/R_{\max}$  where  $R_{\max} = \max_{1 \leq i \leq N} R_i$ . Observe that  $\mathbf{1}^\top \tilde{A}_N = \mathbf{1}^\top$ , and so

$$\begin{aligned} & |(m_i(\boldsymbol{\sigma}) - \bar{m}(\boldsymbol{\sigma})) - \sum_{j=1}^N \tilde{A}_N(i, j) (m_j(\boldsymbol{\sigma}) - \bar{m}(\boldsymbol{\sigma}))| \\ &= |m_i(\boldsymbol{\sigma}) - \sum_{j=1}^N \tilde{A}_N(i, j) m_j(\boldsymbol{\sigma})| \\ &\leq |m_i - \sum_{j=1}^N A_N(i, j) m_j(\boldsymbol{\sigma})| + \sum_{j=1}^N |A_N(i, j) - \tilde{A}_N(i, j)| \\ &\lesssim |m_i - \sum_{j=1}^N A_N(i, j) \tanh(m_j(\boldsymbol{\sigma}))| + \max_{1 \leq i \leq N} |m_i(\boldsymbol{\sigma})|^3 + \max_{1 \leq i \leq N} |R_i - 1|. \end{aligned} \quad (4.8)$$

Using part (b) of Lemma 2.1, a union bound as in the proof of part (a) of Lemma 2.3 shows that for all  $\lambda > 0$  we have  $\mathbb{P}(E_N^c) \leq 2e^{-c\lambda^2}$  for some constant  $c > 0$  free of  $N$ , where

$$E_N := \left\{ \max_{1 \leq i \leq N} \left| m_i(\boldsymbol{\sigma}) - \sum_{j=1}^N A_N(i, j) \tanh(m_j(\boldsymbol{\sigma})) \right| \leq \lambda \sqrt{\alpha_N \log N} \right\} \quad (4.9)$$

for some constant  $c$  free of  $N$ , with  $\alpha_N = \max_{1 \leq i \leq N} \sum_{j=1}^N A_N(i, j)^2$  as in Theorem 1.4. Proceeding to bound the second term in the RHS of (4.8), note that

$$|m_i(\boldsymbol{\sigma})|^3 \lesssim |m_i(\boldsymbol{\sigma}) - \tanh(m_i(\boldsymbol{\sigma}))| \leq |m_i(\boldsymbol{\sigma}) - \sum_{j=1}^N A_N(i, j) \tanh(m_j(\boldsymbol{\sigma}))| + \max_{1 \leq i \leq N} |R_i - 1|. \quad (4.10)$$

Thus, combining (4.8) and (4.10), on the set  $E_N$  we have

$$\max_{1 \leq i \leq N} |(m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) - \sum_{j=1}^N \tilde{A}_N(i, j)(m_j(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))| \leq C \left[ \lambda \sqrt{\alpha_N \log N} + \max_{1 \leq i \leq N} |R_i - 1| \right] \quad (4.11)$$

for some  $C < \infty$  free of  $N$ . Now, for any integer  $\ell \geq 2$  we have

$$\begin{aligned} & |(m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) - \sum_{j=1}^N \tilde{A}_N^\ell(i, j)(m_j(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))| \\ & \leq |(m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) - \sum_{j=1}^N \tilde{A}_N^{\ell-1}(i, j)(m_j(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))| + \left| \sum_{j=1}^N \tilde{A}_N^{\ell-1}(i, j) \left\{ (m_j(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) - \sum_{k=1}^N \tilde{A}_N(j, k)(m_k(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) \right\} \right| \\ & \leq |(m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) - \sum_{j=1}^N \tilde{A}_N^{\ell-1}(i, j)(m_j(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))| + \max_{1 \leq j \leq N} \left| (m_j(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) - \sum_{k=1}^N \tilde{A}_N(j, k)(m_k(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) \right|, \end{aligned}$$

which, via a recursive argument gives

$$\begin{aligned} & \max_{1 \leq i \leq N} \left| (m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) - \sum_{j=1}^N \tilde{A}_N^\ell(i, j)(m_j(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) \right| \\ & \leq \ell \max_{1 \leq i \leq N} \left| (m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) - \sum_{j=1}^N \tilde{A}_N(i, j)(m_j(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) \right| \leq C\ell \left( \lambda \sqrt{\alpha_N \log N} + \max_{1 \leq i \leq N} |R_i - 1| \right), \quad (4.12) \end{aligned}$$

where the last line uses (4.11) on the set  $E_N$ . Using part (a) of Lemma 5.2, we note the existence of  $D$  free of  $N$  such that for the choice  $\ell = D \log N$  we have  $\max_{1 \leq i \leq N} A^\ell(i, i) \leq \frac{3}{N}$ . With this choice of  $\ell$ , we have

$$\begin{aligned} & \mathbb{P} \left( \max_{1 \leq i \leq N} |m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})| \geq 2C\ell \left[ \lambda \sqrt{\alpha_N \log N} + \max_{1 \leq i \leq N} |R_i - 1| \right], E_N \right) \\ & \leq \mathbb{P} \left( \max_{1 \leq i \leq N} \left| \sum_{j=1}^N \tilde{A}_N^\ell(i, j)(m_j(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})) \right| \geq C\ell \left[ \lambda \sqrt{\alpha_N \log N} + \max_{1 \leq i \leq N} |R_i - 1| \right] \right) \\ & \leq \mathbb{P} \left( \sum_{j=1}^N (m_j(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))^2 \geq \frac{C^2 \ell^2 N}{2} \left[ \lambda \sqrt{\alpha_N \log N} + \max_{1 \leq i \leq N} |R_i - 1| \right]^2 \right), \end{aligned}$$

where the last line uses Cauchy-Schwarz inequality. Fixing  $\delta$  small enough and using part (c) of Lemma 2.2, this gives

$$\begin{aligned} & \log \mathbb{P} \left( \max_{1 \leq i \leq N} |m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma})| \geq 2C\ell \left[ \lambda \sqrt{\alpha_N \log N} + \max_{1 \leq i \leq N} |R_i - 1| \right], E_N \right) \\ & \lesssim -N\alpha_N (\log N)^3 \lambda^2 - N(\log N)^2 \max_{1 \leq i \leq N} |R_i - 1|^2 + \log \mathbb{E} e^{\delta \sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))^2} \\ & \lesssim -N\alpha_N (\log N)^3 \lambda^2 - N(\log N)^2 \max_{1 \leq i \leq N} |R_i - 1|^2 + \|A_N\|_F^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1)^2 \right]^2 + \frac{1}{N} \left[ \sum_{i=1}^N (R_i - 1) \right]^2 + \log N, \end{aligned}$$

from which the desired conclusion follows for  $\lambda$  large enough on noting the inequality  $N\alpha_N \geq \|A_N\|_F^2 \gtrsim 1$ .  $\square$

In order to prove Lemma 2.4, parts (b) and (c), we need the following lemma whose proof we defer to the end of this lemma.

**Lemma 4.1.** *Assume that (1.4), (1.5), (1.9) holds, and the RHS of (1.10) is bounded. Then, setting  $\nu_N := \mathbb{E}_1^N[(N^{1/4}\bar{\sigma})^6]$  the following conclusions hold:*

$$\nu_N \lesssim \nu_N^{2/3} + \nu_N^{1/3} + \nu_N^{1/2} + \nu_N^{1/2} \sqrt{\frac{\mathbb{E} \left[ \sum_{i=1}^N (R_i - 1) \sigma_i \right]^2}{N^{1/2}}}, \quad (4.13)$$

$$\mathbb{E} \left[ \sum_{i=1}^N (R_i - 1) \sigma_i \right]^2 \lesssim (\log N)^4 \left( \sum_{i=1}^N (R_i - 1)^2 + N^{-1/2} \left[ \sum_{i=1}^N (R_i - 1) \right]^2 \right) \left( 1 + \mathbb{E}[(N^{1/4}\bar{\sigma})^2] \right). \quad (4.14)$$

*Proof of Lemma 2.4, parts (b) and (c).* Use (4.14) and the fact that the RHS of (1.10) is bounded to get

$$\mathbb{E} \left[ \sum_{i=1}^N (R_i - 1) \sigma_i \right]^2 \lesssim \sqrt{N} (1 + \mathbb{E}[(N^{1/4}\bar{\sigma})^2]) \lesssim \sqrt{N} (1 + \nu_N^{1/3}).$$

Along with (4.13), this gives  $\nu_N \lesssim \nu_N^{2/3} + \nu_N^{1/3} + \nu_N^{1/2} (1 + \nu_N^{1/3}) + 1$ , and so  $\nu_N$  must be bounded, thereby proving part (b). Now, part (c) is an immediate consequence of part (b) and (4.14).  $\square$

*Proof of Lemma 4.1.* (a) Proof of (4.13).

To begin, borrowing notation from the proof of Theorem 1.3 and using (2.18) gives the existence of  $C < \infty$  such that

$$\begin{aligned} & \left| \mathbb{E}[T_N - T'_N | \sigma] - N^{-3/2} T_N^3 / 3 \right| \\ & \leq \frac{2}{15} N^{-2} |T_N|^5 + C \left\{ N^{-3/4} |\bar{\sigma} - \bar{\mathbf{m}}(\sigma)| + N^{-2} |T_N| \sum_{i=1}^N (m_i(\sigma) - \bar{\mathbf{m}}(\sigma))^2 + N^{-7/4} \left| \sum_{i=1}^N (m_i(\sigma) - \bar{\mathbf{m}}(\sigma))^3 \right| \right\}. \end{aligned}$$

On multiplying both sides of the above inequality by  $|T_N|^3$  and taking expectation gives

$$\begin{aligned} \mathbb{E}[T_N^6] & \leq (2/5) N^{-1/2} \mathbb{E}|T_N|^8 + 3C \left\{ N^{3/4} \mathbb{E} [|T_N|^3 |\bar{\sigma} - \bar{\mathbf{m}}(\sigma)|] + N^{-1/2} \mathbb{E} \left[ |T_N|^4 \sum_{i=1}^N (m_i(\sigma) - \bar{\mathbf{m}}(\sigma))^2 \right] \right. \\ & \quad \left. + N^{-1/4} \mathbb{E} \left[ |T_N|^3 \left| \sum_{i=1}^N (m_i(\sigma) - \bar{\mathbf{m}}(\sigma))^3 \right| \right] \right\} + 3N^{3/2} |\mathbb{E}(T_N - T'_N) T_N^3|. \end{aligned} \quad (4.15)$$

We will now bound each of the terms in the RHS of (4.15). To begin, note that that  $|T_N - T'_N| \leq 2N^{-3/4}$  and  $\mathbb{E}[T_N] = \mathbb{E}[T'_N]$ . This, along with the fact that  $(T_N, T'_N)$  is an exchangeable pair gives

$$\begin{aligned} \mathbb{E}(T_N - T'_N) T_N^3 & = (1/2) \mathbb{E}(T_N - T'_N) T_N^3 - (1/2) \mathbb{E}(T_N - T'_N) (T'_N)^3 \\ & = (1/2) \mathbb{E} [(T_N - T'_N)^2 (T_N^2 + T_N T'_N + (T'_N)^2)] \\ & \leq 6N^{-3/2} \mathbb{E}[T_N^2] \leq 6N^{-3/2} \nu_N^{1/6}, \end{aligned} \quad (4.16)$$

where  $\nu_N = \mathbb{E}[(N^{1/4}\bar{\sigma})^6]$  as in the statement of the lemma. Also with  $\varepsilon_N, r_N$  as in the statement of Theorem 1.3, use part (c) of Lemma 2.2, and part (a) of Lemma 2.4 to get that for any positive integer  $p$ , we have

$$\mathbb{E} \left[ \sum_{i=1}^N (m_i(\sigma) - \bar{\mathbf{m}}(\sigma))^2 \right]^p \lesssim \varepsilon_N^p, \quad \mathbb{E} \max_{1 \leq i \leq N} |m_i(\sigma) - \bar{\mathbf{m}}(\sigma)|^p \lesssim r_N^p. \quad (4.17)$$

Finally, since the RHS of (1.10) is bounded, we have

$$\varepsilon_N \lesssim \sqrt{N}, \quad \varepsilon_N r_N \lesssim N^{1/4}, \quad \|\mathbf{c}\|_2^2 + N^{-1/2} \left[ \sum_{i=1}^N c_i \right]^2 \lesssim \sqrt{N}. \quad (4.18)$$

Armed with these estimates and proceeding to bound the second, third and fourth terms in (4.15), use Hölder's inequality to get

$$N^{3/4} \mathbb{E}[|T_N|^3 |\bar{\sigma} - \bar{\mathbf{m}}(\boldsymbol{\sigma})|] \leq N^{-1/4} \sqrt{\nu_N} \sqrt{\mathbb{E} \left[ \sum_{i=1}^N (R_i - 1) \sigma_i \right]^2} \quad (4.19)$$

$$\mathbb{E} \left[ T_N^4 \sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))^2 \right] \leq \nu_N^{2/3} \left( \mathbb{E} \left[ \sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}})^2 \right]^3 \right)^{1/3} \lesssim \nu_N^{2/3} \varepsilon_N \lesssim \nu_N^{2/3} \sqrt{N} \quad (4.20)$$

$$\begin{aligned} \mathbb{E} \left[ |T_N|^3 \left| \sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))^3 \right| \right] &\leq \sqrt{\nu_N} \left( \mathbb{E} \left[ \sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))^2 \right]^4 \right)^{1/4} \left( \mathbb{E} \left[ \max_{1 \leq i \leq N} (m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))^4 \right] \right)^{1/4} \\ &\lesssim \sqrt{\nu_N} \varepsilon_N r_N \lesssim \sqrt{\nu_N} N^{1/4} \end{aligned} \quad (4.21)$$

where the last bounds in (4.20) and (4.21) use (4.17) and (4.18). Finally, for the fifth term in the RHS of (4.15), note that  $|T_N| \leq N^{1/4}$ , and so the first term in the RHS of (4.15) is bounded by  $(2/5) \mathbb{E}[T_N^6]$ . Combining this along with (4.15), (4.16), (4.19), (4.20) and (4.21) gives

$$\nu_N \lesssim \nu_N^{1/3} + \nu_N^{1/2} + \nu_N^{2/3} + \nu_N^{1/2} \sqrt{\frac{\mathbb{E} \left[ \sum_{i=1}^N (R_i - 1) \sigma_i \right]^2}{N^{1/2}}}$$

which completes the proof of (4.13)

(b) Proof of (4.14).

To begin, for any vector  $\mathbf{h} := (h_1, \dots, h_N)$  write

$$\sum_{i=1}^N h_i \sigma_i = \sum_{i=1}^N h_i (\sigma_i - \tanh(m_i(\boldsymbol{\sigma}))) + \sum_{i=1}^N h_i (\tanh(m_i(\boldsymbol{\sigma})) - \tanh(\bar{\mathbf{m}}(\boldsymbol{\sigma}))) + \tanh(\bar{\mathbf{m}}(\boldsymbol{\sigma})) \sum_{i=1}^N h_i,$$

which using part (b) of Lemma 2.1 gives

$$\mathbb{E} \left[ \sum_{i=1}^N h_i \sigma_i \right]^2 \lesssim \|\mathbf{h}\|_2^2 + \|\mathbf{h}\|_2^2 \varepsilon_N + \left[ \sum_{i=1}^N h_i \right]^2 \mathbb{E} \bar{\mathbf{m}}(\boldsymbol{\sigma})^2 \lesssim \|\mathbf{h}\|_2^2 \varepsilon_N + \left[ \sum_{i=1}^N h_i \right]^2 \mathbb{E}(\bar{\mathbf{m}}(\boldsymbol{\sigma})^2), \quad (4.22)$$

where the second line uses part (c) of Lemma 2.2, and  $\varepsilon_N$  equals the RHS of (2.2). Setting  $\mathbf{c} = \mathbf{R} - \mathbf{1}$  and using (4.22) with  $\mathbf{h} = \mathbf{c}$  gives

$$\mathbb{E} \left[ \sum_{i=1}^N c_i \sigma_i \right]^2 \lesssim \|\mathbf{c}\|_2^2 + \|\mathbf{c}\|_2^2 \varepsilon_N + \left[ \sum_{i=1}^N c_i \right]^2 \mathbb{E} \bar{\mathbf{m}}(\boldsymbol{\sigma})^2 \lesssim N, \quad (4.23)$$

where the last line uses (4.18). Along with (4.13) this gives  $\nu_N \lesssim \nu_N^{1/3} + \nu_N^{1/2} + \nu_N^{2/3} + \nu_N^{1/2} \sqrt{N}$ , and so

$$\nu_N \lesssim \sqrt{N} \Rightarrow \mathbb{E} \bar{\sigma}^6 \lesssim N^{-1}. \quad (4.24)$$

Also, an argument similar to the derivation of (4.23) shows that for any positive integer  $p$ , we have

$$\mathbb{E}(\bar{\sigma} - \bar{\mathbf{m}}(\boldsymbol{\sigma}))^{2p} = N^{-2p} \mathbb{E} \left[ \sum_{i=1}^N c_i \sigma_i \right]^{2p} \lesssim N^{-2p} \left( \|\mathbf{c}\|_2^{2p} + \|\mathbf{c}\|_2^{2p} \varepsilon_N^p + \left( \sum_{i=1}^N c_i \right)^{2p} \mathbb{E} \bar{\mathbf{m}}(\boldsymbol{\sigma})^2 \right) \lesssim N^{-p}, \quad (4.25)$$

where the last bound uses (4.18). Combining we have the following conclusions:

$$\mathbb{E} \bar{\mathbf{m}}(\boldsymbol{\sigma})^6 \lesssim \mathbb{E}(\bar{\sigma})^6 + \mathbb{E}(\bar{\sigma} - \bar{\mathbf{m}}(\boldsymbol{\sigma}))^6 \lesssim \frac{1}{N}, \quad (4.26)$$

$$\mathbb{E} \left( \sum_{i=1}^N m_i(\boldsymbol{\sigma})^6 \right) \lesssim N \mathbb{E} \bar{\mathbf{m}}(\boldsymbol{\sigma})^6 + \sqrt{\mathbb{E} \max_{1 \leq i \leq N} (m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))^8} \sqrt{\mathbb{E} \left[ \sum_{i=1}^N (m_i(\boldsymbol{\sigma}) - \bar{\mathbf{m}}(\boldsymbol{\sigma}))^2 \right]^2}$$

$$\lesssim 1 + r_N^4 \varepsilon_N \lesssim 1, \quad (4.27)$$

where (4.26) uses (4.24) and (4.25) with  $p = 3$ , and (4.27) uses (4.26) along with (4.17) and (4.18). Armed with these estimates, we now focus on deriving (4.14).

Let  $\tilde{A}_N$  be as defined in the proof of part (a) of Lemma 2.4, and set  $\mathbf{c}^{(\ell)} := \mathbf{c}^\top \tilde{A}_N^\ell$  and  $x_\ell := \mathbb{E} \left[ \sum_{i=1}^N c_i^{(\ell)} \sigma_i \right]^2$  for  $\ell \geq 0$ . As in the proof of part (b) of Lemma 2.3, we can write  $x_\ell = T_{1,\ell} + T_{2,\ell} + T_{3,\ell}$ , where

$$\begin{aligned} T_{1,\ell} &:= \mathbb{E} \left[ \sum_{i=1}^N c_i^{(\ell)} (\sigma_i - \tanh m_i(\boldsymbol{\sigma})) \right]^2, & T_{2,\ell} &:= \mathbb{E} \left[ \sum_{i=1}^N c_i^{(\ell)} \tanh m_i(\boldsymbol{\sigma}) \right]^2, \\ T_{3,\ell} &:= 2\mathbb{E} \left[ \sum_{i \neq j} c_i^{(\ell)} c_j^{(\ell)} (\sigma_i - \tanh m_i(\boldsymbol{\sigma})) \tanh m_j(\boldsymbol{\sigma}) \right]. \end{aligned}$$

By the argument presented in the proof of part (b) of Lemma 2.3 we have  $T_{1,\ell} \lesssim \|\mathbf{c}^{(\ell)}\|_2^2 \leq \|\mathbf{c}\|_2^2$ , and  $T_{3,\ell} \lesssim \|\mathbf{c}\|_2^2$ . Next, using Taylor Series expansion, we can write  $\tanh(m_i(\boldsymbol{\sigma})) = m_i(\boldsymbol{\sigma}) + \xi_i m_i(\boldsymbol{\sigma})^3$  for random variables  $\{\xi_i\}_{1 \leq i \leq N}$  which are uniformly bounded by 1 in absolute value. Consequently,

$$\begin{aligned} T_{2,\ell} - x_{\ell+1} &= \mathbb{E} \left[ \boldsymbol{\sigma}^\top A_N \mathbf{c}^{(\ell)} + \sum_{i=1}^N c_i^{(\ell)} \xi_i m_i(\boldsymbol{\sigma})^3 \right]^2 - \mathbb{E} \left[ \boldsymbol{\sigma}^\top \tilde{A}_N \mathbf{c}^{(\ell)} \right]^2 \\ &\leq 2\sqrt{x_{\ell+1}} \sqrt{\mathbb{E} \left[ \sum_{i=1}^N |c_i^{(\ell)} m_i(\boldsymbol{\sigma})^3|^2 \right]} + 2\sqrt{x_{\ell+1}} \sqrt{\mathbb{E} \left[ \mathbf{c}^{(\ell)} H_N \boldsymbol{\sigma} \right]^2} + \mathbb{E} \left[ \mathbf{c}^{(\ell)} H_N \boldsymbol{\sigma} \right]^2 \\ &\quad + \mathbb{E} \left[ \sum_{i=1}^N |c_i^{(\ell)} m_i(\boldsymbol{\sigma})^3|^2 \right] + 2\sqrt{\mathbb{E} \left[ \mathbf{c}^{(\ell)} H_N \boldsymbol{\sigma} \right]^2} \sqrt{\mathbb{E} \left[ \sum_{i=1}^N |c_i^{(\ell)} m_i(\boldsymbol{\sigma})^3|^2 \right]} \\ &\leq 2\sqrt{x_{\ell+1}} \|\mathbf{c}\|_2^2 \sqrt{\mathbb{E} \sum_{i=1}^N m_i(\boldsymbol{\sigma})^6} + 2\sqrt{x_{\ell+1}} \sqrt{\mathbb{E} \left[ \mathbf{c}^{(\ell)} H_N \boldsymbol{\sigma} \right]^2} + \mathbb{E} \left[ \mathbf{c}^{(\ell)} H_N \boldsymbol{\sigma} \right]^2 \\ &\quad + \|\mathbf{c}\|_2^2 \mathbb{E} \left[ \sum_{i=1}^N m_i(\boldsymbol{\sigma})^6 \right] + 2\|\mathbf{c}\|_2^2 \sqrt{\mathbb{E} \left[ \mathbf{c}^{(\ell)} H_N \boldsymbol{\sigma} \right]^2} \sqrt{\mathbb{E} \sum_{i=1}^N m_i(\boldsymbol{\sigma})^6}. \end{aligned} \quad (4.28)$$

Proceeding to bound the RHS of (4.28), use (1.9) and (4.25) respectively to note that  $\|H_N\|_{\text{op}} \lesssim N^{-1/4}$ , and  $N\mathbb{E}(\overline{\mathbf{m}}(\boldsymbol{\sigma}))^2 \lesssim N\mathbb{E}(\overline{\boldsymbol{\sigma}})^2 + 1$ , and use and so an application of (4.22) with  $\mathbf{h} = H_N \mathbf{c}^{(\ell)}$  gives

$$\mathbb{E} \left[ (\mathbf{c}^{(\ell)})^\top H_N \boldsymbol{\sigma} \right]^2 \lesssim \|\mathbf{c}^{(\ell)}\|_2^2 \|H_N\|_{\text{op}}^2 \left( \varepsilon_N + N\mathbb{E}(\overline{\mathbf{m}}(\boldsymbol{\sigma}))^2 \right) \lesssim \|\mathbf{c}\|_2^2 \mu_N, \quad (4.29)$$

with  $\mu_N := 1 + \mathbb{E}(N^{1/4} \overline{\boldsymbol{\sigma}})^2$ , where the last line uses (4.17) and (4.18). We now claim that there exists a constant  $D > 0$  such that

$$x_{D(\log N)^2} \lesssim \mu_N \left( \|\mathbf{c}\|_2^2 + N^{-1/2} \left[ \sum_{i=1}^N c_i \right]^2 \right)^2. \quad (4.30)$$

Given this claim, we have the existence of a constant  $C$  free of  $N$  such that

$$x_{D(\log N)^2} \leq C^2 \mu_N \left( \|\mathbf{c}\|_2^2 + N^{-1/2} \left[ \sum_{i=1}^N c_i \right]^2 \right)^2. \quad (4.31)$$

Also, using (4.29) and (4.27), and making  $C$  bigger if needed, for all  $\ell \geq 0$  we have

$$x_\ell \leq x_{\ell+1} + 2C\sqrt{x_{\ell+1}} \|\mathbf{c}\|_2 \sqrt{\mu_N} + C^2 \|\mathbf{c}\|_2^2 \mu_N. \quad (4.32)$$

With  $L = D(\log N)^2$ , we will now show that the bound

$$x_\ell \leq (L - \ell + 1)^2 C^2 \left[ \|\mathbf{c}\|_2^2 + N^{-1/2} \left( \sum_{i=1}^N c_i \right)^2 \right] \quad (4.33)$$

holds for all  $\ell \in [0, L]$  by backwards induction. By (4.31) we have that (4.33) holds for  $\ell = L$ . Suppose (4.33) holds for  $\ell + 1$  for some  $\ell \in [0, L - 1]$ . Using (4.32) gives

$$x_\ell \leq C^2 \mu_N \|\mathbf{c}\|_2^2 \left[ (L - \ell)^2 + 2(L - \ell) + 1 \right] = (L - \ell + 1)^2 C^2 \mu_N \|\mathbf{c}\|_2^2,$$

verifying (4.33) for  $\ell$ , thus verifying (4.33) for all  $\ell \in [0, L]$  by induction. Setting  $\ell = 0$  in (4.33) we get the bound

$$\mathbb{E} \left( \sum_{i=1}^N c_i \sigma_i \right)^2 \leq L^2 C^2 \mu_N \left[ \sum_{i=1}^N c_i^2 + N^{-1/2} \left( \sum_{i=1}^N c_i \right)^2 \right] \leq C^2 D^2 \mu_N (\log N)^4 \left[ \sum_{i=1}^N c_i^2 + N^{-1/2} \left( \sum_{i=1}^N c_i \right)^2 \right],$$

which verifies (4.14), as desired.

It thus remains to verify (4.30), for which using spectral decomposition write  $\tilde{A}_N = \sum_{i=1}^N \tilde{\lambda}_i \tilde{\mathbf{q}}_i \tilde{\mathbf{q}}_i^\top$ , where we set  $\tilde{\lambda}_i := \lambda_i(\tilde{A}_N)$  for convenience of notation. With  $L = D(\log N)^2$ , this gives

$$\mathbf{c}^\top \tilde{A}_N^L \boldsymbol{\sigma} = \bar{\boldsymbol{\sigma}} \sum_{i=1}^N c_i + \tilde{\lambda}_N^L \mathbf{c}^\top \tilde{\mathbf{q}}_N \tilde{\mathbf{q}}_N^\top \boldsymbol{\sigma} + \sum_{i=2}^{N-1} \tilde{\lambda}_i^L \mathbf{c}^\top \tilde{\mathbf{q}}_i \tilde{\mathbf{q}}_i^\top \boldsymbol{\sigma} = \bar{\boldsymbol{\sigma}} \sum_{i=1}^N c_i + \tilde{\lambda}_N^L \mathbf{c}^\top \tilde{\mathbf{q}}_N \tilde{\mathbf{q}}_N^\top \boldsymbol{\sigma} + O(N^{-cD+2}),$$

where the last equality uses Lemma 5.2 to get

$$\max_{2 \leq i \leq N-1} |\tilde{\lambda}_i|^L \leq \left( 1 - \frac{c}{\log N} \right)^\ell \leq N^{-cD}$$

for some  $c > 0$ . Consequently for  $D$  large enough we have

$$\mathbb{E} \left[ \mathbf{c}^\top \tilde{A}_N^L \boldsymbol{\sigma} \right]^2 \lesssim \left[ \sum_{i=1}^N c_i \right]^2 \mathbb{E}[\bar{\boldsymbol{\sigma}}^2] + \|\mathbf{c}\|_2^2 \mathbb{E}[(\tilde{\mathbf{q}}_N^\top \boldsymbol{\sigma})^2]. \quad (4.34)$$

Since  $\tilde{\mathbf{q}}_N^\top \tilde{A}_N = \lambda_N \tilde{\mathbf{q}}_N^\top$  where  $\tilde{\lambda}_N$  is bounded away from 1 by (1.7), we have

$$\begin{aligned} (1 - \tilde{\lambda}_N) \sum_{i=1}^N \tilde{q}_N(i) \sigma_i &= \sum_{i=1}^N \tilde{q}_N(i) (\sigma_i - m_i(\boldsymbol{\sigma})) + \tilde{\mathbf{q}}_N^\top H_N \boldsymbol{\sigma} \\ &= \sum_{i=1}^N \tilde{q}_N(i) (\sigma_i - \tanh(m_i(\boldsymbol{\sigma}))) + \sum_{i=1}^N \tilde{q}_N(i) (\tanh(m_i(\boldsymbol{\sigma})) - m_i(\boldsymbol{\sigma})) + \tilde{\mathbf{q}}_N^\top H_N \boldsymbol{\sigma}. \end{aligned}$$

This immediately gives

$$\begin{aligned} (1 - \tilde{\lambda}_N)^2 \mathbb{E} \left[ \sum_{i=1}^N \tilde{q}_N(i) \sigma_i \right]^2 &\lesssim \mathbb{E} \left[ \sum_{i=1}^N \tilde{q}_N(i) (\sigma_i - \tanh(m_i(\boldsymbol{\sigma}))) \right]^2 + \mathbb{E} \left[ \sum_{i=1}^N |\tilde{q}_N(i)| |m_i(\boldsymbol{\sigma})|^3 \right]^2 + \mathbb{E} \left[ \tilde{\mathbf{q}}_N^\top H_N \boldsymbol{\sigma} \right]^2 \\ &\leq \sum_{i=1}^N \tilde{q}_N(i)^2 + \sqrt{\sum_{i=1}^N \tilde{q}_N(i)^2} \sqrt{\mathbb{E} \left[ \sum_{i=1}^N m_i(\boldsymbol{\sigma})^6 \right]} + \mathbb{E} \left[ \tilde{\mathbf{q}}_N^\top H_N \boldsymbol{\sigma} \right]^2 \\ &\lesssim 1 + \|H_N\|_{\text{op}}^2 \left[ \varepsilon_N + N \mathbb{E}(\bar{\mathbf{m}}(\boldsymbol{\sigma})^2) \right]. \end{aligned}$$

where the last bound uses (4.22) with  $\mathbf{h} = \tilde{\mathbf{q}}_N$ . Since  $N \mathbb{E}(\bar{\mathbf{m}}(\boldsymbol{\sigma})^2) \lesssim N \mathbb{E}(\bar{\boldsymbol{\sigma}}^2) + 1 \lesssim \sqrt{N} \mu_N$ , using the last bound along with (4.34) gives

$$\mathbb{E}(\mathbf{c}^\top \tilde{A}_N^L \boldsymbol{\sigma})^2 \lesssim \mu_N \left( N^{-1/2} \left[ \sum_{i=1}^N c_i \right]^2 + \sum_{i=1}^N c_i^2 \right),$$

thus verifying (4.30), and hence completing the proof of the lemma.  $\square$

**Remark 4.1.** As in the proof of part (b) of Lemma 2.3, the above argument can be modified to bound the moments of general linear combinations  $\sum_{i=1}^N c_i \sigma_i$  for any  $\mathbf{c} \in \mathbb{R}^N$ .



## 5. Proof of supplementary lemmas

### 5.1. Proof of Lemma 3.1 and Lemma 3.2

*Proof of Lemma 3.1.* Noting the presence of  $\text{Tr}^+(D_N)$  in the RHS of the bound, it suffices to prove the result for  $D_N$  with all diagonal entries set to 0. With  $(Z_1, Z_2, \dots, Z_N)$  be i.i.d.  $\mathcal{N}(0, 1)$  random variables, we claim that

$$\left\{ \mathbb{E} \left[ \exp \left( \frac{1}{2} \sum_{i,j=1}^N D_N(i, j) \tilde{X}_i \tilde{X}_j + \sum_{i=1}^N c_i \tilde{X}_i \right) \right] \right\} \leq \left\{ \mathbb{E} \left[ \exp \left( \frac{s_\mu}{2} \sum_{i,j=1}^N D_N(i, j) Z_i Z_j + \sqrt{s_\mu} \sum_{i=1}^N c_i Z_i \right) \right] \right\}. \quad (5.1)$$

Indeed, to see this, recall that the sub-Gaussian norm of  $\tilde{X}_i$  is given by  $s_\mu$  for  $1 \leq i \leq n$ , (see e.g., [33, Theorem 2.1]). Consequently, for every  $\theta \in \mathbb{R}$  we have  $\mathbb{E} \left[ \exp \left( \theta \tilde{X}_i \right) \right] \leq \mathbb{E} \left[ \exp \left( \theta \sqrt{s_\mu} Z_i \right) \right]$ . Using this, (5.1) can be obtained by inductively replacing each  $\tilde{X}_i$  on the left hand side of (5.1) with  $\sqrt{s_\mu} Z_i$ . The RHS of (5.1) can be computed directly to get

$$\log \left\{ \mathbb{E} \left[ \exp \left( \frac{1}{2} \sum_{i,j=1}^N s_\mu D_N(i, j) Z_i Z_j + \sqrt{s_\mu} \sum_{i=1}^N c_i Z_i \right) \right] \right\} = -(1/2) \log \det(I_N - s_\mu D_N) + (1/2) s_\mu \sum_{i=1}^N c_i^2,$$

from which the desired bound follows on noting the existence of  $\rho \in (s_\mu \limsup_{N \rightarrow \infty} \lambda_1(D_N), 1)$ , and using the bound  $-\log(1-x) \lesssim x$  for  $x \in [0, 1-\rho]$ .  $\square$

*Proof of Lemma 3.2.* By Hölder's inequality, for any  $p > 0$  the left hand side of (3.1) can be bounded by

$$\left( \mathbb{E}^{CW} \left[ \exp \left( \frac{\beta(1+p)}{2} \boldsymbol{\sigma}^\top \mathcal{A}_N \boldsymbol{\sigma} \right) \right] \right)^{1/(1+p)} \mathbb{P}(\tilde{W}_N - M(\boldsymbol{\sigma}) | \varepsilon)^{\frac{p}{1+p}}.$$

Since  $\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}(\tilde{W}_N - M(\boldsymbol{\sigma}) | \varepsilon) < 0$  by part (b) of Proposition 5.1, it suffices to show the existence of  $p > 0$  such that

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{E}^{CW} \left[ \exp \left( \frac{\beta(1+p)}{2} \boldsymbol{\sigma}^\top \mathcal{A}_N \boldsymbol{\sigma} \right) \right] \leq 0. \quad (5.2)$$

To this effect, setting

$$g_p(\boldsymbol{\sigma}) := \frac{\beta}{2} \boldsymbol{\sigma}^\top \mathcal{A}_N \boldsymbol{\sigma} + \frac{\beta p}{2} \boldsymbol{\sigma}^\top \mathcal{A}_N \boldsymbol{\sigma}$$

note that

$$\log \mathbb{E}^{CW} \left[ \exp \left( \frac{\beta(1+p)}{2} \boldsymbol{\sigma}^\top \mathcal{A}_N \boldsymbol{\sigma} \right) \right] = \sup_{\boldsymbol{\sigma} \in [-1, 1]^N} \{g_p(\boldsymbol{\sigma}) - \sum_{i=1}^N I(\sigma_i)\} - \log Z_N^{CW}(\beta, B) + o(N), \quad (5.3)$$

where the last line uses [4, Theorem 1.1] along with the observation  $\text{Tr}((A_N + \mathcal{A}_N)^2) = o(N)$ . Using spectral theorem we have  $A_N = \sum_{i=1}^N \lambda_i \mathbf{q}_i \mathbf{q}_i^\top$  with  $\lambda_i = \lambda_i(A_N)$ , and so

$$\begin{aligned} & \sup_{\boldsymbol{\sigma} \in [-1, 1]^N} \left( g_p(\boldsymbol{\sigma}) - \frac{\beta}{2} \sum_{i=1}^N \sigma_i^2 \right) \\ &= \sup_{\boldsymbol{\sigma} \in [-1, 1]^N} \left[ \frac{\beta}{2} \sum_{i=1}^N (\lambda_i - 1) \sigma_i \mathbf{q}_i \mathbf{q}_i^\top \boldsymbol{\sigma} + \frac{\beta p}{2} \boldsymbol{\sigma}^\top \left( \lambda_1 \mathbf{q}_1 \mathbf{q}_1^\top - \frac{\mathbf{1}\mathbf{1}^\top}{N} \right) \boldsymbol{\sigma} + \frac{\beta p}{2} \sum_{i=2}^N \lambda_i \boldsymbol{\sigma}^\top \mathbf{q}_i \mathbf{q}_i^\top \boldsymbol{\sigma} \right] \\ &\lesssim o(N) + \sum_{i=2}^N (\boldsymbol{\sigma}^\top \mathbf{q}_i \mathbf{q}_i^\top \boldsymbol{\sigma}) \left( -\frac{\beta}{2} (1 - \lambda_i) + \frac{\beta p}{2} \lambda_i \right) \end{aligned}$$

where the bound in the last line uses (1.5), and Lemma 5.1. Finally note that (1.7) shows the existence of  $\rho < 1$  such that  $\max_{2 \leq i \leq N} \lambda_i \leq \rho$ , and so there exists  $p = p(\rho)$  such that  $\max_{2 \leq i \leq N} \left( -\frac{\beta}{2} (1 - \lambda_i) + \frac{\beta p}{2} \lambda_i \right) \leq 0$ . Combining we have

$$\sup_{\boldsymbol{\sigma} \in [-1, 1]^N} \left( g_p(\boldsymbol{\sigma}) - \frac{\beta}{2} \sum_{i=1}^N \sigma_i^2 \right) \leq o(N),$$

and so

$$\sup_{\sigma \in [-1,1]^N} (g_p(\sigma) - I(\sigma)) \leq \sup_{\sigma \in [-1,1]^N} \left( g_p(\sigma) - \frac{\beta}{2} \sum_{i=1}^N \sigma_i^2 \right) + \sup_{\sigma \in [-1,1]^N} \left( \frac{\beta}{2} \sum_{i=1}^N \sigma_i^2 - I(\sigma) \right) = o(N) + M_N(\beta, B),$$

where  $M_N(\beta, B)$  is the Mean-Field prediction defined in (1.13). Since  $|\log Z_N^{CW}(\beta, B) - M_N(\beta, B)| \lesssim \log N$  by part (a) of Proposition 5.1, (5.2) follows, thus completing the proof of the lemma.  $\square$

## 5.2. Some results on matrices

**Lemma 5.1.** *Let  $\sum_{i=1}^N \lambda_i(A_N) \mathbf{q}_i \mathbf{q}_i^\top$  be the spectral decomposition of  $A_N$ . Suppose that (1.5) and (1.7) hold, and  $\sum_{i=1}^N (R_i - 1) = o(N)$ .*

(a) *Then  $\|\mathbf{q}_1 - \mathbf{e}\|_2 = o(1)$ , where  $\mathbf{e} := N^{-1/2} \mathbf{1}$ .*

(b) *Further we have  $\limsup_{N \rightarrow \infty} \lambda_1(\mathcal{A}_N) < 1$ , where  $\mathcal{A}_N = A_N - \frac{1}{N} \mathbf{1} \mathbf{1}^\top$ .*

*Proof.* (a) Write  $\mathbf{e} = \sum_{i=1}^N c_i \mathbf{q}_i$  with  $c_1 > 0$  by Perron-Frobenius Theorem, and note that

$$1 + o(1) = \frac{1}{N} \sum_{i=1}^N R_i = \mathbf{e}^\top A_N \mathbf{e} = \sum_{i=1}^N c_i^2 \lambda_i(A_N) \leq \lambda_1(A_N) c_1^2 + \lambda_2(A_N) (1 - c_1^2)$$

Along with (1.5) and (1.7), this gives  $c_1^2 = 1 + o(1)$ , and so  $\langle \mathbf{q}_1, \mathbf{e} \rangle = c_1 = 1 + o(1)$ , thus completing the proof of part (a).

(b) This follows on using part (a) to note that

$$\|\mathcal{A}_N\|_2 \leq \left\| \sum_{i=2}^N \lambda_i(A_N) \mathbf{q}_i \mathbf{q}_i^\top \right\|_2 + \left\| \lambda_1(A_N) \mathbf{q} \mathbf{q}^\top - \mathbf{e} \mathbf{e}^\top \right\|_2 \leq \lambda_2(A_N) + o(1),$$

and using (1.7).  $\square$

**Lemma 5.2.** *Let  $\Gamma_N$  be an  $N \times N$  symmetric matrix with non-negative entries, such that  $\mathbf{1}^\top \Gamma_N = \mathbf{1}^\top$  and  $\Gamma_N$  satisfies (1.7). Then the following conclusions hold:*

(a) *There exists  $c > 0$  such that for all  $\ell \geq 1$  and  $N$  large we have*

$$\max_{1 \leq i \leq N} \Gamma_N^\ell(i, i) \leq \frac{2}{N} + \frac{2}{e^{c\ell}}.$$

(b) *There exists  $\delta > 0$  such that for all  $N$  large enough we have*

$$\max_{2 \leq i \leq N-1} |\lambda_i(\Gamma_N)| \leq 1 - \frac{\delta}{\log N}.$$

*Proof.* (a) Setting  $\lambda_i := \lambda_i(\Gamma_N)$  for simplicity of notation, let  $\mathcal{J}_+ := \{j \in [2, N] : \lambda_j > 0\}$  and  $\mathcal{J}_- := \{j \in [2, N] : \lambda_j < 0\}$ , and use spectral theorem to note that for any positive integer  $\ell$  we have

$$\Gamma_N^\ell = \frac{1}{N} \mathbf{1} \mathbf{1}^\top + \sum_{j \in \mathcal{J}_+} |\lambda_j|^\ell \mathbf{q}_j \mathbf{q}_j^\top + (-1)^\ell \sum_{j \in \mathcal{J}_-} |\lambda_j|^\ell \mathbf{q}_j \mathbf{q}_j^\top,$$

where  $(\mathbf{q}_1, \dots, \mathbf{q}_N)$  are the eigenvectors of  $\Gamma_N$ . To begin, use (1.7) to note the existence of  $c > 0$  such that for all  $N$  large enough we have  $\lambda_2 \leq e^{-c}$ , which gives

$$\sum_{j \in \mathcal{J}_+} |\lambda_j|^\ell p_{ij}^2 \leq \lambda_2^\ell \leq e^{-c\ell} \quad (5.4)$$

For  $\ell$  odd, noting that  $\Gamma_N^\ell(i, i) \geq 0$  gives

$$\sum_{j \in \mathcal{J}_-} |\lambda_j|^\ell q_{ij}^2 \leq \frac{1}{N} + \sum_{j \in \mathcal{J}_+} |\lambda_j|^\ell p_{ij}^2 \leq \frac{1}{N} + \lambda_2^\ell \leq \frac{1}{N} + e^{-c\ell},$$

where the last inequality uses (5.4). Using the fact that  $\max_{2 \leq i \leq N} |\lambda_i| \leq 1$ , for  $\ell \geq 2$  we have

$$\sum_{j \in \mathcal{J}_-} |\lambda_j|^\ell q_{ij}^2 \leq \sum_{j \in \mathcal{J}_-} |\lambda_j|^{\ell-1} q_{ij}^2 \leq \frac{1}{N} + e^{-c\ell}.$$

Combining these two bounds, for all  $\ell \geq 1$  we have

$$|\Gamma_N^\ell(i, i)| \leq \frac{1}{N} + \sum_{j \in \mathcal{J}_+} |\lambda_j|^\ell q_{ij}^2 + \sum_{j \in \mathcal{J}_-} |\lambda_j|^\ell q_{ij}^2 \leq \frac{2}{N} + \frac{2}{e^{c\ell}},$$

thus completing the proof of part (a).

(b) Let  $\delta > 0$  be such that  $3e^{-2\delta/c} > 2$ . Using part (a) with  $\ell = \frac{2 \log N}{c}$  and even, we have

$$\sum_{i=1}^N |\lambda_i|^\ell = \sum_{i=1}^N \Gamma_N^\ell(i, i) \leq 2 + 2Ne^{-2 \log N} \rightarrow 2.$$

On the other hand if  $\max_{2 \leq i \leq N-1} |\lambda_i| > 1 - \frac{\delta}{\log N}$ , then

$$\sum_{i=1}^N |\lambda_i|^\ell \geq 3 \left(1 - \frac{\delta}{\log N}\right)^{\frac{2 \log N}{c}} \rightarrow 3e^{-\frac{2\delta}{c}}.$$

These two together imply  $3e^{-2\delta/c} \leq 2$ , a contradiction.  $\square$

**Remark 5.1.** Note that if  $\Gamma_N$  is the adjacency matrix of a well connected bipartite graph scaled by the average degree, then our lemma implies

$$\lim_{N \rightarrow \infty} \max_{1 \leq i \leq N} \left| N \Gamma_N^{2\ell}(i, i) - 2 \right| = 0$$

for  $\ell = D \log N$  with  $D$  large enough. This highlights the asymptotic optimality of the bound obtained in part (a) of Lemma 5.2. Part (b) quantifies the graph theoretic fact that a connected regular graph, scaled by its degree, the multiplicity of the eigenvalue  $-1$  can be at most 1. It is easy to check that if  $-1$  happens to be an eigenvalue the graph must be a bipartite graph, and all other eigenvalues will be strictly larger than  $-1$  (i.e. there is a unique bipartition for a connected bipartite graph). In fact, our proof can be modified to show the stronger conclusion that for a regular well connected bipartite graphs, the second last eigenvalue is bounded away from  $-1$ , i.e.

$$\liminf_{N \rightarrow \infty} \lambda_{N-1}(\Gamma_N) > -1.$$

### 5.3. Some results for the Curie-Weiss model

The following proposition collects all the results for the Curie-Weiss model which we have used previously.

**Proposition 5.1.** Suppose  $\sigma$  is drawn from the Curie-Weiss model. With  $\widetilde{W}_N$  as in Proposition 3.1, the following conclusions hold:

(a)

$$\begin{aligned} \log Z_N^{CW}(\beta, B) - N \left\{ \frac{\beta}{2} t^2 + Bt - I(t) \right\} &\lesssim 1 && \text{if } (\beta, B) \in \Theta_1 \cup \Theta_2, \\ &\lesssim \log N && \text{if } (\beta, B) \in \Theta_3. \end{aligned}$$

(b) For any  $\lambda > 0$ , we have

$$\log \mathbb{P}^{CW}(|\widetilde{W}_N - M(\sigma)| \geq \lambda) \lesssim -N\lambda^2 \quad \text{if } (\beta, B) \in \Theta_1 \cup \Theta_2,$$

$$\lesssim -N \min(\lambda^2, \lambda^4) \quad \text{if } (\beta, B) \in \Theta_3.$$

Consequently for any sequence  $\delta_N = o(N)$  we have

$$\begin{aligned} \log \mathbb{E}^{CW} e^{\delta_N (\widetilde{W}_N - M(\boldsymbol{\sigma}))^2} &\lesssim 1 \quad \text{if } (\beta, B) \in \Theta_1 \cup \Theta_2, \\ &\lesssim \frac{\delta_N^2}{N} \quad \text{if } (\beta, B) \in \Theta_3. \end{aligned}$$

(c) For  $(\beta, B) \in \Theta_2$ , we have:

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}^{CW} \left( \sum_{i=1}^N \sigma_i \in \{-1, 0, 1\} \right) < 0.$$

*Proof.* (a) With  $f(w) = \frac{\beta w^2}{2} - \log \cosh(\beta w + B)$  as in 3.1, a direct computation gives  $Z_N^{CW}(\beta, B) = e^{-\beta/2} \sqrt{\frac{n\beta}{2\pi}} \int_{\mathbb{R}} e^{-nf(w)} dw$ , where the function  $f(w)$  has a unique global minimum at  $w = t$  for  $(\beta, B) \in \Theta_1 \cup \Theta_3$ , and two global minima at  $\pm t$  for  $(\beta, B) \in \Theta_2$ . Also, it is easy to verify that

$$\begin{aligned} f(w) - f(t) &\cong (w - t)^2, & \text{for all } w \in \mathbb{R}, & \quad \text{if } (\beta, B) \in \Theta_1, \\ f(w) - f(t) &\cong (w - t)^2 & \text{for all } w > 0, & \quad \text{if } (\beta, B) \in \Theta_2, \\ f(w) - f(t) &\cong \min \left[ (w - t)^2, (w - t)^4 \right] & \text{for all } w \in \mathbb{R}, & \quad \text{if } (\beta, B) \in \Theta_3. \end{aligned} \tag{5.5}$$

The desired estimates follow from these bounds and using Laplace method.

(b) Noting that

$$|\widetilde{W}_N - M(\boldsymbol{\sigma})| = |\tanh(\beta W_N + B) - \tanh(\beta M(\boldsymbol{\sigma}) + B)| \leq \beta |W_N - M(\boldsymbol{\sigma})|,$$

it suffices to prove the desired bounds  $W_N$ , which follows from straightforward computations on using (5.5).

(c) This follows on using part (b) to note that, when  $(\beta, B) \in \Theta_2$ , the random variable  $W_N$  has an exponential concentration near the points  $\pm t$ , none of which are near 0.

□

## References

- [1] ADAMCZAK, R. A., KOTOWSKI, M., POLACZYK, B. O. and STRZELECKI, M. (2019). A note on concentration for polynomials in the Ising model. *Electron. J. Probab.* **24** Paper No. 42, 22. [MR3949267](#)
- [2] AUGERI, F. (2019). A transportation approach to the mean-field approximation. *arXiv preprint arXiv:1903.08021*.
- [3] BANDEIRA, A. S. and VAN HANDEL, R. (2016). Sharp nonasymptotic bounds on the norm of random matrices with independent entries. *Ann. Probab.* **44** 2479–2506. [MR3531673](#)
- [4] BASAK, A. and MUKHERJEE, S. (2017). Universality of the mean-field for the Potts model. *Probab. Theory Related Fields* **168** 557–600. [MR3663625](#)
- [5] BERTHET, Q., RIGOLLET, P. and SRIVASTAVA, P. (2019). Exact recovery in the Ising blockmodel. *Ann. Statist.* **47** 1805–1834. [MR3953436](#)
- [6] BORGS, C., CHAYES, J. T., LOVÁSZ, L., SÓS, V. T. and VESZTERGOMBI, K. (2008). Convergent sequences of dense graphs. I. Subgraph frequencies, metric properties and testing. *Adv. Math.* **219** 1801–1851. [MR2455626](#)
- [7] BORGS, C., CHAYES, J. T., LOVÁSZ, L., SÓS, V. T. and VESZTERGOMBI, K. (2012). Convergent sequences of dense graphs II. Multiway cuts and statistical physics. *Ann. of Math. (2)* **176** 151–219. [MR2925382](#)
- [8] BORGS, C., CHAYES, J. T., COHN, H. and ZHAO, Y. (2018). An  $L^p$  theory of sparse graph convergence II: LD convergence, quotients and right convergence. *Ann. Probab.* **46** 337–396. [MR3758733](#)
- [9] BORGS, C., CHAYES, J. T., COHN, H. and ZHAO, Y. (2019). An  $L^p$  theory of sparse graph convergence I: Limits, sparse random graph models, and power law distributions. *Trans. Amer. Math. Soc.* **372** 3019–3062. [MR3988601](#)

- [10] BRESLER, G. and NAGARAJ, D. (2019). Stein’s method for stationary distributions of Markov chains and application to Ising models. *Ann. Appl. Probab.* **29** 3230–3265. [MR4019887](#)
- [11] BRODER, A. and SHAMIR, E. (1987). On the second eigenvalue of random regular graphs. In *28th Annual Symposium on Foundations of Computer Science (sfcs 1987)* 286–294. IEEE.
- [12] CHATTERJEE, S. (2005). *Concentration inequalities with exchangeable pairs*. ProQuest LLC, Ann Arbor, MI Thesis (Ph.D.)–Stanford University. [MR2707160](#)
- [13] CHATTERJEE, S. and DEMBO, A. (2016). Nonlinear large deviations. *Adv. Math.* **299** 396–450. [MR3519474](#)
- [14] CHATTERJEE, S. and SHAO, Q.-M. (2011). Nonnormal approximation by Stein’s method of exchangeable pairs with application to the Curie-Weiss model. *Ann. Appl. Probab.* **21** 464–483. [MR2807964](#)
- [15] CHUANG, H. and OMIDI, G. R. (2009). Graphs with three distinct eigenvalues and largest eigenvalues less than 8. *Linear Algebra Appl.* **430** 2053–2062. [MR2503952](#)
- [16] COMETS, F. and GIDAS, B. (1991). Asymptotics of maximum likelihood estimators for the Curie-Weiss model. *Ann. Statist.* **19** 557–578. [MR1105836](#)
- [17] DEMBO, A. and MONTANARI, A. (2010). Gibbs measures and phase transitions on sparse random graphs. *Braz. J. Probab. Stat.* **24** 137–211. [MR2643563](#)
- [18] DESHPANDE, Y., SEN, S., MONTANARI, A. and MOSSEL, E. (2018). Contextual stochastic block models. In *Advances in Neural Information Processing Systems* 8581–8593.
- [19] EICHELSBACHER, P. and LÖWE, M. (2010). Stein’s method for dependent random variables occurring in statistical mechanics. *Electron. J. Probab.* **15** no. 30, 962–988. [MR2659754](#)
- [20] EL DAN, R. (2018). Taming correlations through entropy-efficient measure decompositions with applications to mean-field approximation. *Probability Theory and Related Fields* 1–19.
- [21] ELLIS, R. S. and NEWMAN, C. M. (1978). The statistics of Curie-Weiss models. *J. Statist. Phys.* **19** 149–161. [MR0503332](#)
- [22] FEIGE, U. and OFEK, E. (2005). Spectral techniques applied to sparse random graphs. *Random Structures Algorithms* **27** 251–275. [MR2155709](#)
- [23] GHEISSARI, R., LUBETZKY, E. and PERES, Y. (2018). Concentration inequalities for polynomials of contracting Ising models. *Electron. Commun. Probab.* **23** Paper No. 76, 12. [MR3873783](#)
- [24] GIARDINÀ, C., GIBERTI, C., VAN DER HOFSTAD, R. and PRIORIELLO, M. L. (2016). Annealed central limit theorems for the Ising model on random graphs. *ALEA Lat. Am. J. Probab. Math. Stat.* **13** 121–161. [MR3476210](#)
- [25] JAIN, V., KOEHLER, F. and RISTESKI, A. (2019). Mean-field approximation, convex hierarchies, and the optimality of correlation rounding: a unified perspective. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing* 1226–1236.
- [26] KABLUCHKO, Z., LÖWE, M. and SCHUBERT, K. (2019). Fluctuations of the Magnetization for Ising Models on Erdős-Rényi Random Graphs—the Regimes of Small  $p$  and the Critical Temperature. *arXiv preprint arXiv:1911.10624*.
- [27] LIU, L. (2017). On the Log Partition Function of Ising Model on Stochastic Block Model. *arXiv preprint arXiv:1710.05287*.
- [28] LOVÁSZ, L. (2012). *Large networks and graph limits*. American Mathematical Society Colloquium Publications **60**. American Mathematical Society, Providence, RI. [MR3012035](#)
- [29] LÖWE, M. and SCHUBERT, K. (2018). Fluctuations for block spin Ising models. *Electron. Commun. Probab.* **23** Paper No. 53, 12. [MR3852267](#)
- [30] MOSSEL, E., NEEMAN, J. and SLY, A. (2012). Stochastic block models and reconstruction. *arXiv preprint arXiv:1202.1499*.
- [31] MUKHERJEE, R., MUKHERJEE, S. and YUAN, M. (2018). Global testing against sparse alternatives under Ising models. *Ann. Statist.* **46** 2062–2093. [MR3845011](#)
- [32] MUKHERJEE, R. and RAY, G. (2019). On testing for parameters in Ising models. *arXiv preprint arXiv:1906.00456*.
- [33] OSTROVSKY, E. and SIROTA, L. (2014). Exact value for subgaussian norm of centered indicator random variable. *arXiv e-prints arXiv:1405.6749*.
- [34] RAVIKUMAR, P., WAINWRIGHT, M. J. and LAFFERTY, J. D. (2010). High-dimensional Ising model selection using  $\ell_1$ -regularized logistic regression. *Ann. Statist.* **38** 1287–1319. [MR2662343](#)
- [35] SLY, A. and SUN, N. (2014). Counting in two-spin models on  $d$ -regular graphs. *Ann. Probab.* **42** 2383–2416. [MR3265170](#)
- [36] XU, M. and MUKHERJEE, S. (2020). Fluctuations in the two-star exponential random graph model. *In preparation*.