

Approximations Related to the Sums of m -dependent Random Variables

Amit N. Kumar¹, Neelesh S. Upadhye², and P. Vellaisamy³

^{1,3}Department of Mathematics, Indian Institute of Technology Bombay,
Powai, Mumbai-400076, India.

²Department of Mathematics, Indian Institute of Technology Madras,
Chennai-600036, India.

¹Email: amit.kumar2703@gmail.com

²Email: neelesh@iitm.ac.in

³Email: pv@math.iitb.ac.in

Abstract

In this paper, we consider the sums of non-negative integer valued m -dependent random variables, and its approximation to the power series distribution. We first discuss some relevant results for power series distribution such as Stein operator, uniform and non-uniform bounds on the solution of Stein equation, and etc. Using Stein's method, we obtain the error bounds for the approximation problem considered. As special cases, we discuss two applications, namely, 2-runs and (k_1, k_2) -runs and compare the bound with the existing bounds.

Keywords: Power series distribution; m -dependent random variables; Stein's method; Runs.

MSC 2010 Subject Classifications : Primary : 62E17, 62E20 ; Secondary : 60F05, 60E05.

1 Introduction and Preliminaries

The sums of m -dependent random variables (rvs) has special attention due to its applicability in many real-life applications such as runs and patterns, DNA sequences, and reliability theory, among many others. However, its distribution is difficult or sometimes intractable, especially if the setup is arising from non-identical rvs concentrated on $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$, the set of non-negative integers. Therefore, there is a need to approximate such a distribution with some known and easy-to-use distributions. In

this article, we consider power series distribution (PSD) approximation to the sums of m -dependent rvs. Approximations related to the sums of locally dependent rvs have been studied by several authors such as Barbour and Xia [4, 5], Fu and Johnson [9], Vellaisamy [26], Wang and Xia [27], and Soon [21], among many others.

A sequence of rvs $\{X_k\}_{k \geq 1}$ is called m -dependent if $\sigma(X_1, X_2, \dots, X_i)$ and $\sigma(X_j, X_{j+1}, \dots)$, for $j - i > m$, are independent, where $\sigma(X)$ denotes the sigma-algebra generated by X . The sums of m -dependent rvs can be reduced to the sums of 1-dependent rvs, using rearrangement of rvs (see Section 3 for details). We mainly focus on the sums of 1-dependent rvs concentrated on \mathbb{Z}_+ , and obtain the error bounds. Of course, the bound can directly apply for special distributions of PSD family. An advantage of approximation to PSD family is that we can obtain the error bounds for approximation to some specific distributions such as Poisson and negative binomial distributions. For some related works, we refer the reader to Lin and Liu [15], Čekanavičius and Vellaisamy [25], and references therein.

For \mathbb{Z}_+ -valued rvs X and X^* , the total variation distance is given by

$$d_{TV}(X, X^*) = \frac{1}{2} \sum_{k=0}^{\infty} |\mathbb{P}(X = k) - \mathbb{P}(X^* = k)|. \quad (1.1)$$

Hereafter, $\mathbf{1}_A$ denotes the indicator function of $A \subseteq \mathbb{Z}_+$. Let X be a rv concentrated on \mathbb{Z}_+ ,

$$\mathcal{G} = \{f : \mathbb{Z}_+ \rightarrow \mathbb{R} \mid f \text{ is bounded}\}$$

and

$$\mathcal{G}_X = \{g \in \mathcal{G} \mid g(0) = 0 \text{ and } g(x) = 0 \text{ for } x \notin \text{supp}(X)\}, \quad (1.2)$$

associated with the rv X , where $\text{supp}(X)$ denotes the support of the rv X . We now briefly discuss Stein's method (Stein [22]) which we use to derive our approximation results in Section 3. The Stein's method can be discussed in following three steps.

- (a) Identify a Stein operator, denoted by \mathcal{A}_X for a rv X , such that $\mathbb{E}[\mathcal{A}_X g(X)] = 0$, for $g \in \mathcal{G}_X$.
- (b) Solve the Stein equation $\mathcal{A}_X g(k) = f(k) - \mathbb{E}f(X)$, for $f \in \mathcal{G}$ and $g \in \mathcal{G}_X$.
- (c) Replace k by a rv Y in Stein equation, and taking expectation and supremum to get

$$d_{TV}(X, Y) := \sup_{f \in \mathcal{H}} |\mathbb{E}f(X) - \mathbb{E}f(Y)| = \sup_{f \in \mathcal{H}} |\mathbb{E}\mathcal{A}_X g_f(Y)|,$$

where g_f is the solution of the Stein equation and $\mathcal{H} = \{\mathbf{1}_A \mid A \subseteq \mathbb{Z}_+\}$.

For additional details on Stein's method, see Barbour *et al.* [3], Barbour and Chen [2], Ley *et al.* [14], Reinert [19], Upadhye *et al.* [24], and the references therein.

This article is organized as follows. In Section 2, we discuss the PSD and its related results to Stein's method. In Section 3, we derive the error bound for PSD approximation to the sums of 1-dependent rvs and discuss some relevant remarks. In Section 4, we discuss two important applications of our results to the sums of 2-runs and (k_1, k_2) -runs.

2 Power Series Distribution and Related Results

Let Z be a \mathbb{Z}_+ -values rv. We say its distribution belongs to the PSD family, denoted by \mathcal{P} , if $\mathbb{P}(Z = k) = p_k$ is of the form

$$p_k = \frac{a_k \theta^k}{\gamma(\theta)}, \quad k \in \mathbb{Z}_+, \quad (2.1)$$

where $\theta > 0$ and $a_k, k \geq 0$, are called series parameter and coefficient function, respectively. Many distributions such as Poisson, binomial, negative binomial, logarithmic series, and inverse sine distributions, among many others, belong to the PSD family. For more details, we refer the reader to Edwin [7], Noack [16], Patil [18], and the references therein.

Next, we give a brief discussion about Stein's method for PSD, in fact, many results follow from Eichelsbacher and Reinert [8]. The following proposition gives a Stein operator for PSD.

Proposition 2.1. *Let the rv Z having distribution belonging to PSD family defined in (2.1). Then a Stein operator for Z is given by*

$$\mathcal{A}_Z g(k) = \theta(k+1) \frac{a_{k+1}}{a_k} g(k+1) - k g(k), \quad g \in \mathcal{G}_Z, \quad k \in \mathbb{Z}_+. \quad (2.2)$$

Proof. From (2.1), it can be easily verified that

$$\theta(k+1) \frac{a_{k+1}}{a_k} p_k - (k+1) p_{k+1} = 0. \quad (2.3)$$

Let $g \in \mathcal{G}_Z$ defined in (1.2), then

$$\sum_{k=0}^{\infty} g(k+1) \left[\theta(k+1) \frac{a_{k+1}}{a_k} p_k - (k+1) p_{k+1} \right] = 0.$$

Rearranging the terms, we have

$$\sum_{k=0}^{\infty} \left[\theta(k+1) \frac{a_{k+1}}{a_k} g(k+1) - k g(k) \right] p_k = 0.$$

This proves the result. □

Now, we discuss the solution of the Stein equation

$$\theta(k+1)\frac{a_{k+1}}{a_k}g(k+1) - kg(k) = f(k) - \mathbb{E}f(Z), \quad f \in \mathcal{G}, g \in \mathcal{G}_Z. \quad (2.4)$$

Next we describe discrete Gibbs measure (DGM), a large class of distributions, studied by Eichelsbacher and Reinert [8]. If a rv U has the distribution of the form

$$\mathbb{P}(U = k) = \frac{1}{\mathcal{C}_w} e^{V(k)} \frac{w^k}{k!}, \quad k \in \mathbb{Z}_+, \quad (2.5)$$

for some function $V : \mathbb{Z}_+ \rightarrow \mathbb{R}$, $w > 0$, and $\mathcal{C}_w = \sum_{k=0}^{\infty} e^{V(k)} \frac{w^k}{k!}$, then we say the rv U belongs to the DGM family. Observe here the support is \mathbb{Z}_+ . Note that if we take $a_k = e^{V(k)}/k! \iff V(k) = \ln(a_k k!)$, $\theta = w$, and $\gamma(\theta) = \mathcal{C}_w$, which are valid choices, then the results derived by Eichelsbacher and Reinert [8] are valid for PSD family. Therefore, the solution of (2.4) can be directly obtained from (2.5) and (2.6) of Eichelsbacher and Reinert [8] and is given by

$$\begin{aligned} g(k) &= \frac{1}{k a_k \theta^k} \sum_{j=0}^{k-1} a_j \theta^j [f(j) - \mathbb{E}f(Z)] \\ &= -\frac{1}{k a_k \theta^k} \sum_{j=k}^{\infty} a_j \theta^j [f(j) - \mathbb{E}f(Z)]. \end{aligned}$$

Also, the Lemma 2.1 of Eichelsbacher and Reinert [8] can be written for PSD family in the following manner.

Lemma 2.1. *Let $\mathcal{G}_1 = \{f : \mathbb{Z}_+ \rightarrow [0, 1]\}$, $F(k) = \sum_{i=0}^k p_i$ and $\bar{F}(k) = \sum_{i=k}^{\infty} p_i$. Assume that*

$$k \frac{F(k)}{F(k-1)} \geq \theta(k+1) \frac{a_{k+1}}{a_k} \geq k \frac{\bar{F}(k+1)}{\bar{F}(k)}.$$

Then, for $f \in \mathcal{G}_1$ and g_f , the solution of (2.4), we have

$$\sup_{f \in \mathcal{B}} |\Delta g_f(k)| = \frac{a_k}{\theta(k+1)a_{k+1}} \bar{F}(k+1) + \frac{1}{k} F(k-1),$$

where $\Delta g_f(k) = g_f(k+1) - g_f(k)$.

Moreover,

$$\sup_{f \in \mathcal{B}} |\Delta g_f(k)| \leq \frac{1}{k} \wedge \frac{a_k}{\theta(k+1)a_{k+1}}, \quad (2.6)$$

where $x \wedge y$ denotes the minimum of x and y .

Now, it is not easy to use direct form of the Stein operator (2.2) as a_k is unknown and depends on k . So, we consider PSD family with Panjer's recursive relation (see Panjer and Wang [17] for details), denoted by \mathcal{P}_1 , which is given by

$$(k+1)\frac{p_{k+1}}{p_k} = a + bk \implies \theta(k+1)\frac{a_{k+1}}{a_k} = a + bk, \quad \text{for some } a, b \in \mathbb{R}. \quad (2.7)$$

Therefore, the stein operator (2.2) can be written as

$$\mathcal{A}_Z g(k) = (a + bk)g(k+1) - kg(k), \quad k \in \mathbb{Z}_+. \quad (2.8)$$

Also, the bound (2.6) becomes

$$\sup_{f \in \mathcal{B}} |\Delta g_f(k)| \leq \frac{1}{k} \wedge \frac{1}{a + bk}, \quad k \geq 1. \quad (2.9)$$

Note that if $a, b \geq 0$ (PSD family satisfies Panjer recursive relation with $a, b \geq 0$, denoted by \mathcal{P}_2) then the bound (2.9) becomes uniform and is given by

$$\sup_{f \in \mathcal{B}} |\Delta g_f(k)| \leq 1 \wedge \frac{1}{a}, \quad k \geq 1. \quad (2.10)$$

Note that $\mathcal{P}_2 \subset \mathcal{P}_1 \subset \mathcal{P}$. Also, observe that $a = \lambda, b = 0$ ($a_k = 1/k!, \theta = \lambda$ and $\gamma(\theta) = e^\theta$) and $a = nq, b = q$ ($a_k = \binom{n+k-1}{k}$, $\theta = q$ and $\gamma(\theta) = (1-\theta)^{-n}$) for Poisson (with parameter λ) and negative binomial (with parameter n and $p = 1 - q$) distributions, respectively, and hence the bounds (from (2.10)) are $1 \wedge \frac{1}{\lambda}$ and $1 \wedge \frac{1}{nq}$, respectively, which are well-known bounds for Poisson and negative binomial distributions. Many distributions satisfy the condition $a, b \geq 0$. However, if the condition is not satisfied, one can still use (2.9) to compute the uniform bound. For example, if $a_k = \binom{n}{k}$, $\theta = p/q$, and $\gamma(\theta) = (1 + \theta)^n$, then $Z \sim \text{Bi}(n, p)$, and $\frac{\theta(k+1)a_{k+1}}{a_k} = \frac{p}{q}(n - k)$, and hence $a = np/q$ and $b = -p/q \leq 0$. Therefore, the bound (2.9) is

$$\begin{aligned} \sup_{f \in \mathcal{B}} |\Delta g_f(k)| &\leq \frac{1}{k} \wedge \frac{q}{p(n-k)} \\ &= \begin{cases} \frac{1}{k} & \text{if } k \geq np \\ \frac{q}{(n-k)p} & \text{if } k \leq np \end{cases} \\ &\leq \begin{cases} \frac{1}{np} & \text{if } k \geq np \\ \frac{1}{np} & \text{if } k \leq np \end{cases} \end{aligned}$$

$$= \frac{1}{np}, \text{ for all } k, \quad (2.11)$$

which leads to a uniform bound for binomial distribution. Note here that the Stein operator (from (2.8)) is

$$\mathcal{A}_Z g(k) = \frac{p}{q}(n-k)g(k+1) - kg(k). \quad (2.12)$$

But, the well-known Stein operator for the binomial distribution is

$$\mathcal{A}_Z g(k) = p(n-k)g(k+1) - qkg(k), \quad (2.13)$$

which follows by multiplying q in (2.12). Also, the uniform bound will be changed and is given by $1/npq$ (that is, divided by q), which is well-known bound with respect to the Stein operator (2.13) (see Upadhye *et al.* [24]). Hence, throughout this article, we use $\|\Delta g\| = \sup_k |\Delta g(k)|$ and the uniform bound for $\|\Delta g\|$ can be obtained from (2.10) or may be computed explicitly for some applications. Next, let $\phi_Z(\cdot)$ be the probability generating function of Z . Then, using (2.7), it can be seen that

$$\phi'_Z(t) = \frac{a\phi_Z(t)}{1-bt}.$$

Hence, mean and variance of the PSD are given by

$$\mathbb{E}(Z) = \frac{a}{1-b} \quad \text{and} \quad \text{Var}(Z) = \frac{a}{(1-b)^2}. \quad (2.14)$$

For more details, we refer the reader to Edwin [7], and Panjer and Wang [17].

3 Approximation Results

Let Y_1, Y_2, \dots, Y_n be a sequence of \mathbb{Z}_+ -valued m dependent rvs and $S_n = \sum_{i=1}^n Y_i$, the sums of m -dependent rvs. Then, grouping the consecutive summations in the following form

$$Y_i^* := \sum_{j=(i-1)m+1}^{\min(im,n)} Y_j, \quad j = 1, 2, \dots, \lfloor n/m \rfloor + 1,$$

where $\lfloor x \rfloor$ denotes the greatest integer function of x , $S_n = \sum_{j=1}^{\lfloor n/m \rfloor + 1} Y_j^*$ become the sums of 1-dependent rvs. In this section, we derive an error bound for PSD approximation to the sums of 1-dependent rvs in total variation distance and discuss some relevant remarks. Throughout this section,

we assume $X_1, X_2, \dots, X_n, n \geq 1$, is a sequence of 1-dependent rvs and

$$W_n = \sum_{i=1}^n X_i. \quad (3.1)$$

For any \mathbb{Z}_+ -valued rv Y , let $D(Y) := 2d_{TV}(Y, Y + 1)$, where $d_{TV}(X, X^*)$ as defined in (1.1). Let

$$N_{i,\ell} := \{j : |j - i| \leq \ell\} \cap \{1, 2, \dots, n\} \quad \text{and} \quad X_{N_{i,\ell}} := \sum_{j \in N_{i,\ell}} X_j, \quad \text{for } \ell = 1, 2.$$

Note that $X_{N_{i,2}} - X_{N_{i,1}} = X_{N_{i,2} - N_{i,1}}$. From (3.1), it can be verified that $\mathbb{E}(W_n) = \sum_{i=1}^n \mathbb{E}(X_i)$ and

$$\text{Var}(W_n) = \sum_{i=1}^n \sum_{|j-i| \leq 1} [\mathbb{E}(X_i X_j) - \mathbb{E}(X_i)\mathbb{E}(X_j)] = \sum_{i=1}^n [\mathbb{E}(X_i X_{N_{i,1}}) - \mathbb{E}(X_i)\mathbb{E}(X_{N_{i,1}})]. \quad (3.2)$$

Now, the following theorem gives the error bound for Z -approximation to W_n .

Theorem 3.1. *Let $Z \in \mathcal{P}_1$ and W_n be defined as in (3.1). Assume that $\mathbb{E}(Z) = \mathbb{E}(W_n)$, and $\tau = \text{Var}(W_n) - \text{Var}(Z)$. Then, for $n \geq 6$,*

$$d_{TV}(W_n, Z) \leq \|\Delta g\| \left\{ \frac{|1-b|}{2} \left[\sum_{i=1}^n \mathbb{E}(X_i) \mathbb{E}[X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)D(W_n | X_{N_{i,1}}, X_{N_{i,2}})] \right. \right. \\ \left. \left. + \sum_{i=1}^n \mathbb{E}[X_i X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)D(W_n | X_{N_{i,1}}, X_{N_{i,2}})] \right] \right. \\ \left. + \sum_{i=1}^n \mathbb{E}[X_i(X_{N_{i,2}} - 1)D(W_n | N_{i,2})] + |\tau(1-b)| \right\}. \quad (3.3)$$

Proof. Consider the Stein operator given in (2.8) and taking expectation with respect to W_n , we have

$$\begin{aligned} \mathbb{E}[\mathcal{A}_Z g(W_n)] &= a\mathbb{E}[g(W_n + 1)] + b\mathbb{E}[W_n g(W_n + 1)] - \mathbb{E}[W_n g(W_n)] \\ &= a\mathbb{E}[g(W_n + 1)] - (1-b)\mathbb{E}[W_n g(W_n + 1)] + \mathbb{E}[W_n \Delta g(W_n)] \\ &= (1-b) \left[\frac{a}{(1-b)} \mathbb{E}[g(W_n + 1)] - \mathbb{E}[W_n g(W_n + 1)] \right] + \mathbb{E}[W_n \Delta g(W_n)]. \end{aligned}$$

Applying the first moment matching condition, $\mathbb{E}(Z) = a/(1-b) = \mathbb{E}(W_n)$, we get

$$\mathbb{E}[\mathcal{A}_Z g(W_n)] = (1-b) \left[\mathbb{E}(W_n) \mathbb{E}[g(W_n + 1)] - \mathbb{E}[W_n g(W_n + 1)] \right] + \mathbb{E}[W_n \Delta g(W_n)]. \quad (3.4)$$

Let now

$$W_{i,n} := W_n - X_{N_{i,1}}$$

so that X_i and $W_{i,n}$ are independent. Consider the following expression from (3.4)

$$\begin{aligned}
\mathbb{E}(W_n)\mathbb{E}[g(W_n + 1)] - \mathbb{E}[W_n g(W_n + 1)] &= \sum_{i=1}^n \mathbb{E}(X_i)\mathbb{E}[g(W_n + 1)] - \sum_{i=1}^n \mathbb{E}[X_i g(W_n + 1)] \\
&= \sum_{i=1}^n \mathbb{E}(X_i)\mathbb{E}[g(W_n + 1)] - \sum_{i=1}^n \mathbb{E}[X_i g(W_n + 1)] \\
&\quad - \sum_{i=1}^n \mathbb{E}[X_i g(W_{i,n} + 1)] + \sum_{i=1}^n \mathbb{E}[X_i g(W_{i,n} + 1)] \\
&= \sum_{i=1}^n \mathbb{E}(X_i)\mathbb{E}[g(W_n + 1) - g(W_{i,n} + 1)] \\
&\quad - \sum_{i=1}^n \mathbb{E}[X_i(g(W_n + 1) - g(W_{i,n} + 1))]. \tag{3.5}
\end{aligned}$$

It can be seen that

$$\begin{aligned}
g(W_n + 1) - g(W_{i,n} + 1) &= g(W_{i,n} + X_{N_{i,1}} + 1) - g(W_{i,n} + 1) \\
&= g(W_{i,n} + X_{N_{i,1}} + 1) - g(W_{i,n} + X_{N_{i,1}}) \\
&\quad + g(W_{i,n} + X_{N_{i,1}}) - g(W_{i,n} + X_{N_{i,1}} - 1) \\
&\quad \vdots \\
&\quad + g(W_{i,n} + 2) - g(W_{i,n} + 1) \\
&= \sum_{j=1}^{X_{N_{i,1}}} \Delta g(W_{i,n} + j). \tag{3.6}
\end{aligned}$$

Using (3.6) in (3.5), we get

$$\begin{aligned}
\mathbb{E}(W_n)\mathbb{E}[g(W_n + 1)] - \mathbb{E}[W_n g(W_n + 1)] &= \sum_{i=1}^n \mathbb{E}(X_i)\mathbb{E}\left[\sum_{j=1}^{X_{N_{i,1}}} \Delta g(W_{i,n} + j)\right] \\
&\quad - \sum_{i=1}^n \mathbb{E}\left[X_i \sum_{j=1}^{X_{N_{i,1}}} \Delta g(W_{i,n} + j)\right]. \tag{3.7}
\end{aligned}$$

Substituting (3.7) in (3.4), we have

$$\begin{aligned} \mathbb{E}[\mathcal{A}_Z g(W_n)] &= (1-b) \left\{ \sum_{i=1}^n \mathbb{E}(X_i) \mathbb{E} \left[\sum_{j=1}^{X_{N_{i,1}}} \Delta g(W_{i,n} + j) \right] - \sum_{i=1}^n \mathbb{E} \left[X_i \sum_{j=1}^{X_{N_{i,1}}} \Delta g(W_{i,n} + j) \right] \right\} \\ &\quad + \sum_{i=1}^n \mathbb{E}[X_i \Delta g(W_n)]. \end{aligned} \quad (3.8)$$

Note that $\mathbb{E}(Z) = a/(1-b) = \mathbb{E}(W_n) = \sum_{i=1}^n \mathbb{E}(X_i)$. Therefore, from (2.14),

$$\text{Var}(Z) = \frac{a}{(1-b)^2} = \frac{1}{(1-b)} \sum_{i=1}^n \mathbb{E}(X_i).$$

Hence,

$$\tau = \text{Var}(W_n) - \text{Var}(Z) = \sum_{i=1}^n \mathbb{E}(X_i X_{N_{i,1}}) - \sum_{i=1}^n \mathbb{E}(X_i) \mathbb{E}(X_{N_{i,1}}) - \frac{1}{(1-b)} \sum_{i=1}^n \mathbb{E}(X_i).$$

This implies

$$(1-b) \left\{ \sum_{i=1}^n \mathbb{E}(X_i X_{N_{i,1}}) - \sum_{i=1}^n \mathbb{E}(X_i) \mathbb{E}(X_{N_{i,1}}) \right\} - \sum_{i=1}^n \mathbb{E}(X_i) - \tau(1-b) = 0. \quad (3.9)$$

Next, define

$$V_{i,n} := W_n - X_{N_{i,1}}$$

so that $X_{N_{i,1}}$ and $V_{i,n}$ are independent, and X_i and $V_{i,n}$ are independent. From (3.9), we get

$$\begin{aligned} (1-b) \left\{ \sum_{i=1}^n \mathbb{E}(X_i X_{N_{i,1}} \Delta g(V_{i,n})) - \sum_{i=1}^n \mathbb{E}(X_i) \mathbb{E}(X_{N_{i,1}} \Delta g(V_{i,n})) \right\} \\ - \sum_{i=1}^n \mathbb{E}(X_i \Delta g(V_{i,n})) - \tau(1-b) \mathbb{E}(\Delta g(V_{i,n})) = 0. \end{aligned}$$

This is equivalent to

$$\begin{aligned} (1-b) \left\{ \sum_{i=1}^n \mathbb{E} \left[X_i \sum_{j=1}^{X_{N_{i,1}}} \Delta g(V_{i,n}) \right] - \sum_{i=1}^n \mathbb{E}(X_i) \mathbb{E} \left[\sum_{j=1}^{X_{N_{i,1}}} \Delta g(V_{i,n}) \right] \right\} \\ - \sum_{i=1}^n \mathbb{E}(X_i \Delta g(V_{i,n})) - \tau(1-b) \mathbb{E}(\Delta g(V_{i,n})) = 0. \end{aligned} \quad (3.10)$$

Using (3.10) in (3.8), we get

$$\begin{aligned}
\mathbb{E}[\mathcal{A}_Z g(W_n)] &= (1-b) \left\{ \sum_{i=1}^n \mathbb{E}(X_i) \mathbb{E} \left[\sum_{j=1}^{X_{N_{i,1}}} \Delta g(W_{i,n} + j) \right] - \sum_{i=1}^n \mathbb{E} \left[X_i \sum_{j=1}^{X_{N_{i,1}}} \Delta g(W_{i,n} + j) \right] \right\} \\
&\quad + (1-b) \left\{ \sum_{i=1}^n \mathbb{E} \left[X_i \sum_{j=1}^{X_{N_{i,1}}} \Delta g(V_{i,n}) \right] - \sum_{i=1}^n \mathbb{E}(X_i) \mathbb{E} \left[\sum_{j=1}^{X_{N_{i,1}}} \Delta g(V_{i,n}) \right] \right\} \\
&\quad + \sum_{i=1}^n \mathbb{E}[X_i \Delta g(W_n)] - \sum_{i=1}^n \mathbb{E}(X_i \Delta g(V_{i,n})) - \tau(1-b) \mathbb{E}(\Delta g(V_{i,n})) \\
&= (1-b) \left\{ \sum_{i=1}^n \mathbb{E}(X_i) \mathbb{E} \left[\sum_{j=1}^{X_{N_{i,1}}} (\Delta g(W_{i,n} + j) - \Delta g(V_{i,n})) \right] \right. \\
&\quad \left. - \sum_{i=1}^n \mathbb{E} \left[X_i \sum_{j=1}^{X_{N_{i,1}}} (\Delta g(W_{i,n} + j) - \Delta g(V_{i,n})) \right] \right\} \\
&\quad + \sum_{i=1}^n \mathbb{E}[X_i (\Delta g(W_n) - \Delta g(V_{i,n}))] - \tau(1-b) \mathbb{E}(\Delta g(V_{i,n})). \tag{3.11}
\end{aligned}$$

Note also that

$$\begin{aligned}
\Delta g(W_{i,n} + j) - \Delta g(V_{i,n}) &= \Delta g(V_{i,n} + X_{N_{i,2}-N_{i,1}} + j) - \Delta g(V_{i,n}) \\
&= \sum_{k=1}^{X_{N_{i,2}-N_{i,1}}+j-1} \Delta^2 g(V_{i,n} + k). \tag{3.12}
\end{aligned}$$

and

$$\begin{aligned}
\Delta g(W_n) - \Delta g(V_{i,n}) &= \Delta g(V_{i,n} + X_{N_{i,2}}) - \Delta g(V_{i,n}) \\
&= \sum_{k=1}^{X_{N_{i,2}}-1} \Delta^2 g(V_{i,n} + k). \tag{3.13}
\end{aligned}$$

Substituting (3.12) and (3.13) in (3.11), we have

$$\begin{aligned}
\mathbb{E}[\mathcal{A}_Z g(W_n)] &= (1-b) \left\{ \sum_{i=1}^n \mathbb{E}(X_i) \mathbb{E} \left[\sum_{j=1}^{X_{N_{i,1}}} \sum_{k=1}^{X_{N_{i,2}-N_{i,1}}+j-1} \Delta^2 g(V_{i,n} + k) \right] \right. \\
&\quad \left. - \sum_{i=1}^n \mathbb{E} \left[X_i \sum_{j=1}^{X_{N_{i,1}}} \sum_{k=1}^{X_{N_{i,2}-N_{i,1}}+j-1} \Delta^2 g(V_{i,n} + k) \right] \right\}
\end{aligned}$$

$$+ \sum_{i=1}^n \mathbb{E} \left[X_i \sum_{j=1}^{X_{N_{i,2}}-1} \Delta^2 g(V_{i,n} + j) \right] - \tau(1-b) \mathbb{E}(\Delta g(V_{i,n})). \quad (3.14)$$

Consider first

$$\begin{aligned} \mathbb{E} \left[X_i \sum_{j=1}^{X_{N_{i,2}}-1} \Delta^2 g(V_{i,n} + j) \right] &= \mathbb{E} \left\{ \mathbb{E} \left[X_i \sum_{j=1}^{X_{N_{i,2}}-1} \Delta^2 g(V_{i,n} + j) \middle| X_{N_{i,2}} \right] \right\} \\ &= \mathbb{E} \left\{ \mathbb{E}(X_i | X_{N_{i,2}}) \mathbb{E} \left[\sum_{j=1}^{X_{N_{i,2}}-1} \Delta^2 g(W_n - X_{N_{i,2}} + j) \middle| X_{N_{i,2}} \right] \right\}, \end{aligned} \quad (3.15)$$

since X_i and $V_{i,n}$ are independent given $X_{N_{i,2}}$. Observe that

$$\left| \mathbb{E} \left[\sum_{j=1}^{X_{N_{i,2}}-1} \Delta^2 g(W_n - X_{N_{i,2}} + j) \middle| X_{N_{i,2}} = n_{i,2} \right] \right| \leq \|\Delta g\| |n_{i,2} - 1| D(W_n | X_{N_{i,2}} = n_{i,2}). \quad (3.16)$$

Using (3.16) in (3.15), we have

$$\begin{aligned} \left| \mathbb{E} \left[X_i \sum_{j=1}^{X_{N_{i,2}}-1} \Delta^2 g(V_{i,n} + j) \right] \right| &\leq \|\Delta g\| \mathbb{E}[X_i | X_{N_{i,2}} - 1 | D(W_n | X_{N_{i,2}})] \\ &= \|\Delta g\| \mathbb{E}[X_i (X_{N_{i,2}} - 1) D(W_n | X_{N_{i,2}})], \end{aligned} \quad (3.17)$$

since $X_i X_{N_{i,2}} \geq X_i \implies X_i (X_{N_{i,2}} - 1) \geq 0$.

Consider next the following expression from (3.14)

$$\mathbb{E} \left[\sum_{j=1}^{X_{N_{i,1}}} \sum_{k=1}^{X_{N_{i,2}}-X_{N_{i,1}}+j-1} \Delta^2 g(V_{i,n} + k) \right] = \mathbb{E} \left\{ \mathbb{E} \left[\sum_{j=1}^{X_{N_{i,1}}} \sum_{k=1}^{X_{N_{i,2}}-X_{N_{i,1}}+j-1} \Delta^2 g(W_n - X_{N_{i,2}} + k) \middle| X_{N_{i,1}}, X_{N_{i,2}} \right] \right\}. \quad (3.18)$$

Then

$$\begin{aligned} \left| \mathbb{E} \left[\sum_{j=1}^{X_{N_{i,1}}} \sum_{k=1}^{X_{N_{i,2}}-X_{N_{i,1}}+j-1} \Delta^2 g(W_n - X_{N_{i,2}} + k) \middle| X_{N_{i,1}} = n_{i,1}, X_{N_{i,2}} = n_{i,2} \right] \right| \\ \leq \frac{\|\Delta g\|}{2} n_{i,1} |2n_{i,2} - n_{i,1} - 1| D(W_n | X_{N_{i,1}} = n_{i,1}, X_{N_{i,2}} = n_{i,2}). \end{aligned} \quad (3.19)$$

Using (3.19) in (3.18), we get

$$\begin{aligned} \left| \mathbb{E} \left[\sum_{j=1}^{X_{N_{i,1}}} \sum_{k=1}^{X_{N_{i,2}} - N_{i,1} + j - 1} \Delta^2 g(V_{i,n} + k) \right] \right| &\leq \frac{\|\Delta g\|}{2} \mathbb{E}[X_{N_{i,1}} |2X_{N_{i,2}} - X_{N_{i,1}} - 1| D(W_n | X_{N_{i,1}}, X_{N_{i,2}})] \\ &= \frac{\|\Delta g\|}{2} \mathbb{E}[X_{N_{i,1}} (2X_{N_{i,2}} - X_{N_{i,1}} - 1) D(W_n | X_{N_{i,1}}, X_{N_{i,2}})], \end{aligned} \quad (3.20)$$

since $X_{N_{i,2}} X_{N_{i,1}} - X_{N_{i,1}}^2 \geq 0$ and $X_{N_{i,2}} X_{N_{i,1}} - X_{N_{i,1}} \geq 0$ which imply $X_{N_{i,1}} (2X_{N_{i,2}} - X_{N_{i,1}} - 1) \geq 0$. Similarly,

$$\left| \mathbb{E} \left[X_i \sum_{j=1}^{X_{N_{i,1}}} \sum_{k=1}^{X_{N_{i,2}} - N_{i,1} + j - 1} \Delta^2 g(V_{i,n} + k) \right] \right| \leq \frac{\|\Delta g\|}{2} \mathbb{E}[X_i X_{N_{i,1}} (2X_{N_{i,2}} - X_{N_{i,1}} - 1) D(W_n | X_{N_{i,1}}, X_{N_{i,2}})]. \quad (3.21)$$

Finally, using (3.17), (3.20) and (3.21) in (3.14), the proof follows. \square

Remarks 3.1. (i) For $n \geq 1$, we can use (3.4) to obtain the following crude upper bound for $d_{TV}(W_n, Z)$.

$$d_{TV}(W_n, Z) \leq (2|1 - b|\|g\| + \|\Delta g\|) \sum_{i=1}^n \mathbb{E}(X_i). \quad (3.22)$$

Note however that for $n \geq 6$, the bound given in (3.3) would be better than the one given in (3.22).

(ii) Assume $D(W_n | X_{N_{i,2}}) \leq c_i(n)$ then $D(W_n | X_{N_{i,1}}, X_{N_{i,2}}) \leq c_i(n)$. Therefore, the bound (3.3) becomes

$$\begin{aligned} d_{TV}(W_n, Z) &\leq \|\Delta g\| \left\{ \sum_{i=1}^n c_i(n) \left[\frac{|1 - b|}{2} \left[\mathbb{E}(X_i) \mathbb{E}[X_{N_{i,1}} (2X_{N_{i,2}} - X_{N_{i,1}} - 1)] \right. \right. \right. \\ &\quad \left. \left. \left. + \mathbb{E}[X_i X_{N_{i,1}} (2X_{N_{i,2}} - X_{N_{i,1}} - 1)] + \mathbb{E}[X_i (X_{N_{i,2}} - 1)] \right] + |\tau(1 - b)| \right\} \\ &=: d_1(n). \end{aligned} \quad (3.23)$$

Furthermore, let us denote $\mathcal{L}(W_{i,n}^*) = \mathcal{L}(W_n | X_{N_{i,2}})$ and $Z_e = \{X_{2m} \mid m \in \{1, \dots, \lfloor n/2 \rfloor\}\} = (X_2, X_4, \dots, X_{2\lfloor n/2 \rfloor})$. Then, $\mathcal{L}(W_{i,n}^* | Z_e = z_e)$ can be written as sum of independent rvs, say

$X_j^{(z_e)}$, for $j = 1, 2, \dots, n_{z_e}$. Therefore, using (5.11) of Röllin [20], we have

$$D(W_{i,n}^*) \leq \mathbb{E}[\mathbb{E}[D(W_{i,n}^*)|Z_e]] \leq \mathbb{E}\left[\frac{2}{V_{i,Z_e}^{1/2}}\right], \quad (3.24)$$

where

$$V_{i,z_e} = \sum_{j=1}^{n_{z_e}} \min\left\{\frac{1}{2}, 1 - D(X_j^{(z_e)})\right\}. \quad (3.25)$$

On the other hand, let

$$m^* = \begin{cases} \lfloor n/2 \rfloor + 1, & \text{if } n \text{ is odd} \\ n/2, & \text{if } n \text{ is even} \end{cases} \quad \text{and} \quad Z_o = \{X_{2m-1} \mid m \in \{1, \dots, m^*\}\}. \quad (3.26)$$

Then, applying the similar argument as above, we get

$$D(W_{i,n}^*) \leq \mathbb{E}[\mathbb{E}[D(W_{i,n}^*)|Z_o]] \leq \mathbb{E}\left[\frac{2}{V_{i,Z_o}^{1/2}}\right], \quad (3.27)$$

where V_{i,z_o} is defined in a similar way as V_{i,z_e} . Hence, from (3.24) and (3.27), we have

$$D(W_{i,n}^*) \leq \min\left\{\mathbb{E}\left[\frac{2}{V_{i,Z_o}^{1/2}}\right], \mathbb{E}\left[\frac{2}{V_{i,Z_e}^{1/2}}\right]\right\} = c_i(n).$$

Note that $c_i(n) = O(n^{-1/2})$ in general. For more details, we refer the reader to Section 5.3 and Section 5.4 of Röllin [20].

(iii) The bound given in Theorem 3.1 can also be used for the case of matching the first two moments (i.e., $\tau = 0$), whenever that is possible with the approximating distribution.

(iv) From (3.8), it can be easily verified that in the case of first moment matching, we have

$$d_{TV}(W_n, Z) \leq \|\Delta g\| \left\{ |1 - b| \sum_{i=1}^n [\mathbb{E}(X_i)\mathbb{E}(X_{N_{i,1}}) + \mathbb{E}(X_i X_{N_{i,1}})] + \sum_{i=1}^n \mathbb{E}(X_i) \right\} =: d_2(n).$$

and then we have $d_{TV}(W_n, Z) \leq \min\{d_1(n), d_2(n)\}$, where $d_1(n)$ is defined in (3.23).

(v) Observe that if $\tau = 0$ then the bound given in (3.3) is of optimal order $O(n^{-1/2})$ and is comparable with the existing bounds (Theorems 3.1, 3.3, and 3.4 for Poisson, negative binomial, and binomial, respectively) given by Čekanavičius and Vellaisamy [25] with the relaxation of the conditions (3.1) – (3.3).

Next, we give a bound for any rv $X \in \mathcal{P}_2$ in the following corollary.

Corollary 3.1. *Assume that the conditions of Theorem 3.1 hold. Then, for any $X \in \mathcal{P}_2$ and $n \geq 6$,*

$$d_{TV}(W_n, X) \leq \min \left\{ 1, \frac{1}{a} \right\} \left\{ \sum_{i=1}^n c_i(n) \left[\frac{|1-b|}{2} \left[\mathbb{E}(X_i) \mathbb{E}[X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)] \right. \right. \right. \\ \left. \left. \left. + \mathbb{E}[X_i X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)] \right] + \mathbb{E}[X_i(X_{N_{i,2}} - 1)] \right] + |\tau(1-b)| \right\}. \quad (3.28)$$

Example 3.1. *Assume that the conditions of Corollary 3.1 hold. Moreover, let $Y \sim \text{Poi}(\lambda)$, the Poisson rv, so that $a = \lambda$ and $b = 0$. Then, for $n \geq 6$,*

$$d_{TV}(W_n, Y) \leq \min \left\{ 1, \frac{1}{\lambda} \right\} \left\{ \sum_{i=1}^n c_i(n) \left[\frac{1}{2} \left[\mathbb{E}(X_i) \mathbb{E}[X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)] \right. \right. \right. \\ \left. \left. \left. + \mathbb{E}[X_i X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)] \right] + \mathbb{E}[X_i(X_{N_{i,2}} - 1)] \right] + |\bar{\tau}_1| \right\},$$

where $\bar{\tau}_1 = \text{Var}(W_n) - \lambda$.

Example 3.2. *Assume that the conditions of Corollary 3.1 hold. Moreover, let $U \sim \text{NB}(\alpha, p)$, the negative binomial rv, so that $a = \alpha(1-p)$ and $b = 1-p$. Then, for $n \geq 6$,*

$$d_{TV}(W_n, U) \leq \min \left\{ 1, \frac{1}{\alpha(1-p)} \right\} \left\{ \sum_{i=1}^n c_i(n) \left[\frac{p}{2} \left[\mathbb{E}(X_i) \mathbb{E}[X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)] \right. \right. \right. \\ \left. \left. \left. + \mathbb{E}[X_i X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)] \right] + \mathbb{E}[X_i(X_{N_{i,2}} - 1)] \right] + |\bar{\tau}_2 p| \right\},$$

where $\bar{\tau}_2 = \text{Var}(W_n) - \alpha(1-p)/p^2$.

4 Applications to Runs

The distribution of runs and patterns has been applied successfully in many areas such as reliability theory, machine maintenance, quality control, and statistical testing, among many others. Also, it is not tractable if the underlying setup is arising from non-identical trials. So, the approximation of the runs has been studied by several researchers which includes, among others, Fu and Johnson [9], Godbole and Schaffner [10], Kumar and Upadhye [13, 23], Vellaisamy [26], and Wang and Xia [27]. In this section, we mainly focus on 2-runs and (k_1, k_2) -runs, however, the results can also be extended to other types of runs.

4.1 2-runs

We consider here the setup similar to the one discussed in Chapter 5 of Balakrishnan and Koutras [1, p. 166] for 2-runs. Let $\eta_1, \eta_2, \dots, \eta_{n+1}$ be a sequence of independent Bernoulli trials with success probability $\mathbb{P}(\eta_i = 1) = p_i = 1 - \mathbb{P}(\eta_i = 0)$, for $i = 1, 2, \dots, n + 1$. Assume that $p_i \leq 1/2$, for all i , and

$$R_n := \sum_{i=1}^n X_i, \quad (4.1)$$

where $X_i = \eta_i \eta_{i+1}$, $1 \leq i \leq n$, is a sequence of 1-dependent rvs. Observe that R_n counts the number of overlapping success runs of length 2 in $n + 1$ trials. It is easy to see that $\mathbb{E}X_i = \mathbb{P}(X_i = 1) = p_i p_{i+1} := a_1(p_i)$. Similarly, $\mathbb{E}(X_i X_{i+1}) = p_i p_{i+1} p_{i+2} := a_2(p_i)$ and $\mathbb{E}(X_i X_{i+1} X_{i+2}) = p_i p_{i+1} p_{i+2} p_{i+3} := a_3(p_i)$. Now, consider the first term in (3.3). Then

$$\begin{aligned} \mathbb{E}(X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)) &= \mathbb{E}(X_{i-1} + X_i + X_{i+1})^2 + \mathbb{E}[(X_{i-1} + X_i + X_{i+1})(2X_{i-2} + 2X_{i+2} - 1)] \\ &= 2 \sum_{j=i-2}^{i+1} a_2(p_j) + 2[a_1(p_{i-1})a_1(p_{i+1}) + a_1(p_{i-2})(a_1(p_i) + a_1(p_{i+1})) \\ &\quad + a_1(p_{i+2})(a_1(p_{i-1}) + a_1(p_i))] := \bar{a}_1(p_i). \end{aligned} \quad (4.2)$$

Similarly,

$$\begin{aligned} \mathbb{E}(X_i X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)) &= \mathbb{E}(X_i(X_{i-1} + X_i + X_{i+1})^2) \\ &\quad + \mathbb{E}[X_i(X_{i-1} + X_i + X_{i+1})(2X_{i-2} + 2X_{i+2} - 1)] \\ &= 2a_1(p_i)(a_1(p_{i-2}) + a_1(p_{i+2})) + 2a_2(p_{i-1})(1 + a_1(p_{i+2})) \\ &\quad + 2a_2(p_i)(1 + a_1(p_{i-2})) + 2 \sum_{j=i-2}^i a_3(p_j) =: \bar{a}_2(p_i). \end{aligned} \quad (4.3)$$

and

$$\begin{aligned} \mathbb{E}(X_i(X_{N_{i,2}} - 1)) &= \mathbb{E}\left(X_i \sum_{j=i-2}^{i+2} X_j\right) - \mathbb{E}(X_i) \\ &= a_1(p_i) \sum_{|j-i|=2} a_1(p_j) + \sum_{j=i-1}^i a_2(p_j) =: \bar{a}_3(p_i). \end{aligned} \quad (4.4)$$

Next, recall from Remarks 3.1 (ii) with $W_n = R_n$ and $W_{i,n}^* = R_{i,n}^*$, $\mathcal{L}(R_{i,n}^* | Z_e = z_e)$ can be written as sum of independent rvs, say $X_j^{(z_e)}$, for $j \in \{1, 2, \dots, n_{z_e}\} \cap \{\ell : |\ell - i| > 2\} =: \mathcal{C}_i$, for $i = 1, 2, \dots, n$. Note that $n_{z_e} = m^*$ defined in (3.26) and $X_j^{(z_e)} = X_{2j-1}$ depends only on X_{2j-2} ($2j \notin \{2, i+4\}$) and X_{2j} ($2j \neq i-2$, $j \leq \lfloor n/2 \rfloor$), $j \in \mathcal{C}_i$, for all values of z_e . So, for simplicity, let us write

$$X_j^{(z_e)} = X_{2j-1}^{(x_{2j-2}, x_{2j})}, \quad j \in \mathcal{C}_i,$$

where x_{2j-2} and x_{2j} are corresponding values of the rvs X_{2j-2} and X_{2j} , respectively. Note that we use the same notation $D(X_{2j-1}^{(x_{2j-2}, x_{2j})})$, for $2j-1 \in \{1, i-3, i+3, m^*\}$, while X_{2j-1} depends either X_{2j-2} or X_{2j} , not both. Therefore, from (3.25), we have

$$V_{i, z_e} = \sum_{j \in \mathcal{C}_i} \min \left\{ \frac{1}{2}, 1 - D \left(X_{2j-1}^{(x_{2j-2}, x_{2j})} \right) \right\}.$$

Note that $D(X_{2j-1}^{(1,1)}) = D(X_{2j-1}^{(1,0)}) = D(X_{2j-1}^{(0,1)}) = D(X_{2j-1}^{(0,0)}) = \frac{1}{2}$, for all $j \in \mathcal{C}_i$, except when $2j-1 \in \{1, i-3, i+3, m^*\}$. For $2j-1 \in \{1, i-3, i+3, m^*\}$, we have $D(X_{2j-1}^{(x_{2j-2}, x_{2j})}) = p_{2j-1}$ or $p_{2j} \leq 1/2 \implies 1 - D(X_{2j-1}^{(x_{2j-2}, x_{2j})}) \geq 1/2$. Hence,

$$V_{i, z_e} = \sum_{j \in \mathcal{C}_i} \min \left\{ \frac{1}{2}, 1 - D \left(X_{2j-1}^{(x_{2j-2}, x_{2j})} \right) \right\} \geq \frac{1}{2}(m^* - 3), \quad \text{for all values of } z_e.$$

Next, from (3.24), we have

$$D(R_{i,n}^*) \leq \mathbb{E} \left[\frac{2}{V_{i, Z_e}^{1/2}} \right] \leq 4(m^* - 3)^{-1/2}, \quad \text{for all } i.$$

Similarly,

$$D(R_{i,n}^*) \leq \mathbb{E} \left[\frac{2}{V_{i, Z_o}^{1/2}} \right] \leq 4(\lfloor n/2 \rfloor - 3)^{-1/2}, \quad \text{for all } i.$$

Therefore,

$$\bar{c}_i(n) = \bar{c}(n) = 4 \min \left\{ (m^* - 3)^{-1/2}, (\lfloor n/2 \rfloor - 3)^{-1/2} \right\} \leq 4(m^* - 3)^{-1/2}, \quad \text{for all } i. \quad (4.5)$$

Hence, using (4.2), (4.3), (4.4), (4.5), and Theorem 3.1 and Remarks 3.1 (ii), we obtain the following theorem.

Theorem 4.1. *Let $Z \in \mathcal{P}_1$ and R_n be defined as in (4.1). Assume that $\mathbb{E}(Z) = \mathbb{E}(R_n)$, and $\tau =$*

$\text{Var}(R_n) - \text{Var}(Z)$. Then, for $n \geq 8$ and $p_i \leq 1/2$,

$$d_{TV}(R_n, Z) \leq \|\Delta g\| \left\{ \bar{c}(n) \sum_{i=1}^n \left[\frac{|1-b|}{2} [a_1(p_i)\bar{a}_1(p_i) + \bar{a}_2(p_i)] + \bar{a}_3(p_i) \right] + |\tau(1-b)| \right\}. \quad (4.6)$$

Remark 4.1. Note that $\|\Delta g\|$ is of $O(n^{-1})$ in general, and hence, if $\text{Var}(Z) = \text{Var}(R_n)$ then the above bound become of order $O(n^{-1/2})$ and is comparable with the bounds given by Barbour and Xia [4], Brown and Xia [5], Daly et al. [6], and Wang and Xia [27]. In fact, if $p_i = p$, for all $1 \leq i \leq n+1$, then $a(p) = p^2$, $\bar{a}_1(p) = 8p^3 + 10p^4$, $\bar{a}_2(p) = 4p^3 + 10p^4 + 4p^5$, and $\bar{a}_3(p) = 2p^3 + 4p^4$, for all $1 \leq i \leq n+1$, and hence, we have from (4.6),

$$d_{TV}(R_n, Z) \leq n\|\Delta g\|\bar{c}(n) \left[\frac{|1-b|}{2} [4p^3 + 10p^4 + 12p^5 + 10p^6] + 2(p^3 + p^4) \right]. \quad (4.7)$$

Now, let $Z \sim \text{NB}(\alpha, \bar{p})$, the negative binomial distribution with parameter α and \bar{p} . Then, $b = 1 - \bar{p}$ with $\bar{p} = 1/(1 + 2p - 3p^3)$ and $\|\Delta g\| \leq \frac{1}{\alpha(1-\bar{p})} = \frac{1+2p-3p^2}{np^2}$, where α and \bar{p} are obtained from first two moments matching condition, and hence, for $n \geq 8$ and $p \leq 1/2$, we get

$$d_{TV}(R_n, \text{NB}(\alpha, \bar{p})) \leq \frac{4p}{(m^* - 3)^{1/2}} [4 + 11p + 4p^2 - p^3]. \quad (4.8)$$

Also, from Theorem 4.2 of Brown and Xia [5], for $n \geq 2$ and $p < 2/3$, we have

$$d_{TV}(R_n, \text{NB}(\alpha, \bar{p})) \leq \frac{32.2p}{\sqrt{(n-1)(1-p)^3}}. \quad (4.9)$$

For $n \geq 8$, we compare our bound with the one given in (4.9), due to Brown and Xia [5]. Some numerical computations are given in the following table.

Table 1: Comparison of bounds.

n	p	From (4.8)	From (4.9)	n	From (4.8)	From (4.9)	n	From (4.8)	From (4.9)
	0.05	0.344694	0.398900		0.303992	0.354924		0.263265	0.322880
20	0.07	0.506847	0.576571	25	0.446997	0.513008	30	0.387111	0.466692
	0.09	0.683285	0.765878		0.602601	0.681445		0.521867	0.619922
	0.11	0.618205	0.723476		0.561012	0.675509		0.493157	0.602650
35	0.13	0.763728	0.884669	40	0.693072	0.826015	50	0.609244	0.736923
	0.15	0.919907	1.057010		0.834802	0.986930		0.733832	0.880482

It is clear from the above table that our bound given in (4.8) is better than the bound given in (4.9), which is due to Brown and Xia [5].

4.2 (k_1, k_2) -runs

In this subsection, we consider the setup similar to Huang and Tsai [12] and Vellaisamy [26]. Let I_1, I_2, \dots be a sequence of independent Bernoulli trials. Here, we consider $I_1, I_2, \dots, I_{(n+1)(k_1+k_2-1)}$ with success probability $\mathbb{P}(I_i = 1) = p_i = 1 - \mathbb{P}(I_i = 0)$, for $i = 1, 2, \dots, (n+1)(k_1 + k_2 - 1)$. Define $m := k_1 + k_2 - 1$ and

$$Y_j := (1 - I_j) \dots (1 - I_{j+k_1-1}) I_{j+k_1} \dots I_{j+k_1+k_2-1}, \quad j = 1, 2, \dots, nm. \quad (4.10)$$

Note that Y_1, Y_2, \dots, Y_{nm} is a sequence of m -dependent rvs. Now, let us also define

$$X_i := \sum_{j=(i-1)m+1}^{im} Y_j, \quad \text{for } i = 1, 2, \dots, n.$$

Then X_1, X_2, \dots, X_n become a sequence of 1-dependent rvs, that is, we reduced m -dependent to 1-dependent sequence of rvs. From (4.10), it is clear that Y_i , for $i = 1, 2, \dots, nm$, are Bernoulli rvs and if $Y_i = 1$ then $Y_j = 0$, for all j such that $|j - i| \leq m$ and $j \neq i$. Therefore, $X_i, i = 1, 2, \dots, n$, are also Bernoulli rvs. Next, let

$$R'_n = \sum_{i=1}^{nm} Y_i = \sum_{i=1}^n X_i, \quad (4.11)$$

the sum of the corresponding 1-dependent rvs (X_i 's). The distribution of R'_n is called the distribution of (k_1, k_2) -runs or modified distribution of order k or distribution of order (k_1, k_2) . For more details, we refer the reader to Balakrishnan and Koutras [1], Huang and Tsai [12], Upadhye and Kumar [23], Vellaisamy [26] and reference therein.

Next, note that

$$\mathbb{E}(Y_j) = \mathbb{P}(Y_j = 1) = (1 - p_j) \dots (1 - p_{j+k_1-1}) p_{j+k_1} \dots p_{j+k_1+k_2-1} =: a(p_j), \quad \text{for } j = 1, 2, \dots, nm,$$

and hence

$$\mathbb{E}X_i = \sum_{j=(i-1)m+1}^{im} \mathbb{E}Y_j = \sum_{j=(i-1)m+1}^{im} a(p_j) =: a^*(p_i), \quad \text{for } i = 1, 2, \dots, n. \quad (4.12)$$

Also,

$$\mathbb{E}(X_i X_{i+1}) = \sum_{\ell_1=(i-1)m+1}^{im-1} a(p_{\ell_1}) \sum_{\ell_2=\ell_1+m+1}^{(i+1)m} a(p_{\ell_2}) + \sum_{\ell_1=im+2}^{(i+1)m} a(p_{\ell_1}) \sum_{\ell_2=(i-1)m+1}^{\ell_1-m-1} a(p_{\ell_2}) =: a^*(p_i p_{i+1}). \quad (4.13)$$

and

$$\begin{aligned} \mathbb{E}(X_i X_{i+1} X_{i+2}) &= \sum_{\ell_1=(i-1)m+1}^{im-1} a(p_{\ell_1}) \sum_{\ell_2=\ell_1+m+1}^{(i+1)m-1} a(p_{\ell_2}) \sum_{\ell_3=\ell_2+m+1}^{(i+2)m} a(p_{\ell_3}) \\ &+ \sum_{\ell_1=im+2}^{(i+1)m-1} a(p_{\ell_1}) \sum_{\ell_2=(i-1)m+1}^{\ell_1-m-1} a(p_{\ell_2}) \sum_{\ell_3=\ell_1+m+1}^{(i+2)m} a(p_{\ell_3}) \\ &+ \sum_{\ell_1=(i+1)m+3}^{(i+2)m} a(p_{\ell_1}) \sum_{\ell_2=im+2}^{\ell_1-m-1} a(p_{\ell_2}) \sum_{\ell_3=(i-1)m+1}^{\ell_2-m-1} a(p_{\ell_3}) \\ &=: a^*(p_i p_{i+1} p_{i+2}). \end{aligned} \quad (4.14)$$

Using the steps similar to (4.2)-(4.4) with (4.12), (4.13) and (4.14), we have

$$\begin{aligned} \mathbb{E}(X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)) &\leq 2 \sum_{j=i-2}^{i+1} a^*(p_j p_{j+1}) + 2[a^*(p_{i-1})a^*(p_{i+1}) \\ &+ a^*(p_{i-2})(a^*(p_i) + a^*(p_{i+1})) + a^*(p_{i+2})(a^*(p_{i-1}) + a^*(p_i))] \\ &:= a_1^*(p_i), \end{aligned} \quad (4.15)$$

$$\begin{aligned} \mathbb{E}(X_i X_{N_{i,1}}(2X_{N_{i,2}} - X_{N_{i,1}} - 1)) &\leq 2a^*(p_i)(a^*(p_{i-2}) + a^*(p_{i+2})) + 2a^*(p_{i-1} p_i)(1 + a^*(p_{i+2})) \\ &+ 2a^*(p_i p_{i+1})(1 + a^*(p_{i-2})) + 2 \sum_{j=i-2}^i a^*(p_j p_{j+1} p_{j+2}) \\ &=: a_2^*(p_i). \end{aligned} \quad (4.16)$$

and

$$\mathbb{E}(X_i(X_{N_{i,2}} - 1)) \leq a^*(p_i) \sum_{|j-i|=2} a^*(p_j) + \sum_{j=i-1}^i a^*(p_j p_{j+1}) =: a_3^*(p_i). \quad (4.17)$$

Next, from Subsection 4.1, following the discussion about Remarks 3.1 (ii), we have

$$V_{i,z_e} = \sum_{j \in \mathcal{C}_i} \min \left\{ \frac{1}{2}, 1 - D \left(X_{2j-1}^{(x_{2j-2}, x_{2j})} \right) \right\}.$$

Note that

$$\begin{aligned} D \left(X_{2j-1}^{(x_{2j-2}, x_{2j})} \right) &= \frac{1}{2} \left[\mathbb{P} \left(X_{2j-1}^{(x_{2j-2}, x_{2j})} = 0 \right) + \left| \mathbb{P} \left(X_{2j-1}^{(x_{2j-2}, x_{2j})} = 0 \right) - \mathbb{P} \left(X_{2j-1}^{(x_{2j-2}, x_{2j})} = 1 \right) \right| \right] \\ &\leq \frac{1}{2} \left[1 + \mathbb{P} \left(X_{2j-1}^{(x_{2j-2}, x_{2j})} = 0 \right) \right] \\ &\leq \frac{1}{2} [1 + \bar{a}(p_{2j-1})], \end{aligned} \tag{4.18}$$

where

$$\bar{a}(p_{2j-1}) = \max_{0 \leq x_{2j-2}, x_{2j} \leq 1} \mathbb{P} \left(X_{2j-1}^{(x_{2j-2}, x_{2j})} = 0 \right). \tag{4.19}$$

Next, using (4.18), we have

$$\frac{1}{V_{i,z_e}^{1/2}} \leq \left(\frac{1}{2} \min \left\{ 1, \sum_{j \in \mathcal{C}_i} (1 - \bar{a}(p_{2j-1})) \right\} \right)^{-1/2}, \text{ for all } z_e$$

Therefore, from (3.24), we have

$$D(R_{i,n}^*) \leq \mathbb{E} \left[\frac{2}{V_{i,Z_e}^{1/2}} \right] \leq 2 \left(\frac{1}{2} \min \left\{ 1, \sum_{j \in \mathcal{C}_i} (1 - \bar{a}(p_{2j-1})) \right\} \right)^{-1/2} =: V_{i,e}^*.$$

Similarly,

$$D(R_{i,n}^*) \leq \mathbb{E} \left[\frac{2}{V_{i,Z_o}^{1/2}} \right] \leq 2 \left(\frac{1}{2} \min \left\{ 1, \sum_{j \in \mathcal{D}_i} (1 - \bar{a}(p_{2j-1})) \right\} \right)^{-1/2} =: V_{i,o}^*,$$

where $\mathcal{D}_i = \{1, 2, \dots, \lfloor n/2 \rfloor\} \cap \{\ell : |\ell - i| > 2\}$. Therefore,

$$c_i^*(n) = \min\{V_{i,e}^*, V_{i,o}^*\}. \tag{4.20}$$

Using (4.12), (4.15)-(4.17), (4.20), Theorem 3.1 and Remarks 3.1 (ii), the following result is established.

Theorem 4.2. *Let $Z \in \mathcal{P}_1$ and R'_n be defined as in (4.11). Assume that $\mathbb{E}(Z) = \mathbb{E}(R'_n)$, and $\tau =$*

$\text{Var}(R'_n) - \text{Var}(Z)$. Then, for $n \geq 3m$, $a(p_{2j-1}) \leq 1/3$ defined in (4.19),

$$d_{TV}(R'_n, Z) \leq \|\Delta g\| \left\{ \sum_{i=1}^n c_i^*(n) \left[\frac{|1-b|}{2} \left[a^*(p_i) a_1^*(p_i) + a_2^*(p_i) \right] + a_3^*(p_i) \right] + |\tau(1-b)| \right\}.$$

Remark 4.2. Note that the above bound is comparable with the existing bounds given by Upadhye and Kumar [23] and order improvement over the bounds given by Barbour et al. [3], Godbole [11], Godbole and Schaffner [10] (with $k_1 = 1$), and Vellaisamy [26]. Also, note that we have used a slightly different form of (k_1, k_2) -runs, that is, we use $I_1, I_2, \dots, I_{(n+1)(k_1+k_2-1)}$ instead of I_1, I_2, \dots, I_n , so that X_1, X_2, \dots, X_n become a sequence of 1-dependent rvs and we can directly apply our result. However, we can also use some other forms and derive the corresponding results.

References

- [1] Balakrishnan, N. and Koutras, M. V. (2002). *Runs and Scans with Applications*. John Wiley, New York.
- [2] Barbour, A. D. and Chen, L. H. Y. (2014). Stein's (magic) method. *Preprint:arXiv:1411.1179*.
- [3] Barbour, A. D., Holst, L. and Janson, S. (1992). *Poisson Approximation*. Oxford Univ. Press, Oxford.
- [4] Barbour, A. D. and Xia, A. (1999). Poisson perturbation. *ESAIM Probab. Statist.* **3**, 131–150.
- [5] Brown, T. C. and Xia, A. (2001). Stein's method and birth-death processes. *Ann. Probab.* **29**, 1373–1403.
- [6] Daly, F., Lefèvre, C. and Utev, S. (2012). Stein's method and stochastic orderings. *Adv. in Appl. Probab.* **44**, 343–372.
- [7] Edwin, T. K. (2014). *Power series distributions and zero-inflated models*. Ph.D. thesis, University of Nairobi.
- [8] Eichelsbacher, P. and Reinert, G. (2008). Stein's method for discrete Gibbs measures. *Ann. Appl. Probab.* **18**, 1588–1618.
- [9] Fu, J. C. and Johnson, B. C. (2009). Approximate probabilities for runs and patterns in I.I.D. and Markov-dependent multistate trials. *Adv. in Appl. Probab.* **41**, 292–308.
- [10] Godbole, A. and Schaffner, A. A. (1993). Improved Poisson approximations for word patterns. *Adv. Appl. Prob.* **25**, 334–347.
- [11] Godbole, A. P. (1993). Approximate reliabilities of m -consecutive- k -out-of- n : failure systems. *Statist. Sinica* **3**, 321–327.

- [12] Huang, W. T. and Tsai, C. S. (1991). On a modified binomial distribution of order k . *Statist. Probab. Lett.* **11**, 125–131.
- [13] Kumar, A. N. and Upadhye, N. S. On discrete Gibbs measure approximation to runs. *arXiv preprint, arXiv:1701.03294*.
- [14] Ley, C., Reinert, G. and Swan, Y. (2017). Stein’s method for comparison of univariate distributions. *Probab. Surv.* **14**, 1–52.
- [15] Lin, Z.-Y. and Liu, W. (2012). m -dependence approximation for dependent random variables. In: *Probability approximations and beyond*, volume 205 of *Lect. Notes Stat.*, Springer, New York, 117–133.
- [16] Noack, A. (1950). A class of random variables with discrete distributions. *Ann. Math. Statistics* **21**, 127–132.
- [17] Panjer, H. H. and Wang, S. (1995). Computational aspects of Sundt’s generalized class. *ASTIN Bull.* **25**, 5–17.
- [18] Patil, G. P. (1962). Certain properties of the generalized power series distribution. *Ann. Inst. Statist. Math.* **14**, 179–182.
- [19] Reinert, G. (2005). *Three general approaches to Stein’s method. An introduction to Stein’s method*. In: A Program in Honour of Charles Stein: Tutorial Lecture Notes. A. D. Barbour, L. H. Y. Chen, eds., 183–221, World Scientific, Singapore.
- [20] Röllin, A. (2008). Symmetric and centered binomial approximation of sums of locally dependent random variables. *Electron. J. Probab.* **13**, no. 24, 756–776.
- [21] Soon, S. Y. T. (1996). Binomial approximation for dependent indicators. *Statist. Sinica* **6**, 703–714.
- [22] Stein, C. (1972). A bound for the error in the normal approximation to the distribution of a sum of dependent random variables. In: *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability (Univ. California, Berkeley, Calif., 1970/1971), Vol. II: Probability theory*, 583–602.
- [23] Upadhye, N. S. and Kumar, A. N. (2018). Pseudo-binomial approximation to (k_1, k_2) -runs. *Statist. Probab. Lett.* **141**, 19–30.
- [24] Upadhye, N. S., Čekanavičius, V. and Vellaisamy, P. (2017). On Stein operators for discrete approximations. *Bernoulli* **23**, 2828–2859.
- [25] Čekanavičius, V. and Vellaisamy, P. (2015). Discrete approximations for sums of m -dependent random variables. *ALEA Lat. Am. J. Probab. Math. Stat.* **12**, 765–792.
- [26] Vellaisamy, P. (2004). Poisson approximation for (k_1, k_2) -events via the Stein–Chen method. *J. Appl. Probab.* **41**, 1081–1092.
- [27] Wang, X. and Xia, A. (2008). On negative approximation to k -runs. *J. Appl. Probab.* **45**, 456–471.