

FLUCTUATION OF LINEAR EIGENVALUE STATISTICS OF REVERSE CIRCULANT MATRICES WITH INDEPENDENT ENTRIES

SHAMBHU NATH MAURYA AND KOUSHIK SAHA

ABSTRACT. In this article, we study the fluctuations of linear eigenvalue statistics of reverse circulant (RC_n) matrices with independent entries which satisfy some moment conditions. We show that $\frac{1}{\sqrt{n}}\text{Tr}\phi(RC_n)$ obey the central limit theorem (CLT) type result, where ϕ is a nice test function.

Keywords : Reverse circulant matrix, linear eigenvalue statistics, weak convergence, central limit theorem, Trace formula, Wick's formula.

1. INTRODUCTION AND MAIN RESULTS

Let A_n be an $n \times n$ matrix with real or complex entries. The linear statistics of eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ of A_n is a function of the form

$$(1) \quad \frac{1}{n} \sum_{k=1}^n \phi(\lambda_k)$$

where ϕ is some fixed function. The function ϕ is known as the test function. The fluctuation of linear statistics of eigenvalues is one of the main area of research in random matrix theory.

First, it was studied by Arharov [4] in 1971 for sample covariance matrices. From then the fluctuations of eigenvalues for various random matrices have been extensively studied by various people. For results on fluctuations of linear eigenvalue statistics of Wigner and sample covariance matrices, see [8], [15], [5], [11], [13]. For band and sparse random matrices, see [3], [7], [9], [14] and for Toeplitz and band Toeplitz matrices, see [6] and [10].

In a recent article [1], the CLT for linear eigenvalue statistics has been established in total variation norm for reverse circulant matrices with Gaussian entries and even degree monomial test functions. In a subsequent article [2], the authors extended their results for independent entries which are smooth functions of Gaussian variables. Here we consider the fluctuation problem for reverse circulant matrices with general entries which are independent and satisfy some moment condition. However we could deal only with the test functions which are even degree monomials or polynomials with even degree terms.

A sequence is said to be an *input sequence* if the matrices are constructed from the given sequence. We consider the input sequence of the form $\{x_i : i \geq 1\}$ and the reverse circulant matrix is defined

Date: February 1, 2022.

The work of Shambhu Nath Maurya is partially supported by UGC Doctoral Fellowship, India and the work of Koushik Saha is partially supported by MATRICS grant of SERB, Department of Science and Technology, Government of India.

as

$$RC_n = \begin{pmatrix} x_1 & x_2 & x_3 & \cdots & x_{n-1} & x_n \\ x_2 & x_3 & x_4 & \cdots & x_n & x_1 \\ x_3 & x_4 & x_5 & \cdots & x_1 & x_2 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ x_n & x_1 & x_2 & \cdots & x_{n-2} & x_{n-1} \end{pmatrix}.$$

For $j = 1, 2, \dots, n-1$, its $(j+1)$ -th row is obtained by giving its j -th row a left circular shift by one positions. Note that the matrix is symmetric and the (i, j) -th element of the matrix is $x_{(i+j-1) \bmod n}$.

Now we consider linear eigenvalue statistics as defined in (1) for RC_n with test function $\phi(x) = x^{2p}$, $p \geq 1$. Therefore

$$\sum_{k=1}^n \phi(\lambda_k) = \sum_{k=1}^n (\lambda_k)^{2p} = \text{Tr}(RC_n)^{2p},$$

where $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of RC_n . We scale and centre $\text{Tr}(RC_n)^{2p}$ to study its fluctuation, and define

$$(2) \quad w_p := \frac{1}{\sqrt{n}} \{ \text{Tr}(RC_n)^{2p} - \mathbb{E}[\text{Tr}(RC_n)^{2p}] \}.$$

Moreover, for a given polynomial

$$Q(x) = \sum_{k=1}^d a_k x^{2k}$$

with degree $2d$, $d \geq 1$ and with even degree terms, we define

$$(3) \quad w_Q := \frac{1}{\sqrt{n}} \{ \text{Tr}(Q(RC_n)) - \mathbb{E}[\text{Tr}(Q(RC_n))] \}.$$

Note that w_Q and w_p depends on n . But we suppress n to keep the notation simple. In our first result, we calculate the covariance between w_p and w_q as $n \rightarrow \infty$.

Theorem 1. *Suppose RC_n is the reverse circulant matrix with independent input sequence $\{\frac{x_i}{\sqrt{n}}\}_{i \geq 1}$ such that*

$$(4) \quad \mathbb{E}(x_i) = 0, \mathbb{E}(x_i^2) = 1, \mathbb{E}(x_i^4) = \mathbb{E}(x_1^4) \quad \forall i \geq 1 \text{ and } \sup_{i \geq 1} \mathbb{E}(|x_i|^k) = \alpha_k < \infty \text{ for } k \geq 3.$$

Then for $p, q \geq 1$,

$$(5) \quad \sigma_{p,q} := \lim_{n \rightarrow \infty} \text{Cov}(w_p, w_q) = \sum_{k=2}^{\min\{p,q\}} c_k g(k) + (\mathbb{E}x_1^4 - 1)c_1,$$

where

$$c_k = \binom{p}{p-k}^2 (p-k)! \binom{q}{q-k}^2 (q-k)!,$$

$$g(k) = \frac{1}{(2k-1)!} \sum_{s=-(k-1)}^{k-1} \sum_{j=0}^{k+s-1} (-1)^j \binom{2k}{j} (k+s-j)^{2k-1} (2 - \mathbf{1}_{\{s=0\}}) k! k!.$$

Moreover, if $p = q$ then we denote $\sigma_{p,q}$ by σ_p^2 .

In our second result, we see the fluctuation of linear eigenvalue statistics of reverse circulant matrices with polynomial test functions with even degree terms.

Theorem 2. Suppose input entry of RC_n is independent sequence $\{\frac{x_i}{\sqrt{n}}\}_{i \geq 1}$ which satisfy (4). Then, as $n \rightarrow \infty$,

$$w_Q \xrightarrow{d} N(0, \sigma_Q^2),$$

where $\sigma_Q^2 = \sum_{\ell=1}^d \sum_{k=1}^d a_\ell a_k \sigma_{\ell,k}$ and $\sigma_{\ell,k}$ is as in (5). In particular, for $Q(x) = x^{2p}$,

$$w_p \xrightarrow{d} N(0, \sigma_p^2),$$

where $\sigma_p^2 = \sigma_{p,p}$.

Remark 3. In the above theorems we have considered the fluctuation of w_p for $p \geq 1$. For $p = 0$,

$$w_0 = \frac{1}{\sqrt{n}} \{ \text{Tr}(I) - \mathbb{E}[\text{Tr}(I)] \} = \frac{1}{\sqrt{n}} [n - n] = 0$$

and hence it has no fluctuation. So we ignore this case, that is, for $p = 0$.

In Section 2 we prove Theorem 1. We derived trace formula and state some results which will be used to prove Theorem 1. In Section 3 we prove Theorem 2. We use method of moments and Wick's formula to prove Theorem 2.

2. PROOF OF THEOREM 1

We first define some notation which will be used in the proof of Theorem 2.

$$(6) \quad \begin{aligned} A_{2p} &= \{(i_1, \dots, i_{2p}) \in \mathbb{N}^{2p} : \sum_{k=1}^{2p} (-1)^k i_k = 0 \pmod{n}, 1 \leq i_1, \dots, i_{2p} \leq n\}, \\ A'_{2p} &= \{(i_1, \dots, i_{2p}) \in \mathbb{N}^{2p} : \sum_{k=1}^{2p} (-1)^k i_k = 0 \pmod{n}, 1 \leq i_1 \neq i_2 \neq \dots \neq i_{2p} \leq n\}, \\ A_{2p,s} &= \{(i_1, \dots, i_{2p}) \in \mathbb{N}^{2p} : \sum_{k=1}^{2p} (-1)^k i_k = sn, 1 \leq i_1, \dots, i_{2p} \leq n\}, \\ A'_{2p,s} &= \{(i_1, \dots, i_{2p}) \in \mathbb{N}^{2p} : \sum_{k=1}^{2p} (-1)^k i_k = sn, 1 \leq i_1 \neq i_2 \neq \dots \neq i_{2p} \leq n\}. \end{aligned}$$

Now recall the trace formula for reverse circulant matrices from [1]. Let e_1, \dots, e_n be the standard unit vectors in \mathbb{R}^n , i.e., $e_i = (0, \dots, 1, \dots, 0)^t$ (1 in i -th place). Therefore we have

$$(RC_n)e_i = i\text{-th column} = \sum_{i_1=1}^n X_{i_1} e_{i_1-i+1 \pmod{n}},$$

for $i = 1, \dots, n$ (we write $e_0 = e_n$). Repeating the procedure we get

$$(RC_n)^2 e_i = \sum_{i_1, i_2=1}^n X_{i_1} X_{i_2} e_{i_2-i_1+i \pmod{n}},$$

for $i = 1, \dots, n$. Therefore in general we get

$$\begin{aligned} (RC_n)^{2p}e_i &= \sum_{i_1, \dots, i_{2p}=1}^n X_{i_1} \dots X_{i_{2p}} e_{i_{2p}-i_{2p-1} \dots -i_1+i \bmod n}, \\ (RC_n)^{2p+1}e_i &= \sum_{i_1, \dots, i_{2p+1}=1}^n X_{i_1} \dots X_{i_{2p+1}} e_{i_{2p+1}-i_{2p} \dots +i_1-i+1 \bmod n}, \end{aligned}$$

for $i = 1, \dots, n$. Therefore the trace of $(RC_n)^{2p}$ can be written as

$$(7) \quad \text{Tr}[(RC_n)^{2p}] = \sum_{i=1}^n e_i^t (RC_n)^{2p} e_i = n \sum_{A_{2p}} X_{i_1} \dots X_{i_{2p}},$$

where A_{2p} as defined in (6).

The following result will be used in the proof of Theorem 1.

Result 4. Suppose $|A_{2p,s}|$ denotes the cardinality of $A_{2p,s}$. Then

$$|A_{2p,s}| = \sum_{k=0}^{p+s-1} (-1)^k \binom{2p}{k} \binom{(p+s-k)n+p-1}{2p-1}, \quad \text{for } s = -(p-1), \dots, 0, \dots, p-1.$$

For the proof of Result 4, we refer to [1, Lemma 14]. Now for a given vector (i_1, i_2, \dots, i_p) , we define *odd-even pair matched* elements of the vector.

Definition 5. Suppose (i_1, i_2, \dots, i_p) is a vector in \mathbb{N}^p . Two elements i_k, i_ℓ of (i_1, i_2, \dots, i_p) are said to be *odd-even pair matched* if $i_k = i_\ell$ and one of the elements appears at an odd position in (i_1, i_2, \dots, i_p) and other one appears at an even position in (i_1, i_2, \dots, i_p) . For example, in $(1, 1, 3, 4)$, entry 1 is *odd-even pair matched* whereas in $(1, 2, 1, 3)$, entry 1 is not *odd-even pair matched*.

We shall call a vector (i_1, i_2, \dots, i_p) is *odd-even pair matched* if all the entries of (i_1, i_2, \dots, i_p) are *odd-even pair matched*. In other words, each entry of (i_1, i_2, \dots, i_p) appears same number of times in odd and even positions.

Observe that, if $(i_1, i_2, \dots, i_{2p}) \in A_{2p}$ and each entry of $\{i_1, i_2, \dots, i_{2p}\}$ has multiplicity greater than or equal to two, then the maximum number of free variables in $(i_1, i_2, \dots, i_{2p})$ will be p , only when $(i_1, i_2, \dots, i_{2p})$ is *odd-even pair matched*. By free variables we mean that these variables can be chosen freely from their range $\{1, 2, \dots, n\}$. We shall use this observation in the proof of Theorem 1.

Now we prove Theorem 1.

Proof of Theorem 1. Since $E(w_p) = E(w_q) = 0$ therefore we get

$$\begin{aligned} \text{Cov}(w_p, w_q) &= E[w_p w_q] \\ &= \frac{1}{n} \left\{ E[\text{Tr}(RC_n)^{2p} \text{Tr}(RC_n)^{2q}] - E[\text{Tr}(RC_n)^{2p}] E[\text{Tr}(RC_n)^{2q}] \right\}. \end{aligned}$$

Using the trace formula (7), we get

$$E[\text{Tr}(RC_n)^{2p}] = E \left[n \sum_{A_{2p}} \frac{x_{i_1}}{\sqrt{n}} \frac{x_{i_2}}{\sqrt{n}} \dots \frac{x_{i_{2p}}}{\sqrt{n}} \right] = \frac{1}{n^{p-1}} E \left[\sum_{A_{2p}} x_{i_1} x_{i_2} \dots x_{i_{2p}} \right].$$

Therefore

$$(8) \quad \begin{aligned} \text{Cov}(w_p, w_q) &= E[w_p w_q] \\ &= \frac{1}{n^{p+q-1}} \sum_{A_{2p}, A_{2q}} \left\{ E[x_{i_1} x_{i_2} \dots x_{i_{2p}} x_{j_1} x_{j_2} \dots x_{j_{2q}}] - E[x_{i_1} x_{i_2} \dots x_{i_{2p}}] E[x_{j_1} x_{j_2} \dots x_{j_{2q}}] \right\}. \end{aligned}$$

First observe that, if $\{i_1, i_2, \dots, i_{2p}\} \cap \{j_1, j_2, \dots, j_{2q}\} = \emptyset$ then from independence of x_i 's

$$\mathbb{E}[x_{i_1} x_{i_2} \cdots x_{i_{2p}} x_{j_1} x_{j_2} \cdots x_{j_{2q}}] - \mathbb{E}[x_{i_1} x_{i_2} \cdots x_{i_{2p}}] \mathbb{E}[x_{j_1} x_{j_2} \cdots x_{j_{2q}}] = 0.$$

Therefore we can get the non-zero contribution in (8) only when there is at least one cross matching among $\{i_1, i_2, \dots, i_{2p}\}$ and $\{j_1, j_2, \dots, j_{2q}\}$, i.e., $\{i_1, i_2, \dots, i_{2p}\} \cap \{j_1, j_2, \dots, j_{2q}\} \neq \emptyset$.

Now using the above observation, (8) can be written as

$$(9) \quad \lim_{n \rightarrow \infty} \text{Cov}(w_p, w_q) = \lim_{n \rightarrow \infty} \frac{1}{n^{p+q-1}} \sum_{m=1}^{\min\{2p, 2q\}} \sum_{I_m} (\mathbb{E}[x_{i_1} x_{i_2} \cdots x_{i_{2p}} x_{j_1} x_{j_2} \cdots x_{j_{2q}}] - \mathbb{E}[x_{i_1} x_{i_2} \cdots x_{i_{2p}}] \mathbb{E}[x_{j_1} x_{j_2} \cdots x_{j_{2q}}]),$$

where for each $m = 1, 2, \dots, \min\{2p, 2q\}$, I_m is defined as

$$(10) \quad I_m := \{((i_1, \dots, i_{2p}), (j_1, \dots, j_{2q})) \in A_{2p} \times A_{2q} : |\{i_1, \dots, i_{2p}\} \cap \{j_1, \dots, j_{2q}\}| = m\}$$

and $|\{\cdot\}|$ denotes cardinality of the set $\{\cdot\}$ (with multiplicity).

Depending on the number of cross matching elements, there are two following cases:

Case I. Odd number of cross matching: Suppose $(i_1, i_2, \dots, i_{2p}) \in A_{2p}$ and $(j_1, j_2, \dots, j_{2q}) \in A_{2q}$, and $m = 2k - 1$, that is, $|\{i_1, i_2, \dots, i_{2p}\} \cap \{j_1, j_2, \dots, j_{2q}\}| = 2k - 1$ for $k = 1, 2, \dots, \min\{p, q\}$. We shall show that such $(i_1, i_2, \dots, i_{2p}) \in A_{2p}$ and $(j_1, j_2, \dots, j_{2q}) \in A_{2q}$ will have zero contribution in (9).

First suppose $k = 1$, that is, $|\{i_1, i_2, \dots, i_{2p}\} \cap \{j_1, j_2, \dots, j_{2q}\}| = 1$. For $k = 1$, the typical term of (9) will look like

$$\sum_{I_1} (\mathbb{E}[x_{\ell_1} x_{i_2} \cdots x_{i_{2p}} x_{\ell_1} x_{j_2} \cdots x_{j_{2q}}] - \mathbb{E}[x_{\ell_1} x_{i_2} \cdots x_{i_{2p}}] \mathbb{E}[x_{\ell_1} x_{j_2} \cdots x_{j_{2q}}]),$$

where $I_1 \subseteq A_{2p} \times A_{2q}$ and it looks like as $((\ell_1, i_2, \dots, i_{2p}), (\ell_1, j_2, \dots, j_{2q}))$.

Since from (4), we have moment conditions on x_i , that is,

$$\mathbb{E}[x_i] = 0, \mathbb{E}(x_i^2) = 1 \text{ and } \sup_{i \geq 1} \mathbb{E}(|x_i|^k) = \alpha_k < \infty \text{ for } k \geq 3.$$

Therefore there exist $\gamma_1 > 0$, which depends only on p, q , such that

$$(11) \quad \sum_{I_1} (\mathbb{E}[x_{\ell_1} x_{i_2} \cdots x_{i_{2p}} x_{\ell_1} x_{j_2} \cdots x_{j_{2q}}] - \mathbb{E}[x_{\ell_1} x_{i_2} \cdots x_{i_{2p}}] \mathbb{E}[x_{\ell_1} x_{j_2} \cdots x_{j_{2q}}]) \leq \gamma_1 |I_1|,$$

where

$$I_1' = \{((\ell_1, i_2, \dots, i_{2p}), (\ell_1, j_2, \dots, j_{2q})) \in I_1 : \text{each elements in } \{\ell_1, i_2, \dots, i_{2p}\} \cup \{\ell_1, j_2, \dots, j_{2q}\} \text{ has multiplicity greater than or equal to two}\}.$$

Now we calculate the cardinality of I_1' . First recall the observation from Definition 5, the maximum number of free variable in $(\ell_1, i_2, \dots, i_{2p})$ will be p , when $(\ell_1, i_2, \dots, i_{2p})$ is *odd-even pair matched*. Once we have chosen $(\ell_1, i_2, \dots, i_{2p})$, the entry ℓ_1 of $(\ell_1, j_2, \dots, j_{2q})$ is also chosen, therefore the maximum possible number of free variables from $(\ell_1, j_2, \dots, j_{2q})$ will be $q - 2$, where one extra (-1) is coming due to the constraint, $-\ell_1 + \sum_{s=2}^{2q} (-1)^s j_s = 0 \pmod{n}$ and hence $|I_1'| = O(n^{p+q-2})$. Therefore from (11),

$$\frac{1}{n^{p+q-1}} \sum_{I_1} (\mathbb{E}[x_{i_1} \cdots x_{i_{2p}} x_{j_1} \cdots x_{j_{2q}}] - \mathbb{E}[x_{i_1} \cdots x_{i_{2p}}] \mathbb{E}[x_{j_1} \cdots x_{j_{2q}}]) = o(1).$$

Now for $k \geq 2$, I_{2k-1} looks like as $((\ell_1, \dots, \ell_{2k-1}, i_{2k}, \dots, i_{2p}), (\ell_1, \dots, \ell_{2k-1}, j_{2k}, \dots, j_{2q}))$.

As we have done for $k = 1$, from (4), there exist $\gamma_{2k-1} > 0$, which depends only on p, q , such that the typical term of (9) will be

$$(12) \quad \left| \sum_{I_{2k-1}} \left(\mathbb{E}[x_{\ell_1} \cdots x_{\ell_{2k-1}} x_{i_{2k}} \cdots x_{i_{2p}} x_{\ell_1} \cdots x_{\ell_{2k-1}} x_{j_{2k}} \cdots x_{j_{2q}}] \right. \right. \\ \left. \left. - \mathbb{E}[x_{\ell_1} \cdots x_{\ell_{2k-1}} x_{i_{2k}} \cdots x_{i_{2p}}] \mathbb{E}[x_{\ell_1} \cdots x_{\ell_{2k-1}} x_{j_{2k}} \cdots x_{j_{2q}}] \right) \right| \\ \leq \gamma_{2k-1} |I'_{2k-1}|,$$

where

$$I'_{2k-1} = \{((\ell_1, \dots, \ell_{2k-1}, i_{2k}, \dots, i_{2p}), (\ell_1, \dots, \ell_{2k-1}, j_{2k}, \dots, j_{2q})) \in I_{2k-1} : \text{each} \\ \text{elements in } \{\ell_1, \dots, \ell_{2k-1}, i_{2k}, \dots, i_{2p}\} \cup \{\ell_1, \dots, \ell_{2k-1}, j_{2k}, \dots, j_{2q}\} \\ \text{has multiplicity greater than or equal to two}\}.$$

Now we calculate the cardinality of I'_{2k-1} . First observe that we shall get maximum number of free variables in I'_{2k-1} , if the following conditions hold:

- (i) Each elements of $\{\ell_1, \ell_2, \dots, \ell_{2k-1}\}$ are distinct.
- (ii) There exist $i^* \in (i_{2k}, i_{2k+1}, \dots, i_{2p})$ such that i^* equal to ℓ_s for $s = 1, 2, \dots, 2k-1$. Without loss of generality, we suppose i^* appears at even position and $\ell_s = \ell_{2k-1}$. That is, $i^* = \ell_{2k-1}$. Similarly, there exist $j^* \in (j_{2k}, j_{2k+1}, \dots, j_{2p})$ at odd position such that $j^* = \ell_{2k-2}$.
- (iii) $\{\ell_1, \ell_2, \dots, \ell_{2k-3}\} \cap \{i_{2k}, \dots, i_{2p}\} \setminus \{i^*\} \cap \{j_{2k}, \dots, j_{2p}\} \setminus \{j^*\} = \emptyset$.
- (iv) Each elements of $\{i_{2k}, i_{2k+1}, \dots, i_{2p}\} \cup \{i^*\}$ and $\{j_{2k}, j_{2k+1}, \dots, j_{2p}\} \cup \{j^*\}$ are *odd-even pair matched*.

Note that, $(\ell_1, \dots, \ell_{2k-1}, i_{2k}, \dots, i_{2p}) \in A_{2p}$ and $(\ell_1, \dots, \ell_{2k-1}, j_{2k}, \dots, j_{2q}) \in A_{2q}$, that is,

$$\sum_{r=1}^{2k-1} (-1)^r \ell_r + \sum_{r=2k}^{2p} (-1)^r i_r = 0 \pmod{n} \quad \text{and} \quad \sum_{r=1}^{2k-1} (-1)^r \ell_r + \sum_{r=2k}^{2q} (-1)^r j_r = 0 \pmod{n}.$$

But each elements of $\{i_{2k}, i_{2k+1}, \dots, i_{2p}\} \cup \{\ell_{2k-1}\}$ and $\{j_{2k}, j_{2k+1}, \dots, j_{2p}\} \cup \{\ell_{2k-2}\}$ are *odd-even pair matched*, therefore the above two constraints will change into new constraints, namely,

$$(13) \quad \sum_{r=1}^{2k-2} (-1)^r \ell_r = 0 \pmod{n} \quad \text{and} \quad \sum_{r=1}^{2k-3} (-1)^r \ell_r - \ell_{2k-1} = 0 \pmod{n}.$$

Hence the maximum number of free entries in I'_{2k-1} are $[(p-k+1) + (q-k+1) + (2k-3) + (-2)] = p+q-3$, where (-2) is coming due to the constraints (13).

Therefore from the above discussion on cardinality of I'_{2k-1} and from (12), we get

$$\frac{1}{n^{p+q-1}} \sum_{I_{2k-1}} \left(\mathbb{E}[x_{\ell_1} \cdots x_{\ell_{2k-1}} x_{i_{2k}} \cdots x_{i_{2p}} x_{\ell_1} \cdots x_{\ell_{2k-1}} x_{j_{2k}} \cdots x_{j_{2q}}] \right. \\ \left. - \mathbb{E}[x_{\ell_1} \cdots x_{\ell_{2k-1}} x_{i_{2k}} \cdots x_{i_{2p}}] \mathbb{E}[x_{\ell_1} \cdots x_{\ell_{2k-1}} x_{j_{2k}} \cdots x_{j_{2q}}] \right) \\ = o(1).$$

Hence for each odd m , we get

$$(14) \quad \lim_{n \rightarrow \infty} \frac{1}{n^{p+q-1}} \sum_{I_m} \left(\mathbb{E}[x_{i_1} \cdots x_{i_{2p}} x_{j_1} \cdots x_{j_{2q}}] - \mathbb{E}[x_{i_1} \cdots x_{i_{2p}}] \mathbb{E}[x_{j_1} \cdots x_{j_{2q}}] \right) = 0.$$

Case II. Even number of cross matching: Suppose $(i_1, i_2, \dots, i_{2p}) \in A_{2p}$ and $(j_1, j_2, \dots, j_{2q}) \in A_{2q}$, and $m = 2k$, that is, $|\{i_1, i_2, \dots, i_{2p}\} \cap \{j_1, j_2, \dots, j_{2q}\}| = 2k$ for some $k = 1, 2, \dots, \min\{p, q\}$.

We shall show that such $(i_1, i_2, \dots, i_{2p}) \in A_{2p}$ and $(j_1, j_2, \dots, j_{2q}) \in A_{2q}$ will have non-zero contribution in (9).

In this case, a typical element of I_{2k} looks like as $((\ell_1, \dots, \ell_{2k}, i_{2k+1}, \dots, i_{2p}), (\ell_1, \dots, \ell_{2k}, j_{2k+1}, \dots, j_{2q}))$. Now we define $H \subseteq I_{2k}$ where I_{2k} is as in (10).

Definition 6. H is a subsets of all $((\ell_1, \dots, \ell_{2k}, i_{2k+1}, \dots, i_{2p}), (\ell_1, \dots, \ell_{2k}, j_{2k+1}, \dots, j_{2q})) \in I_{2k}$ such that

- (i) $\{\ell_1, \ell_2, \dots, \ell_{2k}\} \cap \{i_{2k+1}, i_{2k+2}, \dots, i_{2p}\} \cap \{j_{2k+1}, j_{2k+2}, \dots, j_{2q}\} = \emptyset$,
- (ii) $(i_{2k}, i_{2k+1}, \dots, i_{2p})$ and $(j_{2k}, j_{2k+1}, \dots, j_{2p})$ are *odd-even pair matched*.

By using the similar calculation, as we have done to calculate the number of free entries of I_{2k-1} , we shall get maximum number of free entries in $((\ell_1, \dots, \ell_{2k}, i_{2k+1}, \dots, i_{2p}), (\ell_1, \dots, \ell_{2k}, j_{2k+1}, \dots, j_{2q}))$, when $((\ell_1, \dots, \ell_{2k}, i_{2k+1}, \dots, i_{2p}), (\ell_1, \dots, \ell_{2k}, j_{2k+1}, \dots, j_{2q})) \in H$. Now we calculate the cardinality of H .

Suppose $((\ell_1, \dots, \ell_{2k}, i_{2k+1}, \dots, i_{2p}), (\ell_1, \dots, \ell_{2k}, j_{2k+1}, \dots, j_{2q})) \in H$. Since from (ii) condition of Definition 6, $(i_{2k+1}, i_{2k+2}, \dots, i_{2p})$ and $(j_{2k+1}, j_{2k+2}, \dots, j_{2q})$ are *odd-even pair matched*, therefore $\sum_{r=2k+1}^{2p} (-1)^r i_r = 0 \pmod{n}$ and $\sum_{r=2k+1}^{2q} (-1)^r j_r = 0 \pmod{n}$ are aromatically satisfy, and hence the constraints on $(\ell_1, \dots, \ell_{2k}, i_{2k+1}, \dots, i_{2p})$ and $(\ell_1, \dots, \ell_{2k}, j_{2k+1}, \dots, j_{2q})$ will change into a single constraint, namely,

$$(15) \quad \sum_{r=1}^{2k} (-1)^r \ell_r = 0 \pmod{n}.$$

Now we first deal with $k \geq 2$, later we shall calculate for $k = 1$. For $k = 2, 3, \dots, \min\{p, q\}$, if we assume each elements of $\{\ell_1, \ell_2, \dots, \ell_{2k}\}$ are distinct, then cardinality of H will be $O(n^{p+q-1})$, where (-1) is coming due to the constraint $\sum_{r=1}^{2k} (-1)^r \ell_r = 0 \pmod{n}$. Note that in any other situation, like that any one of the conditions of Definition 6 does not hold or elements of $\{\ell_1, \ell_2, \dots, \ell_{2k}\}$ are not distinct then cardinality of H will be $o(n^{p+q-1})$.

Since each elements of $\{\ell_1, \ell_2, \dots, \ell_{2k}\}$ are distinct, therefore

$$\mathbb{E}[x_{\ell_1} \cdots x_{\ell_{2k}} x_{i_{2k+1}} \cdots x_{i_{2p}}] \mathbb{E}[x_{\ell_1} \cdots x_{\ell_{2k}} x_{j_{2k+1}} \cdots x_{j_{2q}}] = 0.$$

Hence for each $k \geq 2$, we get

$$(16) \quad \begin{aligned} & \lim_{n \rightarrow \infty} \frac{1}{n^{p+q-1}} \sum_{I_{2k}} (\mathbb{E}[x_{i_1} x_{i_2} \cdots x_{i_{2p}} x_{j_1} x_{j_2} \cdots x_{j_{2q}}] - \mathbb{E}[x_{i_1} x_{i_2} \cdots x_{i_{2p}}] \mathbb{E}[x_{j_1} x_{j_2} \cdots x_{j_{2q}}]) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n^{p+q-1}} \sum_{I_{2k}} \mathbb{E}[x_{i_1} x_{i_2} \cdots x_{i_{2p}} x_{j_1} x_{j_2} \cdots x_{j_{2q}}] = O(1). \end{aligned}$$

Now combining both the Cases, (14) and (16), from (9) we get

$$(17) \quad \lim_{n \rightarrow \infty} \text{Cov}(w_p, w_q) = \lim_{n \rightarrow \infty} \frac{1}{n^{p+q-1}} \sum_{k=1}^{\min\{p, q\}} \sum_{I_{2k}} (\mathbb{E}[x_{i_1} x_{i_2} \cdots x_{i_{2p}} x_{j_1} x_{j_2} \cdots x_{j_{2q}}]).$$

Now for fixed $k = 2, 3, \dots, \min\{p, q\}$, we calculate the contribution of each term of (17).

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \frac{1}{n^{p+q-1}} \sum_{I_{2k}} \mathbb{E}[x_{i_1} x_{i_2} \cdots x_{i_{2p}} x_{j_1} x_{j_2} \cdots x_{j_{2q}}] \\
&= \lim_{n \rightarrow \infty} \frac{1}{n^{p+q-1}} \sum_{A'_{2k}} \sum_{A'_{2k}} c_k n^{(p-k)+(q-k)} \mathbb{E}[x_{i_1} x_{i_2} \cdots x_{i_{2k}} x_{j_1} x_{j_2} \cdots x_{j_{2k}}] \\
(18) \quad &= c_k \lim_{n \rightarrow \infty} \frac{1}{n^{2k-1}} \sum_{A'_{2k}} \sum_{A'_{2k}} \mathbb{E}[x_{i_1} x_{i_2} \cdots x_{i_{2k}} x_{j_1} x_{j_2} \cdots x_{j_{2k}}],
\end{aligned}$$

where $c_k = \binom{p}{p-k}^2 (p-k)! \binom{q}{q-k}^2 (q-k)!$. For the odd-even pair matching among $(2p-2k)$ variables from (i_1, \dots, i_{2p}) , first we choose $(p-k)$ odd and $(p-k)$ even position from the available p odd and p even position in $\binom{p}{p-k}^2$ ways. After choosing $(p-k)$ odd, $(p-k)$ even positions and $(p-k)$ free variables in odd positions, the random variables in even positions can permute among themselves (satisfying the condition of odd-even pair matching) in $(p-k)!$ ways. Hence, odd-even pair matching among $(2p-2k)$ variables of (i_1, \dots, i_{2p}) happens in $\binom{p}{p-k}^2 (p-k)! n^{p-k}$ ways. Similarly, odd-even pair matching happens among $(2q-2k)$ variables from $(j_1, j_2, \dots, j_{2q})$ in $\binom{q}{q-k}^2 (q-k)! n^{q-k}$ ways. The rest of the variable $\{i_1, i_2, \dots, i_{2k}\}$ and $\{j_1, j_2, \dots, j_{2k}\}$ will (cross) match completely and both belong to A'_{2k} . Now from (18), we get

$$\begin{aligned}
\lim_{n \rightarrow \infty} \frac{1}{n^{p+q-1}} \sum_{I_{2k}} \mathbb{E}[x_{i_1} \cdots x_{i_{2p}} x_{j_1} \cdots x_{j_{2q}}] &= c_k \lim_{n \rightarrow \infty} \frac{1}{n^{2k-1}} \sum_{A'_{2k}} \sum_{A'_{2k}} \mathbb{E}[x_{i_1} \cdots x_{i_{2k}} x_{j_1} \cdots x_{j_{2k}}] \\
&= c_k \lim_{n \rightarrow \infty} \frac{1}{n^{2k-1}} \sum_{s, t=-(k-1)}^{k-1} \sum_{A'_{2k,s}} \sum_{A'_{2k,t}} \mathbb{E}[x_{i_1} \cdots x_{i_{2k}} x_{j_1} \cdots x_{j_{2k}}] \\
(19) \quad &= c_k \sum_{s=-(k-1)}^{k-1} \lim_{n \rightarrow \infty} \frac{|A'_{2k,s}| (2 - \mathbf{1}_{\{s=0\}}) k! k!}{n^{2k-1}}.
\end{aligned}$$

The constant $(2 - \mathbf{1}_{\{s=0\}})$ appeared because for $s \neq 0$, $A'_{2k,s}$ can (cross) match completely with $A'_{2k,s}$ and $A'_{2k,-s}$. The factor $k!^2$ appeared because for a fixed choice of $(i_1, \dots, i_{2k}) \in A'_{2k,s}$, we can choose the same set of values from $\{j_1, \dots, j_{2k}\}$ in $k!k!$ different ways, permuting the odd positions and the even positions among themselves.

Since from Result 4, we have

$$(20) \quad \lim_{n \rightarrow \infty} \frac{|A'_{2k,s}|}{n^{2k-1}} = \lim_{n \rightarrow \infty} \frac{|A_{2k,s}|}{n^{2k-1}} = \frac{1}{(2k-1)!} \sum_{j=0}^{k+s-1} (-1)^j \binom{2k}{j} (k+s-j)^{2k-1}.$$

Therefore for each fixed $k \geq 2$, from (19) and (20), we get

$$\begin{aligned}
\lim_{n \rightarrow \infty} \frac{1}{n^{p+q-1}} \sum_{I_{2k}} \mathbb{E}[x_{i_1} \cdots x_{i_{2p}} x_{j_1} \cdots x_{j_{2q}}] &= c_k \sum_{s=-(k-1)}^{k-1} \sum_{j=0}^{k+s-1} (-1)^j \binom{2k}{j} (k+s-j)^{2k-1} (2 - \mathbf{1}_{\{s=0\}}) k! k! \\
(21) \quad &= c_k g(k), \text{ say.}
\end{aligned}$$

Now we are left with $k = 1$ in Case II. If $k = 1$ then from (15), we get $\ell_1 = \ell_2$ and therefore

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n^{2p-1}} \sum_{I_2} (\mathbb{E}[x_{i_1} \dots x_{i_{2p}} x_{j_1} \dots x_{j_{2q}}] - \mathbb{E}[x_{i_1} \dots x_{i_{2p}}] \mathbb{E}[x_{j_1} \dots x_{j_{2q}}]) &= c_1 (\mathbb{E}x_1^4 - (\mathbb{E}x_1^2)^2) \\ (22) \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad &= (\mathbb{E}x_1^4 - 1)c_1, \end{aligned}$$

where $c_1 = \binom{p}{p-1}^2 (p-1)! \binom{q}{q-1}^2 (q-1)!$. Hence from (17), (21) and (22), we get

$$\lim_{n \rightarrow \infty} \text{Cov}(w_p, w_q) = \sum_{k=2}^{\min\{p,q\}} c_k g(k) + (\mathbb{E}x_1^4 - 1)c_1,$$

where $c_k = \binom{p}{p-k}^2 (p-k)! \binom{q}{q-k}^2 (q-k)!$ and $g(k)$ is given by

$$g(k) = \frac{1}{(2k-1)!} \sum_{s=-(k-1)}^{k-1} \sum_{j=0}^{k+s-1} (-1)^j \binom{2k}{j} (k+s-j)^{2k-1} (2 - \mathbf{1}_{\{s=0\}}) k! k!.$$

This complete the proof of the theorem 1. ■

3. PROOF OF THEOREM 2

We begin with some notation and definitions which will be used in the proof. First recall A_{2p} from (6) in Section 3,

$$A_{2p} = \{(i_1, \dots, i_{2p}) \in \mathbb{N}^{2p} : \sum_{k=1}^{2p} (-1)^k i_k = 0 \pmod{n}, 1 \leq i_1, \dots, i_{2p} \leq n\}.$$

For a vector $J = (j_1, j_2, \dots, j_{2p}) \in A_{2p}$, we define a multi-set S_J as

$$(23) \qquad \qquad \qquad S_J = \{j_1, j_2, \dots, j_{2p}\}.$$

Definition 7. Two vectors $J = (j_1, j_2, \dots, j_{2p})$ and $J' = (j'_1, j'_2, \dots, j'_{2p})$, where $J \in A_{2p}$ and $J' \in A_{2q}$, are said to be *connected* if $S_J \cap S_{J'} \neq \emptyset$.

For $1 \leq i \leq \ell$, suppose $J_i \in A_{2p_i}$. Now, we define cross-matched and self-matched element in $\cup_{i=1}^{\ell} S_{J_i}$.

Definition 8. An element in $\cup_{i=1}^{\ell} S_{J_i}$ is called *cross-matched* if it appears at least in two distinct S_{J_i} . If it appears in k many S_{J_i} 's, then we say its *cross-multiplicity* is k .

Definition 9. An element in $\cup_{i=1}^{\ell} S_{J_i}$ is called *self-matched* if it appears more than once in one of S_{J_i} . If it appears k many times in S_{J_i} , then we say its *self-multiplicity* in S_{J_i} is k .

An element in $\cup_{i=1}^{\ell} S_{J_i}$ can be both self-matched and cross-matched. If an element of $\cup_{i=1}^{\ell} S_{J_i}$ has *cross-multiplicity* one, that means, it appears only in one of the S_{J_i} .

Definition 10. Given a set of vectors $S = \{J_1, J_2, \dots, J_{\ell}\}$, where $J_i \in A_{2p_i}$ for $1 \leq i \leq \ell$, a subset $T = \{J_{n_1}, J_{n_2}, \dots, J_{n_k}\}$ of S is called a *cluster* if it satisfies the following two conditions:

- (i) For any pair J_{n_i}, J_{n_j} from T one can find a chain of vectors from T , which starts with J_{n_i} and ends with J_{n_j} such that any two neighbouring vectors in the chain are connected.
- (ii) The subset $\{J_{n_1}, J_{n_2}, \dots, J_{n_k}\}$ can not be enlarged to a subset which preserves condition (i).

For more details about cluster, we refer the readers to [12], where they have explained the structure of cluster by using graph.

Now we define $B_{P_{\ell}} \subseteq A_{2p_1} \times A_{2p_2} \times \dots \times A_{2p_{\ell}}$, where A_{2p_i} is as defined in (6).

Definition 11. Let $\ell \geq 2$ and $P_\ell = (2p_1, 2p_2, \dots, 2p_\ell)$. Now B_{P_ℓ} is a subset of $A_{2p_1} \times A_{2p_2} \times \dots \times A_{2p_\ell}$ such that $(J_1, J_2, \dots, J_\ell) \in B_{P_\ell}$ if

- (i) $\{J_1, J_2, \dots, J_\ell\}$ form a cluster,
- (ii) each element in $\cup_{i=1}^\ell S_{J_i}$ has multiplicity greater than or equal to two.

The next lemma gives us the cardinality of B_{P_ℓ} .

Lemma 12. For $\ell \geq 3$,

$$(24) \quad |B_{P_\ell}| = o(n^{p_1+p_2+\dots+p_\ell-\frac{\ell}{2}}).$$

Remark 13. The above lemma is not true if $\ell = 2$ and $p_1 = p_2$. Suppose $(J_1, J_2) \in B_{P_2}$. Then all $2p_1$ entries of J_1 may coincide with $2p_2 (= 2p_1)$ many entries of J_2 and hence

$$|B_{P_2}| = O(n^{2p_1-1}).$$

So in this situation, $|B_{P_2}| > o(n^{p_1+p_2-1})$.

The above lemma is similar to Lemma 15 of [12], where the authors have proved a similar result with a different set of constraints on the sets $A_{2p_1}, \dots, A_{2p_k}$. To prove Lemma 12, we shall use similar ideas. We prove Lemma 12 after the proof of Theorem 2. The following lemma is an easy consequence of Lemma 12.

Lemma 14. Suppose $\{J_1, J_2, \dots, J_\ell\}$ form a cluster where $J_i \in A_{2p_i}$ with $p_i \geq 1$ for $1 \leq i \leq \ell$, $\{x_i\}_{i \geq 1}$ is independent and satisfies (4). Then for $\ell \geq 3$,

$$(25) \quad \frac{1}{n^{p_1+p_2+\dots+p_\ell-\frac{\ell}{2}}} \sum_{A_{2p_1}, A_{2p_2}, \dots, A_{2p_\ell}} E \left[\prod_{k=1}^\ell (x_{J_k} - E(x_{J_k})) \right] = o(1),$$

where

$$J_k = (j_1^k, j_2^k, \dots, j_{2p_k}^k) \text{ and } x_{J_k} = x_{j_1^k} x_{j_2^k} \cdots x_{j_{2p_k}^k}.$$

Proof. First observe that $E \left[\prod_{k=1}^\ell (x_{J_k} - E(x_{J_k})) \right]$ will be non-zero only if each x_i appears at least twice in the collection $\{x_{j_1^k}, x_{j_2^k}, \dots, x_{j_{2p_k}^k}; 1 \leq k \leq \ell\}$, because $E(x_i) = 0$ for each i . Therefore

$$(26) \quad \sum_{A_{2p_1}, \dots, A_{2p_\ell}} E \left[\prod_{k=1}^\ell (x_{J_k} - E(x_{J_k})) \right] = \sum_{(J_1, \dots, J_\ell) \in B_{P_\ell}} E \left[\prod_{k=1}^\ell (x_{J_k} - E(x_{J_k})) \right],$$

where B_{P_ℓ} as in Definition 11. Since from (4), we have

$$E(x_i^2) = 1 \text{ and } \sup_{i \geq 1} E(|x_i|^k) = \alpha_k < \infty \text{ for } k \geq 3.$$

Therefore for $p_1, p_2, \dots, p_\ell \geq 1$, there exists $\beta_\ell > 0$, which depends only on p_1, p_2, \dots, p_ℓ , such that

$$(27) \quad \left| E \left[\prod_{k=1}^\ell (x_{J_k} - E(x_{J_k})) \right] \right| \leq \beta_\ell$$

for all $(J_1, J_2, \dots, J_\ell) \in A_{2p_1} \times A_{2p_2} \times \dots \times A_{2p_\ell}$.

Now using (26) and (27), we have

$$\sum_{A_{2p_1}, A_{2p_2}, \dots, A_{2p_\ell}} \left| E \left[\prod_{k=1}^\ell (x_{J_k} - E(x_{J_k})) \right] \right| \leq \sum_{(J_1, J_2, \dots, J_\ell) \in B_{P_\ell}} \beta_\ell = |B_{P_\ell}| \beta_\ell.$$

By using Lemma 12 in above expression, we get

$$\sum_{A_{2p_1}, A_{2p_2}, \dots, A_{2p_\ell}} \left| \mathbb{E} \left[\prod_{k=1}^{\ell} (x_{J_k} - \mathbb{E}(x_{J_k})) \right] \right| = o(n^{p_1+p_2+\dots+p_\ell-\frac{\ell}{2}}),$$

and hence

$$\frac{1}{n^{p_1+p_2+\dots+p_\ell-\frac{\ell}{2}}} \sum_{A_{2p_1}, A_{2p_2}, \dots, A_{2p_\ell}} \mathbb{E} \left[\prod_{k=1}^{\ell} (x_{J_k} - \mathbb{E}(x_{J_k})) \right] = o(1).$$

This completes the proof of lemma. ■

We shall use the above lemmata and Theorem 1 to prove Theorem 2.

Proof of Theorem 2. We use method of moments and Wick's formula to prove Theorem 2. First recall that from the method of moments, to prove $w_Q \xrightarrow{d} N(0, \sigma_Q^2)$, it is sufficient to show that

$$(28) \quad \lim_{n \rightarrow \infty} \mathbb{E}[(w_Q)^\ell] = \mathbb{E}[(N(0, \sigma_Q^2))^\ell] \quad \forall \ell = 1, 2, \dots$$

So, to prove (28), it is enough to show that, for $p_1, p_2, \dots, p_\ell \geq 1$,

$$\lim_{n \rightarrow \infty} \mathbb{E}[w_{p_1} w_{p_2} \cdots w_{p_\ell}] = \mathbb{E}[N_{p_1} N_{p_2} \cdots N_{p_\ell}],$$

where $\{N_p\}_{p \geq 1}$ is a centered Gaussian family with covariance $\sigma_{p,q}$, as in (5). Now using trace formula (7), we have

$$w_{p_k} = \frac{1}{\sqrt{n}} \left(\text{Tr}(RC_n)^{2p_k} - \mathbb{E}[\text{Tr}(RC_n)^{2p_k}] \right) = \frac{1}{n^{p_k-\frac{1}{2}}} \sum_{A_{2p_k}} \left(x_{j_1^k} \cdots x_{j_{2p_k}^k} - \mathbb{E}[x_{j_1^k} \cdots x_{j_{2p_k}^k}] \right).$$

Note that in the above summation $(j_1^k, j_2^k, \dots, j_{2p_k}^k) \in A_{2p_k}$. Therefore

$$(29) \quad \begin{aligned} & \mathbb{E}[w_{p_1} w_{p_2} \cdots w_{p_\ell}] \\ &= \frac{1}{n^{p_1+p_2+\dots+p_\ell-\frac{\ell}{2}}} \sum_{A_{2p_1}, A_{2p_2}, \dots, A_{2p_\ell}} \mathbb{E}[(x_{J_1} - \mathbb{E}x_{J_1})(x_{J_2} - \mathbb{E}x_{J_2}) \cdots (x_{J_\ell} - \mathbb{E}x_{J_\ell})]. \end{aligned}$$

Now for a fixed J_1, J_2, \dots, J_ℓ , if there exists a $k \in \{1, 2, \dots, \ell\}$ such that J_k is not connected with any J_i for $i \neq k$, then

$$\mathbb{E}[(x_{J_1} - \mathbb{E}x_{J_1})(x_{J_2} - \mathbb{E}x_{J_2}) \cdots (x_{J_\ell} - \mathbb{E}x_{J_\ell})] = 0$$

due to the independence of $\{x_i\}_{i \geq 1}$.

Therefore J_1, J_2, \dots, J_ℓ must form clusters with each cluster length greater than or equal to two, that is, each cluster should contain at least two vectors. Suppose G_1, G_2, \dots, G_s are the clusters formed by vectors J_1, J_2, \dots, J_ℓ and $|G_i| \geq 2$ for all $1 \leq i \leq s$ where $|G_i|$ denotes the length of the cluster G_i . Observe that $\sum_{i=1}^s |G_i| = \ell$.

If there exists a cluster G_j among G_1, G_2, \dots, G_s such that $|G_j| \geq 3$, then from Theorem 1 and Lemma 14, we have

$$\frac{1}{n^{p_1+p_2+\dots+p_\ell-\frac{\ell}{2}}} \sum_{A_{2p_1}, A_{2p_2}, \dots, A_{2p_\ell}} \mathbb{E}[(x_{J_1} - \mathbb{E}x_{J_1})(x_{J_2} - \mathbb{E}x_{J_2}) \cdots (x_{J_\ell} - \mathbb{E}x_{J_\ell})] = o(1).$$

Therefore, if ℓ is odd then there will be a cluster of odd length and hence

$$\lim_{n \rightarrow \infty} \mathbb{E}[w_{p_1} w_{p_2} \cdots w_{p_\ell}] = 0.$$

Similarly, if ℓ is even then the contribution due to $\{J_1, J_2, \dots, J_\ell\}$ to $E[w_{p_1} w_{p_2} \cdots w_{p_\ell}]$ is $O(1)$ only when $\{J_1, J_2, \dots, J_\ell\}$ decomposes into clusters of length 2. Therefore from (29), we get

$$\begin{aligned} & \lim_{n \rightarrow \infty} E[w_{p_1} w_{p_2} \cdots w_{p_\ell}] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n^{p_1 + p_2 + \cdots + p_\ell - \frac{\ell}{2}}} \sum_{A_{2p_1}, A_{2p_2}, \dots, A_{2p_\ell}} E[(x_{J_1} - E x_{J_1})(x_{J_2} - E x_{J_2}) \cdots (x_{J_\ell} - E x_{J_\ell})] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n^{p_1 + p_2 + \cdots + p_\ell - \frac{\ell}{2}}} \sum_{\pi \in \mathcal{P}_2(\ell)} \prod_{i=1}^{\frac{\ell}{2}} \sum_{A_{2p_{y(i)}}, A_{2p_{z(i)}}} E[(x_{J_{y(i)}} - E x_{J_{y(i)}})(x_{J_{z(i)}} - E x_{J_{z(i)}})], \end{aligned}$$

where $\pi = \{\{y(1), z(1)\}, \dots, \{y(\frac{\ell}{2}), z(\frac{\ell}{2})\}\} \in \mathcal{P}_2(\ell)$ and $\mathcal{P}_2(\ell)$ is the set of all pair partition of $\{1, 2, \dots, \ell\}$. Using Theorem 1, from the last equation we get

$$(30) \quad \lim_{n \rightarrow \infty} E[w_{p_1} w_{p_2} \cdots w_{p_\ell}] = \sum_{\pi \in \mathcal{P}_2(\ell)} \prod_{i=1}^{\frac{\ell}{2}} \lim_{n \rightarrow \infty} E[w_{p_{y(i)}} w_{p_{z(i)}}].$$

Since from Theorem 1, we have

$$\lim_{n \rightarrow \infty} E(w_p w_q) = \sigma_{p,q} = E(N_p N_q).$$

Therefore using Wick's formula, from (30) we get

$$\begin{aligned} \lim_{n \rightarrow \infty} E[w_{p_1} w_{p_2} \cdots w_{p_\ell}] &= \sum_{\pi \in \mathcal{P}_2(\ell)} \prod_{i=1}^{\frac{\ell}{2}} \lim_{n \rightarrow \infty} E[w_{p_{y(i)}} w_{p_{z(i)}}] \\ &= \sum_{\pi \in \mathcal{P}_2(\ell)} \prod_{i=1}^{\frac{\ell}{2}} E[N_{p_{y(i)}} N_{p_{z(i)}}] \\ &= E[N_{p_1} N_{p_2} \cdots N_{p_\ell}]. \end{aligned}$$

This completes the proof of Theorem 2. ■

Now we prove Lemma 12.

Proof of Lemma 12. We define subsets B'_{P_ℓ} and B''_{P_ℓ} of B_{P_ℓ} as

$$(31) \quad B'_{P_\ell} = \{(J_1, J_2, \dots, J_\ell) \in B_{P_\ell} : \text{cross-multiplicity of each entry of } \cup_{i=1}^{\ell} S_{J_i} \text{ is less than or equal to two}\}$$

and

$$(32) \quad B''_{P_\ell} = B_{P_\ell} \setminus B'_{P_\ell}.$$

Observe that

$$|B_{P_\ell}| = |B'_{P_\ell}| + |B''_{P_\ell}|.$$

Since in B''_{P_ℓ} there exists at least one entry with *cross-multiplicity* greater than or equal to three, the maximum number of free entries (which can be chosen freely from 1 to n) in B''_{P_ℓ} will be less than or equal to the maximum number of free entries in B'_{P_ℓ} . Therefore

$$|B''_{P_\ell}| \leq |B'_{P_\ell}|$$

and hence

$$(33) \quad |B_{P_\ell}| = O(|B'_{P_\ell}|).$$

So to prove (24), it is enough to prove that

$$(34) \quad |B'_{P_\ell}| = o(n^{p_1+p_2+\dots+p_\ell-\frac{\ell}{2}}).$$

Now we shall prove (34) by using mathematical induction on ℓ . Let us first prove it for $\ell = 3$. Suppose $(J_1, J_2, J_3) \in B_{P_3}$ with $J_i = (j_1^i, j_2^i, \dots, j_{2p_i}^i)$ for $i = 1, 2, 3$, then (J_1, J_2, J_3) form a cluster and they can be connected in the following way:

Suppose r many entries of J_1 matches with r many entries of J_2 and J_3 both; r_1 many entries of J_1 matches with r_1 many entries from J_2 only; r_2 many entries of J_2 matches with r_2 many entries of J_3 only; r_3 many entries of J_3 matches with r_3 many entries of J_1 only, where

$$(35) \quad r, r_1, r_2, r_3 \geq 0, \quad r + r_1 + r_2 \leq 2p_2, \quad r + r_1 + r_3 \leq 2p_1, \quad r + r_2 + r_3 \leq 2p_3.$$

Note that if $r = 0$, that is, all entries of $\cup_{i=1}^3 S_{J_i}$ has *cross-multiplicity* less than or equal to two,




FIGURE 1. Connection between J_1, J_2 and J_3 .

then (J_1, J_2, J_3) belongs to B'_{P_3} as defined in (31). Also if $r \neq 0$, that is, there are r many entries of $\cup_{i=1}^3 S_{J_i}$ which has *cross-multiplicity* three, then (J_1, J_2, J_3) belongs to B''_{P_3} as defined in (32). Now we calculate cardinality of B'_{P_3} , that is, the case $r = 0$, to prove (24).

Recall from the Case I and Case II of proof of Theorem 1, the number of free entries between two vectors which have same k many entries, will be maximum when k is even. Therefore to obtain maximum number of free entries from B'_{P_3} , we shall assume each r_i is even, for $i = 1, 2, 3$. Otherwise, we shall get less contribution.

Since (J_1, J_2, J_3) is forming a cluster, therefore at least two r_i have to be non-zero. Then the following two cases arise.

Case I. Exactly two r_i are non-zero and one r_i is zero. Without loss of generality, we assume $r_3 = 0$. Therefore we can represent J_i as


$$\begin{aligned} J_1 &= (x_1^1, x_2^1, \dots, x_{r_1}^1, j_{r_1+1}^1, j_{r_1+2}^1, \dots, j_{2p_1}^1), \\ J_2 &= (x_1^1, x_2^1, \dots, x_{r_1}^1, x_{r_1+1}^2, x_{r_1+2}^2, \dots, x_{r_2}^2, j_{r_1+r_2+1}^2, j_{r_1+r_2+2}^2, \dots, j_{2p_2}^2), \\ J_3 &= (x_1^2, x_2^2, \dots, x_{r_2}^2, j_{r_2+1}^3, j_{r_2+2}^3, \dots, j_{2p_3}^3), \end{aligned}$$

where (35) will be of the following form

$$r_1 \leq 2p_1, \quad r_1 + r_2 \leq 2p_2, \quad r_2 \leq 2p_3$$

and Figure 1 will look like as Figure 2. As each entry in $\cup_{i=1}^3 S_{J_i}$ has multiplicity at least two, then we shall get maximum contribution from J_1 , if the following holds:

- (i) each elements of $\{x_1^1, x_2^1, \dots, x_{r_1}^1\}$ has to be distinct,
- (ii) $\{x_1^1, x_2^1, \dots, x_{r_1}^1\}$ have no common element with $\{j_{r_1+1}^1, j_{r_1+2}^1, \dots, j_{2p_1}^1\}$,
- (iii) $(j_{r_1+1}^1, j_{r_1+2}^1, \dots, j_{2p_1}^1)$ is *odd-even pair matched*.

FIGURE 2. Connection between J_1, J_2 and J_3 : Case I.

Since $(j_{r_1+1}^1, j_{r_1+2}^1, \dots, j_{2p_1}^1)$ is *odd-even pair matched*, that is, $\sum_{s=r_1+1}^{2p_1} (-1)^s j_s^1 = 0 \pmod{n}$, therefore the constraint on the entries of J_1 , that is, $\sum_{s=1}^{r_1} (-1)^s x_s^1 + \sum_{s=r_1+1}^{p_1} (-1)^s j_s^1 = 0 \pmod{n}$ will change automatically into $\sum_{s=1}^{r_1} (-1)^s x_s^1 = 0 \pmod{n}$. So the maximum contribution from J_1 is $O(n^{(r_1 + \frac{2p_1 - r_1 - 1}{2})})$, where (-1) is coming due to the constraint $\sum_{s=1}^{r_1} (-1)^s x_s^1 = 0 \pmod{n}$. Now after fixing entries in J_1 , by similar argument, the maximum contribution from J_3 will be $O(n^{(r_2 + \frac{2p_3 - r_2 - 1}{2})})$, where (-1) is coming due to the constraint $\sum_{s=1}^{r_2} (-1)^s x_s^2 = 0 \pmod{n}$. After selecting J_1, J_3 ; the contribution from J_2 will be $O(n^{\frac{2p_2 - r_1 - r_2}{2}})$ when $(j_{r_1+r_2+1}^2, j_{r_1+r_2+2}^2, \dots, j_{2p_2}^2)$ is *odd-even pair matched*. Note that the constraint of J_2 is automatically satisfied because we have already considered $\sum_{s=1}^{r_1} (-1)^s x_s^1 = 0 \pmod{n}$ and $\sum_{s=1}^{r_2} (-1)^s x_s^2 = 0 \pmod{n}$. That is why, (-1) is not coming in the contribution of J_2 . So in this case cardinality of B'_{P_3} will be

$$O(n^{(r_1 + p_1 - \frac{r_1}{2} - 1) + (r_2 + p_3 - \frac{r_2}{2} - 1) + (p_2 - \frac{r_1 + r_2}{2})}) = O(n^{p_1 + p_2 + p_3 - 2}).$$

Case II. Each r_i is non-zero for all $i = 1, 2, 3$. Therefore we can represent J_i as

$$\begin{aligned} J_1 &= (x_1^1, x_2^1, \dots, x_{r_1}^1, x_{r_1+1}^3, x_{r_1+2}^3, \dots, x_{r_3}^3, j_{r_1+r_3+1}^1, j_{r_1+r_3+2}^1, \dots, j_{2p_1}^1), \\ J_2 &= (x_1^1, x_2^1, \dots, x_{r_1}^1, x_{r_1+1}^2, x_{r_1+2}^2, \dots, x_{r_2}^2, j_{r_1+r_2+1}^2, j_{r_1+r_2+2}^2, \dots, j_{2p_2}^2), \\ J_3 &= (x_1^2, x_2^2, \dots, x_{r_2}^2, x_{r_2+1}^3, x_{r_2+2}^3, \dots, x_{r_3}^3, j_{r_2+r_3+1}^3, j_{r_2+r_3+2}^3, \dots, j_{2p_3}^3), \end{aligned}$$

where $r_1 + r_3 \leq 2p_1$, $r_1 + r_2 \leq 2p_2$, $r_2 + r_3 \leq 2p_3$.

Observe that each entry in $\cup_{i=1}^3 S_{J_i}$ has multiplicity at least two. Now by the similar arguments, as given in Case I, we shall calculate the contributions due to J_i . For the maximum contribution, $(2p_1 - r_1 - r_3)$ many entries in J_1 have to be *odd-even pair matched*, as we discussed in Case I. So the maximum contribution from J_1 will be $O(n^{(r_1 + r_3 + \frac{2p_1 - r_1 - r_3}{2} - 1)})$, where (-1) is coming due to the new constraint on the entries of J_1 , that is, $\sum_{s=1}^{r_1} (-1)^s x_s^1 + \sum_{s=1}^{r_3} (-1)^{r_1+s} x_s^3 = 0 \pmod{n}$. Now after fixing entries in J_1 , by similar argument the maximum contribution from J_3 will be $O(n^{(r_2 + \frac{2p_3 - r_2 - r_3}{2} - 1)})$, where (-1) is coming due to the new constraint on the entries of J_3 , that is, $\sum_{s=1}^{r_2} (-1)^s x_s^2 + \sum_{s=1}^{r_3} (-1)^{r_2+s} x_s^3 = 0 \pmod{n}$. After selecting J_1, J_3 ; by similar argument, the contribution from J_2 will be $O(n^{(\frac{p_2 - r_1 - r_2}{2})})$. So in this case the cardinality of B'_{P_3} will be

$$O(n^{(r_1 + r_3 + p_1 - \frac{r_1 + r_3}{2} - 1) + (r_2 + p_3 - \frac{r_2 + r_3}{2} - 1) + (p_2 - \frac{r_1 + r_2}{2})}) = O(n^{p_1 + p_2 + p_3 - 2}).$$

Therefore after combining the Case I and Case II, we get

$$|B'_{P_3}| = O(n^{p_1 + p_2 + p_3 - 2}) = o(n^{p_1 + p_2 + p_3 - \frac{3}{2}}).$$

This show that (34) is true for $\ell = 3$. So Lemma 12 is true for $\ell = 3$.

Now for $r \neq 0$, one can show that $|B''_{P_3}| = O(n^{p_1 + p_2 + p_3 - \frac{4+r}{2}})$, which is less than the cardinality of B'_{P_3} . We leave it to the readers to verify.

Suppose (34) is true for $\ell = k$, that is,

$$|B'_{P_k}| = o(n^{p_1 + p_2 + \dots + p_k - \frac{k}{2}}).$$

Now we shall prove it for $\ell = k+1$. Suppose $(J_1, J_2, \dots, J_k, J_{k+1}) \in B_{P_{k+1}}$ with $J_i = (j_1^i, j_2^i, \dots, j_{2p_i}^i)$ for $i = 1, 2, \dots, k+1$, then $\{J_1, J_2, \dots, J_{k+1}\}$ forms a cluster. First consider J_1 , since $\{J_1, J_2, \dots, J_{k+1}\}$ form a cluster therefore there exists at least one $J_i \in \{J_2, J_3, \dots, J_{k+1}\}$ such that J_i is connected with J_1 , and without loss of generality, we suppose $J_i = J_2$. So J_2 is connected with J_1 . Again as $\{J_1, J_2, \dots, J_{k+1}\}$ forms a cluster, there exists at least one $J_i \in \{J_3, J_4, \dots, J_{k+1}\}$ such that J_i is connected either with J_1 or J_2 or both, and without loss of generality, we assume that $J_i = J_3$. Similarly we can arrange $\{J_4, J_5, \dots, J_k\}$. Note that, due to the above arrangement, $\{J_1, J_2, \dots, J_k\}$ forms a cluster. But it is given that $\{J_1, J_2, \dots, J_k, J_{k+1}\}$ forms a cluster, therefore J_{k+1} has to be connected with at least one of $\{J_1, J_2, \dots, J_k\}$. Note that, here $\{J_1, J_2, \dots, J_k, J_{k+1}\}$ are arranged in such a way that if we remove J_{k+1} from the cluster formed by $\{J_1, J_2, \dots, J_k, J_{k+1}\}$, then $\{J_1, J_2, \dots, J_k\}$ also form a cluster.

Now we consider $B'_{P_{k+1}}$, that is, there is no element in $\cup_{i=1}^{k+1} S_{J_i}$ which has *cross-multiplicity* more than two. Suppose for $i = 1, 2, \dots, k$; r_i many entries of J_{k+1} matches with r_i many entries of J_i only. Therefore we can represent J_i as

$$J_i = (x_1^i, x_2^i, \dots, x_{r_i}^i, j_{r_i+1}^i, j_{r_i+2}^i, \dots, j_{2p_i}^i) \text{ for } i = 1, 2, \dots, k,$$

and

$$(36) \quad J_{k+1} = (x_1^1, x_2^1, \dots, x_{r_1}^1, x_1^2, x_2^2, \dots, x_{r_2}^2, \dots, x_1^k, \dots, x_{r_k}^k, j_{r'+1}^{k+1}, j_{r'+2}^{k+1}, \dots, j_{2p_{k+1}}^{k+1}),$$

where

$$0 \leq r_i \leq 2p_i, \quad \sum_{i=1}^k r_i \leq 2p_{k+1} \text{ and } r' = \sum_{i=1}^k r_i.$$

Observe that r_i is non-zero for at least one i , as $\{J_1, J_2, \dots, J_{k+1}\}$ form a cluster. Now we define a multi-set X as

$$(37) \quad X = \{x_1^1, x_2^1, \dots, x_{r_1}^1, x_1^2, x_2^2, \dots, x_{r_2}^2, \dots, x_1^k, x_2^k, \dots, x_{r_k}^k\}.$$

First note that $r_i \neq 2p_i$ for all $i = 1, 2, \dots, k$. Otherwise we will get an element in $\cup_{i=1}^{k+1} S_{J_i}$ which




FIGURE 3. Connection between $J_1, J_2, \dots, J_k, J_{k+1}$

has *cross-multiplicity* greater than two, as $\{J_1, J_2, \dots, J_k\}$ forms a cluster after removing J_{k+1} . That is not possible as we are considering the set $B'_{P_{k+1}}$.

Now let us calculate the contributions of J_i 's to the cardinality of $B'_{P_{k+1}}$, as we have calculated for $\ell = 3$. Note that, we will get maximum number of free entries in $B'_{P_{k+1}}$, when each r_i will be even, as we have assumed for $\ell = 3$ case and we have proved this in the Case I and Case II of proof of Theorem 1. We first calculate the contribution from J_{k+1} . We shall get maximum contribution from J_{k+1} , if the following holds:

- (i) each elements of X has to be distinct,
- (ii) X have no common element with $\{j_{r'+1}^{k+1}, j_{r'+2}^{k+1}, \dots, j_{2p_{k+1}}^{k+1}\}$,
- (iii) $(j_{r'+1}^{k+1}, j_{r'+2}^{k+1}, \dots, j_{2p_{k+1}}^{k+1})$ has to be *odd-even pair matched*,

where $r' = \sum_{i=1}^k r_i \leq 2p_{k+1}$. Recall that, J_{k+1} has a constraint, namely,

$$\sum_{s=1}^{r_1} (-1)^s x_s^1 + \sum_{s=1}^{r_2} (-1)^{r_1+s} x_s^2 + \dots + \sum_{s=1}^{r_k} (-1)^{r_1+r_2+\dots+r_{k-1}+s} x_s^k + \sum_{s=r'+1}^{2p_{k+1}} (-1)^s j_s^{k+1} = 0 \pmod{n}.$$

Since $(j_{r'+1}^{k+1}, j_{r'+2}^{k+1}, \dots, j_{2p_{k+1}}^{k+1})$ is *odd-even pair matched* and each elements of X are distinct. Therefore the above constraint of J_{k+1} will change into the new form

$$(38) \quad \sum_{s=1}^{r_1} (-1)^s x_s^1 + \sum_{s=1}^{r_2} (-1)^{r_1+s} x_s^2 + \dots + \sum_{s=1}^{r_k} (-1)^{r_1+r_2+\dots+r_{k-1}+s} x_s^k = 0 \pmod{n}.$$

So the maximum contribution from J_{k+1} is

$$(39) \quad T_{k+1} = O\left(n^{r' + \frac{2p_{k+1} - r'}{2}} - 1\right),$$

where (-1) is coming in the expression of T_{k+1} because J_{k+1} has the constraint (38). After selecting the entries of J_{k+1} , at least r_i many entries of J_i are also selected (fixed) and hence effective entries (remains to count) from J_i is at most $(2p_i - r_i)$ for $i = 1, 2, \dots, k$. Note that if $\{x_1^i, x_2^i, \dots, x_{r_i}^i\}$ have no common element with $\{j_{r_i+1}^i, j_{r_i+2}^i, \dots, j_{2p_i}^i\}$ for $i = 1, 2, \dots, k$, then after selecting the entries of J_{k+1} , exactly r_i many entries of J_i are also selected (fixed) and hence effective entries (remains to count) from J_i is exactly $(2p_i - r_i)$ for $i = 1, 2, \dots, k$. Observe that the constraint on J_i , $\sum_{s=1}^{r_i} (-1)^s x_s^i + \sum_{s=r_i+1}^{2p_i} (-1)^s j_s^i = 0 \pmod{n}$ will change to

$$\sum_{s=r_i+1}^{P_i} (-1)^s j_s^i = -d_{r_i} \pmod{n},$$

where $d_{r_i} = \sum_{s=1}^{r_i} (-1)^s x_s^i$. Since $|B'_{P_{k+1}}|$ is product of T_{k+1} and the number of free entries of (J_1, J_2, \dots, J_k) which remains to count after selecting J_{k+1} , that is, $|\tilde{B}_{P_k}|$. Therefore the cardinality of $B'_{P_{k+1}}$ will be

$$(40) \quad |B'_{P_{k+1}}| = |\tilde{B}_{P_k}| T_{k+1},$$

where \tilde{B}_{P_k} is defined as,

$$(41) \quad \tilde{B}_{P_k} = \{(J_1, J_2, \dots, J_k) \in B'_{P_k} : r_i \text{ many entries of } J_i \text{ are already chosen for } i = 1, 2, \dots, k\}.$$

Now we calculate the cardinality of \tilde{B}_{P_k} . We denote $(j_{r_i+1}^i, j_{r_i+2}^i, \dots, j_{p_i}^i)$ by J'_i for $i = 1, 2, \dots, k$. Observe that, due to the arrangement of $J_1, J_2, \dots, J_k, J_{k+1}$, if we remove J_{k+1} from $B'_{P_{k+1}}$, then $\{J_1, J_2, \dots, J_k\}$ also form a cluster and hence $\{J'_1, J'_2, \dots, J'_k\}$ also form a cluster. Since each element of $\cup_{i=1}^{k+1} S_{J_i}$ has *cross-multiplicity* less than or equal to two, therefore each element of $\cup_{i=1}^k S_{J'_i}$ also has *cross-multiplicity* less than or equal to two. As for the maximum contribution, $\{x_1^i, x_2^i, \dots, x_{r_i}^i\}$

have no common element with $\{j_{r_i+1}^i, j_{r_i+2}^i, \dots, j_{p_i}^i\}$ for all $i = 1, 2, \dots, k$ and since each elements of $\cup_{i=1}^{k+1} S_{J_i}$ has multiplicity greater than or equal to two, therefore each element of $\cup_{i=1}^k S_{J'_i}$ also has multiplicity greater than or equal to two. From the above discussion on $\{J'_1, J'_2, \dots, J'_k\}$, we get

- (i) $\{J'_1, J'_2, \dots, J'_k\}$ form a cluster,
- (ii) each element of $\cup_{i=1}^k S_{J'_i}$ has multiplicity greater than or equal to two,
- (iii) each element of $\cup_{i=1}^k S_{J'_i}$ has *cross-multiplicity* less than or equal to two,
- (iv) $j_{r_i+1}^i + j_{r_i+2}^i + \dots + j_{p_i}^i = -d_{r_i} \pmod{n}$ for $i = 1, 2, \dots, k$.

So the set \tilde{B}_{P_k} is the collection of the vectors $(J'_1, J'_2, \dots, J'_k)$, which satisfy the above four conditions. Now consider the subset $B'_{P'_k}$ of $B_{P'_k}$, where $B_{P'_k}$ is as in the definition 11 with $P'_k = (p'_1, p'_2, \dots, p'_k)$ and $p'_i = 2p_i - r_i$ for $i = 1, 2, \dots, k$. Therefore

$$(42) \quad B'_{P'_k} = \{(T_1, T_2, \dots, T_k) \in B_{P'_k} : \text{cross-multiplicity of each entry of } \cup_{i=1}^k S_{T_i} \text{ is less than or equal to two}\}.$$

Now by the induction hypothesis, we get

$$|B'_{P'_k}| = o(n^{p'_1+p'_2+\dots+p'_k-\frac{k}{2}}).$$

Now we compare the cardinality of \tilde{B}_{P_k} and $B'_{P'_k}$. Note that both the sets \tilde{B}_{P_k} and $B'_{P'_k}$ are almost same, the only difference is in the constraints. Recall that, if $(J'_1, J'_2, \dots, J'_k) \in \tilde{B}_{P_k}$, then entries of J'_i has constraints

$$(43) \quad j_{r_i+1}^i + j_{r_i+2}^i + \dots + j_{p_i}^i = -d_{r_i} \pmod{n} \text{ for } i = 1, 2, \dots, k,$$

where d_{r_i} are constants for $i = 1, 2, \dots, k$.

Recall that, if $(T_1, T_2, \dots, T_k) \in B'_{P'_k}$, then entries of T_i has constraints

$$(44) \quad t_1^i + t_2^i + \dots + t_{p'_i}^i = 0 \pmod{n} \text{ for } i = 1, 2, \dots, k.$$

Note that, if $d_{r_i} = d_{r_1} \pmod{n}$ for $i = 2, 3, \dots, k$, then the constraints of \tilde{B}_{P_k} , that is, (43) will be

$$j_{r_i+1}^i + j_{r_i+2}^i + \dots + j_{p_i}^i = -d_{r_1} \pmod{n} \text{ for } i = 1, 2, \dots, k,$$

which shows that $|\tilde{B}_{P_k}| = |B'_{P'_k}|$. If $d_{r_i} \neq d_{r_1} \pmod{n}$ for at least one i , then the entries of \tilde{B}_{P_k} have to satisfy more constraints than the constraints on entries of $B'_{P'_k}$, that is, $|\tilde{B}_{P_k}| \leq |B'_{P'_k}|$. Hence

$$(45) \quad |\tilde{B}_{P_k}| \leq |B'_{P'_k}|.$$

Therefore from (40) and (45), we get

$$\begin{aligned} |B'_{P_{k+1}}| &= |\tilde{B}_{P_k}| |T_{k+1}| \\ &\leq |B'_{P'_k}| |T_{k+1}| \\ &= o(n^{p'_1+p'_2+\dots+p'_k-\frac{k}{2}}) O(n^{r'+\frac{2p_{k+1}-r'}{2}}-1) \\ &= o(n^{(p'_1+p'_2+\dots+p'_k-\frac{k}{2})+(r'+\frac{2p_{k+1}-r'}{2})-1}). \end{aligned}$$

Since $p'_i = 2p_i - r_i$ for $i = 1, 2, \dots, k$ and $\sum_{i=1}^k r_i = r'$, from the last equation, we have

$$|B'_{P_{k+1}}| = o(n^{p_1+p_2+\dots+p_k+p_{k+1}-\frac{k+1}{2}-1}) = o(n^{p_1+p_2+\dots+p_k+p_{k+1}-\frac{k+1}{2}}).$$

Therefore from (33), we get

$$|B_{P_{k+1}}| = o\left(n^{p_1+p_2+\dots+p_k+p_{k+1}-\frac{k+1}{2}}\right).$$

So Lemma 12 is true for $\ell = k + 1$. This complete the proof of Lemma 12. ■

REFERENCES

- [1] Kartick Adhikari and Koushik Saha, *Fluctuations of eigenvalues of patterned random matrices*, J. Math. Phys. **58** (2017), no. 6, 063301, 20. MR 3666201
- [2] ———, *Universality in the fluctuation of eigenvalues of random circulant matrices*, Statist. Probab. Lett. **138** (2018), 1–8. MR 3788711
- [3] Greg W. Anderson and Ofer Zeitouni, *A CLT for a band matrix model*, Probab. Theory Related Fields **134** (2006), no. 2, 283–338. MR 2222385
- [4] L. V. Arharov, *Limit theorems for the characteristic roots of a sample covariance matrix*, Dokl. Akad. Nauk SSSR **199** (1971), 994–997. MR 0309171
- [5] Z. D. Bai and Jack W. Silverstein, *CLT for linear spectral statistics of large-dimensional sample covariance matrices*, Ann. Probab. **32** (2004), no. 1A, 553–605. MR 2040792
- [6] Sourav Chatterjee, *Fluctuations of eigenvalues and second order Poincaré inequalities*, Probab. Theory Related Fields **143** (2009), no. 1-2, 1–40. MR 2449121
- [7] I. Jana, K. Saha, and A. Soshnikov, *Fluctuations of linear eigenvalue statistics of random band matrices*, Theory Probab. Appl. **60** (2016), no. 3, 407–443. MR 3568789
- [8] Kurt Johansson, *On fluctuations of eigenvalues of random Hermitian matrices*, Duke Math. J. **91** (1998), no. 1, 151–204. MR 1487983
- [9] Lingyun Li and Alexander Soshnikov, *Central limit theorem for linear statistics of eigenvalues of band random matrices*, Random Matrices Theory Appl. **2** (2013), no. 4, 1350009, 50. MR 3149439
- [10] Dang-Zheng Liu, Xin Sun, and Zheng-Dong Wang, *Fluctuations of eigenvalues for random toeplitz and related matrices*, Electron. J. Probab. **17** (2012), no. 95, 1–22.
- [11] A. Lytova and L. Pastur, *Central limit theorem for linear eigenvalue statistics of random matrices with independent entries*, Ann. Probab. **37** (2009), no. 5, 1778–1840. MR 2561434
- [12] Shambhu Nath Maurya and Koushik Saha, *Process convergence of fluctuations of linear eigenvalue statistics of random circulant matrices*, arXiv preprint arXiv:1909.00686 (2019).
- [13] M. Shcherbina, *Central limit theorem for linear eigenvalue statistics of the Wigner and sample covariance random matrices*, Zh. Mat. Fiz. Anal. Geom. **7** (2011), no. 2, 176–192, 197, 199. MR 2829615
- [14] ———, *On fluctuations of eigenvalues of random band matrices*, J. Stat. Phys. **161** (2015), no. 1, 73–90. MR 3392508
- [15] Ya. Sinai and A. Soshnikov, *Central limit theorem for traces of large random symmetric matrices with independent matrix elements*, Bol. Soc. Brasil. Mat. (N.S.) **29** (1998), no. 1, 1–24. MR 1620151

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI, MAHARASHTRA 400076, INDIA

E-mail address: snmaurya [at] math.iitb.ac.in

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI, MAHARASHTRA 400076, INDIA

E-mail address: koushik.saha [at] iitb.ac.in