

# Linear spectral statistics of eigenvectors of anisotropic sample covariance matrices

Fan Yang <sup>\*1</sup>

<sup>1</sup>Department of Statistics, University of Pennsylvania

March 1, 2022

## Abstract

Consider sample covariance matrices of the form  $Q := \Sigma^{1/2} X X^* \Sigma^{1/2}$ , where  $X = (x_{ij})$  is an  $n \times N$  random matrix whose entries are independent random variables with mean zero and variance  $N^{-1}$ , and  $\Sigma$  is a deterministic positive-definite matrix. We study the limiting behavior of the eigenvectors of  $Q$  through the so-called *eigenvector empirical spectral distribution* (VESD)  $F_{\mathbf{u}}$ , which is an alternate form of empirical spectral distribution with weights given by  $|\mathbf{u}^\top \xi_k|^2$ , where  $\mathbf{u}$  is any deterministic unit vector and  $\xi_k$  are the eigenvectors of  $Q$ . We prove a functional central limit theorem for the linear spectral statistics of  $F_{\mathbf{u}}$ , indexed by functions with Hölder continuous derivatives. We show that the linear spectral statistics converge to universal Gaussian processes both on global scales of order 1, and on local scales that are much smaller than 1 and much larger than the typical eigenvalues spacing  $N^{-1}$ . Moreover, we give explicit expressions for the means and covariance functions of the Gaussian processes, where the exact dependence on  $\Sigma$  and  $\mathbf{u}$  allows for more flexibility in the applications of VESD in statistical estimations of sample covariance matrices.

## 1 Introduction

Consider a centered random vector  $\mathbf{y} \in \mathbb{R}^n$  with population covariance matrix  $\Sigma = \mathbb{E} \mathbf{y} \mathbf{y}^*$ . Given  $N$  i.i.d. samples  $(\mathbf{y}_1, \dots, \mathbf{y}_N)$  of  $\mathbf{y}$ , the simplest estimator for  $\Sigma$  is the sample covariance matrix  $Q := N^{-1} \sum_i \mathbf{y}_i \mathbf{y}_i^*$ . Large dimensional sample covariance matrices have been a central object of study in high-dimensional statistics. In many modern applications, such as statistics [15, 23, 24, 26], economics [37] and population genetics [38], the advance of technology has led to high dimensional data where  $n$  is comparable to or even larger than  $N$ . In this setting, the law of large numbers does not hold and  $\Sigma$  cannot be approximated by  $Q$  directly. However, with more advanced tools in random matrix theory, it is possible to infer some properties of  $\Sigma$  from the eigenvalue and eigenvector statistics of  $Q$ .

In this paper, we consider sample covariance matrices of the form  $Q_1 := \Sigma^{1/2} X X^* \Sigma^{1/2}$ , where  $X = (x_{ij})$  is an  $n \times N$  real data matrix whose entries are independent random variables satisfying

$$\mathbb{E} x_{ij} = 0, \quad \mathbb{E} |x_{ij}|^2 = N^{-1}, \quad 1 \leq i \leq n, \quad 1 \leq j \leq N, \quad (1.1)$$

and the population covariance matrix  $\Sigma$  is an  $n \times n$  deterministic positive-definite matrix. Define the aspect ratio  $d_N := n/N$ . We are interested in the high dimensional setting with  $d_N \rightarrow d \in (0, \infty)$  as  $N \rightarrow \infty$ . We will also use the  $N \times N$  matrix  $Q_2 := X^* \Sigma X$ , which share the same nonzero eigenvalues with  $Q_1$ .

---

\*E-mail: fyang75@wharton.upenn.edu

In the study of eigenvalue statistics of large dimensional sample covariance matrices, one of the most fundamental subjects of study is the asymptotic behavior of empirical spectral distribution (ESD). It is well-known that the ESD of  $Q_1$  converges weakly to the famous Marčenko-Pastur (MP) law  $F_{MP}$  [35]. The convergence rate was first established in [2], and later improved in [20] to  $O(N^{-1/2})$  in probability under the finite 8th moment condition. In [40], the authors proved an almost optimal bound  $O(N^{-1+\varepsilon})$  with high probability for any fixed  $\varepsilon > 0$  under the sub-exponential decay assumption. For the limiting spectral statistics, a functional CLT was proved in [5] for the ESD of  $Q_1$ . Roughly speaking, it was proved that given an analytic functions  $f(x)$ , the random variable

$$\sum_{i=1}^n f(\lambda_i) - n \int f(x) dF_{MP}(x)$$

converges in distribution to a centered Gaussian random variable, where  $\lambda_i$  are the eigenvalues of  $Q_1$ . In fact, [5] proved a more general multivariate statement that for any analytic functions  $f_1(x), \dots, f_k(x)$ , the random vector

$$\left( \sum_{i=1}^n f_s(\lambda_i) - n \int f_s(x) dF_{MP}(x) \right)_{1 \leq s \leq k}$$

converge in distribution to a centered Gaussian vector. This kind of result is usually referred to as “linear eigenvalue statistics”. Recently, in [32] the authors extended it to mesoscopic eigenvalue statistics, that is, for any fixed  $E > 0$  and scale parameter  $n^{-1} \ll \eta \ll 1$ , the random vector

$$\left( \sum_{i=1}^n f_s \left( \frac{\lambda_i - E}{\eta} \right) - n \int f_s \left( \frac{x - E}{\eta} \right) dF_{MP}(x) \right)_{1 \leq s \leq k}$$

also converge in distribution to a centered Gaussian vector. We shall call such a result the “local linear eigenvalue statistics”.

The concept of ESD can be also extended to encode the information of sample eigenvectors. Following [42, 43, 3, 47, 48], we define the following concept of *eigenvector empirical spectral distribution* (VESD). Suppose

$$\Sigma^{1/2} X = \sum_{1 \leq k \leq N \wedge n} \sqrt{\lambda_k} \xi_k \zeta_k^\top \tag{1.2}$$

is a singular value decomposition of  $\Sigma^{1/2} X$ , where

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_{N \wedge n} \geq 0 = \lambda_{N \wedge n + 1} = \dots = \lambda_{N \vee n},$$

$\{\xi_k\}_{k=1}^n$  are the left-singular vectors, and  $\{\zeta_k\}_{k=1}^N$  are the right-singular vectors. Then for any deterministic vector  $\mathbf{u} \in \mathbb{R}^n$ , we define the VESD of  $Q_1$  as

$$F_{\mathbf{u}}(x) := \sum_{k=1}^n |\langle \xi_k, \mathbf{u} \rangle|^2 \mathbf{1}_{\{\lambda_k \leq x\}}, \tag{1.3}$$

Here and throughout the rest of this paper, we shall use the notation  $\langle \mathbf{u}, \mathbf{v} \rangle := \mathbf{u}^* \mathbf{v}$  to denote the inner product of two (possibly complex) vectors, where  $\mathbf{u}^*$  is the conjugate transpose of  $\mathbf{u}$ . In the null case with  $\Sigma = I_n$ , it was proved in [3, 11] that  $F_{\mathbf{u}_n}$  converges weakly to the MP law for any sequence of unit vectors  $\mathbf{u}_n$ . In [48], the convergence rate was shown to be  $O(N^{-1/4+\varepsilon})$  almost surely, which was later improved to  $O(N^{-1/2+\varepsilon})$  in [45]. In fact, [45] considered a more general setting where the population matrix  $\Sigma$  is not

necessarily proportional to identity matrix. In this case, it was found that  $F_{\mathbf{u}_n}(x)$  does not converge to the MP law anymore. Instead, it converges to a distribution depending on the vectors  $\mathbf{u}_n$ :

$$F_{1c, \mathbf{u}_n}(x) := \langle \mathbf{u}_n, \mathbf{F}_{1c}(x) \mathbf{u}_n \rangle,$$

where  $\mathbf{F}_{1c}(x)$  is a matrix-valued function determined by  $\Sigma$ . We shall refer to  $\mathbf{F}_{1c}$  or the class of distributions  $F_{1c, \mathbf{u}}$  as the anisotropic MP laws.

As for the ESD theory, the next piece of the VESD theory is the functional CLT for  $F_{\mathbf{u}}$ . More precisely, we are interested in the CLT for random vectors of the form

$$\left( \sqrt{n} \sum_{k=1}^n |\langle \xi_k, \mathbf{u} \rangle|^2 f_s(\lambda_k) - \sqrt{n} \int f_s(x) dF_{1c, \mathbf{u}}(x) \right)_{1 \leq s \leq k}. \quad (1.4)$$

In this paper, we refer to this kind of result as the “linear eigenvector statistics”. In the null case with  $\Sigma = I_n$ , the linear eigenvector statistics was studied in [43] when  $\mathbf{u}$  takes the form  $(\pm n^{-1/2}, \pm n^{-1/2}, \dots, \pm n^{-1/2})$ . Later it was extended to arbitrary unit vector  $\mathbf{u}$  for general analytic functions  $f_s$  in [3]. In [46], the class of functions is extended to include all functions with continuous third order derivatives. In fact, [3] considered slightly more general  $\Sigma$ , requiring that the sequence of vectors  $\mathbf{v}_n$  satisfy the condition

$$\sup_{z \in \mathcal{D}} \sqrt{n} \left| \mathbf{v}_n^\top \frac{1}{1 + m_{2c}(z)\Sigma} \mathbf{v}_n - \int \frac{1}{1 + m_{2c}(z)t} \pi_\Sigma(dt) \right| \rightarrow 0, \quad (1.5)$$

where  $\pi_\Sigma$  is the ESD of  $\Sigma$ ,  $\mathcal{D}$  is an open neighborhood of the support of the MP law in the complex plane, and  $m_{2c}(z)$  is the Stieltjes transform of the MP law—see (2.9). The condition (1.5) is essentially an isotropic condition, under which the VESD  $F_{\mathbf{v}_n}(x)$  still converges to the MP law  $F_{MP}$ , and the information of the vectors  $\mathbf{v}_n$  is missed in the final result. In general, when (1.5) does not hold, it is still unknown whether the functional CLT holds and, if the CLT indeed holds, how do the mean and covariances of the limiting Gaussian vector depend on the covariance matrix  $\Sigma$  and the vectors  $\mathbf{v}_n$ .

The main goal of this paper is to solve this problem. More precisely, we consider sample covariance matrices with completely general population covariance matrices  $\Sigma$  (up to some technical regularity assumptions). We prove that for any sequences of unit vectors  $\mathbf{u}_s \equiv \mathbf{u}_s^{(n)}$ ,  $1 \leq s \leq k$ , the random vector

$$\left( \sqrt{n} \sum_{k=1}^n |\langle \xi_k, \mathbf{u}_s \rangle|^2 f_s(\lambda_k) - \sqrt{n} \int f_s(x) dF_{1c, \mathbf{u}_s}(x) \right)_{1 \leq s \leq k} \quad (1.6)$$

converges to a centered Gaussian vector. Moreover, we obtain an explicit expression for the covariance matrix of the Gaussian vector, which allows us to characterize precisely how the anisotropy of the covariance matrix  $\Sigma$  affect the linear eigenvector statistics. We also extend the results to “local linear eigenvector statistics”. That is, for any fixed  $E > 0$  and scale parameter  $n^{-1} \ll \eta \ll 1$ , we prove that random vector

$$\left( \sqrt{n\eta} \sum_{k=1}^n |\langle \xi_k, \mathbf{u}_s \rangle|^2 \frac{1}{\eta} f_s \left( \frac{\lambda_k - E}{\eta} \right) - \sqrt{n\eta} \int \frac{1}{\eta} f_s \left( \frac{x - E}{\eta} \right) dF_{1c, \mathbf{u}_s}(x) \right)_{1 \leq s \leq k} \quad (1.7)$$

also converges in distribution to a centered Gaussian vector. As an interesting fact, we found that the covariance matrix in the global linear eigenvector statistics depends on the fourth moments of the  $X$  entries, while in the local linear eigenvector statistics it does not. This suggests that the local eigenvector statistics is a little more universal than the global eigenvector statistics, which is a pretty common phenomenon in random matrix theory.

For any  $z \in \mathbb{C} \setminus \mathbb{R}$ , we define the resolvent (or Green's function) of the sample covariance matrix  $Q_1$  as  $R(z) := (Q_1 - z)^{-1}$ . As a byproduct of our proof, we also obtain a CLT for  $R_{\mathbf{u}\mathbf{v}}(z) := \langle \mathbf{u}, R(z) \mathbf{v} \rangle$ , where  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$  are any deterministic unit vectors. Moreover, we prove the CLT with  $z$  having imaginary part  $\eta$  both of the global scale  $\eta \sim 1$  and of the local scale  $n^{-1} \ll \eta \ll 1$ . In the rest of the paper, we shall call  $R_{\mathbf{u}\mathbf{v}}$  the generalized resolvent entries. Besides pure theoretical interest, the CLT for the generalized resolvent entries seems to be very convenient to use in applications.

The VESD was originally introduced in [42, 43] to study the asymptotic properties of the sample eigenvectors. The study of eigenvectors of large dimensional random matrices is generally harder and much less developed compared with the study of eigenvalues. On the other hand, the eigenvectors play an important role in modern statistics due to its role in principal component analysis (PCA), which is now favorably recognized as a powerful technique for dimensionality reduction. The earlier work on sample eigenvectors goes back to Anderson [1], where the author proved that the eigenvectors of the Wishart matrix are asymptotically normal as  $N \rightarrow \infty$  if  $n$  is fixed. In the high dimensional setting, Johnstone [24] proposed the famous spiked model, which is now a standard model for the study of PCA of large random matrices. Later Paul [39] studied the directions of the sample eigenvectors of the spiked model. One can also refer to [13, 34] and references therein for more recent literatures on sparse PCA and spiked covariance matrices.

PCA focuses on the first couple of eigenvectors corresponding to the largest eigenvalues. On the other hand, studying the asymptotic properties of all the eigenvectors at the same time is much harder. In fact, even formulating the terminology "asymptotic property of the eigenmatrix" is far from trivial, since the sample dimension  $n$  is increasing. For this purpose, the VESD serves as a manageable tool for us to discuss about the asymptotic behavior of all the eigenvectors as a whole. In [3, 47, 48], when  $\Sigma = I_n$ , the VESD was used to characterize the asymptotical Haar property of the eigenmatrix, that is, the eigenmatrix is expected to be asymptotically uniformly distributed over the orthogonal group. In general, when  $\Sigma$  is not isotropic, the eigenmatrix is not asymptotically Haar distributed anymore, and our results describes precisely how the VESD behaves along every direction. In addition, by extending to general  $\Sigma$ , our results provide more flexibility to the applications of VESD in statistical estimations of sample covariance matrices. In particular, we will discuss about three possible applications of our results: the detection of weak signals in spiked model when the signal strengths are below the threshold, the estimation of sample eigenvalues with precision of order  $n^{-1/2}$ , and the sphericity test of the population covariance matrix  $\Sigma$ . These will be discussed in detail in Section 3.

Before concluding the introduction, we summarize the main contributions of our work.

- We extend the function CLT for VESD in [3, 46] to anisotropic sample covariance matrices with general population covariance matrices  $\Sigma$ . This result is presented as Theorem 2.6, and it is stronger than the ones in [3, 46] in several senses; see Remark 2.7 below.
- Besides the global linear eigenvector statistics, we also study the local linear eigenvector statistics, and prove the function CLT for VESD on all the scales  $\eta$  such that  $n^{-1} \ll \eta \ll 1$ ; see Theorem 2.8.
- We prove the CLT for the generalized resolvent entries of  $R(z)$ , for  $\eta := \text{Im } z$  both of the global scale  $\eta \sim 1$  and of the mesoscopic scale  $n^{-1} \ll \eta \ll 1$ ; see Theorems 2.10 and 2.11.
- We discuss about three applications of our results: the weak detection of signals in spiked model, the estimation of population eigenvalues, and the sphericity test. Our estimators or test procedures can attain a precision as small as  $n^{-1/2}$ .

This paper is organized as follows. In Section 2, we state the main results of this paper: Theorems 2.6 and 2.8, which give the functional CLT for VESD, and Theorems 2.10 and 2.11, which give the CLT for the generalized resolvent entries. For these results, we assume that the entries of  $X$  have finite  $(8 + \varepsilon)$ -th moments. In Section 3, we discuss about some applications of our results in statistical estimation of sample

covariance matrices. In Section 4, we collect some basic tools that will be used in the proof. Then in Section 5, we prove Theorems 2.10 and 2.11 under a stronger moment assumption that the entries of  $X$  have finite moments up to any order. Based on the results in Section 5, we prove Theorems 2.6 and 2.8 in Section 6 also under the stronger moment assumption. Finally in Section 7, using a Green's function comparison argument we relax the moment assumption to the finite  $(8 + \varepsilon)$ -th moment assumption in the main theorems.

## 2 Definitions and Main Result

### 2.1 The model

We consider a class of separable covariance matrices of the form  $\mathcal{Q}_1 := \Sigma^{1/2} X X^* \Sigma^{1/2}$ , where  $\Sigma$  is a real deterministic non-negative definite symmetric matrix. We assume that  $X = (x_{ij})$  is an  $n \times N$  random matrix with independent entries  $x_{ij}$ ,  $1 \leq i \leq n$ ,  $1 \leq j \leq N$ , satisfying

$$\mathbb{E}x_{ij} = 0, \quad \mathbb{E}|x_{ij}|^2 = 1. \quad (2.1)$$

We will also use the  $N \times N$  matrix  $\mathcal{Q}_2 := X^* \Sigma X$ . We assume that the aspect ratio  $d_N := n/N$  satisfies

$$\tau \leq d_N \leq \tau^{-1}, \quad (2.2)$$

for some constant  $0 < \tau < 1$ . For simplicity of notations, we will often abbreviate  $d_N$  as  $d$  in this paper. We denote the eigenvalues of  $\mathcal{Q}_1$  and  $\mathcal{Q}_2$  in descending order by  $\lambda_1(\mathcal{Q}_1) \geq \dots \geq \lambda_n(\mathcal{Q}_1)$  and  $\lambda_1(\mathcal{Q}_2) \geq \dots \geq \lambda_N(\mathcal{Q}_2)$ . Since  $\mathcal{Q}_1$  and  $\mathcal{Q}_2$  share the same nonzero eigenvalues, we will for simplicity write  $\lambda_j$ ,  $1 \leq j \leq N \vee n$ , to denote the  $j$ -th eigenvalue while keeping in mind that  $\lambda_j = 0$  for  $j > N \wedge n$ .

We assume that  $\Sigma^{1/2}$  has eigendecomposition

$$\Sigma = O^\top \Lambda O, \quad \Lambda = \text{diag}(\sigma_1, \dots, \sigma_n), \quad (2.3)$$

where  $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n \geq 0$  are the eigenvalues. We denote the empirical spectral density of  $\Sigma$  as

$$\pi_\Sigma \equiv \pi_\Sigma^{(n)} := \frac{1}{n} \sum_{i=1}^n \delta_{\sigma_i}. \quad (2.4)$$

We assume that there exists a small constant  $0 < \tau < 1$  such that for all  $N$  large enough,

$$\sigma_1 \leq \tau^{-1}, \quad \pi_\Sigma^{(n)}([0, \tau]) \leq 1 - \tau. \quad (2.5)$$

The first condition means that the operator norms of  $\Sigma$  is bounded by  $\tau^{-1}$ , and the second condition means that the spectrum of  $\Sigma$  does not concentrate at zero.

### 2.2 Resolvents and limiting law

In this paper, we will study the eigenvalue and eigenvector statistics of  $\mathcal{Q}_1$  and  $\mathcal{Q}_2$  through their *resolvents* (or *Green's functions*). In fact, it is equivalent to study the matrices

$$\tilde{\mathcal{Q}}_1(X) := \Lambda^{1/2} O X X^\top O^\top \Lambda^{1/2}, \quad \tilde{\mathcal{Q}}_2(X) \equiv \mathcal{Q}_2(X) = X^\top \Sigma X. \quad (2.6)$$

In this paper, we shall denote the upper half complex plane and the right half real line by

$$\mathbb{C}_+ := \{z \in \mathbb{C} : \text{Im } z > 0\}, \quad \mathbb{R}_+ := (0, \infty).$$

**Definition 2.1** (Resolvents). For  $z = E + i\eta \in \mathbb{C}_+$ , we define the resolvents for  $\tilde{\mathcal{Q}}_{1,2}$  as

$$\mathcal{G}_1(X, z) := \left( \tilde{\mathcal{Q}}_1(X) - z \right)^{-1}, \quad \mathcal{G}_2(X, z) := \left( \tilde{\mathcal{Q}}_2(X) - z \right)^{-1}. \quad (2.7)$$

We denote the empirical spectral density  $\rho^{(n)}$  of  $\tilde{\mathcal{Q}}_1$  and its Stieltjes transform as

$$\rho \equiv \rho^{(n)} := \frac{1}{n} \sum_{i=1}^n \delta_{\lambda_i(\tilde{\mathcal{Q}}_1)}, \quad m(z) \equiv m^{(n)}(z) := \int \frac{1}{x-z} \rho^{(n)}(dx) = \frac{1}{n} \text{Tr} \mathcal{G}_1(z). \quad (2.8)$$

Note  $\rho^{(n)}$  and  $m^{(n)}$  are also the empirical spectral density and its Stieltjes transform for  $\tilde{\mathcal{Q}}_1$ . We also introduce the following two random quantities:

$$m_1(z) \equiv m_1^{(n)}(z) := \frac{1}{N} \sum_{i=1}^n \sigma_i(\mathcal{G}_1(z))_{ii}, \quad m_2(z) \equiv m_2^{(N)}(z) := \frac{1}{N} \sum_{\mu=1}^N (\mathcal{G}_2(z))_{\mu\mu}.$$

If  $d_N \rightarrow d \in (0, \infty)$  and  $\pi_\Sigma$  converges weakly to some distribution  $\pi$  as  $N \rightarrow \infty$ , then it was shown in [35] that the ESD of  $Q_2$  converges in probability to some deterministic distribution, which is called the (deformed) Marčenko-Pastur law. For any  $N$ , we describe the MP law  $F_{2c}^{(N)}$  through its Stieltjes transform

$$m_{2c}(z) \equiv m_{2c}^{(N)}(z) := \int_{\mathbb{R}} \frac{dF_{2c}^{(N)}(x)}{x-z}, \quad z = E + i\eta \in \mathbb{C}_+.$$

We define  $m_{2c}$  as the unique solution to the self-consistent equation

$$\frac{1}{m_{2c}(z)} = -z + d_N \int \frac{t}{1 + m_{2c}(z)t} \pi_\Sigma(dt), \quad (2.9)$$

subject to the conditions that  $\text{Im} m_{2c}(z) \geq 0$  and  $\text{Im}(zm_{2c}(z)) \geq 0$  for  $z \in \mathbb{C}_+$ . It is well known that the functional equation (2.9) has a unique solution that is uniformly bounded on  $\mathbb{C}_+$  under the assumption (2.5) [35]. Letting  $\eta \downarrow 0$ , we can recover the asymptotic eigenvalue density  $\rho_{2c}$  with the inverse formula

$$\rho_{2c}(E) = \pi^{-1} \lim_{\eta \downarrow 0} \text{Im} m_{2c}(E + i\eta). \quad (2.10)$$

From  $\rho_{2c}$ , we can recover the ESD  $F_{2c}^{(N)}$ . Since  $Q_1$  share the same nonzero eigenvalues with  $Q_2$  and has  $n - N$  more (or  $N - n$  less) zero eigenvalues, we can then obtain the asymptotic ESD for  $Q_1$ :

$$F_{1c}^{(n)} = d_N^{-1} F_{2c}^{(N)} + (1 - d_N^{-1}) \mathbf{1}_{[0, \infty)}.$$

Moreover, in [45], it was shown that the VESD  $F_{\mathbf{u}}$  of  $\mathcal{Q}_1$  converges to the anisotropic MP law  $F_{1c, \mathbf{u}} \equiv F_{1c, \mathbf{u}}^{(n)}$ , whose density  $\rho_{1c, \mathbf{u}}$  is given by

$$\rho_{1c, \mathbf{u}}(E) := \mathbf{u}^T \frac{\rho_{2c}(E) \Sigma}{E(1 + 2 \text{Re} m_{2c}(E) \Sigma + |m_{2c}(E)|^2 \Sigma^2)} \mathbf{u}. \quad (2.11)$$

In the rest of this paper, we will often omit the super-indices  $N$  and  $n$  from our notations. The properties of  $m_{2c}$  and  $\rho_{2c}$  have been studied extensively; see e.g. [4, 6, 10, 21, 29, 41, 44]. The following Lemma 2.2 describes the basic structure of  $\rho_{2c}$ . For its proof, one can refer to [29, Appendix A].

**Lemma 2.2** (Support of the deformed MP law). *The density  $\rho_{2c}$  is a disjoint union of connected components:*

$$\text{supp } \rho_{2c} \cap (0, \infty) = \bigcup_{k=1}^L [a_{2k}, a_{2k-1}] \cap (0, \infty), \quad (2.12)$$

where  $L \in \mathbb{N}$  depends only on  $\pi_\Sigma$ . Moreover,  $N \int_{a_{2k}}^{a_{2k-1}} \rho_{2c}(x) dx$  is an integer for any  $k = 1, \dots, L$ , which gives the classical number of eigenvalues in the bulk component  $[a_{2k}, a_{2k-1}]$ . Finally, we have  $a_1 \leq C$  for some constant  $C > 0$  and  $m_{2c}(a_1) \equiv m_{2c}(a_1 + i0_+) \in (-\sigma_1^{-1}, 0)$ .

We shall call  $a_k$  the edges of  $\rho_{2c}$ . Moreover, following the standard notation in random matrix literature, we shall denote the rightmost and leftmost edges as  $\lambda_+ := a_1$  and  $\lambda_- := a_{2L}$ , respectively. To establish our main result, we need to make some extra assumptions on  $\Sigma$ , which takes the form of the following regularity conditions.

**Definition 2.3** (Regularity). (i) Fix a (small) constant  $\tau > 0$ . We say that the edge  $a_k$ ,  $k = 1, \dots, 2L$ , is  $\tau$ -regular if

$$a_k \geq \tau, \quad \min_{l:l \neq k} |a_k - a_l| \geq \tau, \quad \min_i |1 + m_{2c}(a_k) \sigma_i| \geq \tau, \quad (2.13)$$

where  $m_{2c}(a_k) \equiv m_{2c}(a_k + i0_+)$ .

(ii) We say that the bulk component  $(a_{2k}, a_{2k-1})$  is regular if for any fixed  $\tau' > 0$  there exists a constant  $c \equiv c_{\tau'} > 0$  such that the density of  $\rho_{2c}$  in  $[a_{2k} + \tau', a_{2k-1} - \tau']$  is bounded from below by  $c$ .

*Remark 2.4.* The edge regularity conditions (i) has previously appeared (in slightly different forms) in several works on sample covariance matrices [9, 17, 21, 29, 30, 36]. The condition (2.13) ensures a regular square-root behavior of  $\rho_{2c}$  near  $a_k$ . The bulk regularity condition (ii) was introduced in [29], and it imposes a lower bound on the density of eigenvalues away from the edges. These conditions are satisfied by quite general classes of  $\Sigma$ ; see e.g. [29, Examples 2.8 and 2.9].

## 2.3 Main results

For any fixed  $a, b > 0$ , we define the class of functions  $\mathcal{C}^{1,a,b}(\mathbb{R}_+)$  as

$$\mathcal{C}^{1,a,b}(\mathbb{R}_+) := \left\{ f \in \mathcal{C}_c^1(\mathbb{R}_+) : f' \text{ is } a\text{-H\"older continuous uniformly in } x, \right. \\ \left. \text{and } |f(x)| + |f'(x)| \leq C(1 + |x|)^{-(1+b)} \text{ for some constant } C > 0 \right\}.$$

Similar class has been used in [22] for the study of mesoscopic linear eigenvalue statistics. For  $N^{-1+\tau} \leq \eta \leq 1$ ,  $E \in \mathbb{R}$ ,  $f \in \mathcal{C}^{1,a,b}(\mathbb{R}_+)$  and any deterministic vector  $\mathbf{v} \in \mathbb{R}^n$ , we define

$$Z_{\eta,E}(\mathbf{v}, f) := \sqrt{\frac{N}{\eta}} \int f(\eta^{-1}(x - E)) d(F_{\mathbf{v}}(x) - F_{1c,\mathbf{v}}(x)) \\ = \sqrt{\frac{N}{\eta}} \left( \langle \mathbf{v}, f(\eta^{-1}(\mathcal{Q}_1 - E)) \mathbf{v} \rangle - \int_{\lambda_-}^{\lambda_+} f(\eta^{-1}(x - E)) dF_{1c,\mathbf{v}}(x) \right). \quad (2.14)$$

Before stating the main results on the weak convergence of the process  $Z_{\eta,E}(\mathbf{v}, f)$ , we first give the main assumptions.

**Assumption 2.5.** Fix a small constant  $\tau > 0$ .

(i)  $X = (x_{ij})$  is an  $n \times N$  real matrix whose entries are independent random variables satisfying (2.1).

(ii)  $\tau \leq d_N \leq \tau^{-1}$  and  $|d_N - 1| \geq \tau$ .

(iii)  $\Sigma$  is a deterministic positive-definite matrix satisfying (2.5). Moreover, all the edges of  $\rho_{2c}$  are  $\tau$ -regular, and all the bulk components of  $\rho_{2c}$  are regular in the sense of Definition 2.3.

We also need to introduce several notations. First, we denote

$$\kappa_4(i, j) := \mathbb{E}|\sqrt{N}x_{ij}|^4 - 3, \quad (2.15)$$

which is the fourth cumulant of the entry  $\sqrt{N}x_{ij}$ . Then we define two functions  $\alpha, \beta : \mathbb{R}^{2+2n} \rightarrow \mathbb{R}$  as

$$\begin{aligned} \alpha(x_1, x_2, \mathbf{v}_1, \mathbf{v}_2) &\equiv \alpha^{(N)}(x_1, x_2, \mathbf{v}_1, \mathbf{v}_2) \\ &:= \sum_{i=1}^n \frac{\sum_{j=1}^N \kappa_4(i, j)}{3N} \operatorname{Im} \left[ \frac{m_{2c}(x_1)}{x_1} \left( \frac{\Sigma^{1/2}}{1 + m_{2c}(x_1)\Sigma} \mathbf{v}_1 \right)_i \right]^2 \operatorname{Im} \left[ \frac{m_{2c}(x_2)}{x_2} \left( \frac{\Sigma^{1/2}}{1 + m_{2c}(x_2)\Sigma} \mathbf{v}_2 \right)_i \right]^2, \end{aligned} \quad (2.16)$$

and

$$\begin{aligned} \beta(x_1, x_2, \mathbf{v}_1, \mathbf{v}_2) &\equiv \beta^{(N)}(x_1, x_2, \mathbf{v}_1, \mathbf{v}_2) \\ &:= \operatorname{Re} \left[ \frac{m_{2c}(x_1) - \bar{m}_{2c}(x_2)}{x_1 x_2} \left( \mathbf{v}_1^\top \frac{\Sigma}{(1 + m_{2c}(x_1)\Sigma)(1 + \bar{m}_{2c}(x_2)\Sigma)} \mathbf{v}_2 \right)^2 \right] \\ &\quad - \operatorname{Re} \left[ \frac{m_{2c}(x_1) - m_{2c}(x_2)}{x_1 x_2} \left( \mathbf{v}_1^\top \frac{\Sigma}{(1 + m_{2c}(x_1)\Sigma)(1 + m_{2c}(x_2)\Sigma)} \mathbf{v}_2 \right)^2 \right], \end{aligned} \quad (2.17)$$

for  $x_1, x_2 \in \mathbb{R}$  and  $\mathbf{v}_1, \mathbf{v}_2 \in \mathbb{R}^n$ . Here for  $x \in \mathbb{R}$ , we abbreviated  $m_{2c}(x) \equiv m_{2c}(x + i0_+)$ . It is complex with  $\operatorname{Im} m_{2c}(x) = \pi \rho_{2c}(x)$  if  $x \in \operatorname{supp}(\rho_{2c})$  (see (2.10)); otherwise if  $x \notin \operatorname{supp}(\rho_{2c})$ , then  $m_{2c}(x)$  is real.

We are now ready to state the main results. We first consider the convergence of the process  $Z_{\eta, E}(\mathbf{v}, f)$  with  $\eta = 1$ , i.e. the linear eigenvector statistics on the global scale.

**Theorem 2.6.** *Suppose  $d_N, X$  and  $\Sigma$  satisfy Assumption 2.5, and there exists a constant  $c_0 > 0$  such that*

$$\max_{1 \leq i \leq n, 1 \leq j \leq N} \mathbb{E}|\sqrt{N}x_{ij}|^{8+c_0} \leq c_0^{-1}. \quad (2.18)$$

Fix any  $k \in \mathbb{N}$  and constants  $a, b > 0$ . For any sequences of deterministic unit vectors  $\mathbf{v}_1 \equiv \mathbf{v}_1^{(n)}, \dots, \mathbf{v}_k \equiv \mathbf{v}_k^{(n)} \in \mathbb{R}^n$ , and functions  $f_1, \dots, f_k \in \mathcal{C}^{1, a, b}(\mathbb{R}_+)$ , the random vector

$$(Z_{1,0}(\mathbf{v}_i, f_i))_{1 \leq i \leq k} = \left( \sqrt{N} \left( \langle \mathbf{v}_i, f(Q_1) \mathbf{v}_i \rangle - \int_{\lambda_-}^{\lambda_+} f(x) dF_{1c, \mathbf{v}_i}(x) \right) \right)_{1 \leq i \leq k} \quad (2.19)$$

converges weakly to a Gaussian vector  $(\mathcal{G}_1, \dots, \mathcal{G}_k)$  with mean zero and covariance function

$$\begin{aligned} \mathbb{E}(\mathcal{G}_i \mathcal{G}_j) &= \frac{1}{\pi^2} \iint_{x_1, x_2} f_i(x_1) f_j(x_2) \lim_{N \rightarrow \infty} \alpha^{(N)}(x_1, x_2, \mathbf{v}_i, \mathbf{v}_j) dx_1 dx_2 \\ &\quad + \frac{1}{\pi^2} PV \iint_{x_1, x_2} \frac{f_i(x_1) f_j(x_2)}{x_1 - x_2} \lim_{N \rightarrow \infty} \beta^{(N)}(x_1, x_2, \mathbf{v}_i, \mathbf{v}_j) dx_1 dx_2 \\ &\quad + 2 \int_x f_i(x) f_j(x) \lim_{N \rightarrow \infty} \frac{\rho_{2c}^{(N)}(x)}{x^2} \left( \frac{\Sigma}{(1 + m_{2c}^{(N)}(x)\Sigma)(1 + \bar{m}_{2c}^{(N)}(x)\Sigma)} \right)_{\mathbf{v}_i \mathbf{v}_j}^2 dx, \end{aligned} \quad (2.20)$$



as long as all the limits in (2.20) converge. Here PV stands for “principal value”, that is,

$$PV \iint_{x_1, x_2} \frac{g(x_1, x_2)}{x_1 - x_2} dx_1 dx_2 := \lim_{\delta \downarrow 0} \iint_{x_1, x_2} \frac{g(x_1, x_2) (x_1 - x_2)}{(x_1 - x_2)^2 + \delta^2} dx_1 dx_2$$

for any function  $g$  with sufficient regularity.

*Remark 2.7.* Compared to the results in [3, 46], our results are stronger in the following senses.

- (i) We can deal with very general  $\Sigma$ , without assuming  $\Sigma = I_n$  or (1.5).
- (ii) We require weaker regularity on the functions  $f_i$ .
- (iii) It was assumed that the entries  $x_{ij}$  are i.i.d. with  $E|\sqrt{N}x_{ij}|^4 = 3$  in [3], while we see that the nonzero fourth cumulants of the  $X$  entries lead to an extra term in (2.16).
- (iv) We allow for different choices of vectors  $\mathbf{v}_i$  in the random vector (2.19), while [3, 46] only considered the case with  $\mathbf{v}_i = \mathbf{v}$  for all  $i$ . This generalization is important in applications, since if we want to estimate the difference between, say  $Z_{1,0}(\mathbf{v}_1, f_i)$  and  $Z_{1,0}(\mathbf{v}_2, f_i)$ , then it is crucial to know the covariance between them.

We remark that [3] only requires finite fourth moment for the entries of  $X$ , while we need the stronger moment assumption (2.18). However, we notice that the finite 8th moment condition is assumed in [46].

Next we consider the convergence of the process  $Z_{\eta,E}(\mathbf{v}, f)$  with  $\eta \ll 1$ , i.e. the local linear eigenvector statistics. Here and throughout the rest of the paper, for two (random or deterministic) quantities  $a_N$  and  $b_N$  depending on  $N$ , we shall use the notation  $a_N = o(|b_N|)$  or  $|a_N| \ll |b_N|$  to mean that  $|a_N| \leq c_N |b_N|$  for some positive sequence  $c_N \downarrow 0$  as  $N \rightarrow \infty$ .

**Theorem 2.8.** Fix  $E > 0$  and  $N^{-1+c_1} \leq \eta \ll 1$  for some constant  $c_1 > 0$ . Suppose  $d_N$ ,  $X$  and  $\Sigma$  satisfy Assumption 2.5, and there exist a constant  $c_0 > 0$  such that

$$\max_{1 \leq i \leq n, 1 \leq j \leq N} \mathbb{E} |\sqrt{N}x_{ij}|^{a_\eta + c_0} \leq c_0^{-1}, \quad a_\eta := \frac{8}{1 - \log_N \eta}. \quad (2.21)$$

Fix any  $k \in \mathbb{N}$  and constants  $a, b > 0$ . For any sequences of deterministic unit vectors  $\mathbf{v}_1 \equiv \mathbf{v}_1^{(n)}, \dots, \mathbf{v}_k \equiv \mathbf{v}_k^{(n)} \in \mathbb{R}^n$ , and functions  $f_1, \dots, f_k \in \mathcal{C}^{1,a,b}(\mathbb{R}_+)$ , the random vector

$$(Z_{\eta,E}(\mathbf{v}_i, f_i))_{1 \leq i \leq k} = \left( \sqrt{\frac{N}{\eta}} \left( \langle \mathbf{v}_i, f(\eta^{-1}(\mathcal{Q}_1 - E)) \mathbf{v}_i \rangle - \int_{\lambda_-}^{\lambda_+} f(\eta^{-1}(x - E)) dF_{1c, \mathbf{v}_i}(x) \right) \right)_{1 \leq i \leq k}$$

converges weakly to a Gaussian vector  $(\mathcal{G}_1, \dots, \mathcal{G}_k)$  with mean zero and covariance function

$$\mathbb{E}(\mathcal{G}_i \mathcal{G}_j) = \lim_{N \rightarrow \infty} \frac{2\rho_{2c}^{(N)}(E)}{E^2} \left( \frac{\Sigma}{(1 + m_{2c}^{(N)}(E)\Sigma)(1 + \bar{m}_{2c}^{(N)}(E)\Sigma)} \right)_{\mathbf{v}_i \mathbf{v}_j}^2 \int f_i(x) f_j(x) dx \quad (2.22)$$

as long as the limit in (2.22) converges.

*Remark 2.9.* Note that for  $E$  outside  $\text{supp}(\rho_{2c})$ ,  $(\mathcal{G}_1, \dots, \mathcal{G}_k)$  converges to zero in probability. This is due to the fact that is no eigenvalue around  $E$ , and hence both  $f(\eta^{-1}(\lambda_i - E))$ ,  $1 \leq i \leq N \wedge n$ , and  $f(\eta^{-1}(x - E))$ ,  $x \in \text{supp}(\rho_{2c})$ , are of order  $o(1)$ .

We define the following process of resolvents

$$\mathcal{Y}(\mathbf{v}, w) \equiv \mathcal{Y}_{\eta, E}(\mathbf{v}, w) := \sqrt{N\eta} \mathbf{v}^\top \left( R(E + w\eta) + \frac{(E + w\eta)^{-1}}{1 + m_{2c}(E + w\eta)\Sigma} \right) \mathbf{v}, \quad (2.23)$$

where  $R(z) := (\mathcal{Q}_1 - z)^{-1} = O^\top \mathcal{G}_1(z) O$  (recall (2.7)),  $\mathbf{v}$  is a deterministic unit vector in  $\mathbb{R}^n$  and  $w$  is any fixed complex number in  $\mathbb{C}$ . Note that we have  $\mathcal{Y}(\mathbf{v}, \bar{w}) = \overline{\mathcal{Y}(\mathbf{v}, w)}$ . To prove Theorems 2.6 and 2.8, we will first prove an intermediate CLT on finite dimensional distributions of  $\mathcal{Y}(\mathbf{v}, w)$ . We believe these results will be of independent interest.

To state the results, we define the following functions  $\hat{\alpha}, \hat{\beta} : \mathbb{C}^2 \times \mathbb{R}^{2n} \rightarrow \mathbb{C}$ ,

$$\begin{aligned} \hat{\alpha}(z_1, z_2, \mathbf{v}_1, \mathbf{v}_2) &\equiv \hat{\alpha}^{(N)}(z_1, z_2, \mathbf{v}_1, \mathbf{v}_2) \\ &:= \frac{m_{2c}(z_1)m_{2c}(z_2)}{z_1 z_2} \sum_{i=1}^n \frac{\sum_{j=1}^N \kappa_4(i, j)}{3N} \left( \frac{\Sigma^{1/2}}{1 + m_{2c}(z_1)\Sigma} \mathbf{v}_1 \right)_i^2 \left( \frac{\Sigma^{1/2}}{1 + m_{2c}(z_2)\Sigma} \mathbf{v}_2 \right)_i^2, \end{aligned} \quad (2.24)$$

and

$$\begin{aligned} \hat{\beta}(z_1, z_2, \mathbf{v}_1, \mathbf{v}_2) &\equiv \hat{\beta}^{(N)}(z_1, z_2, \mathbf{v}_1, \mathbf{v}_2) \\ &:= 2 \frac{m_{2c}(z_1) - m_{2c}(z_2)}{z_1 z_2 (z_1 - z_2)} \left( \mathbf{v}_1^\top \frac{\Sigma}{(1 + m_{2c}(z_1)\Sigma)(1 + m_{2c}(z_2)\Sigma)} \mathbf{v}_2 \right)^2, \end{aligned} \quad (2.25)$$

for  $z_{1,2} \in \mathbb{C}$  and  $\mathbf{v}_{1,2} \in \mathbb{R}^n$ , where as a convention,  $(z_1 - z_2)^{-1}(m_{2c}(z_1) - m_{2c}(z_2))$  is understood as  $m'_{2c}(z_1)$  when  $z_1 = z_2$ . Denote  $\mathbb{H} := \{z \in \mathbb{C} : \operatorname{Re} z > 0, z \notin \mathbb{R}\}$ . Now we first give the CLT for (2.23) when  $\eta = 1$ .

**Theorem 2.10.** *Suppose  $d_N, X$  and  $\Sigma$  satisfy Assumption 2.5, and there exists a constant  $c_0 > 0$  such that (2.18) holds. Fix any  $k \in \mathbb{N}$  and complex numbers  $z_1, \dots, z_k \in \mathbb{H}$ . For any sequences of deterministic unit vectors  $\mathbf{v}_1 \equiv \mathbf{v}_1^{(n)}, \dots, \mathbf{v}_k \equiv \mathbf{v}_k^{(n)} \in \mathbb{R}^n$ , the random vector  $(\mathcal{Y}_{1,0}(\mathbf{v}_1, z_1), \dots, \mathcal{Y}_{1,0}(\mathbf{v}_k, z_k))$  converges weakly to a complex Gaussian vector  $(\Upsilon_1, \dots, \Upsilon_k)$  with mean zero and covariance*

$$\mathbb{E}\Upsilon_i \Upsilon_j = \lim_{N \rightarrow \infty} \left[ \hat{\alpha}^{(N)}(z_i, z_j, \mathbf{v}_i, \mathbf{v}_j) + \hat{\beta}^{(N)}(z_i, z_j, \mathbf{v}_i, \mathbf{v}_j) \right], \quad 1 \leq i \leq j \leq k, \quad (2.26)$$

as long as the limit in (2.26) converges.

Then we give the local CLT for (2.23) with  $\eta \ll 1$ . For  $E$  outside the spectrum, that is,

$$E \in S_{out}(\tau) := \{E : \operatorname{dist}(E, \in \operatorname{supp} \rho_{2c}) \geq \tau\},$$

we will give a stronger result.

**Theorem 2.11.** *Fix  $E > 0$  and  $N^{-1+c_1} \leq \eta \ll 1$  for some constant  $c_1 > 0$ . Suppose  $d_N, X$  and  $\Sigma$  satisfy Assumption 2.5, and there exists a constant  $c_0 > 0$  such that (2.21) holds. Fix any  $k \in \mathbb{N}$  and complex numbers  $w_1, \dots, w_k \in \mathbb{H}$ . For any sequences of deterministic unit vectors  $\mathbf{v}_1 \equiv \mathbf{v}_1^{(n)}, \dots, \mathbf{v}_k \equiv \mathbf{v}_k^{(n)} \in \mathbb{R}^n$ , the random vector  $(\mathcal{Y}_{\eta, E}(\mathbf{v}_1, w_1), \dots, \mathcal{Y}_{\eta, E}(\mathbf{v}_k, w_k))$  converges weakly to a complex Gaussian vector  $(\Upsilon_1, \dots, \Upsilon_k)$  with mean zero and covariances*

$$\mathbb{E}\Upsilon_i \Upsilon_j = \mathbf{1}(\operatorname{Im} w_i \cdot \operatorname{Im} w_j < 0) \lim_{N \rightarrow \infty} \frac{4i \cdot \operatorname{Im} m_{2c}^{(N)}(E)}{E^2(w_i - w_j)} \left( \frac{\Sigma}{(1 + m_{2c}^{(N)}(E)\Sigma)(1 + \overline{m_{2c}^{(N)}}(E)\Sigma)} \right)_{\mathbf{v}_i \mathbf{v}_j}^2, \quad (2.27)$$

as long as the limit exists.

In addition, if  $E \in S_{out}(\tau)$  for some constant  $\tau > 0$  and (2.18) holds, then for any  $0 < \eta \ll 1$  the random vector  $\eta^{-1/2}(\mathcal{Y}_{\eta,E}(\mathbf{v}_1, w_1), \dots, \mathcal{Y}_{\eta,E}(\mathbf{v}_k, w_k))$  converges weakly to a real Gaussian vector  $(\Upsilon_1, \dots, \Upsilon_k)$  with mean zero and covariance

$$\mathbb{E}\Upsilon_i\Upsilon_j = \lim_{N \rightarrow \infty} \left[ \hat{\alpha}^{(N)}(E, E, \mathbf{v}_i, \mathbf{v}_j) + \hat{\beta}^{(N)}(E, E, \mathbf{v}_i, \mathbf{v}_j) \right], \quad (2.28)$$

as long as the limit exists.

*Remark 2.12.* The reader may notice that given a vector  $\mathbf{v} \in \mathbb{R}^n$ , the term  $\hat{\alpha}^{(N)}(E, E, \mathbf{v}, \mathbf{v})$  can be negative if the fourth cumulants  $\kappa_4(i, j)$  are negative (e.g. for Rademacher entries). However, using  $\kappa_4(i, j) \geq -2$ , we can bound

$$\begin{aligned} \hat{\alpha}(E, E, \mathbf{v}, \mathbf{v}) &\geq -\frac{2m'_{2c}(E)}{3E^2} \sum_{i=1}^n \left( \frac{\Sigma^{1/2}}{1 + m_{2c}(E)\Sigma} \mathbf{v} \right)_i^2 \left( \frac{\Sigma^{1/2}}{1 + m_{2c}(E)\Sigma} \mathbf{v} \right)_i^2 \\ &\geq -\frac{2m'_{2c}(E)}{3E^2} \left( \mathbf{v}^\top \frac{\Sigma}{(1 + m_{2c}(E)\Sigma)(1 + m_{2c}(E)\Sigma)} \mathbf{v} \right)^2 = -\frac{1}{3} \hat{\beta}(E, E, \mathbf{v}, \mathbf{v}), \end{aligned}$$

where in the second step we used that

$$m'_{2c}(E) = \left( \int \frac{\rho_{2c}(x)}{x - E} dx \right)^2 \leq \int \frac{\rho_{2c}(x)}{(x - E)^2} dx = m'_{2c}(E)$$

by Cauchy-Schwarz inequality. Hence the sum  $\hat{\alpha}^{(N)}(E, E, \mathbf{v}, \mathbf{v}) + \hat{\beta}^{(N)}(E, E, \mathbf{v}, \mathbf{v})$  stays positive (as it should be because it is the asymptotic variance of  $\eta^{-1/2}\mathcal{Y}_{\eta,E}(\mathbf{v}, w)$ ).

*Remark 2.13.* For the local statistics, Theorem 2.8 and Theorem 2.11, to hold, we only need the spectrum  $\rho_{2c}$  to behave well locally around  $E$ . In particular, the assumption  $|d_N - 1| \geq 1$  in Assumption 2.5 is not needed as long as  $E$  is away from zero. Moreover, the regularity of  $\Sigma$  is not required to hold for the full spectrum: if  $E$  is in the bulk, we only need that the density of  $\rho_{2c}$  is of order 1 around  $E$ ; if  $E$  is near an edge, we only need that the corresponding edge is regular; if  $E$  is outside the spectrum, we only need that  $E$  is away from the spectrum by a distance of order 1. However, for simplicity of presentation, we did not try to find the optimal assumptions for Theorem 2.8 and Theorem 2.11.

*Remark 2.14.* The results in Theorems 2.6, 2.8, 2.10 and 2.11 can be used to give the CLT for more general quantities

$$\langle \mathbf{u}, f(\eta^{-1}(\mathcal{Q}_1 - E)) \mathbf{v} \rangle, \quad \langle \mathbf{u}, R(E + w\eta) \mathbf{v} \rangle,$$

by using the polarization identity

$$\langle \mathbf{u}, \mathcal{M} \mathbf{v} \rangle = \frac{1}{2} \langle \langle \mathbf{u} + \mathbf{v}, \mathcal{M}(\mathbf{u} + \mathbf{v}) \rangle \rangle - \frac{1}{2} \langle \langle \mathbf{u} - \mathbf{v}, \mathcal{M}(\mathbf{u} - \mathbf{v}) \rangle \rangle$$

for any symmetric matrix  $\mathcal{M}$ . Moreover, by considering real and imaginary parts separately, we can also extend our results to the case with complex vectors.

*Remark 2.15.* Consider a special case where  $f_i$ 's are analytic functions on an open neighborhood  $\mathcal{D}$  of the complex plane containing the real interval  $[\lambda_-, \lambda_+]$ ,  $d_N \rightarrow d \in (0, \infty) \setminus \{1\}$ , and entries of  $X$  are i.i.d. random variables satisfying (2.18) and  $\mathbb{E}|\sqrt{N}x_{ij}|^4 = 3$ . Moreover, suppose that (1.5) holds for a sequence of deterministic unit vectors  $\mathbf{v}_n$ . Then by Theorem 2.10, we get that for fixed  $z_1, z_2 \in \mathbb{H}$ , the covariance between  $\mathcal{Y}_{1,0}(\mathbf{v}_n, z_1)$  and  $\mathcal{Y}_{1,0}(\mathbf{v}_n, z_2)$  converges to

$$\mathbb{E}\Upsilon_i\Upsilon_j = \lim_{N \rightarrow \infty} \frac{2}{z_1 z_2 (z_1 - z_2) (m_{2c}(z_1) - m_{2c}(z_2))} \left[ \mathbf{v}_n^\top \left( \frac{1}{1 + m_{2c}(z_1)\Sigma} - \frac{1}{1 + m_{2c}(z_2)\Sigma} \right) \mathbf{v}_n \right]^2$$

$$\begin{aligned}
&= \lim_{N \rightarrow \infty} \frac{2}{z_1 z_2 (z_1 - z_2) (m_{2c}(z_1) - m_{2c}(z_2))} \left( \int \frac{\pi_\Sigma(dt)}{1 + m_{2c}(z_1)t} - \int \frac{\pi_\Sigma(dt)}{1 + m_{2c}(z_2)t} \right)^2 \\
&= \lim_{N \rightarrow \infty} \frac{2(z_1 m_{2c}(z_1) - z_2 m_{2c}(z_2))^2}{d^2 z_1 z_2 (z_1 - z_2) (m_{2c}(z_1) - m_{2c}(z_2))}, \tag{2.29}
\end{aligned}$$

where we used (2.9) in the last step.

Now we pick a close contour  $\mathcal{C}$  around  $[\lambda_-, \lambda_+]$  in  $\mathcal{D}$ . Then using Cauchy's integral formula, we get

$$Z_{1,0}(\mathbf{v}_n, f_i) = \frac{1}{2\pi i} \oint_{\mathcal{C}} \mathcal{Y}_{0,1}(\mathbf{v}_n, z) dz.$$

Then using (2.29), the covariance between  $Z_{1,0}(\mathbf{v}_n, f_i)$  and  $Z_{1,0}(\mathbf{v}_n, f_j)$  converges to

$$\begin{aligned}
\lim_{N \rightarrow \infty} \mathbb{E} Z_{1,0}(\mathbf{v}_n, f_i) Z_{1,0}(\mathbf{v}_n, f_j) &= -\frac{1}{4\pi^2} \oint_{\mathcal{C}} \oint_{\mathcal{C}} f_i(z_1) f_j(z_2) \lim_{N \rightarrow \infty} \mathbb{E} Y_{0,1}(z_1) Y_{0,1}(z_2) dz_1 dz_2 \\
&= -\frac{1}{2\pi^2} \oint_{\mathcal{C}} \oint_{\mathcal{C}} f_i(z_1) f_j(z_2) \lim_{N \rightarrow \infty} \frac{(z_1 m_{2c}(z_1) - z_2 m_{2c}(z_2))^2}{d^2 z_1 z_2 (z_1 - z_2) (m_{2c}(z_1) - m_{2c}(z_2))} dz_1 dz_2, \tag{2.30}
\end{aligned}$$

if the function  $m_{2c} \equiv m_{2c}^{(N)}$  converges as  $N \rightarrow \infty$ . Of course, there are some technical details missing in the above derivation, but it can be made rigorous. The formula (2.30) recovers the Theorem 2(b) in [3].

*Remark 2.16.* Suppose the setting of Remark 2.15 holds. In addition, we consider sample covariance matrices with null population covariance matrix  $\Sigma = I_n$ , and assume that the vectors  $\mathbf{v}_1, \dots, \mathbf{v}_k$  are all equal to a unit vector  $\mathbf{v}$ . Then the covariance function in (2.20) can be reduced to

$$\mathbb{E}(\mathcal{G}_i \mathcal{G}_j) = \frac{2}{d} \left[ \int_x f_i(x) f_j(x) \rho_c(x) dx - \int_x f_i(x) \rho_c(x) dx \int_x f_j(x) \rho_c(x) dx \right], \tag{2.31}$$

where  $\rho_c(x)$  is the MP density

$$\rho_c(x) = \frac{\sqrt{(x - \lambda_-)(\lambda_+ - x)}}{2\pi dx}, \quad x \in [\lambda_-, \lambda_+], \quad \lambda_{\pm} := (1 \pm \sqrt{d})^2.$$

In [3], a derivation of (2.31) using (2.30) was given assuming that  $f_i$  are analytic. Later in [46], (2.31) was proved for more general  $f_i$  with continuous third order derivatives. For reader's convenience, we now give a derivation of (2.31) from our result (2.20).

The self-consistent equation (2.9) reduces to

$$\frac{1}{m_{2c}(z)} = -z + \frac{d_N}{1 + m_{2c}(z)}, \tag{2.32}$$

and its solution is

$$m_{2c}(z) = \frac{-(z + 1 - d_N) + \sqrt{(z - \lambda_-^{(N)})(z - \lambda_+^{(N)})}}{2z}, \tag{2.33}$$

where  $\lambda_{\pm}^{(N)} := (1 \pm \sqrt{d_N})^2$ . Then for  $\mathbf{v}_1 = \mathbf{v}_2 = \mathbf{v}$ , using (2.17) and (2.32) we can rewrite

$$\frac{\beta(x_1, x_2, \mathbf{v}, \mathbf{v})}{x_1 - x_2} = \frac{1}{d_N^2 x_1 x_2 (x_1 - x_2)} \operatorname{Re} \left[ \frac{(x_1 m_{2c}(x_1) - x_2 \bar{m}_{2c}(x_2))^2}{m_{2c}(x_1) - \bar{m}_{2c}(x_2)} - \frac{(x_1 m_{2c}(x_1) - x_2 m_{2c}(x_2))^2}{m_{2c}(x_1) - m_{2c}(x_2)} \right]. \tag{2.34}$$

We then use the identity

$$(x_1 m_{2c}(x_1) - x_2 \bar{m}_{2c}(x_2))^2 = m_{2c}(x_1) \bar{m}_{2c}(x_2) (x_1 - x_2)^2 + x_1 x_2 (m_{2c}(x_1) - \bar{m}_{2c}(x_2))^2 + (x_1 m_{2c}(x_1) + x_2 \bar{m}_{2c}(x_2)) (x_1 - x_2) (m_{2c}(x_1) - \bar{m}_{2c}(x_2)).$$

Together with a similar estimate for  $(x_1 m_{2c}(x_1) - x_2 m_{2c}(x_2))^2$ , we can simplify (2.34) as

$$\begin{aligned} \frac{\beta(x_1, x_2, \mathbf{v}, \mathbf{v})}{x_1 - x_2} &= \frac{x_1 - x_2}{d_N^2 x_1 x_2} \operatorname{Re} \left[ \frac{m_{2c}(x_1) \bar{m}_{2c}(x_2)}{m_{2c}(x_1) - \bar{m}_{2c}(x_2)} - \frac{m_{2c}(x_1) m_{2c}(x_2)}{m_{2c}(x_1) - m_{2c}(x_2)} \right] \\ &= \frac{-1}{d_N^3 x_1 x_2} \operatorname{Re} [(1 + x_1 m_{2c}(x_1)) x_2 (\bar{m}_{2c}(x_2) - m_{2c}(x_2))] \\ &= -2d_N^{-3} \operatorname{Im} m_{2c}(x_1) \cdot \operatorname{Im} m_{2c}(x_2) \rightarrow -\frac{2\pi^2}{d} \rho_c(x_1) \rho_c(x_2), \end{aligned} \quad (2.35)$$

where in the second step we used (2.32) to get

$$\frac{(x_1 - x_2) m_{2c}(x_1) \bar{m}_{2c}(x_2)}{m_{2c}(x_1) - \bar{m}_{2c}(x_2)} = 1 - \frac{d_N m_{2c}(x_1) \bar{m}_{2c}(x_2)}{(1 + m_{2c}(x_1))(1 + \bar{m}_{2c}(x_2))} = 1 - d_N^{-1} (1 + x_1 m_{2c}(x_1))(1 + x_2 \bar{m}_{2c}(x_2)),$$

and a similar estimate with  $\bar{m}_{2c}(x_2)$  replaced by  $m_{2c}(x_2)$ . On the other hand, we can check that

$$\frac{\rho_{2c}^{(N)}(x)}{x^2 |1 + m_{2c}^{(N)}(x)|^4} = d_N^{-2} \rho_{2c}^{(N)}(x) \rightarrow d^{-1} \rho_c(x).$$

Together with (2.35), this shows that (2.20) can be reduced to (2.31).

### 3 Applications

In this section, we discuss some possible applications of our results. We shall focus on the local eigenvector statistics with

$$0 < \eta \leq n^{-c_1} \quad (3.1)$$

for some constant  $c_1 > 0$ , because they are relatively more convenient to use compared with the global ones. The main takeaway of this section can be summarized as following: more precise information of population eigenvectors leads to more precise estimations of population eigenvalues.

#### 3.1 Weak detection of signals

We consider the famous *Johnstone's spiked model* [25] with

$$\Sigma = I_{n \times n} + \sum_{i=1}^r \tilde{\sigma}_i \mathbf{v}_i \mathbf{v}_i^\top, \quad (3.2)$$

where  $r$  is a fixed integer,  $\{\tilde{\sigma}_i\}_{1 \leq i \leq r}$  give the strengths of the signals, and  $\{\mathbf{v}_i\}_{1 \leq i \leq k}$  are the eigenvectors. Note that following the notations in (2.3), we have  $\sigma_i = 1 + \tilde{\sigma}_i$  for  $1 \leq i \leq r$  and  $\sigma_i = 1$  for  $i > r$ . Suppose

$$\lim_{N \rightarrow \infty} d_N = d \in (0, \infty) \setminus \{1\}. \quad (3.3)$$

It is well known that if  $\tilde{\sigma}_i > \sqrt{d}$ , then the  $i$ -th signal will give rise to an outlier of the eigenvalue spectrum [7, 8, 12, 16]. In this case, this signal can be readily detected and estimated using the results in those

references. Now we focus on the case with weak signals, that is, we assume that  $\tilde{\sigma}_i < \sqrt{d}$  for  $1 \leq i \leq r$ . Then there will be no outliers in the eigenvalue spectrum [7, 8, 12, 16], and it is impossible to estimate the signal strengths using the sample eigenvalues only. However, if we know the eigenvectors  $\mathbf{v}_i$  of the signals, then it is possible to derive some estimates of the signal strengths using our results. For this purpose, we will use the local CLT for  $\langle \mathbf{v}_i, R(E + i\eta) \mathbf{v}_i \rangle$  for some  $E$  outside the spectrum, because by Theorem 2.11 it has better convergence rate  $N^{-1/2}$  compared with the rate  $(N\eta)^{-1/2}$  for  $\langle \mathbf{v}_i, R(E + i\eta) \mathbf{v}_i \rangle$  with  $E$  inside the spectrum and the linear eigenvector statistics  $\eta^{-1} \langle \mathbf{v}_i, f(\eta^{-1}(\mathcal{Q}_1 - E)) \mathbf{v}_i \rangle$ .

Without loss of generality, we assume that  $d_N \equiv d$ . For  $\Sigma$  in (3.2), we can solve from (2.9) that (recall (2.33))

$$m_{2c}(z) = \tilde{m}_{2c}(z) + O(N^{-1}), \quad \tilde{m}_{2c}(z) \equiv \tilde{m}_{2c}(z) := \frac{-(z+1-d) + \sqrt{(z-\lambda_-)(z-\lambda_+)}}{2z}, \quad (3.4)$$

where  $\lambda_{\pm} := (1 \pm \sqrt{d})^2$ . Pick any fixed  $E > \lambda_+$  and  $\eta = 0_+$ . By Theorem 2.11, we know that

$$Y(\mathbf{v}_i, E) := \sqrt{N} \left( R_{\mathbf{v}_i \mathbf{v}_i}(E) + \frac{E^{-1}}{1 + m_{2c}(E)\sigma_i} \right)$$

converges weakly to a real Gaussian random variable  $\Upsilon$  with

$$\mathbb{E}\Upsilon = 0, \quad \mathbb{E}|\Upsilon|^2 = \lim_{N \rightarrow \infty} \gamma_N^2(E, \sigma_i),$$

where

$$\gamma_N^2(E, \sigma_i) := \frac{\sigma_i^2 \tilde{m}_{2c}^2(E)}{E^2 |1 + \tilde{m}_{2c}(E)\sigma_i|^4} \left( \sum_{k=1}^n \frac{\sum_{j=1}^N \kappa_4(k, j)}{3N} \mathbf{v}_i^4(k) + \frac{2\tilde{m}'_{2c}(E)}{\tilde{m}_{2c}^2(E)} \right). \quad (3.5)$$

Then we propose the following estimator of  $\sigma_i$ :

$$\hat{\sigma}_i \equiv \hat{\sigma}_i^{(N)} := -\frac{1}{\tilde{m}_{2c}(E)} \left( \frac{E^{-1}}{R_{\mathbf{v}_i \mathbf{v}_i}(E)} + 1 \right). \quad (3.6)$$

Moreover, for any  $\alpha > 1$ , we choose  $\delta_\alpha$  such that

$$\begin{aligned} \delta_\alpha(E, \hat{\sigma}_i) &:= \alpha \gamma_N(E, \hat{\sigma}_i) \cdot \frac{E|1 + \tilde{m}_{2c}(E)\hat{\sigma}_i|^2}{|\tilde{m}_{2c}(E)|} \\ &= \alpha \hat{\sigma}_i \left[ \frac{1}{3N} \sum_{1 \leq k \leq n, 1 \leq j \leq N} \kappa_4(k, j) \mathbf{v}_i^4(k) + \frac{2\tilde{m}'_{2c}(E)}{|\tilde{m}_{2c}(E)|^2} \right]^{1/2}. \end{aligned} \quad (3.7)$$

Note that using  $\hat{\sigma}_i = \sigma_i + o(1)$  in probability, we have that

$$\sqrt{N} \left| R_{\mathbf{v}_i \mathbf{v}_i}(E) + \frac{E^{-1}}{1 + \tilde{m}_{2c}(E)\sigma} \right| \geq \alpha \gamma_N(E, \hat{\sigma}_i) - o(1) \quad \text{for} \quad |\sigma - \hat{\sigma}_i| \geq \frac{\delta_\alpha}{\sqrt{N}}$$

in probability. Then we can give a confidence interval  $[\hat{\sigma}_i - \delta_\alpha/\sqrt{N}, \hat{\sigma}_i + \delta_\alpha/\sqrt{N}]$  with confidence level  $2\Phi(\alpha) - 1$ , where  $\Phi$  is the CDF for standard normal distribution. Now we collect the above results into the following proposition.

**Proposition 3.1.** *Suppose Assumption 2.5 (i)-(ii), (2.18) and (3.3) hold. Moreover, assume that  $\Sigma$  takes the form (3.2) with fixed strengths  $\tilde{\sigma}_i < \sqrt{d}$  for  $1 \leq i \leq r$ , and  $E > (1 + \sqrt{d})^2$  is fixed. Then for any  $1 \leq i \leq r$ , we have*

$$\mathbb{P} \left( |\hat{\sigma}_i - \sigma_i| \leq \frac{\delta_\alpha(E)}{\sqrt{N}} \right) \geq 2\Phi(\alpha) - 1 - o(1). \quad (3.8)$$

*Proof.* By Theorem 2.11, we have  $\hat{\sigma}_i = \sigma_i + o(1)$  with probability  $1 - o(1)$ , and that

$$\tilde{Y}(\mathbf{v}_i, E) := \gamma_N^{-1}(E, \hat{\sigma}_i) Y(\mathbf{v}_i, E)$$

converges weakly to a standard normal random variable. Then by the definition of  $\delta_\alpha$ ,  $|\hat{\sigma}_i - \sigma_i| \geq \delta_\alpha / \sqrt{N}$  implies that  $|\tilde{Y}(\mathbf{v}_i, E)| > \alpha - o(1)$ . This concludes (3.8).  $\square$

*Remark 3.2.* In order to apply (3.8), we need to have an estimate on  $\delta_\alpha$  or  $\gamma_N(E, \sigma_i)$ . In particular, we need to deal with the term with fourth cumulants. If the entries of  $X$  are i.i.d. Gaussian, then this term is zero. Otherwise, we need other ways to estimate it.

- (i) Suppose the entries of  $X$  are  $x_{ij} = N^{-1/2} a_{ij}$  for a sequence of i.i.d. random variables  $a_{ij}$  with mean zero and variance one. Then the fourth cumulant  $\kappa_4$  of  $a_{ij}$  can be estimated as

$$\hat{\kappa}_4 = \frac{N}{n} \sum_{1 \leq i \leq n, 1 \leq j \leq N} (\Sigma^{1/2} X)_{ij}^4 - 3. \quad (3.9)$$

It is easy to check that for  $\Sigma$  given by (3.2),  $\hat{\kappa}_4 \rightarrow \kappa_4$  almost surely by LLN.

- (ii) If the vector  $\mathbf{v}_i$  is delocalized in the sense  $\|\mathbf{v}_i\|_\infty = o(1)$ , then the term

$$\frac{1}{3N} \sum_{1 \leq k \leq n, 1 \leq j \leq N} \kappa_4(k, j) \mathbf{v}_i^4(k) = o(1)$$

is negligible.

- (iii) Suppose as in the conventional setting of sample covariance matrices,  $X$  has i.i.d. columns but the random variables in the same sample may have different distributions. Then for some fixed large  $p \in \mathbb{N}$ , we can divide the columns of  $X$  into  $p$  independent copies, each of which is a smaller matrix of dimension  $n \times (N/p)$ . For each copy, we can calculate  $R_{\mathbf{v}_i^{(k)} \mathbf{v}_i^{(k)}}(E)$  ( $k$  denotes the  $k$ -th copy). Then we can use the variance of the  $p$  values  $R_{\mathbf{v}_i^{(k)} \mathbf{v}_i^{(k)}}(E)$ ,  $1 \leq k \leq p$ , to estimate  $\gamma_N^2(E, \sigma_i)$  directly. This is useful when one has sufficiently many datas.

We now use simulations to verify the validity of our estimator (3.6). Consider the setting  $n = 2000$ ,  $N = 2n$ , and  $\Sigma = \text{diag}(\sigma, 1, \dots, 1)$ . We use (3.6) with  $\mathbf{v}_i = \mathbf{e}_1$  and  $E = 4$  to find the estimated  $\hat{\sigma}$  for different values of  $\sigma$ , and use (3.7) to calculate  $\delta_2(\hat{\sigma})$  (which corresponds to a 95% confidence interval). In Figure 1, we plot  $\hat{\sigma}$  versus  $\sigma$ . One can see that our estimator performs quite well.

## 3.2 Estimation of population eigenvalues

The idea in Section 3.1 can be extended beyond the spiked model in (3.2). Suppose  $\Sigma$  is a general matrix as in (2.3), and we know that  $\mathbf{v}_i$  is an eigenvector of  $\Sigma$ . We would like to estimate the corresponding eigenvalue. The idea is still to use the local CLT for  $\langle \mathbf{v}_i, R(E + i\eta) \mathbf{v}_i \rangle$  in Theorem 2.11. Now different from Section 3.1, we do not have an explicit expression for  $m_{2c}(E)$ . However, we can estimate it precisely using the averaged local law in Theorem 4.5 below. By (4.24), for  $E \in S_{out}(\tau)$ , we have that for any constant  $\varepsilon > 0$ ,

$$\left| N^{-1} \text{Tr} (X^\top \Sigma X - E)^{-1} - m_{2c}(E) \right| \leq N^{-1+\varepsilon} \quad (3.10)$$

with probability  $1 - o(1)$ . Hence we define an estimator of  $m_{2c}(E)$  as

$$\hat{m}_{2c}(E) := N^{-1} \text{Tr} (X^\top \Sigma X - E)^{-1}. \quad (3.11)$$




Figure 1: The estimated  $\hat{\sigma}$  v.s.  $\sigma$ . The blue lines plot the straight lines  $y = \sigma$  together with the intervals  $[\sigma - 2\delta_2(\hat{\sigma}), \sigma + 2\delta_2(\hat{\sigma})]$ . The red lines give the estimated values  $\hat{\sigma}$  calculated through (3.6). In the left figure, the entries of  $X$  are i.i.d. Gaussian, while in the right figure, the entries of  $X$  are i.i.d. Rademacher.

Taking derivative of  $\hat{m}_{2c}(E)$ , we can also find an estimator for  $m'_{2c}(E)$ :

$$\hat{m}'_{2c}(E) := N^{-1} \text{Tr} (X^\top \Sigma X - E)^{-2}. \quad (3.12)$$

Combining (4.24) with a Cauchy's integral representation of  $\hat{m}'_{2c}(z)$ , we can get that

$$\left| N^{-1} \text{Tr} (X^\top \Sigma X - E)^{-2} - m'_{2c}(E) \right| \leq N^{-1+\varepsilon}. \quad (3.13)$$

Now similar as in Section 3.1, for any fixed  $E$  outside of the spectrum, we propose the following estimator of  $\sigma_i$ ,

$$\hat{\sigma}_i \equiv \hat{\sigma}_i^{(N)} := -\frac{1}{\hat{m}_{2c}(E)} \left( \frac{E^{-1}}{R_{\mathbf{v}_i \mathbf{v}_i}(E)} + 1 \right), \quad (3.14)$$

and define

$$\hat{\delta}_\alpha(E, \hat{\sigma}_i) = \alpha \hat{\sigma}_i \left[ \frac{1}{3N} \sum_{1 \leq k \leq n, 1 \leq j \leq N} \kappa_4(k, j) \mathbf{v}_i^4(k) + \frac{2\hat{m}'_{2c}(E)}{|\hat{m}_{2c}(E)|^2} \right]^{1/2}. \quad (3.15)$$

for any  $\alpha > 1$ . For the estimation of  $\hat{\delta}_\alpha$ , one can refer to Remark 3.2 above (except for item (i)). Then we have a similar proposition as Proposition 3.1.

**Proposition 3.3.** *Suppose Assumption 2.5 and (2.21) hold. Moreover, assume that  $E > \lambda_1 + c$  for some constant  $c > 0$ , where  $\lambda_1$  is the largest eigenvalue of  $X^\top \Sigma X$ . Then for any small constant  $\varepsilon > 0$ , we have*

$$\mathbb{P} \left( |\hat{\sigma}_i - \sigma_i| \leq \frac{\delta_\alpha(E) + \varepsilon}{\sqrt{N}} \right) \geq 2\Phi(\alpha) - 1 \quad (3.16)$$

as long as  $N$  is large enough.

*Proof.* The proof is the same as the one for Proposition 3.1 using (3.11), (3.12) and Theorem 2.11  $\square$



*Remark 3.4.* The last condition of (2.13) actually rules out the existence of outliers. However, our method can be also extended to the scenario with outliers. In [16], it was shown that if the population covariance matrix  $\Sigma$  has a few large spikes about certain threshold, then these spikes will give rise to outliers that lie around certain deterministic locations with fluctuations of order  $N^{-1/2}$ . Those spiked eigenvalues of  $\Sigma$  can be readily estimated using the corresponding outliers. On the other hand, suppose  $\mathbf{v}_i$  is an eigenvector of  $\Sigma$  corresponding to a non-spiked eigenvalue  $\sigma_i$ . Then we can estimate  $\sigma_i$  using (3.14) and (3.16). More precisely, we will choose the  $E$  in (3.14) such that it is away from the spectrum of  $X^\top \Sigma X$  including the outliers. Then the resolvent  $R$  behaves well locally around  $E$ , which is sufficient for the study of local statistics as discussed in Remark 2.13.

*Remark 3.5.* Suppose one wants to estimate  $m_{2c}(E)$  for  $E$  in the bulk of the eigenvalue spectrum (e.g. in applications of Theorem 2.11 for  $E$  in the bulk). Then we can choose  $\eta \gg N^{-1}$ , and by (4.22) we have that for any constant  $\varepsilon > 0$

$$\left| N^{-1} \operatorname{Tr} (X^\top \Sigma X - E - i\eta)^{-1} - m_{2c}(E + i\eta) \right| \leq \frac{N^\varepsilon}{N\eta}$$

with probability  $1 - o(1)$ . On the other hand, we have  $|m_{2c}(E + i\eta) - m_{2c}(E)| = O(\eta)$ . Hence to minimize the error  $\eta + (N\eta)^{-1}$ , we choose  $\eta = N^{-1/2+\tau}$  for some small constant  $\tau > 0$  and define an estimator of  $m_{2c}(E)$  as

$$\hat{m}_{2c}(E) := N^{-1} \operatorname{Tr} (X^\top \Sigma X - E - i\eta)^{-1},$$

To estimate  $m'_{2c}(E)$ , we can choose the estimator as

$$\hat{m}'_{2c}(E) := N^{-1} \operatorname{Tr} (X^\top \Sigma X - E - i\eta)^{-2}.$$

Combining (4.22) with a Cauchy's integral representation of  $\hat{m}'_{2c}(z)$ , we can get that

$$\left| N^{-1} \operatorname{Tr} (X^\top \Sigma X - E - i\eta)^{-2} - m'_{2c}(E) \right| \leq N^{-1+\varepsilon} \eta^{-2} + C\eta = o(1)$$

with probability  $1 - o(1)$ , as long as  $2\tau > \varepsilon$ .

We use simulations to verify the validity of the estimator (3.14). Consider the setting  $n = 2000$ ,  $N = 2n$ , and

$$\Sigma = \operatorname{diag}(\sigma, \underbrace{1, \dots, 1}_{n/2-1}, \underbrace{2, \dots, 2}_{n/2}).$$

We use (3.14) with  $\mathbf{v}_i = \mathbf{e}_1$  and  $E = 6$  to find the estimated  $\hat{\sigma}$  for different values of  $\sigma$ , and use (3.15) to calculate  $\hat{\delta}_2(E, \hat{\sigma})$ . In calculating  $\hat{\sigma}_i$  and  $\hat{\delta}_2(E, \hat{\sigma})$ , we use (3.11) and (3.12) to find the estimated  $\hat{m}_{2c}(E)$  and  $\hat{m}'_{2c}(E)$ . In Figure 2, we plot  $\hat{\sigma}$  versus  $\sigma$ . One can see our estimator performs quite well.

### 3.3 Sphericity test

Our result is also useful even if we do not have precise information of the population eigenvectors. In this section, we discuss the sphericity test for sample covariance matrices, that is,

$$\mathbf{H}_0 : \Sigma \propto I \quad \text{vs.} \quad \mathbf{H}_1 : \Sigma \not\propto I.$$

More generally, we can test whether  $\Sigma = \Sigma_0$  for some given positive definite matrix  $\Sigma_0$  by using the data  $\Sigma_0^{-1/2}(\Sigma^{1/2}X)$ . We now propose and analyze a new test procedure for sphericity based on our results.

We rephrase the testing problem as

$$\mathbf{H}_0 : \tilde{\Sigma} = I \quad \text{vs.} \quad \mathbf{H}_1 : \tilde{\Sigma} \neq I, \tag{3.17}$$




Figure 2: The estimated  $\hat{\sigma}$  v.s.  $\sigma$ . The blue lines plot the straight lines  $y = \sigma$  together with the intervals  $[\sigma - 2\hat{\delta}_2(\hat{\sigma}), \sigma + 2\hat{\delta}_2(\hat{\sigma})]$ . The red lines give the estimated values  $\hat{\sigma}$  calculated through (3.14). In the left figure, the entries of  $X$  are i.i.d. Gaussian, while in the right figure, the entries of  $X$  are i.i.d. Rademacher.

for the rescaled population covariance matrix  $\tilde{\Sigma} := (n^{-1} \text{Tr} \Sigma)^{-1/2} \Sigma$ . We consider the usual setting where the columns of  $X$  are i.i.d. random vectors. If there exist outliers in the eigenvalue spectrum, then we immediately reject  $\mathbf{H}_0$ . Hence we only study the case without outliers, and use Theorem 2.11 as our tool. Given an allowable error  $\omega$  (say  $\omega = 5\%$ ), our test procedure is as follows.

**Step 1: Rescaling.** Given a data matrix  $\Sigma^{1/2} X$ , we can estimate  $n^{-1} \text{Tr} \Sigma$  as

$$\sigma^2 := \frac{1}{n} \sum_{1 \leq i \leq n, 1 \leq j \leq N} (\Sigma^{1/2} X)_{ij}^2.$$

By concentration, we can check that for any constant  $\varepsilon > 0$ ,

$$\sigma^2 = \frac{1}{n} \text{Tr} \Sigma \cdot (1 + O(N^{-1+\varepsilon})) \quad (3.18)$$

with probability  $1 - o(1)$ . Hence we will use  $\sigma^2$  as an estimator of  $n^{-1} \text{Tr} \Sigma$ , and consider the rescaled data matrix  $W := \sigma^{-1} \Sigma^{1/2} X$ .

**Step 2: Choosing test vectors and parameter  $E$ .** Choose two deterministic unit test vectors  $\mathbf{u}$  and  $\mathbf{v}$ . Choose a fixed  $E > \lambda_1 + c$  for some constant  $c > 0$ , where  $\lambda_1$  is the largest eigenvalue of  $W^\top W$ .

**Step 3: Estimating the variance.** Calculate the estimator  $\hat{m}_{2c}(E)$  and  $\hat{m}'_{2c}(E)$  as (recall (3.11))

$$\hat{m}_{2c}(E) := N^{-1} \text{Tr} (W^\top W - E)^{-1}, \quad \hat{m}'_{2c}(E) := N^{-1} \text{Tr} (W^\top W - E)^{-2}.$$

We estimate the fourth cumulants of the matrix entries of  $W$  by

$$\hat{\kappa}_4(i) = N \sum_{1 \leq j \leq N} W_{ij}^4 - 3, \quad 1 \leq i \leq n.$$

Then we estimate  $\gamma_N^2(E)$  in (3.5) from above as

$$\Gamma_N^2(E) := \frac{\hat{m}_{2c}^2(E)}{E^2 |1 + \hat{m}_{2c}(E)|^4} \left( \frac{1}{3} \max_{1 \leq k \leq n} \max\{\hat{\kappa}_4(k), 0\} + \frac{2\hat{m}'_{2c}(E)}{\hat{m}_{2c}^2(E)} \right).$$

If there is a priori bound on the fourth cumulants, then one can skip the estimation of the fourth cumulants.

**Step 4: Checking the sphericity.** We calculate

$$\mathcal{E} := \sqrt{N} |R_{\mathbf{u}\mathbf{u}}(E) - R_{\mathbf{v}\mathbf{v}}(E)|, \quad R(E) := (WW^\top - E)^{-1}.$$

Let  $\alpha > 0$  be the number such that  $2(1 - \Phi(\alpha)) = \omega$ . If  $\mathcal{E} < \sqrt{2}\alpha\Gamma_N(E)$ , then we accept the hypothesis  $\mathbf{H}_0$ ; otherwise, we reject it.

The idea of the above procedure has been very briefly discussed in [45]. Here we give a more precise analysis of it based on the results of this paper.

**Proposition 3.6.** *Suppose Assumption 2.5 and (2.18) hold. Moreover, assume that the columns of  $X$  are i.i.d. random vectors. Then the type I test error of the above procedure is*

$$\mathbb{P}(\text{rejection of } \mathbf{H}_0 | \tilde{\Sigma} = I) \leq \omega + o(1). \quad (3.19)$$

On the other hand, conditioning on

$$\left| \mathbf{u}^\top \frac{1}{1 + m_{2c}(E)\tilde{\Sigma}} \mathbf{u} - \mathbf{v}^\top \frac{1}{1 + m_{2c}(E)\tilde{\Sigma}} \mathbf{v} \right| \gg N^{-1/2}, \quad (3.20)$$

we have

$$\mathbb{P}(\text{acceptance of } \mathbf{H}_0 | \mathbf{u}, \mathbf{v} \text{ satisfy (3.20)}) = o(1). \quad (3.21)$$

*Proof.* We introduce the matrix  $\tilde{W} := \tilde{\Sigma}^{1/2}X$  and the resolvent

$$\tilde{R}(E) := (\tilde{W}\tilde{W}^\top - E)^{-1}.$$

By (3.18), it is easy to check that

$$\|\tilde{R}(E) - R(E)\| = O(N^{-1+\varepsilon}) \quad (3.22)$$

with probability  $1 - o(1)$ . Hence  $\tilde{R}(E)$  and  $R(E)$  satisfies the same local statistics under the  $\sqrt{N}$  scaling as in Theorem 2.11.

We first prove (3.19). Suppose  $\tilde{\Sigma} = I$ , and  $\sqrt{N} |R_{\mathbf{u}\mathbf{u}}(E) - R_{\mathbf{v}\mathbf{v}}(E)| \geq \sqrt{2}\alpha\Gamma_N(E)$ . Then with (3.22) we have

$$\sqrt{N} \left| \tilde{R}_{\mathbf{u}\mathbf{u}}(E) - \tilde{R}_{\mathbf{v}\mathbf{v}}(E) \right| \geq \sqrt{2}\alpha\Gamma_N(E) - O(N^{-1/2+\varepsilon}) \quad (3.23)$$

with probability  $1 - o(1)$ . For now we assume that  $\mathbf{v}$  is orthogonal to  $\mathbf{u}$ . Then by Theorem 2.11, the two random variables

$$\gamma_N^{-1}(E, \mathbf{u}) \left( \tilde{R}_{\mathbf{u}\mathbf{u}}(E) + \frac{E^{-1}}{1 + \tilde{m}_{2c}(E)} \right), \quad \text{and} \quad \gamma_N^{-1}(E, \mathbf{v}) \left( \tilde{R}_{\mathbf{v}\mathbf{v}}(E) + \frac{E^{-1}}{1 + \tilde{m}_{2c}(E)} \right) \quad (3.24)$$

converge weakly to two independent Gaussian random variables with mean zero and variance one, where  $\tilde{m}_{2c}(E)$  is as in (3.4). Here we defined for any unit vector  $\mathbf{u} \in \mathbb{R}^n$ ,

$$\gamma_N^2(E, \mathbf{u}) := \frac{\tilde{m}_{2c}^2(E)}{E^2 |1 + \tilde{m}_{2c}(E)|^4} \left( \frac{1}{3} \sum_{1 \leq k \leq n} \kappa_4(k) \mathbf{u}^4(k) + \frac{2\tilde{m}'_{2c}(E)}{\tilde{m}_{2c}^2(E)} \right),$$

where  $\kappa_4(k)$  is the fourth cumulant of  $\sqrt{N}x_{kj}$  for the entries in the  $k$ -th row of  $X$ . By (3.22), (3.10) and (3.13),  $\tilde{m}_{2c}(E)$  and  $\tilde{m}'_{2c}(E)$  can be approximated by  $\hat{m}_{2c}(E)$  and  $\hat{m}'_{2c}(E)$  up to a negligible error of order  $O(N^{-1+\varepsilon})$ . Moreover, under the moment condition (2.18), we have

$$\max_{1 \leq k \leq n} |\hat{\kappa}_4(k) - \tilde{\kappa}_4(k)| = o(1) \quad \text{with probability } 1 - o(1),$$

which can be proved using a standard cut-off argument, see Remark 3.7 below. Thus we obtain that with probability  $1 - o(1)$ ,  $\gamma_N^2(E, \mathbf{u}) \leq \Gamma_N^2(E) + o(1)$  for any unit vector  $\mathbf{u} \in \mathbb{R}^n$ . Combining the above estimates, we obtain that  $\Gamma_N^{-1}(E)\sqrt{N} \left| \tilde{R}_{\mathbf{u}\mathbf{u}}(E) - \tilde{R}_{\mathbf{v}\mathbf{v}}(E) \right|$  converges weakly to a Gaussian random variable with mean zero and variance  $\leq 2$ . Hence with (3.23), we get

$$\begin{aligned} \mathbb{P} \left( \sqrt{N} |R_{\mathbf{u}\mathbf{u}}(E) - R_{\mathbf{v}\mathbf{v}}(E)| \geq \sqrt{2}\alpha\Gamma_N(E) \right) &\leq \mathbb{P} \left( \sqrt{N} |(R_0)_{\mathbf{u}\mathbf{u}}(E) - (R_0)_{\mathbf{v}\mathbf{v}}(E)| \geq \sqrt{2}\alpha\Gamma_N(E) - o(1) \right) \\ &\leq 2(1 - \Phi(\alpha)) + o(1) = \omega + o(1). \end{aligned}$$

In general, if  $\mathbf{u}$  and  $\mathbf{v}$  are not orthogonal, then by Theorem 2.10 the two random variables in (3.24) converge weakly to two Gaussian random variables  $\mathcal{G}_1$  and  $\mathcal{G}_2$  with nonzero covariance between them. Then using the same arguments as above, we obtain that

$$\mathbb{P} \left( \sqrt{N} |R_{\mathbf{u}\mathbf{u}}(E) - R_{\mathbf{v}\mathbf{v}}(E)| \geq \sqrt{2}\alpha\Gamma_N(E) \right) \leq \mathbb{P} \left( |\mathcal{G}_1 - \mathcal{G}_2| \geq \sqrt{2}\alpha \right) + o(1).$$

This concludes (3.19) together with the fact that

$$\mathbb{P} \left( |\mathcal{G}_1 - \mathcal{G}_2| \geq \sqrt{2}\alpha \right) \leq \mathbb{P}(|\mathcal{G}| \geq \alpha) = 2(1 - \Phi(\alpha)), \quad (3.25)$$

where  $\mathcal{G}$  is a standard normal random variable.

For (3.21), suppose (3.20) holds. Then by Theorem 2.11 it implies that the mean  $\sqrt{N}(R_{\mathbf{u}\mathbf{u}}(E) - R_{\mathbf{v}\mathbf{v}}(E))$  diverges to infinity, while the variance is always bounded by  $O(1)$ . Hence we conclude (3.21).  $\square$

*Remark 3.7.* Suppose we have a sequence of  $N$  i.i.d. random variables  $x_i$ ,  $1 \leq i \leq N$ , with mean zero and finite  $(2+c)$ -moments:  $\mathbb{E}|x_i|^{2+c} = O(1)$ . We claim that

$$\mathbb{P}(S_N/N \leq N^{-\varepsilon}) \leq N^{-1-\varepsilon} \quad (3.26)$$

for some small constant  $\varepsilon > 0$ , where  $S_N := x_1 + \dots + x_N$ . To show this claim, we introduce a cut-off  $\tilde{x}_i = x_i \mathbf{1}_{|x_i| \leq N^{1-\delta}}$  for some sufficiently small constant  $\delta > 0$  such that  $(1-\delta)(2+c) < 2+\varepsilon$ . Then we have

$$\mathbb{P}(\tilde{x}_i \neq x_i \text{ for some } i) = O(N^{-1-\varepsilon})$$

using Markov's inequality together with a union bound. Then for  $\tilde{S}_N := \tilde{x}_1 + \dots + \tilde{x}_N$ , one can bound  $\mathbb{P}(\tilde{S}_N/N)$  by calculating the fourth moment of  $\tilde{S}_N/N$ . We omit the rest of the details.

One can observe that the efficiency of our procedure depends crucially on the test vectors. If there is a priori information on the possible directions of signals in  $\Sigma$ , then encoding it into the choice of test vectors may greatly improve the efficiency of our procedure. For example, if one knows that the deviation from the sphericity may happen on the  $i$ -th and  $j$ -th features, then it is natural to take the standard basis vector  $\mathbf{e}_i$  and  $\mathbf{e}_j$ , and calculate  $\mathcal{E} = |R_{ii}(E) - R_{jj}(E)|$ . Or one can choose  $(\mathbf{e}_i \pm \mathbf{e}_j)/\sqrt{2}$ , and calculate  $\mathcal{E} = |R_{ij}(E)|$ . The standard basis vector may fail when  $\Sigma$  has some delocalized eigenvectors whose entries have amplitudes of order  $n^{-1/2}$ . For example, if  $\Sigma = I_n + a\mathbf{e}\mathbf{e}^\top$  for some small  $a > 0$  and  $\mathbf{e} = n^{-1/2}(1, \dots, 1)^\top$ , then using  $|R_{ii}(E) - R_{jj}(E)|$  or  $R_{ij}(E)$  may fail to find the signal  $a\mathbf{e}\mathbf{e}^\top$ . Suppose we know the components of certain

$x$	$n^{-1/2}$	0.1	0.125	0.15	0.175	0.20	0.225	0.25	0.5
$\mathbf{e}_1, \mathbf{e}_2$	0.966	0.808	0.571	0.314	0.105	0.032	0.003	0.002	0.000
$(\mathbf{e}_1 \pm \mathbf{e}_2)/\sqrt{2}$	1.000	0.992	0.961	0.947	0.903	0.862	0.815	0.744	0.391
$\mathbf{e}_1, \mathbf{e}$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Table 1: Frequency of misestimation over  $10^3$  simulations for different values of  $x$  in (3.27). Here we take  $a = 1$ ,  $E = 4$  and allowable error  $\omega = 5\%$ .

$a$	0.025	0.05	0.075	0.1	0.125	0.15	0.25	0.50	0.75	1.00
$\mathbf{e}_1, \mathbf{e}_2$	0.985	0.981	0.976	0.972	0.971	0.977	0.952	0.741	0.012	0.032
$\mathbf{e}_1, \mathbf{e}$	0.924	0.778	0.464	0.229	0.068	0.009	0.000	0.000	0.000	0.000

Table 2: Frequency of misestimation over  $10^3$  simulations for different values of  $a$  in (3.27). Here we take  $x = 0.2$ ,  $E = 4$  and allowable error  $\omega = 5\%$ .

signal vector are all positive, then it is natural to choose  $\mathbf{e}$  as one of the test vectors. More generally, if we know certain components (say with indices in  $\mathbb{T}$ ) of the signal vector are positive and the rest are negative, then one can probably choose a test vector  $\mathbf{u}$  such that  $\mathbf{u}(i) = n^{-1/2}\mathbf{1}_{i \in \mathbb{T}} - n^{-1/2}\mathbf{1}_{i \notin \mathbb{T}}$ .

Now we use some Monte-Carlo simulations to verify the validity of our test procedure. Consider the setting  $n = 500$ ,  $N = 2n$ , the entries of  $X$  are i.i.d. Gaussian, and

$$\Sigma = I_n + a \mathbf{v}_x \mathbf{v}_x^\top, \quad \mathbf{v}_x := \left( x, \sqrt{\frac{1-x^2}{n-1}}, \dots, \sqrt{\frac{1-x^2}{n-1}} \right). \quad (3.27)$$

We take  $E = 4$  and allowable error  $\omega = 5\%$ , which corresponds to the threshold  $\sqrt{2}\alpha\Gamma_N(E)$  with  $\alpha = 2$ . In Table 1, we consider the sphericity test for  $\Sigma$  and compute the frequencies of misestimation for different choices of  $x$ . For the first row, we used test vectors  $\mathbf{e}_1$  and  $\mathbf{e}_2$ , and for the second row we used test vectors  $(\mathbf{e}_1 \pm \mathbf{e}_2)/\sqrt{2}$ . If we used the test vectors  $\mathbf{e}_1$  and  $\mathbf{e}$ , then the frequencies of misestimation are all zero as shown by the last row. One can understand the results by checking the difference  $|\langle \mathbf{u}^\top \mathbf{v}_x \rangle|^2 - |\langle \mathbf{v}^\top \mathbf{v}_x \rangle|^2$  for the above three choices of  $\{\mathbf{u}, \mathbf{v}\}$ . In Table 2, we fix  $x = 0.2$  and compute the frequencies of misestimation for different choices of  $a$ . For the first row, we used test vectors  $\mathbf{e}_1$  and  $\mathbf{e}_2$ , and for the second row we used test vectors  $\mathbf{e}_1$  and  $\mathbf{e}$ .

From Table 1, we observe that the choices of the test vectors affect the test efficiency a lot. From Table 2, we see that our procedure can detect very small deviation from the sphericity, as long as we choose the proper test vectors.

## 4 Basic tools

The rest of the paper is devoted to the proof of the main theorems—Theorems 2.6, 2.8, 2.10 and 2.11. In this section, we introduce some notations and collect some basic tools that will be used in the proof.

With the notations in (2.7), the Stieltjes transforms of  $F_{\mathbf{v}}$  are equal to  $\langle \mathbf{u}, \mathcal{G}_1(X, z)\mathbf{u} \rangle$ , where  $\mathbf{u} := O\mathbf{v}$ . One of the most basic tools of the proof is the following asymptotic estimate

$$\langle \mathbf{u}, \mathcal{G}_1(X, z)\mathbf{u} \rangle \approx m_{1c, \mathbf{u}}(z), \quad (4.1)$$

which we shall refer to as the anisotropic local law. More precisely, an *anisotropic local law* is an estimate of the form (4.1) for all  $\text{Im } z \gg N^{-1}$ . Such local law has been established in [11, 28, 29, 49] for sample covariance matrices, assuming certain moment conditions on the matrix entries.

The anisotropic local law can be stated in a simple and unified fashion using the following  $(N+n) \times (N+n)$  symmetric matrix  $H$ :

$$H := \begin{pmatrix} 0 & \Lambda^{1/2} OX \\ (\Lambda^{1/2} OX)^\top & 0 \end{pmatrix}. \quad (4.2)$$

We define the resolvent of  $H$  as

$$G(X, z) := \begin{pmatrix} -I_n & \Lambda^{1/2} OX \\ (\Lambda^{1/2} OX)^\top & -zI_n \end{pmatrix}^{-1}, \quad z \in \mathbb{C}_+. \quad (4.3)$$

Using Schur complement formula, it is easy to check that

$$G = \begin{pmatrix} z\mathcal{G}_1 & \mathcal{G}_1(\Lambda^{1/2} OX) \\ (\Lambda^{1/2} OX)^\top \mathcal{G}_1 & \mathcal{G}_2 \end{pmatrix} = \begin{pmatrix} z\mathcal{G}_1 & (\Lambda^{1/2} OX)\mathcal{G}_2 \\ \mathcal{G}_2(\Lambda^{1/2} OX)^\top & \mathcal{G}_2 \end{pmatrix}. \quad (4.4)$$

Thus a control of  $G$  yields directly a control of the resolvents  $\mathcal{G}_1$  and  $\mathcal{G}_2$ . For simplicity of notations, we define the index sets

$$\mathcal{I}_1 := \{1, \dots, n\}, \quad \mathcal{I}_2 := \{n+1, \dots, n+N\}, \quad \mathcal{I} := \mathcal{I}_1 \cup \mathcal{I}_2.$$

We shall consistently use the latin letters  $i, j \in \mathcal{I}_1$ , greek letters  $\mu, \nu \in \mathcal{I}_2$ , and  $\mathbf{a}, \mathbf{b} \in \mathcal{I}$ . Then we label the indices of  $X$  as  $X = (X_{i\mu} : i \in \mathcal{I}_1, \mu \in \mathcal{I}_2)$ . For simplicity, given a vector  $\mathbf{v} \in \mathbb{C}^{\mathcal{I}_1, 2}$ , we always identify it with its natural embedding in  $\mathbb{C}^{\mathcal{I}}$ . For example, we shall identify  $\mathbf{v} \in \mathbb{C}^{\mathcal{I}_1}$  with  $\begin{pmatrix} \mathbf{v} \\ \mathbf{0}_N \end{pmatrix}$ .

Now we introduce the spectral decomposition of  $G$ . Let

$$\Lambda^{1/2} OX = \sum_{k=1}^{n \wedge N} \sqrt{\lambda_k} \xi_k \zeta_k^\top,$$

be a singular value decomposition of  $\Lambda^{1/2} OX$  (where the notations in (1.2) are slightly abused here). Then using (4.4), we can get that for  $i, j \in \mathcal{I}_1$  and  $\mu, \nu \in \mathcal{I}_2$ ,

$$G_{ij} = \sum_{k=1}^n \frac{z \xi_k(i) \xi_k^\top(j)}{\lambda_k - z}, \quad G_{\mu\nu} = \sum_{k=1}^N \frac{\zeta_k(\mu) \zeta_k^\top(\nu)}{\lambda_k - z}, \quad (4.5)$$

$$G_{i\mu} = \sum_{k=1}^{n \wedge N} \frac{\sqrt{\lambda_k} \xi_k(i) \zeta_k^\top(\mu)}{\lambda_k - z}, \quad G_{\mu i} = \sum_{k=1}^{n \wedge N} \frac{\sqrt{\lambda_k} \zeta_k(\mu) \xi_k^\top(i)}{\lambda_k - z}. \quad (4.6)$$

With these spectral decompositions, one can obtain the bound

$$\|G(z)\| \leq C(\text{Im } z)^{-1} \quad (4.7)$$

for some constant  $C > 0$ . Furthermore, from (4.5) and (4.6) it is also easy to derive the following identities, which we shall refer to as Ward's identities. For the proof, one can refer to Lemma 6.1 of [49].

**Lemma 4.1.** *Let  $\{\mathbf{u}_i\}_{i \in \mathcal{I}_1}$  and  $\{\mathbf{v}_\mu\}_{\mu \in \mathcal{I}_2}$  be orthonormal basis vectors in  $\mathbb{R}^{\mathcal{I}_1}$  and  $\mathbb{R}^{\mathcal{I}_2}$ , respectively. Let  $\mathbf{x} \in \mathbb{C}^{\mathcal{I}_1}$  and  $\mathbf{y} \in \mathbb{C}^{\mathcal{I}_2}$ . Then we have*

$$\sum_{i \in \mathcal{I}_1} |G_{\mathbf{x}\mathbf{u}_i}|^2 = \sum_{i \in \mathcal{I}_1} |G_{\mathbf{u}_i \mathbf{x}}|^2 = \frac{|z|^2}{\eta} \text{Im} \left( \frac{G_{\mathbf{x}\mathbf{x}}}{z} \right), \quad (4.8)$$

$$\sum_{\mu \in \mathcal{I}_2} |G_{\mathbf{y}\mathbf{v}_\mu}|^2 = \sum_{\mu \in \mathcal{I}_2} |G_{\mathbf{v}_\mu \mathbf{y}}|^2 = \frac{\text{Im } G_{\mathbf{y}\mathbf{y}}}{\eta}, \quad (4.9)$$

$$\sum_{i \in \mathcal{I}_1} |G_{\mathbf{y}\mathbf{u}_i}|^2 = \sum_{i \in \mathcal{I}_1} |G_{\mathbf{u}_i \mathbf{y}}|^2 = G_{\mathbf{y}\mathbf{y}} + \frac{\bar{z}}{\eta} \text{Im } G_{\mathbf{y}\mathbf{y}}, \quad (4.10)$$

$$\sum_{\mu \in \mathcal{I}_2} |G_{\mathbf{x}\mathbf{v}_\mu}|^2 = \sum_{\mu \in \mathcal{I}_2} |G_{\mathbf{v}_\mu \mathbf{x}}|^2 = \frac{G_{\mathbf{x}\mathbf{x}}}{z} + \frac{\bar{z}}{\eta} \text{Im} \left( \frac{G_{\mathbf{x}\mathbf{x}}}{z} \right). \quad (4.11)$$

We will use the following notion of stochastic domination, which was first introduced in [18] and subsequently used in many works on random matrix theory. It simplifies the presentation of the results and their proofs by systematizing statements of the form “ $\xi$  is bounded with high probability by  $\zeta$  up to a small power of  $N$ ”.

**Definition 4.2** (Stochastic domination). *(i) Let*

$$\xi = \left( \xi^{(N)}(u) : N \in \mathbb{N}, u \in U^{(N)} \right), \quad \zeta = \left( \zeta^{(N)}(u) : N \in \mathbb{N}, u \in U^{(N)} \right)$$

*be two families of nonnegative random variables, where  $U^{(N)}$  is a possibly  $N$ -dependent parameter set. We say  $\xi$  is stochastically dominated by  $\zeta$ , uniformly in  $u$ , if for any small constant  $\varepsilon > 0$  and large constant  $D > 0$ ,*

$$\sup_{u \in U^{(N)}} \mathbb{P} \left[ \xi^{(N)}(u) > N^\varepsilon \zeta^{(N)}(u) \right] \leq N^{-D}$$

*for large enough  $N \geq N_0(\varepsilon, D)$ .*

*(ii) If  $\xi$  is stochastically dominated by  $\zeta$ , uniformly in  $u$ , we use the notation  $\xi < \zeta$ . Moreover, if for some complex family  $\xi$  we have  $|\xi| < \zeta$ , we also write  $\xi < \zeta$  or  $\xi = \mathbf{O}_<(\zeta)$ .*

*(iii) We say that an event  $\Xi$  holds with high probability if for any constant  $D > 0$ ,  $\mathbb{P}(\Xi) \geq 1 - N^{-D}$  for large enough  $N$ .*

The following lemma collects basic properties of stochastic domination, which will be used tacitly throughout the proof.

**Lemma 4.3** (Lemma 3.2 in [11]). *Let  $\xi$  and  $\zeta$  be families of nonnegative random variables.*

*(i) Suppose that  $\xi(u, v) < \zeta(u, v)$  uniformly in  $u \in U$  and  $v \in V$ . If  $|V| \leq N^C$  for some constant  $C$ , then  $\sum_{v \in V} \xi(u, v) < \sum_{v \in V} \zeta(u, v)$  uniformly in  $u$ .*

*(ii) If  $0 \leq \xi_1(u) < \zeta_1(u)$  and  $0 \leq \xi_2(u) < \zeta_2(u)$  uniformly in  $u \in U$ , then  $\xi_1(u)\xi_2(u) < \zeta_1(u)\zeta_2(u)$  uniformly in  $u \in U$ .*

*(iii) Suppose that  $\Psi(u) \geq N^{-C}$  is deterministic and  $\xi(u)$  satisfies  $\mathbb{E}\xi(u)^2 \leq N^C$  for all  $u$ . Then if  $\xi(u) < \Psi(u)$  uniformly in  $u$ , we have  $\mathbb{E}\xi(u) < \Psi(u)$  uniformly in  $u$ .*

Throughout the rest of this paper, we will consistently use the notation  $z = E + i\eta$  for the spectral parameter  $z$ . We define the spectral domain

$$\mathbf{D} \equiv \mathbf{D}(\omega, N) := \{z \in \mathbb{C}_+ : |z| \geq \omega, N^{-1+\omega} \leq \eta \leq \omega^{-1}\}, \quad (4.12)$$

for some small constant  $\omega > 0$ . We will also consider a domain that is outside  $\text{supp}(\rho_{2c})$ :

$$\mathbf{D}_{\text{out}} \equiv \mathbf{D}_{\text{out}}(\omega, N) := \{z \in \mathbb{C}_+ : |z| \geq \omega, 0 < \eta \leq \omega^{-1}, \text{dist}(E, \text{supp}(\rho_{2c})) \geq \omega\}. \quad (4.13)$$

Recall the condition (2.13), we can take  $\omega$  to be sufficiently small such that  $\omega \leq \lambda_-/2$ . Define the distance to the spectral edges as

$$\kappa := \min_{1 \leq k \leq 2L} |E - a_k|. \quad (4.14)$$

Then we have the following estimates for  $m_{2c}$ :

$$|m_{2c}(z)| \lesssim 1, \quad \text{Im } m_{2c}(z) \lesssim \begin{cases} \eta/\sqrt{\kappa + \eta}, & \text{if } E \notin \text{supp } \rho_{2c} \\ \sqrt{\kappa + \eta}, & \text{if } E \in \text{supp } \rho_{2c} \end{cases}, \quad (4.15)$$

$$|m'_{2c}(z)| \lesssim (\kappa + \eta)^{-1/2}, \quad |m_{2c}(z_1) - m_{2c}(z_2)| \lesssim \sqrt{|z_1 - z_2|}, \quad (4.16)$$

and

$$\max_{i \in \mathcal{I}_1} |(1 + m_{2c}(z)\sigma_i)^{-1}| = \text{O}(1). \quad (4.17)$$

for  $z, z_1, z_2 \in \mathbf{D}(\omega, N)$ . The reader can refer to [29, Appendix A] and [16, Lemma 4.5] for the proof.

We define the deterministic limit

$$\Pi(z) := \begin{pmatrix} -(1 + m_{2c}(z)\Lambda)^{-1} & 0 \\ 0 & m_{2c}(z)I_n \end{pmatrix}, \quad (4.18)$$

and the control parameter

$$\Psi(z) := \sqrt{\frac{\text{Im } m_{2c}(z)}{N\eta}} + \frac{1}{N\eta}. \quad (4.19)$$

Note that by (4.15) and (4.17), we have for  $z \in \mathbf{D}$ ,

$$\|\Pi(z)\| = \text{O}(1), \quad \Psi(z) \lesssim (N\eta)^{-1/2}.$$

Our local law of resolvents will be stated under the following bounded support condition. Note that with a standard cutoff argument, the moment assumptions on the  $X$  entries will imply the following bounded support condition with probability  $1 - o(1)$ .

**Definition 4.4** (Bounded support condition). *We say a random matrix  $X$  satisfies the bounded support condition with  $q$ , if*

$$\max_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |X_{i\mu}| \leq q. \quad (4.20)$$

Here  $q \equiv q_N$  is a deterministic parameter and usually satisfies  $N^{-1/2} \leq q \leq N^{-\phi}$  for some small constant  $\phi > 0$ . Whenever (7.6) holds, we say that  $X$  has support  $q$ .

Now we are ready to state the local laws for the resolvent  $G(X, z)$ , which were proved in [29, 49].

**Theorem 4.5** (Local laws). *Suppose  $d_N, X$  and  $\Sigma$  satisfy Assumption 2.5. Suppose  $X$  satisfies (7.6) with  $q \leq N^{-\phi}$  for some constant  $\phi > 0$ . Then the following estimates hold for  $z \in \mathbf{D}$ :*

- **the anisotropic local law:** for any deterministic unit vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{C}^{\mathcal{I}}$ ,

$$|\langle \mathbf{u}, G(X, z)\mathbf{v} \rangle - \langle \mathbf{u}, \Pi(z)\mathbf{v} \rangle| < q + \Psi(z); \quad (4.21)$$

- **the averaged local law:**

$$|m_2(X, z) - m_{2c}(z)| < (N\eta)^{-1}. \quad (4.22)$$

For  $z \in \mathbf{D}_{out}$ , we have the stronger estimates:



- **the anisotropic local law:** for any deterministic unit vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{C}^{\mathcal{I}}$ ,

$$|\langle \mathbf{u}, G(X, z)\mathbf{v} \rangle - \langle \mathbf{u}, \Pi(z)\mathbf{v} \rangle| < q + N^{-1/2}; \quad (4.23)$$

- **the averaged local law:**

$$|m_2(X, z) - m_{2c}(z)| < N^{-1}. \quad (4.24)$$

All of the above estimates are uniform in the spectral parameter  $z$ .

*Proof.* Under the high moment assumption with  $q < N^{-1/2}$ , the estimates (4.21)-(4.23) are proved in Theorem 3.6 [29]. For more general  $q$ , they are proved in Theorems 3.6 and 3.8 of [49]. It remains to show (4.24). As a corollary of (4.22), we have the following rigidity result for the eigenvalues.

For any  $1 \leq k \leq 2L$ , we define

$$N_k := \sum_{2l \leq k} N \int_{a_{2l}}^{a_{2l-1}} \rho_{2c}(x) dx,$$

which is the classical number of eigenvalues in  $[a_{2k}, \lambda_+]$ . Then we define the classical locations  $\gamma_j$  for the eigenvalues of  $\mathcal{Q}_2$  through

$$1 - F_{2c}(\gamma_j) = \frac{j - 1/2}{N}, \quad 1 \leq j \leq n \wedge N. \quad (4.25)$$

Note that (4.25) is well-defined since the  $N_k$ 's are integers by Lemma 2.2. For convenience, we also denote  $\gamma_0 := +\infty$  and  $\gamma_{K+1} := 0$ .

**Lemma 4.6** (Theorem 3.12 of [29]). *Suppose Theorem 4.5 and the regularity condition (2.13) hold. Then for  $\gamma_j \in [a_{2k}, a_{2k-1}]$ , we have*

$$|\lambda_j - \gamma_j| < [(N_{2k} + 1 - j) \wedge (j + 1 - N_{2k-1})]^{-1/3} N^{-2/3}. \quad (4.26)$$

For  $z \in \mathbf{D}_{out}$ , using the definition (4.25), we get

$$\left| \left( \frac{1}{N} \sum_{j=1}^{N \wedge n} \frac{1}{\gamma_j - z} - \frac{N - N \wedge n}{z} \right) - m_{2c}(z) \right| < N^{-1},$$

and using (4.26), we get

$$\left| \left( \frac{1}{N} \sum_{j=1}^{N \wedge n} \frac{1}{\gamma_j - z} - \frac{N - N \wedge n}{z} \right) - m_2(z) \right| = \left| \frac{1}{N} \sum_{j=1}^K \left( \frac{1}{\gamma_j - z} - \frac{1}{\lambda_j - z} \right) \right| < N^{-1}.$$

There two estimates together imply (4.24).  $\square$

Another ingredient of the proof is the following cumulant expansion formula, whose proof is given in [33, Proposition 3.1] and [27, Section II].

**Lemma 4.7.** *Fix any  $l \in \mathbb{N}$  and let  $f \in \mathcal{C}^{l+1}(\mathbb{R})$ . Let  $h$  be a real valued random variable with finite moments up to order  $l + 2$ . Then we have*

$$\mathbb{E}[f(h)h] = \sum_{k=0}^l \frac{1}{k!} \kappa_{k+1}(h) \mathbb{E}f^{(k)}(h) + R_{l+1},$$

where  $\kappa_k(h)$  is the  $k$ -th cumulant of  $h$  and  $R_{l+1}$  satisfies

$$R_{l+1} \lesssim \mathbb{E} |h|^{l+2} \mathbf{1}_{|h| > N^{\varepsilon-1/2}} \cdot \|f^{(l+1)}\|_{\infty} + \mathbb{E} |h|^{l+2} \cdot \sup_{|x| \leq N^{\varepsilon-1/2}} |f^{(l+1)}(x)|$$

for any constant  $\varepsilon > 0$ .

Finally, we introduce the Helffer-Sjöstrand formula [14], which relates the convergence of the process  $Z_{\eta,E}(\mathbf{v}, f)$  to the CLT of the resolvents  $\sqrt{N\eta}(G - \Pi)_{\mathbf{u}\mathbf{u}}$  with  $\mathbf{u} := O\mathbf{v}$ . It has been used previously to obtain the (almost) sharp convergence rates for the ESD (see e.g. [19, 40]) and VESD (see e.g. [45]) of random matrices, and to study the mesoscopic eigenvalue statistics (see e.g. [22, 32]).

**Lemma 4.8** (Helffer-Sjöstrand formula). *Let  $f \in \mathcal{C}^{1,a,b}$  for some fixed  $a, b > 0$ . Let  $\tilde{f}$  be the almost analytic extension of  $f$  defined by*

$$\tilde{f}(x + iy) = f(x) + i(f(x + y) - f(x)).$$

*Let  $\chi \in \mathcal{C}_c^\infty(\mathbb{R})$  be a smooth cutoff function satisfying  $\chi(0) = 1$ . Then for any  $E \in \mathbb{R}$ , we have*

$$f(E) = \frac{1}{\pi} \int_{\mathbb{R}^2} \frac{\partial_{\bar{z}}(\tilde{f}(z)\chi(y))}{E - x - iy} dx dy,$$

where  $\partial_{\bar{z}} := \frac{1}{2}(\partial_x + i\partial_y)$  is the antiholomorphic derivative.

## 5 CLT for resolvents

In this section, we prove Theorem 2.10 and Theorem 2.11 under a stronger moment assumption: for any  $p \in \mathbb{N}$  there is a constant  $C_p$  such that

$$\max_{i,\mu} \mathbb{E} |\sqrt{N} X_{i\mu}|^p \leq C_p. \quad (5.1)$$

Note by Markov's inequality,  $X$  has bounded support  $q < N^{-1/2}$ . In Section 7, we will discuss how to relax it to (2.18) or (2.21) using a Green's function comparison argument.

**Proposition 5.1.** *Theorem 2.10 and Theorem 2.11 hold under the moment assumption (5.1).*

We rewrite (2.23) using the resolvents in (2.7) and  $\Pi$  in (4.18):

$$Y(\mathbf{u}, w) := \mathcal{Y}_{\eta,E}(\mathbf{v}, w) = \sqrt{N\eta} \mathbf{u}^\top (\mathcal{G}_1(E + w\eta) - (E + w\eta)^{-1} \Pi(E + w\eta)) \mathbf{u}, \quad (5.2)$$

where  $\mathbf{u} = O\mathbf{v}$ . Then Proposition 5.1 follows from the following lemma on the convergence of moments.

**Lemma 5.2.** *Suppose  $d_N$ ,  $X$  and  $\Sigma$  satisfy Assumption 2.5,  $N^{-1+c_1} \leq \eta \leq 1$ , and (5.1) holds. Fix any  $E > 0$  and  $k \in \mathbb{N}$ . Then for any deterministic unit vectors  $\mathbf{v}_1, \dots, \mathbf{v}_k \in \mathbb{R}^n$  and  $w_1, \dots, w_k \in \mathbb{H}$ , we have*

$$\mathbb{E}[Y(\mathbf{u}_1, w_1) \cdots Y(\mathbf{u}_k, w_k)] = \begin{cases} \sum \prod \eta \gamma(z_s, z_t, \mathbf{v}_s, \mathbf{v}_t) + O_{<}((N\eta)^{-1/2}), & \text{if } k \in 2\mathbb{N} \\ O_{<}((N\eta)^{-1/2}), & \text{otherwise} \end{cases}. \quad (5.3)$$

Here we denoted  $\mathbf{u}_i := O\mathbf{v}_i$ ,  $z_i := E + w_i\eta$  and  $\gamma(z_s, z_t, \mathbf{v}_s, \mathbf{v}_t) := \hat{\alpha}(z_s, z_t, \mathbf{v}_s, \mathbf{v}_t) + \hat{\beta}(z_s, z_t, \mathbf{v}_s, \mathbf{v}_t)$ , and  $\sum \prod$  means summing over all distinct ways of partitioning of indices into pairs.

In addition, if  $N^{-C} \leq \eta \ll 1$  for some constant  $C > 1$  and  $E \in S_{\text{out}}(\tau)$ , we have the stronger estimates

$$\mathbb{E} \left[ \frac{Y(\mathbf{u}_1, w_1)}{\sqrt{\eta}} \cdots \frac{Y(\mathbf{u}_k, w_k)}{\sqrt{\eta}} \right] = \begin{cases} \sum \prod \gamma(z_s, z_t, \mathbf{v}_s, \mathbf{v}_t) + O_{<}(N^{-1/2}), & \text{if } k \in 2\mathbb{N} \\ O_{<}(N^{-1/2}), & \text{otherwise} \end{cases}. \quad (5.4)$$

*Remark 5.3.* In the statement of this lemma, we allow that  $\mathbf{u}_s = \mathbf{u}_t$  and  $z_s = z_t$  for  $s \neq t$ . In other words, we are calculating the multivariate moments

$$\mathbb{E}[Y^{r_1}(\mathbf{u}_{i_1}, w_{i_1}) \cdots Y^{r_k}(\mathbf{u}_{i_k}, w_{i_k})], \quad r_1, \dots, r_k \in \mathbb{N},$$

if we combine the terms that are the same.

*Proof of Proposition 5.1.* By Wick's theorem, (5.3) with  $E = 0$  and  $\eta = 1$  shows that the convergence in Theorem 2.10 holds in the sense of moments, which further implies the weak convergence. Similarly, under the setting of Theorem 2.11, (5.3) shows that the random vector  $(\mathcal{Y}_{\eta,E}(\mathbf{v}_1, w_1), \dots, \mathcal{Y}_{\eta,E}(\mathbf{v}_k, w_k))$  converges weakly to a complex centered Gaussian vector  $(\Upsilon_1, \dots, \Upsilon_k)$  with covariances

$$\mathbb{E}\Upsilon_i\Upsilon_j = \lim_{N \rightarrow \infty} \left[ \eta \hat{\alpha}^{(N)}(z_i, z_j, \mathbf{v}_i, \mathbf{v}_j) + \eta \hat{\beta}^{(N)}(z_i, z_j, \mathbf{v}_i, \mathbf{v}_j) \right].$$

With  $\eta \ll 1$ , this expression can be simplified to (2.27).

Finally, under the setting of Theorem 2.11, suppose  $E \in S_{out}(\tau)$  and  $N^{-4} \leq \eta \ll 1$ . By Wick's theorem, (5.4) shows that the random vector  $\eta^{-1/2}(\mathcal{Y}_{\eta,E}(\mathbf{v}_1, w_1), \dots, \mathcal{Y}_{\eta,E}(\mathbf{v}_k, w_k))$  converges weakly to a real centered Gaussian vector  $(\Upsilon_1, \dots, \Upsilon_k)$  with covariances

$$\mathbb{E}\Upsilon_i\Upsilon_j = \lim_{N \rightarrow \infty} \left[ \hat{\alpha}^{(N)}(E, E, \mathbf{v}_i, \mathbf{v}_j) + \hat{\beta}^{(N)}(E, E, \mathbf{v}_i, \mathbf{v}_j) \right].$$

Finally, if  $E \in S_{out}(\tau)$  and  $\eta \leq N^{-4}$ , we can show that the random vector  $\eta^{-1/2}(\mathcal{Y}_{\eta,E}(\mathbf{v}_1, w_1), \dots, \mathcal{Y}_{\eta,E}(\mathbf{v}_k, w_k))$  has the same asymptotic distribution as

$$\left( \eta_0^{-1/2} \mathcal{Y}_{\eta_0,E}(\mathbf{v}_1, w_1), \dots, \eta_0^{-1/2} \mathcal{Y}_{\eta_0,E}(\mathbf{v}_k, w_k) \right), \quad \eta_0 := N^{-4},$$

using the bound

$$\|G(E + w_i\eta) - G(E + w_i\eta_0)\| \lesssim |\eta - \eta_0| \|G(E + w_i\eta)\| \cdot \|G(E + w_i\eta_0)\| \lesssim N^{-4} \quad \text{with high probability.}$$

Here we used that by the rigidity estimate (4.26),  $\|G(z)\| = O(1)$  with high probability for  $z \in S_{out}(\tau)$ .  $\square$

In the rest of this section, we mostly focus on the proof of (5.3). We will discuss how to extend the argument to (5.4) at the end of this section. For simplicity, the bulk of the proof is devoted to the calculation of the moments

$$\mathbb{E} \left[ Y^{k_1}(\mathbf{u}_1, w_1) Y^{k_2}(\mathbf{u}_2, \bar{w}_2) \right], \quad k_1, k_2 \in \mathbb{N}, \quad \mathbf{u}_1, \mathbf{u}_2 \in \mathbb{R}^n, \quad w_1, w_2 \in \mathbb{C}_+. \quad (5.5)$$

The proof for the more general expression in (5.3) is exactly the same, except for some immaterial changes of notations.

In the following calculation, we write  $Y(\mathbf{u}_2, \bar{w}_2)$  as  $\bar{Y}(\mathbf{u}_2, w_2)$ , and abbreviate  $z_1 := E + w_1\eta$ ,  $z_2 := E + w_2\eta$ ,  $G^{(1)} := G(z_1)$ ,  $G^{(2)} := G(z_2)$  and  $T := \Lambda^{1/2}O$  (recall (2.3)). Moreover, we denote

$$\begin{aligned} Y_1 &:= z_1 Y(\mathbf{u}_1, w_1) = \sqrt{N\eta} \langle \mathbf{u}_1, (G(z_1) - \Pi(z_1)) \mathbf{u}_1 \rangle, \\ Y_2 &:= z_2 Y(\mathbf{u}_2, w_2) = \sqrt{N\eta} \langle \mathbf{u}_2, (G(z_2) - \Pi(z_2)) \mathbf{u}_2 \rangle, \end{aligned} \quad (5.6)$$

and  $\mathfrak{G} := Y_1^{k_1} \bar{Y}_2^{k_2}$ . In the following proof, we focus on calculating  $\mathbb{E}\mathfrak{G}$ . Note that by the assumptions of Lemma 5.2, we have  $|z_{1,2}| \sim 1$ . Hence we can easily derive the estimates on (5.5) from  $\mathbb{E}\mathfrak{G}$  using the identity

$$Y^{k_1}(\mathbf{u}_1, w_1) \bar{Y}^{k_2}(\mathbf{u}_2, w_2) = z_1^{-k_1} \bar{z}_2^{-k_2} \mathfrak{G}.$$

Without loss of generality, we assume that  $k_1 \geq k_2$  and  $k_1 + k_2 \geq 1$ . Under the assumption (5.1),  $X$  has bounded support  $q < N^{-1/2}$ . Then by (4.21), we have

$$|Y_1| + |Y_2| < \sqrt{N\eta} \Psi(z_1) + \sqrt{N\eta} \Psi(z_2) = O(1).$$

Then using Lemma 4.3 (iii), we get that for any fixed  $n_1, n_2 \in \mathbb{N}$ ,

$$\mathbb{E}|Y_1|^{n_1}|Y_2|^{n_2} < 1,$$

where the second moment bound on  $|Y_1|^{n_1}|Y_2|^{n_2}$  required by Lemma 4.3 (iii) follows immediately from (4.7).

Now using the definition of  $G$  in (4.3) and  $\Pi$  in (4.18), we can write

$$\begin{aligned} G(z) - \Pi(z) &= G(z) [\Pi^{-1}(z) - G^{-1}(z)] \Pi(z) \\ &= G(z) \left[ \begin{pmatrix} -m_{2c}(z)\Lambda & 0 \\ 0 & (m_{2c}^{-1}(z) + z)I_n \end{pmatrix} - \begin{pmatrix} 0 & TX \\ (TX)^\top & 0 \end{pmatrix} \right] \Pi(z). \end{aligned}$$

Using this identity, for  $\mathbf{u}_1 \in \mathbb{R}^{\mathcal{I}_1}$  we can write

$$\begin{aligned} \mathbb{E}\mathfrak{G} &= \mathbb{E}\sqrt{N\eta} \left\langle \mathbf{u}_1, G^{(1)} \begin{pmatrix} -m_{2c}(z_1)\Lambda & 0 \\ 0 & 0 \end{pmatrix} \Pi(z_1) \mathbf{u}_1 \right\rangle Y_1^{k_1-1} \bar{Y}_2^{k_2} \\ &\quad - \mathbb{E}\sqrt{N\eta} \left\langle \mathbf{u}_1, G^{(1)} \begin{pmatrix} 0 & 0 \\ (TX)^\top & 0 \end{pmatrix} \Pi(z_1) \mathbf{u}_1 \right\rangle Y_1^{k_1-1} \bar{Y}_2^{k_2} =: \mathcal{M}_1 + \mathcal{M}_2. \end{aligned} \quad (5.7)$$

Similar as in (2.15), we denote by  $\kappa_k(i, \mu)$  the  $k$ -th cumulants of  $\sqrt{N}X_{i\mu}$ . Then using Lemma 4.7 with  $h = X_{i\mu}$ , we can express  $\mathcal{M}_2$  as

$$\mathcal{M}_2 = -\sqrt{N\eta} \mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} G_{\mathbf{u}_1 \mu}^{(1)} X_{i\mu} \mathbf{w}_1(i) Y_1^{k_1-1} \bar{Y}_2^{k_2} = \sum_{k=1}^l \mathfrak{G}_k + \mathcal{E}, \quad (5.8)$$

where we denoted  $\mathbf{w}_1 := T^\top \Pi(z_1) \mathbf{u}_1$ . The terms on the right-hand side of (5.8) are

$$\mathfrak{G}_k := -\frac{\sqrt{N\eta}}{k! N^{(k+1)/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_{k+1}(i, \mu) \mathbb{E} \frac{\partial^k (G_{\mathbf{u}_1 \mu}^{(1)} Y_1^{k_1-1} \bar{Y}_2^{k_2})}{\partial (X_{i\mu})^k}, \quad (5.9)$$

and

$$\mathcal{E} := -\sqrt{N\eta} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) R_{l+1}(i\mu), \quad (5.10)$$

where  $R_{l+1}(i\mu)$  satisfies the bound

$$R_{l+1}(i\mu) \lesssim \mathbb{E} \left| X_{i\mu}^{l+2} \mathbf{1}_{|X_{i\mu}| > N^{\varepsilon-1/2}} \right| \cdot \|\partial_{i\mu}^{l+1} f_{i\mu}\|_\infty + \mathbb{E} |X_{i\mu}|^{l+2} \cdot \mathbb{E} \sup_{|x| \leq N^{\varepsilon-1/2}} \left| \partial_{i\mu}^{l+1} f_{i\mu}(H^{(i\mu)} + x\Delta_{i\mu}) \right|.$$

Here we abbreviated  $f_{i\mu} := G_{\mathbf{u}_1 \mu}^{(1)} Y_1^{k_1-1} \bar{Y}_2^{k_2}$ ,  $\partial_{i\mu} := \partial/\partial X_{i\mu}$ ,

$$\Delta_{i\mu} := \begin{pmatrix} 0 & \mathbf{t}_i \mathbf{e}_\mu^\top \\ \mathbf{e}_\mu \mathbf{t}_i^\top & 0 \end{pmatrix} G, \quad \mathbf{t}_i := T \mathbf{e}_i,$$

and  $H^{(i\mu)} = H - X_{i\mu} \Delta_{i\mu}$  such that  $H^{(i\mu)}$  is independent of  $X_{i\mu}$ .

Next we estimate the right-hand side of (5.8) term by term using the formula

$$\frac{\partial^r G}{\partial (X_{i\mu})^r} = (-1)^r r! G(\Delta_{i\mu} G)^r. \quad (5.11)$$

This can be derived from the following resolvent expansion: for any  $x, x' \in \mathbb{R}$  and  $k \in \mathbb{N}$ ,

$$G_{(i\mu)}^{x'} = G_{(i\mu)}^x + \sum_{r=1}^k (x - x')^r G_{(i\mu)}^x \left( \Delta_{i\mu} G_{(i\mu)}^x \right)^r + (x - x')^{k+1} G_{(i\mu)}^{x'} \left( \Delta_{i\mu} G_{(i\mu)}^x \right)^{k+1}, \quad (5.12)$$

where we denoted  $G_{(i\mu)}^x := G(H^{(i\mu)} + x\Delta_{i\mu})$ .

## 5.1 The leading term $\mathfrak{G}_1$

We expand  $\mathfrak{G}_1$  as

$$\mathfrak{G}_1 = -\sqrt{\frac{\eta}{N}}\mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \frac{\partial G_{\mathbf{u}_1 \mu}^{(1)}}{\partial X_{i\mu}} \mathbf{w}_1(i) Y_1^{k_1-1} \bar{Y}_2^{k_2} - \sqrt{\frac{\eta}{N}}\mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} G_{\mathbf{u}_1 \mu}^{(1)} \mathbf{w}_1(i) \frac{\partial (Y_1^{k_1-1} \bar{Y}_2^{k_2})}{\partial X_{i\mu}}. \quad (5.13)$$

For the first term in (5.13), we have

$$\begin{aligned} & -\sqrt{\frac{\eta}{N}}\mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \frac{\partial G_{\mathbf{u}_1 \mu}^{(1)}}{\partial X_{i\mu}} \mathbf{w}_1(i) Y_1^{k_1-1} \bar{Y}_2^{k_2} \\ &= \sqrt{\frac{\eta}{N}}\mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mu}^{(1)} \mathbf{w}_1(i) Y_1^{k_1-1} \bar{Y}_2^{k_2} + \sqrt{\frac{\eta}{N}}\mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} G_{\mu \mu}^{(1)} G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} \mathbf{w}_1(i) Y_1^{k_1-1} \bar{Y}_2^{k_2} \\ &= \sqrt{\frac{\eta}{N}}\mathbb{E} (G^{(1)} J_2 G^{(1)})_{\mathbf{u}_1 \tilde{\mathbf{u}}_1} Y_1^{k_1-1} \bar{Y}_2^{k_2} + \sqrt{N\eta}\mathbb{E} \left( m_2(z_1) G_{\mathbf{u}_1 \tilde{\mathbf{u}}_1}^{(1)} Y_1^{k_1-1} \bar{Y}_2^{k_2} \right), \end{aligned} \quad (5.14)$$

where we denoted  $\tilde{\mathbf{u}}_1 := \sum_{i \in \mathcal{I}_1} \mathbf{w}_1(i) \mathbf{t}_i = \tilde{\Lambda} \Pi(z_1) \mathbf{u}_1$ , and

$$J_2 := \begin{pmatrix} 0 & 0 \\ 0 & I_n \end{pmatrix}, \quad \tilde{\Lambda} := \begin{pmatrix} \Lambda & 0 \\ 0 & 0 \end{pmatrix}.$$

For the first term in (5.14), using Ward's identities, Lemma 4.1, we can bound it as

$$O_{<} \left( \sqrt{\frac{\eta}{N}} \mathbb{E} \left[ |G_{\mathbf{u}_1 \mathbf{u}_1}| + \eta^{-1} \left| \operatorname{Im} \left( \bar{z} G_{\mathbf{u}_1 \mathbf{u}_1}^{(1)} \right) \right| \right]^{1/2} \left[ |G_{\tilde{\mathbf{u}}_1 \tilde{\mathbf{u}}_1}| + \eta^{-1} \left| \operatorname{Im} \left( \bar{z} G_{\tilde{\mathbf{u}}_1 \tilde{\mathbf{u}}_1}^{(1)} \right) \right| \right]^{1/2} \right) < (N\eta)^{-1/2}, \quad (5.15)$$

where in the second step we used (4.21) to bound  $|G_{\mathbf{u}_1 \mathbf{u}_1}| < 1$  and  $|G_{\tilde{\mathbf{u}}_1 \tilde{\mathbf{u}}_1}| < 1$ . On the other hand, using (4.22), we can estimate the second term in (5.14) as

$$\sqrt{N\eta}\mathbb{E} \left( m_{2c}(z_1) (G^{(1)} \tilde{\Lambda} \Pi(z_1))_{\mathbf{u}_1 \mathbf{u}_1} Y_1^{k_1-1} \bar{Y}_2^{k_2} \right) + O_{<} \left( (N\eta)^{-1/2} \right) = -\mathcal{M}_1 + O_{<} \left( (N\eta)^{-1/2} \right). \quad (5.16)$$

Next for the second term in (5.13), using (5.11) we calculate that

$$\begin{aligned} & -\sqrt{\frac{\eta}{N}}\mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} G_{\mathbf{u}_1 \mu}^{(1)} \mathbf{w}_1(i) \frac{\partial (Y_1^{k_1-1} \bar{Y}_2^{k_2})}{\partial X_{i\mu}} \\ &= 2(k_1-1)\eta\mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} G_{\mathbf{u}_1 \mu}^{(1)} \mathbf{w}_1(i) G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mathbf{u}_1}^{(1)} Y_1^{k_1-2} \bar{Y}_2^{k_2} \end{aligned} \quad (5.17)$$

$$+ 2k_2\eta\mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} G_{\mathbf{u}_1 \mu}^{(1)} \mathbf{w}_1(i) \bar{G}_{\mathbf{u}_2 \mu}^{(2)} \bar{G}_{\mathbf{t}_i \mathbf{u}_2}^{(2)} Y_1^{k_1-1} \bar{Y}_2^{k_2-1}, \quad (5.18)$$

where as a convention, if  $k_1 = 1$ , then the first term is zero, and similarly if  $k_2 = 0$ , the second term is zero. For the two terms (5.17) and (5.18), we shall apply the identity

$$\sum_{\mu \in \mathcal{I}_2} G_{\mathbf{u} \mu}(z) G_{\mathbf{u}' \mu}(z') = \frac{G_{\mathbf{u} \mathbf{u}'}(z) - G_{\mathbf{u} \mathbf{u}'}(z')}{z - z'}, \quad z, z' \in \mathbb{C}, \quad \mathbf{u}, \mathbf{u}' \in \mathbb{R}^{\mathcal{I}}, \quad (5.19)$$

which follows directly from the definition (4.3). Applying this identity to (5.18), we can write

$$\begin{aligned} \eta \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} G_{\mathbf{u}_1 \mu}^{(1)} \mathbf{w}_1(i) \overline{G_{\mathbf{u}_2 \mu}^{(2)}} \overline{G_{\mathbf{t}_i \mathbf{u}_2}^{(2)}} &= \eta \left( \frac{G_{\mathbf{u}_1 \mathbf{u}_2}(z_1) - G_{\mathbf{u}_1 \mathbf{u}_2}(\bar{z}_2)}{z_1 - \bar{z}_2} \right) \left( \Pi(z_1) \tilde{\Lambda} \overline{G}^{(2)} \right)_{\mathbf{u}_1 \mathbf{u}_2} \\ &= \eta \left( \frac{\Pi_{\mathbf{u}_1 \mathbf{u}_2}(z_1) - \Pi_{\mathbf{u}_1 \mathbf{u}_2}(\bar{z}_2)}{z_1 - \bar{z}_2} \right) \left( \Pi(z_1) \tilde{\Lambda} \overline{\Pi}(z_2) \right)_{\mathbf{u}_1 \mathbf{u}_2} + O_{<} \left( (N\eta)^{-1/2} \right), \end{aligned} \quad (5.20)$$

where in the last step we used (4.21) and that  $|z_1 - \bar{z}_2| \gtrsim \eta$ .

On the other hand, for (5.17) we develop another version of the identity (5.19) in order to deal with the case where  $z$  is very close to  $z'$  (or even  $z = z'$ ). Suppose  $z, z' \in \mathbb{C}_+$  satisfy that  $\text{Im } z \gtrsim \eta$  and  $\text{Im } z' \gtrsim \eta$ . Then we define the contour  $\Gamma = \partial B_{c\eta}(z) \cup \partial B_{c\eta}(z')$  for some constant  $c > 0$ , where  $\partial B_{c\eta}(z)$  denotes the boundary of the disk around  $z$  with radius  $c\eta$ , and similarly for  $\partial B_{c\eta}(z')$ . We can choose  $c > 0$  small enough such that  $\Gamma \subset \mathbb{C}_+$  and  $\min_{\xi \in \Gamma} \text{Im } \xi \gtrsim \eta$ . Then by Cauchy's integral formula and (4.21), we get that

$$\begin{aligned} \sum_{\mu \in \mathcal{I}_2} G_{\mathbf{u} \mu}(z) G_{\mathbf{u}' \mu}(z') &= \frac{1}{2\pi i} \int_{\Gamma} \frac{G_{\mathbf{u} \mathbf{u}'}(\xi)}{(\xi - z)(\xi - z')} d\xi = \frac{1}{2\pi i} \int_{\Gamma} \frac{\Pi_{\mathbf{u} \mathbf{u}'}(\xi) + O_{<}((N\eta)^{-1/2})}{(\xi - z)(\xi - z')} d\xi \\ &= \frac{\Pi_{\mathbf{u} \mathbf{u}'}(z) - \Pi_{\mathbf{u} \mathbf{u}'}(z')}{z - z'} + O_{<} \left( \eta^{-1} (N\eta)^{-1/2} \right). \end{aligned} \quad (5.21)$$

Applying it to (5.17), we can write

$$\eta \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} G_{\mathbf{u}_1 \mu}^{(1)} \mathbf{w}_1(i) G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mathbf{u}_1}^{(1)} = \eta \Pi'_{\mathbf{u}_1 \mathbf{u}_1}(z_1) \left( \Pi(z_1) \tilde{\Lambda} \Pi(z_1) \right)_{\mathbf{u}_1 \mathbf{u}_1} + O_{<} \left( (N\eta)^{-1/2} \right). \quad (5.22)$$

Plugging (5.20) and (5.22) into (5.17) and (5.18), we obtain that

$$\begin{aligned} & - \sqrt{\frac{\eta}{N}} \mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} G_{\mathbf{u}_1 \mu}^{(1)} \mathbf{w}_1(i) \frac{\partial(Y_1^{k_1-1} \overline{Y}_2^{k_2})}{\partial X_{i\mu}} \\ &= 2(k_1 - 1) \eta \Pi'_{\mathbf{u}_1 \mathbf{u}_1}(z_1) \left( \Pi(z_1) \tilde{\Lambda} \Pi(z_1) \right)_{\mathbf{u}_1 \mathbf{u}_1} \mathbb{E} Y_1^{k_1-2} \overline{Y}_2^{k_2} \\ &+ 2k_2 \eta \left( \frac{\Pi_{\mathbf{u}_1 \mathbf{u}_2}(z_1) - \Pi_{\mathbf{u}_1 \mathbf{u}_2}(\bar{z}_2)}{z_1 - \bar{z}_2} \right) \left( \Pi(z_1) \tilde{\Lambda} \overline{\Pi}(z_2) \right)_{\mathbf{u}_1 \mathbf{u}_2} \mathbb{E} Y_1^{k_1-1} \overline{Y}_2^{k_2-1} + O_{<} \left( (N\eta)^{-1/2} \right). \end{aligned} \quad (5.23)$$

In sum, combining (5.14)-(5.16) and (5.23), we obtain that

$$\begin{aligned} \mathcal{M}_1 + \mathfrak{G}_1 &= 2(k_1 - 1) \eta \Pi'_{\mathbf{u}_1 \mathbf{u}_1}(z_1) \left( \Pi(z_1) \tilde{\Lambda} \Pi(z_1) \right)_{\mathbf{u}_1 \mathbf{u}_1} \mathbb{E} Y_1^{k_1-2} \overline{Y}_2^{k_2} \\ &+ 2k_2 \eta \left( \frac{\Pi_{\mathbf{u}_1 \mathbf{u}_2}(z_1) - \Pi_{\mathbf{u}_1 \mathbf{u}_2}(\bar{z}_2)}{z_1 - \bar{z}_2} \right) \left( \Pi(z_1) \tilde{\Lambda} \overline{\Pi}(z_2) \right)_{\mathbf{u}_1 \mathbf{u}_2} \mathbb{E} Y_1^{k_1-1} \overline{Y}_2^{k_2-1} + O_{<} \left( (N\eta)^{-1/2} \right) \\ &= (k_1 - 1) z_1^2 \eta \hat{\beta}(z_1, z_1, \mathbf{v}_1, \mathbf{v}_1) \mathbb{E} Y_1^{k_1-2} \overline{Y}_2^{k_2} \\ &+ k_2 z_1 \bar{z}_2 \eta \hat{\beta}(z_1, \bar{z}_2, \mathbf{v}_1, \mathbf{v}_2) \mathbb{E} Y_1^{k_1-1} \overline{Y}_2^{k_2-1} + O_{<} \left( (N\eta)^{-1/2} \right), \end{aligned} \quad (5.24)$$

where we used (4.18) to rewrite the coefficients into (2.25) and recall that  $\mathbf{v}_i = O^\top \mathbf{u}_i$ .

Note that if  $k_1 = 1$  and  $k_2 = 0$ , we obtain that

$$\mathbb{E} Y_1 < (N\eta)^{-1/2} \quad (5.25)$$

as a byproduct, which verifies the mean zero condition in Proposition 5.1.

## 5.2 The error term $\mathfrak{G}_2$

For the term

$$\mathfrak{G}_2 := -\frac{\sqrt{\eta}}{2N} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_3(i, \mu) \mathbb{E} \frac{\partial^2 (G_{\mathbf{u}_1 \mu}^{(1)} Y_1^{k_1-1} \bar{Y}_2^{k_2})}{\partial (X_{i\mu})^2},$$

we have the following cases. First, we assume that the two derivatives act on  $G_{\mathbf{u}_1 \mu}^{(1)}$ :

$$\frac{\partial^2 G_{\mathbf{u}_1 \mu}^{(1)}}{\partial X_{i\mu}^2} = 4G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mu \mathbf{t}_i}^{(1)} G_{\mu \mu}^{(1)} + 2G_{\mathbf{u}_1 \mu}^{(1)} (G_{\mu \mathbf{t}_i}^{(1)})^2 + 2G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mathbf{t}_i}^{(1)} G_{\mu \mu}^{(1)}.$$

Inserting these three terms into  $\mathfrak{G}_2$ , we can bound the resulting expressions as follows. First we have

$$2\frac{\sqrt{\eta}}{N} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i) G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mu \mathbf{t}_i}^{(1)} G_{\mu \mu}^{(1)}| < \frac{1}{N^{3/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i)| |G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)}| < \frac{1}{\sqrt{N\eta}}, \quad (5.26)$$

where we used (4.21) in the first step to bound  $G_{\mu \mathbf{t}_i}^{(1)} < (N\eta)^{-1/2}$ , and in the second step we used Lemma 4.1 and (4.21) to bound that

$$\sum_{i \in \mathcal{I}_1} |\mathbf{w}(i)| |G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)}| \leq \left( \sum_{i \in \mathcal{I}_1} |\mathbf{w}(i)|^2 \right)^{1/2} \left( \sum_{i \in \mathcal{I}_1} |G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)}|^2 \right)^{1/2} < \eta^{-1/2}.$$

Similarly, we can bound

$$\frac{\sqrt{\eta}}{N} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i) G_{\mathbf{u}_1 \mu}^{(1)} (G_{\mu \mathbf{t}_i}^{(1)})^2| < \frac{1}{N^{5/2}\eta} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i)| < \frac{1}{N\eta}. \quad (5.27)$$

Finally, we have

$$\begin{aligned} & -\frac{\sqrt{\eta}}{N} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_3(i, \mu) G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mathbf{t}_i}^{(1)} G_{\mu \mu}^{(1)} \\ &= -\frac{\sqrt{\eta}}{N} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_3(i, \mu) G_{\mathbf{u}_1 \mu}^{(1)} \Pi_{\mathbf{t}_i \mathbf{t}_i}(z_1) \Pi_{\mu \mu}(z_1) + O_{<} \left( \frac{1}{N^2 \sqrt{\eta}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i)| \right) \\ &< \frac{\sqrt{\eta}}{N} \sum_{i \in \mathcal{I}_1} |\mathbf{w}_1(i)| \eta^{-1/2} + \frac{1}{\sqrt{N\eta}} < \frac{1}{\sqrt{N\eta}}, \end{aligned} \quad (5.28)$$

where in the second step we apply (4.21) to  $G^{(1)}$  to get

$$\sum_{\mu \in \mathcal{I}_2} \kappa_3(i, \mu) G_{\mathbf{u}_1 \mu}^{(1)} \Pi_{\mu \mu}(z_1) = G_{\mathbf{u}_1 \tilde{\mathbf{w}}_i}^{(1)} < \frac{\sqrt{N}}{\sqrt{N\eta}} = \eta^{-1/2}. \quad (5.29)$$

Here we used that  $\tilde{\mathbf{w}}_i := \sum_{\mu} \kappa_3(i, \mu) \Pi_{\mu \mu}(z_1) \mathbf{e}_{\mu}$  has  $l^2$ -norm  $O(\sqrt{N})$ .

Then we consider the case that one derivative acts on  $G_{\mathbf{u}_1 \mu}^{(1)}$  and the other acts on  $Y_1^{k_1-1} \bar{Y}_2^{k_2}$ . Suppose the other derivative acts on a  $Y_1$  factor, then we need to estimate

$$-\frac{\eta}{\sqrt{N}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_3(i, \mu) \left( G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mu}^{(1)} + G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mu \mu}^{(1)} \right) G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mathbf{u}_1 \mu}^{(1)}.$$

For the first term, we can bound it using (4.21) as

$$\frac{\eta}{\sqrt{N}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| |G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mu}^{(1)} G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mathbf{u}_1 \mu}^{(1)}| < \frac{1}{N^2 \sqrt{\eta}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| < \frac{1}{\sqrt{N\eta}}. \quad (5.30)$$

For the second term, we can apply similar argument as in (5.28) to get that

$$\begin{aligned} & - \frac{\eta}{\sqrt{N}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_3(i, \mu) G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mu \mu}^{(1)} G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mathbf{u}_1 \mu}^{(1)} \\ &= - \frac{\eta}{\sqrt{N}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_3(i, \mu) \Pi_{\mu \mu}(z_1) (G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)})^2 G_{\mathbf{u}_1 \mu}^{(1)} + \text{O} \left( \frac{1}{N^{3/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| |G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)}| \right) \\ &< \frac{1}{\sqrt{N}} \sum_{i \in \mathcal{I}_1} |\mathbf{w}_1(i)| |G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)}| < (N\eta)^{-1/2}, \end{aligned} \quad (5.31)$$

where in the third step we applied (5.29) to the first term, and in the last step we used Lemma 4.1 and (4.21) to bound

$$\sum_{i \in \mathcal{I}_1} |\mathbf{w}_1(i)| |G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)}| < \left( \sum_{i \in \mathcal{I}_1} |\mathbf{w}_1(i)|^2 \right)^{1/2} \left( \sum_{i \in \mathcal{I}_1} |G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)}|^2 \right)^{1/2} < \eta^{-1/2}. \quad (5.32)$$

If the other derivative acts on a  $\bar{Y}_2$  factor, then we have similar estimates.

Finally, we consider the case that there are two derivatives acting on  $Y^{k_1-1} \bar{Y}^{k_2}$ . There are two cases.

**Case 1:** Suppose that the two derivatives act on two different  $Y$  factors. If these are two  $Y_1$  factors, then we need to bound

$$\eta^{3/2} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| |G_{\mathbf{u}_1 \mu}^{(1)}| |G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)}|^2 |G_{\mathbf{u}_1 \mu}^{(1)}|^2 < \frac{N\eta^{3/2}}{(N\eta)^{3/2}} \sum_{i \in \mathcal{I}_1} |\mathbf{w}_1(i)| |G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)}| < \frac{1}{\sqrt{N\eta}}, \quad (5.33)$$

where we used (5.32) in the second step. We have similar estimates if the two derivatives acts on two  $\bar{Y}_2$  factors, or act on a  $Y_1$  factor and a  $\bar{Y}_2$  factor.

**Case 2:** Suppose two derivatives act on one  $Y$  factor. If this is a  $Y_1$  factor, then we need to bound

$$- \frac{\eta}{\sqrt{N}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_3(i, \mu) G_{\mathbf{u}_1 \mu}^{(1)} \left( (G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)})^2 G_{\mu \mu}^{(1)} + (G_{\mathbf{u}_1 \mu}^{(1)})^2 G_{\mathbf{t}_i \mathbf{t}_i}^{(1)} + 2G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mu}^{(1)} \right).$$

The first term has been estimated in (5.31), and the third term has been estimated in (5.30). For the second term, using (4.21) we get

$$\frac{\eta}{\sqrt{N}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| |G_{\mathbf{u}_1 \mu}^{(1)}|^3 |G_{\mathbf{t}_i \mathbf{t}_i}^{(1)}| < \frac{1}{N^2 \sqrt{\eta}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| < \frac{1}{\sqrt{N\eta}}. \quad (5.34)$$

If the two derivatives act on a  $\bar{Y}_2$  factor, then we have similar estimates.

Combining (5.26)-(5.28), (5.30), (5.31), (5.33) and (5.34), we obtain that

$$\mathfrak{G}_2 < \frac{1}{\sqrt{N\eta}}. \quad (5.35)$$



### 5.3 Terms $\mathfrak{G}_k$ with $k \geq 3$

For the terms  $\mathfrak{G}_k$  with  $k \geq 3$ , the expressions for the derivatives begin to become rather complicated. In order to exploit the structures of these term in a systematical way, we introduce the following algebraic objects.

**Definition 5.4** (Words). *Given  $i \in \mathcal{I}_1$  and  $\mu \in \mathcal{I}_2$ , let  $\mathcal{W}$  be the set of words of even length in two letters  $\{\mathbf{i}, \boldsymbol{\mu}\}$ . We denote the length of a word  $w \in \mathcal{W}$  by  $2\mathbf{l}(w)$  with  $\mathbf{l}(w) \in \mathbb{N}$ . We use bold symbols to denote the letters of words. For instance,  $w = \mathbf{a}_1 \mathbf{b}_2 \mathbf{a}_2 \mathbf{b}_3 \cdots \mathbf{a}_r \mathbf{b}_{r+1}$  denotes a word of length  $2r$ . Let  $\mathcal{W}_r := \{w \in \mathcal{W} : \mathbf{l}(w) = r\}$  be the set of words of length  $2r$ , and such that each word  $w \in \mathcal{W}_r$  satisfies that  $\mathbf{a}_l \mathbf{b}_{l+1} \in \{\mathbf{i}\boldsymbol{\mu}, \boldsymbol{\mu}\mathbf{i}\}$  for all  $1 \leq l \leq r$ .*

*Next we assign to each letter a value  $[\cdot]$  through  $[\mathbf{i}] := \mathbf{t}_i$  and  $[\boldsymbol{\mu}] := \mathbf{e}_\mu$ . It is important to distinguish the abstract letter from its value, which is a vector (or can be regarded as a summation index). To each word  $w$  we assign two types of random variables  $A_{i,\mu}^{(1)}(w)$  and  $A_{i,\mu}^{(2)}(w)$  as follows. If  $\mathbf{l}(w) = 0$  we define*

$$A_{i,\mu}^{(1)}(w) := G_{\mathbf{u}_1 \mathbf{u}_1}^{(1)} - \Pi_{\mathbf{u}_1 \mathbf{u}_1}(z_1), \quad A_{i,\mu}^{(2)}(w) := G_{\mathbf{u}_2 \mathbf{u}_2}^{(2)} - \Pi_{\mathbf{u}_2 \mathbf{u}_2}(z_2).$$

*If  $\mathbf{l}(w) \geq 1$ , say  $w = \mathbf{a}_1 \mathbf{b}_2 \mathbf{a}_2 \mathbf{b}_3 \cdots \mathbf{a}_r \mathbf{b}_{r+1}$ , we define*

$$\begin{aligned} A_{i,\mu}^{(1)}(w) &:= G_{\mathbf{u}_1[\mathbf{a}_1]}^{(1)} G_{[\mathbf{b}_2][\mathbf{a}_2]}^{(1)} \cdots G_{[\mathbf{b}_r][\mathbf{a}_r]}^{(1)} G_{[\mathbf{b}_{r+1}]\mathbf{u}_1}^{(1)}, \\ A_{i,\mu}^{(2)}(w) &:= \overline{G}_{\mathbf{u}_2[\mathbf{a}_1]}^{(2)} \overline{G}_{[\mathbf{b}_2][\mathbf{a}_2]}^{(2)} \cdots \overline{G}_{[\mathbf{b}_r][\mathbf{a}_r]}^{(2)} \overline{G}_{[\mathbf{b}_{r+1}]\mathbf{u}_2}^{(2)}. \end{aligned} \quad (5.36)$$

*Finally, for  $w = \mathbf{a}_1 \mathbf{b}_2 \mathbf{a}_2 \mathbf{b}_3 \cdots \mathbf{a}_r \mathbf{b}_{r+1}$  we define another type of word as*

$$\tilde{A}_{i,\mu}(w) := G_{\mathbf{u}_1[\mathbf{a}_1]}^{(1)} G_{[\mathbf{b}_2][\mathbf{a}_2]}^{(1)} \cdots G_{[\mathbf{b}_r][\mathbf{a}_r]}^{(1)} G_{[\mathbf{b}_{r+1}]\mu}^{(1)}. \quad (5.37)$$

Notice the words are constructed such that, by (5.11),

$$\left( \frac{\partial}{\partial X_{i\mu}} \right)^r Y_1 = (-1)^r r! \sqrt{N\eta} \sum_{w \in \mathcal{W}_r} A_{i,\mu}^{(1)}(w), \quad r \in \mathbb{N}.$$

Similarly,  $A_{i,\mu}^{(2)}(w)$  is related to the derivatives of  $\overline{Y}_2$ , and  $\tilde{A}_{i,\mu}(w)$  is related to the derivatives of  $G_{\mathbf{u}_1 \mu}$ . Thus we can write

$$\begin{aligned} \frac{\partial^k (G_{\mathbf{u}_1 \mu}^{(1)} Y_1^{k_1-1} \overline{Y}_2^{k_2})}{\partial (X_{i\mu})^k} &= (-1)^k (N\eta)^{\frac{1}{2}(k_1+k_2-1)} \sum_{l_1+\cdots+l_{k_1+k_2}=k} \left[ l_1! \sum_{w_1 \in \mathcal{W}_{l_1}} \tilde{A}_{i,\mu}(w_1) \right] \\ &\times \prod_{s=2}^{k_1} \left[ l_s! \sum_{w_s \in \mathcal{W}_{l_s}} A_{i,\mu}^{(1)}(w_s) \right] \prod_{s=k_1+1}^{k_1+k_2} \left[ l_s! \sum_{w_s \in \mathcal{W}_{l_s}} A_{i,\mu}^{(2)}(w_s) \right]. \end{aligned} \quad (5.38)$$

In the following proof, for simplicity we shall abbreviate

$$A_{i,\mu}(w_s) \equiv \begin{cases} A_{i,\mu}^{(1)}(w_s), & \text{if } 2 \leq s \leq k_1 \\ A_{i,\mu}^{(2)}(w_s), & \text{if } k_1+1 \leq s \leq k_1+k_2 \end{cases}.$$

Moreover, we introduce the notations

$$a := \#\{2 \leq s \leq k_1+k_2 : l_i \geq 1\}, \quad a_1 := \#\{2 \leq s \leq k_1+k_2 : l_i = 1\}.$$

Without loss of generality, we assume that the words with nonzero length are  $w_{s_1}, \dots, w_{s_a}$ , and the words with length 1 are  $w_{s_1}, \dots, w_{s_{a_1}}$ . Then we have

$$l_{s_1} + \dots + l_{s_a} = k - l_1 \Rightarrow 2a \leq k - l_1 + a_1. \quad (5.39)$$

By the definition, it is easy to verify that

$$|A_{i,\mu}(w_s)| < R_i^2 + R_\mu^2, \quad \text{if } l_s \geq 1, s \geq 2 \quad (5.40)$$

where we abbreviated

$$R_i := |G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)}| + |G_{\mathbf{u}_2 \mathbf{t}_i}^{(2)}|, \quad R_\mu := |G_{\mathbf{u}_1 \mu}^{(1)}| + |G_{\mathbf{u}_2 \mu}^{(2)}| + |G_{\mathbf{t}_i \mu}^{(1)}| + |G_{\mathbf{t}_i \mu}^{(2)}| < (N\eta)^{-1/2}.$$

If  $l_s = 1$  for some  $s \geq 2$ , we have the better bound

$$|A_{i,\mu}(w_s)| < R_i R_\mu < \frac{R_i}{\sqrt{N\eta}}. \quad (5.41)$$

Similarly, we have

$$|\tilde{A}_{i,\mu}(w_1)| < \mathbf{1}(l_1 \geq 1)R_i + R_\mu < \mathbf{1}(l_1 \geq 1)R_i + (N\eta)^{-1/2}. \quad (5.42)$$

Finally, using Lemma 4.1 and (4.21), we can bound

$$\sum_{i \in \mathcal{I}_1} R_i^2 + \sum_{\mu \in \mathcal{I}_2} R_\mu^2 < \eta^{-1}, \quad \sum_{i \in \mathcal{I}_1} |\mathbf{w}_1(i)| R_i < \eta^{-1/2}. \quad (5.43)$$

We will use these bounds tacitly in the following proof.

Now we study the  $k = 3$  case using the above tools. In this case, we will obtain a leading term that depends on the fourth cumulants of the  $X$  entries.

**The leading term  $\mathfrak{G}_3$ .** We insert (5.38) into the term

$$\mathfrak{G}_3 := -\frac{\sqrt{\eta}}{6N^{3/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_4(i, \mu) \mathbb{E} \frac{\partial^3 (G_{\mathbf{u}_1 \mu}^{(1)} Y_1^{k_1-1} \bar{Y}_2^{k_2})}{\partial (X_{i\mu})^3}.$$

Applying (5.40)-(5.42) to (5.38), we need to bound

$$\begin{aligned} & \frac{(N\eta)^{a/2} \sqrt{\eta}}{N^{3/2}} \mathbf{1}(l_1 \geq 1) \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| R_i (R_i R_\mu)^{a_1} (R_i^2 + R_\mu^2)^{a-a_1} \\ & + \frac{(N\eta)^{a/2} \sqrt{\eta}}{N^{3/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| R_\mu (R_i R_\mu)^{a_1} (R_i^2 + R_\mu^2)^{a-a_1} =: \mathcal{K}_1 + \mathcal{K}_2. \end{aligned}$$

Now we bound  $\mathcal{K}_2$ . First we consider the case with  $a_1 = 0$ . Then  $a$  can only be 0 or 1, and we have

$$\begin{aligned} \mathcal{K}_2 & < \frac{\mathbf{1}(a=0)}{N^2} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| + \mathbf{1}(a=1) \frac{(N\eta)^{1/2}}{N^2} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| (R_i^2 + R_\mu^2) \\ & < N^{-1/2} + \frac{(N\eta)^{1/2}}{N^2} \left( \frac{N}{\sqrt{\eta}} + \frac{\sqrt{N}}{\eta} \right) < N^{-1/2}. \end{aligned}$$

Then in the  $a_1 = 1$  case,  $a$  can only be 1 or 2, and we have

$$\begin{aligned} \mathcal{K}_2 &< \mathbf{1}(a=1) \frac{(N\eta)^{1/2}}{N^2} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i)| R_i R_\mu + \mathbf{1}(a=2) \frac{N\eta}{N^2} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i)| (R_i R_\mu) (R_i^2 + R_\mu^2) \\ &< \frac{(N\eta)^{1/2}}{N^2} \frac{\sqrt{N}}{\eta} + \frac{N\eta}{N^2} \left( \frac{\sqrt{N}}{\eta} + \frac{1}{\sqrt{N}\eta^2} \right) < N^{-1/2}. \end{aligned}$$

Finally, for the  $a_1 \geq 2$  case, we have  $a = a_1$  and

$$\mathcal{K}_2 < \frac{(N\eta)^{a_1/2}}{N^2} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| (R_i R_\mu)^{a_1} < \frac{1}{N^2} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| R_i < \frac{1}{N\sqrt{\eta}} \leq N^{-1/2}.$$

Next we estimate  $\mathcal{K}_1$ . If  $a_1 = 0$  and  $l_1 \geq 2$ , then  $a$  can only be 0, and we have that

$$\mathcal{K}_1 < \frac{\sqrt{\eta}}{N^{3/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| R_i < N^{-1/2}.$$

If  $a_1 = 1$  and  $l_1 \geq 1$ , then  $a$  can only be 1, and we have that

$$\mathcal{K}_1 < \frac{(N\eta)^{1/2} \sqrt{\eta}}{N^{3/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| R_i (R_i R_\mu) < \frac{(N\eta)^{1/2} \sqrt{\eta} \sqrt{N}}{N^{3/2} \eta} = N^{-1/2}.$$

If  $a_1 \geq 2$  and  $l_1 \geq 1$ , then  $a = a_1 = 2$  and we have

$$\mathcal{K}_1 < \frac{(N\eta) \sqrt{\eta}}{N^{3/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| R_i (R_i R_\mu)^2 < \frac{(N\eta) \sqrt{\eta}}{N^{3/2}} \frac{1}{\eta^{3/2}} = N^{-1/2}.$$

Finally, we are left with the case  $a_1 = 0$  and  $l_1 = 1$ , which will provide a leading term. In this case we have that one derivative acts on  $G_{\mathbf{u}_1 \mu}^{(1)}$  and two other derivatives act on a  $Y_1$  or  $\bar{Y}_2$  factor, i.e.

$$\begin{aligned} \mathfrak{G}_3 &= (k_1 - 1) \frac{\sqrt{\eta}}{6N^{3/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_4(i, \mu) \mathbb{E} \left( G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mu \mu}^{(1)} + G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mu}^{(1)} \right) \frac{\partial^2 Y_1}{\partial (X_{i\mu})^2} Y_1^{k_1-2} \bar{Y}_2^{k_2} \\ &+ k_2 \frac{\sqrt{\eta}}{6N^{3/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_4(i, \mu) \mathbb{E} \left( G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mu \mu}^{(1)} + G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mu}^{(1)} \right) \frac{\partial^2 \bar{Y}_2}{\partial (X_{i\mu})^2} Y_1^{k_1-1} \bar{Y}_2^{k_2-1} + \mathcal{O}_<(N^{-1/2}) \\ &= (k_1 - 1) \frac{\sqrt{\eta}}{6N^{3/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_4(i, \mu) \mathbb{E} G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mu \mu}^{(1)} \frac{\partial^2 Y_1}{\partial (X_{i\mu})^2} Y_1^{k_1-2} \bar{Y}_2^{k_2} \\ &+ k_2 \frac{\sqrt{\eta}}{6N^{3/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_4(i, \mu) \mathbb{E} G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mu \mu}^{(1)} \frac{\partial^2 \bar{Y}_2}{\partial (X_{i\mu})^2} Y_1^{k_1-1} \bar{Y}_2^{k_2-1} + \mathcal{O}_<(N^{-1/2}), \end{aligned} \quad (5.44)$$

where in the second step we used that the  $G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mu}^{(1)}$  terms have bounded as  $\mathcal{K}_2$  in the above proof. We now calculate the first term on the right-hand side of (5.44), which takes the form

$$\begin{aligned} (k_1 - 1) \frac{\eta}{3N} \mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_4(i, \mu) G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mu \mu}^{(1)} \left( (G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)})^2 G_{\mu \mu}^{(1)} + (G_{\mathbf{u}_1 \mu}^{(1)})^2 G_{\mathbf{t}_i \mathbf{t}_i}^{(1)} + 2G_{\mathbf{u}_1 \mathbf{t}_i}^{(1)} G_{\mathbf{u}_1 \mu}^{(1)} G_{\mathbf{t}_i \mu}^{(1)} \right) Y_1^{k_1-2} \bar{Y}_2^{k_2} \\ =: \mathbb{E} \mathcal{K}_1 + \mathbb{E} \mathcal{K}_2 + \mathbb{E} \mathcal{K}_3, \end{aligned}$$

where we abused the notations  $\mathcal{K}_{1,2}$  a little bit. We can bound

$$\mathcal{K}_2 + \mathcal{K}_3 < \frac{\eta}{N} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| R_i R_\mu^2 < \frac{\eta}{N} \frac{1}{\eta^{3/2}} \leq N^{-1/2}.$$

For  $\mathcal{K}_1$ , we have

$$\begin{aligned} \mathbb{E}\mathcal{K}_1 &= (k_1 - 1) \frac{\eta}{3N} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \mathbf{w}_1(i) \kappa_4(i, \mu) [\Pi_{\mu\mu}(z_1)]^2 [\Pi_{\mathbf{u}_1 \mathbf{t}_i}(z_1)]^3 \mathbb{E}Y_1^{k_1-2} \bar{Y}_2^{k_2} \\ &+ O_{<} \left( \frac{\sqrt{\eta}}{N^{3/2}} \mathbb{E} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| (R_i + |\Pi_{\mathbf{u}_1 \mathbf{t}_i}(z_1)|) \right) \\ &= (k_1 - 1) \frac{\eta m_{2c}^2(z_1)}{3N} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \kappa_4(i, \mu) [\Pi_{\mathbf{u}_1 \mathbf{t}_i}(z_1)]^4 \mathbb{E}Y_1^{k_1-2} \bar{Y}_2^{k_2} + O_{<} \left( N^{-1/2} \right). \end{aligned}$$

where we used that  $\mathbf{w}_1(i) = \Pi_{\mathbf{u}_1 \mathbf{t}_i}(z_1)$  by the definition of  $\mathbf{w}_1$ . We have a similar estimate for the second term on the right-hand side of (5.44). In sum, we obtain that

$$\begin{aligned} \mathfrak{G}_3 &= (k_1 - 1) \frac{\eta m_{2c}^2(z_1)}{3N} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \kappa_4(i, \mu) [\Pi_{\mathbf{u}_1 \mathbf{t}_i}(z_1)]^4 \mathbb{E}Y_1^{k_1-2} \bar{Y}_2^{k_2} \\ &+ k_2 \frac{\eta m_{2c}(z_1) \bar{m}_{2c}(z_2)}{3N} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \kappa_4(i, \mu) [\Pi_{\mathbf{u}_1 \mathbf{t}_i}(z_1)]^2 [\bar{\Pi}_{\mathbf{u}_2 \mathbf{t}_i}(z_2)]^2 \mathbb{E}Y_1^{k_1-1} \bar{Y}_2^{k_2-1} + O_{<} \left( N^{-1/2} \right) \\ &= (k_1 - 1) z_1^2 \eta \hat{\alpha}(z_1, z_1, \mathbf{u}_1, \mathbf{u}_1) \mathbb{E}Y_1^{k_1-2} \bar{Y}_2^{k_2} + k_2 z_1 \bar{z}_2 \eta \hat{\alpha}(z_1, \bar{z}_2, \mathbf{u}_1, \mathbf{u}_2) \mathbb{E}Y_1^{k_1-1} \bar{Y}_2^{k_2-1} + O_{<} \left( N^{-1/2} \right), \quad (5.45) \end{aligned}$$

where we used (4.18) to rewrite the coefficients into (2.24).

Next we deal with the  $k \geq 4$  cases, which only contain error terms.

**The error terms  $\mathfrak{G}_k$ ,  $k \geq 4$ .** The terms  $\mathfrak{G}_k$ ,  $k \geq 4$ , can be estimated in similar ways as  $\mathfrak{G}_3$ . We insert (5.38) into (5.9), and apply (5.40)-(5.42) to get that

$$\begin{aligned} \mathfrak{G}_k &< \frac{(N\eta)^{a/2} \sqrt{\eta}}{N^{k/2}} \mathbf{1}(l_1 \geq 1) \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| R_i (R_i R_\mu)^{a_1} (R_i^2 + R_\mu^2)^{a-a_1} \\ &+ \frac{(N\eta)^{a/2} \sqrt{\eta}}{N^{k/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| R_\mu (R_i R_\mu)^{a_1} (R_i^2 + R_\mu^2)^{a-a_1} =: \mathcal{K}_1 + \mathcal{K}_2. \end{aligned}$$

For the term  $\mathcal{K}_1$ , we have

$$\mathcal{K}_1 < \mathbf{1}(l_1 \geq 1) \frac{(N\eta)^{(a-a_1)/2} \sqrt{\eta}}{N^{k/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i)| R_i < \mathbf{1}(l_1 \geq 1) \frac{(N\eta)^{(a-a_1)/2} \sqrt{\eta}}{N^{k/2}} \frac{N}{\sqrt{\eta}} \leq N^{-(k-a+a_1)/2+1},$$

where in the second step we used (5.43). With (5.39), we obtain that

$$\frac{k-a+a_1}{2} - 1 \geq \frac{1}{2} \left( k + a_1 - \frac{k-l_1+a_1}{2} \right) - 1 \geq \frac{1}{2} \Rightarrow \mathcal{K}_1 < N^{-1/2}, \quad (5.46)$$

if  $k + a_1 + l_1 \geq 6$ . It remain to consider the case  $k = 4$ ,  $l_1 = 1$  and  $a_1 = 0$ . In this case,  $a$  can only be 1 and we still have

$$\frac{k-a+a_1}{2} - 1 = \frac{1}{2} \Rightarrow \mathcal{K}_1 < N^{-1/2}.$$

Then we bound  $\mathcal{K}_2$ . If  $a_1 = 0$ , we have

$$\begin{aligned}\mathcal{K}_2 &< \frac{(N\eta)^{a/2}}{N^{(k+1)/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i)|(R_i^2 + R_\mu^2)^a \\ &< \frac{\mathbf{1}(a=0)}{N^{(k+1)/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i)| + \mathbf{1}(a \geq 1) \frac{(N\eta)^{a/2}}{N^{(k+1)/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i)|(R_i^2 + R_\mu^2) \\ &< \frac{\mathbf{1}(a=0)}{N^{k/2-1}} + \mathbf{1}(a \geq 1) \frac{(N\eta)^{a/2}}{N^{(k+1)/2}} \left( \frac{N}{\sqrt{\eta}} + \frac{\sqrt{N}}{\eta} \right) < \frac{1}{N} + \frac{\mathbf{1}(a \geq 1)}{N^{(k-a-1)/2}} < N^{-1/2}.\end{aligned}$$

where we used (5.43) in the third step,  $k \geq 4$  in the fourth step, and a similar estimate as in (5.46) in the last step:

$$\frac{k-a-1}{2} \geq \frac{k+l_1}{4} - \frac{1}{2} \geq \frac{1}{2}.$$

If  $a_1 \geq 1$ , we have

$$\mathcal{K}_2 < \frac{(N\eta)^{a/2}}{N^{(k+1)/2}} \frac{1}{(N\eta)^{a_1/2}} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}(i)| R_i < \frac{(N\eta)^{(a-a_1)/2}}{N^{(k+1)/2}} \frac{N}{\sqrt{\eta}} < N^{-1/2},$$

where in the last step we used (5.39) to get that

$$\frac{(N\eta)^{(a-a_1)/2}}{N^{(k+1)/2}} \frac{N}{\sqrt{\eta}} < \begin{cases} N^{-(k-1)/2} \eta^{-1/2} \leq N^{-(k-2)/2} \leq N^{-1}, & \text{if } a = a_1 \\ N^{-(k+a_1-a-1)/2} \leq N^{-(k+a_1-2)/4} \leq N^{-1/2}, & \text{if } a > a_1 \end{cases}.$$

In sum, we obtain that

$$\mathfrak{G}_k < N^{-1/2}, \quad k \geq 4. \quad (5.47)$$

## 5.4 The error term $\mathcal{E}$

Finally we show that the term  $\mathcal{E}$  in (5.10) is sufficiently small for large enough  $l$ . We first bound

$$\mathcal{K}_1 := \sqrt{N\eta} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| \mathbb{E} |X_{i\mu}|^{l+2} \cdot \mathbb{E} \sup_{|x| \leq N^{\varepsilon-1/2}} \left| \partial_{i\mu}^{l+1} f_{i\mu}(H^{(i\mu)} + x\Delta_{i\mu}) \right|.$$

We claim that for any deterministic unit vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^{\mathcal{I}}$ ,

$$\sup_{|x| \leq N^{\varepsilon-1/2}} \left( |G_{\mathbf{u}\mathbf{v}}^{(1)}(H^{(i\mu)} + x\Delta_{i\mu})| + |G_{\mathbf{u}\mathbf{v}}^{(2)}(H^{(i\mu)} + x\Delta_{i\mu})| \right) = O(1) \quad \text{with high probability.} \quad (5.48)$$

In fact, for any  $z \in \{z_1, z_2\}$  and  $|x| \leq N^{\varepsilon-1/2}$  we have the following resolvent expansion by (5.12) with  $x' = X_{i\mu}$ :

$$G(H^{(i\mu)} + x\Delta_{i\mu}) = G(z) - (x - X_{i\mu})G(z)\Delta_{i\mu}G(z) + (x - X_{i\mu})^2 G(H^{(i\mu)} + x\Delta_{i\mu}) (\Delta_{i\mu}G(z))^2.$$

Using  $|X_{i\mu}| < N^{-1/2}$ ,  $|x| \leq N^{\varepsilon-1/2}$ , (4.21) for  $G(z)$ , and the rough bound (4.7) for  $G(H^{(i\mu)} + x\Delta_{i\mu})$ , we obtain from the above expansion that

$$G_{\mathbf{u}\mathbf{v}}(H^{(i\mu)} + x\Delta_{i\mu}) = O_{<}(1) + \eta^{-1} N^{-(1-2\varepsilon)} = O_{<}(1),$$

as long as  $\varepsilon$  is small enough such that  $2\varepsilon < c_1$  (recall that  $\eta \geq N^{-1+c_1}$ ). This implies (5.48). With (5.48) and (5.1), we can bound  $|\partial_{i\mu}^{l+1} f_{i\mu}(H^{(i\mu)} + x\Delta_{i\mu})| < (N\eta)^{(k_1+k_2-1)/2}$  and

$$\mathcal{K}_1 < (N\eta)^{(k_1+k_2)/2} N^{3/2} N^{-(l+2)/2} \leq N^{-1/2}$$

as long as  $l \geq k_1 + k_2 + 2$ .

Now fix an  $l \geq k_1 + k_2 + 2$ , we bound the term

$$\mathcal{K}_2 := \sqrt{N\eta} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} |\mathbf{w}_1(i)| \mathbb{E} \left| X_{i\mu}^{l+2} \mathbf{1}_{|X_{i\mu}| > N^{\varepsilon-1/2}} \right| \cdot \|\partial_{i\mu}^{l+1} f_{i\mu}\|_\infty.$$

Recall that the derivatives take the form (5.38). Then using (4.7), we can bound

$$\|\partial_{i\mu}^{l+1} f_{i\mu}\|_\infty \lesssim (N\eta)^{(k_1+k_2-1)/2} \eta^{-(k_1+k_2+l+1)}.$$

On the other hand, by (5.1) we have

$$\mathbb{E} \left| X_{i\mu}^{l+2} \mathbf{1}_{|X_{i\mu}| > N^{\varepsilon-1/2}} \right| \leq N^{-D}$$

for any fixed constant  $D > 0$ . Hence we have

$$\mathcal{K}_2 < (N\eta)^{(k_1+k_2)/2} \eta^{-(k_1+k_2+l+1)} N^{3/2} N^{-D} < N^{-1/2}$$

as long as  $D$  is taken large enough.

In sum, we obtain that

$$\mathcal{E} < N^{-1/2}. \quad (5.49)$$

Combining the estimates (5.24), (5.35), (5.45), (5.47) and (5.49), we conclude that

$$\mathbb{E} Y_1^{k_1} \bar{Y}_2^{k_2} = (k_1 - 1) z_1^2 \eta \gamma(z_1, z_1, \mathbf{v}_1, \mathbf{v}_1) \mathbb{E} Y_1^{k_1-2} \bar{Y}_2^{k_2} + k_2 z_1 \bar{z}_2 \eta \gamma(z_1, \bar{z}_2, \mathbf{v}_1, \mathbf{v}_2) \mathbb{E} Y_1^{k_1-1} \bar{Y}_2^{k_2-1} + O_{<} \left( (N\eta)^{-1/2} \right).$$

Then applying this induction relations repeatedly and using (5.25), we can conclude (5.3) for the expression in (5.5). Now we can extend the proof to the general expression on the left-hand side of (5.3).

*Proof of Lemma 5.2.* We calculate  $\mathbb{E}[Y(\mathbf{u}_1, w_1) \cdots Y(\mathbf{u}_k, w_k)]$  using cumulant expansions as in (5.7) and (5.8). All the leading terms and error terms can be estimated in exactly the same way. For example, if we expand  $Y(\mathbf{u}_1, w_1)$  as in (5.7), we can obtain that

$$\mathbb{E}[Y(\mathbf{u}_1, w_1) \cdots Y(\mathbf{u}_k, w_k)] = \sum_{s=2}^k \eta \gamma(z_1, z_s, \mathbf{u}_1, \mathbf{u}_s) \mathbb{E} \prod_{t \notin \{1, s\}} Y(\mathbf{u}_t, w_t) + O_{<} \left( (N\eta)^{-1/2} \right). \quad (5.50)$$

Using this induction relation and (5.25), we can then conclude (5.3).

For (5.4), the proof is similar and we only explain the differences. First, the local laws (4.21) and (4.22) can be replaced with the stronger ones (4.23) and (4.24). Moreover, by the rigidity estimate (4.26), we have  $\|G(z)\| = O(1)$  with high probability for  $z \in S_{out}(\tau)$ . Thus for all the estimates that used the Ward's identities in Lemma 4.1, we can replace them with a simpler bound: for any deterministic vector  $\mathbf{u} \in \mathbb{R}^{\mathcal{I}}$ ,

$$\sum_{\mathfrak{a} \in \mathcal{I}} |G_{\mathbf{u} \mathfrak{a}}|^2 = (\bar{G}G)_{\mathbf{u} \mathbf{u}} = O(1) \quad \text{with high probability.} \quad (5.51)$$

Finally, in calculating the moments, we need a rough bound

$$\mathbb{E} \left| \sqrt{N} \langle \mathbf{u}, (G(z) - \Pi(z)) \mathbf{u} \rangle \right|^k < 1, \quad (5.52)$$

for any fixed  $k \in \mathbb{N}$  and any deterministic unit vector  $\mathbf{u} \in \mathbb{R}^{\mathcal{I}_1}$ . For  $z \in S_{out}(\tau)$  with  $\text{Im } z \geq N^{-C}$ , this follows from (4.23) and Lemma 4.3 (iii), where the second moment bound on  $|\sqrt{N} \langle \mathbf{u}, (G(z) - \Pi(z)) \mathbf{u} \rangle|^k$  follows from the trivial bound (4.7). This is the only place where we need the condition  $\eta \geq N^{-C}$ . Now plugging (4.23), (4.24), (5.51) and (5.52) into the arguments between (5.7) and (5.49), we can conclude (5.4).  $\square$

## 6 CLT for general functions

In this section, we prove the following weaker version of Theorem 2.6 and Theorem 2.8 under (5.1).

**Proposition 6.1.** *Theorem 2.6 and Theorem 2.8 hold under the moment assumption (5.1).*

As for Proposition 5.1, our proof of Proposition 6.1 is also based on a moment calculation. More precisely, we will prove the following counterpart of Lemma 5.2.

**Lemma 6.2.** *Suppose  $N^{-1+c_1} \leq \eta \leq 1$ ,  $d_N$ ,  $X$  and  $\Sigma$  satisfy Assumption 2.5, and (5.1) holds. Fix any  $E > 0$ ,  $k \in \mathbb{N}$  and constants  $a, b > 0$ . Then for any deterministic unit vectors  $\mathbf{v}_1, \dots, \mathbf{v}_k \in \mathbb{R}^n$  and functions  $f_1, \dots, f_k \in \mathcal{C}^{1,a,b}(\mathbb{R}_+)$ , we have*

$$\mathbb{E} [Z_{\eta,E}(\mathbf{v}_1, f_1) \cdots Z_{\eta,E}(\mathbf{v}_k, f_k)] = \begin{cases} \sum \prod \varpi(f_s, f_t, \mathbf{v}_s, \mathbf{v}_t) + O_{<}(n^{-c}), & \text{if } l \in 2\mathbb{N} \\ O_{<}(n^{-c}), & \text{otherwise} \end{cases}, \quad (6.1)$$

for some constant  $c > 0$ . Here  $\varpi(f_i, f_j, \mathbf{v}_i, \mathbf{v}_j) \equiv \varpi^{(N)}(f_i, f_j, \mathbf{v}_i, \mathbf{v}_j)$  is defined as

$$\begin{aligned} \varpi^{(N)}(f_i, f_j, \mathbf{v}_i, \mathbf{v}_j) &:= \frac{\eta}{\pi^2} \iint_{x_1, x_2} f_i(x_1) f_j(x_2) \alpha(E + x_1\eta, E + x_2\eta, \mathbf{v}_i, \mathbf{v}_j) dx_1 dx_2 \\ &+ \frac{1}{\pi^2} PV \iint_{x_1, x_2} \frac{f_i(x_1) f_j(x_2)}{x_1 - x_2} \beta(E + x_1\eta, E + x_2\eta, \mathbf{v}_i, \mathbf{v}_j) dx_1 dx_2 \\ &+ 2 \int_x f_i(x) f_j(x) \frac{\rho_{2c}(E + x\eta)}{(E + x\eta)^2} \left( \frac{\Sigma}{(1 + m_{2c}(E + x\eta)\Sigma)(1 + \bar{m}_{2c}(E + x\eta)\Sigma)} \right)_{\mathbf{v}_i \mathbf{v}_j}^2 dx, \end{aligned}$$

and  $\sum \prod$  means summing over all distinct ways of partitions into pairs.

*Proof of Proposition 6.1.* By Wick's theorem, (6.1) with  $E = 0$  and  $\eta = 1$  shows that the convergence in Theorem 2.6 holds in the sense of moments, which further implies the weak convergence. The reader may be worried about that in Theorem 2.6,  $E$  was taken to be 0, which is not the case in Proposition 6.1. However, this actually does not cause any trouble. Since  $\text{supp}(f_i) \subset \mathbb{R}_+$ , there exists a constant  $c > 0$  such that  $f_i(x) \equiv 0$  for all  $1 \leq i \leq k$  and  $0 \leq x \leq c$ . Hence we can take  $E = c/2$  and apply Proposition 6.1 with  $\eta = 1$  to the functions  $g_i(x) \in \mathcal{C}^{1,a,b}(\mathbb{R}_+)$  defined through  $g_i(x) = f_i(x + E)$ ,  $x \in \mathbb{R}_+$ .

Under the setting of Theorem 2.8, by Wick's theorem, (6.1) shows that the random vector  $(Z_{\eta,E}(\mathbf{v}_i, f_i))_{1 \leq i \leq k}$  converges weakly (in the sense of moments) to a Gaussian vector. Moreover, the covariance function can be simplified if we take  $\eta = o(1)$  in  $\varpi(f_i, f_j, \mathbf{v}_i, \mathbf{v}_j)$  and use (4.15)-(4.17):

$$\varpi^{(N)}(f_i, f_j, \mathbf{v}_i, \mathbf{v}_j) := \frac{1}{\pi^2} PV \iint_{x_1, x_2} \frac{f_i(x_1) f_j(x_2)}{x_1 - x_2} \beta(E, E, \mathbf{v}_i, \mathbf{v}_j) dx_1 dx_2$$

$$\begin{aligned}
& + 2 \int_x f_i(x) f_j(x) \frac{\rho_{2c}(E)}{E^2} \left( \frac{\Sigma}{(1 + m_{2c}(E)\Sigma)(1 + \bar{m}_{2c}(E)\Sigma)} \right)_{\mathbf{v}_i \mathbf{v}_j}^2 dx + O(\sqrt{\eta}) \\
& = 2 \int_x f_i(x) f_j(x) \frac{\rho_{2c}(E)}{E^2} \left( \frac{\Sigma}{(1 + m_{2c}(E)\Sigma)(1 + \bar{m}_{2c}(E)\Sigma)} \right)_{\mathbf{v}_i \mathbf{v}_j}^2 dx + O(\sqrt{\eta}),
\end{aligned}$$

where in the second step we used  $\beta(E, E, \mathbf{v}_i, \mathbf{v}_i) = 0$ . Taking  $N \rightarrow \infty$ , we get (2.22).  $\square$

The proof of Lemma 6.2 is based on the proof for Lemma 5.2. More precisely, we will use the Helffer-Sjöstrand formula in Lemma 4.8 to reduce the problem into the study of the CLT for the process  $Y(\mathbf{u}, w)$ . Denote  $\tilde{\eta} = N^{-\varepsilon_0} \eta$  for some small constant  $\varepsilon_0 > 0$  and abbreviate

$$f_\eta(x) := f\left(\frac{x-E}{\eta}\right), \quad \tilde{f}_\eta(x+iy) = f_\eta(x) + i(f_\eta(x+y) - f_\eta(x)).$$

Let  $\chi \in C_c^\infty(\mathbb{R})$  be a smooth cutoff function as in Lemma 4.8 satisfying that (i)  $\chi(y) = 1$  for  $|y| \leq 1$ , (ii)  $\chi(y) = 0$  for  $|y| \geq 2$ , and (iii)  $\|\chi^{(k)}\|_\infty = O(1)$  for any fixed  $k \in \mathbb{N}$ . Then using Lemma 4.8, we obtain that

$$\langle \mathbf{u}, f_\eta(\tilde{\mathcal{Q}}_1) \mathbf{u} \rangle = \frac{1}{\pi} \int_{\mathbb{C}} \mathbf{u}^T \frac{\partial_{\bar{z}}(\tilde{f}_\eta(z)\chi(y/\tilde{\eta}))}{\tilde{\mathcal{Q}}_1 - z} \mathbf{u} d^2z = \int_{\mathbb{C}} \phi_f(z) (\mathcal{G}_1)_{\mathbf{u}\mathbf{u}}(z) d^2z, \quad (6.2)$$

where we used (4.4) in the second step, and

$$\phi_f(x+iy) := \frac{1}{2\pi} \left[ (i-1)(f'_\eta(x+y) - f'_\eta(x))\chi(y/\tilde{\eta}) - \frac{1}{\tilde{\eta}}(f_\eta(x+y) - f_\eta(x))\chi'(y/\tilde{\eta}) + \frac{i}{\tilde{\eta}}f_\eta(x)\chi'(y/\tilde{\eta}) \right].$$

For simplicity, the bulk of the proof is devoted to the calculation of the moments

$$\mathbb{E} [Z_{\eta,E}^k(\mathbf{v}, f)], \quad k \in \mathbb{N}, \quad \mathbf{v} \in \mathbb{R}^{\mathcal{I}_1}, \quad f \in C^{1,a,b}(\mathbb{R}_+).$$

The proof for the more general expression in (6.1) is exactly the same, except for some immaterial changes of notations. We will describe it briefly at the end of the proof. Denoting  $\mathbf{u} := O\mathbf{v}$ , we have

$$Z(f) \equiv Z_{\eta,E}(\mathbf{v}, f) := \sqrt{\frac{N}{\eta}} \left( \langle \mathbf{u}, f\left(\eta^{-1}(\tilde{\mathcal{Q}}_1 - E)\right) \mathbf{u} \rangle - \int_{\lambda_-}^{\lambda_+} f\left(\frac{x-E}{\eta}\right) dF_{1c,\mathbf{v}}(x) \right).$$

By (6.2), we can write

$$\mathbb{E} [Z(f)]^k = \frac{1}{\eta^{k/2}} \int \phi_f(z_1) \cdots \phi_f(z_k) \frac{1}{\sqrt{|y_1| \cdots |y_k|}} \mathbb{E} [Y(z_1) \cdots Y(z_k)] d^2z_1 \cdots d^2z_k, \quad (6.3)$$

where we abused the notations a little bit and denote

$$Y(z_i) \equiv Y(\mathbf{u}, z_i) := \sqrt{N|y_i|} \langle \mathbf{u}, (\mathcal{G}_1 - z_i^{-1}\Pi(z_i)) \mathbf{u} \rangle, \quad z_i := x + iy_i, \quad 1 \leq i \leq k.$$

Recall that with the anisotropic local law (4.21), we only have the estimates  $Y(z) < 1$  for  $\text{Im } z \gg N^{-1}$ . In the next lemma, we generalize this bound to  $z$  with  $\text{Im } z \leq N^{-1}$ .

**Lemma 6.3.** *Suppose (4.21) holds for all  $z \in \mathbf{D}$  with  $q < N^{-1/2}$ . Then for any deterministic unit vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^{\mathcal{I}_1}$ , we have*

$$|\langle \mathbf{u}, G(X, z)\mathbf{v} \rangle - \langle \mathbf{u}, \Pi(z)\mathbf{v} \rangle| < (N\eta)^{-1/2} + (N\eta)^{-1}, \quad (6.4)$$

for all  $z \in \mathbf{S}(\omega, N) := \{z \in \mathbb{C}_+ : |z| \geq \omega, 0 < \eta \leq \omega^{-1}\}$ .



*Proof.* By Theorem 4.5, we know that (6.4) holds for  $z \in \mathbf{S}(\omega, N)$  with  $\eta \geq N^{-1+\varepsilon}$  for any small constant  $\varepsilon > 0$ . Then to prove (6.4), it suffices to show that (4.21) holds for  $z \in \mathbf{S}(\omega, N)$  with  $\eta \leq \eta_0 := N^{-1+\varepsilon}$ . For  $z = E + i\eta \in \mathbf{S}(\omega, N)$  with  $\eta \leq \eta_0$ , we denote  $z_0 := E + i\eta_0$ . Then using the spectral decomposition (4.5), we have

$$\begin{aligned}
|G_{\mathbf{u}\mathbf{v}}(z) - G_{\mathbf{u}\mathbf{v}}(z_0)| &\lesssim \sum_{k=1}^n \frac{\eta_0 |\langle \mathbf{u}, \xi_k \rangle| |\langle \mathbf{v}, \xi_k \rangle|}{|(\lambda_k - E - i\eta)(\lambda_k - E - i\eta_0)|} \\
&\leq \eta_0 \left( \sum_{k=1}^n \frac{|\langle \mathbf{u}, \xi_k \rangle|^2}{|(\lambda_k - E - i\eta)|^2} \right)^{1/2} \left( \sum_{k=1}^n \frac{|\langle \mathbf{v}, \xi_k \rangle|^2}{|(\lambda_k - E - i\eta_0)|^2} \right)^{1/2} \\
&\leq \eta_0 \left( \frac{\eta_0^2}{\eta^2} \sum_{k=1}^n \frac{|\langle \mathbf{u}, \xi_k \rangle|^2}{|(\lambda_k - E - i\eta_0)|^2} \right)^{1/2} \sqrt{\frac{\operatorname{Im}[z_0^{-1} G_{\mathbf{v}\mathbf{v}}(z_0)]}{\eta_0}} \\
&= \frac{\eta_0}{\eta} \sqrt{\operatorname{Im}[z_0^{-1} G_{\mathbf{u}\mathbf{u}}(z_0)] \operatorname{Im}[z_0^{-1} G_{\mathbf{v}\mathbf{v}}(z_0)]} < \frac{N^\varepsilon}{N\eta}, \tag{6.5}
\end{aligned}$$

where in the third and fourth steps we used the identity

$$\sum_{k=1}^n \frac{|\langle \mathbf{v}, \xi_k \rangle|^2}{|(\lambda_k - E - i\eta_0)|^2} = \frac{\operatorname{Im}[z_0^{-1} G_{\mathbf{u}\mathbf{u}}(z_0)]}{\eta_0},$$

and in the last step we used (6.4) for  $G(z_0)$ . On the other hand, using (4.15), we get  $|\Pi(z) - \Pi(z_0)| = O(1)$ . Together with (6.5) and (4.21) for  $G(z_0)$ , we get that

$$|G_{\mathbf{u}\mathbf{v}}(z) - \Pi_{\mathbf{u}\mathbf{v}}(z)| < 1 + \frac{N^\varepsilon}{N\eta} + \frac{1}{\sqrt{N\eta_0}}, \quad \eta \leq N^{-1+\varepsilon}.$$

Since  $\varepsilon$  is arbitrary, we get (6.4).  $\square$

With the above lemma, we obtain the following a priori estimates on  $Y(z)$ :

$$|Y(z)| < 1 + (Ny)^{-1/2}, \quad z = x + iy, \quad |z| \geq \omega, \quad 0 < y \leq \omega^{-1}. \tag{6.6}$$

Moreover, by the rough (4.7), we have the deterministic bound  $|y||Y(z)| = O(1)$ . Hence combining (6.6) with Lemma 4.3 (iii), we obtain that for any fixed  $k \in \mathbb{N}$  and  $y > 0$ ,

$$\mathbb{E}|Y(z)|^k = |y|^{-k} \mathbb{E}|yY(z)|^k < \left(1 + (Ny)^{-1/2}\right)^k.$$

We will use this bound tacitly in the proof.

## 6.1 The bad region

In the following proof, we let  $\sigma := N^{-\varepsilon_1}\eta$  for some constant  $\varepsilon_1 > \varepsilon_0$ , which we will choose later. We define the “good” region

$$\mathcal{R} := \{z_1, z_2, \dots, z_k \in \mathbb{C} : |y_1|, \dots, |y_k| \in [\sigma, 2\tilde{\eta}]\}.$$

In this subsection, we show that the integral in (6.3) over the “bad” region  $\mathcal{R}^c$  is negligible. For this purpose, we need to bound the following two integrals

$$\int_{|y| \leq \sigma} \left| \phi_f(z) \left( \frac{1}{\sqrt{\eta}|y|} + \frac{1}{|y|\sqrt{N\eta}} \right) \right| d^2z, \quad \int_{\sigma \leq |y| \leq 2\tilde{\eta}} \left| \phi_f(z) \left( \frac{1}{\sqrt{\eta}|y|} + \frac{1}{|y|\sqrt{N\eta}} \right) \right| d^2z.$$

Note that by definition, we have  $\phi_f(z) = 0$  for  $|y| \geq 2\tilde{\eta}$ .

Since  $\chi'(y/\tilde{\eta}) = 0$  for  $|y| \leq \tilde{\eta}$ , we get that

$$\begin{aligned} \int_{|y| \leq \sigma} \left| \phi_f(z) \left( \frac{1}{\sqrt{\eta|y|}} + \frac{1}{|y|\sqrt{N\eta}} \right) \right| d^2z &\lesssim \int_{|y| \leq \sigma} \frac{|f'_\eta(x+y) - f'_\eta(x)|}{\sqrt{\eta|y|}} d^2z + \frac{1}{\sqrt{N\eta}} \int_{|y| \leq \sigma} \frac{|f'_\eta(x+y) - f'_\eta(x)|}{|y|} d^2z \\ &= \sqrt{\frac{\sigma}{\eta}} \int_{|\tilde{y}| \leq 1} \frac{|f'(\tilde{x} + \tilde{y}N^{-\varepsilon_1}) - f'(\tilde{x})|}{\sqrt{|\tilde{y}|}} d\tilde{x}d\tilde{y} + \frac{1}{\sqrt{N\eta}} \int_{|\tilde{y}| \leq 1} \frac{|f'(\tilde{x} + \tilde{y}N^{-\varepsilon_1}) - f'(\tilde{x})|}{|\tilde{y}|} d\tilde{x}d\tilde{y}, \end{aligned} \quad (6.7)$$

where in the second step we apply the following change of variable:

$$\tilde{x} = \frac{x - E}{\eta}, \quad \tilde{y} := \frac{y}{\sigma}.$$

By the Hölder continuity and decay of  $f$ , we know

$$|f'(\tilde{x} + \tilde{y}N^{-\varepsilon_1}) - f'(\tilde{x})| \leq C \min\{(|\tilde{y}|N^{-\varepsilon_1})^a, (1 + |\tilde{x}|)^{-1-b}\} \leq C \frac{(|\tilde{y}|N^{-\varepsilon_1})^{pa}}{(1 + |\tilde{x}|)^{(1-p)(1+b)}}, \quad (6.8)$$

for all  $p \in [0, 1]$ . Choosing  $p = \frac{b}{2(1+b)}$ , we then have  $(1-p)(1+b) = 1 + b/2 > 1$ . Then the integrals in (6.7) can be bounded as

$$\begin{aligned} \int_{|\tilde{y}| \leq 1} \frac{|f'(\tilde{x} + \tilde{y}N^{-\varepsilon_1}) - f'(\tilde{x})|}{\sqrt{|\tilde{y}|}} d\tilde{x}d\tilde{y} &\leq \int_{|\tilde{y}| \leq 1} \frac{|f'(\tilde{x} + \tilde{y}N^{-\varepsilon_1}) - f'(\tilde{x})|}{|\tilde{y}|} d\tilde{x}d\tilde{y} \\ &\leq CN^{-pa\varepsilon_1} \int_{|\tilde{y}| \leq 1} \frac{|\tilde{y}|^{-1+pa}}{(1 + |\tilde{x}|)^{1+b/2}} d\tilde{x}d\tilde{y} \leq CN^{-pa\varepsilon_1}. \end{aligned}$$

Thus (6.7) gives (recall that  $\eta \geq N^{-1+c_1}$ )

$$\int_{|y| \leq \sigma} \left| \phi_f(z) \left( \frac{1}{\sqrt{\eta|y|}} + \frac{1}{|y|\sqrt{N\eta}} \right) \right| d^2z \leq CN^{-pa\varepsilon_1} (N^{-\varepsilon_1/2} + N^{-c_1/2}). \quad (6.9)$$

Similarly, we can show that

$$\int_{\sigma \leq |y| \leq \tilde{\eta}} |\phi_f(z)| \left| \left( \frac{1}{\sqrt{\eta|y|}} + \frac{1}{|y|\sqrt{N\eta}} \right) \right| d^2z \leq CN^{-pa\varepsilon_0} (N^{-\varepsilon_0/2} + N^{-c_1/2}).$$

On the other hand, we have

$$\begin{aligned} \int_{\tilde{\eta} \leq |y| \leq 2\tilde{\eta}} |\phi_f(z)| \left| \left( \frac{1}{\sqrt{\eta|y|}} + \frac{1}{|y|\sqrt{N\eta}} \right) \right| d^2z &= \int_{1 \leq |\tilde{y}| \leq 2} |\psi_f(\tilde{x}, \tilde{y})| \left| \left( \frac{N^{\varepsilon_0/2}}{\sqrt{\tilde{y}}} + \frac{N^{\varepsilon_0}}{|\tilde{y}|\sqrt{N\eta}} \right) \right| d\tilde{x}d\tilde{y} \\ &\lesssim N^{\varepsilon_0/2} + N^{\varepsilon_0 - c_1/2}, \end{aligned}$$

where

$$\psi_f(\tilde{x}, \tilde{y}) := \frac{1}{2\pi} [N^{-\varepsilon_0}(i-1)(f'(\tilde{x} + N^{-\varepsilon_0}\tilde{y}) - f'(\tilde{x}))\chi(\tilde{y}) - (f(\tilde{x} + N^{-\varepsilon_0}\tilde{y}) - f(\tilde{x}))\chi'(\tilde{y}) + if(\tilde{x})\chi'(\tilde{y})].$$

Combining the above two estimates with (6.9), we get

$$\int \left| \phi_f(z) \left( \frac{1}{\sqrt{\eta|y|}} + \frac{1}{|y|\sqrt{N\eta}} \right) \right| d^2z \leq CN^{\varepsilon_0/2}, \quad (6.10)$$

as long as we choose  $\varepsilon_0 < c_1$ .

Now with (6.6), (6.9) and (6.10), we obtain that

$$\begin{aligned} & \frac{1}{\eta^{k/2}} \int_{\mathcal{R}^c} \phi_f(z_1) \cdots \phi_f(z_k) \frac{1}{\sqrt{|y_1| \cdots |y_k|}} \mathbb{E}[Y(z_1) \cdots Y(z_k)] d^2 z_1 \cdots d^2 z_k \\ & < \frac{1}{\eta^{k/2}} \int_{|y_1| \leq \sigma, \dots, |y_k| \leq \sigma} \prod_{i=1}^k \left| \phi_f(z_i) \left( \frac{1}{\sqrt{\eta|y_i|}} + \frac{1}{|y_i| \sqrt{N\eta}} \right) \right| d^2 z_1 \cdots d^2 z_k \lesssim N^{-\varepsilon_1/2} \cdot N^{(k-1)\varepsilon_0/2} \leq N^{-\varepsilon_0}, \end{aligned}$$

as long as we choose the constants such that  $(k+1)\varepsilon_0 < \varepsilon_1 < c_1/2$ .

## 6.2 The good region

To estimate (6.3), it remains to deal with the integral over the good region  $\mathcal{R}$ , that is,

$$\mathbb{E}[Z(f)]^k = \frac{1}{\eta^{k/2}} \int_{\mathcal{R}} \phi_f(z_1) \cdots \phi_f(z_k) \frac{1}{\sqrt{|y_1| \cdots |y_k|}} \mathbb{E}\mathfrak{G} d^2 z_1 \cdots d^2 z_k + O_{<}(N^{-\varepsilon_0/2}), \quad (6.11)$$

where we have abbreviated  $\mathfrak{G} := Y(z_1) \cdots Y(z_k)$ . Then for  $\mathfrak{G}$ , we can apply the results in Lemma 5.2. Note that on  $\mathcal{R}$ , we can simplify (6.6) as

$$|Y(z)| < 1, \quad z = x + iy, \quad |z| \geq \omega, \quad \sigma < y \leq \omega^{-1}. \quad (6.12)$$

We can perform the same calculations between (5.7) and (5.49) to  $\mathbb{E}\mathfrak{G}$ . The only difference is that for  $\mathfrak{G}$  in (5.7), the imaginary parts of the spectral parameters are all of a fixed scale  $\eta$ , while for  $\mathfrak{G}$  in the current case, the imaginary parts of the spectral parameters are in the range  $\sigma \leq y_i \leq 2\tilde{\eta}$ . However, the calculations after (5.7) can be easily adapted to the current setting, and gives a similar expression as in (5.50):

$$\mathbb{E}\mathfrak{G} = \sqrt{|y_1 y_s|} \gamma(z_1, z_s, \mathbf{u}, \mathbf{u}) \mathbb{E} \prod_{t \notin \{1, s\}} Y(z_t) + O_{<}((N\sigma)^{-1/2}). \quad (6.13)$$

The  $\eta$  factor in (5.50) is replaced with  $\sqrt{|y_1 y_s|}$  because the scaling  $\sqrt{N\eta}$  in  $Y(\mathbf{u}_t, w_t)$  of (5.50) is replaced with  $\sqrt{N|y_s|}$  in  $Y(z_s)$  here. In case the reader is worried about the real parts of  $z_i$ 's, we remark that because  $\text{supp}(f) \subset \mathbb{R}_+$ , the integral in (6.11) is nonzero only when

$$\frac{x_i + y_i - E}{\eta} \geq 0 \quad \text{and} \quad \frac{x_i - E}{\eta} \geq 0 \Rightarrow x_i \geq E - 2\tilde{\eta}, \quad (6.14)$$

for all  $1 \leq i \leq k$ . Thus we have  $x_i \gtrsim 1$  for all  $1 \leq i \leq k$ , which is required in the calculations leading to (5.50).

Plugging (6.13) in (6.11) and using (6.10), we obtain that for  $k \geq 2$ ,

$$\begin{aligned} \mathbb{E}[Z(f)]^k &= (k-1) \left( \frac{1}{\eta} \int_{\sigma \leq |y_1|, |y_s| \leq 2\tilde{\eta}} \phi_f(z_1) \phi_f(z_s) \gamma(z_1, z_s, \mathbf{u}, \mathbf{u}) d^2 z_1 d^2 z_s \right) \mathbb{E}[Z(f)]^{k-2} \\ &+ O_{<}(N^{-\varepsilon_0} + N^{k\varepsilon_0/2} (N\sigma)^{-1/2}), \end{aligned} \quad (6.15)$$

where  $\sigma = N^{-\varepsilon_1} \eta \geq N^{-1+c_1-\varepsilon_1}$ . Recall we have chosen the constants such that  $(k+1)\varepsilon_0 < \varepsilon_1 < c_1/2$ , so we can bound  $N^{k\varepsilon_0/2} (N\sigma)^{-1/2} \leq N^{-\varepsilon_0/2}$ . On the other hand, when  $k = 1$ , by (5.25) we have

$$\mathbb{E}\mathfrak{G} < (N\sigma)^{-1/2}. \quad (6.16)$$

Together with (6.11) and (6.10), we get

$$\mathbb{E}Z(f) < N^{-\varepsilon_0} + N^{k\varepsilon_0/2}(N\sigma)^{-1/2} \lesssim N^{-\varepsilon_0/2}, \quad (6.17)$$

which verifies the mean zero condition of  $\mathcal{G}_i$  in Proposition 6.1.

For (6.15), it remains to study the factor

$$\mathcal{F}(z_1, z_2) := \frac{1}{\eta} \int_{\sigma \leq |y_1|, |y_2| \leq 2\tilde{\eta}} \phi_f(z_1) \phi_f(z_2) \gamma(z_1, z_2) d^2 z_1 d^2 z_2,$$

where we have taken  $s = 2$  and abbreviated  $\gamma(z_1, z_2) \equiv \gamma(z_1, z_2, \mathbf{u}, \mathbf{u}) = \hat{\alpha}(z_1, z_2, \mathbf{u}, \mathbf{u}) + \hat{\beta}(z_1, z_2, \mathbf{u}, \mathbf{u})$ . Here we recall (2.24) and (2.25):

$$\begin{aligned} \hat{\alpha}(z_1, z_2) &\equiv \hat{\alpha}(z_1, z_2, \mathbf{u}, \mathbf{u}) := \frac{m_{2c}(z_1)m_{2c}(z_2)}{3Nz_1z_2} \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \kappa_A(i, \mu) \left( O^\top \frac{\Lambda^{1/2}}{1 + m_{2c}(z_1)\Lambda} \mathbf{u} \right)_i \left( O^\top \frac{\Lambda^{1/2}}{1 + m_{2c}(z_2)\Lambda} \mathbf{u} \right)_i, \\ \hat{\beta}(z_1, z_2) &\equiv \hat{\beta}(z_1, z_2, \mathbf{u}, \mathbf{u}) := 2 \frac{m_{2c}(z_1) - m_{2c}(z_2)}{z_1z_2(z_1 - z_2)} \left( \mathbf{u}^\top \frac{\Lambda}{(1 + m_{2c}(z_1)\Lambda)(1 + m_{2c}(z_2)\Lambda)} \mathbf{u} \right)^2. \end{aligned}$$

We decompose  $\phi_f$  as

$$\phi_f(z) = \phi_1 + \phi_2 + \phi_3,$$

where

$$\phi_1 := \frac{(i-1)}{2\pi} (f'_\eta(x+y) - f'_\eta(x)) \chi(y/\tilde{\eta}), \quad \phi_2 := -\frac{1}{2\pi\tilde{\eta}} (f_\eta(x+y) - f_\eta(x)) \chi'(y/\tilde{\eta}), \quad \phi_3 := \frac{i}{2\pi\tilde{\eta}} f_\eta(x) \chi'(y/\tilde{\eta}).$$

Correspondingly, we decompose

$$\mathcal{F}(z_1, z_2) = \mathcal{F}_{ij}(z_1, z_2), \quad i, j = 1, 2, 3,$$

where

$$\mathcal{F}_{ij} = \mathcal{F}_{ji} := \frac{1}{\eta} \int_{\sigma \leq |y_1|, |y_2| \leq 2\tilde{\eta}} \phi_i(z_1) \phi_j(z_2) \gamma(z_1, z_2) d^2 z_1 d^2 z_2.$$

We will show that  $\mathcal{F}_{33}$  is the main term, while the other  $\mathcal{F}_{ij}$  are all error terms.

### 6.2.1 The error terms

By (6.14), we have  $|z_1| \gtrsim 1$  and  $|z_2| \gtrsim 1$ . Then we can bound  $\gamma(z_1, z_2)$  in the following two cases. If  $|z_1 - z_2| \geq |y_1|/2$ , then using (4.15) and (4.17), we get

$$|\gamma(z_1, z_2)| \lesssim |y_1|^{-1}. \quad (6.18)$$

If  $|z_1 - z_2| < |y_1|/2$ , then using (4.15), (4.16) and (4.17), we get

$$|\gamma(z_1, z_2)| \lesssim \frac{1}{\min_{1 \leq k \leq 2L} |z_1 - a_k|^{1/2}} \leq \min \left\{ \frac{1}{\min_{1 \leq k \leq 2L} |x_1 - a_k|^{1/2}}, |y_1|^{-1/2} \right\} \leq |y_1|^{-1}. \quad (6.19)$$

Then we can bound  $\mathcal{F}_{11}$  as

$$|\mathcal{F}_{11}| \lesssim \frac{1}{\eta} \int_{|y_1| \leq 2\tilde{\eta}, |y_2| \leq 2\tilde{\eta}} \left| \frac{f'_\eta(x_1 + y_1) - f'_\eta(x_1)}{y_1} \right| |f'_\eta(x_2 + y_2) - f'_\eta(x_2)| d^2 z_1 d^2 z_2$$

$$\begin{aligned}
&= N^{-\varepsilon_0} \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2} \left| \frac{f'(\tilde{x}_1 + \tilde{y}_1 N^{-\varepsilon_0}) - f'(\tilde{x}_1)}{\tilde{y}_1} \right| |f'(\tilde{x}_2 + \tilde{y}_2 N^{-\varepsilon_0}) - f'(\tilde{x}_2)| d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\
&\lesssim N^{-\varepsilon_0} \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2} \frac{(|\tilde{y}_1| N^{-\varepsilon_0})^{pa}}{|\tilde{y}_1| (1 + |\tilde{x}_1|)^{(1-p)(1+b)}} \frac{1}{(1 + |\tilde{x}_2|)^{1+b}} d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \leq CN^{-(1+pa)\varepsilon_0},
\end{aligned}$$

where in the second step we apply the following change of variable:

$$\tilde{x}_i = \frac{x_i - E}{\eta}, \quad \tilde{y}_i := \frac{y_i}{\tilde{\eta}}, \quad i = 1, 2,$$

and in the third step we used (6.8) with  $p = \frac{b}{2(1+b)}$ . Similarly, we can bound  $\mathcal{F}_{12}$  and  $\mathcal{F}_{13}$  as following:

$$\begin{aligned}
|\mathcal{F}_{12}| &\lesssim \frac{1}{\eta \tilde{\eta}} \int_{|y_1| \leq 2\tilde{\eta}, |y_2| \leq 2\tilde{\eta}} \left| \frac{f'_\eta(x_1 + y_1) - f'_\eta(x_1)}{y_1} \right| |f_\eta(x_2 + y_2) - f_\eta(x_2)| d^2 z_1 d^2 z_2 \\
&= \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2} \left| \frac{f'(\tilde{x}_1 + \tilde{y}_1 N^{-\varepsilon_0}) - f'_\eta(\tilde{x}_1)}{\tilde{y}_1} \right| |f(\tilde{x}_2 + \tilde{y}_2 N^{-\varepsilon_0}) - f(\tilde{x}_2)| d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\
&\lesssim \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2} \frac{(|\tilde{y}_1| N^{-\varepsilon_0})^{pa}}{|\tilde{y}_1| (1 + |\tilde{x}_1|)^{(1-p)(1+b)}} \frac{1}{(1 + |\tilde{x}_2|)^{1+b}} d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \lesssim N^{-pa\varepsilon_0},
\end{aligned}$$

and

$$\begin{aligned}
|\mathcal{F}_{13}| &\lesssim \frac{1}{\eta \tilde{\eta}} \int_{|y_1| \leq 2\tilde{\eta}, |y_2| \leq 2\tilde{\eta}} \left| \frac{f'_\eta(x_1 + y_1) - f'_\eta(x_1)}{y_1} \right| |f_\eta(x_2)| d^2 z_1 d^2 z_2 \\
&= \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2} \left| \frac{f'(\tilde{x}_1 + \tilde{y}_1 N^{-\varepsilon_0}) - f'_\eta(\tilde{x}_1)}{\tilde{y}_1} \right| |f(\tilde{x}_2)| d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\
&\lesssim \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2} \frac{(|\tilde{y}_1| N^{-\varepsilon_0})^{pa}}{|\tilde{y}_1| (1 + |\tilde{x}_1|)^{(1-p)(1+b)}} \frac{1}{(1 + |\tilde{x}_2|)^{1+b}} d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \lesssim N^{-pa\varepsilon_0}.
\end{aligned}$$

Using (6.18) and (6.19), we can bound  $\mathcal{F}_{22}$  as

$$\begin{aligned}
|\mathcal{F}_{22}| &\lesssim \frac{1}{\eta \tilde{\eta}^2} \int_{|y_1| \leq 2\tilde{\eta}, |y_2| \leq 2\tilde{\eta}} \left| \frac{f_\eta(x_1 + y_1) - f_\eta(x_1)}{y_1} \right| |f_\eta(x_2 + y_2) - f_\eta(x_2)| d^2 z_1 d^2 z_2 \\
&= N^{\varepsilon_0} \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2} \left| \frac{f(\tilde{x}_1 + \tilde{y}_1 N^{-\varepsilon_0}) - f(\tilde{x}_1)}{\tilde{y}_1} \right| |f(\tilde{x}_2 + \tilde{y}_2 N^{-\varepsilon_0}) - f(\tilde{x}_2)| d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\
&\lesssim N^{-\varepsilon_0} \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2} \frac{1}{(1 + |\tilde{x}_1|)^{1+b}} \frac{1}{(1 + |\tilde{x}_2|)^{1+b}} d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \lesssim N^{-\varepsilon_0},
\end{aligned}$$

where in the last step we used

$$|f(\tilde{x}_1 + \tilde{y}_1 N^{-\varepsilon_0}) - f(\tilde{x}_1)| \lesssim \frac{|\tilde{y}_1| N^{-\varepsilon_0}}{(1 + |\tilde{x}_1|)^{1+b}}. \quad (6.20)$$

To bound  $\mathcal{F}_{23}$ , we need better bounds on  $\gamma(z_1, z_2)$ . We decompose the integral in  $\mathcal{F}_{23}$  as

$$\begin{aligned}
\mathcal{F}_{23} &= \frac{1}{\eta} \int_{\sigma \leq |y_1|, |y_2| \leq 2\tilde{\eta}, |x_1 - x_2| \geq \eta N^{-\varepsilon_0/2}} \phi_2(z_1) \phi_3(z_2) \gamma(z_1, z_2) d^2 z_1 d^2 z_2 \\
&\quad + \frac{1}{\eta} \int_{\sigma \leq |y_1|, |y_2| \leq 2\tilde{\eta}, |x_1 - x_2| < \eta N^{-\varepsilon_0/2}} \phi_2(z_1) \phi_3(z_2) \gamma(z_1, z_2) d^2 z_1 d^2 z_2 =: \mathcal{F}_{23}^{(1)} + \mathcal{F}_{23}^{(2)}.
\end{aligned}$$

For  $\mathcal{F}_{23}^{(1)}$ , we use the bound  $|\gamma(z_1, z_2)| \lesssim \eta^{-1} N^{\varepsilon_0/2}$  when  $|x_1 - x_2| > \eta N^{-\varepsilon_0/2}$  to get that

$$\begin{aligned} |\mathcal{F}_{23}^{(1)}| &\lesssim \frac{N^{\varepsilon_0/2}}{\eta^2 \tilde{\eta}^2} \int_{|y_1| \leq 2\tilde{\eta}, |y_2| \leq 2\tilde{\eta}, |x_1 - x_2| \geq \eta N^{-\varepsilon_0/2}} |f_\eta(x_1 + y_1) - f_\eta(x_1)| |f_\eta(x_2)| \, d^2 z_1 d^2 z_2 \\ &\leq N^{\varepsilon_0/2} \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2} |f(\tilde{x}_1 + \tilde{y}_1 N^{-\varepsilon_0}) - f(\tilde{x}_1)| |f(\tilde{x}_2)| \, d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\ &\lesssim N^{\varepsilon_0/2} \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2} \frac{|\tilde{y}_1| N^{-\varepsilon_0}}{(1 + |\tilde{x}_1|)^{1+b}} \frac{1}{(1 + |\tilde{x}_2|)^{1+b}} \, d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \lesssim N^{-\varepsilon_0/2}. \end{aligned}$$

On the other hand, using (6.19) and (6.20) the term  $\mathcal{F}_{23}^{(2)}$  can be bounded as

$$\begin{aligned} |\mathcal{F}_{23}^{(2)}| &\lesssim \frac{1}{\eta \tilde{\eta}^2} \int_{|y_1| \leq 2\tilde{\eta}, |y_2| \leq 2\tilde{\eta}, |x_1 - x_2| < \eta N^{-\varepsilon_0/2}} \left| \frac{f_\eta(x_1 + y_1) - f_\eta(x_1)}{y_1} \right| |f_\eta(x_2)| \, d^2 z_1 d^2 z_2 \\ &= N^{\varepsilon_0} \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2, |\tilde{x}_1 - \tilde{x}_2| < N^{-\varepsilon_0/2}} \left| \frac{f(\tilde{x}_1 + \tilde{y}_1 N^{-\varepsilon_0}) - f(\tilde{x}_1)}{\tilde{y}_1} \right| |f(\tilde{x}_2)| \, d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\ &\lesssim \int_{|\tilde{y}_1| \leq 2, |\tilde{y}_2| \leq 2, |\tilde{x}_1 - \tilde{x}_2| < N^{-\varepsilon_0/2}} \frac{1}{(1 + |\tilde{x}_1|)^{1+b}} \frac{1}{(1 + |\tilde{x}_2|)^{1+b}} \, d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \lesssim N^{-\varepsilon_0/2}. \end{aligned}$$

In sum, we have obtained that

$$\sum_{i=1}^2 \sum_{j=1}^3 |\mathcal{F}_{ij}| \lesssim N^{-\varepsilon_0/2} + N^{-pa\varepsilon_0}. \quad (6.21)$$

## 6.2.2 The main term

It remains to study the main term

$$\begin{aligned} \mathcal{F}_{33}(z_1, z_2) &:= -\frac{1}{4\pi^2 \eta \tilde{\eta}^2} \int_{\tilde{\eta} \leq |y_1|, |y_2| \leq 2\tilde{\eta}} f_\eta(x_1) f_\eta(x_2) \left( \hat{\alpha}(z_1, z_2) + \hat{\beta}(z_1, z_2) \right) \chi'(y_1/\tilde{\eta}) \chi'(y_2/\tilde{\eta}) \, d^2 z_1 d^2 z_2 \\ &=: \mathcal{K}_1 + \mathcal{K}_2. \end{aligned}$$

For term  $\mathcal{K}_1$  containing the  $\hat{\alpha}$  factor, we first consider the integral over the region  $R_{++} := \{\tilde{\eta} \leq y_1 \leq 2\tilde{\eta}, \tilde{\eta} \leq y_2 \leq 2\tilde{\eta}\}$ ,

$$\begin{aligned} (\mathcal{K}_1)_{++} &:= -\frac{1}{4\pi^2 \eta \tilde{\eta}^2} \int_{R_{++}} f_\eta(x_1) f_\eta(x_2) \hat{\alpha}(z_1, z_2) \chi'(y_1/\tilde{\eta}) \chi'(y_2/\tilde{\eta}) \, d^2 z_1 d^2 z_2 \\ &= -\frac{\eta}{4\pi^2} \iint_{1 \leq \tilde{y}_1, 2 \leq \tilde{y}_2} f(\tilde{x}_1) f(\tilde{x}_2) \chi'(\tilde{y}_1) \chi'(\tilde{y}_2) \alpha((E + \tilde{x}_1 \eta) + i\tilde{y}_1 \tilde{\eta}, (E + \tilde{x}_2 \eta) + i\tilde{y}_2 \tilde{\eta}) \, d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2. \end{aligned}$$

With (4.16), we can obtain that

$$|\hat{\alpha}((E + \tilde{x}_1 \eta) + i\tilde{y}_1 \tilde{\eta}, (E + \tilde{x}_2 \eta) + i\tilde{y}_2 \tilde{\eta}) - \hat{\alpha}_{++}(E + \tilde{x}_1 \eta, E + \tilde{x}_2 \eta)| \lesssim \sqrt{\tilde{\eta}}. \quad (6.22)$$

Here for  $x_1, x_2 \in \mathbb{R}_+$ , and  $\mathbf{a}, \mathbf{b} \in \{+, -\}$ , we define

$$\hat{\alpha}_{\mathbf{a}\mathbf{b}}(x_1, x_2) := \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \frac{\kappa_4(i, \mu)}{3N} \left[ \frac{m_{2c}^{\mathbf{a}}(x_1)}{x_1} \left( O^\top \frac{\Lambda^{1/2}}{1 + m_{2c}^{\mathbf{a}}(x_1) \Lambda} \mathbf{u} \right)_i \right]^2 \left[ \frac{m_{2c}^{\mathbf{b}}(x_2)}{x_2} \left( O^\top \frac{\Lambda^{1/2}}{1 + m_{2c}^{\mathbf{b}}(x_2) \Lambda} \mathbf{u} \right)_i \right]^2,$$

where for a complex number  $z \in \mathbb{C}$ , we used the notations  $z^+ := z$  and  $z^- := \bar{z}$ . Thus  $(\mathcal{K}_1)_{++}$  can be reduced to

$$\begin{aligned} (\mathcal{K}_1)_{++} &= -\frac{\eta}{4\pi^2} \iint_{1 \leq \tilde{y}_{1,2} \leq 2} f(\tilde{x}_1) f(\tilde{x}_2) \chi'(\tilde{y}_1) \chi'(\tilde{y}_2) \hat{\alpha}_{++}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 + \mathcal{O}(\sqrt{\tilde{\eta}}) \\ &= -\frac{\eta}{4\pi^2} \iint f(\tilde{x}_1) f(\tilde{x}_2) \hat{\alpha}_{++}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{x}_2 + \mathcal{O}(n^{-\varepsilon_0/2}). \end{aligned}$$

Similarly, we can calculate the integral in  $\mathcal{K}_1$  over the other three regions:  $(\mathcal{K}_1)_{+-}$  for  $R_{+-} := \{\tilde{\eta} \leq y_1 \leq 2\tilde{\eta}, -2\tilde{\eta} \leq y_2 \leq -\tilde{\eta}\}$ ,  $(\mathcal{K}_1)_{-+}$  for  $R_{-+} := \{-2\tilde{\eta} \leq y_1 \leq -\tilde{\eta}, \tilde{\eta} \leq y_2 \leq 2\tilde{\eta}\}$ , and  $(\mathcal{K}_1)_{--}$  for  $R_{--} := \{-2\tilde{\eta} \leq y_1 \leq -\tilde{\eta}, -2\tilde{\eta} \leq y_2 \leq -\tilde{\eta}\}$ . Combining all these four integrals, we obtain that

$$\begin{aligned} \mathcal{K}_1 &= -\frac{\eta}{4\pi^2} \iint f(\tilde{x}_1) f(\tilde{x}_2) (\hat{\alpha}_{++} + \hat{\alpha}_{--} - \hat{\alpha}_{+-} - \hat{\alpha}_{-+})(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{x}_2 + \mathcal{O}(n^{-\varepsilon_0/2}) \\ &= \frac{\eta}{\pi^2} \iint f(\tilde{x}_1) f(\tilde{x}_2) \alpha(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta, O^\top \mathbf{u}, O^\top \mathbf{u}) d\tilde{x}_1 d\tilde{x}_2 + \mathcal{O}(n^{-\varepsilon_0/2}). \end{aligned} \quad (6.23)$$

where for  $x_1, x_2 \in \mathbb{R}$ ,  $\alpha$  is defined as (recall (2.16))

$$\begin{aligned} &\alpha(x_1, x_2, O^\top \mathbf{u}, O^\top \mathbf{u}) \\ &:= \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \frac{\kappa_4(i, \mu)}{3N} \operatorname{Im} \left[ \frac{m_{2c}(x_1)}{x_1} \left( O^\top \frac{\Lambda^{1/2}}{1 + m_{2c}(x_1)\Lambda} \mathbf{u} \right)_i \right]^2 \operatorname{Im} \left[ \frac{m_{2c}(x_2)}{x_2} \left( O^\top \frac{\Lambda^{1/2}}{1 + m_{2c}(x_2)\Lambda} \mathbf{u} \right)_i \right]^2. \end{aligned}$$

Then we study the term  $\mathcal{K}_2$  containing the  $\hat{\beta}$  factor. We introduce the notations

$$\tilde{\beta}(z_1, z_2) := \frac{1}{z_1 z_2} \left( \mathbf{u}^\top \frac{1}{1 + m_{2c}(z_1)\Lambda} \Lambda \frac{1}{1 + m_{2c}(z_2)\Lambda} \mathbf{u} \right)^2,$$

and for  $x_1, x_2 \in \mathbb{R}_+$ ,

$$\tilde{\beta}_{\mathbf{a}\mathbf{b}}(x_1, x_2) := \frac{1}{x_1 x_2} \left( \mathbf{u}^\top \frac{1}{1 + m_{2c}^{\mathbf{a}}(x_1)\Lambda} \Lambda \frac{1}{1 + m_{2c}^{\mathbf{b}}(x_2)\Lambda} \mathbf{u} \right)^2, \quad \mathbf{a}, \mathbf{b} \in \{+, -\}.$$

Then we can write

$$\hat{\beta}(z_1, z_2) := 2 \frac{m_{2c}(z_1) - m_{2c}(z_2)}{z_1 - z_2} \tilde{\beta}(z_1, z_2).$$

Now we first consider the integral in  $\mathcal{K}_2$  over the region  $R_{++}$ :

$$\begin{aligned} (\mathcal{K}_2)_{++} &:= -\frac{1}{4\pi^2 \eta \tilde{\eta}^2} \int_{R_{++}} f_\eta(x_1) f_\eta(x_2) \hat{\beta}(z_1, z_2) \chi'(y_1/\tilde{\eta}) \chi'(y_2/\tilde{\eta}) d^2 z_1 d^2 z_2 \\ &= -\frac{1}{2\pi^2} \iint_{1 \leq \tilde{y}_{1,2} \leq 2} f(\tilde{x}_1) f(\tilde{x}_2) \chi'(\tilde{y}_1) \chi'(\tilde{y}_2) \frac{m_{2c}((E + \tilde{x}_1\eta) + i\tilde{y}_1\tilde{\eta}) - m_{2c}((E + \tilde{x}_2\eta) + i\tilde{y}_2\tilde{\eta})}{(\tilde{x}_1 - \tilde{x}_2) + i(\tilde{y}_1 - \tilde{y}_2)N^{-\varepsilon_0}} \\ &\quad \times \tilde{\beta}((E + \tilde{x}_1\eta) + i\tilde{y}_1\tilde{\eta}, (E + \tilde{x}_2\eta) + i\tilde{y}_2\tilde{\eta}) d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\ &= -\frac{1}{2\pi^2} \iint_{1 \leq \tilde{y}_{1,2} \leq 2} f(\tilde{x}_1) f(\tilde{x}_2) \chi'(\tilde{y}_1) \chi'(\tilde{y}_2) \frac{m_{2c}((E + \tilde{x}_1\eta) + i\tilde{y}_1\tilde{\eta}) - m_{2c}((E + \tilde{x}_2\eta) + i\tilde{y}_2\tilde{\eta})}{(\tilde{x}_1 - \tilde{x}_2) + i(\tilde{y}_1 - \tilde{y}_2)N^{-\varepsilon_0}} \\ &\quad \times \tilde{\beta}_{++}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 + \mathcal{O}(N^{-\varepsilon_0/2}), \end{aligned} \quad (6.24)$$

where we used a similar bound for  $\tilde{\beta}$  as in (6.22):

$$\left| \tilde{\beta}((E + \tilde{x}_1\eta) + i\tilde{y}_1\tilde{\eta}, (E + \tilde{x}_2\eta) + i\tilde{y}_2\tilde{\eta}) - \tilde{\beta}_{++}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) \right| \lesssim \sqrt{\tilde{\eta}}, \quad (6.25)$$

and the bound

$$\begin{aligned} & \iint_{1 \leq \tilde{y}_{1,2} \leq 2} |f(\tilde{x}_1) f(\tilde{x}_2)| \left| \frac{m_{2c}((E + \tilde{x}_1\eta) + i\tilde{y}_1\tilde{\eta}) - m_{2c}((E + \tilde{x}_2\eta) + i\tilde{y}_2\tilde{\eta})}{(\tilde{x}_1 - \tilde{x}_2) + i(\tilde{y}_1 - \tilde{y}_2)N^{-\varepsilon_0}} \right| d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\ & \lesssim \iint_{1 \leq \tilde{y}_{1,2} \leq 2} |f(\tilde{x}_1) f(\tilde{x}_2)| \frac{\sqrt{\tilde{\eta}}}{|\tilde{x}_1 - \tilde{x}_2|^{1/2} + |\tilde{y}_1 - \tilde{y}_2|^{1/2} N^{-\varepsilon_0/2}} d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 = O(1) \end{aligned}$$

using (4.16).

For the integral in (6.24), we decompose it as  $(\mathcal{K}_2)_{++} = (\mathcal{K}_2^{(1)})_{++} + (\mathcal{K}_2^{(2)})_{++}$ , where  $(\mathcal{K}_2^{(1)})_{++}$  contains the integral over the region with  $|\tilde{x}_1 - \tilde{x}_2| \leq N^{-\varepsilon}$ , and  $(\mathcal{K}_2^{(2)})_{++}$  contains the integral over the region with  $|\tilde{x}_1 - \tilde{x}_2| > N^{-\varepsilon}$ . Here  $\varepsilon$  is a sufficiently small constant such that

$$0 < \varepsilon < \varepsilon_0/10.$$

For  $(\mathcal{K}_2^{(1)})_{++}$ , we have

$$\begin{aligned} |(\mathcal{K}_2^{(1)})_{++}| & \lesssim \iint_{1 \leq \tilde{y}_{1,2} \leq 2, |\tilde{x}_1 - \tilde{x}_2| \leq N^{-\varepsilon}} \frac{\eta^{1/2} |f(\tilde{x}_1) f(\tilde{x}_2)|}{|\tilde{x}_1 - \tilde{x}_2|^{1/2} + |\tilde{y}_1 - \tilde{y}_2|^{1/2} N^{-\varepsilon_0/2}} \tilde{\beta}_{++}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 + O(N^{-\varepsilon_0}) \\ & \lesssim \iint_{1 \leq \tilde{y}_{1,2} \leq 2, |\tilde{x}_1 - \tilde{x}_2| \leq N^{-\varepsilon}} \frac{|f(\tilde{x}_1) f(\tilde{x}_2)|}{|\tilde{x}_1 - \tilde{x}_2|^{1/2}} \tilde{\beta}_{++}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \lesssim N^{-\varepsilon/2}, \end{aligned}$$

where we used (4.16) in the first step. For  $(\mathcal{K}_2^{(2)})_{++}$ , we have

$$\begin{aligned} (\mathcal{K}_2^{(2)})_{++} & = -\frac{1}{2\pi^2} \iint_{1 \leq \tilde{y}_{1,2} \leq 2, |\tilde{x}_1 - \tilde{x}_2| > N^{-\varepsilon}} f(\tilde{x}_1) f(\tilde{x}_2) \chi'(\tilde{y}_1) \chi'(\tilde{y}_2) \frac{m_{2c}(E + \tilde{x}_1\eta) - m_{2c}(E + \tilde{x}_2\eta)}{\tilde{x}_1 - \tilde{x}_2} \\ & \quad \times \tilde{\beta}_{++}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 + O(N^{-\varepsilon_0/2+\varepsilon}) \\ & = -\frac{1}{2\pi^2} \iint_{|\tilde{x}_1 - \tilde{x}_2| > N^{-\varepsilon}} f(\tilde{x}_1) f(\tilde{x}_2) \frac{m_{2c}(E + \tilde{x}_1\eta) - m_{2c}(E + \tilde{x}_2\eta)}{\tilde{x}_1 - \tilde{x}_2} \tilde{\beta}_{++}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{x}_2 + O(N^{-\varepsilon_0/4}) \\ & = -\frac{1}{2\pi^2} \iint f(\tilde{x}_1) f(\tilde{x}_2) \frac{m_{2c}(E + \tilde{x}_1\eta) - m_{2c}(E + \tilde{x}_2\eta)}{\tilde{x}_1 - \tilde{x}_2} \tilde{\beta}_{++}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{x}_2 + O(N^{-\varepsilon/2}), \end{aligned}$$

where in the first step we used

$$\frac{1}{(\tilde{x}_1 - \tilde{x}_2) + i(\tilde{y}_1 - \tilde{y}_2)N^{-\varepsilon_0}} = \frac{1}{\tilde{x}_1 - \tilde{x}_2} + O(N^{-\varepsilon_0+2\varepsilon}).$$

and

$$m_{2c}((E + \tilde{x}_i\eta) + i\tilde{y}_i\tilde{\eta}) - m_{2c}(E + \tilde{x}_i\eta) = O(N^{-\varepsilon_0/2}), \quad i = 1, 2,$$



by (4.16), and in the last step we used

$$\begin{aligned} & \iint_{|\tilde{x}_1 - \tilde{x}_2| \leq N^{-\varepsilon}} |f(\tilde{x}_1) f(\tilde{x}_2)| \left| \frac{m_{2c}(E + \tilde{x}_1\eta) - m_{2c}(E + \tilde{x}_2\eta)}{\tilde{x}_1 - \tilde{x}_2} \tilde{\beta}_{+++}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) \right| d\tilde{x}_1 d\tilde{x}_2 \\ & \lesssim \iint_{|\tilde{x}_1 - \tilde{x}_2| \leq N^{-\varepsilon}} \frac{|f(\tilde{x}_1) f(\tilde{x}_2)|}{|\tilde{x}_1 - \tilde{x}_2|^{1/2}} d\tilde{x}_1 d\tilde{x}_2 \lesssim N^{-\varepsilon/2}. \end{aligned}$$

In sum, we get that

$$(\mathcal{K}_2)_{++} = -\frac{1}{2\pi^2} \iint f(\tilde{x}_1) f(\tilde{x}_2) \frac{m_{2c}(E + \tilde{x}_1\eta) - m_{2c}(E + \tilde{x}_2\eta)}{\tilde{x}_1 - \tilde{x}_2} \tilde{\beta}_{+++}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{x}_2 + O(N^{-\varepsilon/2}). \quad (6.26)$$

Then we study the integrals  $(\mathcal{K}_2)_{+-}$ . Using (6.25) and (4.16), we can simplify that

$$\begin{aligned} (\mathcal{K}_2)_{+-} &= -\frac{1}{2\pi^2} \iint_{1 \leq \tilde{y}_1 \leq 2, -2 \leq \tilde{y}_2 \leq -1} f(\tilde{x}_1) f(\tilde{x}_2) \chi'(\tilde{y}_1) \chi'(\tilde{y}_2) \\ & \times \frac{m_{2c}((E + \tilde{x}_1\eta) + i\tilde{y}_1\tilde{\eta}) - m_{2c}((E + \tilde{x}_2\eta) + i\tilde{y}_2\tilde{\eta})}{(\tilde{x}_1 - \tilde{x}_2) + i(\tilde{y}_1 - \tilde{y}_2)N^{-\varepsilon_0}} \tilde{\beta}((E + \tilde{x}_1\eta) + i\tilde{y}_1\tilde{\eta}, (E + \tilde{x}_2\eta) + i\tilde{y}_2\tilde{\eta}) d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\ &= -\frac{1}{2\pi^2} \iint_{1 \leq \tilde{y}_1 \leq 2, -2 \leq \tilde{y}_2 \leq -1} f(\tilde{x}_1) f(\tilde{x}_2) \chi'(\tilde{y}_1) \chi'(\tilde{y}_2) \\ & \times \frac{m_{2c}(E + \tilde{x}_1\eta) - \overline{m}_{2c}(E + \tilde{x}_2\eta)}{(\tilde{x}_1 - \tilde{x}_2) + i(\tilde{y}_1 - \tilde{y}_2)N^{-\varepsilon_0}} \tilde{\beta}_{+-}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\ & + O\left( \iint_{1 \leq \tilde{y}_1 \leq 2, -2 \leq \tilde{y}_2 \leq -1} \frac{N^{-\varepsilon_0/2} |f(\tilde{x}_1) f(\tilde{x}_2)|}{|\tilde{x}_1 - \tilde{x}_2| + |\tilde{y}_1 - \tilde{y}_2| N^{-\varepsilon_0}} d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \right). \end{aligned}$$

We can bound the second term as

$$\begin{aligned} & \iint_{1 \leq \tilde{y}_1 \leq 2, -2 \leq \tilde{y}_2 \leq -1} \frac{N^{-\varepsilon_0/2} |f(\tilde{x}_1) f(\tilde{x}_2)|}{|\tilde{x}_1 - \tilde{x}_2| + |\tilde{y}_1 - \tilde{y}_2| N^{-\varepsilon_0}} d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\ & \lesssim \iint_{1 \leq \tilde{y}_1 \leq 2, -2 \leq \tilde{y}_2 \leq -1} \frac{N^{-\varepsilon_0/2}}{|\tilde{x}_1 - \tilde{x}_2| + 2N^{-\varepsilon_0}} \frac{1}{(1 + |\tilde{x}_1|)^{1+b}} \frac{1}{(1 + |\tilde{x}_2|)^{1+b}} d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \lesssim N^{-\varepsilon_0/2} \log N. \end{aligned}$$

Then we can write

$$\begin{aligned} (\mathcal{K}_2)_{+-} &= -\frac{1}{2\pi^2} \iint_{1 \leq \tilde{y}_1 \leq 2, -2 \leq \tilde{y}_2 \leq -1} f(\tilde{x}_1) f(\tilde{x}_2) \chi'(\tilde{y}_1) \chi'(\tilde{y}_2) [m_{2c}(E + \tilde{x}_1\eta) - \overline{m}_{2c}(E + \tilde{x}_2\eta)] \tilde{\beta}_{+-}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) \\ & \times \operatorname{Re} \frac{1}{(\tilde{x}_1 - \tilde{x}_2) + i(\tilde{y}_1 - \tilde{y}_2)N^{-\varepsilon_0}} d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 \\ & + \frac{i}{2\pi^2} \iint_{1 \leq \tilde{y}_1 \leq 2, -2 \leq \tilde{y}_2 \leq -1} f(\tilde{x}_1) f(\tilde{x}_2) \chi'(\tilde{y}_1) \chi'(\tilde{y}_2) [m_{2c}(E + \tilde{x}_1\eta) - \overline{m}_{2c}(E + \tilde{x}_2\eta)] \tilde{\beta}_{+-}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) \\ & \times \operatorname{Im} \frac{1}{(\tilde{x}_1 - \tilde{x}_2) - i(\tilde{y}_1 - \tilde{y}_2)N^{-\varepsilon_0}} d\tilde{x}_1 d\tilde{y}_1 d\tilde{x}_2 d\tilde{y}_2 + O\left(N^{-\varepsilon_0/2}\right) =: (\mathcal{K}_2^{(1)})_{+-} + (\mathcal{K}_2^{(2)})_{+-} + O\left(N^{-\varepsilon_0/2} \log N\right). \end{aligned}$$

For the term  $(\mathcal{K}_2^{(1)})_{+-}$ , we observe that the integral converges to the Cauchy principal value, while for the term  $(\mathcal{K}_2^{(1)})_{+-}$ ,  $\pi^{-1} \operatorname{Im}[(\tilde{x}_1 - \tilde{x}_2) + i(\tilde{y}_1 - \tilde{y}_2)N^{-\varepsilon_0}]^{-1}$  is an approximate delta function. More precisely, we have the following estimates. The proof is pretty standard, so we omit the details.

**Lemma 6.4.** *Suppose  $g(x_1, x_2)$  is 1/2-Hölder continuous uniformly in  $x_1$  and  $x_2$ , and  $|g(x_1, x_2)| \leq C(1 + |x_1|)^{-(1+b)}(1 + |x_2|)^{-(1+b)}$  for some constant  $C > 0$ . Then for any  $0 < \delta \ll 1$ , we have*

$$\left| \frac{1}{\pi} \iint_{x_1, x_2} g(x_1, x_2) \operatorname{Im} \frac{1}{(x_1 - x_2) - i\delta} dx_1 dx_2 - \int_{x_1} g(x_1, x_1) dx_1 \right| \lesssim \delta^{1/2},$$

and

$$\left| \iint_{x_1, x_2} g(x_1, x_2) \operatorname{Re} \frac{1}{(x_1 - x_2) + i\delta} dx_1 dx_2 - PV \iint_{x_1, x_2} \frac{g(x_1, x_2)}{x_1 - x_2} dx_1 dx_2 \right| \lesssim \delta^{1/3},$$

where

$$PV \iint_{x_1, x_2} \frac{g(x_1, x_2)}{x_1 - x_2} dx_1 dx_2 := \lim_{\delta \downarrow 0} \iint_{x_1, x_2} g(x_1, x_2) \operatorname{Re} \frac{1}{(x_1 - x_2) + i\delta} dx_1 dx_2.$$

With Lemma 6.4, we obtain that

$$\begin{aligned} (\mathcal{K}_2^{(1)})_{+-} &= \frac{1}{2\pi^2} PV \iint \frac{f(\tilde{x}_1) f(\tilde{x}_2)}{\tilde{x}_1 - \tilde{x}_2} [m_{2c}(E + \tilde{x}_1\eta) - \overline{m_{2c}(E + \tilde{x}_2\eta)}] \tilde{\beta}_{+-}(E + \tilde{x}_1\eta, E + \tilde{x}_2\eta) d\tilde{x}_1 d\tilde{x}_2 \\ &\quad + O(N^{-\varepsilon_0/3}). \end{aligned} \quad (6.27)$$

and

$$\begin{aligned} (\mathcal{K}_2^{(2)})_{+-} &= -\frac{i}{2\pi} \int f^2(\tilde{x}_1) [m_{2c}(E + \tilde{x}_1\eta) - \overline{m_{2c}(E + \tilde{x}_1\eta)}] \tilde{\beta}_{+-}(E + \tilde{x}_1\eta, E + \tilde{x}_1\eta) d\tilde{x}_1 + O(N^{-\varepsilon_0/2}) \\ &= \int f^2(\tilde{x}_1) \rho_{2c}(E + \tilde{x}_1\eta) \cdot \tilde{\beta}_{+-}(E + \tilde{x}_1\eta, E + \tilde{x}_1\eta) d\tilde{x}_1 + O(N^{-\varepsilon_0/2}), \end{aligned} \quad (6.28)$$

where we used  $\rho_{2c}(x) = \pi^{-1} \operatorname{Im} m_{2c}(x)$ .

Using (6.26), (6.27), (6.28),  $(\mathcal{K}_2)_{--} = \overline{(\mathcal{K}_2)_{++}}$  and  $(\mathcal{K}_2)_{-+} = \overline{(\mathcal{K}_2)_{+-}}$ , we obtain that

$$\begin{aligned} \mathcal{K}_2 &= (\mathcal{K}_2)_{++} + (\mathcal{K}_2)_{+-} + (\mathcal{K}_2)_{-+} + (\mathcal{K}_2)_{--} \\ &= \frac{1}{\pi^2} PV \iint \frac{f(x_1) f(x_2)}{x_1 - x_2} \beta(x_1, x_2, \mathbf{v}, \mathbf{v}) dx_1 dx_2 \\ &\quad + 2 \int f^2(x) \frac{\rho_{2c}(E + x\eta)}{(E + x\eta)^2} \left( \mathbf{u}^\top \frac{\Lambda}{(1 + m_{2c}(x)\Lambda)(1 + \overline{m_{2c}(x)\Lambda})} \mathbf{u} \right)^2 dx + O(N^{-\varepsilon/2}), \end{aligned} \quad (6.29)$$

where recall that  $\mathbf{v} = O^\top \mathbf{u}$ , and for  $x_1, x_2 \in \mathbb{R}$ ,  $\beta$  is defined in (2.17).

Finally, plugging (6.21), (6.23) and (6.29) into (6.15), we obtain that

$$\mathbb{E}[Z(f)]^k = (k-1) \varpi(f, f, \mathbf{v}, \mathbf{v}) \mathbb{E}[Z(f)]^{k-2} + O_\prec(N^{-c})$$

for some constant  $c > 0$ . In general, we can extend this induction relation to more general expression in (6.1) and hence prove Lemma 6.2.

*Proof of Lemma 6.2.* We expand the left-hand side of (6.1) using Helffer-Sjöstrand formula, Lemma 4.8, and obtain the similar expression as in (6.3):

$$\mathbb{E}[Z_{\eta,E}(\mathbf{v}_1, f_1) \cdots Z_{\eta,E}(\mathbf{v}_k, f_k)] = \frac{1}{\eta^{k/2}} \int \frac{\phi_{f_1}(z_1) \cdots \phi_{f_k}(z_k)}{\sqrt{|y_1| \cdots |y_k|}} \mathbb{E}[Y(\mathbf{u}_1, z_1) \cdots Y(\mathbf{u}_k, z_k)] d^2 z_1 \cdots d^2 z_k.$$

Then applying the same arguments between (6.6) and (6.29), we can obtain that

$$\mathbb{E}[Z_{\eta,E}(\mathbf{v}_1, f_1) \cdots Z_{\eta,E}(\mathbf{v}_k, f_k)] = \sum_{s=2}^k \varpi(f_1, f_s, \mathbf{v}_1, \mathbf{v}_s) \mathbb{E} \prod_{t \notin \{1, s\}} Z_{\eta,E}(\mathbf{v}_t, f_t) + O_{<}(N^{-c})$$

for some constant  $c > 0$ . Using this induction relation and (6.17), we then conclude (6.1).  $\square$

## 7 Weaker moment assumptions

In this section we use a Green's function comparison argument to relax the moment assumptions in Proposition 5.1 and Proposition 6.1, and hence complete the proof of Theorems 2.6, 2.8, 2.10 and 2.11. In this section, we focus on the proof of Theorems 2.10 and 2.11. Later we will explain how to extend the arguments to Theorems 2.6 and 2.8.

For any fixed  $c_0 > 0$ , we can choose a constant  $0 < c_\phi < 1/2$  small enough such that

$$\left( (N/\eta)^{1/4} N^{-c_\phi} \right)^{a_\eta + c_0} \geq N^{2+\varepsilon_0}$$

for some constant  $\varepsilon_0 > 0$ . Then we introduce the following truncation  $X'$ , where

$$X'_{i\mu} = \mathbf{1}_{|X_{i\mu}| \leq \phi_N} X_{i\mu}, \quad \phi_N := \frac{N^{-c_\phi}}{(N\eta)^{1/4}}. \quad (7.1)$$

Without loss of generality, we can choose  $c_\phi$  small enough such that

$$\phi_N \geq (N\eta)^{-1/2}.$$

By the moment condition (2.21) and a simple union bound, we have

$$\mathbb{P}(X' \neq X) = O(N^{-\varepsilon_0}). \quad (7.2)$$

Using (2.21) and integration by parts, it is easy to verify that

$$\mathbb{E}|X_{i\mu}| \mathbf{1}_{|X_{i\mu}| > \phi_N} = O(N^{-2-\varepsilon_0}), \quad \mathbb{E}|X_{i\mu}|^2 \mathbf{1}_{|X_{i\mu}| > \phi_N} = O(N^{-2-\varepsilon_0}),$$

which imply that

$$|\mathbb{E}X'_{i\mu}| = O(N^{-2-\varepsilon_0}), \quad \mathbb{E}|X'_{i\mu}|^2 = N^{-1} + O(N^{-2-\varepsilon_0}). \quad (7.3)$$

Moreover, we trivially have

$$\mathbb{E}|X'_{i\mu}|^4 \leq \mathbb{E}|X_{i\mu}|^4 = O(N^{-4}).$$

Then we introduce the matrices

$$\mathring{X} = X' - \mathbb{E}X',$$

where by (7.3) we have

$$\|\mathbb{E}X'\| = O(N^{-1-\varepsilon_0}), \quad \text{Var}(\mathring{X}_{i\mu}) = N^{-1} (1 + O(N^{-1-\varepsilon_0})). \quad (7.4)$$

Then we can define  $\mathring{\mathcal{G}}_{1,2}(\mathring{X}, z)$  (see (2.7)) and  $\mathring{G}(\mathring{X}, z)$  (see (4.3)) by replacing  $X$  with  $\mathring{X}$ .

**Claim 7.1.** *Under the above setting, we have for any deterministic unit vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{C}^{\mathcal{I}}$ ,*

$$\left| \langle \mathbf{u}, G(X, z)\mathbf{v} \rangle - \langle \mathbf{u}, \mathring{G}(\mathring{X}, z)\mathbf{v} \rangle \right| < N^{-1-\varepsilon_0}\eta^{-1/2}$$

*uniformly in  $z \in \mathbf{D}$ .*

*Proof.* See the proof of Lemma 4.4 in Section A.1 of [45] □

Note that under the scaling  $\sqrt{N\eta}$  in the definition (5.2),  $N^{-1-\varepsilon_0}\eta^{-1/2}$  is a negligible error. Hence it suffices to prove that Theorem 2.10 holds under the following assumptions on  $X$ , which correspond to the above setting for  $\mathring{X}$ .

**Assumption 7.2.** *Fix a small constant  $\tau > 0$ .*

(i)  $X = (X_{i\mu})$  is a real  $n \times N$  matrix, whose entries are independent random variables that satisfy

$$\mathbb{E}X_{i\mu} = 0, \quad \mathbb{E}X_{i\mu}^2 = N^{-1} + \mathcal{O}(N^{-2-\varepsilon_0}), \quad (7.5)$$

and the following bounded support condition with  $\phi_N$ :

$$\max_{i,\mu} |X_{i\mu}| \leq \phi_N. \quad (7.6)$$

Moreover, we assume the matrix entries have bounded fourth moments

$$\max_{i,\mu} \mathbb{E}|X_{i\mu}|^4 \leq CN^{-4}. \quad (7.7)$$

(ii) Assumption 2.5 (ii)-(iii) hold.

Then the results in Section 4 can be extended to the setting with the above assumption. In particular, we have the following version of Theorem 4.5.

**Theorem 7.3** (Theorem 3.6 of [49]). *Suppose Assumption 7.2 holds. For any fixed  $\varepsilon > 0$  and deterministic unit vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{C}^{\mathcal{I}}$ , the following anisotropic local laws holds. For any  $z \in \mathbf{D}$ , we have*

$$|\langle \mathbf{u}, G(z)\mathbf{v} \rangle - \langle \mathbf{u}, \Pi(z)\mathbf{v} \rangle| < \phi_N + \Psi(z). \quad (7.8)$$

Given any random matrices  $X$  satisfying Assumption 7.2, we can construct matrices  $\tilde{X}$  and  $\tilde{Y}$  that match the first four moments as  $X$  and  $Y$  but with smaller support  $\phi_N < N^{-1/2}$ , which is the content of the next lemma.

**Lemma 7.4** (Lemma 5.1 of [31]). *Suppose  $X$  satisfies Assumption 7.2. Then there exists another matrix  $\tilde{X} = (\tilde{X}_{i\mu})$  such that  $\tilde{X}$  satisfies (2.1), (5.1) and the following moments matching conditions*

$$\mathbb{E}X_{i\mu}^k = [1 + \mathcal{O}(N^{-1-\varepsilon_0})] \mathbb{E}\tilde{X}_{i\mu}^k, \quad k = 2, 3, 4. \quad (7.9)$$

Note that  $\tilde{X}$  has bounded support  $\mathcal{O}_{<}(N^{-1/2})$  under the moment condition (5.1). We can define  $\tilde{G}(z) := G(\tilde{X}, z)$  and  $\tilde{Y}_{\eta,E}$  by replacing  $X$  with  $\tilde{X}$ . We have shown that Lemma 5.2 holds for  $\tilde{Y}_{\eta,E}$ . Then it remains to prove that  $(Y_{\eta,E}(\mathbf{u}_1, w_1), \dots, Y_{\eta,E}(\mathbf{u}_k, w_k))$  has the same moments as  $(\tilde{Y}_{\eta,E}(\mathbf{u}_1, w_1), \dots, \tilde{Y}_{\eta,E}(\mathbf{u}_k, w_k))$  asymptotically.

**Proposition 7.5.** *Under the setting of Theorem 2.10, for any deterministic unit vectors  $\mathbf{u}_1, \dots, \mathbf{u}_r \in \mathbb{R}^{\mathcal{I}_1}$  and fixed  $w_1, \dots, w_r \in \mathbb{H}$ ,*

$$\mathbb{E} \prod_{i=1}^r Y_{\eta, E}(\mathbf{u}_i, w_i) = \mathbb{E} \prod_{i=1}^r \tilde{Y}_{\eta, E}(\mathbf{u}_i, w_i) + O(n^{-\varepsilon}), \quad (7.10)$$

for some constant  $\varepsilon > 0$ .

*Proof.* To prove this proposition, we will use the continuous comparison method introduced in [29]. We first introduce the following interpolation.

**Definition 7.6** (Interpolating matrices). *Introduce the notations  $X^0 := \tilde{X}$  and  $X^1 := X$ . Let  $\rho_{i\mu}^0$  and  $\rho_{i\mu}^1$  be the laws of  $\tilde{X}_{i\mu}$  and  $X_{i\mu}$ , respectively. For  $\theta \in [0, 1]$ , we define the interpolated law*

$$\rho_{i\mu}^\theta := (1 - \theta)\rho_{i\mu}^0 + \theta\rho_{i\mu}^1.$$

Let  $\{X^\theta : \theta \in (0, 1)\}$  be a collection of random matrices such that the following properties hold. For any fixed  $\theta \in (0, 1)$ ,  $(X^0, X^\theta, X^1)$  is a triple of independent  $\mathcal{I}_1 \times \mathcal{I}_2$  random matrices, and the matrix  $X^\theta = (X_{i\mu}^\theta)$  has law

$$\prod_{i \in \mathcal{I}_1} \prod_{\mu \in \mathcal{I}_2} \rho_{i\mu}^\theta(dX_{i\mu}^\theta). \quad (7.11)$$

Note that we do not require  $X^{\theta_1}$  to be independent of  $X^{\theta_2}$  for  $\theta_1 \neq \theta_2 \in (0, 1)$ . For  $\lambda \in \mathbb{R}$ ,  $i \in \mathcal{I}_1$  and  $\mu \in \mathcal{I}_2$ , we define the matrix  $X_{(i\mu)}^{\theta, \lambda}$  as

$$\left( X_{(i\mu)}^{\theta, \lambda} \right)_{j\nu} := \begin{cases} X_{i\mu}^\theta, & \text{if } (j, \nu) \neq (i, \mu) \\ \lambda, & \text{if } (j, \nu) = (i, \mu) \end{cases}. \quad (7.12)$$

Correspondingly, we define the resolvents

$$G^\theta(z) := G(X^\theta, z), \quad G_{(i\mu)}^{\theta, \lambda}(z) := G(X_{(i\mu)}^{\theta, \lambda}, z).$$

and for  $1 \leq s \leq k$  (recall (5.6)),

$$Y_s^\theta := z_s Y_{\eta, E}(\mathbf{u}_s, w_s, X^\theta) = \sqrt{N\eta} \langle \mathbf{u}_s, (G^\theta(z_s) - \Pi(z_s)) \mathbf{u}_s \rangle, \quad (Y_s)_{(i\mu)}^{\theta, \lambda} := z_s Y_{\eta, E}(\mathbf{u}_s, w_s, X_{(i\mu)}^{\theta, \lambda}).$$

where  $z_s := E + w_s \eta$ .

Using (7.11) and fundamental calculus, we get the following basic interpolation formula.

**Lemma 7.7.** *For  $F : \mathbb{R}^{\mathcal{I}_1 \times \mathcal{I}_2} \rightarrow \mathbb{C}$  we have*

$$\frac{d}{d\theta} \mathbb{E} F(X^\theta) = \sum_{i \in \mathcal{I}_1, \mu \in \mathcal{I}_2} \left[ \mathbb{E} F \left( X_{(i\mu)}^{\theta, X_{i\mu}^1} \right) - \mathbb{E} F \left( X_{(i\mu)}^{\theta, X_{i\mu}^0} \right) \right], \quad (7.13)$$

provided all the expectations exist.

Then the main work is devoted to proving the following estimate for the right-hand side of (7.13). Note that combining Lemma 7.7 and Lemma 7.8, we can conclude Proposition 7.5.

**Lemma 7.8.** *Under the assumptions of Proposition 7.5, there exists a constant  $\varepsilon > 0$  such that*

$$\sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} \left[ \mathbb{E} F \left( X_{(i\mu)}^{\theta, X_{i\mu}^1} \right) - \mathbb{E} F \left( X_{(i\mu)}^{\theta, X_{i\mu}^0} \right) \right] \leq N^{-\varepsilon}, \quad (7.14)$$

for all  $\theta \in [0, 1]$ , where  $F(X^\theta) := \prod_{s=1}^k Y_s^\theta$ .

Underlying the proof of (7.14) is an expansion approach which we will describe below. We first rewrite the resolvent expansion (5.12) using the new notations: for any  $\lambda, \lambda' \in \mathbb{R}$  and  $K \in \mathbb{N}$ ,

$$G_{(i\mu)}^{\theta, \lambda'} = G_{(i\mu)}^{\theta, \lambda} + \sum_{k=1}^K (\lambda - \lambda')^k G_{(i\mu)}^{\theta, \lambda} \left( \Delta_{i\mu} G_{(i\mu)}^{\theta, \lambda} \right)^k + (\lambda - \lambda')^{K+1} G_{(i\mu)}^{\theta, \lambda'} \left( \Delta_{i\mu} G_{(i\mu)}^{\theta, \lambda} \right)^{K+1}. \quad (7.15)$$

With this expansion, we can prove the following estimate: suppose that  $y$  is a random variable satisfying  $|y| \leq \phi_N$ , then for any deterministic unit vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{C}^{\mathcal{I}}$  and  $z \in \mathbf{D}$ ,

$$\langle \mathbf{u}, \left( G_{(i\mu)}^{\theta, y}(z) - \Pi(z) \right) \mathbf{v} \rangle < \phi_N + \Psi(z), \quad i \in \mathcal{I}_1, \mu \in \mathcal{I}_2. \quad (7.16)$$

In fact, to prove this estimate, we will apply the expansion (7.15) with  $\lambda' = y$  and  $\lambda = X_{i\mu}^\theta$ , so that  $G_{(i\mu)}^{\theta, \lambda} = G^\theta$ . To bound the right-hand side of (7.16), we will use  $y \leq \phi_N$ ,  $|X_{i\mu}^\theta| \leq \phi_N$ , the anisotropic local law in Theorem 7.3 for  $G^\theta$ , and the trivial bound  $\|G_{(i\mu)}^{\theta, y}\| \leq C\eta^{-1}$ . Here we choose  $K$  such that  $\phi_N^K \eta^{-1} \leq 1$ , and hence the last term in (7.15) can be bounded by

$$(\lambda - \lambda')^{K+1} \left( G_{(i\mu)}^{\theta, \lambda'} \left( \Delta_{i\mu} G_{(i\mu)}^{\theta, \lambda} \right)^{K+1} \right)_{\mathbf{u}\mathbf{v}} < \phi_N^{K+1} \eta^{-1} < \phi_N.$$

Next we give the proof of (7.14) using (7.15) and (7.16).

*Proof of (7.14).* For simplicity, we only give the estimate for the case  $Y_s^\theta = Y^\theta$  for all  $1 \leq s \leq r$ , where

$$Y^\theta := \sqrt{N\eta} \langle \mathbf{u}, (G^\theta(z) - \Pi(z)) \mathbf{u} \rangle, \quad z = E + w\eta,$$

for any deterministic unit vector  $\mathbf{u} \in \mathbb{R}^{\mathcal{I}_1}$  and fixed  $w \in \mathbb{C}_+$ . In other words, we will show

$$\sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} \left[ \mathbb{E} \left( Y_{(i\mu)}^{\theta, X_{i\mu}^1} \right)^r - \mathbb{E} \left( Y_{(i\mu)}^{\theta, X_{i\mu}^0} \right)^r \right] \leq n^{-\varepsilon}. \quad (7.17)$$

The general multi-variable case can be dealt with in exactly the same way, except that the notations are a little more complicated.

Using (7.15) and (7.16), we get that for any random variable  $y$  satisfying  $|y| \leq \phi_N$  and any fixed  $K \in \mathbb{N}$ ,

$$Y_{(i\mu)}^{\theta, y} - Y_{(i\mu)}^{\theta, 0} = \sum_{k=1}^K \sqrt{N\eta} (-y)^k x_k(i, \mu) + O_{<}(\sqrt{N\eta} \phi_N^{K+1}), \quad (7.18)$$

where

$$x_k(i, \mu) := \langle \mathbf{u}, G_{(i\mu)}^{\theta, 0} \left( \Delta_{i\mu} G_{(i\mu)}^{\theta, 0} \right)^k \mathbf{u} \rangle. \quad (7.19)$$

In the following proof, we choose  $K \geq 3/c_\phi$  such that

$$\sqrt{N\eta} \phi_N^{K+1} \leq N^{-3}.$$

With (7.16), we trivially have  $x_k(i, \mu) < 1$  for  $k \geq 1$ . Moreover, we have the better bound for odd  $k$ :

$$x_k(i, \mu) < \phi_N, \quad k \in 2\mathbb{N} + 1. \quad (7.20)$$

This is because if  $k$  is odd, then there exists at least one  $(G_{(i\mu)}^{\theta,0})_{\mathbf{u}\mu}$  or  $(G_{(i\mu)}^{\theta,0})_{i\mu}$  factor in the expansion of  $x_k(i, \mu)$ . Using (7.20) for  $k = 1$  and the bound  $|y| \leq \phi_N$ , we can obtain the rough bound

$$\sqrt{N\eta}(-y)^k x_k(i, \mu) < N^{-kc_\phi}, \quad k \geq 1. \quad (7.21)$$

Now using (7.18) and (7.21), the Taylor expansion of  $(Y_{(i\mu)}^{\theta, X_{i\mu}^a})^r$  up to  $K$ -th order gives that for  $a \in \{0, 1\}$ ,

$$\begin{aligned} \mathbb{E} \left( Y_{(i\mu)}^{\theta, X_{i\mu}^a} \right)^r - \mathbb{E} \left( Y_{(i\mu)}^{\theta, 0} \right)^r &= \sum_{k=1}^{K \wedge r} \binom{r}{k} \mathbb{E} \left( Y_{(i\mu)}^{\theta, 0} \right)^{r-k} \left[ \sum_{l=1}^K \sqrt{N\eta} (-X_{i\mu}^{\theta, a})^l x_l(i, \mu) \right]^k + O_{<}(N^{-3}) \\ &= \sum_{s=1}^{K \wedge r} \sum_{k=1}^s \sum_{\mathbf{s}}^* \binom{r}{k} \mathbb{E} (-X_{i\mu}^{\theta, a})^{s\mathbb{E}} \left( Y_{(i\mu)}^{\theta, 0} \right)^{r-k} \prod_{l=1}^k \sqrt{N\eta} x_{s_l}(i, \mu) + O_{<}(N^{-3}), \end{aligned} \quad (7.22)$$

where the sum  $\sum_{\mathbf{s}}^*$  means the sum over  $\mathbf{s} = (s_1, \dots, s_k) \in \mathbb{N}^k$  satisfying

$$1 \leq s_i \leq K \wedge r, \quad \sum_{l=1}^k l \cdot s_l = s. \quad (7.23)$$

Here we only keep  $s \leq K$ , because otherwise by (7.21),

$$\prod_{l=1}^k \sqrt{N\eta} (-X_{i\mu}^{\theta, a})^{s_l} x_{s_l}(i, \mu) < N^{-Kc_\phi} \leq N^{-3}.$$

Then combining (7.22) with (7.9), we get that

$$\begin{aligned} \left| \mathbb{E} \left( Y_{(i\mu)}^{\theta, X_{i\mu}^1} \right)^r - \mathbb{E} \left( Y_{(i\mu)}^{\theta, X_{i\mu}^0} \right)^r \right| &< N^{-1-\varepsilon_0} \sum_{s=2}^4 \sum_{k=1}^s \sum_{\mathbf{s}}^* N^{-s/2} \mathbb{E} \left| \prod_{l=1}^k \sqrt{N\eta} x_{s_l}(i, \mu) \right| \\ &+ \sum_{s=5}^K \sum_{k=1}^s \sum_{\mathbf{s}}^* N^{-2} \phi_N^{s-4} \mathbb{E} \left| \prod_{l=1}^k \sqrt{N\eta} x_{s_l}(i, \mu) \right| + O_{<}(N^{-3}), \end{aligned}$$

where we used the moment bound  $\mathbb{E} |X_{i\mu}^{\theta, a}|^s \leq \phi_N^{s-4} \mathbb{E} |X_{i\mu}^{\theta, a}|^4 \lesssim \phi_N^{s-4} N^{-2}$  for  $s \geq 4$ . Thus to show (7.17), we only need to prove that there exists a constant  $\varepsilon > 0$  such that for  $s = 2, 3, 4$ ,

$$N^{-1-\varepsilon_0} \sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} N^{-s/2} \mathbb{E} \left| \prod_{l=1}^k \sqrt{N\eta} x_{s_l}(i, \mu) \right| < N^{-\varepsilon}, \quad (7.24)$$

and for any fixed  $s \geq 5$  and  $\mathbf{s}$  such that (7.23) holds,

$$\sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} N^{-2} \phi_N^{s-4} \mathbb{E} \left| \prod_{l=1}^k \sqrt{N\eta} x_{s_l}(i, \mu) \right| < N^{-\varepsilon}. \quad (7.25)$$

To prove these two estimates, we shall use the following bounds:

$$|x_s(i, \mu)| < \begin{cases} R_i^2 + R_\mu^2, & \text{if } s \geq 2, \\ R_i R_\mu + \phi_N (R_i^2 + R_\mu^2), & \text{if } s = 1, \end{cases} \quad (7.26)$$

where

$$R_i := |\langle \mathbf{u}, G^\theta \mathbf{t}_i \rangle|, \quad R_\mu := |\langle \mathbf{u}, G^\theta \mathbf{e}_\mu \rangle|.$$

In fact, by the definition (7.19), we have

$$x_k(i, \mu) < \begin{cases} |\langle \mathbf{u}, G_{(i\mu)}^{\theta,0} \mathbf{t}_i \rangle|^2 + |\langle \mathbf{u}, G_{(i\mu)}^{\theta,0} \mathbf{e}_\mu \rangle|^2, & \text{if } s \geq 2, \\ |\langle \mathbf{u}, G_{(i\mu)}^{\theta,0} \mathbf{t}_i \rangle| |\langle \mathbf{u}, G_{(i\mu)}^{\theta,0} \mathbf{e}_\mu \rangle|, & \text{if } s = 1. \end{cases} \quad (7.27)$$

On the other hand, using (7.15) and (7.16) we get

$$|\langle \mathbf{u}, G_{(i\mu)}^{\theta,0} \mathbf{t}_i \rangle| \leq |G_{\mathbf{u}\mathbf{t}_i}^\theta| + |X_{i\mu}^\theta| \left( |G_{\mathbf{u}\mu}^\theta| |\langle \mathbf{t}_i, G_{(i\mu)}^{\theta,0} \mathbf{t}_i \rangle| + |G_{\mathbf{u}\mathbf{t}_i}^\theta| |\langle \mathbf{e}_\mu, G_{(i\mu)}^{\theta,0} \mathbf{t}_i \rangle| \right) < R_i + \phi_N R_\mu, \quad (7.28)$$

and

$$|\langle \mathbf{u}, G_{(i\mu)}^{\theta,0} \mathbf{e}_\mu \rangle| \leq |G_{\mathbf{u}\mu}^\theta| + |X_{i\mu}^\theta| \left( |G_{\mathbf{u}\mu}^\theta| |\langle \mathbf{t}_i, G_{(i\mu)}^{\theta,0} \mathbf{e}_\mu \rangle| + |G_{\mathbf{u}\mathbf{t}_i}^\theta| |\langle \mathbf{e}_\mu, G_{(i\mu)}^{\theta,0} \mathbf{e}_\mu \rangle| \right) < R_\mu + \phi_N R_i. \quad (7.29)$$

Plugging (7.28) and (7.29) into (7.27), we obtain (7.26).

Note that by (4.1) and (7.8), the following estimates holds:

$$R_\mu < \phi_N + \Psi(z) \lesssim \phi_N, \quad \sum_{i \in \mathcal{I}_1} R_i^2 + \sum_{\mu \in \mathcal{I}_2} R_\mu^2 < \eta^{-1}. \quad (7.30)$$

where we used  $\phi_N \geq (N\eta)^{-1/2} \gtrsim \Psi(z)$  for the first estimate. Then with (7.26) and (7.30), we can bound the left-hand side of (7.24) by

$$\begin{aligned} N^{-1-\varepsilon_0} \sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} N^{-s/2} \mathbb{E} \left| \prod_{l=1}^k \sqrt{N\eta} x_{s_l}(i, \mu) \right| &< N^{-1-\varepsilon_0} \sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} N^{-s/2} (N\eta)^{k/2} (R_i^2 + R_\mu^2) \\ &< N^{-\varepsilon_0} N^{-(s-k)/2} \eta^{(k-2)/2} \leq N^{-\varepsilon_0}. \end{aligned}$$

This concludes (7.24). For the proof of (7.25), we consider the following three cases.

**Case 1:**  $s_l \geq 2$  for  $1 \leq l \leq k$ , which gives  $k \leq s/2$ . Then using (7.26) and (7.30), we can obtain that

$$\begin{aligned} \sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} N^{-2} \phi_N^{s-4} \mathbb{E} \left| \prod_{l=1}^k \sqrt{N\eta} x_{s_l}(i, \mu) \right| &< \sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} (N\eta)^{k/2} N^{-2} \phi_N^{s-4} (R_i^2 + R_\mu^2) \\ &< (N\eta)^{k/2-1} \phi_N^{s-4} \leq (N\eta \phi_N^4)^{s/4-1} \leq N^{-c_\phi}, \end{aligned}$$

where we used the definition of  $\phi_N$  in (7.1) and  $s \geq 5$  in the last step.

**Case 2:** There is only one  $l$  such that  $s_l = 1$ . Without loss of generality, we assume that  $s_1 = 1$  and  $s_l \geq 2$  for  $2 \leq l \leq k$ . Thus we have  $s \geq 2k - 1$ . Then using (7.26) and (7.30), we can obtain that

$$\begin{aligned} \sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} N^{-2} \phi_N^{s-4} \mathbb{E} \left| \prod_{l=1}^k \sqrt{N\eta} x_{s_l}(i, \mu) \right| &< \sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} (N\eta)^{k/2} N^{-2} \phi_N^{s-4} \phi_N (R_i^2 + R_\mu^2) \\ &< (N\eta)^{k/2-1} \phi_N^{s-3} \leq (N\eta \phi_N^4)^{(s+1)/4-1} \leq N^{-c_\phi}. \end{aligned}$$

**Case 3:** There are at least two  $l$ 's such that  $s_l = 1$ . Without loss of generality, we assume that  $s_1 = s_2 = \dots = s_j = 1$  for some  $2 \leq j \leq k$ . Thus we have  $s \geq 2(k-j) + j = 2k - j$ . Then using (7.26) and (7.30), we can obtain that

$$\sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} N^{-2} \phi_N^{s-4} \mathbb{E} \left| \prod_{l=1}^k \sqrt{N\eta} x_{s_l}(i, \mu) \right| < \sum_{i \in \mathcal{I}_1} \sum_{\mu \in \mathcal{I}_2} (N\eta)^{k/2} N^{-2} \phi_N^{s-4} \phi_N^{j-2} (R_i^2 R_\mu^2 + \phi_N^2 (R_i^4 + R_\mu^4))$$



$$< (N\eta)^{k/2} N^{-2} \phi_N^{s-4} \phi_N^{j-2} \left( \frac{1}{\eta} + \frac{N}{\eta} \phi_N^2 \right) \lesssim (N\eta)^{k/2-1} \phi_N^{s-4+j} \leq (N\eta \phi_N^4)^{(s+j)/4-1} \leq N^{-c_\phi}.$$

Combining the above three cases, we conclude (7.25). Then (7.24) and (7.25) together imply (7.14).  $\square$

Combining (7.13) and (7.14), we conclude the proof of Proposition 7.5.  $\square$

Finally, we complete the proof of the main theorems.

*Proof of Theorems 2.6, 2.8, 2.10 and 2.11.* Combining Proposition 7.5 with Lemma 5.2 for  $\tilde{Y}_{\eta,E}$ , we get that (5.3) holds even under the weaker moment assumption (2.21):

$$\mathbb{E}[Y(\mathbf{u}_1, w_1) \cdots Y(\mathbf{u}_k, w_k)] = \begin{cases} \sum \prod \eta \gamma(z_s, z_t, \mathbf{v}_s, \mathbf{v}_t) + O_{<}(N^{-\varepsilon}), & \text{if } k \in 2\mathbb{N}, \\ O_{<}(N^{-\varepsilon}), & \text{otherwise,} \end{cases} \quad (7.31)$$

for some constant  $\varepsilon > 0$ . By Wick's theorem, (7.31) shows that the convergence of  $(\mathcal{Y}_{\eta,E}(\mathbf{v}_1, w_1), \dots, \mathcal{Y}_{\eta,E}(\mathbf{v}_k, w_k))$  in Theorems 2.10 and 2.11 holds in the sense of moments, which further implies the weak convergence.

For the convergence of  $\eta^{-1/2}(\mathcal{Y}_{\eta,E}(\mathbf{v}_1, w_1), \dots, \mathcal{Y}_{\eta,E}(\mathbf{v}_k, w_k))$  in Theorem 2.11 for  $E \in S_{out}(\tau)$ , we can prove a similar comparison estimates as in (7.10):

$$\mathbb{E} \prod_{i=1}^r \eta^{-1/2} Y_{\eta,E}(\mathbf{u}_i, w_i) = \mathbb{E} \prod_{i=1}^r \eta^{-1/2} \tilde{Y}_{\eta,E}(\mathbf{u}_i, w_i) + O(n^{-\varepsilon}). \quad (7.32)$$

Its proof is similar to that of (7.10), so we omit the details. Then (7.32) together with Lemma 5.2 conclude the convergence of  $\eta^{-1/2}(\mathcal{Y}_{\eta,E}(\mathbf{v}_1, w_1), \dots, \mathcal{Y}_{\eta,E}(\mathbf{v}_k, w_k))$  for  $E \in S_{out}(\tau)$ .

Next Theorems 2.6 and 2.8 can be derived from (7.31) in the same way that Proposition 6.1 is derived from Lemma 5.2. As in Section 6, we apply the Helffer-Sjöstrand formula to get a similar expression as in (6.3). The only difference from the arguments in Section 6 is the local law for the  $Y(z)$  terms: under the weaker moment assumption (2.21), we only have the bound

$$|Y(z)| < \phi_N + (N\eta)^{-1/2}, \quad z \in \mathbf{D}.$$

We let  $\eta_1$  be such that  $(N\eta_1)^{-1/2} = \phi_N$ . Then for  $\text{Im } z \leq \eta_1$ , the local law (6.4) holds as before. For  $\text{Im } z > \eta_1$ , we do not have the high probability bound  $|Y(z)| < 1$ . However, by (7.31), we still have  $|\mathbb{E}[Y(z_1) \cdots Y(z_k)]| < 1$ . Then all the arguments after (6.3) will work and lead to Theorems 2.6 and 2.8.  $\square$

## References

- [1] T. W. Anderson. Asymptotic theory for principal component analysis. *The Annals of Mathematical Statistics*, 34(1):122–148, 1963.
- [2] Z. D. Bai. Convergence rate of expected spectral distributions of large random matrices. part II. sample covariance matrices. *Ann. Probab.*, 21(2):649–672, 1993.
- [3] Z. D. Bai, B. Q. Miao, and G. M. Pan. On asymptotics of eigenvectors of large sample covariance matrix. *Ann. Probab.*, 35(4):1532–1572, 2007.
- [4] Z. D. Bai and J. W. Silverstein. No eigenvalues outside the support of the limiting spectral distribution of large-dimensional sample covariance matrices. *Ann. Probab.*, 26:316–345, 1998.

- [5] Z. D. Bai and J. W. Silverstein. CLT for linear spectral statistics of large-dimensional sample covariance matrices. *Ann. Probab.*, 32(1A):553–605, 2004.
- [6] Z. D. Bai and J. W. Silverstein. *Spectral Analysis of Large Dimensional Random Matrices*, volume 2 of *Mathematics Monograph Series*. Science Press, Beijing, 2006.
- [7] J. Baik, G. Ben Arous, and S. Péché. Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices. *Annals of Probability*, 33(5):1643–1697, 2005.
- [8] J. Baik and J. W. Silverstein. Eigenvalues of large sample covariance matrices of spiked population models. *Journal of Multivariate Analysis*, 97(6):1382–1408, 2006.
- [9] Z. Bao, G. Pan, and W. Zhou. Universality for the largest eigenvalue of sample covariance matrices with general population. *Ann. Statist.*, 43:382–421, 2015.
- [10] Z. G. Bao, G. M. Pan, and W. Zhou. Local density of the spectrum on the edge for sample covariance matrices with general population. *Preprint*, 2013.
- [11] A. Bloemendal, L. Erdős, A. Knowles, H.-T. Yau, and J. Yin. Isotropic local laws for sample covariance and generalized Wigner matrices. *Electron. J. Probab.*, 19(33):1–53, 2014.
- [12] A. Bloemendal, A. Knowles, H.-T. Yau, and J. Yin. On the principal components of sample covariance matrices. *Probability Theory and Related Fields*, 164(1):459–552, 2016.
- [13] T. T. Cai, Z. Ma, and Y. Wu. Sparse PCA: Optimal rates and adaptive estimation. *Ann. Statist.*, 41(6):3074–3110, 2013.
- [14] E. B. Davies. The functional calculus. *J. London Math. Soc. (2)*, 52:166–176, 1995.
- [15] M. Dieng and C. A. Tracy. *Application of Random Matrix Theory to Multivariate Statistics*, pages 443–507. Springer New York, New York, NY, 2011.
- [16] X. Ding and F. Yang. Spiked separable covariance matrices and principal components. *arXiv preprint arXiv:1905.13060*, 2019.
- [17] N. El Karoui. Tracy-Widom limit for the largest eigenvalue of a large class of complex sample covariance matrices. *Ann. Probab.*, 35(2):663–714, 2007.
- [18] L. Erdős, A. Knowles, and H.-T. Yau. Averaging fluctuations in resolvents of random band matrices. *Ann. Henri Poincaré*, 14:1837–1926, 2013.
- [19] L. Erdős, H.-T. Yau, and J. Yin. Universality for generalized Wigner matrices with Bernoulli distribution. *J. of Combinatorics*, 2(1):15–81, 2011.
- [20] F. Götze and A. Tikhomirov. Rate of convergence in probability to the Marchenko-Pastur law. *Bernoulli*, 10(3):503–548, 2004.
- [21] W. Hachem, A. Hardy, and J. Najim. Large complex correlated Wishart matrices: Fluctuations and asymptotic independence at the edges. *Ann. Probab.*, 44(3):2264–2348, 2016.
- [22] Y. He and A. Knowles. Mesoscopic eigenvalue statistics of Wigner matrices. *Ann. Appl. Probab.*, 27(3):1510–1550, 2017.
- [23] I. M. Johnstone. High dimensional statistical inference and random matrices. *arXiv:0611589*.

- [24] I. M. Johnstone. On the distribution of the largest eigenvalue in principal components analysis. *Ann. Statist.*, 29:295–327, 2001.
- [25] I. M. Johnstone. On the distribution of the largest eigenvalue in principal components analysis. *Ann. Statist.*, 29(2):295–327, 2001.
- [26] I. M. Johnstone. Multivariate analysis and Jacobi ensembles: Largest eigenvalue, Tracy-Widom limits and rates of convergence. *Ann. Statist.*, 36(6):2638–2716, 2008.
- [27] A. M. Khorunzhy, B. A. Khoruzhenko, and L. A. Pastur. Asymptotic properties of large random matrices with independent entries. *Journal of Mathematical Physics*, 37(10):5033–5060, 1996.
- [28] A. Knowles and J. Yin. The isotropic semicircle law and deformation of Wigner matrices. *Communications on Pure and Applied Mathematics*, 66(11):1663–1749, 2013.
- [29] A. Knowles and J. Yin. Anisotropic local laws for random matrices. *Probability Theory and Related Fields*, pages 1–96, 2016.
- [30] J. O. Lee and K. Schnelli. Tracy-Widom distribution for the largest eigenvalue of real sample covariance matrices with general population. *Ann. Appl. Probab.*, 26(6):3786–3839, 2016.
- [31] J. O. Lee and J. Yin. A necessary and sufficient condition for edge universality of Wigner matrices. *Duke Math. J.*, 163:117–173, 2014.
- [32] Y. Li, K. Schnelli, and Y. Xu. Central limit theorem for mesoscopic eigenvalue statistics of deformed Wigner matrices and sample covariance matrices. *arXiv preprint arXiv:1909.12821*, 2019.
- [33] A. Lytova and L. Pastur. Central limit theorem for linear eigenvalue statistics of random matrices with independent entries. *Ann. Probab.*, 37(5):1778–1840, 09 2009.
- [34] Z. Ma. Sparse principal component analysis and iterative thresholding. *Ann. Statist.*, 41(2):772–801, 2013.
- [35] V. A. Marčenko and L. A. Pastur. Distribution of eigenvalues for some sets of random matrices. *Mathematics of the USSR-Sbornik*, 1:457, 1967.
- [36] A. Onatski. The Tracy-Widom limit for the largest eigenvalues of singular complex Wishart matrices. *Ann. Appl. Probab.*, 18(2):470–490, 2008.
- [37] A. Onatski. Testing hypotheses about the number of factors in large factor models. *Econometrica*, 77(5):1447–1479, 2009.
- [38] N. Patterson, A. L. Price, and D. Reich. Population structure and eigenanalysis. *PLOS Genetics*, 2(12):1–20, 2006.
- [39] D. Paul. Asymptotics of sample eigenstructure for a large dimensional spiked covariance model. *Statistica Sinica*, 17(4):1617–1642, 2007.
- [40] N. S. Pillai and J. Yin. Universality of covariance matrices. *Ann. Appl. Probab.*, 24:935–1001, 2014.
- [41] J. Silverstein and Z. Bai. On the empirical distribution of eigenvalues of a class of large dimensional random matrices. *Journal of Multivariate Analysis*, 54(2):175 – 192, 1995.
- [42] J. W. Silverstein. On the eigenvectors of large dimensional sample covariance matrices. *Journal of Multivariate Analysis*, 30(1):1 – 16, 1989.

- [43] J. W. Silverstein. Weak convergence of random functions defined by the eigenvectors of sample covariance matrices. *Ann. Probab.*, 18(3):1174–1194, 1990.
- [44] J. W. Silverstein and S. I. Choi. Analysis of the limiting spectral distribution of large dimensional random matrices. *Journal of Multivariate Analysis*, 54(2):295 – 309, 1995.
- [45] H. Xi, F. Yang, and J. Yin. Convergence of eigenvector empirical spectral distribution of sample covariance matrices. *arXiv:1705.03954*.
- [46] N. Xia and Z. Bai. Functional CLT of eigenvectors for large sample covariance matrices. *Statistical Papers*, 56(1):23–60, 2015.
- [47] N. Xia and Z. Bai. Convergence rate of eigenvector empirical spectral distribution of large Wigner matrices. *Statistical Papers*, pages 1–33, 2016.
- [48] N. Xia, Y. Qin, and Z. Bai. Convergence rates of eigenvector empirical spectral distribution of large dimensional sample covariance matrix. *Ann. Statist.*, 41(5):2572–2607, 2013.
- [49] F. Yang. Edge universality of separable covariance matrices. *Electron. J. Probab.*, 24:57 pp., 2019.