

Uncertainty quantification in the stochastic block model with an unknown number of classes

J. van Waaij^{*1} and B.J.K. Kleijn²

¹Department of Statistics, University of Padova

²Korteweg-de Vries Institute for Mathematics, University of Amsterdam

¹jvanwaaij@gmail.com

²b.j.k.kleijn@uva.nl

August 16, 2021

Abstract

We study the frequentist properties of Bayesian statistical inference for the stochastic block model, with an unknown number of classes of varying sizes. We equip the space of vertex labellings with a prior on the number of classes and, conditionally, a prior on the labels. The number of classes may grow to infinity as a function of the number of vertices, depending on the sparsity of the graph. We derive non-asymptotic posterior contraction rates of the form $P_{\theta_{0,n}} \Pi_n(B_n | X^n) \leq \varepsilon_n$, where X^n is the observed graph, generated according to $P_{\theta_{0,n}}$, B_n is either $\{\theta_{0,n}\}$ or, in the very sparse case, a ball around $\theta_{0,n}$ of known extent, and ε_n is an explicit rate of convergence.

These results enable conversion of credible sets to confidence sets. In the sparse case, credible tests are shown to be confidence sets. In the very sparse case, credible sets are enlarged to form confidence sets. Confidence levels are explicit, for each n , as a function of the credible level and the rate of convergence. Hypothesis testing between the number of classes is considered with the help of posterior odds, and is shown to be consistent. Explicit upper bounds on errors of the first and second type and an explicit lower bound on the power of the tests are given.

1 Communities in random graphs

Networks are more present than ever in history. The emergence of the Internet, which is only about 30 years old, is clearly one of the most eye-catching examples, but also

^{*}The first author is supported by the University of Padova under the STARS Grant.

developments in biology giving rise to enormous networks, all waiting for the statistician to be analysed.

Under a network or graph we understand a collection of vertices (also called nodes) and edges between vertices, which can be directed or undirected. Very often data is associated with the vertices and edges. For example in the Facebook network, the vertices are users and edges are formed when two users are friends. With the vertex the user's name, age, etc. are associated, and with the edges for example the date that the connected users became friends could be stored.

There are many things that can be measured from a network, however, the most intrinsic to a graph is its geometric form, emerging from the connections between the vertices. One interesting question that may arise is, are there communities in a graph? Communities are generally understood to be sets of vertices that are more densely connected among each other, than to vertices outside the set. Communities are an indication of shared common properties, for example in the yeast protein-protein interaction network, communities (so-called "functional modules") are "cellular entities that perform certain biological functions, which are relative independent of each other" (Chen and Yuan, 2006).

We use the stochastic block model (SBM), as a mathematical model for communities in a network. The SBM was first considered in Holland, Laskey, and Leinhardt, 1983. In the SBM we observe a graph X^n with n vertices, in which vertices belong to one of a finite number of classes, while edges occur independently, with probabilities that depend on the classes of the vertices they connect.

SBMs have many applications in science and machine learning applications. One interesting example is modelling gene expression (Cline et al., 2007). Another example is the already mentioned yeast protein-protein interaction network studied in Chen and Yuan, 2006. More examples and references of applications are provided in the first section of Abbe, 2018.

There is now a substantial body of literature on the SBM that deals with recovery of communities in a network. Transition phases for the SBM for different number of blocks are studied in Abbe and Sandon, 2015, Abbe and Sandon, 2018, and Zhang and Zhou, 2016. Rates of posterior convergence for SBMs are examined in Mariadassou and Matias, 2015, van der Pas and van der Vaart, 2018, Ghosh, Pati, and Bhattacharya, 2019, and Geng, Bhattacharya, and Pati, 2019. A Bayesian framework for estimating the parameters of a SBM (number of classes, connecting probabilities) is considered in Gao, van der Vaart, and Zhou, 2019. For a comprehensive overview one consults Abbe, 2018.

Recent work mainly considers the SBM under the assumption that the number of blocks

is known, or unknown but bounded. For example Zhang and Zhou, 2016, allows the number of blocks to be as large as $\mathcal{O}(n/\log n)$, but still assumes it to be known. Geng, Bhattacharya, and Pati, 2019 allows the unknown number to be two or three, but state that the general problem (four or more classes) is unsolved.

In the non-Bayesian approach this is less a problem, as one can first estimate the parameters of the SBM (using for instance one of the algorithms mentioned in Abbe, 2018, sec. 7), and then recover the clustering. From a Bayesian perspective this is unnatural (unless one uses an empirical Bayes approach). Here we give a partial answer to this problem. We allow the number of blocks to be unknown and even as large as n^α , for some $0 < \alpha < 1/2$ (in the dense phase, see example 3.4).

This also gives a partial answer to the open problem in Abbe, 2018, sec. 8, to whether exact recovery is possible in classes of sub-linear size, in this case of size $n^{1-\alpha}$. Another question that he raises (p. 4) and tries to find an answer to is:

Are there really communities? Algorithms may output community structures, but are these meaningful or artefacts?

We do this here in several ways. One is by providing confidence sets for our estimators, and another is by providing frequentist guarantees for testing with posterior odds, whether the data comes from a Erdős-Renì graph model (there are no communities), or a SBM (there are communities).

Outline of our results We study posterior rates of convergence for large classes of priors. Posterior rates of convergence are balls B_n of labellings centred around the true labelling $\theta_{0,n}$ of a certain radius r_n (in a metric to be specified) and a sequence (ε_n) such that

$$P_{\theta_{0,n}} \Pi^n(B_n | X_n) \geq 1 - \varepsilon_n.$$

For graphs that are not too sparse, we take $B_n = \{\theta_{0,n}\}$, which leads to exact detection.

This unusual precise description of the posterior convergence rates with ε_n and r_n allows us to construct confidence sets from the credible sets and to derive explicit bounds on the errors for testing with posterior odds.

In the sparse and dense cases, credible sets are shown to be confidence sets with an exact confidence level, which is a function of the credible level and the posterior rate of convergence. In the very sparse case, credible sets are enlarged to convert them into confidence sets.

For symmetric testing with posterior odds, we show that posterior convergence in a parameter set A_n allows us to consistently test between A_n and another disjoint set B_n .

This enables us for instance to test for the number of classes and in particular to test between the SBM and the Erdős-Rényi graph model (basically the SBM with one class).

In the next section we describe our model. In sections 3 and 4 we derive posterior convergence and apply that to some interesting examples. Confidence sets are examined in section 5 and symmetric testing in section 6. Proofs are mostly deferred to appendices B and C. A brief review of the most relevant Bayesian theory can be found in appendix A. Notation and conventions are also explained in this appendix.

2 The stochastic block model

In a SBM for a random graph X^n of order n , each vertex is assigned to a class through an unobserved *class assignment vector* θ'_n . The space in which the random graph X^n takes its values is denoted \mathcal{X}_n (e.g. represented by its (random) adjacency matrix with entries $\{X_{ij} : 1 \leq i < j \leq n\}$). Each vertex belongs to a class and any edge occurs (independently of others) with a probability depending on the classes of the vertices that it connects. We study the *planted multi-section model*, in which the vertices are divided into $\ell \geq 1$ groups of $1 \leq m_{n,1,1} \leq \dots \leq m_{n,\ell,\ell}$ vertices, for a total of $n = m_{n,\ell,1} + \dots + m_{n,\ell,\ell}$ vertices (with labels $1, \dots, \ell$). The corresponding ℓ -vector is denoted by $\mathbf{m}_{n,\ell} = (m_{n,\ell,1}, \dots, m_{n,\ell,\ell})$. We define the set $\Theta'_{\mathbf{m}_{n,\ell}}$ to be the subset of $\{1, \dots, \ell\}^n$ consisting of all class assignment vectors $\theta' = (\theta'_1, \dots, \theta'_n)$, with labels $\theta'_i \in \{1, \dots, \ell\}$, $i = 1, \dots, n$, that have exactly $m_{n,\ell,1}$ elements of one label, and $m_{n,\ell,2}$ elements of another label, *etcetera*.

In the SBM for a graph X^n with class assignment vector $\theta'_n \in \Theta'_n$, the probability of an edge occurring between vertices $1 \leq i, j \leq n, i \neq j$ is denoted $Q_{i,j;n}$ and depends on $n, \theta'_{n,i}$ and $\theta'_{n,j}$ only. If edge probabilities vary (that is, if edge probabilities within different classes vary or if, for example, the edge probability between classes one and two is different from that between three and four) vertices are classified correctly by analysis of their asymptotic degrees (Channaron, Daudin, and Robin, 2012). In the planted multi-section model it is assumed that the probability that an edge occurs, depends only on whether the vertices it connects belong to the same class or to different classes: the probability of an edge *within the same class* is denoted $p_n \in (0, 1)$; the probability of an edge *between classes* is denoted $q_n \in (0, 1)$,

$$Q_{i,j;n}(\theta') = \begin{cases} p_n, & \text{if } \theta'_i = \theta'_j, \\ q_n, & \text{if } \theta'_i \neq \theta'_j. \end{cases} \quad (1)$$

Note that if $p_n = q_n$, X^n is a Erdős-Rényi graph and the class assignment $\theta'_n \in \Theta'_n$ is not identifiable. Indeed asymptotic proximity to the Erdős-Rényi graph (in the sense that the

difference between p_n and q_n decreases too fast to zero as $n \rightarrow \infty$) can make consistent community detection impossible, see Mossel, Neeman, and Sly, 2016 for example in the case of two blocks. We distinguish between the following three cases:

Dense phase: The average degree is proportional to n , so $\liminf_{n \rightarrow \infty} p_n > 0$ and $\liminf_{n \rightarrow \infty} q_n > 0$.

Chernoff-Hellinger phase: The average degrees grows logarithmically. So $p_n = \frac{a_n \log n}{n}$ and $q_n = \frac{b_n \log n}{n}$ for certain bounded positive sequences so that $\liminf_{n \rightarrow \infty} a_n > 0$ and $\liminf_{n \rightarrow \infty} b_n > 0$.

Kesten-Stigum phase: The average degree is bounded from below, so $p_n = c_n/n$ and $q_n = d_n/n$, for bounded sequences of positive numbers so that $\liminf_{n \rightarrow \infty} c_n > 0$ and $\liminf_{n \rightarrow \infty} d_n > 0$.

The ability to detect the correct labelling, depends of course on the differences between p_n, q_n , and a_n, b_n and c_n, d_n , respectively. Exact thresholds for these are studied in Abbe and Sandon, 2015, and Abbe and Sandon, 2018.

Identifiability is also lost due to invariance of the model under permutation of class labels: for example with $\ell = 3$, the class assignment vectors $\theta'_1 = (112233)$, $\theta'_2 = (223311)$ and $\theta'_3 = (331122)$ all give rise to the same distribution for X^6 . This is expressed via an equivalence relation \sim on $\Theta'_{n,\ell}$. Two labels θ'_n, θ''_n are said to be equivalent, if there is a permutation π of the labels $\{1, \dots, \ell\}$ such that for all $1 \leq i \leq n$,

$$\theta'_{1,i} = \theta''_{2,\pi(i)}.$$

We define for an ℓ -vector of class sizes $\mathbf{m}_{n,\ell} = (m_{n,\ell,1}, \dots, m_{n,\ell,\ell})$ satisfying, $1 \leq m_{n,\ell,1} \leq \dots \leq m_{n,\ell,\ell}$ and $\sum_{i=1}^{\ell} m_{n,\ell,i} = n$, the model $\Theta_{\mathbf{m}_{n,\ell}} = \Theta'_{\mathbf{m}_{n,\ell}} / \sim$, which can be identified with the space of all partitions of n elements into ℓ sets of sizes $m_{n,\ell,1}, \dots, m_{n,\ell,\ell}$. One can see that eq. (1) does not depend on the chosen representation θ' of θ , hence we may define

$$Q_{i,j,n}(\theta) = Q_{i,j,n}(\theta'). \quad (2)$$

In our set-up, we have to put some constraints on the $\mathbf{m}_{n,\ell}$. Hence, for given ℓ , we let $M_{n,\ell}$ be the set of all allowed ℓ -vectors $\mathbf{m}_{n,\ell}$. By \mathcal{L}_n we denote all ℓ for which $M_{n,\ell}$ is not empty. We define $M_n = \bigcup_{\ell=1}^n M_{n,\ell}$. We set

$$\Theta_{n,\ell} = \bigcup_{\mathbf{m}_{n,\ell} \in M_{n,\ell}} \Theta_{\mathbf{m}_{n,\ell}}.$$

The full parameter space is denoted

$$\Theta_n = \bigcup_{\mathbf{m}_n \in M_n} \Theta_{\mathbf{m}_n} = \bigcup_{\ell \in \mathcal{L}_n} \Theta_{n,\ell}.$$

For $\ell \in \mathcal{L}_n$, we define

$$m_{n,\ell,\min} = \min_{\mathbf{m}_{n,\ell} \in M_{n,\ell}} m_{n,\ell,1}, \quad \text{and} \quad m_{n,\ell,\max} = \max_{\mathbf{m}_{n,\ell} \in M_{n,\ell}} m_{n,\ell,\ell}.$$

(Recall that $m_{n,\ell,1} = \min_{1 \leq i \leq \ell} m_{n,\ell,i}$ and $m_{n,\ell,\ell} = \max_{1 \leq i \leq \ell} m_{n,\ell,i}$.)

It is noted explicitly that $\ell = 1$ and $\mathbf{m}_{n,1} = (n)$ is also allowed, which allows us to test between the Erdős-Rényi graph model and a SBM with at least two classes. In case $\ell = 1$, $\Theta'_{\mathbf{m}_{n,1}}$ consist of only one element: the n -vector $(1, \dots, 1)$.

In order to distinguish between the number of classes, in our approach, the minimum and maximum sizes of the classes are required to adhere to the following assumption:

ASSUMPTION 2.1 *For all $\ell_1, \ell_2 \in \mathcal{L}_n$, whenever $\ell_1 < \ell_2$, $m_{n,\ell_1,\min} \geq m_{n,\ell_2,\max}$.*

This assumption organises class sizes for decreasing number of classes in consecutive intervals, concentrated around the mean number of elements $m := n/\ell$ per class. For instance, if we impose an upper bound L_n for ℓ given n , then any vector $\mathbf{m}_{n,\ell}$ that satisfies $n/\ell - \frac{1}{4} \frac{n}{L_n^2} \leq m_{n,\ell,i} \leq n/\ell + \frac{1}{4} \frac{n}{L_n^2}$, satisfies assumption 2.1, see eq. (10). A condition like this is quite common in the literature, see for instance Zhang and Zhou, 2016, p. 2254 or the consistency results in Geng, Bhattacharya, and Pati, 2019, assumption (A1), p. 897. The difference $m_{n,\ell_1,\min} - m_{n,\ell_2,\max}$ determines the power to test between Θ_{n,ℓ_1} and Θ_{n,ℓ_2} , see theorem 6.1 in combination with proposition 3.1.

In the planted multi-section model one observes a sequence of graphs X^n of order n and the statistical question is to reconstruct the unobserved class assignment vectors θ_n consistently, that is, correctly with probability growing to one as $n \rightarrow \infty$.

DEFINITION 2.2 *Let $\theta_{0,n} \in \Theta_n$ for all $n \geq 1$ be given. An estimator sequence $\hat{\theta}_n : \mathcal{X}_n \rightarrow \Theta_n$ is said to recover the class assignment $\theta_{0,n}$ exactly if,*

$$P_{\theta_{0,n}}(\hat{\theta}_n(X^n) = \theta_{0,n}) \rightarrow 1,$$

as $n \rightarrow \infty$. That is, $\hat{\theta}_n$ is the true class assignment (up to a permutation) with high probability.

In the very sparse regime where the average degree is large enough, or grows arbitrarily slowly to infinity, exact recovery is not possible, however weaker forms of recovery are still possible, for which we define:

DEFINITION 2.3 *Let $\theta_{0,n} \in \Theta_n$ for all $n \geq 1$ be given. Let $\hat{\theta}_n : \mathcal{X}_n \rightarrow \Theta_n$ be an estimator and s_n be an integer such that there are representations $\theta'_{0,n}$ of $\theta_{0,n}$ and $\hat{\theta}'_n$ of $\hat{\theta}_n$ such that*

$$P_{\theta_{0,n}}\left(\#\left\{i : \hat{\theta}'_{n,i} = \theta'_{0,n,i}\right\} \geq s_n\right) \rightarrow 1, \quad \text{as } n \rightarrow \infty.$$

that is, if $\hat{\theta}_n$ recovers s_n labels correctly, with high probability. When $s_n/n \rightarrow 1$, we say that $\hat{\theta}_n$ recovers the class assignment $\theta_{0,n}$ almost exactly, that is, the estimator recovers

the labelling correct, up to an arbitrarily small fraction and up to a permutation of the labels. When $s_n/n \geq a_n > 1/L_n$, for some deterministic sequence a_n , where L_n is the maximum number of classes in Θ_n , then we say that $\hat{\theta}_n$ recovers the labelling partially. Which means that $\hat{\theta}_n$ performs on average better than a random guess.

Below we specialize to the Bayesian approach in the planted multi-section model: with given $n \geq 1$, $\theta_n \in \Theta_n$ and random X^n , the likelihood is given by,

$$p_{\theta_n}(X^n) = \prod_{1 \leq i < j \leq n} Q_{i,j;n}(\theta_n)^{X_{ij}} (1 - Q_{i,j;n}(\theta_n))^{1-X_{ij}},$$

where $Q_{i,j;n}$ is defined in eq. (2).

We make the following convenient assumption on the prior (which always holds after removing parameters with zero prior mass from the parameter space).

ASSUMPTION 2.4 *The prior mass function π_n of the prior Π_n on Θ_n satisfies $\pi_n(\theta_n) > 0$, for all $\theta_n \in \Theta_n$.*

The posterior distribution of a subset $S_n \subseteq \Theta_n$ is given by

$$\Pi_n(S_n | X^n) = \frac{\sum_{\theta_n \in S_n} \pi_n(\theta_n) p_{\theta_n}(X^n)}{\sum_{\theta_n \in \Theta_n} \pi_n(\theta_n) p_{\theta_n}(X^n)}.$$

PROPOSITION 2.5 *For fixed n , consider an prior probability mass function π_n on Θ_n satisfying assumption 2.4. Suppose that for some $\theta_{0,n} \in \Theta_n$, we observe a graph X^n with n vertices, distributed according to $P_{\theta_{0,n}}$. Let $S_n \subseteq \Theta_n \setminus \{\theta_{0,n}\}$ be non-empty. For $\theta_n \in S_n$, and representations $\theta'_{0,n} \in \theta_{0,n}$ and $\theta'_n \in \theta_n$ define*

$$\begin{aligned} D_{1,n}(\theta_n) &= \{(i, j) \in \{1, \dots, n\}^2 : i < j, \theta'_{0,n,i} = \theta'_{0,n,j}, \theta'_{n,i} \neq \theta'_{n,j}\}, \\ D_{2,n}(\theta_n) &= \{(i, j) \in \{1, \dots, n\}^2 : i < j, \theta'_{0,n,i} \neq \theta'_{0,n,j}, \theta'_{n,i} = \theta'_{n,j}\}. \end{aligned}$$

Then $D_{1,n}(\theta_n)$ and $D_{2,n}(\theta_n)$ are well-defined and disjoint, and if

$$0 < B_n \leq \inf_{\theta_n \in S_n} |D_{1,n}(\theta_n) \cup D_{2,n}(\theta_n)|,$$

then

$$P_{\theta_{0,n}} \Pi_n(S_n | X^n) \leq 2 \left(\frac{\pi_n(S_n)}{\pi_n(\theta_{0,n})} \sqrt{|S_n|} \right) \rho(p_n, q_n)^{B_n}. \quad (3)$$

PROOF Obviously, $D_{1,n}(\theta_n)$ and $D_{2,n}(\theta_n)$ are disjoint and do not depend on the chosen representations. According to lemma A.3 (with $B_n = \{\theta_0\}$), for any tests $\phi_{S_n} : \mathcal{X}_N \rightarrow [0, 1]$, we have,

$$P_{\theta_{0,n}} \Pi_n(S_n | X^n) \leq P_{\theta_{0,n}} \phi_{S_n}(X^n) + \frac{1}{\pi_{\theta_{0,n}}^n} \sum_{\theta_n \in S_n} \pi_n(\theta_n) P_{\theta_n}(1 - \phi_{S_n}(X^n)).$$

Lemma B.1 proves that for any $\theta_n \in S$ there is a test function ϕ_{θ_n} that distinguishes $\theta_{0,n}$ from θ_n as follows,

$$P_{\theta_{0,n}}\phi_{\theta_n}(X^n) + P_{\theta_n}(1 - \phi_{\theta_n}(X^n)) \leq \rho(p_n, q_n)^{|D_{1,n}(\theta_n) \cup D_{2,n}(\theta_n)|} \leq \rho(p_n, q_n)^{B_n},$$

where the last inequality follows from the fact that $\rho(p_n, q_n) \leq 1$ and the assumption $|D_{1,n}(\theta_n) \cup D_{2,n}(\theta_n)| \geq B_n$, for all $\theta_n \in S_n$. Then using test functions $\phi_{S_n}(X^n) = \max\{\phi_{\theta_n}(X^n) : \theta_n \in S_n\}$, we have,

$$P_{\theta_{0,n}}\phi_{S_n}(X^n) \leq \sum_{\theta_n \in S_n} P_{\theta_{0,n}}\phi_{\theta_n}(X^n),$$

so that,

$$\begin{aligned} P_{\theta_{0,n}}\Pi_n(S_n|X^n) &\leq \sum_{\theta_n \in S_n} \left(\frac{\pi_n(\theta_n)}{\pi_n} \vee 1 \right) (P_{\theta_{0,n}}\phi_{\theta_n}(X^n) + P_{\theta_n}(1 - \phi_{\theta_n}(X^n))) \\ &\leq 2 \left(\frac{\pi_n(S_n)}{\pi_n(\theta_{0,n})} \vee |S_n| \right) \rho(p_n, q_n)^{B_n}, \end{aligned}$$

where we use that for a finite index set I and a sequence of non-negative numbers $(x_i : i \in I)$, $\sum_{i \in I} (x_i \vee 1) \leq 2(|I| \vee \sum_{i \in I} x_i)$. \square

3 Selection of the number of classes

Consider the sequence of experiments in which we observe random graphs $X^n \in \mathcal{X}^n$ generated by the SBM of eq. (1). We first provide a general condition under which the posterior selects the true model $\Theta_{n,\ell}$ consistently. Next we apply this to several interesting examples. The Hellinger-affinity between two Bernoulli-distributions with parameters p and q , is given by

$$\rho(p, q) = p^{1/2}q^{1/2} + (1-p)^{1/2}(1-q)^{1/2}.$$

This quantity determines the power of our tests, lemma B.1, similar as in Kleijn and van Waaij, 2018. (See also Zhang and Zhou, 2016, equation 1.2 and theorem 1.1 where minus the log of this quantity (i.e. the Rényi divergence of order 1/2) emerges in the minimax rate of convergence.)

PROPOSITION 3.1 *For fixed n , let the set of models M_n satisfy assumption 2.1 and let Π_n a prior on Θ_n satisfying assumption 2.4. Suppose that for some $\ell_0 \in \mathcal{L}_n$, we observe a graph X^n with n vertices, distributed according to $P_{\theta_{0,n}}$ for some class assignment $\theta_{0,n} \in \Theta_{n,\ell_0}$. For $\ell \in \mathcal{L}_n$ unequal to ℓ_0 , we have,*

$$P_{\theta_{0,n}}\Pi(\Theta_{n,\ell} | X^n) \leq 2 \left(\frac{\pi_n(\Theta_{n,\ell})}{\pi_n(\theta_{0,n})} \vee |\Theta_{n,\ell}| \right) \rho(p_n, q_n)^{\frac{1}{2}n(m_{n,\ell_0 \wedge \ell, \min} - m_{n,\ell_0 \vee \ell, \max})}. \quad (4)$$

PROOF This follows from proposition 2.5 and lemma B.2. \square

The number of elements in $\Theta_{n,\ell}$ is bounded by the number of partitions of $\{1, \dots, n\}$ into ℓ sets, which is known as the Stirling number $V(n, \ell)$ of second kind, which satisfies

$$V(n, \ell) \leq \frac{1}{2} \binom{n}{\ell} \ell^{n-\ell} \leq \frac{1}{2} e^\ell n^\ell \ell^{n-2\ell}. \quad (5)$$

3.1 Examples: convergence to the true model

We consider several examples of posterior contraction in one model Θ_{n,ℓ_0} , whose priors and models are defined below.

EXAMPLE 3.2 (Priors) The first prior is defined hierarchically by taking the uniform prior on \mathcal{L}_n , and conditionally on ℓ , we choose the uniform prior on $\Theta_{n,\ell}$. So

$$\begin{aligned} \ell &\sim \frac{1}{|\mathcal{L}_n|}, \\ \theta_{0,n} | \ell &\sim \frac{1}{|\Theta_{n,\ell}|}. \end{aligned} \quad (6)$$

A second option for a prior is the uniform prior on Θ_n , so

$$\theta_{0,n} \sim \frac{1}{|\Theta_n|}. \quad (7)$$

In both cases, applying proposition 3.1 gives, for $\ell \in \mathcal{L}_n, \ell \neq \ell_0$, the upper bound,

$$P_{\theta_{0,n}} \Pi_n(\Theta_{n,\ell} | X^n) \leq 2 \left(\max_{\ell \in \mathcal{L}_n} |\Theta_{n,\ell}| \right) \rho(p_n, q_n)^{\frac{1}{2}n(m_{n,\ell_0 \wedge \ell, \min} - m_{n,\ell_0 \vee \ell, \max})}. \quad (8)$$

EXAMPLE 3.3 (Models) We take $\mathcal{L}_n = \{1, \dots, L_n\}$, where $2 \leq L_n \leq n$ depends on the degree of sparsity and is specified in the examples. We let $\Theta_{n,\ell}$ be all parameters with ℓ classes and satisfying

$$m_{n,\ell, \min} \geq n/\ell - \frac{1}{4} \frac{n}{L_n^2} \quad \text{and} \quad m_{n,\ell, \max} \leq n/\ell + \frac{1}{4} \frac{n}{L_n^2}. \quad (9)$$

One verifies easily that assumption 2.1 is satisfied and that for $\ell, \ell_0 \in \mathcal{L}_n, \ell \neq \ell_0$,

$$m_{n,\ell_0 \wedge \ell, \min} - m_{n,\ell_0 \vee \ell, \max} \geq \frac{1}{2} \frac{n}{L_n^2}. \quad (10)$$

Using eq. (5), the number of elements of $\Theta_{n,\ell}$ is upper bounded by $\frac{1}{2} e^{2n \log L_n}$, which is trivial for $\ell = 1$, and when $\ell \geq 2$ and $n \geq 4$, we have,

$$\frac{1}{2} e^\ell n^\ell \ell^{n-2\ell} \leq \frac{1}{2} n^\ell \ell^n \leq \frac{1}{2} \ell^{2n} \leq \frac{1}{2} e^{2n \log L_n}, \quad (11)$$

using that for integers $2 \leq x \leq y$ with either $x \geq 3$ or $y \geq 4$, $y^x \leq x^y$ and using that $\ell \leq L_n \leq n$. It follows that,

$$P_{\theta_{0,n}} \Pi_n(\Theta_{n,\ell} | X^n) \leq e^{2n \log L_n} \rho(p_n, q_n)^{\frac{1}{4}n^2/(L_n^2)} \quad (12)$$

EXAMPLE 3.4 (*Dense phase*) Define $b_n = -\log \rho(p_n, q_n) > 0$. Let $L_n = \mathbf{o}(\sqrt{n/\log n})$, for example $L_n = \lfloor n^\alpha \rfloor$, for some $\alpha < 1/2$. When $\frac{nb_n}{L_n^2 \log L_n} \geq 12$, $P_{\theta_{0,n}} \Pi(\Theta_{n,\ell} | X^n) \leq e^{-b_n n^2/(12L_n^2)}$. It follows that $P_{\theta_{0,n}} \Pi(\Theta_{n,\ell_0}^c | X^n) \leq L_n e^{-b_n n^2/(12L_n^2)} \leq L_n^{1-n} \rightarrow 0$, as $n \rightarrow \infty$. \square

In the sparse regimes another approximation strategy is needed.

EXAMPLE 3.5 (*Chernoff-Hellinger phase*) Recall that in the Chernoff-Hellinger phase, we assume that the edge probabilities satisfy $np_n = a_n \log(n)$ and $nq_n = b_n \log(n)$ for sequences a_n, b_n bounded away from zero. Suppose L_n satisfies

$$48L_n^2 \log L_n \leq (\sqrt{a_n} - \sqrt{b_n})^2 \log n. \quad (13)$$

Equation (13) is satisfied for sufficiently large n , when $L_n \leq (\log n)^a$ with $a < 1/2$.

Using lemma C.2, we find $\sqrt{1-p}\sqrt{1-q} \leq 1 - \frac{1}{2}(p+q) + \frac{1}{4}pq$, and using the fact that $(1+x/r)^r \leq e^x$ (see lemma C.3), for all positive integers r and $x > -r$, we obtain

$$\begin{aligned} & \rho(p_n, q_n)^{\frac{1}{2}n|m_{n,\ell_0 \wedge \ell, \min} - m_{n,\ell_0 \vee \ell, \max}|} \\ & \leq \left[1 - \frac{1}{2}(\sqrt{p_n} - \sqrt{q_n})^2 + p_n q_n / 4 \right]^{\frac{1}{2}n|m_{n,\ell_0 \wedge \ell, \min} - m_{n,\ell_0 \vee \ell, \max}|} \\ & = \left[1 - \frac{\log n}{n} \left\{ \frac{1}{2}(\sqrt{a_n} - \sqrt{b_n})^2 - \frac{a_n b_n \log n}{4n} \right\} \right]^{\frac{1}{2}n(m_{n,\ell_0 \wedge \ell, \min} - m_{n,\ell_0 \vee \ell, \max})} \\ & \leq \exp \left(-\frac{1}{2}(m_{n,\ell_0 \wedge \ell, \min} - m_{n,\ell_0 \vee \ell, \max}) \left\{ \frac{1}{2}(\sqrt{a_n} - \sqrt{b_n})^2 - \frac{a_n b_n \log n}{4n} \right\} \log n \right) \\ & \leq \exp \left(-\frac{1}{16}(\sqrt{a_n} - \sqrt{b_n})^2 \frac{n \log n}{L_n^2} \right), \end{aligned}$$

for n sufficiently large. It follows from eq. (12) that

$$\begin{aligned} P_{\theta_{0,n}} \Pi(\Theta_{n,\ell} | X^n) & \leq \exp \left(-n \left(\frac{1}{16}(\sqrt{a_n} - \sqrt{b_n})^2 \frac{\log n}{L_n^2} - 2 \log L_n \right) \right) \\ & \leq \exp \left(-(\sqrt{a_n} - \sqrt{b_n})^2 \frac{n \log n}{48L_n^2} \right), \end{aligned}$$

using eq. (13) in the last inequality. It follows that

$$P_{\theta_{0,n}} \Pi(\Theta_{\mathbf{m}_{n,\ell_0}}^c | X^n) \leq L_n \exp \left(-(\sqrt{a_n} - \sqrt{b_n})^2 \frac{n \log n}{48L_n^2} \right) \leq L_n^{1-n} \rightarrow 0,$$

as $n \rightarrow \infty$. \square

Even in the Kesten-Stigum phase, there is testing power enough to decide between the classes.

EXAMPLE 3.6 (*Kesten-Stigum phase*) Recall that in the Kesten-Stigum phase, $np_n = c_n$ and $nq_n = d_n$ for sequences of positive numbers c_n and d_n bounded away from zero. We obtain in similar manner as in example 3.5 that,

$$(c_n^{1/2} - d_n^{1/2})^2 \geq 48L_n^2 \log L_n,$$

guarantees that,

$$P_{\theta_{0,n}} \Pi(\Theta_{n,\ell_0}^c | X^n) \leq L_n \exp\left(-\left(\sqrt{c_n} - \sqrt{d_n}\right)^2 \frac{n}{48L_n^2}\right) \leq L_n^{1-n} \rightarrow 0,$$

as $n \rightarrow \infty$. □

4 Posterior concentration at the parameter

Posterior convergence at the true parameter $\theta_{0,n} \in \Theta_{n,\ell_0}$, where ℓ_0 is the true number of classes, is derived in the following way. Noting that

$$P_{\theta_0} \Pi(\{\theta_{n,0}\}^c | X^n) = P_{\theta_0} \Pi(\Theta_{n,\ell_0}^c | X^n) + P_{\theta_{n,0}} \Pi(\Theta_{n,\ell_0} \setminus \{\theta_{0,n}\} | X^n),$$

posterior consistency is established once both expectations on the right converge to zero as $n \rightarrow \infty$. The first was the content of the previous section, the second we treat here. Obviously, when $\ell_0 = 1$, then $\Theta_{n,\ell_0} = \{\theta_{n,0}\}$ and convergence at the true parameter follows from section 3. Hence, we assume that $\ell_0 > 1$. In the previous section posterior convergence in a set Θ_{n,ℓ_0} was established by showing that there was enough testing power between Θ_{n,ℓ_0} and $\Theta_{n,\ell}$, for $\ell \neq \ell_0$. Here we do something similar within Θ_{n,ℓ_0} , by constructing tests between $\theta_{0,n}$ (in the dense and sparse case) and rings $V_{n,k} \subseteq \Theta_{n,\ell_0}$ of radius k around $\theta_{0,n}$ in the distance r_n .

Define the distance r_n on Θ_n as follows: first define r'_n on Θ'_n by

$$r'_n(\theta', \eta') = \max_{a \neq b} \# \{i : \theta'_i = a, \eta'_i = b\}.$$

This does depend on the particular choice of the representation. So we define for $\theta, \eta \in \Theta_n$,

$$r_n(\theta, \eta) = \min_{\theta' \in \theta, \eta' \in \eta} r'_n(\theta', \eta'). \quad (14)$$

The function r_n takes values in $\{0, \dots, \lfloor n/2 \rfloor\}$ and according to lemma B.5, when $\theta, \eta \in \Theta_{n,\ell}$, $r_n(\theta, \eta) \leq m_{n,\ell,\max}/2$. The function r_n is symmetric and $r_n(\theta, \eta) = 0$ if and only if $\theta = \eta$.

For a given $\ell_0 \in \mathcal{L}_n$ and $\theta_{0,n} \in \Theta_{n,\ell_0}$ define subsets of Θ_{n,ℓ_0} , by

$$V_{n,\ell_0,k} = \{\theta_n \in \Theta_{n,\ell_0} : r_n(\theta_n, \theta_{0,n}) = k\}.$$

These sets resemble the sets $V_{n,k}$ in Kleijn and van Waaij, 2018 and are rings of radius k around $\theta_{0,n}$ in Θ_{n,ℓ_0} . When $k \leq m_{n,\ell_0,\min}$, lemma B.6 bounds the number of elements in $V_{n,\ell_0,k}$ by $2^{k\ell_0(\ell_0-1)} \binom{n(\ell_0-1)}{\ell_0(\ell_0-1)k}$, which in turn is bounded by $\left(\frac{2en}{\ell_0 k}\right)^{\ell_0(\ell_0-1)k}$.

PROPOSITION 4.1 Consider a prior on Θ_n satisfying assumption 2.4. Suppose that for some $\ell_0 \in \mathcal{L}_n$, $\ell_0 > 1$, we observe a graph X^n with n vertices, distributed according to $P_{\theta_{0,n}}$, where $\theta_{0,n} \in \Theta_{n,\ell_0}$. For all $k \in \{0, \dots, \lfloor m_{n,\ell_0,\max}/2 \rfloor\}$, we have,

$$P_{\theta_{0,n}} \Pi(V_{n,\ell_0,k} \mid X^n) \leq 2 \left(|V_{n,\ell_0,k}| \vee \frac{\pi_n(V_{n,\ell_0,k})}{\pi_n(\theta_{0,n})} \right) \rho(p_n, q_n)^{2k(m_{n,\ell_0,\min} - k)^+}. \quad (15)$$

PROOF This follows from proposition 2.5 and the lower bound on $|D_{1,n}(\theta_n) \cup D_{2,n}(\theta_n)|$ in lemma B.4. \square

Define

$$W_{n,\ell_0,k_n} = \bigcup_{k=k_n}^{\lfloor m_{n,\ell_0,\max}/2 \rfloor} V_{n,\ell_0,k},$$

to be the set of all elements with r_n distance at least k_n from $\theta_{0,n}$. Taking $k_n = 1$, $W_{n,\ell_0,1} = \Theta_{n,\ell_0} \setminus \{\theta_{0,n}\}$.

Applying proposition 4.1 to W_{n,ℓ_0,k_n} gives

$$P_{\theta_{0,n}} \Pi(W_{n,k_n} \mid X^n) \leq \sum_{k=k_n}^{\lfloor m_{n,\ell_0,\max}/2 \rfloor} 2 \left(|V_{n,\ell_0,k}| \vee \frac{\pi_n(V_{n,\ell_0,k})}{\pi_n(\theta_{0,n})} \right) \rho(p_n, q_n)^{2k(m_{n,\ell_0,\min} - k)^+}$$

For the next corollary we need the following assumption

ASSUMPTION 4.2 For all $\ell \in \mathcal{L}_n$, $m_{n,\ell,\min} \geq m_{n,\ell,\max}/2$.

COROLLARY 4.3 Assume assumption 4.2. Suppose that for some $\ell_0 \in \mathcal{L}_n$, $\ell_0 > 1$, and $\theta_{0,n} \in \Theta_{n,\ell_0}$, we observe a graph X^n with n vertices, distributed according to $P_{\theta_{0,n}}$. Furthermore, assume that the prior π_n gives positive mass to every element of Θ_{n,ℓ_0} and satisfies

$$\max_{\theta_n, \eta_n \in \Theta_{n,\ell_0}} \frac{\pi_n(\theta_n)}{\pi_n(\eta_n)} \leq K_{n,\ell_0}.$$

Define

$$B_n = 2n \rho(p_n, q_n)^{\frac{2m_{n,\ell_0,\min} - m_{n,\ell_0,\max}}{\ell_0(\ell_0 - 1)}},$$

then

$$P_{\theta_{n,0}} \Pi_n(\Theta_{n,\ell_0} \setminus \{\theta_{n,0}\} \mid X^n) \leq 2K_{n,\ell_0} B_n^{\ell_0(\ell_0 - 1)} e^{(\ell_0 - 1)B_n}. \quad (16)$$

PROOF Take $k_n \equiv 1$ for all n in proposition 4.1. Using proposition 4.1, the upper bound on the number of elements in $V_{n,\ell,k}$ eq. (27) and the condition on the prior mass function

π_n we have the bound

$$\begin{aligned}
& P_{\theta_{n,0}} \Pi(\Theta_{n,\ell_0} \setminus \{\theta_{n,0}\} \mid X^n) \\
& \leq 2K_{n,\ell_0} \sum_{k=1}^{\lfloor m_{n,\ell_0,\max}/2 \rfloor} 2^{k\ell_0(\ell_0-1)} \binom{n(\ell_0-1)}{\ell_0(\ell_0-1)k} \rho(p_n, q_n)^{2k(m_{n,\ell_0,\min}-k)} \\
& \leq 2K_{n,\ell_0} \sum_{k=1}^{\lfloor m_{n,\ell_0,\max}/2 \rfloor} \frac{(2n(\ell_0-1))^{\ell_0(\ell_0-1)k}}{(\ell_0(\ell_0-1)k)!} \rho(p_n, q_n)^{2k(m_{n,\ell_0,\min}-m_{n,\ell_0,\max}/2)} \\
& \leq 2K_{n,\ell_0} \sum_{k=\ell_0(\ell_0-1)}^{\infty} \frac{(2n(\ell_0-1))^k}{k!} \rho(p_n, q_n)^{k \frac{2m_{n,\ell_0,\min}-m_{n,\ell_0,\max}}{\ell_0(\ell_0-1)}} \\
& \leq 2K_{n,\ell_0} \frac{(2n(\ell_0-1))^{\ell_0(\ell_0-1)}}{(\ell_0(\ell_0-1))!} \rho(p_n, q_n)^{2m_{n,\ell_0,\min}-m_{n,\ell_0,\max}} \\
& \quad \times \sum_{k=0}^{\infty} \frac{(2n(\ell_0-1))^k}{k!} \rho(p_n, q_n)^{k \frac{2m_{n,\ell_0,\min}-m_{n,\ell_0,\max}}{\ell_0(\ell_0-1)}} \\
& = 2K_{n,\ell_0} \frac{1}{(\ell_0(\ell_0-1))!} \left((2n(\ell_0-1)) \rho(p_n, q_n)^{\frac{2m_{n,\ell_0,\min}-m_{n,\ell_0,\max}}{\ell_0(\ell_0-1)}} \right)^{\ell_0(\ell_0-1)} \\
& \quad \times \exp \left(2n(\ell_0-1) \rho(p_n, q_n)^{\frac{2m_{n,\ell_0,\min}-m_{n,\ell_0,\max}}{\ell_0(\ell_0-1)}} \right) \\
& = \frac{2K_{n,\ell_0}}{(\ell_0(\ell_0-1))!} ((\ell_0-1)B_n)^{\ell_0(\ell_0-1)} e^{(\ell_0-1)B_n}.
\end{aligned}$$

With the lower bound $k! \geq \sqrt{2\pi}k^{k+1/2}e^{-k}$ one can easily see that for $\ell_0 \geq 3$, $\frac{(\ell_0-1)^{\ell_0(\ell_0-1)}}{(\ell_0(\ell_0-1))!} \leq 1$, but it also holds for $\ell_0 = 2$. Hence $P_{\theta_{n,0}} \Pi(\Theta_{n,\ell_0} \setminus \{\theta_{n,0}\} \mid X^n) \leq 2K_{n,\ell_0} B_n^{\ell_0(\ell_0-1)} e^{(\ell_0-1)B_n}$. \square

4.1 Examples: recovery of the parameters

In this section we are interested in exact and almost exact recovery for some interesting examples. We consider the same priors and models as in examples 3.2 and 3.3. Note that in this case

$$m_{n,\ell,\min} - m_{n,\ell,\max}/2 \geq \frac{n}{8\ell}, \quad (17)$$

so assumption 4.2 is satisfied for all $\ell \in \mathcal{L}_n$.

EXAMPLE 4.4 (*Dense phase, continuation of example 3.4*)

Recall that $b_n = -\log \rho(p_n, q_n)$. The quantity B_n in corollary 4.3 is bounded as follows

$$B_n = 2n \rho(p_n, q_n)^{\frac{2m_{n,\ell_0,\min}-m_{n,\ell_0,\max}}{\ell_0(\ell_0-1)}} \leq 2n \rho(p_n, q_n)^{\frac{n}{4\ell_0^2(\ell_0-1)}} = e^{-\frac{nb_n}{4\ell_0^2(\ell_0-1)} + \log(2n)}.$$

So with corollary 4.3 and the result of example 3.4, when $L_n = \mathfrak{o}(\sqrt{n/\log n})$ and

$$nb_n \geq 8L_n^2(L_n - 1) \log(2n),$$

we obtain

$$P_{\theta_{0,n}} \Pi_n(\theta_0 | X^n) \geq 1 - L_n e^{-b_n n^2 / (12L_n^2)} - 2\sqrt{e} e^{-nb_n / (8L_n)}.$$

In particular $P_{\theta_{0,n}} \Pi_n(\theta_0 | X^n) \rightarrow 1$, as $n \rightarrow \infty$. \square

So in the dense phase we have exponentially fast convergence of the expected posterior mass of $\theta_{0,n}$ to one, under the true distribution. In the sparse phases we get only a polynomial rate of convergence.

EXAMPLE 4.5 (*Chernoff-Hellinger phase, continuation of example 3.5*) With a similar calculation as in example 3.5, we find,

$$\begin{aligned} & \rho(p_n, q_n)^{\frac{2m_{n,\ell_0,\min} - m_{n,\ell_0,\max}}{\ell_0(\ell_0-1)}} \\ &= \left[1 - \frac{\log n}{n} \left(\frac{1}{2} (\sqrt{a_n} - \sqrt{b_n})^2 - \frac{a_n b_n \log n}{4n} \right) \right]^{n \frac{2m_{n,\ell_0,\min} - m_{n,\ell_0,\max}}{\ell_0(\ell_0-1)n}} \\ &\leq \exp \left(-\frac{2m_{n,\ell_0,\min} - m_{n,\ell_0,\max}}{\ell_0(\ell_0-1)n} \left(\frac{1}{2} (\sqrt{a_n} - \sqrt{b_n})^2 - \frac{a_n b_n \log n}{4n} \right) \log n \right). \end{aligned}$$

It follows that

$$B_n \leq e^{\log(2n) - \frac{(a_n^{1/2} - b_n^{1/2})^2}{16\ell_0^2(\ell_0-1)} \log n}.$$

So when $(a_n^{1/2} - b_n^{1/2})^2 \geq 32L_n^2(L_n - 1) \frac{\log(2n)}{\log n}$,

$$P_{\theta_{n,0}} \Pi_n(\Theta_{n,\ell_0} \setminus \{\theta_{n,0}\} | X^n) \leq 2\sqrt{e} n^{-(a_n^{1/2} - b_n^{1/2})^2 / (32L_n)}.$$

It follows that when $(a_n^{1/2} - b_n^{1/2})^2 \geq 48L_n^2(L_n - 1)$, then with the result of example 3.5

$$P_{\theta_{0,n}} \Pi_n(\theta_0 | X^n) \geq 1 - L_n \exp \left(-n \frac{1}{48} (\sqrt{a_n} - \sqrt{b_n})^2 \frac{\log n}{L_n^2} \right) - 2\sqrt{e} n^{-(a_n^{1/2} - b_n^{1/2})^2 / (32L_n)}.$$

In particular $P_{\theta_{0,n}} \Pi_n(\theta_{0,n} | X^n) \rightarrow 1$, as $n \rightarrow \infty$. \square

In the even sparser Kesten-Stigum phase exact recovery is not possible anymore (Abbe and Sandon, 2015; Abbe and Sandon, 2018). Instead we obtain weak recovery, in which all but a small fraction of the labels is recovered.

EXAMPLE 4.6 (*Kesten-Stigum phase, continuation of example 3.6*) With a calculation similar to that of example 3.5, we find that,

$$\rho(p_n, q_n)^{2k(m_{n,\ell_0,\min} - k)} \leq e^{-k(c_n^{1/2} - a_n^{1/2})^2 / (16\ell_0)}.$$

Using the bound $2^{\ell_0(\ell_0-1)k} \binom{n(\ell_0-1)}{\ell_0(\ell_0-1)k} \leq \left(\frac{2en}{\ell_0 k}\right)^{\ell_0(\ell_0-1)k}$, we see that the posterior

$$\begin{aligned} P_{\theta_{0,n}} \Pi_n(W_{n,\ell_0,k_n} | X^n) &\leq 2 \sum_{k=k_n}^{\lfloor m_{n,\ell_0,\max}/2 \rfloor} \left(\frac{2en}{\ell_0 k} e^{-(c_n^{1/2}-d_n^{1/2})^2/(16\ell_0^2(\ell_0-1))}\right)^{\ell_0(\ell_0-1)k} \\ &\leq 2 \frac{\left(\frac{2en}{\ell_0 k_n} e^{-(c_n^{1/2}-d_n^{1/2})^2/(16\ell_0^2(\ell_0-1))}\right)^{\ell_0(\ell_0-1)k_n}}{1 - \left(\frac{2en}{\ell_0 k_n} e^{-(c_n^{1/2}-d_n^{1/2})^2/(16\ell_0^2(\ell_0-1))}\right)^{\ell_0(\ell_0-1)}}. \end{aligned}$$

Let $(\delta_n)_{n \geq 1}$ a sequence in $(0, 1)$, such that $\delta_n n$ is an integer and δ_n converges to a number $\delta \in [0, 1)$. For $k_n = \delta_n n$, we see

$$P_{\theta_{0,n}} \Pi_n(W_{n,\ell_0,k_n} | X^n) \leq 2 \frac{\left(2e\delta_n^{-1}\ell_0^{-1} e^{-(c_n^{1/2}-d_n^{1/2})^2/(16\ell_0^2(\ell_0-1))}\right)^{\delta_n \ell_0(\ell_0-1)n}}{1 - \left(2e\delta_n^{-1}\ell_0^{-1} e^{-(c_n^{1/2}-d_n^{1/2})^2/(16\ell_0^2(\ell_0-1))}\right)^{\ell_0(\ell_0-1)}}.$$

It follows from lemma C.1 that when $\delta_n n \geq 2$, and

$$\ell_0(\ell_0-1) \left[-1 - \log 2 + \log \delta_n + \log \ell_0 + (c_n^{1/2} - d_n^{1/2})^2/(16\ell_0^2(\ell_0-1))\right] \geq \sqrt{\frac{2}{\delta_n n}},$$

$$P_{\theta_{0,n}} \Pi_n(W_{n,\ell_0,k_n} | X^n) \leq 2e^{-\delta_n \ell_0(\ell_0-1) \left[-1 - \log 2 + \log \delta_n + \log \ell_0 + (c_n^{1/2} - d_n^{1/2})^2/(16\ell_0^2(\ell_0-1))\right] n/4}.$$

We define the Hamming metric m_n , closely related to r_n , on the parameter space, again stepwise via a metric m'_n on Θ'_n . Define m'_n on Θ'_n , by

$$m'_n(\theta', \eta') = \sum_{i=1}^n \mathbb{I}_{\theta'_i \neq \eta'_i},$$

and next we define

$$m_n(\theta, \eta) = \min_{\theta' \in \Theta, \eta' \in \Theta} m'_n(\theta', \eta') \quad (18)$$

on Θ_n . The quantity $m_n(\hat{\theta}_n, \theta_{0,n})$ counts the number of misspecified labels of an estimator $\hat{\theta}_n$. Let

$$\bar{B}_{n,k_n}(\theta_n) = \{\eta_n \in \Theta_{n,\ell_0} : r_n(\theta_n, \eta_n) \leq k_n\}$$

and

$$B_{n,k_n}(\theta_n) = \{\eta_n \in \Theta_n : m_n(\theta_n, \eta_n) \leq k_n\}. \quad (19)$$

It follows from lemma B.7 that

$$\bar{B}_{n,k_n/(\ell_0(\ell_0-1))}(\theta_n) \subseteq B_{n,k_n}(\theta_n) \quad \text{and clearly} \quad W_{n,\ell_0,k_n} = \Theta_{n,\ell_0} \setminus \bar{B}_{n,k_n-1}(\theta_{0,n}). \quad (20)$$

Let now δ_n be a sequence of positive numbers such that $\delta_n n / [\ell_0(\ell_0 - 1)]$ are integers, $\delta_n / [\ell_0(\ell_0 - 1)] \in (0, 1)$ and decreases to a number δ in $[0, 1)$, then

$$\begin{aligned} & P_{\theta_{0,n}} \Pi_n(\Theta_{m_n, \ell_0} \setminus B_{n, \delta_n} \mid X^n) \\ & \leq P_{\theta_{0,n}} \Pi_n(\Theta_{m_n, \ell_0} \setminus \bar{B}_{n, \delta_n n / [\ell_0(\ell_0 - 1)] - 1} \mid X^n) \\ & = P_{\theta_{0,n}} \Pi_n(W_{n, \ell_0, \delta_n n / [\ell_0(\ell_0 - 1)]} \mid X^n) \\ & \leq 2e^{-\delta_n [-1 - \log 2 + \log \delta_n - \log(\ell_0 - 1) + (c_n^{1/2} - d_n^{1/2})^2 / (16\ell_0^2(\ell_0 - 1))]} n^{1/4}. \end{aligned}$$

When $\delta_n \rightarrow 0$, $-\log \delta_n \rightarrow \infty$, so with the result of example 3.6 almost exact convergence is achieved once $(c_n^{1/2} - d_n^{1/2})^2 \rightarrow \infty$ (however slowly). To be exact, for $(c_n^{1/2} - d_n^{1/2})^2 \geq 48L^2 \log L$, $\frac{\delta_n n}{L(L-1)} \geq 2$, and

$$-1 - \log 2 + \log \delta_n - \log(L-1) + (c_n^{1/2} - d_n^{1/2})^2 / (16L^2(L-1)) \geq \sqrt{\frac{2L(L-1)}{\delta_n n}}, \quad (21)$$

$$\begin{aligned} P_{\theta_{0,n}} \Pi_n(B_{n, \delta_n} \mid X^n) & \leq 1 - L_n \exp\left(-n \frac{1}{48L^2} (\sqrt{c_n} - \sqrt{d_n})^2\right) \\ & \quad - 2e^{-\delta_n (-1 - \log 2 + \log \delta_n - \log(L-1) + (c_n^{1/2} - d_n^{1/2})^2 / (16L^2(L-1)))} n^{1/4}. \end{aligned}$$

Note that condition eq. (21) is in particular satisfied when

$$\delta_n := 2(L-1)e^{2 - (c_n^{1/2} - d_n^{1/2})^2 / (16L^2(L-1))}$$

and $(c_n^{1/2} - d_n^{1/2})^2$ converges to infinity. Then $\delta_n \rightarrow 0$ and

$$P_{\theta_{0,n}} \Pi_n(B_{n, \delta_n} \mid X^n) \geq 1 - L_n \exp\left(-n \frac{1}{48L^2} (\sqrt{c_n} - \sqrt{d_n})^2\right) - 2e^{-\delta_n n^{1/4}}.$$

For fixed $\delta_0 < 1/L$, setting $(c_n^{1/2} - d_n^{1/2})^2$ so that $\delta_0 = 2(L-1)e^{2 - (c_n^{1/2} - d_n^{1/2})^2 / (16L^2(L-1))}$, and $\delta_n = \left\lfloor 2(L-1)n e^{2 - (c_n^{1/2} - d_n^{1/2})^2 / (16L^2(L-1))} \right\rfloor / n$ gives concentration in a ball of radius $\delta_0 n$. \square

5 Coverage of credible sets with examples

Conditionally on an observation X^n , a credible set of credible level $1 - \alpha_n$ is a measurable subset $D_n(X^n)$ of the parameter set with posterior mass at least $1 - \alpha_n$:

$$\Pi_n(D_n(X^n) \mid X^n) \geq 1 - \alpha_n.$$

In our (discrete, finite) setting any set-valued map $x^n \mapsto B_n(x^n) \subseteq \Theta_n$, the corresponding map $x^n \mapsto \Pi_n(B_n(x^n) \mid x^n)$ is measurable and positive, and hence the integral

$P_{\theta_0}\Pi_n(B_n(X^n) | X^n)$ is well-defined, see appendix A for details. From this perspective, a credible set (of confidence level $1 - \alpha_n$) is a set-valued map $x^n \mapsto D_n(x^n)$ satisfying $\Pi_n(D_n(x^n) | x^n) \geq 1 - \alpha_n$, for every $x^n \in \mathcal{X}_n$. In nonparametric setting, credible sets can have bad coverage: Freedman, 1999 provides us with examples. However in this section we show that in the case of exact recovery credible sets cover $\theta_{0,n}$ with high probability. In case of almost exact recovery we make the credible sets larger in order to guarantee asymptotic coverage, using ideas of Kleijn, 2020 and Kleijn and van Waaij, 2018.

LEMMA 5.1 *Let $n \geq 1$. Let $x^n \mapsto B_n(x^n) \subset \Theta_n$ be a set valued map, such that $P_{\theta_0}\Pi_n(B_n(X^n) | X^n) \geq 1 - a_n$, with $0 < a_n < 1$. Then, for every $0 < r_n < 1$,*

$$P_{\theta_0}[\Pi_n(B_n(X^n) | X^n) \geq 1 - r_n] \geq 1 - \frac{1}{r_n}a_n.$$

PROOF Let $E_n = \{\omega : \Pi_n(B_n | X^n(\omega)) \geq 1 - r_n\}$ be the event that the posterior mass of $B_n(X^n)$ is at least $1 - r_n$. Let $\delta > 0$. Suppose that $P_{\theta_0}(E_n) \leq 1 - \frac{1}{r_n}a_n - \delta$. Then

$$\begin{aligned} P_{\theta_0}\Pi_n(B_n(X^n) | X^n) &\leq P_{\theta_0}(E_n) + (1 - r_n)P_{\theta_0}(E_n^c) = P_{\theta_0}(E_n) + (1 - r_n)(1 - P_{\theta_0}(E_n)) \\ &= r_n P_{\theta_0}(E_n) + 1 - r_n \leq r_n \left(1 - \frac{1}{r_n}a_n - \delta\right) + 1 - r_n = 1 - a_n - \delta r_n < 1 - a_n, \end{aligned}$$

which contradicts with our assumption that $P_{\theta_0}\Pi_n(B_n(X^n) | X^n) \geq 1 - a_n$. Hence $P_{\theta_0}(E_n) > 1 - \frac{1}{r_n}a_n - \delta$. As this holds for every $\delta > 0$, it follows that $P_{\theta_0}(E_n) \geq 1 - \frac{1}{r_n}a_n$. \square

LEMMA 5.2 *Suppose $P_{\theta_0}\Pi_n(\{\theta_{0,n}\} | X^n) \geq 1 - x_n$, where $0 < x_n < 1$. Let $\alpha_n \in (0, 1)$ and $D_n(X^n)$ a $1 - \alpha_n$ credible set, i.e. $\Pi_n(D_n(X^n) | X^n) \geq 1 - \alpha_n$. Then*

$$P_{\theta_{0,n}}(\theta_{0,n} \in D_n(X^n)) \geq 1 - \frac{1}{1 - \alpha_n}x_n.$$

PROOF Let $E_n = \{\Pi(\{\theta_0\} | X^n) \geq r\}$ be the event that $\{\theta_0\}$ has posterior mass at least r , $r > \alpha_n$. It follows from lemma 5.1 that $P_{\theta_{0,n}}(E_n) \geq 1 - \frac{1}{1-r}x_n$. As $D_n(X^n)$ has at least $1 - \alpha_n$ posterior mass, $D_n(X^n)$ and $\{\theta_0\}$ cannot be disjoint on the event E_n , as $1 - \alpha_n + r > 1$. In other words, $\theta_0 \in D_n(X^n)$ on E_n . So $P_{\theta_{0,n}}(\theta_0 \in D_n(X^n)) \geq P_{\theta_{0,n}}(E_n) \geq 1 - \frac{1}{1-r}x_n$. As this holds for any $r > \alpha_n$ we have $P_{\theta_{0,n}}(\theta_0 \in D_n(X^n)) \geq 1 - \frac{1}{1-\alpha_n}x_n$. \square

EXAMPLE 5.3 (*Dense phase, continuation of examples 3.4 and 4.4*) Let $D_n(X^n)$ be a $1 - \alpha_n$ credible set, $0 < \alpha_n < 1$. With example 4.4 and lemma 5.2 we get

$$P_{\theta_{0,n}}(\theta_{0,n} \in D_n(X^n)) \geq 1 - \frac{1}{1 - \alpha_n} \left(L_n e^{-b_n n^2 / (12L_n^2)} + 2\sqrt{e} e^{-nb_n / (8L_n)} \right).$$

\square

EXAMPLE 5.4 (*Chernoff-Hellinger phase, continuation of examples 3.5 and 4.5*) Let $D_n(X^n)$ be a $1 - \alpha_n$ credible set, $0 < \alpha_n < 1$. With example 4.5 and lemma 5.2 we get

$$P_{\theta_0}(\theta_{0,n} \in D_n(X^n)) \geq 1 - \frac{1}{1 - \alpha_n} \left[L_n \exp \left(-n \frac{1}{48} (\sqrt{a_n} + \sqrt{b_n})^2 \frac{\log n}{L_n^2} \right) + 2\sqrt{e} n^{-(a_n^{1/2} - b_n^{1/2})^2 / (32L_n)} \right].$$

□

5.1 Enlarged credible sets

In the case of almost exact convergence, credible sets need to be enlarged, in order to make them asymptotic confidence sets.

Let $n \geq 1$, let $D_n(X^n)$ be a credible set. For a nonnegative integer k_n , we define the k_n -enlargement of $D_n(X^n)$ to be the set

$$C_n(X^n) = \{\theta_n \in \Theta_n : \exists \eta_n \in D_n(X^n), m_n(\theta_n, \eta_n) \leq k_n\}.$$

Recall the definition of $B_{n,k_n}(\theta_{0,n})$ in eq. (19),

$$B_{n,k_n}(\theta_n) = \{\eta_n \in \Theta_n : m_n(\theta_n, \eta_n) \leq k_n\}.$$

We have the following result

LEMMA 5.5 *Suppose $P_{\theta_0} \Pi_n(B_{n,k_n}(\theta_{0,n}) | X^n) \geq 1 - x_n$, $0 < x_n < 1$. Let $\alpha_n \in (0, 1)$ and $D_n(X^n)$ a $1 - \alpha_n$ -credible set, with k_n -enlargement C_n , then*

$$P_{\theta_{0,n}}(\theta_{0,n} \in C_n(X^n)) \geq 1 - \frac{1}{1 - \alpha_n} x_n.$$

.

PROOF Let $E_n = \{\Pi(B_{n,k_n}(\theta_{0,n}) | X^n) \geq r\}$ be the event that $B_{n,k_n}(\theta_{0,n})$ has posterior mass at least r , $r > \alpha_n$. It follows from lemma 5.1 that $P_{\theta_{0,n}}(E_n) \geq 1 - \frac{1}{1-r} x_n$. As $D_n(X^n)$ has at least $1 - \alpha_n$ posterior mass, $D_n(X^n)$ and $B_{n,k_n}(\theta_{0,n})$ cannot be disjoint on the event E_n , as $1 - \alpha_n + r > 1$. Hence $\theta_0 \in C_n(X^n)$ on E_n . So $P_{\theta_0}(\theta_0 \in C_n(X^n)) \geq P_{\theta_0}(E_n) \geq 1 - \frac{1}{1-r} x_n$. As this holds for any $r > \alpha_n$ we have $P_{\theta_0}(\theta_0 \in C_n(X^n)) \geq 1 - \frac{1}{1-\alpha_n} x_n$. □

EXAMPLE 5.6 (*Kesten-Stigum phase, continuation of examples 3.6 and 4.6.*) Let $D_n(X^n)$ a $1 - \alpha_n$ -credible set and $C_n(X^n)$ the $a_n n$ enlargement of $D_n(X^n)$, with $a_n := 2(L-1)e^{2 - \frac{1}{4L}(c_n^{1/2} - d_n^{1/2})^2}$. With example 4.6 and lemma 5.5 we get

$$P_{\theta_{0,n}}(\theta_{0,n} \in C_n) \geq 1 - \frac{1}{1 - \alpha_n} \left(L_n \exp \left(-n \frac{1}{48L^2} (\sqrt{c_n} - \sqrt{d_n})^2 \right) + 2e^{-a_n n/4} \right).$$

□

6 Consistent hypothesis testing with posterior odds

Besides parameter estimation, an interesting question is testing between two alternatives, whether the true parameter $\theta_{0,n}$ is in the set A_n or in the set B_n , where $A_n, B_n \subseteq \Theta_n$ are disjoint non-random sets. In particular we consider symmetric testing between two alternatives

$$H_0 : \theta_{0,n} \in A_n \quad \text{versus} \quad H_1 : \theta_{0,n} \in B_n.$$

Taking, for example, $A_n = \Theta_{n,\ell_0}$ and $B_n = \Theta_n \setminus \Theta_{n,\ell_0}$ allows us to test whether the true parameter has ℓ_0 classes or not. The case $\ell_0 = 1$ is testing between the Erdős-Rényi model and the stochastic block model. We establish frequentist results for posterior odds testing between A_n and B_n .

We use posterior odds to test between the models, which is defined by

$$F_n = \frac{\Pi_n(B_n | X^n)}{\Pi_n(A_n | X^n)}.$$

Obviously, $F_n < 1$ counts as evidence in favour of H_0 and $F_n > 1$ as evidence in favour of H_1 . In the following theorem we give sufficient conditions for this Bayesian test to be valid in a frequentist sense.

THEOREM 6.1 *Let $\theta_{0,n} \in \Theta_n$. When $P_{\theta_{0,n}} \Pi_n(A_n | X^n) \geq 1 - a_n$, with $0 < a_n < 1$, then*

$$P_{\theta_{0,n}}(F_n > r_n) \leq 2a_n \left(1 + \frac{1}{r_n}\right).$$

If, in addition, $P_{\theta_{0,n}} \Pi_n(B_n | X^n) \leq b_n$, then

$$P_{\theta_{0,n}}(F_n > r_n) \leq 2a_n + \frac{2b_n}{r_n}.$$

PROOF From the posterior convergence condition on A_n it follows that $P_{\theta_{0,n}} \Pi_n(B_n | X^n) \leq a_n$. Hence the first result follows from the second, so we assume $P_{\theta_{0,n}} \Pi_n(B_n | X^n) \leq b_n$ in what follows. Let $E_n = \{\Pi_n(A_n | X^n) \geq 1/2\}$ be the event that the posterior gives at least mass 1/2 to A_n . It follows from lemma 5.1 that $P_{\theta_{0,n}}(E_n) \geq 1 - 2a_n$. So

$$P_{\theta_{0,n}}(F_n > r_n) \leq P_{\theta_{0,n}}(\Pi_n(B_n | X^n) \geq r_n/2) + 2a_n.$$

The probability on the right is by the Markov inequality bounded by

$$\frac{2}{r_n} P_{\theta_{0,n}} \Pi_n(B_n | X^n) \leq \frac{2b_n}{r_n}.$$

We thus arrive at the result

$$P_{\theta_{0,n}}(F_n > r_n) \leq 2a_n + \frac{2b_n}{r_n}.$$

□

Suppose one rejects the null-hypothesis when $F_n > r_n$, for some $r_n > 0$. The first order error is when H_0 is true, so $\theta_{0,n}$ is in fact in A_n , but H_0 is rejected (so $F_n > r_n$). The probability of this error is bounded by the theorem above. The error of second kind is when in fact H_1 is true, but H_0 is not rejected. This probability is given by $P_{\theta_{0,n}}(F_n < r_n)$, $\theta_{0,n} \in B_n$. As $P_{\theta_{0,n}}(F_n < r_n) = P_{\theta_{0,n}}(F_n^{-1} > 1/r_n)$, and reversing the roles of A_n and B_n in theorem 6.1, the probability of this event is also covered by the theorem, using posterior convergence results for $\theta_{0,n} \in B_n$. The power of the test is defined as the probability of rejecting the null hypothesis when H_1 is true. As $P_{\theta_{0,n}}(F_n > r_n) = P_{\theta_{0,n}}(F_n^{-1} < 1/r_n) = 1 - P_{\theta_{0,n}}(F_n^{-1} \geq 1/r_n)$, this probability can be lower bounded with the theorem above.

6.1 Examples: consistent Bayesian testing

In this section we determine conditions for Bayesian testing for different sparsity regimes. We consider the same priors and models as in example section 3.1.

We consider testing

$$H_0 : \theta_{0,n} \in \Theta_{n,\ell_0} \quad \text{versus} \quad H_1 : \theta_{0,n} \in \Theta_n \setminus \Theta_{n,\ell_0}.$$

So we consider testing whether the true parameter has ℓ_0 classes or not. The case $\ell_0 = 1$ is testing between the Erdős-Rényi graph and the stochastic block model.

EXAMPLE 6.2 (*Dense phase, continuation of example 3.4*) It follows from example 3.4 and theorem 6.1 that

$$P_{\theta_{0,n}}(F_n > r_n) \leq 2L_n \left(1 + \frac{1}{r_n}\right) e^{-b_n n^2 / (12L_n^2)}.$$

Taking $r_n \equiv 1$, the error of first kind is bounded by $4L_n e^{-b_n n^2 / (12L_n^2)}$. Reversing the roles of A_n and B_n in theorem 6.1 gives the error of second kind is bounded by $4e^{-b_n n^2 / (12L_n^2)}$, and the power of the test is lower bounded by $1 - 4e^{-b_n n^2 / (12L_n^2)}$. \square

EXAMPLE 6.3 (*Chernoff-Hellinger phase, continuation of example 3.5*) With example 3.5 and theorem 6.1 we get for $\theta_{0,n} \in \Theta_{n,\ell_0}$

$$P_{\theta_{0,n}}(F_n > r_n) \leq 2L_n \left(1 + \frac{1}{r_n}\right) \exp\left(-\frac{n}{48} \left(\sqrt{a_n} - \sqrt{b_n}\right)^2 \frac{\log n}{L_n^2}\right)$$

and for $\theta_{0,n} \in \Theta_n \setminus \Theta_{n,\ell_0}$,

$$P_{\theta_{0,n}}(F_n^{-1} > r_n^{-1}) \leq 2(1 + r_n) \exp\left(-\frac{n}{48} \left(\sqrt{a_n} - \sqrt{b_n}\right)^2 \frac{\log n}{L_n^2}\right),$$

which is the bound for the second order error. The power for the test is lower bounded by

$$1 - 2(1 + r_n) \exp\left(-\frac{n}{48} \left(\sqrt{a_n} - \sqrt{b_n}\right)^2 \frac{\log n}{L_n^2}\right).$$

□

EXAMPLE 6.4 (*Kesten-Stigum phase, continuation example 3.6*) It follows from example 3.6 and theorem 6.1 that

$$P_{\theta_{0,n}}(F_n > r_n) \leq 2L \left(1 + \frac{1}{r_n}\right) \exp\left(-\frac{n}{48L^2} \left(\sqrt{c_n} - \sqrt{d_n}\right)^2\right).$$

And similar results as in the example above hold for the second order error and the power of the test. □

A Definitions and conventions

Notation When S is a set, $|S|$ denotes the cardinality of S .

We assume for every $n \geq 1$, a random graph X^n taking values in the (finite) space \mathcal{X}_n of all undirected simple graphs (i.e. no self-loops or multiple edges) with n vertices. Let \mathcal{B}_n be the powerset of \mathcal{X}_n , be the σ -algebra corresponding to \mathcal{X}_n and let \mathcal{P}_n be the set of all probability distributions $P_n : \mathcal{B}_n \rightarrow [0, 1]$. A model is a subset \mathcal{P}_n of \mathcal{P}_n , which is parametrized by a bijective mapping $\Theta_n \rightarrow \mathcal{P}_n : \theta \mapsto P_{\theta_n}$. We equip Θ_n with a σ -algebra \mathcal{G}_n and a probability measure $\Pi_n : \mathcal{G}_n \rightarrow [0, 1]$ (the so-called prior). As we only consider finite parameters sets Θ_n , we set \mathcal{G}_n to be the powerset of Θ_n . As frequentists, we assume that there exists a ‘true, underlying distribution for the data’; that means that for every $n \geq 1$, the n -th graph X^n is drawn from $P_{\theta_{0,n}}$, for some $\theta_{0,n} \in \Theta_n$. We call $\theta_{0,n}$ the ‘true parameter’.

DEFINITION A.1 *Given $n \geq 1$ and a prior probability measure Π_n on Θ_n , define the n -th prior predictive distribution as:*

$$P_n^{\Pi_n}(A) = \int_{\Theta} P_{\theta_n}(A) d\Pi_n(\theta), \quad (22)$$

for all $A \in \mathcal{B}_n$.

The prior predictive distribution $P_n^{\Pi_n}$ is the marginal distribution for X^n in the Bayesian perspective that considers parameter and sample jointly $(\theta, X^n) \in \Theta \times \mathcal{X}_n$ as the random quantity of interest.

DEFINITION A.2 *Given $n \geq 1$, a (version of) the posterior is any map $\Pi_n(\cdot | X^n = \cdot) : \mathcal{G}_n \times \mathcal{X}_n \rightarrow [0, 1]$ such that,*

1. for every $B_n \in \mathcal{G}_n$, the map $\mathcal{X}_n \rightarrow [0, 1] : x^n \mapsto \Pi(B_n | X^n = x^n)$ is \mathcal{B}_n -measurable,
2. for all $A_n \in \mathcal{B}_n$ and $V_n \in \mathcal{G}_n$,

$$\int_{A_n} \Pi_n(V_n | X^n) dP_n^{\Pi_n} = \int_{V_n} P_{\theta_n}(A_n) d\Pi_n(\theta_n). \quad (23)$$

Bayes's Rule is expressed through equality (23) and is sometimes referred to as a 'disintegration' (of the joint distribution of (θ_n, X^n)). Because the models \mathcal{P}_n are dominated (denote the density of P_{θ_n} by p_{θ_n}), the fraction of integrated likelihoods,

$$\Pi(V_n|X^n) = \int_{V_n} p_{\theta_n}(X^n) d\Pi_n(\theta_n) \Big/ \int_{\Theta_n} p_{\theta_n}(X^n) d\Pi_n(\theta_n), \quad (24)$$

for all $V_n \in \mathcal{G}_n$, $n \geq 1$.

For completeness sake, we include Kleijn, 2020, lemma 2.2, which plays an essential role in our theorems on posterior consistency.

LEMMA A.3 *For any $B_n, V_n \in \mathcal{G}_n$ with $\Pi_n(B_n) > 0$ and any measurable map $\phi_n : \mathcal{X}_n \rightarrow [0, 1]$,*

$$\begin{aligned} \int P_{\theta_n} \Pi_n(V_n | X^n) d\Pi_n(\theta_n | B_n) &\leq \int P_{\theta_n}[\phi_n(X^n)] d\Pi_n(\theta_n | B_n) \\ &\quad + \frac{1}{\Pi_n(B_n)} \int_{V_n} P_{\theta_n}[1 - \phi_n(X^n)] d\Pi_n(\theta_n). \end{aligned}$$

B Existence of suitable tests

Given $n \geq 1$, and two class assignment vectors $\theta_{0,n}, \theta_n \in \Theta_n$, we are interested in determining testing power, for which we need the likelihood ratio $dP_{\theta_n}/dP_{\theta_{0,n}}$.

Fix $n \geq 1$, and let X^n denote the random graph associated with $\theta_{0,n} \in \Theta_n$, and let ℓ_0 be the number of different labels of $\theta_{0,n}$, so $\theta_{0,n} \in \Theta_{n,\ell_0}$. Let θ_n denote another element of Θ_n and suppose $\theta_n \in \Theta_{n,\ell}$, for some $\ell \in \mathcal{L}_n$ (which might or might not be equal to ℓ_0). Compare $p_{\theta_{0,n}}(X^n)$ with $p_{\theta_n}(X^n)$ in the likelihood ratio. Recall that the likelihood of θ_n is given by,

$$p_{\theta_n}(X^n) = \prod_{i < j} Q_{i,j;n}(\theta'_n)^{X_{ij}} (1 - Q_{i,j;n}(\theta'_n))^{1-X_{ij}},$$

where θ'_n is a representation of θ_n and

$$Q_{i,j;n}(\theta'_n) = \begin{cases} p_n, & \text{if } \theta'_i = \theta'_j, \\ q_n, & \text{if } \theta'_i \neq \theta'_j. \end{cases}$$

Let $\theta'_{0,n}$ be a representation of $\theta_{0,n}$ and θ'_n a representation of θ_n . Define two sets of edges, one consisting of all edges that connect within a class under $\theta'_{0,n}$ and between classes under θ'_n , and another consisting of all edges that connect between classes under $\theta'_{0,n}$ and within a class under θ'_n :

$$\begin{aligned} D_{1,n} &= \{(i, j) \in \{1, \dots, n\}^2 : i < j, \theta'_{0,n,i} = \theta'_{0,n,j}, \theta'_{n,i} \neq \theta'_{n,j}\}, \\ D_{2,n} &= \{(i, j) \in \{1, \dots, n\}^2 : i < j, \theta'_{0,n,i} \neq \theta'_{0,n,j}, \theta'_{n,i} = \theta'_{n,j}\}. \end{aligned}$$

Note that $D_{1,n}$ and $D_{2,n}$ do not depend on the chosen representations of $\theta_{0,n}$ and θ_n . Also define,

$$(S_n, T_n) := \left(\sum \{X_{ij} : (i, j) \in D_{1,n}\}, \sum \{X_{ij} : (i, j) \in D_{2,n}\} \right),$$

and note that, under $P_{\theta_{0,n}}$ and P_{θ_n} ,

$$(S_n, T_n) \sim \begin{cases} \text{Bin}(|D_{1,n}|, p_n) \times \text{Bin}(|D_{2,n}|, q_n), & \text{if } X^n \sim P_{\theta_{0,n}}, \\ \text{Bin}(|D_{1,n}|, q_n) \times \text{Bin}(|D_{2,n}|, p_n), & \text{if } X^n \sim P_{\theta_n}. \end{cases} \quad (25)$$

Since S_n and T_n are independent, the likelihood ratio is fixed as a product two exponentiated binomial random variables:

$$\frac{p_{\theta_n}}{p_{\theta_{0,n}}}(X^n) = \left(\frac{1-p_n}{p_n} \frac{q_n}{1-q_n} \right)^{S_n-T_n} \left(\frac{1-q_n}{1-p_n} \right)^{|D_{1,n}|-|D_{2,n}|} \quad (26)$$

This gives rise to the following lemma:

LEMMA B.1 *Let $n \geq 1$, $\theta_{0,n}, \theta_n \in \Theta_n$ be given. Then there exists a test function $\phi_n : \mathcal{X}_n \rightarrow [0, 1]$ such that,*

$$P_{\theta_{0,n}}\phi_n(X^n) + P_{\theta_n}(1 - \phi_n(X^n)) \leq \rho(p_n, q_n)^{|D_{1,n}|+|D_{2,n}|}.$$

PROOF The likelihood ratio test $\phi_n(X^n)$ has testing power bounded by the Hellinger transform,

$$P_{\theta_{0,n}}\phi_n(X^n) + P_{\theta_n}(1 - \phi_n(X^n)) \leq P_{\theta_{0,n}}\left(\frac{p_{\theta_n}}{p_{\theta_{0,n}}}(X^n)\right)^{1/2},$$

(see, *e.g.* Le Cam, 1986 and Kleijn, 2020, lemma 2.7). Then

$$\begin{aligned} P_{\theta_{0,n}}\left(\frac{p_{\theta_n}}{p_{\theta_{0,n}}}(X^n)\right)^{1/2} &= P_{\theta_{0,n}}\left(\frac{p_n}{1-p_n} \frac{1-q_n}{q_n}\right)^{\frac{1}{2}(T_n-S_n)} \left(\frac{1-q_n}{1-p_n}\right)^{\frac{1}{2}(|D_{1,n}|-|D_{2,n}|)} \\ &= P e^{\frac{1}{2}\lambda_n S_n} P e^{-\frac{1}{2}\lambda_n T_n} \left(\frac{1-q_n}{1-p_n}\right)^{\frac{1}{2}(|D_{1,n}|-|D_{2,n}|)}, \end{aligned}$$

where $\lambda_n := \log(1-p_n) - \log(p_n) + \log(q_n) - \log(1-q_n)$ and (S_n, T_n) are distributed binomially, as in the first part of (25). Using the moment-generating function of the binomial distribution, we conclude that,

$$\begin{aligned} P_{\theta_{0,n}}\left(\frac{p_{\theta_n}}{p_{\theta_{0,n}}}(X^n)\right)^{1/2} &= \left(1-p_n + p_n \left(\frac{1-p_n}{p_n} \frac{q_n}{1-q_n}\right)^{1/2}\right)^{|D_{1,n}|} \\ &\quad \times \left(1-q_n + q_n \left(\frac{p_n}{1-p_n} \frac{1-q_n}{q_n}\right)^{1/2}\right)^{|D_{2,n}|} \left(\frac{1-q_n}{1-p_n}\right)^{\frac{1}{2}(|D_{1,n}|-|D_{2,n}|)} \\ &= \rho(p_n, q_n)^{|D_{1,n}|+|D_{2,n}|}, \end{aligned}$$

which proves the assertion. \square

B.1 The sizes of $D_{1,n}$ and $D_{2,n}$

Let $\theta_n \in \Theta_{\mathbf{m}_{n,\ell}}$. There are $\frac{1}{2} \sum_{i=1}^{\ell} m_{n,\ell,i}(m_{n,\ell,i} - 1)$ pairs $(i, j), i < j$ with $\theta_{n,i} = \theta_{n,j}$.

We have

$$\frac{1}{2}n(m_{n,\ell,\min} - 1) \leq \frac{1}{2} \sum_{i=1}^{\ell} m_{n,\ell,i}(m_{n,\ell,i} - 1) \leq \frac{1}{2}n(m_{n,\ell,\max} - 1).$$

LEMMA B.2 Under assumption 2.1,

$$|D_{1,n} \cup D_{2,n}| \geq \frac{1}{2}n(m_{n,\ell_0 \wedge \ell, \min} - m_{n,\ell_0 \vee \ell, \max}).$$

PROOF As there are at least $\frac{1}{2}n(m_{n,\ell_0, \min} - 1)$ elements $(i, j), i < j$ for which $\theta_{0,n,i} = \theta_{0,n,j}$ and at most $\frac{1}{2}n(m_{n,\ell, \max} - 1)$ elements $(i, j), i < j$ for which $\theta_{n,i} = \theta_{n,j}$, in case $\frac{1}{2}n(m_{n,\ell_0, \min} - 1) > \frac{1}{2}n(m_{n,\ell, \max} - 1)$ going from $\theta_{0,n}$ to θ_n there are at least

$$\frac{1}{2}n(m_{n,\ell_0, \min} - 1) - \frac{1}{2}n(m_{n,\ell, \max} - 1) = \frac{1}{2}n(m_{n,\ell_0, \min} - m_{n,\ell, \max})$$

elements contributed to $D_{1,n}$.

In case $\frac{1}{2}n(m_{n,\ell, \min} - 1) \geq \frac{1}{2}n(m_{n,\ell_0, \max} - 1)$ going from θ_n to $\theta_{0,n}$ there are at least

$$\frac{1}{2}n(m_{n,\ell, \min} - 1) - \frac{1}{2}n(m_{n,\ell_0, \max} - 1) = \frac{1}{2}n(m_{n,\ell, \min} - m_{n,\ell_0, \max}).$$

elements contributed to $D_{2,n}$. It follows from assumption 2.1 that exactly one of these two options occurs. Hence we have that $D_{1,n} \cup D_{2,n}$ consist of at least

$$\frac{1}{2}n(m_{n,\ell_0 \wedge \ell, \min} - m_{n,\ell_0 \vee \ell, \max})$$

elements. □

REMARK B.3 The lower bound may be obtained. (Let $1^{k_1}2^{k_2} \dots \ell^{k_\ell}$ denote the label vector with k_1 consecutive 1's, followed by k_2 consecutive 2's, etc.) Consider an even ℓ_0 and let $\ell = \ell_0/2$ and $m = 2m_0$, \mathbf{m}_{n,ℓ_0} is the ℓ_0 -vector (m_0, \dots, m_0) and $\mathbf{m}_{n,\ell}$ is the ℓ -vector (m, \dots, m) . Let $\theta_{0,n} = (1^{m_0}2^{m_0} \dots \ell_0^{m_0}) \in \Theta_{\mathbf{m}_{n,\ell_0}}$ and $\theta_n = (1^{2m_0}2^{2m_0} \dots (\ell_0/2)^{2m_0}) \in \Theta_{\mathbf{m}_{n,\ell}}$. Note that when $\theta_{0,n,i} = k$, then $\theta_{n,i} = \lceil k/2 \rceil$. It follows that $D_{1,n}$ is an empty set in this case. On the other hand, if $\theta_{0,n,i} = 2k + 1$ and $\theta_{0,n,j} = 2k + 2$ with $k \in \{0, \dots, \ell_0/2 - 1\}$, then $\theta_{n,i} = \theta_{n,j} = k + 1$. There are $\frac{1}{2}\ell_0 m_0^2 = \frac{1}{2}n m_0 = \frac{1}{2}n|m_0 - m|$ of such combinations. Hence $\#D_{2,n} = \frac{1}{2}n|m_0 - m|$ and the lower bound is actually achieved.

Obviously, the bound above is useless when θ_n and $\theta_{0,n}$ both belong to the same set $\Theta_{n,\ell}$.

The following lemma relates the number of elements in $D_{1,n} \cup D_{2,n}$ to the r_n -distance (eq. (14)) between $\theta_{0,n}$ and θ_n .

LEMMA B.4 For $\theta_{0,n}, \theta_n \in \Theta_{n,\ell_0}$ with $r = r_n(\theta_{0,n}, \theta_n)$, we have

$$|D_{1,n} \cup D_{2,n}| \geq 2r(m_{n,\ell_0,\min} - r)^+,$$

where for a real number x , $x^+ = \max\{0, x\}$.

PROOF Let $\theta'_{0,n}$ and θ'_n be representations of $\theta_{0,n}$ and θ_n , respectively, so that $r'_n(\theta'_{0,n}, \theta'_n) = r$. There are $s, t \in \{1, \dots, \ell_0\}$, $s \neq t$ so that there are exactly r vertices with $\theta'_{0,n}$ -label s and θ'_n -label t . Clearly, when this happens, $m_{n,\ell_0,s} \geq r$, and trivially, $m_{n,\ell_0,s} \geq m_{n,\ell_0,1} \geq m_{n,\ell_0,\min}$. It follows that $D_{1,n}$ has at least $r(m_{n,\ell_0,s} - r) \geq r(m_{n,\ell_0,\min} - r)^+$ elements. Using that $D_{1,n}$ and $D_{2,n}$ are disjoint and reversing the role of $\theta_{0,n}$ and θ_n and $D_{1,n}$ and $D_{2,n}$ gives $|D_{1,n} \cup D_{2,n}| \geq 2r(m_{n,\ell_0,\min} - r)^+$. \square

B.2 The distance r_n and the sets $V_{n,\ell,k}$

LEMMA B.5 Let $\ell \leq \ell'$. Let $\theta \in \Theta_{\mathbf{m}_{n,\ell}}$ and $\eta \in \Theta_{\tilde{\mathbf{m}}_{n,\ell'}}$. The distance $r_n(\theta, \eta)$ is bounded by $m_{n,\ell,\ell}/2$, which in turn is bounded by $m_{n,\ell,\max}/2$.

PROOF Note that θ (η , resp.) is characterised by the partition of $\{1, \dots, n\}$ in ℓ (ℓ' , resp.) sets of sizes $m_{n,\ell,1}, \dots, m_{n,\ell,\ell}$ ($\tilde{m}_{n,\ell',1}, \dots, \tilde{m}_{n,\ell',\ell'}$, resp.). Vice versa, for such a partition, one can define a labelling, by assigning a label to each set in the partition, and giving each element of the set that label. Define

$$\Xi_\theta = \left\{ \{i : \theta'_i = a\} : a \in \{1, \dots, \ell\} \right\},$$

note that Ξ_θ is independent of the representation, and define Ξ_η similarly. Now note that there are at most ℓ pairs of sets (A, B) , $A \in \Xi_\theta, B \in \Xi_\eta$ with $|A \cap B| > m_{n,\ell,\ell}/2$, because Ξ_θ consists of ℓ elements and Ξ_η consists of $\ell' \geq \ell$ elements, and if there are more than ℓ such pairs, then there is an A in Ξ_θ and two different B_1, B_2 in Ξ_η such that $A \cap B_1$ and $A \cap B_2$ have more than $m_{n,\ell,\ell}/2$ elements. But B_1 and B_2 are disjoint, so A has more than $m_{n,\ell,\ell}$ elements, which is a contradiction.

Let $(A_1, B_1), \dots, (A_k, B_k)$ the pairs with $|A_i \cap B_i| > m_{n,\ell,\ell}/2$, with $k \leq \ell (\leq \ell')$. Define a label $\theta' \in \theta$ by assigning label i to the sets A_i , $i \in \{1, \dots, k\}$ and the remaining $\ell - k$ labels to the remaining sets in the partition Ξ_θ and define $\eta' \in \eta$ similarly by assigning label i to the sets B_i , $i \in \{1, \dots, k\}$ and the remaining $\ell' - k$ labels to the remaining sets in Ξ_η . It follows that $r(\theta, \eta) \leq r(\theta', \eta') \leq m_{n,\ell,\ell}/2$. By the definition of $m_{n,\ell,\max}$, $m_{n,\ell,\ell} \leq m_{n,\ell,\max}$. \square

Let $\theta \in \Theta_{\mathbf{m}_{n,\ell}}$ and define $V_{n,\ell,k}(\theta) = \{\eta \in \Theta_{n,\ell} : r_n(\theta, \eta) = k\}$.

LEMMA B.6 Let $\theta \in \Theta_{\mathbf{m}_{n,\ell}}$, and $k \leq m_{n,\ell,1}$. Then

$$|V_{n,\ell,k}(\theta)| \leq 2^{k\ell(\ell-1)} \binom{n(\ell-1)}{\ell(\ell-1)k} \leq \left(\frac{2en}{k\ell}\right)^{\ell(\ell-1)k}. \quad (27)$$

Note that eq. (27) is well defined when $k\ell \leq n$, which is the case when $k \leq m_{n,\ell,1}$.

PROOF When $\ell = 1$, the lemma is trivial. Let $\ell > 1$. Choose a representation θ' of θ so that exactly $m_{n,\ell,i}$ labels of θ' are i , $i \in \{1, \dots, \ell\}$. Let us count the number of ways in which we can change the labelling θ' into a label η' , while keeping $r_n(\theta', \eta') \leq k$. Let $i, j \in \{1, \dots, \ell\}$, $i \neq j$. We can choose $\binom{m_{n,\ell,i}}{k}$ vertices with θ' -label i and either retain the label i or give them the label j . This is possible in 2^k ways. There are $\ell - 1$ labels $j \neq i$, and there are ℓ labels i . Hence $V_{n,\ell,k}$ has at most

$$2^{k\ell(\ell-1)} \prod_{i=1}^{\ell} \binom{m_{n,\ell,i}}{k}^{\ell-1}$$

elements.

Using $\binom{a}{b} \binom{c}{d} \leq \binom{a+c}{b+d}$, the number of elements in $V_{n,\ell,k}$ is bounded by

$$2^{k\ell(\ell-1)} \binom{n(\ell-1)}{(\ell-1)k}.$$

Equation (27) is in turn bounded by

$$\left(\frac{2en}{k\ell}\right)^{\ell(\ell-1)k}.$$

□

B.3 The distance m_n

The distance m_n , restricted to $\Theta_{n,\ell}$ has the following relationship with r_n :

LEMMA B.7 For all $\theta, \eta \in \Theta_{n,\ell}$,

$$r_n(\theta, \eta) \leq m(\theta, \eta) \leq \ell(\ell-1)r_n(\theta, \eta). \quad (28)$$

PROOF When $\ell = 1$, $\Theta_{n,\ell}$ consist of only one element, and eq. (28) is trivial. Now suppose $\ell > 1$. Every representation θ' of θ and η' of η satisfies $r'_n(\theta', \eta') \geq r_n(\theta, \eta) =: r$. Hence, for fixed representations θ' and η' , there are at least r vertices with θ' label a have η' label b , for some $a, b \in \{1, \dots, \ell\}$, $a \neq b$. Hence $m'_n(\theta', \eta') \geq r$. So $m_n(\theta, \eta) \geq r$. On the other hand, if θ' and η' are representations of θ and η , respectively, such that $r'_{n,\ell}(\theta', \eta') = r_{n,\ell}(\theta, \eta) =: r$, then there are at most r vertices with θ' label 1, that have η' label 2, and at most r vertices with θ' label 1 that have η' label 3, etc. As there are $\ell - 1$ labels not equal to 1, and θ' has ℓ labels, $m(\theta, \eta) \leq m'(\theta', \eta') \leq \ell(\ell-1)r$. □

C Auxiliary lemmas

LEMMA C.1 *Let $C \geq 2$. For all $x \geq \sqrt{2/C}$,*

$$\frac{e^{-Cx}}{1 - e^{-x}} \leq e^{-Cx/4}.$$

PROOF Note that

$$\begin{aligned} 1 - e^{-x} &= \int_0^x e^{-y} dy \\ &\geq xe^{-x}. \end{aligned}$$

Using this and the fact that $C - 1 \geq C/2$, we see that

$$\frac{e^{-Cx}}{1 - e^{-x}} \leq \frac{e^{-Cx}}{xe^{-x}} = \frac{e^{-(C-1)x}}{x} \leq \frac{e^{-Cx/2}}{x} = \frac{e^{-Cx/4}}{x} e^{-Cx/4}.$$

As $x \geq \sqrt{2/C}$ is equivalent to $Cx/4 \geq x^{-1}/2$ and $x > 0$, we have

$$\frac{e^{-Cx}}{1 - e^{-x}} \leq x^{-1} e^{-x^{-1}/2} e^{-Cx/4}.$$

One verifies that $f(y) = ye^{-y/2}$ attains its maximum on \mathbb{R} at $y = 2$ and $f(2) = 2/e < 1$.

It now follows that

$$\frac{e^{-Cx}}{1 - e^{-x}} \leq e^{-Cx/4}.$$

□

LEMMA C.2 *For $x \in [0, 1]$, $\sqrt{1-x} \leq 1 - x/2$.*

PROOF Define $f(x) = \sqrt{1-x}$ and $g(x) = 1 - x/2$. Note that $f(0) = g(0)$ and $f'(x) = -\frac{1}{2\sqrt{1-x}} \leq -\frac{1}{2} = g'(x)$, for all $x \in [0, 1]$. It follows that $g(x) \geq f(x)$ for all $x \in [0, 1]$. □

LEMMA C.3 *For all positive integers r and real numbers $x > -r$, $(1 + x/r)^r \leq e^x$.*

PROOF Let for $x > -r$, $f(x) = r \log(1 + x/r)$ and $g(x) = x$. Then $f'(x) = \frac{1}{1+x/r}$ and $g'(x) = 1$. It follows that $f'(x) \leq g'(x)$, when $x \geq 0$, $f'(x) > g'(x)$ when $-n < x < 0$ and $f(0) = g(0)$. It follows that $f(x) \leq g(x)$ for all $x > -r$. As $y \rightarrow e^y$ is increasing, for all real y , it follows that for all $x > -n$, $(1 + x/r)^r = e^{f(x)} \leq e^{g(x)} = e^x$. □

References

Abbe, E. (2018). “Community Detection and Stochastic Block Models: Recent Developments”. In: *Journal of Machine Learning Research* 18.177, pp. 1–86. URL: <http://jmlr.org/papers/v18/16-480.html>.

- Abbe, E. and C. Sandon (2015). “Community Detection in General Stochastic Block models: Fundamental Limits and Efficient Algorithms for Recovery”. In: *2015 IEEE 56th Annual Symposium on Foundations of Computer Science*, pp. 670–688. DOI: [10.1109/FOCS.2015.47](https://doi.org/10.1109/FOCS.2015.47).
- Abbe, Emmanuel and Colin Sandon (2018). “Proof of the Achievability Conjectures for the General Stochastic Block Model”. In: *Communications on Pure and Applied Mathematics* 71.7, pp. 1334–1406. DOI: [10.1002/cpa.21719](https://doi.org/10.1002/cpa.21719).
- Channarond, A., J.-J. Daudin, and S. Robin (2012). “Classification and estimation in the Stochastic Blockmodel based on the empirical degrees”. In: *Electron. J. Statist.* 6, pp. 2574–2601. DOI: [10.1214/12-EJS753](https://doi.org/10.1214/12-EJS753).
- Chen, Jingchun and Bo Yuan (2006). “Detecting functional modules in the yeast protein-protein interaction network”. In: *Bioinformatics* 22.18, pp. 2283–2290. DOI: [10.1093/bioinformatics/btl370](https://doi.org/10.1093/bioinformatics/btl370).
- Cline, M.S. et al. (2007). “Integration of biological networks and gene expression data using Cytoscape”. In: *Nature protocols* 2.10, pp. 2366–2382. ISSN: 1750-2799. DOI: [10.1038/nprot.2007.324](https://doi.org/10.1038/nprot.2007.324).
- Freedman, D. (1999). “On the Bernstein-Von Mises Theorem with Infinite-Dimensional Parameters”. In: *The Annals of Statistics* 27.4, pp. 1119–1140. ISSN: 00905364. URL: <http://www.jstor.org/stable/120155>.
- Gao, C., A.W. van der Vaart, and H.H. Zhou (2019). “A General Framework for Bayes Structured Linear Models”. In: *Annals of Statistics (accepted)*. URL: <https://www.e-publications.org/ims/submission/AOS/user/submissionFile/27372?confirm=67ba2787>.
- Geng, J., A. Bhattacharya, and D. Pati (2019). “Probabilistic Community Detection With Unknown Number of Communities”. In: *Journal of the American Statistical Association* 114.526, pp. 893–905. DOI: [10.1080/01621459.2018.1458618](https://doi.org/10.1080/01621459.2018.1458618).
- Ghosh, P., D. Pati, and A. Bhattacharya (2019). “Posterior Contraction Rates for Stochastic Block Models”. In: *Sankhya A*. ISSN: 0976-8378. DOI: [10.1007/s13171-019-00180-5](https://doi.org/10.1007/s13171-019-00180-5).
- Holland, P. W., K. B. Laskey, and S. Leinhardt (1983). “Stochastic blockmodels: First steps”. In: *Social Networks* 5.2, pp. 109–137. ISSN: 0378-8733. DOI: [https://doi.org/10.1016/0378-8733\(83\)90021-7](https://doi.org/10.1016/0378-8733(83)90021-7).
- Kleijn, B. J. K. (2020). “Frequentist validity of Bayesian limits”. In: *Annals of Statistics (accepted)*. URL: <https://www.e-publications.org/ims/submission/AOS/user/submissionFile/34072?confirm=0095591e>.
- Kleijn, B. J. K. and J. van Waaij (2018). “Recovery, detection and confidence sets of communities in a sparse stochastic block model”. In: *arXiv e-prints*. arXiv: [1810.09533](https://arxiv.org/abs/1810.09533).

- Le Cam, L. (1986). *Asymptotic methods in statistical decision theory*. Springer-Verlag New York. ISBN: 978-1-4612-4946-7. DOI: [10.1007/978-1-4612-4946-7](https://doi.org/10.1007/978-1-4612-4946-7).
- Mariadassou, M. and C. Matias (2015). “Convergence of the groups posterior distribution in latent or stochastic block models”. In: *Bernoulli* 21.1, pp. 537–573. DOI: [10.3150/13-BEJ579](https://doi.org/10.3150/13-BEJ579).
- Mossel, E., J. Neeman, and A. Sly (2016). “Consistency thresholds for the planted bisection model”. In: *Electron. J. Probab.* 21, 24 pp. DOI: [10.1214/16-EJP4185](https://doi.org/10.1214/16-EJP4185).
- van der Pas, S.L. and A.W. van der Vaart (2018). “Bayesian Community Detection”. In: *Bayesian Anal.* 13.3, pp. 767–796. DOI: [10.1214/17-BA1078](https://doi.org/10.1214/17-BA1078).
- Zhang, A.Y. and H.H. Zhou (2016). “Minimax rates of community detection in stochastic block models”. In: *Ann. Statist.* 44.5, pp. 2252–2280. DOI: [10.1214/15-AOS1428](https://doi.org/10.1214/15-AOS1428).