

Global lower mass-bound for critical configuration models in the heavy-tailed regime

Shankar Bhamidi¹, Souvik Dhara^{2,3}, Remco van der Hofstad⁴, Sanchayan Sen⁵

March 18, 2022

Abstract

We establish the global lower mass-bound property for the largest connected components in the critical window for the configuration model when the degree distribution has an infinite third moment. The scaling limit of the critical percolation clusters, viewed as measured metric spaces, was established in [7] with respect to the Gromov-weak topology. Our result extends those scaling limit results to the stronger Gromov-Hausdorff-Prokhorov topology under slightly stronger assumptions on the degree distribution. This implies the distributional convergence of global functionals such as the diameter of the largest critical components. Further, our result gives a sufficient condition for compactness of the random metric spaces that arise as scaling limits of critical clusters in the heavy-tailed regime.

1 Introduction

Any connected graph \mathcal{C} can be viewed as a metric space with the distance between points given by $ad(\cdot, \cdot)$ for some constant $a > 0$, where $d(\cdot, \cdot)$ is used as a generic notation to denote the graph-distance (i.e., number of edges in the shortest path between vertices). There is a natural (counting) probability measure associated to (\mathcal{C}, ad) given by $\mu(A) = |A|/|\mathcal{C}|$ for any $A \subset \mathcal{C}$, where $|A|$ denotes the number of vertices in A . We denote the above metric space with a measure by (\mathcal{C}, ad) . Fix any $\delta > 0$ and define the δ -lower mass of (\mathcal{C}, ad) by

$$m(\delta) := \frac{\inf_{u \in \mathcal{C}} |\{v \in \mathcal{C} : ad(v, u) \leq \delta\}|}{|\mathcal{C}|}. \quad (1.1)$$

Thus, $m(\delta)$ is the least *mass* in any δ -neighborhood of a vertex in (\mathcal{C}, ad) . For a sequence $(\mathcal{C}_n, a_n)_{n \geq 1}$ of graphs viewed as metric spaces endowed with a measure, the global lower mass-bound property is defined as follows:

Emails: bhamidi@email.unc.edu, sdhara@mit.edu, r.w.v.d.hofstad@tue.nl, sanchayan.sen1@gmail.com

¹Department of Statistics and Operations Research, University of North Carolina

²Department of Mathematics, Massachusetts Institute of Technology

³Microsoft Research Lab – New England

⁴Department of Mathematics and Computer Science, Eindhoven University of Technology

⁵Department of Mathematics, Indian Institute of Science

2010 *Mathematics Subject Classification.* Primary: 60C05, 05C80.

Keywords and phrases. Global lower mass-bound, critical configuration model, heavy-tailed degrees

Acknowledgment. SB was partially supported by NSF grants DMS-1613072, DMS-1606839 and ARO grant W911NF-17-1-0010. SD, RvdH was supported by the Netherlands Organisation for Scientific Research (NWO) through Gravitation Networks grant 024.002.003. In addition, RvdH was supported by VICI grant 639.033.806. SS has been supported in part by the Infosys foundation, Bangalore, and by a MATRICS grant from SERB. SD thanks Eindhoven University of Technology, where a major part of this work was done.

Definition 1 (Global lower mass-bound property [5]). For $\delta > 0$, let $m_n(\delta)$ denote the δ -lower mass of (\mathcal{C}_n, a_n) . Then $(\mathcal{C}_n, a_n)_{n \geq 1}$ is said to satisfy the global lower mass-bound property if and only if $\sup_{n \geq 1} m_n(\delta)^{-1} < \infty$ for any $\delta > 0$. When $(\mathcal{C}_n)_{n \geq 1}$ is a collection of random graphs, $(\mathcal{C}_n, a_n)_{n \geq 1}$ is said to satisfy the global lower mass-bound property if and only if $(m_n(\delta)^{-1})_{n \geq 1}$ is a tight sequence of random variables for any $\delta > 0$.

The aim of this paper is to prove the global lower mass-bound property for connected components of random graphs with given degrees (configuration model) at criticality, when the third moment of the empirical degree distribution tends to infinity (Theorem 1.1). Informally speaking, the global lower mass-bound property ensures that all the small neighborhoods of vertices in the ‘large’ critical component have mass bounded away from zero, so that the component does not have any *light spots* and the total mass is well-distributed over the whole component. This has several interesting consequences in the theory of critical random graphs. Our main motivation comes from the work of Athreya, Löhner and Winter [5], who have shown that the global lower mass-bound property can be used to prove Gromov-Hausdorff-Prokhorov (GHP) convergence of random metric spaces. In a previous paper [7], we have studied the critical percolation clusters for the configuration model in the heavy-tailed universality class. We have proved that the ordered vector of components converges in distribution to suitable random objects in the Gromov-weak topology. The global lower mass-bound in this paper shows that the result of [7] in fact holds with respect to the stronger GHP-topology. One motivating reason for proving the GHP-convergence is that it yields the scaling limit of global functionals like the diameter of large critical components. Finding the scaling limit for the diameter of critical components is a daunting task even for the Erdős-Rényi random graph. Nachmias and Peres [31] estimated the tail probabilities of the diameter, but showing a distributional convergence result was a difficult question, until the seminal paper by Addario-Berry, Broutin and Goldschmidt [2] that proved the GHP-convergence for critical Erdős-Rényi random graphs. As a corollary of Theorem 1.1, we also get distributional convergence of the suitably rescaled diameter of the critical percolation clusters in the heavy-tailed regime (Theorem 1.4), where the scaling limit and exponents turn out to be different than those for the Erdős-Rényi case.

We will further discuss the applications and the scope of this work as well as its technical contributions after stating our results in Section 1.3. We start by defining the configuration model and state the precise assumptions.

1.1 The configuration model

Consider a non-increasing sequence of degrees $\mathbf{d} = (d_i)_{i \in [n]}$ such that $\ell_n = \sum_{i \in [n]} d_i$ is even. For notational convenience, we suppress the dependence of the degree sequence on n . The configuration model on n vertices having degree sequence \mathbf{d} is constructed as follows [6, 10]:

Equip vertex j with d_j stubs, or *half-edges*. Two half-edges create an edge once they are paired. Therefore, initially we have $\ell_n = \sum_{i \in [n]} d_i$ half-edges. Pick any one half-edge and pair it with another uniformly chosen half-edge from the remaining unpaired half-edges and keep repeating the above procedure until all the unpaired half-edges are exhausted.

Let $\text{CM}_n(\mathbf{d})$ denote the graph constructed by the above procedure. Note that $\text{CM}_n(\mathbf{d})$ may contain self-loops or multiple edges. Given any degree sequence, let $\text{UM}_n(\mathbf{d})$ denote the graph chosen uniformly at random from the collection of all simple graphs with degree sequence \mathbf{d} . It can be shown that the conditional law of $\text{CM}_n(\mathbf{d})$, conditioned on it being simple, is the same as $\text{UM}_n(\mathbf{d})$ (see e.g. [23, Proposition 7.13]).

1.2 Main results

Fix a constant $\tau \in (3, 4)$, which will denote the power-law exponent of the asymptotic degree distribution of $\text{CM}_n(\mathbf{d})$. Throughout this paper we will use the shorthand notation

$$\alpha = 1/(\tau - 1), \quad \rho = (\tau - 2)/(\tau - 1), \quad \eta = (\tau - 3)/(\tau - 1). \quad (1.2)$$

We use the standard notation of $\xrightarrow{\mathbb{P}}$ and \xrightarrow{d} to denote convergence in probability and in distribution, respectively. Also, we use a generic notation C to denote a positive universal constant whose exact value may change from line to line. We use Bachmann–Landau asymptotic notation $o(\cdot)$, $O(\cdot)$, $\Theta(\cdot)$, $\omega(\cdot)$, $\Omega(\cdot)$. A sequence of events $(\mathcal{E}_n)_{n \geq 1}$ is said to occur with high probability (whp) with respect to the probability measures $(\mathbb{P}_n)_{n \geq 1}$ when $\mathbb{P}_n(\mathcal{E}_n) \rightarrow 1$. For (random) variables X_n and Y_n , define $X_n = O_{\mathbb{P}}(Y_n)$ when $(|X_n|/|Y_n|)_{n \geq 1}$ is a tight sequence; $X_n = o_{\mathbb{P}}(Y_n)$ when $X_n/Y_n \xrightarrow{\mathbb{P}} 0$; $X_n = \Theta_{\mathbb{P}}(Y_n)$ if both $X_n = O_{\mathbb{P}}(Y_n)$ and $Y_n = O_{\mathbb{P}}(X_n)$.

We first state the general assumptions that are used to prove scaling limits for critical configuration models with heavy-tailed degree distributions as identified previously in [7, 16]:

Assumption 1 (General assumptions). For each $n \geq 1$, let $\mathbf{d} = \mathbf{d}_n = (d_1, \dots, d_n)$ be a degree sequence satisfying $d_1 \geq d_2 \geq \dots \geq d_n$. We assume the following about $(\mathbf{d}_n)_{n \geq 1}$ as $n \rightarrow \infty$:

- (i) (*High-degree vertices*) For each fixed $i \geq 1$,

$$n^{-\alpha} d_i \rightarrow \theta_i, \quad (1.3)$$

where $\boldsymbol{\theta} = (\theta_1, \theta_2, \dots) \in \ell_{\downarrow}^3 \setminus \ell_{\downarrow}^2$, where $\ell^p := \{(x_i)_{i \geq 1} : x_1 \geq x_2 \geq \dots \text{ and } \sum_i x_i^p < \infty\}$.

- (ii) (*Moment assumptions*) Let D_n denote the degree of a typical vertex, i.e., a vertex chosen uniformly at random from the vertex set $[n]$, independently of $\text{CM}_n(\mathbf{d})$. Then, D_n converges in distribution to some discrete random variable D and

$$\frac{1}{n} \sum_{i \in [n]} d_i \rightarrow \mu := \mathbb{E}[D], \quad \frac{1}{n} \sum_{i \in [n]} d_i^2 \rightarrow \mu_2 := \mathbb{E}[D^2], \quad (1.4)$$

$$\lim_{K \rightarrow \infty} \limsup_{n \rightarrow \infty} n^{-3\alpha} \sum_{i=K+1}^n d_i^3 = 0. \quad (1.5)$$

- (iii) Let n_1 be the number of degree-one vertices. Then $n_1 = \Theta(n)$, which is equivalent to assuming that $\mathbb{P}(D = 1) > 0$.

We further assume that the configuration model lies within the critical window of the phase transition, i.e., for some $\lambda \in \mathbb{R}$,

$$\nu_n = \frac{\sum_{i \in [n]} d_i(d_i - 1)}{\sum_{i \in [n]} d_i} = 1 + \lambda n^{-\eta} + o(n^{-\eta}). \quad (1.6)$$

Denote the i -th largest connected component of $\text{CM}_n(\mathbf{d})$ by $\mathcal{C}_{(i)}$, breaking ties arbitrarily. For each $v \in [n]$ and $\delta > 0$, let $\mathcal{N}_v(\delta)$ denote the δn^η neighborhood of v in $\text{CM}_n(\mathbf{d})$ in the graph distance. For each $i \geq 1$, define

$$\mathbf{m}_i^n(\delta) = \inf_{v \in \mathcal{C}_{(i)}} n^{-\rho} |\mathcal{N}_v(\delta)|. \quad (1.7)$$

Our goal is to prove the global lower mass-bound property for the critical components $\mathcal{C}_{(i)}$. For $\text{CM}_n(\mathbf{d})$ satisfying Assumption 1 and (1.6), it was shown in [16, Theorem 1] that

$$(n^{-\rho} |\mathcal{C}_{(i)}|)_{i \geq 1} \xrightarrow{d} (\xi_i)_{i \geq 1}, \quad (1.8)$$

with respect to the ℓ_{\downarrow}^2 -topology, where the ξ_i 's are non-degenerate random variables with support $(0, \infty)$. Therefore, it is enough to rescale by n^ρ in (1.7) instead of the component sizes as given in (1.1). In order to prove tightness of $\mathfrak{m}_i^n(\delta)$, we will need a further technical assumption on the degrees.

Assumption 2. (i) Let V_n^* be the degree of a vertex chosen in a size-biased manner with sizes being $(d_i/\ell_n)_{i \in [n]}$, i.e., $\mathbb{P}(V_n^* = i) = d_i/\ell_n$, and let D_n^* be the degree of V_n^* . There exists a constant $c_0 > 0$ such that for all $1 \leq l \leq d_1$ and $n \geq 1$,

$$\mathbb{P}(D_n^* \geq l) \geq \frac{c_0}{l^{\tau-2}}. \quad (1.9)$$

(ii) Let $\beta_i^n = n^{-2\alpha} \sum_{j=1}^{i-1} d_j^2$. There exists a sequence $(K_n)_{n \geq 1}$ with $K_n \rightarrow \infty$, and $K_n = o(n^\alpha)$ such that $\beta_{K_n}^n = \omega(\log n)$.

(iii) For all $\varepsilon > 0$,

$$\lim_{K \rightarrow \infty} \limsup_{n \rightarrow \infty} \sum_{i \in (K, K_n)} \left(\frac{d_i}{n^\alpha} \right) \times e^{-\varepsilon \beta_i^n} = 0. \quad (1.10)$$

The following theorem is the main result of this paper:

Theorem 1.1 (Global lower mass-bound for $\text{CM}_n(\mathbf{d})$). *Suppose that Assumptions 1, 2 and the criticality condition (1.6) hold. Then, for each fixed $i \geq 1$, $(\mathcal{C}_{(i), n^{-\eta}})_{n \geq 1}$ satisfies the global lower mass-bound, i.e., for any $\delta > 0$, the sequence $(\mathfrak{m}_i^n(\delta)^{-1})_{n \geq 1}$ is tight.*

By [24, Theorem 1.1], under the condition (1.4) in Assumption 1,

$$\liminf_{n \rightarrow \infty} \mathbb{P}(\text{CM}_n(\mathbf{d}) \text{ is simple}) > 0. \quad (1.11)$$

This immediately implies the following:

Theorem 1.2 (Global lower mass-bound for $\text{UM}_n(\mathbf{d})$). *Under Assumption 1, 2 and (1.6), the largest components of $\text{UM}_n(\mathbf{d})$ also satisfy the global lower mass-bound property.*

Next we state another important corollary, which says that the global lower mass-bound property is also satisfied by critical percolation clusters of $\text{CM}_n(\mathbf{d})$ and $\text{UM}_n(\mathbf{d})$. To this end, let us assume that

$$\lim_{n \rightarrow \infty} \frac{\sum_{i \in [n]} d_i(d_i - 1)}{\sum_{i \in [n]} d_i} = \nu > 1. \quad (1.12)$$

In this regime, $\text{CM}_n(\mathbf{d})$ is supercritical in the sense that there exists a unique *giant* component whp for $\nu > 1$, and when $\nu < 1$, all the components have size $o_{\mathbb{P}}(n)$ [26, 30]. Percolation refers to deleting each edge of a graph independently with probability $1 - p$. The critical window for percolation on $\text{CM}_n(\mathbf{d})$ in the heavy-tailed setting was studied in [7, 16], and is defined by the values of p given by

$$p_c(\lambda) = \frac{1}{\nu_n} + \frac{\lambda}{n^\eta} + o(n^{-\eta}). \quad (1.13)$$

Let $\mathcal{C}_{(i)}(p_c(\lambda))$ denote the i -th largest component of the graph obtained by percolation with probability $p_c(\lambda)$ on the graph $\text{CM}_n(\mathbf{d})$. Then the following result holds:

Theorem 1.3 (Global lower mass-bound for critical percolation). *Under Assumptions 1, 2 and (1.13), $(\mathcal{C}_{(i)}(p_c(\lambda)), n^{-\eta})_{n \geq 1}$ satisfies the global lower mass-bound property, for each fixed $i \geq 1$. This result also holds for percolation on $\text{UM}_n(\mathbf{d})$.*

Let \mathcal{G}_n denote the graph obtained by doing percolation with retention probability $p_c(\lambda)$ (defined in (1.13)) on $\text{CM}_n(\mathbf{d})$. Let $\mathbf{d}^p = (d_i^p)_{i \in [n]}$ denote the degree sequence of \mathcal{G}_n . By [19, Lemma 3.2], the conditional law of \mathcal{G}_n , conditionally on \mathbf{d}^p , is same as the law of $\text{CM}_n(\mathbf{d}^p)$. Thus, Theorem 1.3 follows from Theorem 1.1 if we can show that the percolated degree sequence satisfies (with possibly different parameters) Assumptions 1 and 2 with high probability when the original degree sequence $(d_i)_{i \in [n]}$ does so, and also (1.6) holds. The verification of these assumptions are provided in Section 5.

Remark 1. It is worthwhile to point out that Theorem 1.1 can be proved when the $\mathcal{C}_{(i)}$'s are endowed with a more general measure rather than the counting measure. To be precise, for any sequence of vertex weights $(w_v)_{v \in [n]}$, the component $\mathcal{C}_{(i)}$ can be equipped with the measure $\mu_{(i)}(A) = \sum_{v \in A} w_v / \sum_{v \in \mathcal{C}_{(i)}} w_v$, for any $A \subset \mathcal{C}_{(i)}$. Then Theorem 1.1 can also be proved using identical methods as in this paper, with the additional assumptions that

$$\lim_{n \rightarrow \infty} \frac{1}{\ell_n} \sum_{i \in [n]} d_i w_i = \mu_w, \quad \max \left\{ \sum_{i \in [n]} d_i w_i^2, \sum_{i \in [n]} d_i^2 w_i \right\} = O(n^{3\alpha}).$$

These additional assumptions are required when we apply the results from [16] (see [16, Theorem 21]). We adopted the simpler version of the counting measure here because it relates directly to [7, Theorem 2.1].

1.3 Discussion

Scaling limit of critical percolation clusters. We write $n^{-\eta} \mathcal{C}_{(i)}(p_c(\lambda))$ to denote the i -th largest component of $\text{CM}_n(\mathbf{d}, p_c(\lambda))$, viewed as a measured metric space with the metric being the graph distance re-scaled by n^η , and the measure being proportional to the counting measure. Athreya, Löhner and Winter [5] showed that the global lower mass-bound property forms a crucial ingredient to prove convergence of random metric spaces such as $n^{-\eta} \mathcal{C}_{(i)}(p_c(\lambda))$ with respect to the Gromov-Hausdorff-Prokhorov (GHP) topology on the space of compact metric spaces. The other key ingredient is the scaling limit for $n^{-\eta} \mathcal{C}_{(i)}(p_c(\lambda))$ with respect to the Gromov-weak topology, which was established in [7, Theorem 2.1]. The Gromov-weak topology is an analogue of finite-dimensional convergence, since it considers distances between a finite number of sampled points from the underlying metric space. Thus, global functionals such as the diameter are not continuous with respect to this topology. Indeed, it may be the case that there is a long path of growing length, that has asymptotically negligible mass. In our context, the problem could arise due to paths of length much larger than n^η . The global lower mass-bound property ensures that the components have sufficient mass everywhere. This forbids the existence of long thin paths, when the total mass of the component converges. For this reason, Gromov-weak convergence and global lower mass-bound together imply GHP-convergence when the support of the limiting measure is the entire limiting space [5, Theorem 6.1]. For formal definitions of the Gromov-weak topology, and the GHP-topology on the space of compact measured metric spaces, we refer the reader to [5, 8, 22].

Following the above discussion, the next theorem is a direct consequence of Theorem 1.3, [7, Theorem 2.1] and [5, Theorem 6.1]: Let \mathbb{M} denote the space of measured compact metric spaces equipped with the GHP-topology, and let $\mathbb{M}^{\mathbb{N}}$ denote the product space with the associated product topology.

Theorem 1.4 (GHP convergence of critical percolation clusters). *There exists a sequence of measured metric spaces $(\mathcal{M}_i)_{i \geq 1} = ((M_i, d_i, \mu_i))_{i \geq 1} \in \mathbb{M}^{\mathbb{N}}$ such that, under Assumptions 1, 2 and (1.13), as $n \rightarrow \infty$,*

$$(n^{-\eta} \mathcal{C}_{(i)}(p_c(\lambda))) \xrightarrow{d} (\mathcal{M}_i)_{i \geq 1} \quad \text{in } \mathbb{M}^{\mathbb{N}}. \quad (1.14)$$

Moreover, the results also hold for $\text{UM}_n(\mathbf{d}, p_c(\lambda))$.

The exact description of the metric space \mathcal{M}_i can be found in [7]. It is worthwhile mentioning a recent work by Conchon-Kerjan and Goldschmidt [15] which is closely related to Theorem 1.4. Conchon-Kerjan and Goldschmidt [15] deduce scaling limits for the vector of components in GHP-topology for critical configuration models having i.i.d power law degrees with exponent $\tau \in (3, 4)$. Their method relies on an alternative approach showing convergence of the height processes corresponding to the components. The associated limiting object was studied in [21], which turns out to have a quite different description than those in [7, 8]. Below, we will see that Assumptions 1 and 2 also hold when degrees are i.i.d samples from a power-law distribution with exponent $\tau \in (3, 4)$.

Scaling limit of maximal distances. For any metric space (X, d) and a point $x \in X$, define the radius of x in X and the diameter of X by

$$\text{Rad}(x, X) = \sup_{y \in X} d(x, y) \quad \text{and} \quad \text{diam}(X) = \sup_{x \in X} \text{Rad}(x, X) = \sup_{x, y \in X} d(x, y). \quad (1.15)$$

An important corollary of Theorem 1.4 is the convergence of the radius and the diameter of the critical components: Let $V_{n,i}$ be a uniformly chosen vertex in $\mathcal{C}_{(i)}(p_c(\lambda))$, where $(V_{n,i})_{i \geq 1}$ is an independent collection conditionally on $(\mathcal{C}_{(i)}(p_c(\lambda)))_{i \geq 1}$. Similarly, using the notation of the scaling limits in Theorem 1.4, let V_i be chosen from M_i according to the measure μ_i and let $(V_i)_{i \geq 1}$ be an independent collection conditionally on $(\mathcal{M}_i)_{i \geq 1}$.

Corollary 1.5 (Convergence of radius and diameter). *Under Assumptions 1, 2 and (1.13), as $n \rightarrow \infty$,*

$$\begin{aligned} (n^{-\eta} \text{Rad}(V_{n,i}, \mathcal{C}_{(i)}(p_c(\lambda))))_{i \geq 1} &\xrightarrow{d} (\text{Rad}(V_i, \mathcal{M}_i))_{i \geq 1}, \\ (n^{-\eta} \text{diam}(\mathcal{C}_{(i)}(p_c(\lambda))))_{i \geq 1} &\xrightarrow{d} (\text{diam}(\mathcal{M}_i))_{i \geq 1}, \end{aligned} \quad (1.16)$$

with respect to the product topology, where $(\mathcal{M}_i)_{i \geq 1}$ is given by Theorem 1.4. Moreover, the result also holds for $\text{UM}_n(\mathbf{d})$.

Proving scaling limits for the diameter of the critical tree-like objects is often a difficult task. In [32], Szekeres proved that, for the uniform random rooted labelled tree on m vertices, the diameter, rescaled by \sqrt{m} , converges in distribution. Szekeres also provided an explicit formula for the density of the limiting distribution in [32, Page 395, (12)]. Szekeres' method was based on generating functions. Łuczak [28] also considered enumeration of trees with diameter $\gg \sqrt{m}$. On the other hand, Aldous [3] (see [3, Section 3.4]) noted that the GHP-convergence can be used as an effective tool to prove scaling limit results for the diameter. This is the motivating idea behind Corollary 1.5. Aldous [3] also raised a natural question whether it is possible to obtain an explicit formula from a result such as Corollary 1.5. In a recent paper, Wang [33] showed that it is indeed possible to get such a formula for the Brownian tree. In the context of Corollary 1.5, the difficulty is two-fold: First, the critical components have surplus edges. For the scaling limits of critical Erdős-Rényi random graphs, Miermont and Sen [29] recently gave a breadth-first construction, which yields an alternative description of the scaling limit of the radius function from a fixed point (rescaled by $n^{1/3}$). However, the description for the diameter and an explicit formula such as the one by Wang [33] is still an open question. Second, the scaling limit in Corollary 1.5 is in the heavy-tailed universality class. Even for \mathbf{p} -trees (see [14]) that satisfy $p_i / (\sum_i p_i^2)^{1/2} \rightarrow \beta_i > 0$, with $(\beta_i)_{i \geq 1} \in \ell_{\downarrow}^2 \setminus \ell_{\downarrow}^1$, obtaining an explicit description for the distribution of the diameter is an interesting question.

Compactness of the limiting metric space. Establishing the almost sure compactness of random metric spaces in the heavy-tailed setting (such as \mathcal{M}_i) is not straightforward, and in fact

the compactness may not hold for all $\theta \in \ell_\downarrow^3 \setminus \ell_\downarrow^2$. It is interesting to find an explicit criterion for the compactness of the limiting objects \mathcal{M}_i in terms of the underlying parameters.

Indeed, in the context of compactness of limits of \mathbf{p} -trees, Aldous, Miermont and Pitman [4, Section 7] states an additional condition, which was conjectured to be necessary and sufficient for the compactness of limits of \mathbf{p} -trees. In the context of critical random graphs, a recent paper by Broutin, Duquesne, and Wang [13] shows that the following criterion, analogous to [4], is sufficient for the almost sure compactness of \mathcal{M}_i :

$$\int_1^\infty \frac{du}{\Psi_\theta(u)} < \infty, \quad \text{where} \quad \Psi_\theta(u) = \sum_{i \geq 1} \theta_i (e^{-u\theta_i} - 1 + u\theta_i). \quad (1.17)$$

Our GHP convergence indirectly yields a sufficient condition for the compactness of the limiting metric space almost surely, given as follows: $\theta \in \ell_\downarrow^3 \setminus \ell_\downarrow^2$,

$$\inf_i \theta_{i+1}^{\tau-2} \sum_{j=1}^i \theta_j > 0, \quad \text{and} \quad \sum_{i \geq 1} \theta_i e^{-\varepsilon \sum_{j \leq i} \theta_j^2} < \infty, \quad \forall \varepsilon > 0. \quad (1.18)$$

To see the first condition, take $d_{i+1} < l \leq d_i$, where i is fixed, and note that $\mathbb{P}(D_n^* \geq l) = \frac{1}{\ell_n} \sum_{j=1}^i d_j \geq c_0 d_{i+1}^{-(\tau-2)}$ and that $n^{-\alpha} d_i \rightarrow \theta_i$. The second condition is an asymptotic version of (1.10). The condition in (1.18) is weaker than assuming $\theta_i = i^{-\alpha}$ as in [8]. For example, $\theta_i \geq i^{-\alpha}$ satisfies (1.18). Let us now show that only the first condition in (1.18) implies the condition in (1.17). Fix $M \geq 1$ such that $\theta_{i+1}^{\tau-2} \sum_{j=1}^i \theta_j \geq 1/M$ for all $i \geq 1$. Now, for $u \in (\frac{1}{\theta_i}, \frac{1}{\theta_{i+1}}]$,

$$\begin{aligned} \Psi_\theta(u) &\geq C \left[\sum_{k=1}^i u\theta_k^2 + \sum_{k=i+1}^\infty u^2\theta_k^2 \right] \geq Cu\theta_{i+1} \sum_{k=1}^i \theta_k \\ &\geq \frac{Cu\theta_{i+1}^{\tau-2} \sum_{k=1}^i \theta_k}{\theta_i^{\tau-3}} \geq \frac{Cu}{M\theta_i^{\tau-3}} \geq \frac{Cu^{\tau-2}}{M}, \end{aligned} \quad (1.19)$$

where $C > 0$ is a constant. Thus, $\int_{\theta_1^{-1}}^\infty \frac{du}{\Psi_\theta(u)} \leq \frac{C}{M} \int_1^\infty u^{-(\tau-2)} du < \infty$ (since $\tau > 3$), which yields (1.17).

Remark 2. The discussion about the compactness of the limiting metric space shows that the GHP-convergence cannot possibly be established under Assumption 1 from [7], because that would imply the almost sure compactness of the limiting metric space only assuming that $\theta \in \ell_\downarrow^3 \setminus \ell_\downarrow^2$. To see this, note that (1.17) is strictly stronger than assuming $\theta \in \ell_\downarrow^3 \setminus \ell_\downarrow^2$, which can be seen for example by taking $\theta_i = i^{-1/2}$. Indeed, for $u \in (\theta_2^{-1}, \infty)$, let $i_0 = i_0(u)$ be such that $\theta_{i_0}^{-1} < u \leq \theta_{i_0+1}^{-1}$, i.e., $\sqrt{i_0} < u \leq \sqrt{i_0+1}$. Then, as $u \rightarrow \infty$,

$$\Psi_\theta(u) \leq C \left[\sum_{i \leq i_0} \theta_i(u\theta_i) + \sum_{i > i_0} \theta_i(u\theta_i)^2 \right] = C \left[u \sum_{i \leq u^2} \frac{1}{i} + u^2 \sum_{i > u^2} \frac{1}{i^{3/2}} \right] \leq C[u \log u + u], \quad (1.20)$$

and thus (1.17) cannot hold.

Verifying the assumptions in the power-law case. Assumption 1 is the general set of assumptions on the degree distribution under which the scaling limit for the component sizes, surplus edges and the metric structure of critical configuration model was identified in [7, 16] in the heavy-tailed universality class. As important examples, these assumptions were shown to apply to configuration models with a power-law degree distribution with exponent $\tau \in (3, 4)$ [16, Section 2]. More precisely, if F is a distribution function on the non-negative integers satisfying $[1 - F](x) = (1 + o(1))Cx^{-(\tau-1)}$ as $x \rightarrow \infty$, then Assumptions 1 is satisfied when (i) $d_i = [1 - F]^{-1}(i/n)$, and when (ii) d_i are the order statistics of an i.i.d. sample from F (we add a dummy half-edge to vertex 1 if $\sum_{i \in [n]} d_i$ is odd). Let us now show

that Assumption 2 also holds in both settings. For case (i), let $G = 1 - F$ and note that $\frac{1}{n} \sum_{i \in [n]} G^{-1}(i/n) = \int_0^1 G^{-1}(x) dx + o(1) = \mathbb{E}[D] + o(1)$, where D has distribution function F . Similarly,

$$\begin{aligned} \frac{1}{n} \sum_{i \in [n]} G^{-1}(i/n) \mathbb{1} \{G^{-1}(i/n) \geq l\} &= \int_0^{G(l)} G^{-1}(x) dx + o(1) \\ &= \int_l^\infty z dF(z) + o(1) = \mathbb{E}[D] \mathbb{P}(D^* \geq l) + o(1), \end{aligned} \quad (1.21)$$

where D^* is the size-biased version of D , i.e., $\mathbb{P}(D^* = k) = k\mathbb{P}(D = k)/\mathbb{E}[D]$ for all $k \geq 1$. Assumption 2 (i) follows using the fact that D has a power-law distribution with exponent τ . Next, note that

$$d_i = \left(\frac{n^\alpha C_F^\alpha}{i^\alpha} \right) (1 + o(1)), \quad (1.22)$$

uniformly over $i = o(n)$ (see [16, Section 2.1]). Taking $K_n \gg (\log(n))^{1/(1-2\alpha)}$, it follows that

$$n^{-2\alpha} \sum_{i \leq K_n} d_i^2 = C(1 + o(1)) \sum_{i \leq K_n} i^{-2\alpha} \gg \log(n). \quad (1.23)$$

Moreover, (1.22), together with the fact that $n^{-2\alpha} \sum_{j < i} d_j^2 = C(1 + o(1))i^{1-2\alpha}$ uniformly over $i \leq K_n$, verifies Assumption 2 (iii).

For case (ii), consider the following construction: Let $(E_j)_{j \geq 1}$ be i.i.d. Exponential(1) random variables, let $\Gamma_i = \sum_{j=1}^i E_j$ be a Gamma random variable, and define $d_i = (1-F)^{-1}(\Gamma_i/\Gamma_{n+1})$. Then $(d_i)_{i \in [n]}$ have the same distribution as the order-statistics of an i.i.d. sample from F (see [12, Section 13.6]). Thus,

$$d_i = \left(\frac{\Gamma_{n+1}^\alpha C_F^\alpha}{\Gamma_i^\alpha} \right) (1 + o(1)), \quad (1.24)$$

uniformly over $i = o(n)$ (see [16, Section 2.1]). Moreover, $\Gamma_i/i \rightarrow 1$ almost surely as $i \rightarrow \infty$ by the law of large numbers. Thus, for any $\varepsilon > 0$, there exists $M = M(\varepsilon) > 0$ such that

$$\mathbb{P}(\exists i \geq 1 : \Gamma_i \notin [i/M, iM]) < \varepsilon. \quad (1.25)$$

On the event $\Gamma_i \in [i/M, iM]$, we can verify Assumption 2 using identical argument as in case (i).

Proof ideas and technical motivation for this work. The proof of Theorem 1.1 consists of two main steps, that form the key ideas in the argument. The first step is to show that the neighborhoods of the high-degree vertices, called *hubs*, have mass $\Theta_{\mathbb{P}}(n^\rho)$. Secondly, the probability of any small εn^η neighborhood not containing hubs is small. These two facts, summarized in Propositions 2.1 and 2.2 below, together ensure that the total mass of any neighborhood of $\mathcal{C}_{(i)}$ of radius εn^η is bounded away from zero. These two facts were proved in [8] in the context of rank-one inhomogeneous random graphs. However, the proof techniques are completely different here. The main advantage in [8] was that the breadth-first exploration of components could be dominated by a branching process with *mixed Poisson* progeny distribution that is *independent of n* . This allows one to use existing literature to estimate the probabilities that a long path exists in the branching process. However, such a technique is specific to rank-one inhomogeneous random graphs and does not work in the cases where the above stochastic domination does not hold. This was one of the technical motivations for this work. Moreover, the final section contains results about exponential tail-bounds for the number of edges in large critical components (Proposition 4.1), as well as a coupling of the neighborhood exploration with a branching process with stochastically larger progeny distribution (Section 4.2), which are both interesting in their own right.

Organization of this paper. The rest of this paper is organized as follows: In Section 2, we state two key propositions, the first involving the total mass of small neighborhoods, and the second involving a bound on the diameter of a *slightly subcritical* $\text{CM}_n(\mathbf{d})$. The proof of Theorem 1.1 is completed in Section 2. In Section 3, we derive the required bounds on the total mass of small neighborhoods. In Section 4, we obtain bounds on the diameter of the connected components after removing the high-degree vertices.

2 Proof of the global lower mass-bound

In this section, we first state the two key ingredients in Propositions 2.1 and 2.2, and then complete the proof of Theorem 1.1. The proofs of Propositions 2.1 and 2.2 are given in the subsequent sections. The first ingredient shows that hub i has sufficient mass close to it with high probability:

Proposition 2.1. *Assume that Assumptions 1 and (1.6) hold. Recall that $\mathcal{N}_v(\delta)$ denotes the δn^η neighborhood of v . For each fixed $i \geq 1$ and $\varepsilon_2 > 0$, there exists $\delta_{i,\varepsilon_2} > 0$ and $n_{i,\varepsilon_2} \geq 1$ such that, for any $\delta \in (0, \delta_{i,\varepsilon_2}]$ and $n \geq n_{i,\varepsilon_2}$,*

$$\mathbb{P}(|\mathcal{N}_i(\delta)| \leq \theta_i \delta n^\rho) \leq \frac{\varepsilon_2}{2^{i+1}}. \quad (2.1)$$

Next, we need some control on the diameter of the graph after removing the hubs. Denote by $\mathcal{G}_n^{>K}$ the graph obtained by removing the vertices $[K] = \{1, \dots, K\}$ having the largest degrees and the edges incident to them from $\text{CM}_n(\mathbf{d})$. Note that $\mathcal{G}_n^{>K}$ is a configuration model conditionally on its degree sequence. Let $\Delta^{>K}$ denote the maximum of the diameters of the connected components of $\mathcal{G}_n^{>K}$. The following proposition shows that, for large K , $\Delta^{>K}$ is small with high probability:

Proposition 2.2. *Assume that Assumptions 1, 2 and (1.6) hold. Then, for any $\varepsilon_1, \varepsilon_2 > 0$, there exists $K = K(\varepsilon_1, \varepsilon_2)$ and $n_0 = n_0(\varepsilon_1, \varepsilon_2)$ such that for all $n \geq n_0$,*

$$\mathbb{P}(\Delta^{>K} > \varepsilon_1 n^\eta) \leq \frac{\varepsilon_2}{4}. \quad (2.2)$$

We now prove Theorem 1.1 assuming Propositions 2.1 and 2.2:

Proof of Theorem 1.1. Fix $i \geq 1$ and $\varepsilon_1, \varepsilon_2 > 0$. For a component $\mathcal{C} \subset \text{CM}_n(\mathbf{d})$, we write $\Delta(\mathcal{C})$ to denote its diameter. Let us choose K and n_0 so that (2.2) holds for all $n \geq n_0$. In view of Proposition 2.1, let $\delta_0 = \min\{\varepsilon_1, \delta_{1,\varepsilon_2}, \dots, \delta_{K,\varepsilon_2}\}/2$, and $n'_0 = \max\{n_0, n_{1,\varepsilon_2}, \dots, n_{K,\varepsilon_2}\}$. Thus, for all $n \geq n'_0$, (2.1) is satisfied for all $i \in [K]$. Define

$$F_1 := \{\Delta^{>K} < \varepsilon_1 n^\eta / 2\}, \quad F_2 := \{\Delta(\mathcal{C}_{(i)}) > \varepsilon_1 n^\eta / 2\}. \quad (2.3)$$

Notice that, on the event $F_1 \cap F_2$, it must be the case that one of the vertices in $[K]$ belongs to $\mathcal{C}_{(i)}$, and that the union of the neighborhoods of $[K]$ of radius $\lceil \varepsilon_1 n^\eta / 2 \rceil + 1 \approx \varepsilon_1 n^\eta / 2$ covers $\mathcal{C}_{(i)}$. Therefore, given any vertex $v \in \mathcal{C}_{(i)}$, $\mathcal{N}_v(\varepsilon_1)$ contains at least one of the neighborhoods $(\mathcal{N}_j(\varepsilon_1/2))_{j \in [K]}$. This observation yields that

$$\inf_{v \in \mathcal{C}_{(i)}} n^{-\rho} |\mathcal{N}_v(\varepsilon_1)| \geq \min_{j \in [K]} n^{-\rho} |\mathcal{N}_j(\varepsilon_1/2)| \geq \min_{j \in [K]} n^{-\rho} |\mathcal{N}_j(\delta_0)|. \quad (2.4)$$

Thus, for all $n \geq n'_0$

$$\begin{aligned} & \mathbb{P}\left(F_1 \cap F_2 \cap \left\{ \inf_{v \in \mathcal{C}_{(i)}} n^{-\rho} |\mathcal{N}_v(\varepsilon_1)| \leq \theta_K \delta_0 \right\}\right) \\ & \leq \sum_{j \in [K]} \mathbb{P}(|\mathcal{N}_j(\delta)| \leq \theta_j \delta_0 n^\rho) \leq \sum_{j=1}^K \frac{\varepsilon_2}{2^{j+1}} \leq \frac{\varepsilon_2}{2}, \end{aligned} \quad (2.5)$$

where the one-but-last step follows from Proposition 2.1. Further, on the event F_2^c , $|\mathcal{N}_v(\varepsilon_1)| = |\mathcal{C}_{(i)}|$ for all $v \in \mathcal{C}_{(i)}$. Moreover, using (1.8), it follows that $n^{-\rho}|\mathcal{C}_{(i)}|$ converges in distribution to a random variable with strictly positive support. Using the Portmanteau theorem, the above implies that for any $\delta'_0 > 0$, there exists $\tilde{n}_0 = \tilde{n}_0(\varepsilon_2, \delta'_0)$ such that, for all $n \geq \tilde{n}_0$,

$$\mathbb{P}(n^{-\rho}|\mathcal{C}_{(i)}| \leq \delta'_0) \leq \frac{\varepsilon_2}{4}. \quad (2.6)$$

Therefore,

$$\mathbb{P}\left(F_2^c \cap \left\{ \inf_{v \in \mathcal{C}_{(i)}} n^{-\rho}|\mathcal{N}_v(\varepsilon_1)| \leq \delta'_0 \right\}\right) \leq \frac{\varepsilon_2}{4}. \quad (2.7)$$

Now, using (2.5), (2.7) together with Proposition 2.2, it follows that, for any $n \geq \max\{n'_0, \tilde{n}_0\}$, and K chosen as above,

$$\mathbb{P}\left(\inf_{v \in \mathcal{C}_{(i)}} n^{-\rho}|\mathcal{N}_v(\varepsilon_1)| \leq \min\{\delta'_0, \theta_K \delta_0\}\right) \leq \varepsilon_2. \quad (2.8)$$

This completes the proof of Theorem 1.1. \square

3 Lower bound on the total mass of neighborhoods of hubs

In this section, we prove Proposition 2.1.

Proof of Proposition 2.1. Let us denote the component of $\text{CM}_n(\mathbf{d})$ containing vertex i by $\mathcal{C}(i)$. Consider the breadth-first exploration of $\mathcal{C}(i)$ starting from vertex i , given by the following exploration algorithm [16]:

Algorithm 1 (Exploring the graph). The algorithm carries along vertices that can be alive, active, exploring and killed, and half-edges that can be alive, active or killed. We sequentially explore the graph as follows:

- (S0) At stage $l = 0$, all the vertices and the half-edges are *alive*, and only the half-edges associated to vertex i are *active*. Also, there are no *exploring* vertices except i .
- (S1) At each stage l , take an active half-edge e of an exploring vertex v and pair it uniformly to another alive half-edge f . Kill e, f . If f is incident to a vertex v' that has not been discovered before, then declare all the half-edges incident to v' (if any) active, except f . If $\text{degree}(v') = 1$ (i.e. the only half-edge incident to v' is f) then kill v' . Otherwise, declare v' to be active and larger than all other vertices that are alive. After killing e , if v does not have another active half-edge, then kill v also.
- (S2) Repeat (S1) until there is no active half-edges left.

Call a vertex *discovered* if it is either active or killed. Let \mathcal{V}_l denote the set of vertices discovered up to time l and $\mathcal{I}_j^n(l) := \mathbb{1}\{j \in \mathcal{V}_l\}$. Define the exploration process by

$$S_n(l) = d_i + \sum_{j \neq i} d_j \mathcal{I}_j^n(l) - 2l = d_i + \sum_{j \neq i} d_j \left(\mathcal{I}_j^n(l) - \frac{d_j}{\ell_n} l \right) + \left(\frac{1}{\ell_n} \sum_{j \neq i} d_j^2 - 2 \right) l. \quad (3.1)$$

Note that the exploration process keeps track of the number of active half-edges. Thus, $\mathcal{C}(i)$ is explored when S_n hits zero. Moreover, since one edge is explored at each step, the hitting time of zero is the total number of edges in $\mathcal{C}(i)$. Define the re-scaled version \bar{S}_n of S_n by $\bar{S}_n(t) = n^{-\alpha} S_n(\lfloor tn^\rho \rfloor)$. Then, by Assumption 1 and (1.6),

$$\bar{S}_n(t) = \theta_i - \frac{\theta_i^2 t}{\mu} + n^{-\alpha} \sum_{j \neq i} d_j \left(\mathcal{I}_j^n(tn^\rho) - \frac{d_j}{\ell_n} tn^\rho \right) + \lambda t + o(1). \quad (3.2)$$

The convergence of this exploration process was considered in [16, Theorem 8] except for the fact that the exploration process started at zero in [16]. However, using identical arguments to [16, Theorem 8], it can be shown that

$$\bar{S}_n \xrightarrow{d} S_\infty, \quad (3.3)$$

with respect to the Skorohod J_1 -topology, where

$$S_\infty(t) = \theta_i - \frac{\theta_i^2 t}{\mu} + \sum_{j \neq i} \theta_j \left(\mathcal{I}_j(t) - \frac{\theta_j t}{\mu} \right) + \lambda t, \quad (3.4)$$

with $\mathcal{I}_j(s) := \mathbb{1} \{ \xi_j \leq s \}$ and $\xi_j \sim \text{Exponential}(\theta_j/\mu)$ independently of each other.

Let $h_n(u)$ (respectively $h_\infty(u)$) denote the first hitting time of \bar{S}_n (respectively S_∞) of u . More precisely,

$$h_n(u) := \inf \left\{ t : \bar{S}_n(t) \leq u \text{ or } \lim_{t' \nearrow t} \bar{S}_n(t') \leq u \right\}, \quad (3.5)$$

and define $h_\infty(u)$ similarly by replacing $\bar{S}_n(t)$ by $\bar{S}_\infty(t)$. Note that using [16, Lemma 36], the distribution of $h_\infty(u)$ does not have any atoms and therefore, for any $\varepsilon_2 > 0$, there exists $\beta_{\varepsilon_2, i} > 0$ such that

$$\mathbb{P}(h_\infty(\theta_i/2) \leq \beta_{\varepsilon_2, i}) \leq \frac{\varepsilon_2}{2^{i+1}}.$$

Now we use the following fact:

Fact 1. *Let $(X_n(t))_{t \geq 0} \xrightarrow{d} (X(t))_{t \geq 0}$ in Skorohod J_1 -topology and let $h(X_n)$ (respectively $h(X)$) denote the hitting time to zero of X_n (respectively X). Then, $\liminf_{n \rightarrow \infty} \mathbb{P}(h(X_n) > a) \geq \mathbb{P}(h(X) > a)$, for all $a > 0$.*

Proof. Let $(f_n)_{n \geq 1}$ be such that $h(f_n) \leq a$ for all $n \geq 1$ and $f_n \rightarrow f$ in Skorohod J_1 -topology as $n \rightarrow \infty$. Now, $h(f_n) \leq a$ implies that $\inf_{t \in [0, a]} f_n(t) = 0$. Using [34, Theorem 13.4.1], it follows that $\inf_{t \in [0, a]} f(t) = 0$ and thus $h(f) \leq a$. Thus, we have shown that $\{f : h(f) \leq a\}$ is a closed set in the Skorohod J_1 -topology, and therefore $\{f : h(f) > a\}$ is an open set. The proof follows using the Portmanteau theorem [9, Theorem 2.1 (iv)]. \square

Using (3.3) and Fact 1, there exists $n_{i, \varepsilon_2} \geq 1$ such that, for all $n \geq n_{i, \varepsilon_2}$,

$$\mathbb{P}(h_n(\theta_i/2) \leq n^\rho \beta_{\varepsilon_2, i}) \leq \frac{\varepsilon_2}{2^i}. \quad (3.6)$$

Firstly the goal is to show that there exists a $\delta_{i, \varepsilon}$ such that for any $\delta \in (0, \delta_{i, \varepsilon}]$,

$$\sum_{k \in \mathcal{N}_i(\delta)} d_k \leq \theta_i \delta n^\rho \implies h_n(\theta_i/2) \leq n^\rho \beta_{\varepsilon_2, i}. \quad (3.7)$$

Recall that $\mathcal{N}_v(\delta)$ denotes the δn^η neighborhood of v in $\text{CM}_n(\mathbf{d})$. To prove (3.7), let $\partial(j)$ denote the set of vertices at distance j from i . Let E_{j1} denote the total number of edges between vertices in $\partial(j)$ and $\partial(j-1)$, and let E_{j2} denote the number of edges within $\partial(j-1)$. Define $E_j = E_{j1} + E_{j2}$. Fix any $\delta < 2\beta_{\varepsilon_2, i}/\theta_i$. Note that if $\sum_{k \in \mathcal{N}_i(\delta)} d_k \leq \theta_i \delta n^\rho$, then the total number of edges in $\mathcal{N}_i(\delta)$ is at most $\theta_i \delta n^\rho / 2$. Thus there exists $j \leq \delta n^\eta$ such that $E_j \leq \theta_i \delta n^\rho / 2\delta n^\eta = \theta_i n^\alpha / 2$. This implies that S_n must go below $\theta_i n^\alpha / 2$ before exploring all the vertices in $\mathcal{N}_i(\delta)$. This is because we are exploring the components in a breadth-first manner and \bar{S}_n keeps track of the number of active half-edges which are the potential connections to vertices at the next level. Since one edge is explored in each time step, and we rescale time by n^ρ , this implies that

$$h_n(\theta_i/2) \leq \frac{1}{2} n^{-\rho} \sum_{k \in \mathcal{N}_i(\delta)} d_k \leq \theta_i \delta / 2 \leq \beta_{\varepsilon_2, i}. \quad (3.8)$$

Therefore, for all $n \geq n_{i,\varepsilon_2}$,

$$\mathbb{P}\left(\sum_{k \in \mathcal{N}_i(\delta)} d_k \leq \theta_i \delta n^\rho\right) \leq \mathbb{P}(h_n(\theta_i/2) \leq \beta_{\varepsilon_2,i}) \leq \frac{\varepsilon_2}{2^i}. \quad (3.9)$$

Finally, to conclude Proposition 2.1 from (3.9), we use the following result from [16, Lemma 22]: For any $T > 0$,

$$\sup_{u \leq T} \left| \sum_{i \in [n]} \mathcal{I}_i^n(un^\rho) - un^\rho \right| = o_{\mathbb{P}}(n^\rho). \quad (3.10)$$

This implies that the difference between the number of edges and the number of vertices explored upto time un^ρ is $o_{\mathbb{P}}(n^\rho)$ uniformly over $u \leq T$. The proof of Proposition 2.1 now follows. \square

4 Diameter after removing hubs

Throughout the remainder of the paper, we fix the convention that $C, C', C'' > 0$ etc. denote constants whose value can change from line to line. Recall the definition of the graph $\mathcal{G}_n^{>K}$ from Proposition 2.2. If we keep on exploring $\mathcal{G}_n^{>K}$ in a breadth-first manner using Algorithm 1 and ignore the cycles created, then we get a random tree. The idea is to couple neighborhoods in $\mathcal{G}_n^{>K}$ with a suitable branching process such that the progeny distribution of the branching process dominates the number of children of each vertex in the breadth-first tree. Therefore, when there is a long path in $\mathcal{G}_n^{>K}$ that makes the diameter large, that long path must be present in the branching process as well under the above coupling. In this way, the question about the diameter of $\mathcal{G}_n^{>K}$ reduces to the question about the height of a branching process. To estimate the height suitably, we use a recent beautiful proof technique by Addario-Berry [1] which allows one to relate the height of a branching process to the sum of inverses of the associated breadth-first random walk.

In Section 4.1, we establish large deviation bounds for the number of edges within components. This allows us to formulate the desired coupling in Section 4.2. In Section 4.3, we analyze the breadth-first random walk to show that it is unlikely that the height of the branching process is larger than εn^η . These bounds are different from those derived in [1] since our branching process depends on n and there is a joint scaling involved between the distances and the law of the branching process.

4.1 Asymptotics for the number of edges

For a graph G , let $E(G)$ denote the number of edges in G .

Proposition 4.1. *Suppose that Assumption 1 and (1.6) hold. There exists $\varepsilon_0 > 0$ such that for all $\varepsilon \in (0, \varepsilon_0)$, and for all sufficiently large n ,*

$$\mathbb{P}(E(\mathcal{C}(i)) > n^{\rho+\varepsilon}) \leq C e^{-C' n^{\varepsilon/2}}, \quad (4.1)$$

for some absolute constants $C, C' > 0$ and all $i \in [n]$.

The proof of Proposition 4.1 relies on concentration techniques for martingales. We start by defining the relevant notation. Consider exploring $\text{CM}_n(\mathbf{d})$ with Algorithm 1, and let the associated exploration process be defined in (3.1). Let us denote the degree of the vertex found at step l by $d_{(l)}$. If no new vertex is found at step l , then $d_{(l)} = 0$. Also, let \mathcal{F}_l denote the sigma-algebra containing all the information revealed by the exploration process upto time l . Thus,

$$S_n(0) = d_i, \quad \text{and} \quad S_n(l) = S_n(l-1) + (d_{(l)} - 2). \quad (4.2)$$

Using the Doob-Meyer decomposition, one can write

$$S_n(l) = S_n(0) + M_n(l) + A_n(l), \quad (4.3)$$

where M_n is a martingale with respect to $(\mathcal{F}_l)_{l \geq 1}$. The drift A_n and the quadratic variation $\langle M_n \rangle$ of M_n are given by

$$A_n(l) = \sum_{j=1}^l \mathbb{E}[d_{(j)} - 2 | \mathcal{F}_{j-1}], \quad \langle M_n \rangle(l) = \sum_{j=1}^l \text{Var}(d_{(j)} | \mathcal{F}_{j-1}). \quad (4.4)$$

Fix $\varepsilon_0 = (4 - \tau)/(\tau - 1)$. We will show that for any $\varepsilon \in (0, \varepsilon_0)$, the following two lemmas hold:

Lemma 4.2. *Suppose that Assumption 1 and (1.6) hold. For all sufficiently large n ,*

$$\mathbb{P}(n^{-(\alpha+\varepsilon)} M_n(n^{\rho+\varepsilon}) > 1) \leq C e^{-C' n^\varepsilon}, \quad (4.5)$$

for some absolute constants $C, C' > 0$.

Lemma 4.3. *Suppose that Assumption 1 and (1.6) hold. For all fixed $K \geq 1$, and for all sufficiently large n ,*

$$\mathbb{P}\left(n^{-(\alpha+\varepsilon)} A_n(n^{\rho+\varepsilon}) \geq -C \sum_{i=1}^K \theta_i^2\right) \leq C e^{-C' n^{\varepsilon/2}}, \quad (4.6)$$

for some absolute constants $C, C' > 0$.

Proof of Proposition 4.1 subject to Lemmas 4.2, 4.3. Throughout, we write $t_n := n^{\rho+\varepsilon}$. Note that we can choose $K \geq 1$ such that $\sum_{i=1}^K \theta_i^2$ is arbitrarily large since $\theta \notin \ell_v^2$. Thus, if $n^{-(\alpha+\varepsilon)} M_n(t_n) \leq 1$ and $n^{-(\alpha+\varepsilon)} A_n(t_n) \leq -C \sum_{i=1}^K \theta_i^2$, then $n^{-(\alpha+\varepsilon)} S_n(t_n) < 0$, and therefore $\mathcal{C}(i)$ must be explored before time t_n , and thus $\mathbb{E}(\mathcal{C}(i)) \leq t_n$. Thus, Lemmas 4.2 and 4.3 together complete the proof of Proposition 4.1. \square

Proof of Lemma 4.2. First note that $\varepsilon_0 + \rho < 1$ and therefore $t_n = o(n)$. Thus, uniformly over $j \leq t_n$,

$$\text{Var}(d_{(j)} | \mathcal{F}_{j-1}) \leq \mathbb{E}[d_{(j)}^2 | \mathcal{F}_{j-1}] = \frac{\sum_{j \notin \mathcal{V}_{j-1}} d_j^3}{\ell_n - 2j + 2} \leq \frac{\sum_{j \in [n]} d_j^3}{\ell_n - 2t_n + 2} \leq C n^{3\alpha-1}, \quad (4.7)$$

so that, almost surely,

$$\langle M_n \rangle(t_n) \leq t_n C n^{3\alpha-1} = C n^{2\alpha+\varepsilon}. \quad (4.8)$$

Also, $d_{(j)} \leq C n^\alpha$ almost surely. We can now use Freedman's inequality [20, Proposition 2.1] which says that if $Y(k) = \sum_{j \leq k} X_j$ with $\mathbb{E}[X_j | \mathcal{F}_{j-1}] = 0$ (for some filtration $(\mathcal{F}_j)_{j \geq 1}$) and $\mathbb{P}(|X_j| \leq R, \forall j \geq 1) = 1$, then, for any $a, b > 0$,

$$\mathbb{P}(Y(k) \geq a, \text{ and } \langle Y \rangle(k) \leq b) \leq \exp\left(\frac{-a^2}{2(Ra + b)}\right). \quad (4.9)$$

We apply (4.9) with $a = n^{\alpha+\varepsilon}$, $b = C n^{2\alpha+\varepsilon}$ and $R = C n^\alpha$. Note that $\langle M_n \rangle(t_n) \leq b$ with probability 1 by (4.8). It follows that

$$\mathbb{P}(M_n(t_n) > n^{\alpha+\varepsilon}) \leq \exp\left(-\frac{n^{2\alpha+2\varepsilon}}{2(n^\alpha n^{\alpha+\varepsilon} + C n^{2\alpha+\varepsilon})}\right) \leq C e^{-C' n^\varepsilon}, \quad (4.10)$$

and the proof follows. \square

Proof of Lemma 4.3. Note that

$$\begin{aligned}
\mathbb{E}[d_{(i)} - 2 | \mathcal{F}_{i-1}] &= \frac{\sum_{j \notin \mathcal{V}_{i-1}} d_j^2}{\ell_n - 2i + 1} - 2 \\
&= \frac{1}{\ell_n} \sum_{j \in [n]} d_j(d_j - 2) - \frac{1}{\ell_n} \sum_{j \in \mathcal{V}_{i-1}} d_j^2 + \frac{(2i-1) \sum_{j \notin \mathcal{V}_{i-1}} d_j^2}{\ell_n(\ell_n - 2i + 1)} \\
&\leq \lambda n^{-\eta} - \frac{1}{\ell_n} \sum_{j \in \mathcal{V}_{i-1}} d_j^2 + \frac{(2i-1)}{(\ell_n - 2i + 1)^2} \sum_{j \in [n]} d_j^2 + o(n^{-\eta})
\end{aligned} \tag{4.11}$$

uniformly over $i \leq t_n$. Therefore, for all sufficiently large n ,

$$\begin{aligned}
A_n(t_n) &= \sum_{j=1}^{t_n} \mathbb{E}[d_{(j)} - 2 | \mathcal{F}_{j-1}] \leq \lambda t_n n^{-\eta} - \frac{1}{\ell_n} \sum_{i=1}^{t_n} \sum_{j \in \mathcal{V}_{i-1}} d_j^2 + \frac{C t_n^2}{\ell_n} + o(n^{\alpha+\varepsilon}) \\
&= \lambda n^{\alpha+\varepsilon} - \frac{1}{\ell_n} \sum_{i=1}^{t_n} \sum_{j \in \mathcal{V}_{i-1}} d_j^2 + o(n^{\alpha+\varepsilon}),
\end{aligned} \tag{4.12}$$

where in the second step we have used $\sum_{i \in [n]} d_i^2 / \ell_n = O(1)$, and in the last step we have used that $t_n^2 / \ell_n = O(n^{2\rho+2\varepsilon-1}) = o(n^{\alpha+\varepsilon})$ for $\varepsilon < 1 + \alpha - 2\rho = (4 - \tau) / (\tau - 1) = \varepsilon_0$. Let us denote the second term in (4.12) by (A). To analyze (A), define the event

$$\mathcal{A}_n := \{\exists j : d_j > n^{\alpha-\varepsilon/2}, j \notin \mathcal{V}_{t_n/2}\}. \tag{4.13}$$

Thus, for all sufficiently large n ,

$$\mathbb{P}(\mathcal{A}_n) \leq \sum_{j: d_j > n^{\alpha-\varepsilon/2}} \left(1 - \frac{d_j}{\ell_n - 2t_n}\right)^{t_n} \leq n e^{-C n^{\varepsilon/2}}. \tag{4.14}$$

On the event \mathcal{A}_n^c ,

$$(A) = \frac{1}{\ell_n} \sum_{i=1}^{t_n} \sum_{j \in [n]} d_j^2 \mathbb{1}\{j \in \mathcal{V}_{i-1}\} \geq \frac{1}{\ell_n} \sum_{i=\frac{t_n}{2}+1}^{t_n} \sum_{j=1}^K d_j^2 \mathbb{1}\{j \in \mathcal{V}_{i-1}\} \geq C n^{\alpha+\varepsilon} \sum_{j=1}^K \theta_j^2. \tag{4.15}$$

Combining (4.12), (4.14) and (4.15) now completes the proof. \square

4.2 Coupling with branching processes

Recall that $\mathcal{C}(i)$ is the connected component in $\text{CM}_n(\mathbf{d})$ containing vertex i . Define the event $\mathcal{K}_n := \{\mathbb{E}(\mathcal{C}(i)) > n^{\rho+\varepsilon}\}$. Proposition 4.1 implies that the probability of \mathcal{K}_n happening is stretched exponentially small in n . On the event \mathcal{K}_n^c , we can couple the breadth-first exploration starting from vertex i with a suitable branching process. Let n_k denote the number of vertices of degree k and consider the branching process $\mathcal{X}_n(i)$ starting with d_i individuals, and the progeny distribution $\bar{\xi}_n$ given by

$$\mathbb{P}(\bar{\xi}_n = k) = \bar{p}_k = \begin{cases} \frac{(k+1)n_{k+1}}{\ell_n} & \text{for } k \geq 1, \\ \frac{n_1 - 2n^{\rho+\varepsilon}}{\ell_n} & \text{for } k = 0, \end{cases} \tag{4.16}$$

where $\ell_n = \ell_n - 2n^{\rho+\varepsilon}$. Note that, at each step of the exploration, we have at most $(k+1)n_{k+1}$ half-edges that are incident to vertices having k further unpaired half-edges. Further, on the event \mathcal{K}_n^c , we have at least ℓ_n choices for pairing. Therefore, the number of active

half-edges discovered at each step in the breadth-first exploration of the neighborhoods of i is stochastically dominated by $\bar{\xi}_n$. This proves the next proposition, which we state after setting up some further notation. Recall that $\mathcal{G}_n^{>i-1}$ denotes the graph obtained by deleting vertices in $[i-1]$ and the associated edges from $\text{CM}_n(\mathbf{d})$. Let $\partial_i(r)$ denote the number of vertices at distance r from vertex i in the graph $\mathcal{G}_n^{>i-1}$. Let $\bar{\xi}_n(i)$ denote the random variable with the distribution in (4.16) truncated in such a way that $\{d_1, \dots, d_{i-1}\}$ are excluded from the support. More precisely,

$$\mathbb{P}(\bar{\xi}_n(i) = k) = \begin{cases} 0 & \text{for } k > d_i, \\ \frac{(k+1) \#\{j \geq i : d_j = k+1\}}{L} & \text{for } 1 \leq k \leq d_i, \\ \frac{n_1 - 2n^{\rho+\varepsilon}}{L} & \text{for } k = 0, \end{cases} \quad (4.17)$$

where $L = \ell_n - \sum_{j=1}^{i-1} d_j$ is the appropriate normalizing constant. Let $\mathcal{X}_{n,\text{res}}(i)$ denote the branching process starting with d_i individuals and progeny distribution $\bar{\xi}_n(i)$ and let $\bar{\partial}_i(r)$ denote the number of individuals in generation r of $\mathcal{X}_{n,\text{res}}(i)$. Then the above stochastic domination argument immediately yields the next proposition:

Proposition 4.4. *Suppose that Assumption 1 and (1.6) hold. Let K_n be as described in Assumption 2. For all $r \geq 1$, $i \leq K_n$, and $n \geq 1$,*

$$\mathbb{P}(\partial_i(r) \neq \emptyset) \leq \mathbb{P}(\bar{\partial}_i(r) \neq \emptyset) + \mathbb{P}(\mathbb{E}(\mathcal{C}(i)) > n^{\rho+\varepsilon}). \quad (4.18)$$

Before proceeding with the next section in which we investigate $\mathbb{P}(\bar{\partial}_i(r) \neq \emptyset)$, we estimate the expectation and variance of the branching process $\mathcal{X}_{n,\text{res}}(i)$ using Assumptions 1, 2 and (1.6). Using $\sum_i \theta_i^2 = \infty$, we can choose $i_2(\lambda)$ (depending only on λ) such that

$$\frac{1}{\mu} \sum_{i=1}^{i_2(\lambda)} \theta_i^2 \geq 5\lambda. \quad (4.19)$$

Also note that the normalizing constant in (4.17) satisfies $L = \ell_n(1 + o(n^{-\eta}))$ uniformly over $1 \leq i \leq K_n$, since $\sum_{j \leq i} d_j = o(n^{2\alpha})$ by Assumption 2 and $2\alpha - 1 = -\eta$. Now using Assumption 1 (iii), note that there exists $N_\lambda \geq 1$ such that for all $n \geq N_\lambda$ and $i_2(\lambda) \leq i \leq K_n$,

$$\begin{aligned} \mathbb{E}[\bar{\xi}_n(i)] &= \frac{1}{L} \sum_{j \geq i} d_j(d_j - 1) = \frac{1}{\ell_n} \sum_{j \geq i} d_j(d_j - 1) + o(n^{-\eta}) \\ &= 1 + \lambda n^{-\eta} - \frac{1}{\ell_n} \sum_{j < i} d_j(d_j - 1) + o(n^{-\eta}) \\ &\leq 1 + \lambda n^{-\eta} - \frac{1}{2\ell_n} \sum_{j < i} d_j^2 + o(n^{-\eta}) \\ &\leq 1 - \left(C n^{-2\alpha} \sum_{j < i} d_j^2 \right) n^{-\eta} + o(n^{-\eta}), \end{aligned} \quad (4.20)$$

where the third step uses (1.6), the penultimate step uses the fact that $d_i \geq 2$ so that $\sum_{j < i} d_j \leq \sum_{j < i} d_j^2/2$ for $i \leq K_n$, and the last step uses (4.19). Thus, for $n \geq N_\lambda$ and $i_2(\lambda) \leq i \leq K_n$,

$$\mathbb{E}[\bar{\xi}_n(i)] \leq 1 - \beta_i^n n^{-\eta} \quad \text{where} \quad \beta_i^n = C n^{-2\alpha} \sum_{j < i} d_j^2. \quad (4.21)$$

The estimate in (4.21) will be crucial in the next section.

4.3 Estimating heights of trees via random walks

We will prove the following theorem in this section:

Theorem 4.5. *Suppose that Assumptions 1, 2 and (1.6) hold. Fix $\varepsilon > 0$. Then, for all $i_2(\lambda) \leq i \leq K_n$ (where $i_2(\lambda)$ is given by (4.19)) and $n \geq N_\lambda$,*

$$\mathbb{P}(\bar{\partial}_i(\varepsilon n^\eta) \neq \emptyset) \leq \frac{C d_i}{n^\alpha} e^{-\frac{\varepsilon \beta_i^n}{2}}, \quad (4.22)$$

for some constant $C = C(\varepsilon, \lambda) > 0$.

Define $\mathcal{X}_n^1(i)$ to be the Galton-Watson tree starting with one offspring and progeny distribution $\bar{\xi}_n(i)$ and let $\bar{\partial}_i^1(r)$ denote the number of individuals in generation r of $\mathcal{X}_n^1(i)$. The crucial ingredient for the proof of Theorem 4.5 is the following:

Proposition 4.6. *Under identical conditions as in Theorem 4.5, for all $n \geq N_\lambda$,*

$$\mathbb{P}(\bar{\partial}_{i_2(\lambda)}^1(\varepsilon n^\eta) \neq \emptyset) \leq \frac{C}{n^\alpha}, \quad (4.23)$$

for some constant $C = C(\varepsilon, \lambda) > 0$.

Proof of Theorem 4.5 using Proposition 4.6. Let M_r denote the number of children at generation r of $\mathcal{X}_{n, \text{res}}(i)$, and note that

$$\mathbb{P}(\bar{\partial}_i(\varepsilon n^\eta) \neq \emptyset) \leq \mathbb{E}[M_{\varepsilon n^\eta/2}] \times \mathbb{P}(\bar{\partial}_i^1(\varepsilon n^\eta/2) \neq \emptyset). \quad (4.24)$$

Now, using (4.21),

$$\mathbb{E}[M_{\varepsilon n^\eta/2}] \leq d_i (1 - \beta_i^n n^{-\eta})^{\varepsilon n^\eta/2} \leq d_i e^{-\frac{\varepsilon \beta_i^n}{2}}, \quad (4.25)$$

and $\bar{\xi}_n(i) \preceq \bar{\xi}_n(i-1) \preceq \dots \preceq \bar{\xi}_n(i_2(\lambda))$, where \preceq denotes stochastic domination. Thus,

$$\mathbb{P}(\bar{\partial}_i(\varepsilon n^\eta) \neq \emptyset) \leq d_i e^{-\frac{\varepsilon \beta_i^n}{2}} \times \mathbb{P}(\bar{\partial}_{i_2(\lambda)}^1(\varepsilon n^\eta/2) \neq \emptyset), \quad (4.26)$$

and the proof of Theorem 4.5 follows using Proposition 4.6. \square

The rest of this section is devoted to the proof of Proposition 4.6. We leverage some key ideas from [1]. Define the breadth-first random walk by

$$s_n(u) = s_n(u-1) + \zeta_u - 1, \quad s_n(0) = 1, \quad (4.27)$$

where $(\zeta_u)_{u \geq 1}$ are i.i.d. observations from the distribution of $\bar{\xi}_n(i_2(\lambda))$. Define the function

$$H_n(t) = \sum_{u \in [0, t)} \frac{1}{s_n(u)}, \quad (4.28)$$

and $\sigma = \inf\{u : s_n(u) = 0\}$. A remarkable fact, which was observed in [1, Proposition 1.7], states that the height of a tree with breadth-first exploration process s_n is at most $3H_n(\sigma)$. Thus Proposition 4.6 can be concluded directly from the following estimate:

Proposition 4.7. *Under identical conditions as in Theorem 4.5, for all $n \geq N_\lambda$,*

$$\mathbb{P}(H_n(\sigma) > \varepsilon n^\eta) \leq \frac{C}{n^\alpha}, \quad (4.29)$$

for some constant $C = C(\varepsilon, \lambda) > 0$.

In what follows, we fix $\delta > 0$ such that $\delta n^\alpha + 2 < d_{i_2(\lambda)}/100$. Let $m = \max\{l \geq 1 : 2^{l+1} \leq \delta n^\alpha\}$. Define $I_l := [2^{l-1}, 2^{l+1})$ for $l \geq 1$. Let \mathbb{P}_x denote the law of the random walk s_n , starting from x and satisfying the recurrence relation in (4.27). Let $\sigma_{nl} := \min\{t \geq 1 : s_n(t) \notin I_l\}$ and $r_{nl} := \min\{t \geq 1 : \sup_{x \in I_l} \mathbb{P}_x(\sigma_{nl} > t) \leq 1/2\}$. Let us first obtain the following bound on r_{nl} :

Lemma 4.8. *For all $n \geq N_\lambda$ and $1 \leq l \leq d_{i_2(\lambda)}$, $r_{nl} \leq C2^{(\tau-2)l}$, for some (sufficiently large) constant $C = C(\varepsilon, \lambda) > 0$.*

Proof. Fix any $1 \leq l \leq d_{i_2(\lambda)}$, and suppose that $d_{k+1} < l \leq d_k$. Using (4.17) and Assumption 2 (i), it follows that

$$\mathbb{P}(\bar{\xi}_n(i_2(\lambda)) \geq l) = \frac{\sum_{j=i_2(\lambda)}^k d_j}{L} \geq C \frac{\sum_{j=i_2(\lambda)}^k d_j}{\sum_{j=1}^k d_j} \times \frac{\sum_{j=1}^k d_j}{\ell_n} \geq Cl^{-(\tau-2)}, \quad (4.30)$$

where we have used that $\sum_{j=i_2(\lambda)}^k d_j = \Theta(\sum_{j=1}^k d_j)$ since i_λ is fixed. Next, in order to estimate σ_{nl} , we bound $\sup_{x \in I_l} \mathbb{P}_x(s_n(t) \in I_l)$ using an upper bound on Lévy's concentration function due to Esseen [18]. For a random variable Z , define Lévy's concentration function

$$Q(Z, L) = \sup_{x \in \mathbb{R}} \mathbb{P}(Z \in [x, x + L]). \quad (4.31)$$

By [18, Theorem 3.1], for any $u > 0$,

$$Q(s_n(t), u) \leq \frac{Cu}{(t \times \mathbb{E}[|\zeta_1 - \zeta_2|^2 \mathbb{1}\{|\zeta_1 - \zeta_2| \leq u\}])^{1/2}}, \quad (4.32)$$

where ζ_1, ζ_2 are i.i.d. realizations from the distribution of $\bar{\xi}_n(i_2(\lambda))$. To lower bound (4.32), first observe that, for any random variable Y supported on \mathbb{N} ,

$$\begin{aligned} \mathbb{E}[Y^2 \mathbb{1}\{Y \leq u\}] &= \sum_{1 \leq y \leq u} y^2 \mathbb{P}(Y = y) = \sum_{1 \leq y \leq u} \sum_{1 \leq x \leq y} y \mathbb{P}(Y = y) \\ &= \sum_{1 \leq x \leq u} \sum_{y \geq x} y \mathbb{P}(Y = y) \geq \sum_{1 \leq x \leq u} x \mathbb{P}(Y \geq x). \end{aligned} \quad (4.33)$$

Thus, for any $0 \leq u \leq \delta n^\alpha$,

$$\begin{aligned} \mathbb{E}[|\zeta_1 - \zeta_2|^2 \mathbb{1}\{|\zeta_1 - \zeta_2| \leq u\}] &\geq \sum_{1 \leq x \leq u} x \mathbb{P}(|\zeta_1 - \zeta_2| \geq x) \\ &\geq \sum_{1 \leq x \leq u} x \mathbb{P}(\zeta_1 \geq x + 1) \mathbb{P}(\zeta_2 = -1) \geq Cu^{4-\tau}, \end{aligned} \quad (4.34)$$

where the final step follows using (4.30) and the fact from (4.17) that $\liminf_{n \rightarrow \infty} \mathbb{P}(\zeta_2 = -1) > 0$. Thus, (4.32) yields

$$\sup_{x \in I_l} \mathbb{P}_x(\sigma_{nl} > t) \leq \sup_{x \in I_l} \mathbb{P}_x(s_n(t) \in I_l) \leq Q(s_n(t), 2^{l+2}) \leq \frac{C2^l}{t2^{l(4-\tau)}}, \quad (4.35)$$

which is at most $1/2$ by choosing $t = C2^{l(\tau-2)}$, for some large constant $C > 0$. Thus, the proof of Lemma 4.8 follows. \square

We now decompose the possible values of the random walk (4.27) into different scales. Recall that $I_l := [2^{l-1}, 2^{l+1})$. At each time t , the scale of $s_n(t)$, denoted by $\text{scl}(s_n(t))$, is an integer. Let $\text{scl}(s_n(0)) = 1$. Suppose that $\text{scl}(s_n(u)) = l$ for some $u > 0$. A change of scale occurs when s_n leaves I_l , i.e., at time $T := \inf\{t > u : s_n(t) \notin I_l\}$, and the new scale is given by

$\text{scl}(s_n(T)) = l'$, where l' is such that $s_n(T) \in [2^{l'-1}, 2^{l'})$. Now, the next change of scale occurs at time $T' := \inf\{t > T : s_n(t) \notin I_{l'}\}$, and the scale remains the same until T' , i.e., $\text{scl}(s_n(t)) = l'$ for all $T \leq t < T'$. Define

$$H_{nl}(t) := \sum_{u \in [0, t], \text{scl}(s_n(u))=l} \frac{1}{s_n(u)}, \quad \text{so that} \quad H_n(t) = \sum_{l \geq 1} H_{nl}(t). \quad (4.36)$$

Let $T_{nl}(t) := \#\{u \in [0, t) : \text{scl}(s_n(u)) = l\}$, and note that

$$2^{l-1} H_{nl}(t) \leq T_{nl}(t) \leq 2^{l+1} H_{nl}(t). \quad (4.37)$$

Therefore, for any $x \in \mathbb{N}$,

$$\mathbb{P}\left(H_{nl}(\sigma) \geq \frac{xr_{nl}}{2^{l-1}}\right) \leq \mathbb{P}(T_{nl}(\sigma) \geq xr_{nl}). \quad (4.38)$$

The next lemma estimates $\mathbb{P}(T_{nl}(\sigma) \geq xr_{nl})$:

Lemma 4.9. *For all $n \geq 1, l \geq 1$, and $x \in \mathbb{N}$,*

$$\mathbb{P}(T_{nl}(\sigma) \geq xr_{nl}) \leq C2^{-l-C'x}, \quad (4.39)$$

for some absolute constants $C, C' > 0$.

Proof. Let us first show that there exists a constant $C > 0$ such that for any $l \geq 2$,

$$\mathbb{P}(T_{nl}(\sigma) \neq 0) \leq 2^{-(l-1)}. \quad (4.40)$$

For any $t \geq 0$, let \mathcal{F}_t denote the minimum sigma-algebra with respect to which $(\zeta_u)_{u=0}^t$ is measurable, where we take $\zeta_0 = 1$. Note that if $T_{nl}(\sigma) \neq 0$, then $s_n(u)$ hits 2^{l-1} before hitting zero. For $H > 1$, let $\gamma_H := \min\{t : s_n(t) \geq H, \text{ or } s_n(t) = 0\}$. Since $(s_n(t))_{t \geq 0}$ is a supermartingale with respect to the filtration $(\mathcal{F}_t)_{t \geq 0}$, as $\mathbb{E}[\zeta_u - 1] < 0$ by (4.21), an application of the optional stopping theorem yields

$$H\mathbb{P}(s_n(\gamma_H) \geq H) \leq \mathbb{E}[s_n(\gamma_H)] \leq \mathbb{E}[s_n(0)] = 1, \quad (4.41)$$

and therefore,

$$\mathbb{P}(s_n(\gamma_H) \geq H) \leq \frac{1}{H}. \quad (4.42)$$

Thus, (4.40) follows by taking $H = 2^{l-1}$ together with the fact that $T_{nl}(\sigma) \neq 0$ implies that $s_n(\gamma_H) \geq H$.

Next, the number of upcrossings of an interval $[a, b]$ by s_n is defined to be the supremum of all integers k such that there exist times $(u_j, t_j)_{j=1}^k$ satisfying $u_1 < t_1 < u_2 < \dots < t_k$, and $s_n(u_j) < a < b < s_n(t_j)$ for all $j \in [k]$. We will use the following simple fact (see [1, Proposition 3.2]). Let $U_n(t, [a, b])$ denote the number of upcrossings of $[a, b]$ by s_n before time t . Then, for any positive integers k, a, b with $a < b$,

$$\mathbb{P}(U_n(\sigma, [a, b]) \geq k) \leq \left(\frac{a-1}{b}\right)^k. \quad (4.43)$$

Next define $\text{visit}(l, t)$ to be the number of visits to scale l upto time t , i.e., this is the supremum over $k \in \mathbb{N}$ such that one can find $(u_j, t_j)_{j=1}^k$ with $u_1 < t_1 < \dots < u_k < t_k \leq t$ satisfying $\text{scl}(s_n(u_j)) \neq l$ but $\text{scl}(s_n(t_j)) = l$. Set $\text{visit}(1, 0) = 1$ and $\text{visit}(l, t) = 0$ if $\text{scl}(s_n(t)) \neq l$. Further, define $M_{nl} = \text{visit}(l, \sigma)$ (the total number of visits to scale l) and $t_{jl} = \#\{t < \sigma : \text{scl}(s_n(t)) =$

l , $\text{visit}(l, t) = j$ (the time spent at scale l during the j -th visit). Thus $T_{nl}(\sigma) = \sum_{j=1}^{M_{nl}} t_{jl}$, and for $m \geq 2$,

$$\mathbb{P}\left(\sum_{j=1}^{M_{nl}} t_{jl} > ar_{nl}\right) \leq \mathbb{P}(M_{nl} > m) + \mathbb{P}\left(\sum_{j=1}^m t_{jl} > ar_{nl}\right). \quad (4.44)$$

Now s_n can enter scale l from below, which yields an upcrossing of the interval $[2^{l-1}, 2^l]$. Otherwise, s_n can enter scale l from above, whence it must be the case that while leaving the scale l during the previous visit, the walk went from scale l to $l+1$. The latter case yields an upcrossing of $[2^l, 2^{l+1}]$. Therefore,

$$M_{nl} \leq U_n(\sigma, [2^{l-1}, 2^l]) + U_n(\sigma, [2^l, 2^{l+1}]), \quad (4.45)$$

and using (4.43), it follows that

$$\begin{aligned} & \mathbb{P}(M_{nl} > m) \\ & \leq \mathbb{P}\left(U_n(\sigma, [2^{l-1}, 2^l]) \geq (m+1)/2\right) + \mathbb{P}\left(U_n(\sigma, [2^l, 2^{l+1}]) \geq (m+1)/2\right) \\ & \leq \frac{1}{2^{(m-1)/2}}. \end{aligned} \quad (4.46)$$

On the other hand, after each time r_{nl} , the probability of exiting from scale l is at most $1/2$, by definition. Now, by the definition of r_{nl} just above Lemma 4.9, $\mathbb{P}(t_{jl} > kr_{nl}) \leq 2^{-k}$, which implies that $\lfloor t_{jl}/r_{nl} \rfloor$ can be stochastically dominated by a Geometric($1/2$) random variable. Also t_{jl} are independent for different j since they depend on disjoint sets of ζ -values. Thus, if $(g_i)_{i \geq 1}$ denotes an i.i.d. collection of Geometric($1/2$) random variables,

$$\begin{aligned} \mathbb{P}\left(\sum_{j=1}^m t_{jl} \geq (k+m)r_{nl}\right) & \leq \mathbb{P}\left(\sum_{j=1}^m \lfloor \frac{t_{jl}}{r_{nl}} \rfloor \geq k\right) \leq \mathbb{P}\left(\sum_{i=1}^m g_i > k\right) \\ & = \mathbb{P}(\text{Bin}(k, 1/2) < m) \leq e^{-(k-2m)^2/2k}, \end{aligned} \quad (4.47)$$

for $2m < k$, where the penultimate step uses the fact that $\sum_{i=1}^m g_i$ has a negative binomial distribution, and last step follows using standard concentration inequalities such as [27, Theorem 2.1]. Therefore, the proof follows by taking $k = x$ and $m = x/4$. \square

We are now ready to prove Proposition 4.7.

Proof of Proposition 4.7. Recall the definition of s_n from (4.27) starting from one. We first estimate the probability of the event \mathcal{B}_n that s_n hits $\delta n^\alpha/2$ before hitting zero. Let $\gamma := \min\{t : s_n(t) \geq \delta n^\alpha/2, \text{ or } s_n(t) = 0\}$. Using (4.42),

$$\mathbb{P}(\mathcal{B}_n) = \mathbb{P}\left(s_n(\gamma) \geq \frac{\delta n^\alpha}{2}\right) \leq \frac{2}{\delta n^\alpha}. \quad (4.48)$$

Next, for any sequence of positive numbers $(b_l)_{l \geq 1}$,

$$\begin{aligned} \mathbb{P}\left(H_n(\sigma) \geq \sum_{l=1}^m \frac{b_l r_{nl}}{2^{l-1}}\right) & \leq \frac{2}{\delta n^\alpha} + \mathbb{P}\left(H_n(\sigma) \geq \sum_{l=1}^m \frac{b_l r_{nl}}{2^{l-1}}, \text{ and } \mathcal{B}_n^c \text{ occurs}\right) \\ & \leq \frac{2}{\delta n^\alpha} + \mathbb{P}\left(H_{nl}(\sigma) \geq \frac{b_l r_{nl}}{2^{l-1}} \text{ for some } 1 \leq l \leq m\right), \end{aligned} \quad (4.49)$$

where the last step uses the fact that $H_n(\sigma) = \sum_{l=1}^m H_{nl}(\sigma)$ on \mathcal{B}_n^c . Using (4.38) and Lemma 4.9, (4.49) yields

$$\mathbb{P}\left(H_n(\sigma) \geq \sum_{l=1}^m \frac{b_l r_{nl}}{2^l}\right) \leq \frac{2}{\delta n^\alpha} + \sum_{l=1}^m \mathbb{P}(T_{nl}(\sigma) \geq b_l r_{nl}) \leq \frac{2}{\delta n^\alpha} + C \sum_{l=1}^m 2^{-l-C' b_l}. \quad (4.50)$$

Putting $b_l = \frac{1}{\tau}(m-l+1+2\log_2(m-l+1))$ for $1 \leq l \leq m$, and using Lemma 4.8,

$$\begin{aligned} \sum_{l=1}^m \frac{b_l r_{nl}}{2^{l-1}} &\leq C \sum_{l=1}^m (m-l+1+2\log_2(m-l+1)) 2^{l(\tau-3)} \\ &= C \sum_{j=1}^m \frac{(j+2\log_2 j) 2^{(m+1)(\tau-3)}}{2^{j(\tau-3)}} \leq C(\delta n^\alpha)^{\tau-3} \sum_{j=1}^m \frac{(j+2\log_2 j)}{2^{j(\tau-3)}} \\ &\leq C n^\eta \delta^{\tau-3}, \end{aligned} \quad (4.51)$$

where we have used $\sum_{j=1}^{\infty} \frac{(j+2\log_2 j)}{2^{j(\tau-3)}} < \infty$ in the last step. Also,

$$\sum_{l=1}^m 2^{-l-C'b_l} = \sum_{l=1}^m 2^{-(m+1)} (m-l+1)^{-2} \leq \frac{4}{\delta n^\alpha} \sum_{l=1}^{\infty} \frac{1}{l^2}. \quad (4.52)$$

Thus the proof of Proposition 4.7 follows by applying (4.51) and (4.52) to (4.50). \square

4.4 Proof of Proposition 2.2

Let us now complete the proof of Proposition 2.2 using Proposition 4.4 and Theorem 4.5. Note that these bounds work for $i_2(\lambda) \leq i \leq K_n$, and we will use path counting from [7, 25] to bound the diameter for $i > K_n$. Define $\mathcal{C}_{\text{res}}(i)$ to be the connected component containing vertex i in the graph $\mathcal{G}_n^{>i-1} = \text{CM}_n(\mathbf{d}) \setminus [i-1]$. Note that if $\Delta^{>K} > \varepsilon n^\eta$, then there exists a path of length εn^η in $\text{CM}_n(\mathbf{d})$ avoiding all the vertices in $[K]$. Suppose that the minimum index among vertices on that path is i_0 . Then $\text{diam}(\mathcal{C}_{\text{res}}(i_0)) > \varepsilon n^\eta$. Therefore, $\Delta^{>K} > \varepsilon n^\eta$ implies that there exists $i \in (K, K_n)$ satisfying $\text{diam}(\mathcal{C}_{\text{res}}(i)) > \varepsilon n^\eta$, or $\text{diam}(\text{CM}_n(\mathbf{d}) \setminus [K_n]) > \varepsilon n^\eta$. We will use the following lemma first to complete the proof of Proposition 2.2 and prove the lemma subsequently:

Lemma 4.10. *Under Assumptions 1, 2, for any $\varepsilon > 0$, $\lim_{n \rightarrow \infty} \mathbb{P}(\text{diam}(\text{CM}_n(\mathbf{d}) \setminus [K_n]) > \varepsilon n^\eta) = 0$.*

Proof of Proposition 2.2. Let $\partial_i(r)$ denote the number of vertices at distance r starting from vertex i in the graph $\mathcal{G}_n^{>i-1}$. Recall the definition of $\bar{\partial}$ in Proposition 4.4. Thus, using Proposition 4.4, and Theorem 4.5 together with Lemma 4.10, yields that

$$\begin{aligned} \mathbb{P}(\Delta^{>K} > \varepsilon n^\eta) &\leq \sum_{i \in (K, K_n)} \mathbb{P}(\bar{\partial}_i(\varepsilon n^\eta) \neq \emptyset) + \mathbb{P}(\text{diam}(\text{CM}_n(\mathbf{d}) \setminus [K_n]) > \varepsilon n^\eta) \\ &\leq C \sum_{i \in (K, K_n)} \left(\frac{d_i}{n^\alpha} \right) e^{-\varepsilon \beta_i^n} + o(1), \end{aligned} \quad (4.53)$$

where the final term tends to zero if we first take $n \rightarrow \infty$ and then take $K \rightarrow \infty$ using (1.10) in Assumption 2. Thus the proof of Proposition 2.2 follows. \square

Proof of Lemma 4.10. Note that $\text{CM}_n(\mathbf{d}) \setminus [K_n]$ is again distributed as a configuration model conditionally on its degree sequence \mathbf{d}' , with the criticality parameter at most

$$\nu'_n = \frac{\sum_{i > K_n} d'_i (d'_i - 1)}{\sum_{i > K_n} d'_i} \leq \frac{\sum_{i > K_n} d_i (d_i - 1)}{\ell_n - 2 \sum_{i=1}^{K_n} d_i} \leq 1 - R_n n^{-\eta}, \quad R_n = \omega(\log n), \quad (4.54)$$

where the one-but-last step follows using $d'_i \leq d_i$ and $\ell'_n = \sum_{i > K_n} d'_i = \ell_n - 2 \sum_{i=1}^{K_n} d_i$, and the last bound follows using Assumption 2 (ii) and an identical argument as in (4.20). Let $\mathbb{P}'(\cdot)$ denote the probability measure conditionally on \mathbf{d}' . We will use path-counting techniques

for subcritical configuration models. An identical argument as [25, Lemma 6.1] also shows that, for any $l \geq 1$, the expected number of paths of length l starting from vertex i is at most $\frac{\ell_n' d_i (\nu_n')^{l-1}}{\ell_n' - 2l + 3} \leq \ell_n'^2 (\nu_n')^{l-1}$. Thus, for any $i > K_n$,

$$\mathbb{P}'(\exists \text{ path of length at least } \varepsilon n^\eta \text{ from } i \text{ in } \text{CM}_n(\mathbf{d}) \setminus [K_n]) \leq C \ell_n'^2 \sum_{l > \varepsilon n^\eta} (\nu_n')^l, \quad (4.55)$$

Thus, for $i > K_n$, the probability in (4.55) is at most

$$C n^2 (1 - R_n n^{-\eta})^{\varepsilon n^\eta} \leq C n^2 e^{-\varepsilon R_n} = o(1/n). \quad (4.56)$$

Therefore,

$$\mathbb{P}'(\exists i > K_n : \exists \text{ path of length at least } \varepsilon n^\eta \text{ from } i \text{ in } \text{CM}_n(\mathbf{d}) \setminus [K_n]) = o(1), \quad (4.57)$$

and the proof of Lemma 4.10 follows. \square

5 Verification of the assumptions for percolated degrees: Proof of Theorem 1.3

Let \mathcal{G}_n denote the graph obtained by doing percolation with retention probability $p_c(\lambda)$ (defined in (1.13)) on $\text{CM}_n(\mathbf{d})$. Let $\mathbf{d}^p = (d_i^p)_{i \in [n]}$ denote the degree sequence of \mathcal{G}_n . By [19, Lemma 3.2], the law of \mathcal{G}_n , conditionally on \mathbf{d}^p , is same as the law of $\text{CM}_n(\mathbf{d}^p)$. Thus, it is enough to show that the percolated degree sequence satisfies (with possibly different parameters) Assumptions 1 and 2 with high probability when the original degree sequence $(d_i)_{i \in [n]}$ does so. Then the proof of Theorem 1.3 follows from Theorem 1.1.

First, note that $d_i^p \sim \text{Binomial}(d_i, p_c(\lambda))$, and thus, using standard concentration inequalities [27, Corollary 2.3], it follows that, for each fixed $i \geq 1$, and for any $\varepsilon > 0$,

$$\mathbb{P}(|d_i^p - d_i p_c(\lambda)| > \varepsilon d_i p_c(\lambda)) \leq 2e^{-\frac{\varepsilon^2}{3} d_i p_c(\lambda)}, \quad (5.1)$$

which verifies Assumption 1 (i). Next, let $M_r^p = \sum_{i \in [n]} (d_i^p)_r$ and $M_r = \sum_{i \in [n]} (d_i)_r$, where $(x)_r := x(x-1) \cdots (x-r+1)$. To verify the moment conditions in Assumption 1 (ii), note that (1.5) holds for \mathbf{d}^p since $\sum_{i > K} (d_i^p)_3 \leq \sum_{i > K} (d_i)_3$, where $(x)_r := x(x-1) \cdots (x-r+1)$. We will show that

$$M_1^p = (1 + O_{\mathbb{P}}(n^{-1/2})) p_c(\lambda) M_1, \quad M_2^p = (1 + O_{\mathbb{P}}(n^{\frac{3\alpha}{2}-1})) p_c(\lambda)^2 M_2. \quad (5.2)$$

Using (5.2), the first and second moment assumptions in Assumption 1 (ii) holds for the percolated degree sequence. The estimate (5.2) also shows that (1.6) holds. Indeed,

$$\frac{M_2^p}{M_1^p} = \frac{p_c(\lambda) M_2}{M_1} (1 + O_{\mathbb{P}}(n^{\frac{3\alpha}{2}-1})) = 1 + C \lambda n^{-\eta} + o(n^{-\eta}), \quad (5.3)$$

where $C = \lim_{n \rightarrow \infty} \frac{M_2}{M_1}$, and the last step follows using (1.13) and the fact that $-1 + 3\alpha/2 < -1 + 2\alpha = -\eta$.

It remains to prove (5.2). Since $\frac{1}{2} \sum_{i \in [n]} d_i^p$ has a binomial distribution with parameter $\ell_n/2$ and $p_c(\lambda)$, the first asymptotics follows from Chebyshev's inequality. For the asymptotics of M_2^p , we use the following construction of the percolated configuration model from [19].

Algorithm 2. $\mathbf{d}^p = (d_i^p)_{i \in [n]}$ can be generated as follows:

(S0) Sample $R_n \sim \text{Bin}(\ell_n/2, p_c(\lambda))$;

(S1) Conditionally on R_n , sample a uniform subset of $2R_n$ half-edges from the set of ℓ_n half-edges. Let $I_j^{(i)}$ denote the indicator that j -th half-edge of i is selected. Then $d_i^p = \sum_{j=1}^{d_i} I_j^{(i)}$ for all $i \in [n]$.

Using the above construction, note that

$$M_2^p = \sum_{i \in [n]} \sum_{1 \leq j_1 \neq j_2 \leq d_i} I_{j_1}^{(i)} I_{j_2}^{(i)}. \quad (5.4)$$

Let $\mathbb{P}_1(\cdot) = \mathbb{P}(\cdot | R_n)$ and similarly define $\mathbb{E}_1[\cdot]$, $\text{Var}_1(\cdot)$, and $\text{Cov}_1(\cdot, \cdot)$. Then,

$$\begin{aligned} \mathbb{E}_1[M_2^p] &= \sum_{i \in [n]} \sum_{1 \leq j_1 \neq j_2 \leq d_i} \mathbb{P}_1(I_{j_1}^{(i)} = 1, I_{j_2}^{(i)} = 1) = \sum_{i \in [n]} \sum_{1 \leq j_1 \neq j_2 \leq d_i} \frac{\binom{\ell_n - 2}{2R_n - 2}}{\binom{\ell_n}{2R_n}} \\ &= \sum_{i \in [n]} \sum_{1 \leq j_1 \neq j_2 \leq d_i} \frac{2R_n(2R_n - 1)}{\ell_n(\ell_n - 1)} = (1 + O_{\mathbb{P}}(n^{-1/2}))p_c(\lambda)^2 M_2, \end{aligned} \quad (5.5)$$

where the last step follows using $R_n = (1 + O_{\mathbb{P}}(n^{-1/2}))p_c(\lambda)\ell_n/2$. Also, conditionally on R_n , $I_j^{(i)}$, $j = 1, \dots, d_i$, $i \in [n]$ are negatively associated (cf. [17, Theorem 10]), which yields

$$\begin{aligned} \text{Var}_1(M_2^p) &\leq \sum_{i \in [n]} \sum_{1 \leq j_1 \neq j_2 \leq d_i} \text{Var}_1(I_{j_1}^{(i)} I_{j_2}^{(i)}) + \sum_{i \in [n]} \sum_{\substack{1 \leq j_1 \neq j_2 \leq d_i \\ 1 \leq j_3 \neq j_4 \leq d_i \\ |\{j_1, j_2\} \cap \{j_3, j_4\}| = 1}} \text{Cov}_1(I_{j_1}^{(i)} I_{j_2}^{(i)}, I_{j_3}^{(i)} I_{j_4}^{(i)}) \\ &\leq \sum_{i \in [n]} d_i^2 + 4 \sum_{i \in [n]} d_i^3 = O(n^{3\alpha}). \end{aligned} \quad (5.6)$$

Thus, for any $A > 0$,

$$\mathbb{P}(|M_2^p - \mathbb{E}_1[M_2^p]| > An^{3\alpha/2}) = \mathbb{E}[\mathbb{P}_1(|M_2^p - \mathbb{E}_1[M_2^p]| > An^{3\alpha/2})] \leq \frac{\mathbb{E}[\text{Var}_1(M_2^p)]}{A^2 n^{3\alpha}}, \quad (5.7)$$

which can be made arbitrarily small by choosing $A > 0$ large. Thus, we conclude the asymptotics of M_2^p in (5.2) by using (5.5) and (5.7).

Finally, we need to show convergence of empirical measure of d^p to finish verifying Assumption 1 (ii). Let $n_k^p = \#\{i : d_i^p = k\}$, and $n_{\geq k}^p = \sum_{r \geq k} n_r^p$. It suffices to show that

$$\frac{n_{\geq k}^p}{n} \xrightarrow{\mathbb{P}} \mathbb{P}(D^p \geq k) \quad \forall k \geq 1, \quad (5.8)$$

where D^p satisfies $D^p | D = l \sim \text{Bin}(l, 1/\nu)$, for all $l \geq 1$. Let V_n be a uniformly chosen vertex and $D_n^p = d_{V_n}^p$ and $D_n = d_{V_n}$. By the construction in Algorithm 2,

$$\begin{aligned} \mathbb{P}(D_n^p = k | D_n = l, R_n) &= \frac{\binom{l}{k} \binom{\ell_n - l}{2R_n - k}}{\binom{\ell_n}{2R_n}} = (1 + o(1)) \binom{l}{k} \left(\frac{2R_n}{\ell_n}\right)^k \left(1 - \frac{2R_n}{\ell_n}\right)^{l-k} \\ &= (1 + o_{\mathbb{P}}(1)) \binom{l}{k} \left(\frac{1}{\nu}\right)^k \left(1 - \frac{1}{\nu}\right)^{l-k}, \end{aligned} \quad (5.9)$$

where in the final step we have used that $R_n = p_c(\lambda)\ell_n/2(1 + o_{\mathbb{P}}(1))$ and $p_c(\lambda) = \nu^{-1}(1 + o(1))$. Thus,

$$\mathbb{E} \left[\frac{n_k^p}{n} \mid R_n \right] = \mathbb{P}(D_n^p = k | R_n) \xrightarrow{\mathbb{P}} \mathbb{P}(D^p = k). \quad (5.10)$$

Moreover, using the negative association above,

$$\begin{aligned}\text{Var}(n_{\geq k}^p | R_n) &= \text{Var}\left(\sum_{i:d_i \geq k} \mathbb{1}_{\{d_i^p \geq k\}} \mid R_n\right) \\ &\leq \sum_{i:d_i \geq k} \text{Var}(\mathbb{1}_{\{d_i^p \geq k\}} | R_n) = \mathbb{E}[n_{\geq k}^p | R_n] \leq n,\end{aligned}\tag{5.11}$$

and thus $\mathbb{E}[\text{Var}(n_{\geq k}^p/n | R_n)] = O(1/n)$. This together with (5.10) concludes (5.8).

We now verify here that Assumption 2 holds with high probability. To verify Assumption 2 (i), it is enough to show that there exists a constant $c_1 > 0$ such that with high probability

$$\frac{1}{n} \sum_{i \in [n]} d_i^p \mathbb{1}_{\{d_i^p \geq l\}} \geq \frac{c_1}{l^{\tau-2}}, \quad \text{for all } 1 \leq l \leq d_1.\tag{5.12}$$

First, for any random variable X , any a with $\mathbb{P}(X \geq a) \geq 1/2$, we have that $\mathbb{E}[X \mathbb{1}_{\{X \geq a\}}] \geq \mathbb{E}[X] \mathbb{P}(X \geq a) \geq \mathbb{E}[X]/2$. The condition on a is clearly satisfied by the binomial distribution with $a = \lfloor \mathbb{E}[X] \rfloor$. Therefore,

$$\begin{aligned}\frac{1}{n} \mathbb{E}\left[\sum_{i \in [n]} d_i^p \mathbb{1}_{\{d_i^p \geq l\}}\right] &\geq \frac{1}{n} \sum_{i:d_i p_c(\lambda) \geq l} \mathbb{E}[d_i^p \mathbb{1}_{\{d_i^p \geq l\}}] \\ &\geq \frac{1}{n} \sum_{i:d_i p_c(\lambda) \geq l} \mathbb{E}[d_i^p \mathbb{1}_{\{d_i^p \geq \lfloor d_i p_c(\lambda) \rfloor\}}] \geq \frac{1}{2n} \sum_{i:d_i \geq l/p_c(\lambda)} d_i p_c(\lambda) \geq \frac{C}{l^{\tau-2}}.\end{aligned}\tag{5.13}$$

Let $F_1 = \sum_{i \in [n]} d_i^p \mathbb{1}_{\{d_i^p \geq l\}}$, $F_2 = \mathbb{E}[F_1 | \text{CM}_n(\mathbf{d})]$. We will apply the bounded difference inequality from [27, Corollary 2.27]. Given the graph $\text{CM}_n(\mathbf{d})$, if we keep one extra edge in the percolated graph, then F_1 can change by at most $2l$. Thus, for any $\varepsilon > 0$,

$$\mathbb{P}\left(|F_1 - F_2| > \frac{n\varepsilon}{l^{\tau-2}} \mid \text{CM}_n(\mathbf{d})\right) \leq 2 \exp\left(-\frac{n^2 \varepsilon^2}{l^{2(\tau-2)} 4l^2 \frac{\ell_n}{2}}\right) \leq 2e^{-C\varepsilon^2 n l^{-2(\tau-1)}},\tag{5.14}$$

almost surely. Also, we can apply concentration inequalities such as [11, Lemma 2.5] to conclude that

$$\mathbb{P}\left(|F_2 - \mathbb{E}[F_2]| > \frac{n\varepsilon}{l^{\tau-2}}\right) \leq 2e^{-C\varepsilon^2 n l^{-2(\tau-1)}}.\tag{5.15}$$

Combining (5.14), (5.15) together with (5.13) shows that there exists an $\varepsilon_0 > 0$ such that with high probability (5.12) holds for all $1 \leq l \leq n^{\varepsilon_0}$. For $l \geq n^{\varepsilon_0}$, we can use (5.1) together with a union bound, and thus we conclude (5.12) for all $1 \leq l \leq d_1$. Thus, Assumption 2 (i) holds with high probability for the percolated degrees. To verify Assumptions 2 (ii), and (iii), let $K'_n := \min\{j \leq K_n : d_j \leq n^{\alpha/2}\}$, and $K'_n = K_n$ if $d_j \geq n^{\alpha/2}$ for all $j \leq K_n$. Then note that $(d_i)_{i \in [n]}$ satisfies Assumptions 2 (ii) and (iii) with K_n replaced by K'_n . This is because

$$\beta_{K_n}^n = n^{-2\alpha} \sum_{j \leq K'_n} d_j^2 + n^{-2\alpha} \sum_{K'_n \leq j < K_n} d_j^2 = \beta_{K'_n}^n + O(K_n n^{-\alpha}) = \beta_{K'_n}^n + o(1),\tag{5.16}$$

using $K_n = o(n^\alpha)$, and thus $\beta_{K'_n}^n = \omega(\log n)$. Also $K'_n \rightarrow \infty$, since if $K'_n = O(1)$, then $\beta_{K'_n}^n = O(1)$. Now, using (5.1) again, it follows that, for any $\varepsilon > 0$,

$$\mathbb{P}(\exists i \leq K'_n : |d_i^p - d_i p_c(\lambda)| > \varepsilon d_i p_c(\lambda)) \leq \sum_{i=1}^{K'_n} 2e^{-\frac{\varepsilon^2}{3} d_i p_c(\lambda)} \leq K'_n e^{-Cn^{\alpha/2}},\tag{5.17}$$

for some constant $C > 0$. If $|d_i^p - d_i p_c(\lambda)| \leq \varepsilon d_i p_c(\lambda)$, then Assumptions 2 (ii) and (iii) are clearly satisfied for d^p and thus the proof follows.

References

- [1] Addario-Berry, L. (2019). Most trees are short and fat. *Probab. Theory Relat. Fields*, 173(1-2):1–26.
- [2] Addario-Berry, L., Broutin, N., and Goldschmidt, C. (2012). The continuum limit of critical random graphs. *Probab. Theory Relat. Fields*, 152(3):367–406.
- [3] Aldous, D. (1991). The continuum random tree II: an overview. In Barlow, M. T. and Bingham, N. H., editors, *Stochastic Analysis: Proc. Durham Symp. Stochastic Analysis, 1990*, London Mathematical Society Lecture Note Series, pages 23–70. Cambridge University Press, Cambridge.
- [4] Aldous, D., Miermont, G., and Pitman, J. (2004). The exploration process of inhomogeneous continuum random trees, and an extension of Jeulin’s local time identity. *Probab. Theory Relat. Fields*, 129(2):182–218.
- [5] Athreya, S., Löhr, W., and Winter, A. (2016). The gap between Gromov-vague and Gromov–Hausdorff-vague topology. *Stoch. Proc. Appl.*, 126(9):2527–2553.
- [6] Bender, E. A. and Canfield, E. R. (1978). The asymptotic number of labeled graphs with given degree sequences. *J. Combin. Theory Ser. A*, 24(3):296–307.
- [7] Bhamidi, S., Dhara, S., van der Hofstad, R., and Sen, S. (2020). Universality for critical heavy-tailed random graphs: Metric structure of maximal components. *Electron. J. Probab.*, 25(47):1–57.
- [8] Bhamidi, S., van der Hofstad, R., and Sen, S. (2018). The multiplicative coalescent, inhomogeneous continuum random trees, and new universality classes for critical random graphs. *Probab. Theory Relat. Fields*, 170(1):387–474.
- [9] Billingsley, P. (1999). *Convergence of Probability Measures*. John Wiley & Sons, Inc.
- [10] Bollobás, B. (1980). A probabilistic proof of an asymptotic formula for the number of labelled regular graphs. *European J. Combin.*, 1(4):311–316.
- [11] Borgs, C., Chayes, J. T., Dhara, S., and Sen, S. (2019). Limits of sparse configuration models and beyond: Graphexes and Multi-Graphexes. *arXiv:1907.01605*.
- [12] Breiman, L. (1968). *Probability*. Classics in Applied Mathematics. SIAM: Society for Industrial and Applied Mathematics.
- [13] Broutin, N., Duquesne, T., and Wang, M. (2020). Limits of multiplicative inhomogeneous random graphs and Lévy trees: Limit theorems. *arXiv:2002.02769*.
- [14] Camarri, M. and Pitman, J. (2000). Limit distributions and random trees derived from the birthday problem with unequal probabilities. *Electron. J. Probab.*, 5:1–18.
- [15] Conchon-Kerjan, G. and Goldschmidt, C. (2020). The stable graph: the metric space scaling limit of a critical random graph with i.i.d. power-law degrees. *arXiv:2002.04954*.
- [16] Dhara, S., van der Hofstad, R., van Leeuwen, J. S. H., and Sen, S. (2016). Heavy-tailed configuration models at criticality. *arXiv:1612.00650*, To appear in *Ann. Inst. H. Poincaré (B) Probab. Statist.*
- [17] Dubhashi, D., Priebe, V., and Ranjan, D. (1996). Negative dependence through the FKG inequality. *Basic Research in Computer Science (BRICS)*.
- [18] Esseen, C. G. (1968). On the concentration function of a sum of independent random variables. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 9(4):290–308.
- [19] Fountoulakis, N. (2007). Percolation on sparse random graphs with given degree sequence. *Internet Math.*, 4(1):329–356.
- [20] Freedman, D. A. (1975). On tail probabilities for martingales. *Ann. Probab.*, 3(1):100–118.
- [21] Goldschmidt, C., Haas, B., and Sénizergues, D. (2018). Stable graphs: distributions and line-breaking construction. *arXiv:1811.06940*.
- [22] Greven, A., Pfaffelhuber, P., and Winter, A. (2009). Convergence in distribution of random metric measure spaces (Λ -coalescent measure trees). *Probab. Theory Relat. Fields*, 145(1):285–322.
- [23] van der Hofstad, R. (2017). *Random Graphs and Complex Networks*, volume 1. Cambridge University Press, Cambridge.
- [24] Janson, S. (2009). The probability that a random multigraph is simple. *Comb. Probab. Comp.*, 18(1-2):205–225.
- [25] Janson, S. (2010). Susceptibility of random graphs with given vertex degrees. *J. Combin.*, 1(3-4):357–387.
- [26] Janson, S. and Łuczak, M. J. (2009). A new approach to the giant component problem. *Random Struct. Algor.*, 34(2):197–216.
- [27] Janson, S., Łuczak, T., and Rucinski, A. (2000). *Random Graphs*. Wiley, New York.
- [28] Łuczak, T. (1995). The number of trees with large diameter. *J. Aust. Math. Soc. Series A. Pure Mathematics and Statistics*, 58(3):298–311.

- [29] Miermont, G. and Sen, S. (2019). On breadth-first constructions of scaling limits of random graphs and random unicellular maps. *arXiv:1908.04403*.
- [30] Molloy, M. and Reed, B. (1995). A critical-point for random graphs with a given degree sequence. *Random Struct. Algor.*, 6(2-3):161–179.
- [31] Nachmias, A. and Peres, Y. (2008). Critical random graphs: Diameter and mixing time. *Ann. Probab.*, 36(4):1267–1286.
- [32] Szekeres, G. (1983). Distribution of labelled trees by diameter. In Casse, L. R. A., editor, *Combinatorial Mathematics X*, pages 392–397, Berlin, Heidelberg. Springer Berlin Heidelberg.
- [33] Wang, M. (2015). Height and diameter of Brownian tree. *Electron. Commun. Probab.*, 20(88):1–15.
- [34] Whitt, W. (2002). *Stochastic-Process Limits: An Introduction to Stochastic-Process Limits and Their Application to Queues*. Springer-Verlag, New York.