

ON q -DEFORMED LEVI-CIVITA CONNECTIONS

JOAKIM ARNLIND, KWALOMBOTA ILWALE AND GIOVANNI LANDI

ABSTRACT. We explore the possibility of introducing q -deformed connections on the quantum 2-sphere and 3-sphere, satisfying a twisted Leibniz' rule in analogy with q -deformed derivations. We show that such connections always exist on projective modules. Furthermore, a condition for metric compatibility is introduced, and an explicit formula is given, parametrizing all metric connections on a free module. For the module of 1-forms on the quantum 3-sphere, a q -deformed torsion freeness condition is introduced and we derive explicit expressions for the Christoffel symbols of a Levi-Civita connection for a general class of metrics satisfying a certain reality condition. Finally, we construct metric connections on a class of projective modules over the quantum 2-sphere.

CONTENTS

1. Introduction	1
2. The quantum 3-sphere	3
2.1. q -deformed derivations	4
2.2. A left covariant calculus on S_q^3	5
3. q -affine connections	5
3.1. q -affine connections on projective modules	9
4. A q -affine Levi-Civita connection on $\Omega^1(S_q^3)$	10
5. The quantum 2-sphere	11
5.1. A left covariant calculus on S_q^2	12
5.2. Connections on projective modules over S_q^2	14
6. Computation of Christoffel symbols	17
References	22

1. INTRODUCTION

In recent years, a lot of progress has been made in understanding the Riemannian aspects of noncommutative geometry. These are not only mathematically interesting, but also important in physics where noncommutative geometry is expected to play a key role, notably in a theory of quantum gravity. In Riemannian geometry the Levi-Civita connection and its curvature have a central role, and it turns out that there are several different ways of approaching these objects in the noncommutative setting (see e.g. [CFF93, DVMMM96, Maj05, AC10, BM11, Ros13, AW17, BGM19, BGL20]).

From an algebraic perspective, the set of vector fields and the set of differential forms are (finitely generated projective) modules over the algebra of functions, a viewpoint which is

Date: 5 May 2020.

also adopted in noncommutative geometry. However, considering vector fields as derivations does not immediately carry over to noncommutative geometry, since the set of derivations of a (noncommutative) algebra is in general not a module over the algebra but only a module over the center of the algebra. Therefore, one is lead naturally to focus on differential forms and define a connection on a general module as taking values in the tensor product of the module with the module of differential forms. More precisely, let M be a (left) \mathcal{A} -module and let $\Omega^1(\mathcal{A})$ denote a module of differential forms together with a differential $d : \mathcal{A} \rightarrow \Omega^1(\mathcal{A})$. A connection on M is a linear map $\nabla : M \rightarrow \Omega^1(\mathcal{A}) \otimes M$ satisfying a version of Leibniz rule

$$(1.1) \quad \nabla(fm) = f\nabla m + df \otimes m$$

for $f \in \mathcal{A}$ and $m \in M$. In differential geometry, for a vector field X one obtains a covariant derivative $\nabla_X : M \rightarrow M$, by pairing differential forms with X (as differential forms are dual to vector fields). In a noncommutative version of the above, there is in general no canonical way of obtaining a ‘‘covariant derivative’’ $\nabla_X : M \rightarrow M$. In a derivation based approach to noncommutative geometry (see e.g. [DV88, DVMMM96]), one puts emphasis on the choice of a Lie algebra \mathfrak{g} of derivations of the algebra \mathcal{A} . Given a (left) \mathcal{A} -module M one defines a connection as a map $\nabla : \mathfrak{g} \times M \rightarrow M$, usually writing $\nabla(\partial, m) = \nabla_\partial m$ for $\partial \in \mathfrak{g}$ and $m \in M$, satisfying

$$(1.2) \quad \nabla_\partial(fm) = f\nabla_\partial m + \partial(f)m$$

for $f \in \mathcal{A}$ and $m \in M$, in parallel with (1.1).

For quantum groups, it turns out that natural analogues of vector fields are not quite derivations, but rather maps satisfying a twisted Leibniz rule. For instance, as we shall see, for the quantum 3-sphere S_q^3 one defines maps $X_a : S_q^3 \rightarrow S_q^3$ satisfying

$$(1.3) \quad X_a(fg) = fX_a(g) + X_a(f)\sigma_a(g)$$

for $f, g \in S_q^3$, where $\sigma_a : S_q^3 \rightarrow S_q^3$, for $a = 1, 2, 3$, are algebra morphisms. Thus, in this note, we explore the possibility of introducing a corresponding q -affine connection on a (left) S_q^3 -module M . That is, motivated by (1.3) we introduce a covariant derivative $\nabla_{X_a} : M \rightarrow M$ such that

$$(1.4) \quad \nabla_{X_a}(fm) = f\nabla_{X_a}m + X_a(f)\hat{\sigma}_a(m)$$

for $f \in S_q^3$ and $m \in M$, where $\hat{\sigma}_a$ denotes an extension of σ_a to the module M (cf. Section 3). In the following, we make these ideas precise and prove that there exist q -affine connections on projective modules. Furthermore, we introduce a condition for metric compatibility, and in the particular case of a left covariant calculus over S_q^3 , we investigate a derivation based definition of torsion. Then we explicitly construct a Levi-Civita connection, that is a torsion free and metric compatible connection. Moreover, we construct metric connections on a class of projective modules over the quantum 2-sphere. We note that the Riemannian geometry of quantum spheres can be studied [BM11] from the point of view of a bimodule connection on differential forms satisfying (1.1) as well as a right Leibniz rule twisted by a braiding map.

2. THE QUANTUM 3-SPHERE

In this section we recall a few basic properties of the quantum 3-sphere [Wor87]. The algebra S_q^3 is a unital $*$ -algebra generated by a, a^*, c, c^* fulfilling

$$\begin{aligned} ac &= qca & c^*a^* &= qa^*c^* & ac^* &= qc^*a \\ ca^* &= qa^*c & cc^* &= c^*c & a^*a + c^*c &= aa^* + q^2cc^* = \mathbb{1} \end{aligned}$$

for a real parameter q . The identification of S_q^3 with the quantum group $SEE_q(2)$ is via the Hopf algebra structure given by

$$\begin{aligned} \Delta(a) &= a \otimes a - qc^* \otimes c & \Delta(c) &= c \otimes a + a^* \otimes c \\ \Delta(a^*) &= -qc \otimes c^* + a^* \otimes a^* & \Delta(c^*) &= a \otimes c^* + c^* \otimes a^* \end{aligned}$$

with antipode and counit

$$\begin{aligned} S(a) &= a^* & S(c) &= -qc & \epsilon(a) &= 1 & \epsilon(c) &= 0 \\ S(a^*) &= a & S(c^*) &= -q^{-1}c^* & \epsilon(a^*) &= 1 & \epsilon(c^*) &= 0. \end{aligned}$$

Furthermore, the quantum enveloping algebra $\mathcal{U}_q(\mathfrak{su}(2))$ is the $*$ -algebra with generators E, F, K, K^{-1} satisfying

$$K^{\pm 1}E = q^{\pm 1}EK^{\pm 1} \quad K^{\pm 1}F = q^{\mp 1}FK^{\pm 1} \quad [E, F] = \frac{K^2 - K^{-2}}{q - q^{-1}}.$$

The corresponding Hopf algebra structure is given by the coproduct,

$$\Delta(E) = E \otimes K + K^{-1} \otimes E \quad \Delta(F) = F \otimes K + K^{-1} \otimes F \quad \Delta(K^{\pm 1}) = K^{\pm 1} \otimes K^{\pm 1}$$

together with antipode and counit

$$\begin{aligned} S(K) &= K^{-1} & S(E) &= -qE & S(F) &= -q^{-1}F \\ \epsilon(K) &= 1 & \epsilon(E) &= 0 & \epsilon(F) &= 0. \end{aligned}$$

We recall that there is a unique bilinear pairing between $\mathcal{U}_q(\mathfrak{su}(2))$ and S_q^3 given by

$$\begin{aligned} \langle K^{\pm 1}, a \rangle &= q^{\mp 1/2} & \langle K^{\pm 1}, a^* \rangle &= q^{\mp 1/2} \\ \langle E, c \rangle &= 1 & \langle F, c^* \rangle &= -q^{-1}, \end{aligned}$$

with the remaining pairings being zero. The pairing induces a $\mathcal{U}_q(\mathfrak{su}(2))$ -bimodule structure on S_q^3 given by

$$(2.1) \quad h \triangleright f = f_{(1)} \langle h, f_{(2)} \rangle \quad \text{and} \quad f \triangleleft h = \langle h, f_{(1)} \rangle f_{(2)}$$

for $h \in \mathcal{U}_q(\mathfrak{su}(2))$ and $f \in S_q^3$ with Sweedler's notation $\Delta(f) = f_{(1)} \otimes f_{(2)}$ (and implicit sum). The $*$ -structure on $\mathcal{U}_q(\mathfrak{su}(2))$, unconventionally denoted here by \dagger (for reasons that will become clear momentarily), is given by $(K^{\pm 1})^\dagger = K^{\pm 1}$ and $E^\dagger = F$. The action of $\mathcal{U}_q(\mathfrak{su}(2))$ is compatible with the $*$ -algebra structures in the following sense

$$h \triangleright f^* = (S(h)^\dagger \triangleright f)^* \quad f^* \triangleleft h = (f \triangleleft S(h)^\dagger)^*.$$

Let us for convenience list the left and right actions of the generators:

$$\begin{aligned}
K^{\pm 1} \triangleright a^n &= q^{\mp \frac{n}{2}} a^n & K^{\pm 1} \triangleright c^n &= q^{\mp \frac{n}{2}} c^n \\
K^{\pm 1} \triangleright a^{*n} &= q^{\pm \frac{n}{2}} (a^*)^n & K^{\pm 1} \triangleright c^{*n} &= q^{\pm \frac{n}{2}} (c^*)^n \\
E \triangleright a^n &= -q^{(3-n)/2} [n] a^{n-1} c^* & E \triangleright c^n &= q^{(1-n)/2} [n] c^{n-1} a^* \\
E \triangleright (a^*)^n &= 0 & E \triangleright (c^*)^n &= 0. \\
F \triangleright a^n &= 0 & F \triangleright c^n &= 0 \\
F \triangleright (a^*)^n &= q^{(1-n)/2} [n] c (a^*)^{n-1} & F \triangleright (c^*)^n &= -q^{-(1+n)/2} [n] a (c^*)^{n-1}
\end{aligned}$$

and

$$\begin{aligned}
a^n \triangleleft K^{\pm 1} &= q^{\mp \frac{n}{2}} a^n & (a^*)^n \triangleleft K^{\pm 1} &= q^{\pm \frac{n}{2}} (a^*)^n \\
c^n \triangleleft K^{\pm 1} &= q^{\pm \frac{n}{2}} c^n & (c^*)^n \triangleleft K^{\pm 1} &= q^{\mp \frac{n}{2}} (c^*)^n \\
a^n \triangleleft F &= q^{\frac{n-1}{2}} [n] c a^{n-1} & (a^*)^n \triangleleft F &= 0 \\
c^n \triangleleft F &= 0 & (c^*)^n \triangleleft F &= -q^{\frac{n-3}{2}} [n] a^* (c^*)^{n-1} \\
a^n \triangleleft E &= 0 & (a^*)^n \triangleleft E &= -q^{\frac{n-3}{2}} [n] c^* (a^*)^{n-1} \\
c^n \triangleleft E &= q^{\frac{n-1}{2}} [n] c^{n-1} a & (c^*)^n \triangleleft E &= 0
\end{aligned}$$

where $[n] = (q^n - q^{-n}) / (q - q^{-1})$.

2.1. q -deformed derivations. The algebra S_q^3 comes with a standard set of three q -deformed derivations (which can be used to generate a left covariant differential calculus, see Section 2.2). Namely, defining X_a for $a = 1, 2, 3$ as

$$X_1 \equiv X_+ = \sqrt{q} EK \quad X_2 \equiv X_- = \frac{1}{\sqrt{q}} FK \quad X_3 \equiv X_z = \frac{1 - K^4}{1 - q^{-2}}$$

it follows that for $f, g \in S_q^3$ (where $X_a(f)$ denotes either $X_a \triangleright f$ or $f \triangleleft X_a$)

$$\begin{aligned}
X_+(fg) &= fX_+(g) + X_+(f)\sigma_+(g) \\
X_-(fg) &= fX_-(g) + X_-(f)\sigma_-(g) \\
X_z(fg) &= fX_z(g) + X_z(f)\sigma_z(g),
\end{aligned}$$

with

$$(2.2) \quad \sigma_+ = \sigma_- = K^2 \quad \text{and} \quad \sigma_z = K^4.$$

Furthermore, these maps satisfy the following q -deformed commutation relations

$$(2.3) \quad X_- X_+ - q^2 X_+ X_- = X_z$$

$$(2.4) \quad q^2 X_z X_- - q^{-2} X_- X_z = (1 + q^2) X_-$$

$$(2.5) \quad q^2 X_+ X_z - q^{-2} X_z X_+ = (1 + q^2) X_+.$$

For an arbitrary map $X : S_q^3 \rightarrow S_q^3$ one defines $X^* : S_q^3 \rightarrow S_q^3$ as

$$X^*(f) = (X(f^*))^*$$

and it follows that

$$(2.6) \quad X_+^* = -K^{-2}X_- \quad X_-^* = -K^{-2}X_+ \quad X_z^* = -K^{-4}X_z,$$

satisfying

$$(2.7) \quad X_+^*(fg) = \sigma_+^*(f)X_+^*(g) + X_+^*(f)g$$

$$(2.8) \quad X_-^*(fg) = \sigma_-^*(f)X_-^*(g) + X_-^*(f)g$$

$$(2.9) \quad X_z^*(fg) = \sigma_z^*(f)X_z^*(g) + X_z^*(f)g$$

with

$$(2.10) \quad \sigma_+^* = \sigma_+^{-1} = K^{-2}, \quad \sigma_-^* = \sigma_-^{-1} = K^{-2}, \quad \sigma_z^* = \sigma_z^{-1} = K^{-4}.$$

We stress that X_a^* is different from X_a^\dagger , as defined above on $\mathcal{U}_q(\text{su}(2))$.

2.2. A left covariant calculus on S_q^3 . It is well known that there is a left covariant (first order) differential calculus on S_q^3 , denoted by $\Omega^1(S_q^3)$, generated as a left S_q^3 -module by

$$\omega_1 = \omega_+ = a dc - qc da \quad \omega_2 = \omega_- = c^* da^* - qa^* dc^* \quad \omega_3 = \omega_z = a^* da + c^* dc,$$

together with the differential $d : S_q^3 \rightarrow \Omega^1(S_q^3)$

$$(2.11) \quad df = (X_+ \triangleright f)\omega_+ + (X_- \triangleright f)\omega_- + (X_z \triangleright f)\omega_z$$

for $f \in S_q^3$ [Wor87]. In fact, $\Omega^1(S_q^3)$ is a free left module with a basis given by $\{\omega_+, \omega_-, \omega_z\}$.

Moreover, $\Omega^1(S_q^3)$ is a bimodule with respect to the relations

$$\begin{aligned} \omega_z a &= q^{-2} a \omega_z & \omega_z a^* &= q^2 a^* \omega_z & \omega_z c &= q^{-2} c \omega_z & \omega_z c^* &= q^2 c^* \omega_z \\ \omega_\pm a &= q^{-1} a \omega_\pm & \omega_\pm a^* &= q a^* \omega_\pm & \omega_\pm c &= q^{-1} c \omega_\pm & \omega_\pm c^* &= q c^* \omega_\pm, \end{aligned}$$

and, furthermore, $\Omega^1(S_q^3)$ is a $*$ -bimodule with

$$\omega_+^\dagger = -\omega_- \quad \omega_z^\dagger = -\omega_z$$

satisfying $(f\omega g)^\dagger = g^* \omega^\dagger f^*$ for $f, g \in S_q^3$ and $\omega \in \Omega^1(S_q^3)$.

3. q -AFFINE CONNECTIONS

In differential geometry, a connection extends the action of derivatives to vector fields, and for S_q^3 a natural set of (q -deformed) derivations is given by $\{X_a\}_{a=1}^3 = \{X_+, X_-, X_z\}$. In this section, we will introduce a framework extending the action of X_a to a connection on S_q^3 -modules. Let us first define the set of derivations we shall be interested in.

Definition 3.1. The quantum tangent space of S_q^3 is defined as

$$TS_q^3 = \mathbb{C} \langle X_+, X_+, X_-, X_-, X_z, X_z^* \rangle,$$

that is the complex vector space generated by X_a and X_a^* for $a = 1, 2, 3$.

Considering TS_q^3 to be the analogue of a (complexified) tangent space of S_q^3 , we would like to introduce a covariant derivative ∇_X on a (left) S_q^3 -module M , for $X \in TS_q^3$. Since the basis elements of TS_q^3 act as q -deformed derivations, the connection should obey an analogous q -deformed Leibniz rule. The motivating example is when $M = S_q^3$ and the action of TS_q^3 is simply $\nabla_X f = X(f)$ for $X \in TS_q^3$ and $f \in S_q^3$. In fact, let us be slightly more general and

consider the action on a free module of rank n . Thus, we let M be a free left S_q^3 -module with basis $\{e_i\}_{i=1}^n$, and write an arbitrary element $m \in M$ as $m = m^i e_i$ for $m^i \in S_q^3$, implicitly assuming a summation over i from 1 to n . Moreover, we assume there exist \mathbb{C} -linear maps $\hat{\sigma}_a, \hat{\sigma}_a^* : M \rightarrow M$ such that

$$\hat{\sigma}_a(fm) = \sigma_a(f)\hat{\sigma}_a(m) \quad \text{and} \quad \hat{\sigma}_a^*(fm) = \sigma_a^*(f)\hat{\sigma}_a^*(m)$$

for $f \in S_q^3$, $m \in M$ and $a = 1, 2, 3$ (for instance, one may choose $\hat{\sigma}_a(m^i e_i) = \sigma_a(m^i) e_i$ and similarly for $\hat{\sigma}_a^*$). Let us define $\nabla^0 : TS_q^3 \times M \rightarrow M$ by setting

$$(3.1) \quad \nabla_{X_a}^0(m) = X_a(m^i)\hat{\sigma}_a(e_i) \quad \text{and} \quad \nabla_{X_a^*}^0(m) = X_a^*(m^i)e_i$$

for $m = m^i e_i \in M$ (and extending it linearly to all of TS_q^3). Now, it is easy to check that

$$(3.2) \quad \nabla_{X_a}^0(fm) = f\nabla_{X_a}^0 m + X_a(f)\hat{\sigma}_a(m)$$

$$(3.3) \quad \nabla_{X_a^*}^0(fm) = \sigma_a^*(f)\nabla_{X_a^*}^0 m + X_a^*(f)m$$

for $f \in S_q^3$ and $m \in M$. Let us generalize these concepts to arbitrary S_q^3 -modules supporting an action of σ_a .

Definition 3.2. Let M be a left S_q^3 -module and let $\hat{\sigma}_a, \hat{\sigma}_a^* : M \rightarrow M$ be maps such that

$$\begin{aligned} \hat{\sigma}_a(\lambda_1 m_1 + \lambda_2 m_2) &= \lambda_1 \hat{\sigma}_a(m_1) + \lambda_2 \hat{\sigma}_a(m_2) & \hat{\sigma}_a(fm) &= \sigma_a(f)\hat{\sigma}_a(m) \\ \hat{\sigma}_a^*(\lambda_1 m_1 + \lambda_2 m_2) &= \lambda_1 \hat{\sigma}_a^*(m_1) + \lambda_2 \hat{\sigma}_a^*(m_2) & \hat{\sigma}_a^*(fm) &= \sigma_a^*(f)\hat{\sigma}_a^*(m) \end{aligned}$$

for $\lambda_1, \lambda_2 \in \mathbb{C}$, $f \in S_q^3$, $m_1, m_2, m \in M$ and $a = 1, 2, 3$. Then $(M, \hat{\sigma}_a, \hat{\sigma}_a^*)$ is called a σ -module. Moreover, given the σ -modules $(M, \hat{\sigma}_a, \hat{\sigma}_a^*)$ and $(\tilde{M}, \tilde{\sigma}_a, \tilde{\sigma}_a^*)$, a left module homomorphism $\phi : M \rightarrow \tilde{M}$ is called a σ -module homomorphism if

$$\phi(\hat{\sigma}_a(m)) = \tilde{\sigma}_a(\phi(m))z \quad \text{and} \quad \phi(\hat{\sigma}_a^*(m)) = \tilde{\sigma}_a^*(\phi(m))$$

for $m \in M$ and $a = 1, 2, 3$.

For notational convenience, when the maps $\hat{\sigma}_a, \hat{\sigma}_a^*$ are clear from the context, we shall simply write M for the σ -module $(M, \hat{\sigma}, \hat{\sigma}^*)$. Next, motivated by (3.2) and (3.3), we introduce connections on σ -modules.

Definition 3.3. Let M be a left σ -module. A left q -affine connection on M is a map $\nabla : TS_q^3 \times M \rightarrow M$ such that

- (1) $\nabla_X(\lambda_1 m_1 + \lambda_2 m_2) = \lambda_1 \nabla_X m_1 + \lambda_2 \nabla_X m_2$,
- (2) $\nabla_{\lambda_1 X + \lambda_2 Y} m = \lambda_1 \nabla_X m + \lambda_2 \nabla_Y m$,
- (3) $\nabla_{X_a}(fm) = f\nabla_{X_a} m + X_a(f)\hat{\sigma}_a(m)$,
- (4) $\nabla_{X_a^*}(fm) = \sigma_a^*(f)\nabla_{X_a^*} m + X_a^*(f)m$,

for $a = 1, 2, 3$, $m, m_1, m_2 \in M$, $f \in S_q^3$, $X \in TS_q^3$ and $\lambda_1, \lambda_2 \in \mathbb{C}$.

Remark 3.4. Note that given ∇_{X_a} for $a = 1, 2, 3$, one can set

$$\nabla_{X_+^*} = -\hat{\sigma}_-^* \circ \nabla_{X_-} \quad \nabla_{X_-^*} = -\hat{\sigma}_+^* \circ \nabla_{X_+} \quad \nabla_{X_z^*} = -\hat{\sigma}_z^* \circ \nabla_{X_z}$$

satisfying (4) in Definition 3.3, due to (2.6) and (2.10).

Next, assume that the module M comes with a hermitian form $h : M \times M \rightarrow S_q^3$ satisfying

$$\begin{aligned} h(fm_1, m_2) &= fh(m_1, m_2) & h(m_1, m_2)^* &= h(m_2, m_1) \\ h(m_1 + m_2, m_3) &= h(m_1, m_3) + h(m_2, m_3) \end{aligned}$$

for $f \in S_q^3$ and $m_1, m_2, m_3 \in M$. On a free module with basis $\{e_i\}_{i=1}^n$, a hermitian form is given by $h_{ij} = h_{ji}^* \in S_q^3$ by setting

$$(3.4) \quad h(m_1, m_2) = m_1^i h_{ij} (m_2^j)^*$$

for $m_1 = m_1^i e_i \in (S_q^3)^n$ and $m_2 = m_2^j e_j \in (S_q^3)^n$. In the case of the q -affine connection ∇^0 in (3.1), one finds that

$$\begin{aligned} X_+(h(m_1, m_2)) &= X_+(m_1^i h_{ij} (m_2^j)^*) = m_1^i X_+(h_{ij} (m_2^j)^*) + X_+(m_1^i) \sigma_+(h_{ij} (m_2^j)^*) \\ &= m_1^i h_{ij} X_+((m_2^j)^*) + m_1^i X_+(h_{ij}) \sigma_+((m_2^j)^*) + X_+(m_1^i) \sigma_+(h_{ij} (m_2^j)^*), \end{aligned}$$

and assuming that $X_+(h_{ij}) = 0$ one obtains

$$\begin{aligned} X_+(h(m_1, m_2)) &= m_1^i h_{ij} (X_+^*(m_2^j))^* + \sigma_+((\sigma_+^{-1} \circ X_+)(m_1^i) h_{ij} (m_2^j)^*) \\ &= h(m_1, \nabla_{X_+^*}^0(m_2)) - \sigma_+(h(\nabla_{X_+^*}^0 m_1, m_2)), \end{aligned}$$

by using that $X_-^* = -\sigma_+^{-1} \circ X_+$. Corresponding formulas are easily worked out for $\nabla_{X_-}^0, \nabla_{X_z}^0$, and we shall take this as a motivation for the following definition.

Definition 3.5. A q -affine connection ∇ on a left σ -module M is compatible with the hermitian form $h : M \times M \rightarrow S_q^3$ if

$$(3.5) \quad X_+(h(m_1, m_2)) = -\sigma_+(h(\nabla_{X_-^*} m_1, m_2)) + h(m_1, \nabla_{X_+^*} m_2)$$

$$(3.6) \quad X_-(h(m_1, m_2)) = -\sigma_-(h(\nabla_{X_+^*} m_1, m_2)) + h(m_1, \nabla_{X_-^*} m_2)$$

$$(3.7) \quad X_z(h(m_1, m_2)) = -\sigma_z(h(\nabla_{X_z^*} m_1, m_2)) + h(m_1, \nabla_{X_z^*} m_2),$$

for $m_1, m_2 \in M$.

Note that (3.5) and (3.6) are equivalent since

$$\begin{aligned} &\left(X_+(h(m_2, m_1)) + \sigma_+(h(\nabla_{X_-^*} m_2, m_1)) - h(m_2, \nabla_{X_+^*} m_1) \right)^* \\ &= -K^{-2} \left(X_-(h(m_1, m_2)) + \sigma_-(h(\nabla_{X_+^*} m_1, m_2)) - h(m_1, \nabla_{X_-^*} m_2) \right). \end{aligned}$$

In the case of a q -affine connection on a free module, one can derive a convenient parametrization of all connections that are compatible with a given hermitian form. To this end, let us introduce some notation. Let $(S_q^3)^n$ be a free σ -module with basis $\{e_i\}_{i=1}^n$. A q -affine connection ∇ on $(S_q^3)^n$ can be determined by specifying the Christoffel symbols

$$\nabla_{X_a} e_i = \Gamma_{ai}^j e_j,$$

with $\Gamma_{ai}^j \in S_q^3$ for $a = 1, 2, 3$ and $i, j = 1, \dots, n$, and setting

$$\begin{aligned} \nabla_{X_+^*} e_i &= -\hat{\sigma}_-^*(\nabla_{X_-} e_i) & \nabla_{X_-^*} e_i &= -\hat{\sigma}_+^*(\nabla_{X_+} e_i) & \nabla_{X_z^*} e_i &= -\hat{\sigma}_z^*(\nabla_{X_z} e_i) \\ \nabla_{X_a} (m^i e_i) &= m^i \nabla_{X_a} e_i + X_a(m^i) \hat{\sigma}_a(e_i) & \nabla_{X_a^*} (m^i e_i) &= \sigma_a^*(m^i) \nabla_{X_a^*} e_i + X_a^*(m^i) e_i. \end{aligned}$$

As we shall see, the metric compatibility of ∇ is conveniently formulated in terms of

$$(3.8) \quad \tilde{\Gamma}_{ai,j} = \Gamma_{ai}^k \sigma_a(\tilde{h}_{akj}) \quad \tilde{h}_{aij} = h(\hat{\sigma}_a^*(e_i), e_j).$$

The hermitian form h is assumed to be invertible (i.e. inducing an isomorphism of the module and its dual) which implies that there exists h^{ij} such that $h_{ij}h^{jk} = h^{kj}h_{ji} = \delta_i^k \mathbf{1}$. In this case, one finds that \tilde{h}_{aij} (for $a = 1, 2, 3$) is invertible as well, implying that one may invert (3.8) as

$$\Gamma_{aj}^i = \tilde{\Gamma}_{aj,k} \sigma_a(\tilde{h}^{ki}).$$

Proposition 3.6. *Let $(S_q^3)^n$ be a free σ -module with a basis $\{e_i\}_{i=1}^n$ and let ∇ be a q -affine connection on $(S_q^3)^n$ given by the Christoffel symbols $\nabla_a e_i = \Gamma_{ai}^j e_j$. Furthermore, assume that h is an invertible hermitian form on $(S_q^3)^n$. Then ∇ is compatible with h if and only if there exist hermitian matrices $\alpha, \beta, \rho \in \text{Mat}_n(S_q^3)$ such that*

$$(3.9) \quad \tilde{\Gamma}_{+i,j} = \frac{1}{2}X_+(h_{ij}) + K(\alpha_{ij}) + iK(\beta_{ij})$$

$$(3.10) \quad \tilde{\Gamma}_{-i,j} = \frac{1}{2}X_-(h_{ij}) + K(\alpha_{ij}) - iK(\beta_{ij})$$

$$(3.11) \quad \tilde{\Gamma}_{zi,j} = \frac{1}{2}X_z(h_{ij}) + K^2(\rho_{ij}),$$

with $h_{ij} = h(e_i, e_j)$.

Proof. Starting from $\nabla_{X_a} e_i = \Gamma_{ai}^j e_j$ one obtains

$$\begin{aligned} -\sigma_+(h(\nabla_{X_+^*} e_i, e_j)) + h(e_i, \nabla_{X_+^*} e_j) &= \sigma_+(h(\hat{\sigma}_+^*(\Gamma_{+i}^k e_k), e_j)) - h(e_i, \hat{\sigma}_+^*(\Gamma_{-j}^k e_k)) \\ &= \Gamma_{+i}^k \sigma_+(h(\hat{\sigma}_+^*(e_k), e_j)) - h(e_i, \hat{\sigma}_+^*(e_k)) \sigma_-^*(\Gamma_{-j}^k)^* \\ &= \tilde{\Gamma}_{+i,j} - \sigma_-^*(\tilde{\Gamma}_{-j,i})^* \end{aligned}$$

giving the metric compatibility equation (3.5) as

$$(3.12) \quad \tilde{\Gamma}_{+i,j} = X_+(h_{ij}) + K^{-2}(\tilde{\Gamma}_{-j,i})^*.$$

Similarly, equation (3.7) may be written as

$$(3.13) \quad \tilde{\Gamma}_{zi,j} = X_z(h_{ij}) + K^{-4}(\tilde{\Gamma}_{zj,i})^*.$$

To solve (3.12) one may freely choose $\tilde{\Gamma}_{-i,j}$ and define $\tilde{\Gamma}_{+i,j}$ accordingly. Without loss of generality, let us write $\tilde{\Gamma}_{-i,j}$ in the following form

$$\tilde{\Gamma}_{-i,j} = \frac{1}{2}X_-(h_{ij}) + K(\alpha_{ij}) - iK(\beta_{ij})$$

for arbitrary $\alpha_{ij} = \alpha_{ji}^*$ and $\beta_{ij} = \beta_{ji}^*$. Then one obtains

$$\begin{aligned} \tilde{\Gamma}_{+i,j} &= X_+(h_{ij}) + K^{-2}(\tilde{\Gamma}_{-j,i})^* \\ &= X_+(h_{ij}) + \frac{1}{2}K^2(X_-(h_{ji})^*) + K^2(K(\alpha_{ji})^*) + iK^2(K(\beta_{ji})^*) \\ &= X_+(h_{ij}) - \frac{1}{2}K^2(K^{-2}X_+(h_{ij})) + K^2(K^{-1}(\alpha_{ij})) + iK^2(K^{-1}(\beta_{ij})) \\ &= \frac{1}{2}X_+(h_{ij}) + K(\alpha_{ij}) + iK(\beta_{ij}) \end{aligned}$$

Hence, every solution of (3.5) may be written in the above form. For the metric equation (3.7) one writes $\tilde{\Gamma}_{zi,j} = K^2(\rho_{ij} + ig_{ij})$, with $\rho_{ij} = \rho_{ji}^*$ and $g_{ij} = g_{ji}^*$, and notes that (3.13) is equivalent to

$$K^2(\rho_{ij} + ig_{ij}) = X_z(h_{ij}) + K^2(\rho_{ij} - ig_{ij}) \Leftrightarrow iK^2(g_{ij}) = \frac{1}{2}X_z(h_{ij}).$$

This is compatible with the requirement that $g_{ij} = g_{ji}^*$ since

$$(K^{-2}X_z(h_{ji}))^* = K^2(X_z(h_{ji})^*) = -K^2(K^{-4}X_z(h_{ji})) = -K^{-2}X_z(h_{ij}).$$

Hence, the general solution to (3.13) can be written as

$$\tilde{\Gamma}_{zi,j} = \frac{1}{2}X_z(h_{ij}) + K^2(\rho_{ij})$$

for arbitrary $\rho_{ij} = \rho_{ji}^*$, which concludes the proof. \square

3.1. q -affine connections on projective modules. As expected, q -affine connections exist on projective modules. More precisely, one proves the following result.

Proposition 3.7. *Let $M = ((S_q^3)^n, \hat{\sigma}_a^0, (\sigma_a^0)^*)$ be a free σ -module and let ∇^0 be a q -affine connection on M . If $p : (S_q^3)^n \rightarrow (S_q^3)^n$ is a projection and $\hat{\sigma}_a = p \circ \hat{\sigma}_a^0$, $\hat{\sigma}_a^* = p \circ (\hat{\sigma}_a^0)^*$, then*

$$(p(S_q^3)^n, \hat{\sigma}_a, \hat{\sigma}_a^*),$$

is a σ -module and $p \circ \nabla^0$ is a q -affine connection on $(p(S_q^3)^n, \hat{\sigma}_a, \hat{\sigma}_a^)$.*

Proof. It follows immediately that $(p(S_q^3)^n, p \circ \hat{\sigma}_a^0, p \circ (\hat{\sigma}_a^0)^*)$ satisfy the requirements of Definition 3.2, since $((S_q^3)^n, \hat{\sigma}_a^0, (\sigma_a^0)^*)$ is a σ -module. For instance,

$$\hat{\sigma}_a(fm) = p(\hat{\sigma}_a^0(fm)) = p(\sigma_a(f)\hat{\sigma}_a^0(m)) = \sigma_a(f)p(\hat{\sigma}_a^0(m)) = \sigma_a(f)\hat{\sigma}_a(m).$$

Since ∇^0 is a q -affine connection, it is immediate that $\nabla = p \circ \nabla^0$ satisfies properties (1) and (2) in Definition 3.3. Moreover, for $m \in p(S_q^3)^n$

$$\begin{aligned} \nabla_{X_a}(fm) &= p\nabla_{X_a}^0(fm) = fp(\nabla_{X_a}^0 m) + X_a(f)(p \circ \hat{\sigma}_a^0)(m) \\ &= f\nabla_{X_a} m + X_a(f)\hat{\sigma}_a(m) \\ \nabla_{X_a^*}(fm) &= \sigma_a^*(f)p(\nabla_{X_a^*}^0 m) + X_a^*(f)p(m) \\ &= \sigma_a^*(f)\nabla_{X_a^*} m + X_a^*(f)m, \end{aligned}$$

from which we conclude that ∇ is a q -affine connection on $p(S_q^3)^n$. \square

Since we have shown in the previous section that one can construct q -affine connections on free modules, Proposition 3.7 shows that q -affine connections exist on projective modules. Moreover, Let ∇ and $\tilde{\nabla}$ be q -affine connections on a σ -module M and define

$$\alpha(X, m) = \nabla_X m - \tilde{\nabla}_X m.$$

Then $\alpha : TS_q^3 \times M \rightarrow M$ satisfies

$$(3.14) \quad \alpha(\lambda X + Y, m_1) = \lambda\alpha(X, m_1) + \alpha(Y, m_1)$$

$$(3.15) \quad \alpha(X, fm_1 + m_2) = f\alpha(X, m_1) + \alpha(X, m_2)$$

for $m_1, m_2 \in M$, $X \in TS_q^3$, $f \in S_q^3$ and $\lambda \in \mathbb{C}$. Conversely, every q -affine connection on a projective module M can be written as

$$\nabla_X m = p(\nabla_X^0 m) + \alpha(X, m).$$

where ∇^0 is the connection defined in (3.1) and $\alpha : TS_q^3 \times M \rightarrow M$ is an arbitrary map satisfying (3.14) and (3.15). Next, let us show that a connection on a projective module is compatible with the restricted metric if the projection is orthogonal.

Proposition 3.8. *Let ∇ be a q -affine connection on the free σ -module $(S_q^3)^n$ and assume furthermore that ∇ is compatible with a hermitian form h on $(S_q^3)^n$. If $p : (S_q^3)^n \rightarrow (S_q^3)^n$ is an orthogonal projection, i.e.*

$$h(p(m_1), m_2) = h(m_1, p(m_2))$$

for all $m_1, m_2 \in (S_q^3)^n$, then $\tilde{\nabla} = p \circ \nabla$ is compatible with h restricted to $p(S_q^3)^n$.

Proof. Let us explicitly check one of the conditions in Definition 3.5 for $m_1, m_2 \in p(S_q^3)^n$:

$$\begin{aligned} -\sigma_+(h(\tilde{\nabla}_{X_-} m_1, m_2)) + h(m_1, \tilde{\nabla}_{X_+} m_2) &= -\sigma_+(h(p\nabla_{X_-} m_1, m_2)) + h(m_1, p\nabla_{X_+} m_2) \\ &= -\sigma_+(h(\nabla_{X_-} m_1, p(m_2))) + h(p(m_1), \nabla_{X_+} m_2) \\ &= -\sigma_+(h(\nabla_{X_-} m_1, m_2)) + h(m_1, \nabla_{X_+} m_2) = X_+(h(m_1, m_2)). \end{aligned}$$

The remaining conditions are checked in an analogous way. \square

4. A q -AFFINE LEVI-CIVITA CONNECTION ON $\Omega^1(S_q^3)$

In this section we shall construct a q -affine connection on the free left module $\Omega^1(S_q^3)$, compatible with a hermitian form h , satisfying a certain torsion freeness condition. The module $\Omega^1(S_q^3)$ is a free S_q^3 -module of rank 3 with basis $\omega_+, \omega_-, \omega_z$ which implies that the results of Proposition 3.6 may be used. To start with, one has to endow $\Omega^1(S_q^3)$ with the structure of a σ -module. Firstly, the actions (2.1) are extended to forms by requiring they commute with the differential d . Then, one checks directly that $K \triangleright \omega_+ = q^{-1}\omega_+$, $K \triangleright \omega_- = q\omega_-$, $K \triangleright \omega_z = \omega_z$, while $K \triangleleft \omega_a = \omega_a$, for $a = 1, 2, 3$. Let us work with the right action (without indicating it explicitly). The σ -module structure is then introduced as

$$K(\omega_a) = \omega_a \quad \Rightarrow \quad \hat{\sigma}_a(m^b \omega_b) = \sigma_a(m^b) \omega_b \quad \hat{\sigma}_a^*(m^b \omega_b) = \sigma_a^*(m^b) \omega_b.$$

Furthermore, we assume that h is an invertible hermitian form on $\Omega^1(S_q^3)$, such that $K(h_{ab}) = h_{ab}$, with $h_{ab} = h(\omega_a, \omega_b)$, for $a, b = 1, 2, 3$. (Note that for such a metric one has $X_z(h_{ab}) = 0$.) With these choices, one finds that $\tilde{\Gamma}_{ab,c} = \Gamma_{ab}^p h_{pc}$ where the Christoffel symbols are defined as $\nabla_{X_a} \omega_b = \Gamma_{ab}^c \omega_c$ for $a, b = 1, 2, 3$. In the case of a q -affine connection on $\Omega^1(S_q^3)$, there is a natural definition of torsion, motivated by (2.3)–(2.5).

Definition 4.1. A q -affine connection ∇ on $\Omega^1(S_q^3)$ is *torsion free* if

$$(4.1) \quad \nabla_- \omega_+ - q^2 \nabla_+ \omega_- = \omega_z$$

$$(4.2) \quad q^2 \nabla_z \omega_- - q^{-2} \nabla_- \omega_z = (1 + q^2) \omega_-$$

$$(4.3) \quad q^2 \nabla_+ \omega_z - q^{-2} \nabla_z \omega_+ = (1 + q^2) \omega_+.$$

In terms of $\tilde{\Gamma}_{ab,c}$, the conditions for a torsion free connection may be reformulated as

$$(4.4) \quad \tilde{\Gamma}_{-,+,a} - q^2 \tilde{\Gamma}_{+-,a} = K^2(\tilde{h}_{+za})$$

$$(4.5) \quad q^2 \tilde{\Gamma}_{z-,a} - q^{-2} \tilde{\Gamma}_{-,z,b} \sigma_-(\tilde{h}_-^{bc}) \sigma_z(\tilde{h}_{zca}) = (1 + q^2) K^4(\tilde{h}_{z-a})$$

$$(4.6) \quad q^{-2} \tilde{\Gamma}_{z+,a} - q^2 \tilde{\Gamma}_{+,z,b} \sigma_+(\tilde{h}_+^{bc}) \sigma_z(\tilde{h}_{zca}) = -(1 + q^2) K^4(\tilde{h}_{z+a})$$

for $a = 1, 2, 3$ and an arbitrary σ -module structure on $\Omega^1(S_q^3)$. For the particular case when the metric is invariant by K , that is $K(h_{ab}) = h_{ab}$, and $K(\omega_a) = \omega_a$ for $a, b = 1, 2, 3$, the torsion free equations become

$$(4.7) \quad \tilde{\Gamma}_{-,+,a} - q^2 \tilde{\Gamma}_{+-,a} = h_{za}$$

$$(4.8) \quad q^2 \tilde{\Gamma}_{z-,a} - q^{-2} \tilde{\Gamma}_{-,z,a} = (1 + q^2) h_{-a}$$

$$(4.9) \quad q^2 \tilde{\Gamma}_{z+,a} - q^{-2} \tilde{\Gamma}_{+,z,a} = (1 + q^2) h_{+a}.$$

Out of these, in Section 6, we derive an explicit expression for a torsion free and metric connection on $\Omega^1(S_q^3)$, and show that such a connection exists if the following reality condition is satisfied (cf. eq. (6.32)):

$$(4.10) \quad (X_+(h_{-z} - q^{-4} h_{z+}))^* = X_+(h_{-z} - q^{-4} h_{z+}).$$

The connection is not unique, and the solution depends on 6 parameters. For the particular case when $h_{ab} = h \delta_{ab}$ (obviously satisfying (4.10)), one can set all parameters to be zero: $\tau_1 = \tau_4 = \gamma_{+-} = f_0 = \mu_2 = \rho_{zz} = 0$, in the notation of Section 6. It follows that a torsion free and metric connection is given by

$$\begin{aligned} \nabla_+ \omega_+ &= X_+(h) h^{-1} \omega_+ \\ \nabla_+ \omega_- &= -\frac{1}{2} q^{-2} \omega_z \\ \nabla_+ \omega_z &= \frac{1}{2} \omega_+ + \frac{1}{2} X_+(h) h^{-1} \omega_z \\ \nabla_- \omega_+ &= \frac{1}{2} \omega_z \\ \nabla_- \omega_- &= X_-(h) h^{-1} \omega_- \\ \nabla_- \omega_z &= -\frac{1}{2} q^{-2} \omega_- + \frac{1}{2} X_-(h) h^{-1} \omega_z \\ \nabla_z \omega_+ &= -\frac{1}{2} q^2 (2 + q^2) \omega_+ + \frac{1}{2} q^4 X_+(h) h^{-1} \omega_z \\ \nabla_z \omega_- &= \frac{1}{2} (2 + 2q^{-2} - q^{-6}) \omega_- + \frac{1}{2} q^{-4} X_-(h) h^{-1} \omega_z \\ \nabla_z \omega_z &= -\frac{1}{2} q^2 X_-(h) h^{-1} \omega_+ - \frac{1}{2} q^{-2} X_+(h) h^{-1} \omega_- . \end{aligned}$$

5. THE QUANTUM 2-SPHERE

The noncommutative (standard) Podleś sphere S_q^2 [Pod87] can be considered as a subalgebra of S_q^3 by identifying the generators B_0, B_+, B_- of S_q^2 as

$$B_0 = cc^* \quad B_+ = ca^* \quad B_- = ac^* = B_+^*,$$

satisfying then the relations

$$\begin{aligned} B_- B_0 &= q^2 B_0 B_- & B_+ B_0 &= q^{-2} B_0 B_+ \\ B_- B_+ &= q^2 B_0 (\mathbb{1} - q^2 B_0) & B_+ B_- &= B_0 (\mathbb{1} - B_0). \end{aligned}$$

These elements generate the fix-point algebra of the right $U(1)$ -action

$$\alpha_z(a) = az \quad \alpha_z(a^*) = a^* \bar{z} \quad \alpha_z(c) = cz \quad \alpha_z(c^*) = c^* \bar{z}$$

for $z \in U(1)$ and $a \in S_q^3$, related to the $U(1)$ -Hopf-fibration $S_q^2 \hookrightarrow S_q^3$.

Now, the left action of the X_a does not preserve the algebra S_q^2 : one readily computes,

$$\begin{aligned} X_+ \triangleright B_0 &= qa^*c^* & X_- \triangleright B_0 &= -q^{-1}ca & X_z \triangleright B_0 &= 0 \\ X_+ \triangleright B_+ &= q(a^*)^2 & X_- \triangleright B_+ &= c^2 & X_z \triangleright B_+ &= 0 \\ X_+ \triangleright B_- &= q^2(c^*)^2 & X_- \triangleright B_- &= -q^{-1}(a)^2 & X_z \triangleright B_- &= 0. \end{aligned}$$

On the other hand, the right action of X_a does preserve the algebra S_q^2 . Let us denote $Y_a = X_a$ for the right action. Then, it is easy to check that

$$\begin{aligned} B_0 \triangleleft Y_+ &= q^{-1}B_- & B_0 \triangleleft Y_- &= -q^{-1}B_+ & B_0 \triangleleft Y_z &= 0 \\ B_+ \triangleleft Y_+ &= q\mathbb{1} - q(1+q^2)B_0 & B_+ \triangleleft Y_- &= 0 & B_+ \triangleleft Y_z &= -q^2(1+q^2)B_+ \\ B_- \triangleleft Y_+ &= 0 & B_- \triangleleft Y_- &= -q^{-1}\mathbb{1} + q^{-1}(1+q^2)B_0 & B_- \triangleleft Y_z &= (1+q^{-2})B_-. \end{aligned}$$

Note that when restricted to S_q^2 the Y_a are not independent. A long but straightforward computation shows that they are indeed related as

$$\begin{aligned} & ((f \triangleleft Y_+)B_+ q + (f \triangleleft Y_-)B_- q^{-1})(1+q^2) + (f \triangleleft Y_z) \left(1 - 2\frac{1+q^2}{1+q^4}B_0\right) \\ (5.1) \quad & = (f \triangleleft Y_z^2) q^{-2} \left(\frac{1-q^2}{1+q^4}(2q^4+q^2+1)B_0 - (1-q^6)B_0^2\right) \\ & + (f \triangleleft K^4) q^{-2}(1+q^2)((q^4-1)B_0 + (1-q^6)B_0^2), \end{aligned}$$

for $f \in S_q^2$. This can be checked on a vector space basis for the algebra S_q^2 , a basis which can be taken as $X(m)(B_0)^n$ for $m \in \mathbb{Z}$, $n \in \mathbb{N}$ with $X(m) = (B_+)^m$ for $m \geq 0$ and $X(m) = (B_-)^{-m}$ for $m < 0$ (cf. [MNW91]).

5.1. A left covariant calculus on S_q^2 . Since the element K acts (on the left) as the identity on S_q^2 , the differential (2.11) when restricted to $f \in S_q^2$ becomes

$$(5.2) \quad df = (X_- \triangleright f) \omega_- + (X_+ \triangleright f) \omega_+.$$

Note that $X_{\pm} \triangleright f \notin S_q^2$. In particular one finds

$$\begin{aligned} dB_+ &= q(a^*)^2 \omega_+ + c^2 \omega_-, \\ dB_- &= -q^2(c^*)^2 \omega_+ - q^{-1}a^2 \omega_-, \\ dB_0 &= c^*a^* \omega_+ - q^{-1}ca \omega_- \end{aligned}$$

which can be inverted to yield

$$\begin{aligned} \omega_+ &= q^{-1}a^2 dB_+ - q^2c^2 dB_- + (1+q^2)ac dB_0 \\ \omega_- &= (c^*)^2 dB_+ - q(a^*)^2 dB_- - (1+q^2)c^*a^* dB_0, \end{aligned}$$

implying that the differential in (5.2) can be expressed as

$$\begin{aligned} df &= (q^{-1}(X_+ \triangleright f) a^2 + (X_- \triangleright f) (c^*)^2) dB_+ \\ &\quad - (q^2(X_+ \triangleright f) + c^2 q(X_- \triangleright f) (a^*)^2) dB_- \\ &\quad + (1 + q^2)((X_+ \triangleright f) ac - (X_- \triangleright f) c^* a^*) dB_0. \end{aligned}$$

From this expression one finds that the differential d on S_q^2 can be written in terms of the right acting operators Y_a .

Lemma 5.1. *For $f \in S_q^2$, the differential in (5.2) can be written as*

$$(5.3) \quad df = (f \triangleleft V_+) dB_+ + (f \triangleleft V_-) dB_- + (f \triangleleft V_0) dB_0$$

where

$$V_+ = Y_+(1 - q^{-2}(1 + q^2)B_0)q^{-1} - Y_z B_- \frac{q^{-2}(1 + q^6)}{1 + q^4} + Y_z^2 B_- \frac{1 - q^2}{(1 + q^2)(1 + q^4)}$$

$$V_- = -Y_-(1 - q^2(1 + q^2)B_0)q + Y_z B_+ \frac{q^{-2}(1 + q^6)}{1 + q^4} - Y_z^2 B_+ \frac{1 - q^2}{(1 + q^2)(1 + q^4)}$$

$$V_0 = (Y_+ B_+ q^{-1} - Y_- B_- q)(1 + q^2) + Y_z B_0 \frac{(1 - q^4)(1 + q^6)}{1 + q^4} - Y_z^2 B_0 \frac{1 - q^2}{1 + q^4}.$$

Proof. By acting on the vector space basis $X(m)(B_0)^n$ (as introduced previously), one explicitly checks the equality of (5.2) and (5.3) via a tedious but straightforward computation. \square

Remark 5.2. When $q = 1$ the derivative (5.3) reduces to

$$(5.4) \quad \begin{aligned} df &= 2((f \triangleleft Y_+) B_+ - (f \triangleleft Y_-) B_-) dB_0 \\ &\quad + ((f \triangleleft Y_+) (1 - 2B_0) - (f \triangleleft Y_z) B_-) dB_+ \\ &\quad + (- (f \triangleleft Y_-) (1 - 2B_0) + (f \triangleleft Y_z) B_+) dB_-. \end{aligned}$$

Classically, the vector field X_a are the left invariant vector fields on $S^3 = SU(2)$ with dual left invariant forms ω_a . Thus they do not project to vector fields on the base space S^2 with commuting coordinates (B_+, B_-, B_0) and relation $B_+ B_- = B_0(1 - B_0)$: $X_a \triangleright f$ is not a function on S^2 even when f is. On the other hand, the vector fields Y_a are the right invariant vector fields on $SU(2)$ and thus they project to vector fields on S^2 , where they are not independent any longer and are related by

$$(5.5) \quad 2(B_+ Y_+ + B_- Y_-) + (1 - 2B_0) Y_z = 0,$$

which is just the relation to which (5.1) reduces when $q = 1$.

By changing coordinates $B_0 = \frac{1}{2}(1 - x)$ so that the radius condition for S^2 is written as $r^2 = 4B_+ B_- + x^2$, the exterior derivative operator in (5.4) becomes

$$df = \partial_x f dx + \partial_+ f dB_+ + \partial_- f dB_- - (\Delta f) (x dx + 2B_- dB_+ + 2B_+ dB_-)$$

where $\Delta = x \partial_x + B_+ \partial_+ + B_- \partial_-$ is the Euler (dilatation) vector field. One then computes $dr^2 = 2(1 - r^2)(x dx + 2B_- dB_+ + 2B_+ dB_-)$, which vanishes when restricting to S^2 : $r^2 - 1 = 0$.

The form (5.2) of the differential that uses left invariant vector fields and forms can be seen as identifying the cotangent bundle of S^2 with the direct sum of the line bundles of ‘charge’ ± 2 , that is $\Omega^1(S^2) \simeq \mathcal{L}_{-2}\omega_- \oplus \mathcal{L}_{+2}\omega_+$. This identification can be used also for the quantum sphere S_q^2 with the line bundles defined as in (5.6) below.

5.2. Connections on projective modules over S_q^2 . The definitions of σ -modules and q -affine connections apply equally well to the algebra S_q^2 . Note that the right action of K preserve S_q^2

$$B_0 \triangleleft K = B_0 \quad B_+ \triangleleft K = qB_+ \quad B_- \triangleleft K = q^{-1}B_-.$$

implying that σ_a, σ_a^* leave S_q^2 invariant.

In this section, we will construct q -affine connections on a class of projective modules over S_q^2 (cf. [BM98, HM99, Lan18]). For $n \geq 0$ and $\mu = 0, 1, \dots, n$, let $(\Psi_n)_\mu, (\Phi_n)_\mu \in S_q^3$ be given as

$$(\Phi_n)_\mu = \sqrt{\alpha_{n\mu}} c^{n-\mu} a^\mu \quad (\Psi_n)_\mu = \sqrt{\beta_{n\mu}} (c^*)^\mu (a^*)^{n-\mu}$$

with

$$\alpha_{n\mu} = \prod_{k=0}^{n-\mu-1} \frac{1 - q^{2(n-k)}}{1 - q^{2(k+1)}} \quad \beta_{n\mu} = q^{2\mu} \prod_{k=0}^{\mu-1} \frac{1 - q^{-2(n-k)}}{1 - q^{-2(k+1)}}.$$

It is straight-forward to check that

$$\sum_{\mu=0}^n (\Phi_n)_\mu^* (\Phi_n)_\mu = \sum_{\mu=0}^n (\Psi_n)_\mu^* (\Psi_n)_\mu = \mathbf{1},$$

implying that

$$(p_n)_\mu^\nu = (\Psi_n)_\mu (\Psi_n)_\nu^* = \sqrt{\beta_{n\mu}\beta_{n\nu}} (c^*)^\mu (a^*)^{n-\mu} a^{n-\nu} c^\nu$$

$$(p_{-n})_\mu^\nu = (\Phi_n)_\mu (\Phi_n)_\nu^* = \sqrt{\alpha_{n\mu}\alpha_{n\nu}} c^{n-\mu} a^\mu (a^*)^\nu (c^*)^{n-\nu}$$

satisfy $p_n^2 = p_n$ and $p_{-n}^2 = p_{-n}$. Moreover, it is easy to see that $(p_n)_\mu^\nu, (p_{-n})_\mu^\nu \in S_q^2$, which implies that

$$M_n = \begin{cases} p_n(S_q^2)^{n+1} & \text{if } n \geq 0 \\ p_{-|n|}(S_q^2)^{|n|+1} & \text{if } n < 0 \end{cases}$$

are finitely generated projective S_q^2 -modules for $n \in \mathbb{Z}$. Let us recall that these modules are isomorphic to the components in a (vector space) decomposition of S_q^3

$$S_q^3 = \bigoplus_{n \in \mathbb{Z}} \mathcal{L}_n$$

with

$$(5.6) \quad \mathcal{L}_n = \{f \in S_q^3 : \alpha_z(f) = \bar{z}^n f\},$$

and it follows that $\mathcal{L}_0 = S_q^2$, as well as $\mathcal{L}_n \mathcal{L}_m \subseteq \mathcal{L}_{n+m}$. For $f \in S_q^2$ and $f_n \in \mathcal{L}_n$ one has

$$\alpha_z(ff_n) = \alpha_z(f)\alpha_z(f_n) = \bar{z}^n ff_n,$$

which implies that \mathcal{L}_n is a left S_q^2 -module. Furthermore, it is easy to see that the right action of $\mathcal{U}_q(\mathfrak{su}(2))$ leaves each \mathcal{L}_n invariant. Let $\{e_\mu\}_{\mu=0}^n$ be a basis of $(S_q^2)^{n+1}$ and let $\phi_n^0 : (S_q^2)^{n+1} \rightarrow \mathcal{L}_n$ be defined as

$$\phi_n^0(m^\mu e_\mu) = \begin{cases} m^\mu (\Psi_n)_\mu & n \geq 0 \\ m^\mu (\Phi_{|n|})_\mu & n < 0 \end{cases},$$

and we note that $\phi_n^0(m) \in \mathcal{L}_n$ since $(\Psi_n)_\mu \in \mathcal{L}_n$ and $(\Phi_n)_\mu \in L_{-n}$ (for $n \geq 0$).

Lemma 5.3. *Let $m \in (S_q^2)^{n+1}$. If $p_n(m) = 0$ then $\phi_n^0(m) = 0$.*

Proof. If $p_n(m) = 0$ then (for $n \geq 0$)

$$\begin{aligned} \sum_{\mu=0}^n m^\mu (p_n)_\mu^\nu = 0 \quad \forall \nu &\Rightarrow \sum_{\mu,\nu=0}^n m^\mu (p_n)_\mu^\nu (\Psi_n)_\nu = 0 \Rightarrow \\ \left(\sum_{\mu=0}^n m^\mu (\Psi_n)_\mu \right) \left(\sum_{\nu=0}^n (\Psi_n)_\nu^* (\Psi_n)_\nu \right) = 0 &\Rightarrow \sum_{\mu=0}^n m^\mu (\Psi_n)_\mu = 0 \end{aligned}$$

which is equivalent to $\phi_n^0(m) = 0$. The proof for $n < 0$ is analogous. \square

The above result implies that ϕ_n^0 descends to a module homomorphism $\phi_n : M_n \rightarrow \mathcal{L}_n$, and one can show that ϕ_n is in fact an isomorphism. To simplify the presentation, let us in the following assume that $n \geq 0$. As generators of M_n one can choose $\hat{e}_\mu = p_n(e_\mu)$ for $\mu = 0, 1, \dots, n$, and one notes that

$$\phi_n(\hat{e}_\mu) = (p_n)_\mu^\nu (\Psi_n)_\nu = (\Psi_n)_\mu,$$

implying that $\{(\Psi_n)_\mu\}_{\mu=0}^n$ generates \mathcal{L}_n . Since the S_q^2 -module \mathcal{L}_n is a subset of S_q^3 which is also invariant under the right action of $\mathcal{U}_q(\mathfrak{su}(2))$, it is naturally a σ -module with respect to the right action of σ_a and σ_a^* ; one finds that

$$(\Psi_n)_\mu \triangleleft K = q^{\frac{1}{2}(n-2\mu)} (\Psi_n)_\mu \quad (\Psi_n)_\mu^* \triangleleft K = q^{-\frac{1}{2}(n-2\mu)} (\Psi_n)_\mu^*$$

giving

$$\begin{aligned} \sigma_\pm((\Psi_n)_\mu) &= q^{n-2\mu} (\Psi_n)_\mu & \sigma_z((\Psi_n)_\mu) &= q^{2(n-2\mu)} (\Psi_n)_\mu \\ \sigma_\pm^*((\Psi_n)_\mu) &= q^{-(n-2\mu)} (\Psi_n)_\mu & \sigma_z^*((\Psi_n)_\mu) &= q^{-2(n-2\mu)} (\Psi_n)_\mu. \end{aligned}$$

Correspondingly, we would like to define a σ -module structure on M_n such that ϕ_n is a morphism of σ -modules. To this end, we start by introducing a σ -module structure on $(S_q^2)^{n+1}$ in analogy with Proposition 3.7; namely, starting from

$$\begin{aligned} \hat{\sigma}_\pm^0(e_\mu) &= q^{n-2\mu} e_\mu & \hat{\sigma}_z^0(e_\mu) &= q^{2(n-2\mu)} e_\mu \\ (\hat{\sigma}_\pm^0)^*(e_\mu) &= q^{-(n-2\mu)} e_\mu & (\hat{\sigma}_z^0)^*(e_\mu) &= q^{-2(n-2\mu)} e_\mu \end{aligned}$$

we set $\hat{\sigma}_a = p \circ \hat{\sigma}_a^0 : M_n \rightarrow M_n$ and $\hat{\sigma}_a^* = p \circ (\hat{\sigma}_a^0)^* : M_n \rightarrow M_n$. Since

$$(p_n)_\mu^\nu \triangleleft K = ((\Psi_n)_\mu (\Psi_n)_\nu) \triangleleft K = q^{-(\mu-\nu)} (p_n)_\mu^\nu$$

one finds that

$$\begin{aligned}\hat{\sigma}_\pm(\hat{e}_\mu) &= (p_n \circ \hat{\sigma}_\pm^0)((p_n)_\mu^\nu e_\nu) = \sigma_\pm((p_n)_\mu^\nu)(p_n \circ \hat{\sigma}_\pm^0)(e_\nu) \\ &= q^{-2(\mu-\nu)} q^{n-2\nu} p_n(e_\nu) = q^{n-2\mu} \hat{e}_\mu \\ \hat{\sigma}_z(\hat{e}_\mu) &= q^{2(n-2\mu)} \hat{e}_\mu\end{aligned}$$

(and similarly for $\hat{\sigma}_a^*$) implying that ϕ_n is a σ -module isomorphism. Finally, let us now turn to the question of finding q -affine connections on M_n that are compatible with a given hermitian form. Thus, assume that h is a hermitian form on $(S_q^2)^{n+1}$ for which p_n is an orthogonal projection. According to Proposition 3.6, every q -affine connection ∇_0 on $(S_q^2)^{n+1}$ may be written in the form $\nabla_{X_a}^0 e_\mu = \Gamma_{a\mu}^\nu e_\nu$ with

$$(5.7) \quad \begin{aligned}\tilde{\Gamma}_{+\mu,\nu} &= \frac{1}{2}X_+(h_{\mu\nu}) + K(\alpha_{\mu\nu}) + iK(\beta_{\mu\nu}) \\ \tilde{\Gamma}_{-\mu,\nu} &= \frac{1}{2}X_-(h_{\mu\nu}) + K(\alpha_{\mu\nu}) - iK(\beta_{\mu\nu}) \\ \tilde{\Gamma}_{z\mu,\nu} &= \frac{1}{2}X_z(h_{\mu\nu}) + K^2(\rho_{\mu\nu}),\end{aligned}$$

for an arbitrary choice of hermitian $\alpha, \beta, \rho \in \text{Mat}_{n+1}(S_q^2)$ and, furthermore, Proposition 3.7 shows that $\nabla = p_n \circ \nabla^0$ is a q -affine connection on M_n , compatible with the restriction of h to M_n . For the generators $\{\hat{e}_\mu\}_{\mu=0}^n$ one finds that

$$\begin{aligned}\nabla_{X_+} \hat{e}_\mu &= (p_n \circ \nabla^0)((p_n)_\mu^\nu e_\nu) = p((p_n)_\mu^\nu \Gamma_{+\nu}^\kappa e_\kappa + X_+((p_n)_\mu^\nu) \hat{\sigma}_+(e_\nu)) \\ &= \left((p_n)_\mu^\nu \Gamma_{+\nu}^\kappa + q^{n-2\mu} X_+((p_n)_\mu^\kappa) \right) \hat{e}_\kappa \\ \nabla_{X_-} \hat{e}_\mu &= \left((p_n)_\mu^\nu \Gamma_{-\nu}^\kappa + q^{n-2\mu} X_-((p_n)_\mu^\kappa) \right) \hat{e}_\kappa \\ \nabla_{X_z} \hat{e}_\mu &= \left((p_n)_\mu^\nu \Gamma_{z\nu}^\kappa + q^{2(n-2\mu)} X_z((p_n)_\mu^\kappa) \right) \hat{e}_\kappa.\end{aligned}$$

In the current case, one finds that

$$\begin{aligned}\tilde{h}_{\pm\mu\nu} &= h((\hat{\sigma}_\pm^0)^*(e_\mu), e_\nu) = q^{2\mu-n} h_{\mu\nu} \quad \Rightarrow \quad \tilde{h}_\pm^{\mu\nu} = h^{\mu\nu} q^{n-2\nu} \\ \tilde{h}_{z\mu\nu} &= h((\hat{\sigma}_z^0)^*(e_\mu), e_\nu) = q^{2(2\mu-n)} h_{\mu\nu} \quad \Rightarrow \quad \tilde{h}_z^{\mu\nu} = h^{\mu\nu} q^{2(n-2\nu)}\end{aligned}$$

giving

$$\begin{aligned}\Gamma_{\pm\mu}^\nu &= \tilde{\Gamma}_{\pm\mu,\rho} \sigma_\pm(\tilde{h}_\pm^{\rho\mu}) = q^{2\nu-n} \tilde{\Gamma}_{\pm\mu,\rho} K^2(h^{\rho\nu}) \\ \Gamma_{z\mu}^\nu &= \tilde{\Gamma}_{z\mu,\rho} \sigma_z(\tilde{h}_z^{\rho\nu}) = q^{2(2\nu-m)} \tilde{\Gamma}_{z\mu,\rho} K^4(h^{\rho\nu})\end{aligned}$$

Thus, for the choice $\alpha = \beta = \rho = 0$ in (5.7), one obtains the formulas

$$\begin{aligned}\nabla_{X_\pm} \hat{e}_\mu &= \left(\frac{1}{2} q^{2\kappa-n} (p_n)_\mu^\nu X_\pm(h_{\nu\rho}) K^2(h^{\rho\kappa}) + q^{n-2\mu} X_\pm((p_n)_\mu^\kappa) \right) \hat{e}_\kappa \\ \nabla_{X_z} \hat{e}_\mu &= \left(\frac{1}{2} q^{2(2\kappa-n)} (p_n)_\mu^\nu X_z(h_{\nu\rho}) K^4(h^{\rho\kappa}) + q^{2(n-2\mu)} X_z((p_n)_\mu^\kappa) \right) \hat{e}_\kappa.\end{aligned}$$

6. COMPUTATION OF CHRISTOFFEL SYMBOLS

In this section we work out an explicit expression for a metric and torsion free connection on $\Omega^1(S_q^3)$, as presented in Section 4, for the case when the metric is invariant by K , that is $K(h_{ab}) = h_{ab}$, and $K(\omega_a) = \omega_a$ for $a, b = 1, 2, 3$. As previously noted, for such a metric, $\tilde{\Gamma}_{ab,c} = \Gamma_{ab}^p h_{pc}$ and the torsion free equations become (cf. (4.7)-(4.9)):

$$\begin{aligned}\tilde{\Gamma}_{-,a} - q^2 \tilde{\Gamma}_{+,-,a} &= h_{za} \\ q^2 \tilde{\Gamma}_{z-,a} - q^{-2} \tilde{\Gamma}_{-z,a} &= (1 + q^2) h_{-a} \\ q^2 \tilde{\Gamma}_{+z,a} - q^{-2} \tilde{\Gamma}_{z+,a} &= (1 + q^2) h_{+a}.\end{aligned}$$

These can be solved as

$$(6.1) \quad \tilde{\Gamma}_{-,a} = \frac{1}{2} h_{za} + qK(\gamma_a^{(1)})$$

$$(6.2) \quad \tilde{\Gamma}_{+,-,a} = -\frac{1}{2} q^{-2} h_{za} + q^{-1} K(\gamma_a^{(1)})$$

$$(6.3) \quad \tilde{\Gamma}_{z-,a} = \frac{1}{2} q^{-2} (1 + q^2) h_{-a} + q^{-2} K(\gamma_a^{(2)})$$

$$(6.4) \quad \tilde{\Gamma}_{-z,a} = -\frac{1}{2} q^2 (1 + q^2) h_{-a} + q^2 K(\gamma_a^{(2)})$$

$$(6.5) \quad \tilde{\Gamma}_{+z,a} = \frac{1}{2} q^{-2} (1 + q^2) h_{+a} + q^{-2} K(\gamma_a^{(3)})$$

$$(6.6) \quad \tilde{\Gamma}_{z+,a} = -\frac{1}{2} q^2 (1 + q^2) h_{+a} + q^2 K(\gamma_a^{(3)})$$

for arbitrary $\gamma_a^{(1)}, \gamma_a^{(2)}, \gamma_a^{(3)} \in S_q^3$. On the other hand, Proposition 3.6 gives general expressions for $\tilde{\Gamma}_{ab,c}$ for a metric connection, and combining the two results yields the following set of equations to be solved:

$$(6.7) \quad \frac{1}{2} X_-(h_{+a}) + K(\gamma_{a+}^*) = \frac{1}{2} h_{za} + qK(\gamma_a^{(1)})$$

$$(6.8) \quad \frac{1}{2} X_+(h_{-a}) + K(\gamma_{-a}) = -\frac{1}{2} q^{-2} h_{za} + q^{-1} K(\gamma_a^{(1)})$$

$$(6.9) \quad \frac{1}{2} X_z(h_{-a}) + K^2(\rho_{-a}) = \frac{1}{2} q^{-2} (1 + q^2) h_{-a} + q^{-2} K(\gamma_a^{(2)})$$

$$(6.10) \quad \frac{1}{2} X_-(h_{za}) + K(\gamma_{az}^*) = -\frac{1}{2} q^2 (1 + q^2) h_{-a} + q^2 K(\gamma_a^{(2)})$$

$$(6.11) \quad \frac{1}{2} X_+(h_{za}) + K(\gamma_{za}) = \frac{1}{2} q^{-2} (1 + q^2) h_{+a} + q^{-2} K(\gamma_a^{(3)})$$

$$(6.12) \quad \frac{1}{2} X_z(h_{+a}) + K^2(\rho_{+a}) = -\frac{1}{2} q^2 (1 + q^2) h_{+a} + q^2 K(\gamma_a^{(3)})$$

where $\gamma_{ab} = \alpha_{ab} + i\beta_{ab}$. These equations can be rewritten as

$$(6.13) \quad \gamma_{-a} = -\frac{1}{2} K^{-1} X_+(h_{-a}) - \frac{1}{2} q^{-2} h_{za} + q^{-1} \gamma_a^{(1)}$$

$$(6.14) \quad \gamma_{a+} = \frac{1}{2} K^{-1} X_+(h_{a+}) + \frac{1}{2} h_{az} + q(\gamma_a^{(1)})^*$$

$$(6.15) \quad \rho_{-a} = -\frac{1}{2} K^{-2} X_z(h_{-a}) + \frac{1}{2} q^{-2} (1 + q^2) h_{-a} + q^{-2} K^{-1}(\gamma_a^{(2)})$$

$$(6.16) \quad \gamma_{az} = \frac{1}{2} K^{-1} X_+(h_{az}) - \frac{1}{2} q^2 (1 + q^2) h_{a-} + q^2 (\gamma_a^{(2)})^*$$

$$(6.17) \quad \gamma_{za} = -\frac{1}{2} K^{-1} X_+(h_{za}) + \frac{1}{2} q^{-2} (1 + q^2) h_{+a} + q^{-2} \gamma_a^{(3)}$$

$$(6.18) \quad \rho_{+a} = -\frac{1}{2} K^{-2} X_z(h_{+a}) - \frac{1}{2} q^2 (1 + q^2) h_{+a} + q^2 K^{-1}(\gamma_a^{(3)}).$$

When solving for γ_{ab} and ρ_{ab} from the above equations, one finds several ambiguities and constraints. Namely, there are multiple expressions for γ_{-+} , γ_{-z} , γ_{z+} and γ_{zz} , and we require

(from Proposition 3.6) that $\rho_{++}^* = \rho_{++}$, $\rho_{--}^* = \rho_{--}$ and $\rho_{+-}^* = \rho_{-+}$. Let us consider these constraints one by one.

γ_{-+} : Equations (6.13) and (6.14) give two expressions for γ_{-+} :

$$\begin{aligned}\gamma_{-+} &= -\frac{1}{2}K^{-1}X_+(h_{-+}) - \frac{1}{2}q^{-2}h_{z+} + q^{-1}\gamma_+^{(1)} \\ \gamma_{-+} &= \frac{1}{2}K^{-1}X_+(h_{-+}) + \frac{1}{2}h_{-z} + q(\gamma_-^{(1)})^*\end{aligned}$$

which coincide if

$$\begin{aligned}\gamma_+^{(1)} &= \frac{1}{2}qK^{-1}X_+(h_{-+}) + \frac{1}{2}q^{-1}h_{z+} + q\tau_1 \\ \gamma_-^{(1)} &= -\frac{1}{2}q^{-1}K^{-1}X_-(h_{+-}) - \frac{1}{2}q^{-1}h_{z-} + q^{-1}\tau_1^*\end{aligned}$$

for arbitrary $\tau_1 \in S_q^3$, giving $\gamma_{-+} = \tau_1$.

γ_{-z} : Equations (6.13) and (6.16) give two expressions for γ_{-z} :

$$\begin{aligned}\gamma_{-z} &= -\frac{1}{2}K^{-1}X_+(h_{-z}) - \frac{1}{2}q^{-2}h_{zz} + q^{-1}\gamma_z^{(1)} \\ \gamma_{-z} &= \frac{1}{2}K^{-1}X_+(h_{-z}) - \frac{1}{2}q^2(1+q^2)h_{--} + q^2(\gamma_-^{(2)})^*\end{aligned}$$

which coincide if

$$(6.19) \quad \begin{aligned}\gamma_z^{(1)} &= \frac{1}{2}qK^{-1}X_+(h_{-z}) + \frac{1}{2}q^{-1}h_{zz} + q\tau_2 \\ \gamma_-^{(2)} &= \frac{1}{2}q^{-2}K^{-1}X_-(h_{z-}) + \frac{1}{2}(1+q^2)h_{--} + q^{-2}\tau_2^*\end{aligned}$$

for arbitrary $\tau_2 \in S_q^3$, giving $\gamma_{-z} = \tau_2$.

γ_{z+} : Equations (6.14) and (6.17) give two expressions for γ_{z+} :

$$\begin{aligned}\gamma_{z+} &= \frac{1}{2}K^{-1}X_+(h_{z+}) + \frac{1}{2}h_{zz} + q(\gamma_z^{(1)})^* \\ \gamma_{z+} &= -\frac{1}{2}K^{-1}X_+(h_{z+}) + \frac{1}{2}q^{-2}(1+q^2)h_{++} + q^{-2}\gamma_+^{(3)}\end{aligned}$$

which coincide if

$$(6.20) \quad \begin{aligned}\gamma_z^{(1)} &= \frac{1}{2}q^{-1}K^{-1}X_-(h_{+z}) - \frac{1}{2}q^{-1}h_{zz} + q^{-1}\tau_3^* \\ \gamma_+^{(3)} &= \frac{1}{2}q^2K^{-1}X_+(h_{z+}) - \frac{1}{2}(1+q^2)h_{++} + q^2\tau_3\end{aligned}$$

for arbitrary $\tau_3 \in S_q^3$, giving $\gamma_{z+} = \tau_3$.

γ_{zz} : Equations (6.16) and (6.17) give two expressions for γ_{zz} :

$$\begin{aligned}\gamma_{zz} &= \frac{1}{2}K^{-1}X_+(h_{zz}) - \frac{1}{2}q^2(1+q^2)h_{z-} + q^2(\gamma_z^{(2)})^* \\ \gamma_{zz} &= -\frac{1}{2}K^{-1}X_+(h_{zz}) + \frac{1}{2}q^{-2}(1+q^2)h_{+z} + q^{-2}\gamma_z^{(3)}\end{aligned}$$

which coincide if

$$\begin{aligned}\gamma_z^{(2)} &= \frac{1}{2}q^{-2}K^{-1}X_-(h_{zz}) + \frac{1}{2}(1+q^2)h_{-z} + q^{-2}\tau_4^* \\ \gamma_z^{(3)} &= \frac{1}{2}q^2K^{-1}X_+(h_{zz}) - \frac{1}{2}(1+q^2)h_{+z} + q^2\tau_4\end{aligned}$$

for arbitrary $\tau_4 \in S_q^3$, giving $\gamma_{zz} = \tau_4$.

The above considerations determine $\gamma_+^{(1)}$, $\gamma_-^{(1)}$, $\gamma_-^{(2)}$, $\gamma_+^{(3)}$, $\gamma_z^{(2)}$, $\gamma_z^{(3)}$, $\gamma_z^{(1)}$, but give two expressions for $\gamma_z^{(1)}$; equating them gives the following.

$\gamma_z^{(1)}$: Equations (6.19) and (6.20) give two expressions for $\gamma_z^{(1)}$:

$$\begin{aligned}\gamma_z^{(1)} &= \frac{1}{2}qK^{-1}X_+(h_{-z}) + \frac{1}{2}q^{-1}h_{zz} + q\tau_2 \\ \gamma_z^{(1)} &= \frac{1}{2}q^{-1}K^{-1}X_-(h_{+z}) - \frac{1}{2}q^{-1}h_{zz} + q^{-1}\tau_3^*\end{aligned}$$

which coincide if

$$\begin{aligned}\tau_2 &= -\frac{1}{2}K^{-1}X_+(h_{-z}) - \frac{1}{2}q^{-2}h_{zz} + q^{-1}\mu_1 \\ \tau_3 &= \frac{1}{2}K^{-1}X_+(h_{z+}) + \frac{1}{2}h_{zz} + q\mu_1^*\end{aligned}$$

for arbitrary $\mu_1 \in S_q^3$, giving $\gamma_z^{(1)} = \mu_1$ and

$$\begin{aligned}\gamma_-^{(2)} &= q^{-2}K^{-1}X_-(h_{z-}) + \frac{1}{2}(1+q^2)h_{--} - \frac{1}{2}q^{-4}h_{zz} + q^{-3}\mu_1^* \\ \gamma_+^{(3)} &= q^2K^{-1}X_+(h_{z+}) - \frac{1}{2}(1+q^2)h_{++} + \frac{1}{2}q^2h_{zz} + q^3\mu_1^*.\end{aligned}$$

Let us summarize what we have obtained so far:

$$(6.21) \quad \gamma_+^{(1)} = \frac{1}{2}qK^{-1}X_+(h_{-+}) + \frac{1}{2}q^{-1}h_{z+} + q\tau_1$$

$$(6.22) \quad \gamma_-^{(1)} = -\frac{1}{2}q^{-1}K^{-1}X_-(h_{+-}) - \frac{1}{2}q^{-1}h_{z-} + q^{-1}\tau_1^*$$

$$(6.23) \quad \gamma_z^{(1)} = \mu_1$$

$$(6.24) \quad \gamma_-^{(2)} = q^{-2}K^{-1}X_-(h_{z-}) + \frac{1}{2}(1+q^2)h_{--} - \frac{1}{2}q^{-4}h_{zz} + q^{-3}\mu_1^*$$

$$(6.25) \quad \gamma_+^{(3)} = q^2K^{-1}X_+(h_{z+}) - \frac{1}{2}(1+q^2)h_{++} + \frac{1}{2}q^2h_{zz} + q^3\mu_1^*$$

$$(6.26) \quad \gamma_z^{(2)} = \frac{1}{2}q^{-2}K^{-1}X_-(h_{zz}) + \frac{1}{2}(1+q^2)h_{-z} + q^{-2}\tau_4^*$$

$$(6.27) \quad \gamma_z^{(3)} = \frac{1}{2}q^2K^{-1}X_+(h_{zz}) - \frac{1}{2}(1+q^2)h_{+z} + q^2\tau_4,$$

for arbitrary μ_1, τ_1, τ_4 . It remains to solve the constraints $\rho_{++}^* = \rho_{++}$, $\rho_{--}^* = \rho_{--}$ and $\rho_{+-}^* = \rho_{-+}$.

ρ_{++} : From equation (6.18) one obtains

$$\rho_{++} = -\frac{1}{2}q^2(1+q^2)h_{++} + q^2K^{-1}(\gamma_+^{(3)})$$

by using that $X_z(h_{ab}) = 0$. Requiring $\rho_{++}^* = \rho_{++}$ gives

$$\gamma_+^{(3)} = K^2((\gamma_+^{(3)})^*)$$

and inserting (6.25) yields

$$K^2(\mu_1) - \mu_1^* = q^{-1}K^{-1}X_+(h_{z+}) + q^{-1}KX_-(h_{+z})$$

which is solved by

$$(6.28) \quad \mu_1 = q^{-1}K^{-1}X_-(h_{+z}) + K^{-1}(f_1)$$

for arbitrary hermitian $f_1 \in S_q^3$.

ρ_{--} : From equation (6.15) one obtains

$$\rho_{--} = \frac{1}{2}q^{-2}(1+q^2)h_{--} + q^{-2}K^{-1}(\gamma_-^{(2)})$$

by using that $X_z(h_{ab}) = 0$. Requiring $\rho_{--}^* = \rho_{--}$ gives

$$\gamma_-^{(2)} = K^2((\gamma_-^{(2)})^*)$$

and inserting (6.24) gives

$$K^2(\mu_1) - \mu_1^* = qK^{-1}X_-(h_{z-}) + qKX_+(h_{-z})$$

which is solved by

$$(6.29) \quad \mu_1 = qK^{-1}X_+(h_{-z}) + K^{-1}(f_2)$$

for arbitrary hermitian $f_2 \in S_q^3$.

Equations (6.28) and (6.29) give two expressions for μ_1 , which coincide if

$$(6.30) \quad f_1 - f_2 = qX_+(h_{-z}) - q^{-1}X_-(h_{+z}).$$

Since $f_1 - f_2$ is hermitian, a necessary condition for this equation to have a solution is that

$$H = qX_+(h_{-z}) - q^{-1}X_-(h_{+z})$$

is hermitian. Using $K^{-1}X_{\pm} = q^{\mp 1}X_{\pm}K^{-1}$ and $K^{-1}(h_{ab}) = h_{ab}$, $H = H^*$ is equivalent to

$$(6.31) \quad X_+(h_{-z} - q^{-4}h_{z+}) = -q^2X_-(h_{z-} - q^{-4}h_{+z}).$$

which can also be written as the reality condition

$$(6.32) \quad (X_+(h_{-z} - q^{-4}h_{z+}))^* = X_+(h_{-z} - q^{-4}h_{z+}).$$

If the above condition is fulfilled, then (6.30) is solved by

$$\begin{aligned} f_1 &= \frac{1}{2}qX_+(h_{-z}) - \frac{1}{2}q^{-1}X_-(h_{+z}) + f_0 \\ f_2 &= -\frac{1}{2}qX_+(h_{-z}) + \frac{1}{2}q^{-1}X_-(h_{+z}) + f_0 \end{aligned}$$

for arbitrary hermitian $f_0 \in S_q^3$. This gives

$$(6.33) \quad \mu_1 = \frac{1}{2}qK^{-1}X_+(h_{-z}) + \frac{1}{2}q^{-1}K^{-1}X_-(h_{+z}) + K^{-1}(f_0).$$

ρ_{+-} : Equations (6.15) and (6.18) give

$$\begin{aligned} \rho_{+-} &= -\frac{1}{2}q^2(1+q^2)h_{+-} + q^2K^{-1}(\gamma_-^{(3)}) \\ \rho_{-+} &= \frac{1}{2}q^{-2}(1+q^2)h_{-+} + q^{-2}K^{-1}(\gamma_+^{(2)}) \end{aligned}$$

and requiring $\rho_{+-}^* = \rho_{-+}$ yields

$$-\frac{1}{2}q^2(1+q^2)h_{-+} + q^2K((\gamma_-^{(3)})^*) = \frac{1}{2}q^{-2}(1+q^2)h_{-+} + q^{-2}K^{-1}(\gamma_+^{(2)})$$

which is solved by

$$\begin{aligned} \gamma_-^{(3)} &= \frac{1}{2}(1+q^2)h_{+-} + q^{-2}K(\mu_2^*) \\ \gamma_+^{(2)} &= -\frac{1}{2}(1+q^2)h_{-+} + q^2K(\mu_2) \end{aligned}$$

for arbitrary $\mu_2 \in S_q^3$.

Thus, we have obtained the following equations

$$\begin{aligned}
\gamma_+^{(1)} &= \frac{1}{2}qK^{-1}X_+(h_{-+}) + \frac{1}{2}q^{-1}h_{z+} + q\tau_1 \\
\gamma_-^{(1)} &= -\frac{1}{2}q^{-1}K^{-1}X_-(h_{+-}) - \frac{1}{2}q^{-1}h_{z-} + q^{-1}\tau_1^* \\
\gamma_z^{(1)} &= \frac{1}{2}qK^{-1}X_+(h_{-z}) + \frac{1}{2}q^{-1}K^{-1}X_-(h_{+z}) + K^{-1}(f_0) \\
\gamma_+^{(2)} &= -\frac{1}{2}(1+q^2)h_{-+} + q^2K(\mu_2) \\
\gamma_-^{(2)} &= \frac{1}{2}q^{-2}K^{-1}X_-(h_{z-}) - \frac{1}{2}q^{-4}K^{-1}X_+(h_{z+}) + \frac{1}{2}(1+q^2)h_{--} - \frac{1}{2}q^{-4}h_{zz} + q^{-3}K(f_0) \\
\gamma_z^{(2)} &= \frac{1}{2}q^{-2}K^{-1}X_-(h_{zz}) + \frac{1}{2}(1+q^2)h_{-z} + q^{-2}\tau_4^* \\
\gamma_+^{(3)} &= \frac{1}{2}q^2K^{-1}X_+(h_{z+}) - \frac{1}{2}q^4K^{-1}X_-(h_{z-}) - \frac{1}{2}(1+q^2)h_{++} + \frac{1}{2}q^2h_{zz} + q^3K(f_0) \\
\gamma_-^{(3)} &= \frac{1}{2}(1+q^2)h_{+-} + q^{-2}K(\mu_2^*) \\
\gamma_z^{(3)} &= \frac{1}{2}q^2K^{-1}X_+(h_{zz}) - \frac{1}{2}(1+q^2)h_{+z} + q^2\tau_4
\end{aligned}$$

giving all 27 Christoffel symbols as

$$\begin{aligned}
\tilde{\Gamma}_{++,+} &= X_+(h_{++}) - \frac{1}{2}q^2X_-(h_{+-}) + h_{+z} + q^2K(\tau_1) \\
\tilde{\Gamma}_{+,-} &= -\frac{1}{2}q^{-2}X_-(h_{+-}) - q^{-2}h_{z-} + q^{-2}K(\tau_1^*) \\
\tilde{\Gamma}_{+z,z} &= \frac{1}{2}X_+(h_{zz}) + K(\tau_4) \\
\tilde{\Gamma}_{++,-} &= \frac{1}{2}X_+(h_{+-}) + K(\gamma_{+-}) \\
\tilde{\Gamma}_{+-,+} &= \frac{1}{2}X_+(h_{-+}) + K(\tau_1) \\
\tilde{\Gamma}_{+z,+} &= \frac{1}{2}X_+(h_{z+}) - \frac{1}{2}q^2X_-(h_{z-}) + \frac{1}{2}h_{zz} + qK^2(f_0) \\
\tilde{\Gamma}_{++z} &= X_+(h_{+z}) - q^2(1+q^2)h_{+-} + q^4\mu_2^* \\
\tilde{\Gamma}_{+z,-} &= q^{-2}(1+q^2)h_{+-} + q^{-4}K^2(\mu_2^*) \\
\tilde{\Gamma}_{+-,z} &= \frac{1}{2}X_+(h_{-z}) + \frac{1}{2}q^{-2}X_-(h_{+z}) - \frac{1}{2}q^{-2}h_{zz} + q^{-1}f_0 \\
\tilde{\Gamma}_{-+,+} &= \frac{1}{2}q^2X_+(h_{-+}) + h_{z+} + q^2K(\tau_1^*) \\
\tilde{\Gamma}_{--,-} &= X_-(h_{--}) + \frac{1}{2}q^{-2}X_+(h_{-+}) - q^{-2}h_{-z} + q^{-2}K(\tau_1) \\
\tilde{\Gamma}_{-z,z} &= \frac{1}{2}X_-(h_{zz}) + K(\tau_4^*) \\
\tilde{\Gamma}_{-+,-} &= \frac{1}{2}X_-(h_{+-}) + K(\tau_1^*) \\
\tilde{\Gamma}_{--,+} &= \frac{1}{2}X_-(h_{-+}) + K(\gamma_{+-}^*) \\
\tilde{\Gamma}_{-+,z} &= \frac{1}{2}X_-(h_{+z}) + \frac{1}{2}q^2X_+(h_{-z}) + \frac{1}{2}h_{zz} + qf_0 \\
\tilde{\Gamma}_{-z,+} &= -q^2(1+q^2)h_{-+} + q^{-1}K^2(\mu_2) \\
\tilde{\Gamma}_{--z} &= X_-(h_{-z}) + q^2(1+q^2)h_{-+} + q^{-4}\mu_2 \\
\tilde{\Gamma}_{-z,-} &= \frac{1}{2}X_-(h_{z-}) - \frac{1}{2}q^{-2}X_+(h_{z+}) - \frac{1}{2}q^{-2}h_{zz} + q^{-1}K^2(f_0)
\end{aligned}$$

$$\begin{aligned}
\tilde{\Gamma}_{z+,+} &= \frac{1}{2}q^4 X_+(h_{z+}) - \frac{1}{2}q^6 X_-(h_{z-}) - q^2(1+q^2)h_{++} + \frac{1}{2}q^4 h_{zz} + q^5 K^2(f_0) \\
\tilde{\Gamma}_{z-,-} &= \frac{1}{2}q^{-4} X_-(h_{z-}) - \frac{1}{2}q^{-6} X_+(h_{z+}) + q^{-2}(1+q^2)h_{--} - \frac{1}{2}q^{-6} h_{zz} + q^{-5} K^2(f_0) \\
\tilde{\Gamma}_{zz,z} &= K^2(\rho_{zz}) \\
\tilde{\Gamma}_{z+,-} &= K^2(\mu_2^*) \\
\tilde{\Gamma}_{z-,+} &= K^2(\mu_2) \\
\tilde{\Gamma}_{z+,z} &= \frac{1}{2}q^4 X_+(h_{zz}) - q^2(1+q^2)h_{+z} + q^4 K(\tau_4) \\
\tilde{\Gamma}_{zz,+} &= -\frac{1}{2}q^2 X_-(h_{zz}) - q^2(1+q^2)h_{z+} + q^4 K^3(\tau_4^*) \\
\tilde{\Gamma}_{z-,z} &= \frac{1}{2}q^{-4} X_-(h_{zz}) + q^{-2}(1+q^2)h_{-z} + q^{-4} K(\tau_4^*) \\
\tilde{\Gamma}_{zz,-} &= -\frac{1}{2}q^{-2} X_+(h_{zz}) + q^{-2}(1+q^2)h_{z-} + q^{-4} K^3(\tau_4)
\end{aligned}$$

depending on the parameters $\tau_1, \tau_4, \mu_2, \gamma_{+-}, \rho_{zz}, f_0 \in S_q^3$ (with $f_0^* = f_0$).

Acknowledgements: The paper is partially supported by INFN-Trieste. GL is supported by INFN, Iniziativa Specifica GAST, and by INDAM - GNSAGA. JA is supported by grant 2017-03710 from the Swedish Research Council. Furthermore, JA would like to thank the Department of Mathematics and Geosciences, University of Trieste for hospitality.

REFERENCES

- [AC10] P. Aschieri and L. Castellani. Noncommutative gravity solutions. *J. Geom. Phys.*, 60(3):375–393, 2010.
- [AW17] J. Arnlind and M. Wilson. Riemannian curvature of the noncommutative 3-sphere. *J. Noncommut. Geom.*, 11(2):507–536, 2017.
- [BGM19] J. Bhowmick, D. Goswami, and S. Mukhopadhyay. Levi-civita connections for a class of spectral triples. *Lett. Math. Phys.*, 110:835–884, 2019.
- [BGL20] J. Bhowmick, D. Goswami, G. Landi. On the Koszul formula in noncommutative geometry. *Rev. Math. Phys.* in press; arXiv:1910.09306
- [BM98] T. Brzeziński and S. Majid. Line bundles on quantum spheres. In *Particles, fields, and gravitation (Łódź, 1998)*, vol. 453 *AIP Conf. Proc.*, pp 3–8. AIP, Woodbury, NY, 1998.
- [BM11] E. J. Beggs and S. Majid. *-compatible connections in noncommutative Riemannian geometry. *J. Geom. Phys.*, 61(1):95–124, 2011.
- [CFF93] A. H. Chamseddine, G. Felder, and J. Fröhlich. Gravity in noncommutative geometry. *Comm. Math. Phys.*, 155(1):205–217, 1993.
- [DV88] M. Dubois-Violette. Dérivations et calcul différentiel non commutatif. *C. R. Acad. Sci. Paris Sér. I Math.*, 307(8):403–408, 1988.
- [DVMMM96] M. Dubois-Violette, J. Madore, T. Masson, and J. Mourad. On curvature in noncommutative geometry. *J. Math. Phys.*, 37(8):4089–4102, 1996.
- [HM99] P. M. Hajac and S. Majid. Projective module description of the q -monopole. *Comm. Math. Phys.*, 206(2):247–264, 1999.
- [Lan18] G. Landi. Twisted sigma-model solitons on the quantum projective line. *Lett. Math. Phys.*, 108(8):1955–1983, 2018.
- [Maj05] S. Majid. Noncommutative Riemannian and spin geometry of the standard q -sphere. *Comm. Math. Phys.*, 256(2):255–285, 2005.
- [MNW91] T. Masuda, Y. Nakagami, and J. Watanabe. Noncommutative differential geometry on the quantum two sphere of Podleś. I. An algebraic viewpoint. *K-Theory*, 5(2):151–175, 1991.

- [Pod87] P. Podleś. Quantum spheres. *Lett. Math. Phys.*, 14(3):193–202, 1987.
[Ros13] J. Rosenberg. Levi-Civita’s theorem for noncommutative tori. *SIGMA*, 9:071, 2013.
[Wor87] S. L. Woronowicz. Twisted $SU(2)$ group. An example of a noncommutative differential calculus. *Publ. Res. Inst. Math. Sci.*, 23(1):117–181, 1987.

(Joakim Arnlind) DEPT. OF MATH., LINKÖPING UNIVERSITY, 581 83 LINKÖPING, SWEDEN

E-mail address: joakim.arnlind@liu.se

(Kwalombota Ilwale) DEPT. OF MATH., LINKÖPING UNIVERSITY, 581 83 LINKÖPING, SWEDEN

E-mail address: kwalombota.ilwale@liu.se

(Giovanni Landi) MATEMATICA, UNIVERSITÀ DI TRIESTE,

VIA A. VALERIO, 12/1, 34127 TRIESTE, ITALY

INSTITUTE FOR GEOMETRY AND PHYSICS (IGAP) TRIESTE, ITALY AND INFN, TRIESTE, ITALY

E-mail address: landi@units.it