

Universal Coding and Prediction on Ergodic Martin-Löf Random Points

Łukasz Dębowski and Tomasz Steifer*

Abstract

Suppose that we have a method which estimates the conditional probabilities of some unknown stochastic source and we use it to guess which of the outcomes will happen. We want to make a correct guess as often as it is possible. What estimators are good for this? In this work, we consider estimators given by a familiar notion of universal coding for stationary ergodic measures, while working in the framework of algorithmic randomness, i.e, we are particularly interested in prediction of Martin-Löf random points. We outline the general theory and exhibit some counterexamples. Completing a result of Ryabko from 2009 we also show that universal probability measure in the sense of universal coding induces a universal predictor in the prequential sense. Surprisingly, this implication holds true provided the universal measure does not ascribe too low conditional probabilities to individual symbols. As an example, we show that the Prediction by Partial Matching (PPM) measure satisfies this requirement with a large reserve.

Keywords: algorithmic randomness; stationary ergodic processes; universal coding; universal prediction; prediction by partial matching

MSC 2000: 94A29, 62M20, 03D32

1 Introduction

A sequence of outcomes X_1, X_2, \dots coming from a finite alphabet is drawn in a sequential manner from an unknown stochastic source P . At each moment a finite prefix $X_1^n = (X_1, X_2, \dots, X_n)$ is available. The forecaster has to predict the next outcome using this information. The task may take one of the two following forms. In the first scenario, the forecaster simply makes a guess about the next outcome. The forecaster's performance is then assessed by comparing the guess with the outcome. This scenario satisfies the weak prequential principle of Dawid [12]. In the second case, we allow the forecaster to be uncertain, namely, we ask them to assign a probability value for each of the outcomes. These values may be interpreted as estimates of the conditional probabilities $P(X_{n+1}|X_1^n)$. Various criteria of success may be chosen here such as the quadratic difference of distributions or the Kullback-Leibler divergence. The key aspect of both

*T. Steifer is with the Institute of Fundamental Technological Research, Polish Academy of Sciences, ul. Pawińskiego 5B, 02-106 Warszawa, Poland (e-mail: tsteifer@ippt.pan.pl). Ł. Dębowski is with the Institute of Computer Science, Polish Academy of Sciences, ul. Jana Kazimierza 5, 01-248 Warszawa, Poland (e-mail: ldebowsk@ipipan.waw.pl).

problems is that we assume limited knowledge about the true probabilities governing the process that we want to forecast. Thus, an admissible solution should achieve the optimal results for an arbitrary process from some general class. For clarity, term “universal predictor” will be used to denote the solution of guessing the outcome, while the solution of estimation of probabilities will be referred to as “universal estimator”, “universal measure”, or “universal code” depending on the exact meaning.

In the literature, it is frequently assumed without explanation that universal predictors and universal estimators reduce to another in some way. Such simplification is not without a rationale. It is known that for a fixed stochastic source P , the optimal prediction is given by the predictor induced by P , i.e., the informed scheme which predicts the outcome with the largest conditional probability $P(X_{n+1}|X_1^n)$ [1]. In particular, a good universal estimator should induce a good universal predictor. That being said, the devil is hidden in the details such as what is meant by a “good” universal code, measure, estimator, or predictor.

In this paper, we will assume that the unknown stochastic source P lies in the class of stationary ergodic measures. Moreover, we are concerned with universal measures which are universal in the information-theoretic sense of universal coding, i.e., the rate of Kullback-Leibler divergence of the estimate and the true measure P vanishes for any stationary ergodic measure. As for universal predictors, we assume that the rate of correct guesses is equal to the respective rate for the predictor induced by measure P . In this setting, a universal measure need not belong to the class of stationary ergodic measures and can be computable, which makes the problem utterly practical. Our framework should be contrasted with universal prediction à la Solomonoff for left-c.e. semimeasures where the universal semimeasure belongs to the class and is not computable [42]. In general, existence of a universal measure for an arbitrary class of probability measures can be linked to separability of the considered class [40].

Now, we can ask the question whether a universal measure in the above sense of universal coding induces a universal predictor. Curiously, this simple question has not been unambiguously answered in the literature, although a host of related propositions were compiled by Suzuki [45] and Ryabko [36, 37], see also [38]. It was shown by Ryabko [37], see also [38], that the expected value of the average absolute difference between the conditional probability for a universal measure and the true value $P(X_{n+1}|X_1^n)$ converges to zero almost surely for any stationary ergodic measure P . Ryabko [37] showed also that there exists a universal measure that induces a universal predictor. As we argue in this paper, this result does not solve the general problem.

Completing works [45, 36, 37], in this paper we will show that any universal measure R in the sense of universal coding that additionally satisfies a uniform bound

$$-\log R(X_{n+1}|X_1^n) \leq \epsilon_n \sqrt{n/\ln n}, \quad \lim_{n \rightarrow \infty} \epsilon_n = 0 \quad (1)$$

induces a universal predictor, indeed. On our way, we will use the Breiman ergodic theorem [7] and the Azuma inequality for martingales with bounded increments [3], which is the source of condition (1). It is left open whether this condition is necessary. Fortunately, condition (1) is satisfied by reasonable universal measures such as the Prediction by Partial Matching (PPM) measure

[10, 39, 36], which we also show in this paper. It may be interesting to exhibit universal measures for which this condition fails. There is a large gap between bound (1) and the respective bound for the PPM measure, which begs for further research.

To add more weight and to make the problem interesting from a computational perspective, we consider this topic in the context of algorithmic randomness and we seek for effective versions of probabilistic statements. Effectivization is meant as the research program of reformulating almost sure statements into respective statements about algorithmically random points, i.e., algorithmically random infinite sequences. Any plausible class of random points is of measure one, see [18], and the effective versions of theorems substitute phrase “almost surely” with “on all algorithmically random points”. Usually, randomness in the Martin-Löf sense is the desired goal [31]. In many cases, the standard proofs are already constructive, effectivization of some theorems asks for developing new proofs, but sometimes the effective versions are false.

In this paper, we will successfully show that the algorithmic randomness theory is mature enough to make the theory of universal coding and prediction for stationary ergodic sources effective in the Martin-Löf sense. The main keys to this success are: the framework for randomness with respect to uncomputable measures by Reimann and Slaman [34, 35], the effective Birkhoff ergodic theorem [47, 6, 21], generalized by us to the version resembling Breiman’s ergodic theorem [7], and an effective Azuma theorem, which follows from a result of Solovay (unpublished, see [18])—which we call here the effective Borel-Cantelli lemma—and the Azuma inequality [3]. As a little surprise there is also a negative result concerning universal forward estimators—Theorem 9. Not everything can be made effective.

The organization of the paper is as follows. In Section 2, we discuss preliminaries: notation (Subsection 2.1), stationary and ergodic measures (Subsection 2.2), algorithmic randomness (Subsection 2.3), and some known effectivizations (Subsection 2.4). Section 3 contains main results concerning: universal coding (Subsection 3.1), universal prediction (Subsection 3.2), universal predictors induced by universal backward estimators (Subsection 3.3) and by universal codes (Subsection 3.4), as well as the PPM measure (Subsection 3.5), which constitutes a simple example of a universal code and a universal predictor.

2 Preliminaries

In this section we familiarize the readers with our notation, we recall the concepts of stationary and ergodic measures, we discuss various sorts of algorithmic randomness, and we recall known facts from the effectivization program.

2.1 Notation

Throughout this paper, we consider the standard measurable space $(\mathbb{X}^{\mathbb{Z}}, \mathcal{X}^{\mathbb{Z}})$ of two-sided infinite sequences over a finite alphabet $\mathbb{X} = \{a_1, \dots, a_D\}$, where $D \geq 2$. (Occasionally, we also apply the space of one-sided infinite sequences $(\mathbb{X}^{\mathbb{N}}, \mathcal{X}^{\mathbb{N}})$.) The points of the space are (infinite) sequences $x = (x_i)_{i \in \mathbb{Z}} \in \mathbb{X}^{\mathbb{Z}}$. We also denote (finite) strings $x_j^k = (x_i)_{j \leq i \leq k}$, where $x_j^{j-1} = \lambda$ equals the empty string. By $\mathbb{X}^* = \bigcup_{n \geq 0} \mathbb{X}^n$ we denote the set of strings of an arbitrary length

including the singleton $\mathbb{X}^0 = \{\lambda\}$. We use random variables $X_k((x_i)_{i \in \mathbb{Z}}) := x_k$. Having these, the σ -field $\mathcal{X}^{\mathbb{Z}}$ is generated by cylinder sets $(X_{-|\sigma|+1}^{|\tau|} = \sigma\tau)$ for all $\sigma, \tau \in \mathbb{X}^*$. We tacitly assume that P and R denote probability measures on $(\mathbb{X}^{\mathbb{Z}}, \mathcal{X}^{\mathbb{Z}})$. For any probability measure P , we use the shorthand notations $P(x_1^n) := P(X_1^n = x_1^n)$ and $P(x_j^n | x_1^{j-1}) := P(X_j^n = x_j^n | X_1^{j-1} = x_1^{j-1})$. Notation $\log x$ denotes the binary logarithm, whereas $\ln x$ is the natural logarithm.

2.2 Stationary and ergodic measures

Let us denote the measurable shift operation $T((x_i)_{i \in \mathbb{Z}}) := (x_{i+1})_{i \in \mathbb{Z}}$ for two-sided infinite sequences $(x_i)_{i \in \mathbb{Z}} \in \mathbb{X}^{\mathbb{Z}}$.

Definition 1 (stationary measures). *A probability measure P on $(\mathbb{X}^{\mathbb{Z}}, \mathcal{X}^{\mathbb{Z}})$ is called stationary if $P(T^{-1}(A)) = P(A)$ for all events $A \in \mathcal{X}^{\mathbb{Z}}$.*

Definition 2 (ergodic measures). *A probability measure P on $(\mathbb{X}^{\mathbb{Z}}, \mathcal{X}^{\mathbb{Z}})$ is called ergodic if for each event $A \in \mathcal{X}^{\mathbb{Z}}$ such that $T^{-1}(A) = A$ we have either $P(A) = 1$ or $P(A) = 0$.*

The class of stationary ergodic probability measures has various nice properties guaranteed by the collection of fundamental results called ergodic theorems. Typically stationary ergodic measures are not computable (e.g., consider independent biased coin tosses with a common uncomputable bias) but they allow for computable universal coding and prediction on algorithmically random points, as it will be explained in Section 3.

2.3 Sorts of randomness

Now let us discuss some computability notions. In the following, *computably enumerable* is abbreviated as *c.e.* Given a real r , the set $\{q \in \mathbb{Q} : q < r\}$ is called the left cut of r . A real function f with arguments in a countable set is called computable or left-c.e. respectively if the left cuts of $f(\sigma)$ are uniformly computable or c.e. given an enumeration of σ . For an infinite sequence $s \in \mathbb{X}^{\mathbb{Z}}$, we say that real functions f are s -computable or s -left-c.e. if they are computable or left-c.e. with oracle s . Similarly, for a real function f taking arguments in $\mathbb{X}^{\mathbb{Z}}$, we will say that f is s -computable or s -left-c.e. if left cuts of $f(x)$ are uniformly computable or c.e. with oracles $x \oplus s := (\dots, x_{-1}, s_{-1}, x_0, s_0, x_1, s_1, \dots)$. This induces in effect s -computable and s -left-c.e. random variables on $(\mathbb{X}^{\mathbb{Z}}, \mathcal{X}^{\mathbb{Z}})$.

For stationary ergodic measures, we need a definition of algorithmically random points with respect to an arbitrary, i.e., not necessarily computable probability measure on $(\mathbb{X}^{\mathbb{Z}}, \mathcal{X}^{\mathbb{Z}})$. A simple definition thereof was proposed by Reimann [34] and Reimann and Slaman [35]. This definition is equivalent to earlier approaches by Levin [28, 29, 30] and Gács [22] as shown by Day and Miller [13] and we will use it since it leads to straightforward generalizations of the results in Section 2.4. The definition is based on measure representations. Let $\mathcal{P}(\mathbb{X}^{\mathbb{Z}})$ be the space of probability measures on $(\mathbb{X}^{\mathbb{Z}}, \mathcal{X}^{\mathbb{Z}})$. A measure $P \in \mathcal{P}(\mathbb{X}^{\mathbb{Z}})$ is called s -computable if real function $(\sigma, \tau) \mapsto P(X_{-|\sigma|+1}^{|\tau|} = \sigma\tau)$ is s -computable. Similarly, a representation function is a function $\rho : \mathbb{X}^{\mathbb{Z}} \rightarrow \mathcal{P}(\mathbb{X}^{\mathbb{Z}})$ such that real function $(\sigma, \tau, s) \mapsto \rho(s)(X_{-|\sigma|+1}^{|\tau|} = \sigma\tau)$ is computable. Subsequently, we say that an infinite sequence $s \in \mathbb{X}^{\mathbb{Z}}$ is a representation of measure

P if there exists a representation function ρ such that $\rho(s) = P$. We note that any measure P is s -computable for any representation s of P .

We will consider two important sorts of algorithmically random points: Martin-Löf or 1-random points and weakly 2-random points with respect to an arbitrary stationary ergodic measure P on $(\mathbb{X}^{\mathbb{Z}}, \mathcal{X}^{\mathbb{Z}})$. Note that the following notions are typically defined for one-sided infinite sequences over the binary alphabet and computable measures P . In the following parts of this paper, let an infinite sequence $s \in \mathbb{X}^{\mathbb{Z}}$ be a representation of measure P .

Definition 3. A collection of events $U_1, U_2, \dots \in \mathcal{X}^{\mathbb{Z}}$ is called uniformly s -c.e. if and only if there is a collection of sets $V_1, V_2, \dots \subset \mathbb{X}^* \times \mathbb{X}^*$ such that

$$U_i = \left\{ x \in \mathbb{X}^{\mathbb{Z}} : \exists(\sigma, \tau) \in V_i : x_{-|\sigma|+1}^{|\tau|} = \sigma\tau \right\}$$

and sets V_1, V_2, \dots are uniformly s -c.e.

Definition 4 (Martin-Löf test). A uniformly s -c.e. collection of events $U_1, U_2, \dots \in \mathcal{X}^{\mathbb{Z}}$ is called a Martin-Löf (s, P) -test if $P(U_n) \leq 2^{-n}$ for every $n \in \mathbb{N}$.

Definition 5 (Martin-Löf or 1-randomness). A point $x \in \mathbb{X}^{\mathbb{Z}}$ is called Martin-Löf (s, P) -random or 1- (s, P) -random if for each Martin-Löf (s, P) -test U_1, U_2, \dots we have $x \notin \bigcap_{i \geq 1} U_i$. A point is called Martin-Löf P -random or 1- P -random if it is 1- (s, P) -random for some representation s of P .

Subsequently, an event $C \in \mathcal{X}^{\mathbb{Z}}$ is called a $\Sigma_2^0(s)$ event if there exists a uniformly s -c.e. sequence of events U_1, U_2, \dots such that $\mathbb{X}^{\mathbb{Z}} \setminus C = \bigcap_{i \geq 1} U_i$.

Definition 6 (weak 2-randomness). A point $x \in \mathbb{X}^{\mathbb{Z}}$ is called weakly 2- (s, P) -random if x is contained in every $\Sigma_2^0(s)$ event C such that $P(C) = 1$. A point is called weakly 2- P -random if it is weakly 2- (s, P) -random for some representation s of P .

The sets of weakly 2-random points are strictly smaller than the respective sets of 1-random points, see [18].

In general, there is a whole hierarchy of algorithmically random points, such as (weakly) n -random points, where n runs over natural numbers. For our purposes, however, only 1-random points and weakly 2-random points matter since the following proposition sets the baseline for effectivization:

Proposition 1 (folklore). Let Y_1, Y_2, \dots be a sequence of uniformly s -computable random variables. If limit $\lim_{n \rightarrow \infty} Y_n$ exists P -almost surely, then it exists on all weakly 2- (s, P) -random points.

The above proposition is obvious since the set of points on which limit $\lim_{n \rightarrow \infty} Y_n$ exists is a $\Sigma_2^0(s)$ event. The effectivization program aims to strengthen the above claim to 1- P -random points (or even weaker notions such as Schnorr randomness) but this need not always be feasible. In particular, one can observe that:

Proposition 2 (folklore). Let P be a non-atomic computable measure on $\mathbb{X}^{\mathbb{N}}$. Then there exists a computable function $f : \mathbb{X}^* \rightarrow \{0, 1\}$ such that the limit $\lim_{n \rightarrow \infty} f(X_1^n)$ exists and is equal zero P -almost surely but it is not defined on exactly one point, which is 1- P -random.

This fact is a simple consequence of the existence of Δ_2^0 1- P -random sequences (for a computable P) and may be also interpreted in terms of learning theory (cf. [33] and the upcoming paper [43]).

2.4 Known effectivizations

Many probabilistic theorems have been effectivized so far. Usually they were stated for computable measures but their generalizations for uncomputable measures follow easily by relativization, i.e., putting a representation s of measure P into the oracle. In this section, we list several known effectivizations of almost sure theorems which we will use further.

As shown by Solovay (unpublished, see [18]), we have this effective version of the Borel-Cantelli lemma:

Proposition 3 (effective Borel-Cantelli lemma). *Let P be a probability measure. If a uniformly s -c.e. sequence of events $U_0, U_1, \dots \in \mathcal{X}^{\mathbb{Z}}$ satisfies $\sum_{i=1}^{\infty} P(U_n) < \infty$ then $\sum_{i=1}^{\infty} \mathbf{1}\{x \in U_n\} < \infty$ on each 1- (s, P) -random point x .*

By the effective Borel-Cantelli lemma, Proposition 3, follows the effective version of the Barron lemma [5, Theorem 3.1]:

Proposition 4 (effective Barron lemma). *For any probability measure P and any s -computable probability measure R , on 1- (s, P) -random points we have*

$$\lim_{n \rightarrow \infty} [-\log R(X_1^n) + \log P(X_1^n) + 2 \log n] = \infty. \quad (2)$$

Random variables $P(x_0 | X_{-n}^{-1})$ for $n \geq 1$ form a martingale process. Thus applying the effective martingale convergence [46], we obtain the effective Lévy law in particular:

Proposition 5 (effective Lévy law). *For a stationary probability measure P , on 1- P -random points there exist limits*

$$P(x_0 | X_{-\infty}^{-1}) := \lim_{n \rightarrow \infty} P(x_0 | X_{-n}^{-1}). \quad (3)$$

Some celebrated result is the effective Birkhoff ergodic theorem [47, 6, 21]. In the following, $\mathbf{E} X := \int X dP$ stands for the expectation of a random variable X with respect to measure P .

Proposition 6 (effective Birkhoff ergodic theorem [6, Theorem 10]). *For a stationary ergodic probability measure P and an s -left-c.e. real random variable G such that $G \geq 0$ and $\mathbf{E} G < \infty$, on 1- (s, P) -random points we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} G \circ T^i = \mathbf{E} G. \quad (4)$$

The proof of the next proposition is an easy application of Proposition 6 and properties of left-c.e. functions.

Proposition 7 (effective Breiman ergodic theorem [44]). *For a stationary ergodic probability measure P and uniformly s -computable real random variables*

$(G_i)_{i \geq 0}$ such that $G_n \geq 0$, $\mathbf{E} \sup_n G_n < \infty$, and limit $\lim_{n \rightarrow \infty} G_n$ exists P -almost surely, on 1 - (s, P) -random points we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} G_i \circ T^i = \mathbf{E} \lim_{n \rightarrow \infty} G_n. \quad (5)$$

Proof. Let $H_k := \sup_{t > k} G_t \geq 0$. Then $G_t \leq H_k$ for all $t > k$ and consequently,

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} G_i \circ T^i \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} H_k \circ T^i. \quad (6)$$

Observe that the supremum H_k of uniformly s -computable functions G_{k+1}, G_{k+2}, \dots is s -left-c.e. Indeed, to enumerate the left cut of the supremum $H_k(x)$ we simultaneously enumerate the left cuts of $G_{k+1}(x), G_{k+2}(x), \dots$. This is possible since every s -computable function is also s -left-c.e. Moreover, we are considering only countably many functions, hence we can guarantee that an element of each left cut appears in the enumeration at least once.

Now, since for all $k \geq 0$ random variables H_k are s -left-c.e. then by Theorem 6 (effective Birkhoff ergodic theorem), on 1 - (s, P) -random points we have

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} G_i \circ T^i \leq \mathbf{E} H_k. \quad (7)$$

Since $H_k \geq 0$ and $\mathbf{E} \sup_k H_k < \infty$ then by the dominated convergence,

$$\inf_{k \geq 0} \mathbf{E} H_k = \lim_{k \rightarrow \infty} \mathbf{E} H_k = \mathbf{E} \lim_{k \rightarrow \infty} H_k = \mathbf{E} \lim_{n \rightarrow \infty} G_n. \quad (8)$$

Thus,

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} G_i \circ T^i \leq \mathbf{E} \lim_{n \rightarrow \infty} G_n. \quad (9)$$

For the converse inequality, consider a natural number M and put random variables $\bar{H}_k := M - \inf_{t > k} \min \{G_t, M\} \in [0, M]$. We observe that \bar{H}_k are also s -left-c.e. since G_t are uniformly s -computable by the hypothesis. By Theorem 6 (effective Birkhoff ergodic theorem), on 1 - (s, P) -random points we have

$$M - \liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \min \{G_i, M\} \circ T^i \leq \mathbf{E} \bar{H}_k. \quad (10)$$

Since $0 \leq \bar{H}_k \leq M$ then by the dominated convergence,

$$\inf_{k \geq 0} \mathbf{E} \bar{H}_k = \lim_{k \rightarrow \infty} \mathbf{E} \bar{H}_k = \mathbf{E} \lim_{k \rightarrow \infty} \bar{H}_k = M - \mathbf{E} \min \left\{ \lim_{n \rightarrow \infty} G_n, M \right\}. \quad (11)$$

Hence, regrouping the terms we obtain

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} G_i \circ T^i &\geq \liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \min \{G_i, M\} \circ T^i \\ &\geq \mathbf{E} \min \left\{ \lim_{n \rightarrow \infty} G_n, M \right\} \xrightarrow{M \rightarrow \infty} \mathbf{E} \lim_{n \rightarrow \infty} G_n, \end{aligned} \quad (12)$$

where the last transition follows by the monotone convergence. By (9) and (12) we derive the claim. \square

The almost sure versions of Propositions 6 and 7 concern random variables which need not be nonnegative [7].

An important result for universal prediction is the Azuma inequality [3], whose following corollary will be used Subsections 3.2 and 3.4.

Theorem 1 (effective Azuma theorem). *For a probability measure P and uniformly s -computable real random variables $(Z_n)_{n \geq 1}$ such that $|Z_n| \leq \epsilon_n \sqrt{n/\ln n}$ with $\lim_{n \rightarrow \infty} \epsilon_n = 0$, on 1 - (s, P) -random points we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n [Z_i - \mathbf{E}(Z_i | X_1^{i-1})] = 0. \quad (13)$$

Proof. Define

$$Y_n := \sum_{i=1}^n [Z_i - \mathbf{E}(Z_i | X_1^{i-1})]. \quad (14)$$

Process $(Y_n)_{n \geq 1}$ is a martingale with respect to process $(X_n)_{n \geq 1}$ with increments bounded by inequality

$$|Z_n - \mathbf{E}(Z_n | X_1^{n-1})| \leq c_n := 2\epsilon_n \sqrt{n/\ln n}. \quad (15)$$

By the Azuma inequality [3] for any $\epsilon > 0$ we obtain

$$P(|Y_n| \geq n\epsilon) \leq 2 \exp\left(-\frac{\epsilon^2 n^2}{2 \sum_{i=1}^n c_i^2}\right) = n^{-\alpha_n}, \quad (16)$$

where

$$\alpha_n := \frac{\epsilon^2 n}{8 \sum_{i=1}^n \epsilon_i^2}. \quad (17)$$

Since $\alpha_n \rightarrow \infty$, we have $\sum_{n=1}^{\infty} P(|Y_n| \geq n\epsilon) < \infty$ and by Proposition 3 (effective Borel-Cantelli lemma), we obtain (13) on 1 - (s, P) -random points. \square

3 Main results

This section contains results concerning effective universal coding and prediction, predictors induced by universal measures and some examples of universal measures and universal predictors.

3.1 Universal coding

Let us begin our considerations with the problem of universal measures, which is related to the problem of universal coding. Suppose that we want to compress losslessly a typical sequence generated by a stationary probability measure P . We can reasonably ask what is the lower limit of such a compression, i.e., what is the minimal ratio of the encoded string length divided by the original string length. In information theory, it is well known that the greatest lower bound of such ratios is given by the entropy rate of measure P . For a stationary probability measure P , we denote its entropy rate as

$$h_P := \lim_{n \rightarrow \infty} \frac{1}{n} \mathbf{E}[-\log P(X_1^n)] = \lim_{k \rightarrow \infty} \mathbf{E}[-\log P(X_{k+1} | X_1^k)]. \quad (18)$$

The entropy rate has the interpretation of the minimal asymptotic rate of lossless encoding of sequences emitted by measure P in various senses: in expectation, almost surely, or on algorithmically random points, where the last interpretation will be pursued in this subsection.

To furnish some theoretical background for universal coding let us recall the Kraft inequality $\sum_{w \in A} 2^{-|w|} \leq 1$, which holds for any prefix-free subset of strings $A \subset \{0, 1\}^*$. The Kraft inequality implies in particular that lossless compression procedures, called uniquely decodable codes, can be mapped one-to-one to semi-measures. In particular, if we are seeking for a universal code, i.e., a uniquely decodable code $w \mapsto C(w) \in \{0, 1\}^*$ which is optimal for some class of stochastic sources P , we can equivalently seek for a universal semi-measure of form $w \mapsto R(w) := 2^{-|C(w)|}$. Consequently, the problem of universal coding would be solved if we point out such a semi-measure R that

$$\lim_{n \rightarrow \infty} \frac{1}{n} [-\log R(X_1^n)] = \lim_{n \rightarrow \infty} \frac{1}{n} |C(X_1^n)| = h_P \quad (19)$$

for some points that are typical of P .

As it is well established in information theory, some initial insight into the problem of universal coding or universal measures is given by the Shannon-McMillan-Breiman (SMB) theorem, which states that function $\frac{1}{n} [-\log P(X_1^n)]$ tends P -almost surely to the entropy rate h_P . The classical proofs of this result were given by Algoet and Cover [2] and Chung [9]. An effective version of the SMB theorem was presented by Hochman [25] and Hoyrup [26].

Theorem 2 (effective SMB theorem [25, 26]). *For a stationary ergodic probability measure P , on 1- P -random points we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} [-\log P(X_1^n)] = h_P. \quad (20)$$

The essential idea of Hoyrup's proof, which is a bit more complicated, can be retold using tools developed in Subsection 2.4. Observe first that we have

$$\frac{1}{n} [-\log P(X_1^n)] = \frac{1}{n} \sum_{i=1}^n [-\log P(X_i | X_1^{i-1})]. \quad (21)$$

Moreover, we have the uniform bound

$$\mathbf{E} \sup_{n \geq 0} [-\log P(X_0 | X_{-n}^{-1})] \leq \mathbf{E} [-\log P(X_0)] + \log e \leq \log eD < \infty, \quad (22)$$

see [41, Lemma 4.26]—invoked by Hoyrup as well. Consequently, the effective SMB theorem follows by Proposition 7 (effective Breiman ergodic theorem) and Proposition 5 (effective Lévy law). In contrast, the reasoning by Hoyrup was more casuistic—he did not mention the Breiman ergodic theorem and invoked some more specific statements.

We note in passing that it could be also interesting to check whether one can effectivize the textbook sandwich proof of the SMB theorem by Algoet and Cover [2] using the decomposition of conditionally algorithmically random sequences by Takahashi [46]. However, this step would require some novel theoretical considerations about conditional algorithmic randomness for uncomputable

measures. We mention this only to point out a possible direction for future research.

As a direct consequence of the effective SMB theorem and Proposition 4 (effective Barron lemma), we obtain this effectivization of another well-known almost sure statement:

Theorem 3 (effective source coding). *For any stationary ergodic measure P and any s -computable probability measure R , on 1 -(s, P)-random points we have*

$$\liminf_{n \rightarrow \infty} \frac{1}{n} [-\log R(X_1^n)] \geq h_P. \quad (23)$$

In the almost sure setting, relationship (23) holds P -almost surely for any stationary ergodic measure P and any (not necessarily computable) probability measure R .

Now we can define universal measures.

Definition 7 (universal measure). *A computable (not necessarily stationary) probability measure R is called (weakly) n -universal if for any stationary ergodic probability measure P , on (weakly) n - P -random points we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} [-\log R(X_1^n)] = h_P. \quad (24)$$

In the almost sure setting, we say that a probability measure R is almost surely universal if (24) holds P -almost surely for any stationary ergodic probability measure P . By Proposition 1, there are only two practically interesting cases of computable universal measures: weakly 2-universal ones and 1-universal ones, since every computable almost surely universal probability measure is automatically weakly 2-universal. We stress that we impose computability of (weakly) n -universal measures by definition since it simplifies statements of some theorems. This should be contrasted with universal prediction à la Solomonoff for left-c.e. semimeasures where the universal element belongs to the class and is not computable [42].

Computable almost surely universal measures exist if the alphabet \mathbb{X} is finite. An important example of an almost surely universal and, as we will see in Section 3.5, also 1-universal measure is the Prediction by Partial Matching (PPM) measure [10, 39, 36]. As we have mentioned, universal measures are closely related to the problem of universal coding (data compression) and more examples of universal measures can be constructed from universal codes, for instance given in [48, 27, 8, 15], using the normalization by Ryabko [37]. This normalization is not completely straightforward, since we need to forge semi-measures into probability measures.

3.2 Universal prediction

Universal prediction is a problem similar to universal coding. In this problem, we also seek for a single procedure that would be optimal within a class of probabilistic sources but we apply a different loss function, namely, we impose the error rate being the density of incorrect guesses of the next output given previous ones. In spite of this difference, we will try to state the problem of universal prediction analogously to universal coding. A predictor is an arbitrary

total function $f : \mathbb{X}^* \rightarrow \mathbb{X}$. The predictor induced by a probability measure P will be defined as

$$f_P(x_1^n) := \arg \max_{x_{n+1} \in \mathbb{X}} P(x_{n+1} | x_1^n), \quad (25)$$

where $\arg \max_{x \in \mathbb{X}} g(x) := \min \{a \in \mathbb{X} : g(a) \geq g(x) \text{ for all } x \in \mathbb{X}\}$ for the total order $a_1 < \dots < a_D$ on $\mathbb{X} = \{a_1, \dots, a_D\}$. Moreover, for a stationary measure P , we define the unpredictability rate

$$u_P := \lim_{n \rightarrow \infty} \mathbf{E} \left[1 - \max_{x_0 \in \mathbb{X}} P(x_0 | X_{-n}^{-1}) \right]. \quad (26)$$

It is natural to ask whether the unpredictability rate can be related to entropy rate. Using the Fano inequality [19], a classical result of information theory, and its converse [17], both independently brought to algorithmic randomness theory by Fortnow and Lutz [20], yields this bound:

Theorem 4. *For a stationary measure P over a D -element alphabet,*

$$\frac{D}{D-1} \eta \left(\frac{1}{D} \right) u_P \leq h_P \leq \eta(u_P) + u_P \log(D-1), \quad (27)$$

where $\eta(p) := -p \log p - (1-p) \log(1-p)$.

Moreover, Fortnow and Lutz [20] found out some stronger inequalities, sandwich-bounding the unpredictability of an arbitrary sequence in terms of its effective dimension. The effective dimension turns out to be a generalization of the entropy rate to arbitrary sequences [26], which are not necessarily random with respect to stationary ergodic measures.

In the less general framework of stationary ergodic measures, using the Azuma theorem, we can show that no predictor can beat the induced predictor and the error rate committed by the latter equals the unpredictability rate u_P . The following proposition concerning the error rates effectivizes the well-known almost sure proposition (the proof in the almost sure setting is available in [1]).

Theorem 5 (effective source prediction). *For any stationary ergodic measure P and any s -computable predictor f , on 1 -(s, P)-random points we have*

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \mathbf{1}\{X_{i+1} \neq f(X_1^i)\} \geq u_P. \quad (28)$$

Moreover, if the induced predictor f_P is s -computable then (28) holds with the equality for $f = f_P$.

Proof. Let measure P be stationary ergodic. In view of Theorem 1 (effective Azuma theorem), for any s -computable predictor f , on 1 -(s, P)-random points we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} [\mathbf{1}\{X_{i+1} \neq f(X_1^i)\} - P(X_{i+1} \neq f(X_1^i) | X_1^i)] = 0. \quad (29)$$

Moreover, we have

$$P(X_{i+1} \neq f(X_1^i)|X_1^i) \geq 1 - \max_{x_{i+1} \in \mathbb{X}} P(x_{i+1}|X_1^i) \quad (30)$$

Subsequently, we observe that limits $\lim_{n \rightarrow \infty} P(x_0|X_{-n}^{-1})$ exist on 1- (s, P) -random points by Proposition 5 (effective Lévy law). Thus by Proposition 7 (effective Breiman ergodic theorem) and the dominated convergence, on 1- (s, P) -random points we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \left[1 - \max_{x_{i+1} \in \mathbb{X}} P(x_{i+1}|X_1^i) \right] &= \mathbf{E} \lim_{n \rightarrow \infty} \left[1 - \max_{x_0 \in \mathbb{X}} P(x_0|X_{-n}^{-1}) \right] \\ &= u_P. \end{aligned} \quad (31)$$

Hence inequality (28) follows by (29), (30) and (31). Similarly, the equality in (28) for $f = f_P$ follows by noticing that inequality (30) turns out to be the equality in this case. \square

In the almost sure setting, relationship (28) holds P -almost surely for any stationary ergodic measure P and any (not necessarily computable) predictor f .

We can see that there can be some problem in the effectivization of relationship (28) caused by the induced predictor f_P possibly not being s -computable for certain representations s of measure P —since sometimes testing the equality of two real numbers cannot be done in a finite time. However, probabilities $P(X_{i+1} \neq f_P(X_1^i)|X_1^i)$ are always s -computable. Thus, we can try to define universal predictors in the following way.

Definition 8 (universal predictor). *A computable predictor f is called (weakly) n -universal if for any stationary ergodic probability measure P , on (weakly) n - P -random points we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \mathbf{1}\{X_{i+1} \neq f(X_1^i)\} = u_P. \quad (32)$$

In the almost sure setting, we say that a predictor f is almost surely universal if (32) holds P -almost surely for any stationary ergodic probability measure P . Almost surely universal predictors exist if the alphabet \mathbb{X} is finite [4, 32, 1, 24, 23]. In [44] it was proved that the almost sure predictor by [24] is also 1-universal.

3.3 Predictors induced by backward estimators

The almost surely universal predictors by [4, 32, 1, 24, 23] were constructed without a reference to universal measures. Nevertheless, these constructions are all based on estimation of conditional probabilities. For a stationary ergodic process one can consider two separate problems: backward and forward estimation. The first problem is naturally connected to prediction. We want to estimate the conditional probability of $(n + 1)$ -th bit given the first n bits. Is it possible that, as we increase n , our estimates converge to the true value at

some point? To be precise, we ask whether there exists a probability measure R such that for every stationary ergodic measure P we have P -almost surely

$$\lim_{n \rightarrow \infty} \sum_{x_{n+1} \in \mathbb{X}} |R(x_{n+1}|X_1^{n-1}) - P(x_{n+1}|X_1^{n-1})| = 0. \quad (33)$$

It was shown by Bailey [4] that this is not possible. As we are about to see, we can get something a bit weaker, namely, the convergence in Cesaro averages. But to get there, it will be helpful to consider a bit different problem.

Suppose again that we want to estimate a conditional probability but the bit that we are interested in is fixed and we are looking more and more into the past. In this scenario, we want to estimate the conditional probability $P(x_0|X_{-\infty}^{-1})$ and we ask whether increasing the knowledge of the past can help us achieve the perfect guess. Precisely, we ask if there exists a probability measure R such that for every stationary ergodic measure P we have P -almost surely

$$\lim_{n \rightarrow \infty} \sum_{x_0 \in \mathbb{X}} |R(x_0|X_{-n}^{-1}) - P(x_0|X_{-\infty}^{-1})| = 0. \quad (34)$$

It was famously shown by Ornstein that such estimators exist. (Ornstein proved this for binary-valued processes but the technique can be generalized to finite-valued processes.)

Theorem 6 (Ornstein theorem [32]). *Let the alphabet be finite. There exists a computable measure R such that for every stationary ergodic measure P we have P -almost surely that*

$$\lim_{n \rightarrow \infty} \sum_{x_0 \in \mathbb{X}} |R(x_0|X_{-n}^{-1}) - P(x_0|X_{-\infty}^{-1})| = 0. \quad (35)$$

Definition 9. *We call a measure R an almost surely universal backward estimator when it satisfies condition (35) P -almost surely for every stationary ergodic measure P , whereas it is called a (weakly) n -universal backward estimator if R is computable and convergence (35) holds on all respective (weakly) n - P -random points.*

One can come up with a naive idea: What if we take a universal backward estimator and use it in a forward fashion? Surprisingly, this simple trick gives us almost everything we can get, i.e., a forward estimator that converges to the conditional probability on average. Bailey [4] showed that for an almost surely universal backward estimator R and for every stationary ergodic measure P we have P -almost surely

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \sum_{x_{i+1} \in \mathbb{X}} |R(x_{i+1}|X_1^i) - P(x_{i+1}|X_1^i)| = 0. \quad (36)$$

The proof of this fact is a direct application of the Breiman ergodic theorem. Since we have a stronger effective version of the Breiman theorem (Theorem 7), we can strengthen Bailey's result to an effective version as well. It turns out that even if we take a backward estimator that is good only almost surely (possibly failing on some random points), then the respective result for the forward estimation will hold in the strong sense—on every 1- P -random point.

Theorem 7 (effective Bailey theorem). *Let R be a computable almost surely universal backward estimator. For every stationary ergodic measure P on 1- P -random points we have (36).*

Proof. Let R be a computable almost surely universal backward estimator. Fix an $x \in \mathbb{X}$. By Proposition 5 (effective Lévy law), for every stationary ergodic probability measure P we have P -almost surely

$$\lim_{n \rightarrow \infty} |R(x|X_{-n}^{-1}) - P(x|X_{-n}^{-1})| = 0. \quad (37)$$

Note that the bound $0 \leq |R(x|X_{-n}^{-1}) - P(x|X_{-n}^{-1})| \leq 1$ holds uniformly. Moreover, variables $R(x|X_{-n}^{-1}) - P(x|X_{-n}^{-1})$ are uniformly s -computable for any representation s of P . Hence, we can apply Theorem 7 (effective Breiman ergodic theorem) to obtain

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} |R(x|X_1^i) - P(x|X_1^i)| = \mathbf{E} 0 = 0. \quad (38)$$

for 1- P -random points. The claim follows from this immediately. \square

Definition 10. *We call a measure R an almost surely universal forward estimator when it satisfies condition (36) P -almost surely for every stationary ergodic measure P , whereas it is called a (weakly) n -universal forward estimator if R is computable and convergence (36) holds on all respective (weakly) n - P -random points.*

One can expect that the predictor f_R induced by a universal forward estimator R in the sense of Definition 10 is also universal in the sense of Definition 8. This is indeed true. To show this fact, we will first prove a certain inequality for induced predictors, which generalizes the result from [14, Theorem 2.2] for binary classifiers. This particular observation seems to be new.

Proposition 8 (prediction inequality). *Let p and q be two probability distributions over a countable alphabet \mathbb{X} . For $x_p = \arg \max_{x \in \mathbb{X}} p(x)$ and $x_q = \arg \max_{x \in \mathbb{X}} q(x)$, we have inequality*

$$0 \leq p(x_p) - p(x_q) \leq \sum_{x \in \mathbb{X}} |p(x) - q(x)|. \quad (39)$$

Proof. Without loss of generality, assume $x_p \neq x_q$. By the definition of x_p and x_q , we have $p(x_p) - p(x_q) \geq 0$ and $q(x_q) - q(x_p) \geq 0$. Hence we obtain

$$\begin{aligned} 0 \leq p(x_p) - p(x_q) &\leq p(x_p) - p(x_q) - q(x_p) + q(x_q) \\ &\leq |p(x_p) - q(x_p)| + |p(x_q) - q(x_q)| \leq \sum_x |p(x) - q(x)|. \end{aligned} \quad (40)$$

\square

Now we can show a general result about universal predictors induced by forward estimators of conditional probabilities.

Theorem 8 (effective induced prediction I). *For a 1-universal forward estimator R , the induced predictor f_R is 1-universal if f_R is computable.*

Proof. Let R be 1-universal forward estimator. By the definition, for every stationary ergodic measure P and all 1- P -random points

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \sum_{x_{i+1} \in \mathbb{X}} |P(x_{i+1}|X_1^i) - R(x_{i+1}|X_1^i)| = 0. \quad (41)$$

Consequently, combining this with Proposition 8 (prediction inequality) yields on 1- P -random points

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} [P(X_{i+1} \neq f_R(X_1^i)|X_1^i) - P(X_{i+1} \neq f_P(X_1^i)|X_1^i)] = 0. \quad (42)$$

Now, we notice that by (29), we have on 1- P -random points

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} [\mathbf{1}\{X_{i+1} \neq f_R(X_1^i)\} - P(X_{i+1} \neq f_R(X_1^i)|X_1^i)] = 0, \quad (43)$$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} [\mathbf{1}\{X_{i+1} \neq f_P(X_1^i)\} - P(X_{i+1} \neq f_P(X_1^i)|X_1^i)] = 0. \quad (44)$$

Combining the three above observations completes the proof. \square

Interestingly, it suffices for a measure to be a computable almost surely universal backward estimator to yield a 1-universal forward estimator and, consequently, a 1-universal predictor. In contrast, we can easily see that a computable almost surely universal forward estimator does not necessarily induce a 1-universal predictor.

Theorem 9. *There exists a computable almost surely universal forward estimator R such that the induced predictor f_R is not 1-universal.*

Proof. Let us take $\mathbb{X} = \{0, 1\}$ and restrict ourselves to one-sided space $\mathbb{X}^{\mathbb{N}}$ without loss of generality. Fix a computable almost surely universal forward estimator Q . Let P_0 be the computable measure of a Bernoulli(θ) process, i.e., $P_0(x_1^n) = \prod_{i=1}^n \theta^{x_i} (1-\theta)^{1-x_i}$, where $\theta > 1/2$ is rational. Observe that by Proposition 2 there exists a point $y \in \mathbb{X}^{\mathbb{N}}$ which is 1- P_0 -random and a computable function $g : \mathbb{X}^* \rightarrow \{0, 1\}$ such that $P_0(A) = 0$ and $A = \{y\}$ for event

$$A := (\# \{i \in \mathbb{N} : g(X_1^i) = 1\} = \infty). \quad (45)$$

In other words, there is a computable method to single out some 1- P_0 -random point y out of the set of sequences $\mathbb{X}^{\mathbb{N}}$. In particular, we can use function g to spoil measure Q on that point y while preserving the property of an almost surely universal forward estimator. We will denote the spoiled version of measure Q by R . Conditional distributions $R(X_{m+1}|X_1^m)$ will differ from $Q(X_{m+1}|X_1^m)$ for infinitely many m on point y and for finitely many m elsewhere.

Let $K(x_1^n) := \# \{i \leq n : g(x_1^i) = 1\}$. The construction of measure R proceeds by induction on the string length together with an auxiliary counter U . We let $R(x_1) := Q(x_1)$ and $U(x_1) := 0$. Suppose that $R(x_1^n)$ and $U(x_1^n)$ are defined but $R(x_1^{n+1})$ is not. If $U(x_1^n) \geq K(x_1^n)$ then we put $R(x_{n+1}|x_1^n) :=$

$Q(x_{n+1}|x_1^n)$ and $U(x_1^{n+1}) := U(x_1^n)$. Else, if $U(x_1^n) < K(x_1^n)$ then we put $R(x_{n+1}^{n+N}|x_1^n) := \prod_{i=n+1}^{n+N} \theta^{1-x_i} (1-\theta)^{x_i}$ (reverted compared to the definition of $P_0!$) and $U(x_{n+N}) := K(x_1^n)$ where N is the smallest number such that

$$\frac{1}{n+N} \left[\sum_{i=0}^{n-1} P_0(X_{i+1} \neq f_R(x_1^i) | X_1^i = x_1^i) + N\theta \right] \geq \frac{1}{2}. \quad (46)$$

Such number N exists since $P_0(X_{i+1} \neq f_R(x_1^i) | X_1^i = x_1^i) > 1-\theta$. This completes the construction of R .

The sets of 1- P -random sequences are disjoint for distinct stationary ergodic P by Theorem 6 (effective Birkhoff ergodic theorem). Hence $K(X_1^n)$ is bounded P -almost surely for any stationary ergodic P . Consequently, since $U(X_1^n)$ is non-decreasing then P -almost surely there exists a random number $M < \infty$ such that for all $m > M$ we have $R(X_{m+1}|X_1^m) = Q(X_{m+1}|X_1^m)$. Hence R inherits the property of an almost surely universal forward estimator from Q .

Now let us inspect what happens on y . Since $K(X_1^n)$ is unbounded on y then by the construction of R , we obtain on y that $U(X_1^n) < K(X_1^n)$ holds infinitely often and

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^n P_0(X_{i+1} \neq f_R(X_1^i) | X_1^i) \geq \frac{1}{2} > u_{P_0} = 1 - \theta. \quad (47)$$

Hence predictor f_R is not 1-universal. \square

3.4 Predictors induced by universal measures

Following the work of Ryabko [37], see also [38], we can ask a natural question whether predictors induced by some universal measures in the sense of Definition 7, such as the PPM measure [10, 39, 36] to be discussed in Section 3.5, are also universal. Ryabko was close to demonstrate the analogous implication in the almost sure setting but did not provide the complete proof. He has shown this proposition:

Theorem 10 ([36]). *Let R be an almost surely universal measure and P be a stationary ergodic measure. We have P -almost surely*

$$\lim_{n \rightarrow \infty} \mathbf{E} \frac{1}{n} \sum_{i=0}^{n-1} |P(X_{i+1}|X_0^i) - R(X_{i+1}|X_0^i)| = 0. \quad (48)$$

Condition (48) is quite close to condition (36), i.e., universal forward estimator, which—as we have shown in Theorem 8—implies universality of the induced predictor. Close does not mean exactly the same. If we would like to derive universality of the induced predictor directly from (48), there are two problems on the way (where $Y_n \geq 0$ stands for the expression under the expectation): Firstly, $\lim_{n \rightarrow \infty} \mathbf{E} Y_n = 0$ does not necessarily imply $\mathbf{E} \lim_{n \rightarrow \infty} Y_n = 0$ since the limit may not exist almost surely and, secondly, if $\mathbf{E} \lim_{n \rightarrow \infty} Y_n = 0$ then $\lim_{n \rightarrow \infty} Y_n = 0$ holds almost surely but this equality may fail on some 1-random points.

In spite of these difficulties, Ryabko [37] showed in the almost sure setting that there exist universal forward estimators which are generated by some universal measures. In view of Theorem 8, this solves the question of existence of

universal measures that induce universal predictors. But it does not solve yet the problem of effectivization to algorithmically random points and of systematicity of the construction.

In this section, we will show that each 1-universal measure, under a relatively mild condition (1), satisfied by the PPM measure, is a 1-universal forward estimator and hence, in the light of the previous section, it induces a 1-universal predictor. We do not know yet whether this condition is necessary. We will circumvent Theorem 10 by applying Proposition 7 (effective Breiman ergodic theorem) and Theorem 1 (effective Azuma theorem). The first stage of our preparations includes two statements which can be called the effective conditional SMB theorem and the effective conditional universality.

Proposition 9 (effective conditional SMB theorem). *Let the alphabet be finite and let P be a stationary ergodic probability measure. On 1- P -random points we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \left[- \sum_{x_{i+1} \in \mathbb{X}} P(x_{i+1} | X_1^i) \log P(x_{i+1} | X_1^i) \right] = h_P. \quad (49)$$

Proof. Let us write the conditional entropy

$$W_i := \left[- \sum_{x_{i+1} \in \mathbb{X}} P(x_{i+1} | X_1^i) \log P(x_{i+1} | X_1^i) \right]. \quad (50)$$

We have $0 \leq W_i \leq \log D$ with D being the cardinality of the alphabet. Moreover by Proposition 5 (effective Lévy law), on 1- P -random points there exists limit

$$\lim_{n \rightarrow \infty} W_n \circ T^{-n-1} = \left[- \sum_{x_0 \in \mathbb{X}} P(x_0 | X_{-\infty}^{-1}) \log P(x_0 | X_{-\infty}^{-1}) \right]. \quad (51)$$

Hence by Proposition 7 (effective Breiman ergodic theorem), on 1- (s, P) -random points

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} W_i = \mathbf{E} \left[- \sum_{x_0 \in \mathbb{X}} P(x_0 | X_{-\infty}^{-1}) \log P(x_0 | X_{-\infty}^{-1}) \right] = h_P \quad (52)$$

since $\mathbf{E} [-\log P(X_0 | X_{-\infty}^{-1})] = \lim_{n \rightarrow \infty} [-\log P(X_1^n)] / n = h_P$. \square

Proposition 10 (effective conditional universality). *Let the alphabet be finite and let P be a stationary ergodic probability measure.*

$$-\log R(x_{n+1} | x_1^n) \leq \epsilon_n \sqrt{n / \ln n}, \quad \lim_{n \rightarrow \infty} \epsilon_n = 0 \quad (53)$$

then on 1- P -random points we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \left[- \sum_{x_{i+1} \in \mathbb{X}} P(x_{i+1} | X_1^i) \log R(x_{i+1} | X_1^i) \right] = h_P. \quad (54)$$

Proof. Let us write the conditional pointwise entropy $Z_i := -\log R(X_{i+1}|X_1^i)$. Now suppose that measure R is 1-universal and satisfies (53). Then by Theorem 1 (effective Azuma theorem), on 1- P -random points we obtain

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \mathbf{E}(Z_i | X_1^i) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} Z_i = \lim_{n \rightarrow \infty} \frac{1}{n} [-\log R(X_1^n)] = h_P, \quad (55)$$

which is the claim of Proposition 10. \square

In the second stage of our preparations, we recall the famous Pinsker inequality used by Ryabko [37] to prove Theorem 10.

Proposition 11 (Pinsker inequality [11]). *Let p and q be probability distributions over a countable alphabet \mathbb{X} . We have*

$$\left[\sum_{x \in \mathbb{X}} |p(x) - q(x)| \right]^2 \leq (2 \ln 2) \sum_{x \in \mathbb{X}} p(x) \log \frac{p(x)}{q(x)}. \quad (56)$$

Now we can show the main result of this section, namely, that every universal measure which satisfies a mild condition induces a universal predictor.

Theorem 11 (effective induced prediction II). *If measure R is 1-universal and satisfies (53) then it is a 1-universal forward estimator.*

Proof. Let R be a 1-universal measure, whereas P be the stationary ergodic measure. By Propositions 9 (effective conditional SMB theorem) and 10 (effective conditional universality), on 1- P -random points we obtain

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \left[\sum_{x_{i+1}} P(x_{i+1} | X_1^i) \log \frac{P(x_{i+1} | X_1^i)}{R(x_{i+1} | X_1^i)} \right] = 0. \quad (57)$$

Hence by Proposition 11 (Pinsker inequality), we derive on 1- P -random points

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \left[\sum_{x_{i+1}} |P(x_{i+1} | X_1^i) - R(x_{i+1} | X_1^i)| \right]^2 = 0. \quad (58)$$

Subsequently, the Cauchy-Schwarz inequality $\mathbf{E} Y^2 \geq (\mathbf{E} Y)^2$ yields on 1- P -random points

$$\begin{aligned} 0 &\geq \lim_{n \rightarrow \infty} \left[\frac{1}{n} \sum_{i=0}^{n-1} \sum_{x_{i+1}} |P(x_{i+1} | X_1^i) - R(x_{i+1} | X_1^i)| \right]^2 \\ &= \left[\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \sum_{x_{i+1}} |P(x_{i+1} | X_1^i) - R(x_{i+1} | X_1^i)| \right]^2 \geq 0. \end{aligned} \quad (59)$$

Consequently, R is a 1-universal forward estimator. \square

Combining Theorem 11 and Theorem 8, we obtain that predictor f_R is 1-universal provided measure R is 1-universal and satisfies condition (53)—if predictor f_R is computable itself. Condition (53) does not seem to have been discussed in the literature of universal prediction.

3.5 PPM measure

In this section, we will discuss the Prediction by Partial Matching (PPM) measure. The PPM measure comes in several flavors and was discovered gradually. Cleary and Witten [10] coined the name PPM, which we prefer since it is more distinctive, and considered the adaptive Markov approximations PPM_k defined roughly in equation (60). Later, Ryabko [39, 36] considered the infinite series PPM defined in equation (62), called it the measure R , and proved that it is a universal measure. Precisely, Ryabko used the Krichevsky-Trofimov smoothing (+1/2) rather than the Laplace smoothing (+1) applied in (60). This difference does not affect universality. As we will show now, series PPM provides an example of a 1-universal measure that satisfies condition (53) and thus yields a natural 1-universal predictor.

Upon the first reading, the definition of the PPM measure may appear cumbersome but it is roughly a Bayesian mixture of all Markov chains of all orders. Its universality can be then motivated by the fact that Markov chains with rational transition probabilities are both countable and dense in the class of stationary ergodic measures [40]. Our specific definition of measure PPM is as follows.

Definition 11 (PPM measure). *Let the alphabet be $\mathbb{X} = \{a_1, \dots, a_D\}$, where $D \geq 2$. Adapting definitions by [10, 39, 36, 16], the PPM measure of order $k \geq 0$ is defined as*

$$\text{PPM}_k(x_1^n) := \begin{cases} D^{-k-1} \prod_{i=k+2}^n \frac{N(x_{i-k}^i | x_1^{i-1}) + 1}{N(x_{i-k}^{i-1} | x_1^{i-2}) + D}, & k \leq n-2, \\ D^{-n}, & k \geq n-1, \end{cases} \quad (60)$$

where the frequency of a substring w_1^k in a string x_1^n is

$$N(w_1^k | x_1^n) := \sum_{i=1}^{n-k+1} \mathbf{1}\{x_i^{i+k-1} = w_1^k\}. \quad (61)$$

Subsequently, we define the total PPM measure

$$\text{PPM}(x_1^n) := \sum_{k=0}^{\infty} \left[\frac{1}{k+1} - \frac{1}{k+2} \right] \text{PPM}_k(x_1^n). \quad (62)$$

Infinite series (62) is computable since $\text{PPM}_k(x_1^n) = D^{-n}$ for $k \geq n-1$. The almost sure universality of the total PPM measure follows by the Stirling approximation and the Birkhoff ergodic theorem, see [39, 36, 16]. Since the Birkhoff ergodic theorem can be effectivized for 1-random points in form of Proposition 6, we obtain in turn this effectivization.

Theorem 12 (effective PPM universality, cf. [36]). *Measure PPM is 1-universal.*

Proof. Computability of the PPM measure follows since series (62) can be truncated with the constant term $\text{PPM}_k(x_1^n) = D^{-n}$ for $k \geq n-1$ and thus values $\text{PPM}(x_1^n)$ are rational. To show 1-universality of the PPM measure, we observe the following. Using the Stirling approximation, the PPM measure can be

related to the empirical entropy of the respective string. The empirical (conditional) entropy of string x_1^n of order $k \geq 0$ is defined as

$$h_k(x_1^n) := \sum_{w_1^{k+1} \in \mathbb{X}^{k+1}} \frac{N(w_1^{k+1}|x_1^n)}{n-k} \log \frac{N(w_1^k|x_1^{n-1})}{N(w_1^{k+1}|x_1^n)}. \quad (63)$$

In particular, by Theorem A4 in [16], we have the bound

$$0 \leq -\log \text{PPM}_k(x_1^n) - k \log D - (n-k)h_k(x_1^n) \leq D^{k+1} \log[e^2 n]. \quad (64)$$

Subsequently, by Proposition 6 (effective Birkhoff ergodic theorem), on 1- P -random points we have

$$\lim_{n \rightarrow \infty} \frac{N(w_1^{k+1}|X_1^n)}{n-k} = P(w_1^{k+1}). \quad (65)$$

Then by (64),

$$\lim_{n \rightarrow \infty} \frac{1}{n} [-\log \text{PPM}_k(X_1^n)] = h_{k,P} := \mathbf{E} [-\log P(X_{k+1}|X_1^k)]. \quad (66)$$

Since

$$-\log \text{PPM}(x_1^n) \leq 2 \log(k+2) - \log \text{PPM}_k(x_1^n) \quad (67)$$

then

$$\limsup_{n \rightarrow \infty} \frac{1}{n} [-\log \text{PPM}(X_1^n)] \leq \inf_{k \geq 0} h_{k,P} = h_P \quad (68)$$

on 1- P -random points, whereas the reverse inequality for the lower limit follows by Proposition 4 (effective Barron lemma). \square

Finally, we can show that predictor f_{PPM} induced by the PPM measure is 1-universal. First, we notice explicitly these bounds:

Theorem 13 (PPM bounds). *We have*

$$-\log \text{PPM}(x_1^n) \leq 2 \log(n+1) + n \log D, \quad (69)$$

$$-\log \text{PPM}(x_{n+1}|x_1^n) \leq 3 \log(n+D). \quad (70)$$

Proof. Observe that $\text{PPM}_k(x_1^n) = D^{-n}$ for $k \geq n-1$. Hence by (67), we obtain claim (69). The derivation of claim (70) is slightly longer. First, by the definition of PPM_k , we have

$$-\log \text{PPM}_k(x_{n+1}|x_1^n) \leq \log [N(x_{n-k+1}^n|x_1^{n-1}) + D] \leq \log(n+D). \quad (71)$$

Now let $G(x_1^n)$ be the minimal k such that $\text{PPM}_k(x_1^n)$ is maximal. We have $G(x_1^n) \leq n-1$, since $\text{PPM}_k(x_1^n) = D^{-n}$ for $k \geq n-1$. Moreover, we have a bound reverse to (67), namely

$$-\log \text{PPM}(x_1^n) \geq -\log \text{PPM}_{G(x_1^n)}(x_1^n). \quad (72)$$

Combining the above with (67) yields

$$\begin{aligned} -\log \text{PPM}(x_{n+1}|x_1^n) &= -\log \text{PPM}(x_1^{n+1}) + \log \text{PPM}(x_1^n) \\ &\leq 2 \log(G(x_1^n) + 2) - \log \text{PPM}_{G(x_1^n)}(x_1^{n+1}) + \log \text{PPM}_{G(x_1^n)}(x_1^n) \\ &= 2 \log(G(x_1^n) + 2) + \log \text{PPM}_{G(x_1^n)}(x_{n+1}|x_1^n) \leq 3 \log(n+D). \end{aligned} \quad (73)$$

\square

Now comes the main theorem.

Theorem 14. *Predictor f_{PPM} is 1-universal.*

Proof. Computability of predictor f_{PPM} follows since values $\text{PPM}(x_1^n)$ are rational so the least symbol of those having the maximal conditional probability can be computed in a finite time. Consequently, the claim follows by Theorems 11, 12, and 13. \square

We think that 1-universality of predictor f_{PPM} is quite expected and intuitive. But as we can see, the PPM measure satisfies condition (53) with a large reserve. It is an open question whether there are 1-universal measures such that conditional probabilities $R(x_{n+1}|x_1^n)$ converge to zero much faster than for the PPM measure but they still induce 1-universal predictors. It would be interesting to find such measures. Maybe they have some other desirable properties, also from a practical point of view.

Funding

This work was supported by the National Science Centre Poland grant 2018/31/B/HS1/04018.

Acknowledgments

The authors are grateful to the anonymous reviewers of unaccepted earlier conference versions of the paper, who provided a very stimulating and encouraging feedback. Additional improvements to the paper were inspired by participants of the Kolmogorov seminar in Moscow at which this work was presented by the first author. Finally, the authors express their gratitude to Dariusz Kalociński for his comments and proofreading. Both authors declare an equal contribution to the paper.

References

- [1] Paul H. Algoet. The strong law of large numbers for sequential decisions under uncertainty. *IEEE Trans. Inform. Theory*, 40(3):609–633, 1994.
- [2] Paul H. Algoet and Thomas M. Cover. A sandwich proof of the Shannon-McMillan-Breiman theorem. *Ann. Probab.*, 16:899–909, 1988.
- [3] Kazuoki Azuma. Weighted sums of certain dependent random variables. *Tohoku Math. J. Sec. Ser.*, 19(3):357–367, 1967.
- [4] David Harold Bailey. *Sequential schemes for classifying and predicting ergodic processes*. PhD thesis, Stanford University, 1976.
- [5] Andrew R. Barron. *Logically Smooth Density Estimation*. PhD thesis, Stanford University, 1985.
- [6] Laurent Bienvenu, Adam R. Day, Mathieu Hoyrup, Ilya Mezhirov, and Alexander Shen. A constructive version of Birkhoff’s ergodic theorem for Martin-Löf random points. *Inform. and Comput.*, 210:21–30, 2012.

- [7] Leo Breiman. The individual ergodic theorem of information theory. *Ann. Math. Statist.*, 28:809–811, 1957.
- [8] Moses Charikar, Eric Lehman, April Lehman, Ding Liu, Rina Panigrahy, Manoj Prabhakaran, Amit Sahai, and Abhi Shelat. The smallest grammar problem. *IEEE Trans. Inform. Theory*, 51:2554–2576, 2005.
- [9] K. L. Chung. A note on the ergodic theorem of information theory. *Ann. Math. Statist.*, 32:612–614, 1961.
- [10] John Cleary and Ian Witten. Data compression using adaptive coding and partial string matching. *IEEE Trans. Comm.*, 32:396–402, 1984.
- [11] Imre Csiszár and János Körner. *Information Theory: Coding Theorems for Discrete Memoryless Systems*. Cambridge University Press, 2011.
- [12] A. Philip Dawid and Vladimir G. Vovk. Prequential probability: principles and properties. *Bernoulli*, 5(1):125–162, 1999.
- [13] Adam D. Day and Joseph S. Miller. Randomness for non-computable measures. *Trans. Amer. Math. Soc.*, 365(7):3575–3591, 2013.
- [14] Luc Devroye, László Györfi, and Gábor Lugosi. *A Probabilistic Theory of Pattern Recognition*. Springer, 1996.
- [15] Łukasz Dębowski. On the vocabulary of grammar-based codes and the logical consistency of texts. *IEEE Trans. Inform. Theory*, 57:4589–4599, 2011.
- [16] Łukasz Dębowski. Is natural language a perigraphic process? The theorem about facts and words revisited. *Entropy*, 20(2):85, 2018.
- [17] Łukasz Dębowski. *Information Theory Meets Power Laws: Stochastic Processes and Language Models*. Wiley and Sons, 2021.
- [18] Rodney G. Downey and Denis R. Hirschfeldt. *Algorithmic Randomness and Complexity*. Springer, 2010.
- [19] Robert M. Fano. *Transmission of Information*. The MIT Press, 1961.
- [20] Lance Fortnow and Jack H. Lutz. Prediction and dimension. *J. Comp. System Sci.*, 70(4):570–589, 2005.
- [21] Johanna N. Y. Franklin, Noam Greenberg, Joseph S. Miller, and Keng Meng Ng. Martin-Löf random points satisfy Birkhoff’s ergodic theorem for effectively closed sets. *Proc. Amer. Math. Soc.*, 140:3623–3628, 2012.
- [22] Peter Gács. Uniform test of algorithmic randomness over a general space. *Theor. Comp. Sci.*, 341:91–137, 2005.
- [23] László Györfi and Gábor Lugosi. Strategies for sequential prediction of stationary time series. In M. Dror, P. L’Ecuyer, and F. Szidarovszky, editors, *Modeling Uncertainty: An examination of its theory, methods, and applications*. Kluwer Academic Publishers, 2001.

- [24] László Györfi, Gábor Lugosi, and Gusztáv Morvai. A simple randomized algorithm for consistent sequential prediction of ergodic time series. *IEEE Trans. Inform. Theory*, 45(7):2642–2650, 1999.
- [25] Michael Hochman. Upcrossing inequalities for stationary sequences and applications. *Ann. Probab.*, 37(6):2135–2149, 2009.
- [26] Mathieu Hoyrup. The dimension of ergodic random sequences. In *29th International Symposium on Theoretical Aspects of Computer Science*, page 567, 2012.
- [27] John C. Kieffer and Enhui Yang. Grammar-based codes: A new class of universal lossless source codes. *IEEE Trans. Inform. Theory*, 46:737–754, 2000.
- [28] Leonid A. Levin. The concept of a random sequence. *Dokl. Akad. Nauk SSSR*, 21:548–550, 1973.
- [29] Leonid A. Levin. Uniform tests for randomness. *Dokl. Akad. Nauk SSSR*, 227(1):33–35, 1976.
- [30] Leonid A. Levin. Randomness conservation inequalities: information and independence in mathematical theories. *Inform. and Control*, 61(1):15–37, 1984.
- [31] Per Martin-Löf. The definition of random sequences. *Inform. and Control*, 9:602–619, 1966.
- [32] Donald S. Ornstein. Guessing the next output of a stationary process. *Israel J. Math.*, 30(3):292–296, 1978.
- [33] Daniel Osherson and Scott Weinstein. Recognizing strong random reals. *Rev. Symbol. Logic*, 1(1):56–63, 2008.
- [34] Jan Reimann. Effectively closed sets of measures and randomness. *Ann. Pure Appl. Logic*, 156(1):170–182, 2008.
- [35] Jan Reimann and Theodore A. Slaman. Measures and their random reals. *Trans. Amer. Math. Soc.*, 367:5081–5097, 2015.
- [36] Boris Ryabko. Applications of Kolmogorov complexity and universal codes to nonparametric estimation of characteristics of time series. *Fund. Informat.*, 83(1-2):177–196, 2008.
- [37] Boris Ryabko. Compression-based methods for nonparametric prediction and estimation of some characteristics of time series. *IEEE Trans. Inform. Theory*, 55(9):4309–4315, 2009.
- [38] Boris Ryabko, Jaakko Astola, and Mikhail Malyutov. *Compression-Based Methods of Statistical Analysis and Prediction of Time Series*. Springer, 2016.
- [39] Boris Y. Ryabko. Prediction of random sequences and universal coding. *Probl. Inform. Transm.*, 24(2):87–96, 1988.

- [40] Daniil Ryabko. On finding predictors for arbitrary families of processes. *J. Machine Learn. Res.*, 11:581–602, 2010.
- [41] Meir Smorodinsky. *Ergodic Theory, Entropy*, volume 214 of *Lecture Notes in Mathematics*. Springer, 1971.
- [42] R. J. Solomonoff. A formal theory of inductive inference, part 1 and part 2. *Inform. and Control*, 7:1–22, 224–254, 1964.
- [43] Tomasz Steifer. A note on learning-theoretic characterizations of randomness and convergence. submitted.
- [44] Tomasz Steifer. *Computable Prediction of Infinite Binary Sequences with Zero-One Loss*. PhD thesis, Institute of Computer Science, Polish Academy of Sciences, 2020.
- [45] Joe Suzuki. Universal prediction and universal coding. *Syst. Comput. Japan*, 34(6):1–11, 2003.
- [46] Hayato Takahashi. On a definition of random sequences with respect to conditional probability. *Inform. and Comput.*, 206(12):1375–1382, 2008.
- [47] Vladimir V. V’yugin. Ergodic theorems for individual random sequences. *Theor. Comput. Sci.*, 207(2):343–361, 1998.
- [48] Jacob Ziv and Abraham Lempel. A universal algorithm for sequential data compression. *IEEE Trans. Inform. Theory*, 23:337–343, 1977.