

TWO METHODS OF ESTIMATION OF THE DRIFT PARAMETERS OF THE COX–INGERSOLL–ROSS PROCESS: CONTINUOUS OBSERVATIONS

Olena Dehtiar, Yuliya Mishura and Kostiantyn Ralchenko

Department of Probability, Statistics and Actuarial Mathematics

Mechanics and Mathematics Faculty

Taras Shevchenko National University of Kyiv

64 Volodymyrska, 01601 Kiev, Ukraine

Key Words: Cox–Ingersoll–Ross process, continuous observations, parameter estimation, strong consistency.

ABSTRACT

We consider a stochastic differential equation of the form $dr_t = (a - br_t)dt + \sigma\sqrt{r_t}dW_t$, where a , b and σ are positive constants. The solution corresponds to the Cox–Ingersoll–Ross process. We study the estimation of an unknown drift parameter (a, b) by continuous observations of a sample path $\{r_t, t \in [0, T]\}$. First, we prove the strong consistency of the maximum likelihood estimator. Since this estimator is well-defined only in the case $2a > \sigma^2$, we propose another estimator that is defined and strongly consistent for all positive a , b , σ . The quality of the estimators is illustrated by simulation results.

1. INTRODUCTION

The Cox–Ingersoll–Ross (CIR) process is a very famous object and is a unique solution of the following stochastic differential equation

$$dr_t = (a - br_t)dt + \sigma\sqrt{r_t}dW_t, \quad r_t|_{t=0} = r_0 > 0 \quad (1)$$

where $W = \{W_t, t \geq 0\}$ is a Wiener process and a, b, σ are positive constants.

There are many papers devoted to the construction and the asymptotic properties of the parameter estimators of the CIR process, based on conditional least squares and the

maximum likelihood estimator see, e. g., Overbeck and Rydén (1997); Li and Ma (2015); De Rossi (2010); Barczy et al. (2019); Ben Alaya and Kebaier (2012, 2013). More precisely, Overbeck and Rydén (1997) used conditional least squares as a basic method and derived two estimators, which differ by the method of estimating of parameter σ : pseudo likelihood method and unweighted least squares method based on squared residuals. Moreover, strong consistency of these estimators was proved for both observations at equidistant time point and continuous observations. Nevertheless, the simulation study demonstrated that the maximum likelihood estimator outperforms the conditional least-squares estimators and the pseudo likelihood approach in most cases.

Ben Alaya and Kebaier (2012, 2013) presented a new approach to the investigation of the asymptotic behavior of the maximum likelihood estimator, depending on the values of the parameters, both for continuous and discrete observations. Using an exact simulation algorithm, the authors illustrated practical behavior of these estimators' errors covering ergodic and nonergodic situations. Asymptotic properties of maximum likelihood estimator for the stable CIR process were studied in Li and Ma (2015). In Barczy et al. (2019) the authors investigated the asymptotic behavior of the maximum likelihood estimator for so-called jump-type CIR process, driven by a standard Wiener process and a subordinator. They distinguish three cases: subcritical, critical and supercritical.

The paper De Rossi (2010) contains the method that is based on the sequential Monte Carlo techniques and shows how to construct a simulated maximum likelihood procedure. This paper also describes two methods of computing the likelihood: sampling and re-sampling algorithm which solve the problem of degeneracy for realistic sample sizes. In order to maximize the likelihood, the author applies the genetic algorithm that relies on the survival of the fittest in determining the optimal parameter vector. Testing on simulated data confirmed that this approach allows not to undermine the accuracy of the estimation procedure by the effect of simulation errors and copes with larger parameter dimensions at a modest computational cost. Avoiding the computational burden the MATLAB implementation of the estimation routine is provided and tested in Kladívko (2007). Moreover, the

simulation algorithm for the approximation of the CIR process trajectories was described in Milstein and Schoenmakers (2015).

Concerning the discretization, due to the square-root diffusion coefficient the classical Euler–Maruyama scheme does not preserve the non-negativity of the process. Cozma and Reisinger (2016) provided a brief discussion of the discretization schemes often encountered in the finance literature. They also investigated the exponential integrability properties which play an essential role in deriving strong convergence of Euler discretization schemes. The high priority to the convergence results and its applications were given in the paper Deelstra and Delbaen (1995). The Euler approach to constructing discretization schemes was also introduced in Dereich et al. (2012). Here the using of the drift-implicit square-root Euler method gives a strictly positive approximation of the original CIR process and some global convergence results. The paper Mishura and Munchak (2016) is the example of constructing additive and multiplicative discrete approximation schemes taking the Euler approximations of the CIR process itself but replacing the increments of the Wiener process with i. i. d. bounded vanishing symmetric random variables.

In this paper we investigate two estimators of the parameter (a, b) by continuous observations of a sample path of CIR process $r = \{r_t, t \in [0, T]\}$ and prove their strong consistency as $T \rightarrow \infty$. The first one is the standard maximum likelihood estimator, which was constructed and studied in Ben Alaya and Kebaier (2012, 2013). Compared to the known results, we establish the strong consistency instead of weak one. Note that the maximum likelihood estimator is well-defined only if $2a \geq \sigma^2$, because, in particular, it contains the integral $\int_0^T \frac{1}{r_t} dt$, which exists with probability one if and only if $2a \geq \sigma^2$, see (Ben Alaya and Kebaier, 2012, Prop. 4). For this reason, we decided to create some statistics that converges regardless of whether $2a \geq \sigma^2$ or not. On this way we created a different estimator of the vector parameter (a, b) , which is strongly consistent for all positive a, σ and b . Another advantage of the new alternative estimator is that it has simpler form, therefore, it is computationally faster. It includes only two statistics of the process r , namely the Lebesgue integrals $\int_0^T r_t dt$ and $\int_0^T r_t^2 dt$, see Theorem 8 below. At the same time the maximum likelihood estimator depends

on two Lebesgue integrals, $\int_0^T r_t dt$ and $\int_0^T \frac{dt}{r_t}$, on the stochastic integral $\int_0^T \frac{dr_t}{r_t}$ and on the process itself. Particular attention in the paper is paid to the a. s. asymptotic behavior of the integral $\int_0^T r_t^2 dt$, which is crucial for the construction of the alternative estimator. We would like to emphasize that in the present paper we restrict ourselves to the case $b > 0$. The boundary case $b = 0$ was investigated in Ben Alaya and Kebaier (2013), where the consistency and asymptotic distribution on the maximum likelihood estimator was derived, assuming that $2a \geq \sigma^2$. To the best of our knowledge, parameter estimation for the case $b < 0$ remains an open problem.

The paper is organized as follows. In Section 2 we recall some basic properties of CIR process. Section 3 is devoted to the strong consistency of the maximum likelihood estimator. In Section 4 we introduce an alternative estimator and prove its strong consistency.

2. PRELIMINARIES

In this section we consider the properties of the CIR process. Most of them will be useful for statistical parameter estimation, but some facts are of independent interest. These facts are well-known, but we combined them into one statement for the reader's convenience.

Proposition 1. *Assume that $2a > \sigma^2$. Then*

- 1) *The unique solution $r = \{r_t, t \geq 0\}$ of equation (1) is positive with probability 1:*

$$\inf \{t \geq 0 : r_t = 0\} = +\infty \quad a. s. \quad (2)$$

(with convention $\inf \emptyset = +\infty$). Moreover,

$$\mathbf{P} \left\{ \limsup_{t \rightarrow \infty} r_t = +\infty \right\} = \mathbf{P} \left\{ \liminf_{t \rightarrow \infty} r_t = 0 \right\} = 1. \quad (3)$$

- 2) *The process r is ergodic and it has continuous stationary density that corresponds to gamma distribution and has the following form:*

$$p_\infty(x) = \frac{\beta^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\beta x} \mathbf{1}_{x>0}, \quad \alpha = \frac{2a}{\sigma^2}, \quad \beta = \frac{2b}{\sigma^2}. \quad (4)$$

- 3) *For any function $f: \mathbb{R}^+ \rightarrow \mathbb{R}$ such that $\int_{\mathbb{R}^+} |f(x)| p_\infty(x) dx < \infty$ we have that*

$$\frac{1}{T} \int_0^T f(r_t) dt \rightarrow \int_{\mathbb{R}} f(x) p_\infty(x) dx, \quad a. s., \quad \text{as } T \rightarrow \infty. \quad (5)$$

The results of Proposition 1 follow from the general theory of homogeneous diffusions. In particular, (2) follows from the Feller's test for explosions (see, e. g., Karatzas and Shreve, 1991, Thm. 5.29, p. 348), and (3) can be deduced from Karatzas and Shreve (1991, Prop. 5.22, p. 345) (see also Mijatović and Urusov, 2012, Sec. 2.3). Statements 2)–3) are based on the ergodic theory for homogeneous diffusions, see, e. g., Kutoyants (2004, Thm. 1.16) or Skorokhod (1989, Ch. 1, § 3). The conditions of these general results for the case of the CIR process can be easily verified, see, e. g., Bel Hadj Khlifa et al. (2016, Example 4, p. 280). A direct proof of (2) for the CIR process is given in Alfonsi (2015, Sec. 1.2.4). The derivation of the stationary distribution (4) can be found, e. g., in Alfonsi (2015, Eq. (1.24)).

Corollary 2. *It follows immediately from (5) (see also Remark after Proposition 4 in Ben Alaya and Kebaier (2012)) that in the case $2a > \sigma^2$ we have the following asymptotic relations:*

$$\frac{1}{T} \int_0^T r_t dt \rightarrow \int_{\mathbb{R}} xp_{\infty}(x) dx = \frac{\alpha}{\beta} = \frac{a}{b}, \quad a. s., \text{ as } T \rightarrow \infty, \quad (6)$$

$$\frac{1}{T} \int_0^T \frac{1}{r_t} dt \rightarrow \int_{\mathbb{R}} \frac{1}{x} p_{\infty}(x) dx = \frac{\beta}{\alpha - 1} = \frac{b}{a - \sigma^2/2}, \quad a. s., \text{ as } T \rightarrow \infty, \quad (7)$$

$$\frac{1}{T} \int_0^T r_t^2 dt \rightarrow \int_{\mathbb{R}} x^2 p_{\infty}(x) dx = \left(\frac{\alpha}{\beta}\right)^2 + \frac{\alpha}{\beta^2} = \frac{a^2}{b^2} + \frac{a\sigma^2}{2b^2}, \quad a. s., \text{ as } T \rightarrow \infty. \quad (8)$$

Remark 1. It was proved in (Deelstra and Delbaen, 1995, Thm. 1) that the convergence (6) holds also in the case $0 < 2a \leq \sigma^2$. The convergence (8) is valid for all positive a and σ , this will be justified in Theorem 7 below.

3. MAXIMUM LIKELIHOOD ESTIMATION

Let us recall the construction of the maximum likelihood estimator of the couple of unknown parameters (a, b) by the continuous observations of r over the interval $[0, T]$. We assume that $2a > \sigma^2$ throughout this section.

Dividing (1) by $\sqrt{r_t}$ and integrating (1) over the time interval $[0, s]$ we get the equality:

$$\int_0^s \frac{dr_t}{\sqrt{r_t}} = \int_0^s \left(\frac{a}{\sqrt{r_t}} - b\sqrt{r_t} \right) dt + \sigma W_s.$$

In order to construct likelihood function for the estimation of couple (a, b) of parameters, we use the Girsanov theorem for the Wiener process with the drift that equals

$$\frac{1}{\sigma} \int_0^s \left(\frac{a}{\sqrt{r_t}} - b\sqrt{r_t} \right) dt.$$

Then the likelihood function that corresponds to the likelihood $dQ_{(0,0)}/dQ_{(a,b)}$, where $Q_{(0,0)}$ is a probability measure responsible for zero values of parameters, and $Q_{(a,b)}$ is responsible for the couple (a, b) , gets the following form:

$$\begin{aligned} \mathcal{L} &= \exp \left\{ - \int_0^T \frac{a - br_t}{\sigma\sqrt{r_t}} dW_t - \frac{1}{2} \int_0^T \frac{(a - br_t)^2}{\sigma^2 r_t} dt \right\} \\ &= \exp \left\{ - \int_0^T \frac{a - br_t}{\sigma^2 r_t} dr_t + \frac{1}{2} \int_0^T \frac{(a - br_t)^2}{\sigma^2 r_t} dt \right\}. \end{aligned}$$

Since we are interested in maximization of the function that corresponds to $dQ_{(a,b)}/dQ_{(0,0)}$, our likelihood function has the form

$$\tilde{\mathcal{L}} = \exp \left\{ \int_0^T \frac{a - br_t}{\sigma^2 r_t} dr_t - \frac{1}{2} \int_0^T \frac{(a - br_t)^2}{\sigma^2 r_t} dt \right\}.$$

The maximum likelihood estimator for the couple (a, b) is constructed by maximizing of $\tilde{\mathcal{L}}$ with respect to (a, b) . It has the following form:

$$\hat{a}_T = \frac{\int_0^T r_t dt \int_0^T \frac{dr_t}{r_t} - T \cdot (r_T - r_0)}{\int_0^T r_t dt \cdot \int_0^T \frac{dt}{r_t} - T^2}; \quad (9)$$

$$\hat{b}_T = \frac{(r_0 - r_T) \int_0^T \frac{dt}{r_t} + T \int_0^T \frac{dr_t}{r_t}}{\int_0^T r_t dt \cdot \int_0^T \frac{dt}{r_t} - T^2}. \quad (10)$$

Theorem 3. *Assume that $2a > \sigma^2$. Then the estimator (\hat{a}_T, \hat{b}_T) is strongly consistent.*

Proof. Taking into consideration that $\int_0^T \frac{dr_t}{r_t} = a \int_0^T \frac{dt}{r_t} - bT + \sigma \int_0^T \frac{dW_t}{\sqrt{r_t}}$ we can represent (9) in the following form:

$$\begin{aligned} \hat{a}_T &= \frac{a \int_0^T \frac{dt}{r_t} \int_0^T r_t dt - bT \int_0^T r_t dt + \sigma \int_0^T \frac{dW_t}{\sqrt{r_t}} \int_0^T r_t dt - T(r_T - r_0)}{\int_0^T r_t dt \cdot \int_0^T \frac{dt}{r_t} - T^2} \\ &= \frac{a \int_0^T \frac{dt}{r_t} \int_0^T r_t dt - bT \int_0^T r_t dt + \sigma \int_0^T \frac{dW_t}{\sqrt{r_t}} \int_0^T r_t dt - T \int_0^T (a - br_t) dt - \sigma T \int_0^T \sqrt{r_t} dW_t}{\int_0^T r_t dt \cdot \int_0^T \frac{dt}{r_t} - T^2} \end{aligned}$$

$$= a + \frac{\sigma \int_0^T \frac{dW_t}{\sqrt{r_t}} \cdot \int_0^T r_t dt - \sigma T \int_0^T \sqrt{r_t} dW_t}{\int_0^T r_t dt \cdot \int_0^T \frac{dt}{r_t} - T^2}, \quad (11)$$

and similarly, estimator from (10) can be presented as

$$\begin{aligned} \hat{b}_T &= \frac{(r_0 - r_T) \int_0^T \frac{dt}{r_t} + aT \int_0^T \frac{dt}{r_t} - bT^2 + \sigma T \int_0^T \frac{dW_t}{\sqrt{r_t}}}{\int_0^T r_t dt \cdot \int_0^T \frac{dt}{r_t} - T^2} \\ &= \frac{-\int_0^T (a - br_t) dt \cdot \int_0^T \frac{dt}{r_t} - \sigma \int_0^T \sqrt{r_t} dW_t \cdot \int_0^T \frac{dt}{r_t} + aT \int_0^T \frac{dt}{r_t} - bT^2 + \sigma T \int_0^T \frac{dW_t}{\sqrt{r_t}}}{\int_0^T r_t dt \cdot \int_0^T \frac{dt}{r_t} - T^2} \\ &= b + \frac{-\sigma \int_0^T \sqrt{r_t} dW_t \cdot \int_0^T \frac{dt}{r_t} + \sigma T \int_0^T \frac{dW_t}{\sqrt{r_t}}}{\int_0^T r_t dt \cdot \int_0^T \frac{dt}{r_t} - T^2}. \end{aligned} \quad (12)$$

Consider the remainder

$$R_T^a = \frac{\sigma \int_0^T \frac{dW_t}{\sqrt{r_t}} \cdot \int_0^T r_t dt - \sigma T \int_0^T \sqrt{r_t} dW_t}{\int_0^T r_t dt \cdot \int_0^T \frac{dt}{r_t} - T^2}$$

from (11), and the remainder

$$R_T^b = \frac{-\sigma \int_0^T \sqrt{r_t} dW_t \cdot \int_0^T \frac{dt}{r_t} + \sigma T \int_0^T \frac{dW_t}{\sqrt{r_t}}}{\int_0^T r_t dt \cdot \int_0^T \frac{dt}{r_t} - T^2}$$

from (12) can be considered in the same lines. So,

$$R_T^a = \frac{\sigma \left(\frac{1}{T} \int_0^T \frac{dW_t}{\sqrt{r_t}} \right) \cdot \left(\frac{1}{T} \int_0^T r_t dt \right) - \sigma \left(\frac{1}{T} \int_0^T \sqrt{r_t} dW_t \right)}{\frac{1}{T^2} \int_0^T r_t dt \cdot \int_0^T \frac{dt}{r_t} - 1} \quad (13)$$

According to Corollary 2, relations (5) and (6), the denominator in (13) tends to $\frac{a}{a - \frac{\sigma^2}{2}} - 1 > 0$.

Moreover, both values in the numerator, $\frac{1}{T} \int_0^T \frac{dW_t}{\sqrt{r_t}}$ and $\frac{1}{T} \int_0^T \sqrt{r_t} dW_t$ tends to zero a.s. as $T \rightarrow \infty$. Indeed,

$$\frac{1}{T} \int_0^T \frac{dW_t}{\sqrt{r_t}} = \frac{\int_0^T \frac{dt}{r_t}}{T} \cdot \frac{\int_0^T \frac{dW_t}{\sqrt{r_t}}}{\int_0^T \frac{dt}{r_t}},$$

and

$$\frac{\int_0^T \frac{dt}{r_t}}{T} \rightarrow \frac{b}{a - \frac{\sigma^2}{2}}.$$

This means that $\int_0^T \frac{dt}{r_t} \rightarrow \infty, T \rightarrow \infty$, but $\int_0^T \frac{dt}{r_t}$ is a square characteristics of the locally square integrable martingale $\int_0^T \frac{dW_t}{\sqrt{r_t}}$. According to the strong law of large numbers for locally

square integrable martingales (Liptser and Shiryaev (2012)), $\left(\int_0^T \frac{dt}{r_t}\right)^{-1} \cdot \int_0^T \frac{dW_t}{\sqrt{r_t}} \rightarrow 0$ a.s. as $T \rightarrow \infty$. Therefore, $\frac{1}{T} \int_0^T \frac{dW_t}{\sqrt{r_t}} \rightarrow 0$ a.s. as $T \rightarrow \infty$, and similar relation holds for $\frac{1}{T} \int_0^T \sqrt{r_t} dW_t$. Together with (13), this means that \hat{a}_T is strongly consistent. \square

Remark 2. The weak consistency and asymptotic normality of the maximum likelihood estimator (\hat{a}, \hat{b}) was obtained in Ben Alaya and Kebaier (2012, 2013). Mention that the weak consistency holds also in the boundary cases, when $2a = \sigma^2$ and/or $b = 0$ (however, joint asymptotic distribution is not normal in these cases). If $2a < \sigma^2$, then the maximum likelihood estimator is not well-defined. If $2a \geq \sigma^2$ and $b < 0$, then it is not consistent.

Remark 3. An alternative parametrization of the CIR process is

$$dr_t = \alpha(\mu - r_t)dt + \sigma\sqrt{r_t}dW_t, \quad r_t|_{t=0} = r_0 > 0, \quad (14)$$

where $W = \{W_t, t \geq 0\}$ is a Wiener process and α, μ, σ are positive constants. Assume that $2\alpha\mu > \sigma^2$. Then the maximum likelihood estimator for the couple (α, μ) constructed by the continuous observations of r over the interval $[0, T]$ has the following form

$$\hat{\alpha}_T = \frac{T \int_0^T \frac{dr_t}{r_t} - \int_0^T \frac{dt}{r_t} \int_0^T dr_t}{\int_0^T r_t dt \int_0^T \frac{dt}{r_t} - T^2}, \quad \hat{\mu}_T = \frac{\int_0^T r_t dt \int_0^T \frac{dr_t}{r_t} - T(r_T - r_0)}{T \int_0^T \frac{dr_t}{r_t} - \int_0^T \frac{dt}{r_t} (r_T - r_0)}.$$

The strong consistency of $(\hat{\alpha}_T, \hat{\mu}_T)$ follows from Theorem 3, if we take into account the relations

$$\alpha = b, \quad \mu = \frac{a}{b}, \quad \hat{\alpha}_T = \hat{b}_T, \quad \hat{\mu}_T = \frac{\hat{a}_T}{\hat{b}_T}. \quad (15)$$

4. AN ALTERNATIVE APPROACH TO DRIFT PARAMETERS ESTIMATION

The disadvantage of the maximum likelihood estimators is that they work only if $a > \sigma^2/2$, however, a priori we do not know if this relation holds for the observed process. To avoid this circumstance, in this section we will introduce a new estimator for the parameter (a, b) based on the statistics $\int_0^T r_t dt$ and $\int_0^T r_t^2 dt$. First, we will prove that the convergence (8) remains valid in the case $0 < a \leq \sigma^2/2$. To this end, we start with several auxiliary results. The first result gives the asymptotics of two normalized Lebesgue integrals

Lemma 4. *Let $a > 0$, $b > 0$, $\sigma > 0$. Then the following normalized integrals asymptotically vanish as $T \rightarrow \infty$:*

$$\frac{1}{T} \int_0^T e^{-bt} r_t dt \rightarrow 0 \quad \text{a. s.}, \quad (16)$$

$$\frac{1}{T} e^{-2bT} \int_0^T e^{2bt} r_t dt \rightarrow 0 \quad \text{a. s.} \quad (17)$$

Proof. 1. In order to prove the asymptotic relation (16), we rewrite the normalized integral as follows:

$$\frac{1}{T} \int_0^T e^{-bt} r_t dt = \frac{1}{T} \int_0^{\sqrt{T}} e^{-bt} r_t dt + \frac{1}{T} \int_{\sqrt{T}}^T e^{-bt} r_t dt.$$

Then the first integral can be bounded in the following way:

$$\frac{1}{T} \int_0^{\sqrt{T}} e^{-bt} r_t dt \leq \frac{1}{T} \int_0^{\sqrt{T}} r_t dt = \frac{1}{\sqrt{T}} \cdot \frac{1}{\sqrt{T}} \int_0^{\sqrt{T}} r_t dt \rightarrow 0 \quad \text{a. s.}, \text{ as } T \rightarrow \infty,$$

since $\frac{1}{\sqrt{T}} \int_0^{\sqrt{T}} r_t dt \rightarrow \frac{a}{b}$ a. s., as $T \rightarrow \infty$, by (6), see Remark 1. Furthermore, the second integral can be bounded as

$$\frac{1}{T} \int_{\sqrt{T}}^T e^{-bt} r_t dt \leq e^{-b\sqrt{T}} \frac{1}{T} \int_{\sqrt{T}}^T r_t dt \leq e^{-b\sqrt{T}} \frac{1}{T} \int_0^T r_t dt \rightarrow 0 \quad \text{a. s.}, \text{ as } T \rightarrow \infty,$$

where the convergence follows from (6). Thus, relation (16) is proved.

2. Note that $\int_0^T e^{2bt} r_t dt \geq \int_0^T r_t dt \rightarrow \infty$ a. s., as $T \rightarrow \infty$, by (6). Therefore, applying the L'Hôpital's rule, we conclude that

$$\lim_{T \rightarrow \infty} \frac{\int_0^T e^{2bt} r_t dt}{T e^{2bT}} = \lim_{T \rightarrow \infty} \frac{e^{2bT} r_T}{e^{2bT} + 2be^{2bT} T} = \lim_{T \rightarrow \infty} \frac{r_T}{1 + 2bT} \quad \text{a. s.}$$

Now (17) follows from the a. s. convergence $\frac{r_T}{T} \rightarrow 0$, $T \rightarrow \infty$. The latter convergence was established in the proof of Theorem 1 of Deelstra and Delbaen (1995). \square

The next result presents the bounds for the moments of the related stochastic integral.

Lemma 5. *Denote*

$$Z_t = \int_0^t e^{bu} \sqrt{r_u} dW_u, \quad t \geq 0. \quad (18)$$

There exists a constant $C > 0$ such that for all $t \geq 0$,

$$\mathbf{E} [Z_t^2] \leq C e^{2bt}, \quad \mathbf{E} [Z_t^3] \leq C e^{3bt}, \quad \mathbf{E} [Z_t^2 r_t] \leq C e^{2bt}. \quad (19)$$

Proof. 1. According to (Deelstra and Delbaen, 1995, Eq. (1)), the process r satisfies the following relations

$$\begin{aligned} r_t &= e^{-bt} \left(r_0 + a \int_0^t e^{bu} du + \sigma \int_0^t e^{bu} \sqrt{r_u} dW_u \right) \\ &= e^{-bt} \left(r_0 - \frac{a}{b} + \frac{a}{b} e^{bt} + \sigma \int_0^t e^{bu} \sqrt{r_u} dW_u \right). \end{aligned} \quad (20)$$

Therefore, in particular, its expectation equals

$$\mathbf{E}r_t = \left(r_0 - \frac{a}{b} \right) e^{-bt} + \frac{a}{b}.$$

Then for the 2nd moment we have the following representation:

$$\begin{aligned} \mathbf{E}Z_t^2 &= \mathbf{E} \left[\left(\int_0^t e^{bu} \sqrt{r_u} dW_u \right)^2 \right] = \mathbf{E} \left[\int_0^t e^{2bu} r_u du \right] = \int_0^t \left(\left(r_0 - \frac{a}{b} \right) e^{bu} + \frac{a}{b} e^{2bu} \right) du \\ &= \left(r_0 - \frac{a}{b} \right) \frac{e^{bt} - 1}{b} + \frac{a}{b} \cdot \frac{e^{2bt} - 1}{2b} = e^{2bt} \left(\frac{a}{2b^2} + \frac{1}{b} \left(r_0 - \frac{a}{b} \right) e^{-bt} - \frac{1}{b} \left(r_0 - \frac{a}{2b} \right) e^{-2bt} \right). \end{aligned}$$

Consequently, $\mathbf{E}Z_t^2 \leq C e^{2bt}$ with $C = \frac{a}{2b^2} + \frac{1}{b} \left| r_0 - \frac{a}{b} \right| + \frac{1}{b} \left| r_0 - \frac{a}{2b} \right|$.

2. Let us consider $\mathbf{E}Z_t^3$. By Itô's formula, from (18) we have

$$Z_t^3 = 3 \int_0^t Z_s e^{2bs} r_s ds + 3 \int_0^t Z_s^2 e^{bs} \sqrt{r_s} dW_s. \quad (21)$$

Note that according to Karatzas and Shreve (1991, Problem 3.15, p. 306) (see also Mishura and Shevchenko 2017, Thm. 9.3), for any $p \geq 1$, $\mathbf{E} \left[\sup_{t \in [0, T]} |r_t|^{2p} \right] < \infty$. It follows from (20) and (18) that

$$r_s = e^{-bs} \left(r_0 - \frac{a}{b} + \frac{a}{b} e^{bs} + \sigma Z_s \right) \quad (22)$$

Therefore, we have also that $\mathbf{E} \left[\sup_{t \in [0, T]} |Z_t|^{2p} \right] < \infty$ for any $p \geq 1$. Consequently, all the terms in the equality (21) have bounded expectations.

Hence,

$$\mathbf{E} [Z_t^3] = 3\mathbf{E} \int_0^t Z_s e^{2bs} r_s ds. \quad (23)$$

We insert (22) into (23) and obtain

$$\mathbf{E} [Z_t^3] = 3\mathbf{E} \int_0^t Z_s e^{bs} \left(r_0 - \frac{a}{b} + \frac{a}{b} e^{bs} + \sigma Z_s \right) ds = 3\sigma \int_0^t e^{bs} \mathbf{E} [Z_s^2] ds,$$

since $\mathbf{E}Z_s = 0$. Applying the bound $\mathbf{E}[Z_s^2] \leq Ce^{2bs}$ from (19), we get

$$\mathbf{E}[Z_t^3] \leq 3\sigma C \int_0^t e^{3bs} ds = \frac{C\sigma}{b} (e^{3bt} - 1) \leq \frac{C\sigma}{b} e^{3bt}.$$

3. We express r_t through Z_t by (22), and get

$$\mathbf{E}[Z_t^2 r_t] = \left(\left(r_0 - \frac{a}{b} \right) e^{-bt} + \frac{a}{b} \right) \mathbf{E}Z_t^2 + \sigma e^{-bt} \mathbf{E}Z_t^3 \leq Ce^{2bt},$$

where the inequality follows from the first two bounds in (19). \square

Lemma 6. *Let the process Z be defined by (18). Then the following normalized stochastic integrals vanish as $T \rightarrow \infty$:*

$$\frac{1}{T} \int_0^T e^{-bt} Z_t \sqrt{r_t} dW_t \rightarrow 0 \quad \text{a. s.}, \quad (24)$$

$$\frac{1}{T} e^{-2bT} \int_0^T e^{bt} Z_t \sqrt{r_t} dW_t \rightarrow 0 \quad \text{a. s.} \quad (25)$$

Proof. 1. Obviously, the convergence (24) is equivalent to

$$\frac{1}{T+1} \int_0^T e^{-bt} Z_t \sqrt{r_t} dW_t \rightarrow 0 \quad \text{a. s.}, \text{ as } T \rightarrow \infty.$$

By Kronecker's lemma (see, e. g., Deelstra and Delbaen, 1995), in order to prove this convergence it suffices to show that

$$\int_0^\infty \frac{e^{-bt} Z_t \sqrt{r_t}}{t+1} dW_t < \infty \quad \text{a. s.} \quad (26)$$

Since the process $M_T = \int_0^T \frac{e^{-bt} Z_t \sqrt{r_t}}{t+1} dW_t$ is a martingale, it suffices to prove that

$$\mathbf{E}\langle M \rangle_T = \mathbf{E} \left[\int_0^T \frac{e^{-2bt} Z_t^2 r_t}{(t+1)^2} dt \right] < \infty. \quad (27)$$

But (27) follows immediately from the third bound of (19):

$$\mathbf{E} \left[\int_0^T \frac{e^{-2bt} Z_t^2 r_t}{(t+1)^2} dt \right] \leq C \int_0^T \frac{1}{(t+1)^2} dt = C \left(1 - \frac{1}{T+1} \right) \leq C.$$

Thus, (24) is proved.

2. Similarly, by Kronecker's lemma, the convergence (25) follows from the existence a. s. of the integral

$$\int_0^\infty \frac{e^{-bt} Z_t \sqrt{r_t}}{(t+1)e^{2bt}} dW_t = \int_0^\infty \frac{e^{-bt} Z_t \sqrt{r_t}}{t+1} dW_t.$$

This integral is finite, because it coincides with the integral (26) considered in the first part of the proof. \square

Now we are ready to prove the announced asymptotic result for $\frac{1}{T} \int_0^T r_t^2 dt$.

Theorem 7. *The following convergence holds:*

$$\frac{1}{T} \int_0^T r_t^2 dt \rightarrow \frac{a^2}{b^2} + \frac{a\sigma^2}{2b^2} \quad \text{a. s., as } T \rightarrow \infty.$$

Proof. Using relation (20), we get the following equalities:

$$\begin{aligned} \frac{1}{T} \int_0^T r_t^2 dt &= \frac{1}{T} \int_0^T e^{-2bt} \left(r_0 - \frac{a}{b} + \frac{a}{b} e^{bt} + \sigma \int_0^t e^{bu} \sqrt{r_u} dW_u \right)^2 dt \\ &= \frac{1}{T} \int_0^T e^{-2bt} \left(r_0 - \frac{a}{b} + \frac{a}{b} e^{bt} \right)^2 dt \\ &\quad + \frac{2\sigma}{T} \int_0^T e^{-2bt} \left(r_0 - \frac{a}{b} + \frac{a}{b} e^{bt} \right) \left(\int_0^t e^{bu} \sqrt{r_u} dW_u \right) dt \\ &\quad + \frac{\sigma^2}{T} \int_0^T e^{-2bt} \left(\int_0^t e^{bu} \sqrt{r_u} dW_u \right)^2 dt \\ &=: I_1 + I_2 + I_3. \end{aligned} \tag{28}$$

The term I_1 is the subject of straightforward calculations:

$$\begin{aligned} I_1 &= \frac{1}{T} \int_0^T e^{-2bt} \left(r_0 - \frac{a}{b} + \frac{a}{b} e^{bt} \right)^2 dt \\ &= \frac{1}{T} \int_0^T \left(\left(r_0 - \frac{a}{b} \right)^2 e^{-2bt} + \frac{2a}{b} \left(r_0 - \frac{a}{b} \right) e^{-bt} + \frac{a^2}{b^2} \right) dt. \\ &= \left(r_0 - \frac{a}{b} \right)^2 \frac{1 - e^{-2bT}}{2bT} + \frac{2a}{b} \left(r_0 - \frac{a}{b} \right) \frac{1 - e^{-bT}}{bT} + \frac{a^2}{b^2}. \end{aligned}$$

Evidently, the following convergence holds:

$$I_1 \rightarrow \frac{a^2}{b^2}, \quad T \rightarrow \infty. \tag{29}$$

Let us consider I_2 . The equality (20) implies that

$$\sigma \int_0^t e^{bu} \sqrt{r_u} dW_u = e^{bt} r_t - r_0 + \frac{a}{b} - \frac{a}{b} e^{bt}.$$

Consequently, integral I_2 can be transformed as follows:

$$\begin{aligned} I_2 &= \frac{2\sigma}{T} \int_0^T e^{-2bt} \left(r_0 - \frac{a}{b} + \frac{a}{b} e^{bt} \right) \left(\int_0^t e^{bu} \sqrt{r_u} dW_u \right) dt \\ &= \frac{2}{T} \int_0^T e^{-2bt} \left(r_0 - \frac{a}{b} + \frac{a}{b} e^{bt} \right) \left(e^{bt} r_t - r_0 + \frac{a}{b} - \frac{a}{b} e^{bt} \right) dt \\ &= \frac{2}{T} \left(r_0 - \frac{a}{b} \right) \int_0^T e^{-bt} r_t dt + \frac{2a}{bT} \int_0^T r_t dt - \frac{2}{T} \int_0^T e^{-2bt} \left(r_0 - \frac{a}{b} + \frac{a}{b} e^{bt} \right)^2 dt. \end{aligned} \quad (30)$$

Asymptotic relation (16) implies that the first term in the right-hand side of (30) converges to zero a. s. as $T \rightarrow \infty$. By (6), for the second term we have

$$\frac{2a}{bT} \int_0^T r_t dt \rightarrow \frac{2a^2}{b^2} \quad \text{a. s., as } T \rightarrow \infty.$$

Finally, the third term equals $-2I_1$, therefore, by (29),

$$-\frac{2}{T} \int_0^T e^{-2bt} \left(r_0 - \frac{a}{b} + \frac{a}{b} e^{bt} \right)^2 dt = -2I_1 \rightarrow -\frac{2a^2}{b^2}, \quad \text{as } T \rightarrow \infty.$$

Thus,

$$I_2 \rightarrow 0 \quad \text{a. s., as } T \rightarrow \infty. \quad (31)$$

It remains to study the asymptotic behavior of I_3 . Note that

$$I_3 = \frac{\sigma^2}{T} \int_0^T e^{-2bt} Z_t^2 dt,$$

where Z_t is defined in (18). By Itô's formula, from (18) we get

$$Z_t^2 = \int_0^t e^{2bs} r_s ds + 2 \int_0^t Z_s e^{bs} \sqrt{r_s} dW_s.$$

Therefore we can present I_3 as the sum

$$I_3 = \frac{\sigma^2}{T} \int_0^T e^{-2bt} \int_0^t e^{2bs} r_s ds dt + \frac{2\sigma^2}{T} \int_0^T e^{-2bt} \int_0^t Z_s e^{bs} \sqrt{r_s} dW_s dt =: I_{3,1} + I_{3,2}. \quad (32)$$

Using the Fubini theorem, we transform $I_{3,1}$ as follows:

$$\begin{aligned} I_{3,1} &= \frac{\sigma^2}{T} \int_0^T e^{2bs} r_s \int_s^T e^{-2bt} dt ds = \frac{\sigma^2}{T} \int_0^T e^{2bs} r_s \frac{e^{-2bs} - e^{-2bT}}{2b} ds \\ &= \frac{\sigma^2}{2bT} \int_0^T r_s ds - \frac{\sigma^2}{2bT} e^{-2bT} \int_0^T e^{2bs} r_s ds. \end{aligned}$$

Using asymptotic relations (6) and (17), we obtain

$$I_{3,1} \rightarrow \frac{a\sigma^2}{2b^2}, \quad \text{a. s., as } T \rightarrow \infty. \quad (33)$$

Changing the order of integration, we rewrite $I_{3,2}$ in the following way:

$$\begin{aligned} I_{3,2} &= \frac{2\sigma^2}{T} \int_0^T Z_s e^{bs} \sqrt{r_s} \int_s^T e^{-2bt} dt dW_s = \frac{2\sigma^2}{T} \int_0^T Z_s e^{bs} \sqrt{r_s} \frac{e^{-2bs} - e^{-2bT}}{2b} dt dW_s \\ &= \frac{\sigma^2}{bT} \int_0^T e^{-bs} Z_s \sqrt{r_s} dt dW_s - \frac{\sigma^2}{bT} e^{-2bT} \int_0^T e^{bs} Z_s \sqrt{r_s} dt dW_s. \end{aligned}$$

According to Lemma 6, both stochastic integrals in the right-hand side converge to zero a. s.

Therefore, $I_{3,2} \rightarrow 0$ a. s., as $T \rightarrow \infty$. Combining this with (32) and (33), we see that

$$I_3 \rightarrow \frac{a\sigma^2}{2b^2} \quad \text{a. s., as } T \rightarrow \infty. \quad (34)$$

Finally, inserting the convergences (29), (31), (34) into the equality (28), we conclude the proof. \square

Theorem 7 enables to construct a strongly consistent estimator for the parameter (a, b) .

Theorem 8. *Define*

$$\begin{aligned} \tilde{a}_T &= \frac{\sigma^2}{2} \cdot \frac{\left(\int_0^T r_t dt \right)^2}{T \int_0^T r_t^2 dt - \left(\int_0^T r_t dt \right)^2}, \\ \tilde{b}_T &= \frac{\sigma^2}{2} \cdot \frac{T \int_0^T r_t dt}{T \int_0^T r_t^2 dt - \left(\int_0^T r_t dt \right)^2}. \end{aligned}$$

Then vector $(\tilde{a}_T, \tilde{b}_T)$ is a strongly consistent estimator of (a, b) .

Proof. It follows from the convergence (6) and Theorem 7 that

$$\tilde{b}_T = \frac{\sigma^2}{2} \cdot \frac{\frac{1}{T} \int_0^T r_t dt}{\frac{1}{T} \int_0^T r_t^2 dt - \left(\frac{1}{T} \int_0^T r_t dt\right)^2} \rightarrow \frac{\sigma^2}{2} \cdot \frac{\frac{a}{b}}{\frac{a^2}{b^2} + \frac{a\sigma^2}{2b^2} - \frac{a^2}{b^2}} = b, \quad \text{a. s., as } T \rightarrow \infty.$$

Further, from (6) we have the convergence

$$\tilde{a}_T = \tilde{b}_T \cdot \frac{1}{T} \int_0^T r_t dt \rightarrow b \cdot \frac{a}{b} = a, \quad \text{a. s., as } T \rightarrow \infty. \quad \square$$

Remark 4. For the model (14) the alternative estimator of (α, μ) can be defined as follows:

$$\tilde{\alpha}_T = \frac{\sigma^2}{2} \cdot \frac{T \int_0^T r_t dt}{T \int_0^T r_t^2 dt - \left(\int_0^T r_t dt\right)^2}, \quad \tilde{\mu}_T = \frac{1}{T} \int_0^T r_t dt.$$

The strong consistency of the estimator $(\tilde{\alpha}_T, \tilde{\mu}_T)$ follows from Theorem 8 and the relations (15).

5. SIMULATIONS

In this section we illustrate the quality of the estimators, using Monte Carlo simulation. For each set of parameters (a, b, σ) , we generate 100 sample paths of the solution $r = \{r_t, t \in [0, T]\}$ to the equation (1) using Euler's approximation. We choose the initial value $r_0 = 1$ for all simulations. Then we compute means and standard deviations of the estimators at the times $T = 10, 50, 100, 150, 200$.

In the ergodic case $2a > \sigma^2$ we compare two estimators, (\hat{a}_T, \hat{b}_T) and $(\tilde{a}_T, \tilde{b}_T)$. Tables 1 and 2 report the simulation results concerning the estimation of a and b respectively. We see that both estimators are consistent and behave similarly, however, the maximum likelihood estimator (\hat{a}_T, \hat{b}_T) slightly outperforms the alternative estimator $(\tilde{a}_T, \tilde{b}_T)$.

For the non-ergodic case $2a < \sigma^2$ the maximum likelihood estimator does not make sense, so only the means and variances for $(\tilde{a}_T, \tilde{b}_T)$ are reported, see Tables 3 and 4. Clearly, the numerical results confirm the consistency of this estimator. However, in many cases the rate of convergence is quite slow (compared to the ergodic case), especially for \tilde{b}_T .

Table 1: Means and standard deviations of \hat{a}_T and \tilde{a}_T for $2a > \sigma^2$

a	b	σ		T				
				10	50	100	150	200
1	1	1	$\mathbf{E}[\hat{a}_T]$	1.3520	1.0507	1.0187	1.0105	1.0073
			$\boldsymbol{\sigma}[\hat{a}_T]$	0.5357	0.1628	0.0983	0.0818	0.0723
			$\mathbf{E}[\tilde{a}_T]$	1.5328	1.1152	1.0535	1.0300	1.0226
			$\boldsymbol{\sigma}[\tilde{a}_T]$	0.5804	0.2435	0.1701	0.1306	0.1203
1	2	1	$\mathbf{E}[\hat{a}_T]$	1.2013	1.0401	1.0144	1.0082	1.0046
			$\boldsymbol{\sigma}[\hat{a}_T]$	0.3201	0.1077	0.0700	0.0557	0.0495
			$\mathbf{E}[\tilde{a}_T]$	1.3028	1.0766	1.0270	1.0206	1.0140
			$\boldsymbol{\sigma}[\tilde{a}_T]$	0.4304	0.1618	0.1051	0.0974	0.0872
1	3	1	$\mathbf{E}[\hat{a}_T]$	1.0915	1.0201	1.0192	1.0129	1.0141
			$\boldsymbol{\sigma}[\hat{a}_T]$	0.2029	0.0821	0.0559	0.0470	0.0459
			$\mathbf{E}[\tilde{a}_T]$	1.1142	1.0411	1.0241	1.0252	1.0239
			$\boldsymbol{\sigma}[\tilde{a}_T]$	0.2961	0.1282	0.0944	0.0810	0.0778
2	1	1	$\mathbf{E}[\hat{a}_T]$	2.8227	2.0724	1.9885	1.9909	1.9963
			$\boldsymbol{\sigma}[\hat{a}_T]$	1.1649	0.3163	0.2316	0.2087	0.1908
			$\mathbf{E}[\tilde{a}_T]$	2.8584	2.1097	2.0035	2.0030	2.0157
			$\boldsymbol{\sigma}[\tilde{a}_T]$	1.1089	0.4139	0.3142	0.2724	0.2446
2	2	1	$\mathbf{E}[\hat{a}_T]$	2.2837	2.0511	2.0353	2.0170	2.0216
			$\boldsymbol{\sigma}[\hat{a}_T]$	0.6633	0.2375	0.1662	0.1365	0.1173
			$\mathbf{E}[\tilde{a}_T]$	2.4734	2.1057	2.0447	2.0057	2.0065
			$\boldsymbol{\sigma}[\tilde{a}_T]$	0.7880	0.2990	0.2134	0.1784	0.1560
2	3	1	$\mathbf{E}[\hat{a}_T]$	2.2114	2.0113	2.0120	2.0098	2.0091
			$\boldsymbol{\sigma}[\hat{a}_T]$	0.5337	0.1986	0.1509	0.1057	0.0915
			$\mathbf{E}[\tilde{a}_T]$	2.2874	2.0208	2.0224	2.0195	2.0103
			$\boldsymbol{\sigma}[\tilde{a}_T]$	0.5829	0.2602	0.1825	0.1379	0.1217
3	1	1	$\mathbf{E}[\hat{a}_T]$	3.9869	3.1923	3.0761	3.0475	3.0538
			$\boldsymbol{\sigma}[\hat{a}_T]$	1.2874	0.5061	0.4039	0.3421	0.2991
			$\mathbf{E}[\tilde{a}_T]$	3.7772	3.1524	3.0538	3.0347	3.0475
			$\boldsymbol{\sigma}[\tilde{a}_T]$	1.3190	0.5808	0.4591	0.4042	0.3519
3	1	2	$\mathbf{E}[\hat{a}_T]$	3.7755	3.1447	3.0612	3.0358	3.0359
			$\boldsymbol{\sigma}[\hat{a}_T]$	1.2919	0.3776	0.2618	0.2124	0.1919
			$\mathbf{E}[\tilde{a}_T]$	4.1227	3.3441	3.1022	3.0743	3.0872
			$\boldsymbol{\sigma}[\tilde{a}_T]$	1.5027	0.6588	0.5050	0.4323	0.4014

Table 2: Means and standard deviations of \hat{b}_T and \tilde{b}_T for $2a > \sigma^2$

a	b	σ		T				
				10	50	100	150	200
1	1	1	$\mathbf{E}[\hat{b}_T]$	1.4514	1.0772	1.0244	1.0176	1.0132
			$\boldsymbol{\sigma}[\hat{b}_T]$	0.5969	0.2184	0.1398	0.1086	0.0986
			$\mathbf{E}[\tilde{b}_T]$	1.6350	1.1459	1.0554	1.0363	1.0271
			$\boldsymbol{\sigma}[\tilde{b}_T]$	0.6687	0.2983	0.1829	0.1437	0.1293
1	2	1	$\mathbf{E}[\hat{b}_T]$	2.4399	2.1229	2.0677	2.0335	2.0159
			$\boldsymbol{\sigma}[\hat{b}_T]$	0.6993	0.2658	0.1941	0.1693	0.1587
			$\mathbf{E}[\tilde{b}_T]$	2.5135	2.1743	2.0851	2.0526	2.0303
			$\boldsymbol{\sigma}[\tilde{b}_T]$	0.7686	0.3495	0.2628	0.2393	0.2178
1	3	1	$\mathbf{E}[\hat{b}_T]$	3.3578	3.0316	3.0287	3.0094	3.0162
			$\boldsymbol{\sigma}[\hat{b}_T]$	0.8546	0.3701	0.2479	0.2071	0.1949
			$\mathbf{E}[\tilde{b}_T]$	3.1754	3.0495	3.0243	3.0313	3.0339
			$\boldsymbol{\sigma}[\tilde{b}_T]$	0.8816	0.4501	0.3351	0.2731	0.2578
2	1	1	$\mathbf{E}[\hat{b}_T]$	1.4767	1.0521	0.9980	0.9967	0.9998
			$\boldsymbol{\sigma}[\hat{b}_T]$	0.6254	0.1944	0.1323	0.1214	0.1101
			$\mathbf{E}[\tilde{b}_T]$	1.5466	1.0813	1.0102	1.0056	1.0120
			$\boldsymbol{\sigma}[\tilde{b}_T]$	0.5977	0.2338	0.1702	0.1475	0.1333
2	2	1	$\mathbf{E}[\hat{b}_T]$	2.3087	2.0703	2.0437	2.0210	2.0216
			$\boldsymbol{\sigma}[\hat{b}_T]$	0.6851	0.2865	0.1996	0.1600	0.1324
			$\mathbf{E}[\tilde{b}_T]$	2.4894	2.1255	2.0536	2.0104	2.0074
			$\boldsymbol{\sigma}[\tilde{b}_T]$	0.7913	0.3422	0.2441	0.2007	0.1755
2	3	1	$\mathbf{E}[\hat{b}_T]$	3.3561	3.0417	3.0420	3.0354	3.0311
			$\boldsymbol{\sigma}[\hat{b}_T]$	0.8854	0.3546	0.2705	0.1811	0.1619
			$\mathbf{E}[\tilde{b}_T]$	3.4176	3.0446	3.0535	3.0465	3.0305
			$\boldsymbol{\sigma}[\tilde{b}_T]$	0.9314	0.4384	0.3185	0.2254	0.1988
3	1	1	$\mathbf{E}[\hat{b}_T]$	1.3899	1.0798	1.0337	1.0264	1.0262
			$\boldsymbol{\sigma}[\hat{b}_T]$	0.5563	0.1882	0.1525	0.1283	0.1109
			$\mathbf{E}[\tilde{b}_T]$	1.3878	1.0797	1.0326	1.0264	1.0268
			$\boldsymbol{\sigma}[\tilde{b}_T]$	0.5825	0.2175	0.1688	0.1451	0.1250
3	1	2	$\mathbf{E}[\hat{b}_T]$	1.3616	1.0617	1.0209	1.0038	0.9989
			$\boldsymbol{\sigma}[\hat{b}_T]$	0.4884	0.2102	0.1361	0.1011	0.0977
			$\mathbf{E}[\tilde{b}_T]$	1.6055	1.1483	1.0447	1.0220	1.0185
			$\boldsymbol{\sigma}[\tilde{b}_T]$	0.6558	0.3057	0.2179	0.1673	0.1519

Table 3: Means and standard deviations \tilde{a}_T for $2a < \sigma^2$

a	b	σ		T				
				10	50	100	150	200
1	1	2	$\mathbf{E}[\tilde{a}_T]$	1.6220	1.1125	1.0583	1.0384	1.0348
			$\boldsymbol{\sigma}[\tilde{a}_T]$	0.6134	0.2720	0.2164	0.1826	0.1589
1	1	3	$\mathbf{E}[\tilde{a}_T]$	1.3276	1.1253	1.1256	1.1407	1.1091
			$\boldsymbol{\sigma}[\tilde{a}_T]$	2.5786	1.0947	0.6717	0.5451	0.4372
1	2	2	$\mathbf{E}[\tilde{a}_T]$	1.3309	1.0907	1.0570	1.0357	1.0212
			$\boldsymbol{\sigma}[\tilde{a}_T]$	0.3762	0.2141	0.1442	0.1317	0.1144
1	2	3	$\mathbf{E}[\tilde{a}_T]$	1.2227	1.1028	1.1034	1.0695	1.0367
			$\boldsymbol{\sigma}[\tilde{a}_T]$	2.8947	0.6768	0.4528	0.3581	0.2840
1	3	2	$\mathbf{E}[\tilde{a}_T]$	1.3113	1.1099	1.0618	1.0432	1.0330
			$\boldsymbol{\sigma}[\tilde{a}_T]$	0.3936	0.1917	0.1263	0.1137	0.1020
1	3	3	$\mathbf{E}[\tilde{a}_T]$	1.3829	1.2046	1.0947	1.0544	1.0404
			$\boldsymbol{\sigma}[\tilde{a}_T]$	2.2021	0.8893	0.3813	0.2773	0.2507
2	1	3	$\mathbf{E}[\tilde{a}_T]$	3.3746	2.3050	2.1699	2.1496	2.1363
			$\boldsymbol{\sigma}[\tilde{a}_T]$	1.5845	0.5720	0.4390	0.3586	0.3095
2	2	3	$\mathbf{E}[\tilde{a}_T]$	2.7734	2.2172	2.1665	2.1346	2.1137
			$\boldsymbol{\sigma}[\tilde{a}_T]$	0.9826	0.4067	0.3020	0.2526	0.2273
2	3	3	$\mathbf{E}[\tilde{a}_T]$	2.5827	2.1628	2.1024	2.0647	2.0350
			$\boldsymbol{\sigma}[\tilde{a}_T]$	0.7142	0.3657	0.2816	0.2316	0.1994
3	1	3	$\mathbf{E}[\tilde{a}_T]$	4.5037	3.4455	3.2534	3.1348	3.0772
			$\boldsymbol{\sigma}[\tilde{a}_T]$	1.6354	0.7223	0.5435	0.5096	0.4290

Table 4: Means and standard deviations \tilde{b}_T for $2a < \sigma^2$

a	b	σ		T				
				10	50	100	150	200
1	1	2	$\mathbf{E}[\tilde{b}_T]$	2.0929	1.2595	1.1232	1.0907	1.0621
			$\sigma[\tilde{b}_T]$	1.2187	0.4458	0.3275	0.2782	0.2326
1	1	3	$\mathbf{E}[\tilde{b}_T]$	1.4055	1.0156	1.1587	1.2190	1.1632
			$\sigma[\tilde{b}_T]$	2.6746	1.9533	0.9132	0.7432	0.6196
1	2	2	$\mathbf{E}[\tilde{b}_T]$	2.9370	2.2790	2.1923	2.1499	2.1059
			$\sigma[\tilde{b}_T]$	1.2131	0.5691	0.3801	0.3290	0.2883
1	2	3	$\mathbf{E}[\tilde{b}_T]$	1.7045	2.2836	2.3097	2.2555	2.1756
			$\sigma[\tilde{b}_T]$	9.6195	2.1124	1.1064	0.8718	0.6762
1	3	2	$\mathbf{E}[\tilde{b}_T]$	3.9779	3.3292	3.2273	3.1416	3.1023
			$\sigma[\tilde{b}_T]$	1.5206	0.7385	0.5665	0.5010	0.4566
1	3	3	$\mathbf{E}[\tilde{b}_T]$	3.8650	3.8265	3.4710	3.2583	3.1462
			$\sigma[\tilde{b}_T]$	13.9606	4.6146	1.5811	0.9728	0.8789
2	1	3	$\mathbf{E}[\tilde{b}_T]$	2.2511	1.2859	1.1613	1.1223	1.1047
			$\sigma[\tilde{b}_T]$	1.6015	0.4753	0.3403	0.2929	0.2429
2	2	3	$\mathbf{E}[\tilde{b}_T]$	3.2242	2.2986	2.2252	2.1454	2.1101
			$\sigma[\tilde{b}_T]$	1.5489	0.5355	0.3979	0.3355	0.2954
2	3	3	$\mathbf{E}[\tilde{b}_T]$	4.1619	3.3906	3.2490	3.1841	3.1132
			$\sigma[\tilde{b}_T]$	1.7422	0.7886	0.5821	0.4935	0.4223
3	1	3	$\mathbf{E}[\tilde{b}_T]$	2.0684	1.3138	1.1622	1.1039	1.0754
			$\sigma[\tilde{b}_T]$	0.9192	0.3868	0.2481	0.2418	0.2081

BIBLIOGRAPHY

- A. Alfonsi. *Affine diffusions and related processes: simulation, theory and applications*, volume 6 of *Bocconi & Springer Series*. Springer, Cham; Bocconi University Press, Milan, 2015.
- M. Barczy, M. Ben Alaya, A. Kebaier, and G. Pap. Asymptotic properties of maximum likelihood estimator for the growth rate of a stable CIR process based on continuous time observations. *Statistics*, 53(3):533–568, 2019.
- M. Bel Hadj Khelifa, Y. Mishura, K. Ralchenko, and M. Zili. Drift parameter estimation in stochastic differential equation with multiplicative stochastic volatility. *Mod. Stoch. Theory Appl.*, 3(4):269–285, 2016.
- M. Ben Alaya and A. Kebaier. Parameter estimation for the square-root diffusions: ergodic and nonergodic cases. *Stoch. Models*, 28(4):609–634, 2012.
- M. Ben Alaya and A. Kebaier. Asymptotic behavior of the maximum likelihood estimator for ergodic and nonergodic square-root diffusions. *Stoch. Anal. Appl.*, 31(4):552–573, 2013.
- A. Cozma and C. Reisinger. Exponential integrability properties of Euler discretization schemes for the Cox–Ingersoll–Ross process. *Discrete Contin. Dyn. Syst. Ser. B*, 21(10):3359–3377, 2016.
- G. De Rossi. Maximum likelihood estimation of the Cox–Ingersoll–Ross model using particle filters. *Computational Economics*, 36(1):1–16, 2010.
- G. Deelstra and F. Delbaen. Long-term returns in stochastic interest rate models. *Insurance Math. Econom.*, 17(2):163–169, 1995.
- S. Dereich, A. Neuenkirch, and L. Szpruch. An Euler-type method for the strong approximation of the Cox–Ingersoll–Ross process. *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 468(2140):1105–1115, 2012.

- I. Karatzas and S. E. Shreve. *Brownian motion and stochastic calculus*, volume 113 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991.
- K. Kládívko. Maximum likelihood estimation of the cox–ingersoll–ross process: the matlab implementation. *Technical Computing Prague*, 7, 2007.
- Y. A. Kutoyants. *Statistical inference for ergodic diffusion processes*. Springer Series in Statistics. Springer-Verlag London, Ltd., London, 2004.
- Z. Li and C. Ma. Asymptotic properties of estimators in a stable Cox–Ingersoll–Ross model. *Stochastic Process. Appl.*, 125(8):3196–3233, 2015.
- R. Liptser and A. N. Shiriyayev. *Theory of martingales*, volume 49. Springer Science & Business Media, 2012.
- A. Mijatović and M. Urusov. Convergence of integral functionals of one-dimensional diffusions. *Electron. Commun. Probab.*, 17:no. 61, 13, 2012.
- G. N. Milstein and J. Schoenmakers. Uniform approximation of the Cox–Ingersoll–Ross process. *Adv. in Appl. Probab.*, 47(4):1132–1156, 2015.
- Y. Mishura and Y. Munchak. Functional limit theorems for additive and multiplicative schemes in the Cox–Ingersoll–Ross model. *Mod. Stoch. Theory Appl.*, 3(1):1–17, 2016.
- Y. Mishura and G. Shevchenko. *Theory and Statistical Applications of Stochastic Processes*. ISTE, 2017.
- L. Overbeck and T. Rydén. Estimation in the Cox–Ingersoll–Ross model. *Econometric Theory*, 13(3):430–461, 1997.
- A. V. Skorokhod. *Asymptotic methods in the theory of stochastic differential equations*, volume 78 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1989.