

NON-LINEAR LOG-SOBOLEV INEQUALITIES FOR THE POTTS SEMIGROUP AND APPLICATIONS TO RECONSTRUCTION PROBLEMS

YUZHOU GU, YURY POLYANSKIY

ABSTRACT. Consider a Markov process with state space $[k]$, which jumps (after an exponentially distributed time) to a new state chosen uniformly at random and regardless of the previous state. The collection of transition kernels (indexed by time $t \geq 0$) is the Potts semigroup. Diaconis and Saloff-Coste [DSC96] computed the maximum of the ratio of the relative entropy and the Dirichlet form obtaining the constant α_2 in the 2-log-Sobolev inequality (2-LSI). In this paper, we obtain the best possible *non-linear* inequality relating entropy and the Dirichlet form (i.e., p -NLSI, $p \geq 1$). As an example, we show $\alpha_1 = 1 + \frac{1+o(1)}{\log k}$. The more precise NLSIs have been shown by Polyanskiy and Samorodnitsky [PS19] to imply various geometric and Fourier-analytic results.

Beyond the Potts semigroup, we also analyze Potts channels – Markov transition matrices $[k] \times [k]$ constant on and off diagonal. (Potts semigroup corresponds to a (ferromagnetic) subset of matrices with positive second eigenvalue). By integrating the 1-NLSI we obtain the new strong data processing inequality (SDPI), which in turn allows us to improve results on reconstruction thresholds for Potts models on trees. A special case is the problem of reconstructing color of the root of a k -colored tree given knowledge of colors of all the leaves. We show that to have a non-trivial reconstruction probability the branching number of the tree should be at least

$$\frac{\log k}{\log k - \log(k-1)} = (1 - o(1))k \log k.$$

This extends previous results (of Sly [Sly09b] and Bhatnagar et al. [BVVW11]) to general trees, and avoids the need for any specialized arguments. Similarly, we improve the state-of-the-art on reconstruction threshold for the stochastic block model with k balanced groups, for all $k \geq 3$. These improvements advocate information-theoretic methods as a useful complement to the conventional techniques originating from the statistical physics.

CONTENTS

1.	Introduction	2
2.	Non-linear p -log-Sobolev inequalities for the Potts semigroup	8
2.1.	Non-linear p -log-Sobolev inequality for $p > 1$	9
2.2.	Non-linear 1-log-Sobolev inequality	12
2.3.	Input-restricted non-linear SDPI for Potts channels	14
2.4.	Behavior for $k \rightarrow \infty$	18
3.	Product spaces	20

Y.G. and Y.P. {emails: yuzhougu@mit.edu, yp@mit.edu} are with the Institute for Data, Systems, and Society and the Department of Electrical Engineering and Computer Science, Massachusetts Institute of Technology, Cambridge, MA 02139, USA.

3.1. Tensorization	21
3.2. Linear piece	23
3.3. Edge isoperimetric inequalities	25
4. Non-reconstruction for broadcast models on trees	26
5. Potts model on a tree	29
5.1. Binary symmetric channel	30
5.2. Random coloring	30
6. Stochastic block model	32
6.1. Impossibility of weak recovery via information percolation	33
6.2. Impossibility of weak recovery via Potts channel	34
Appendix A. Input-unrestricted contraction coefficient for Potts channels	35
Appendix B. An upper bound for input-restricted contraction coefficient for Potts channels	39
Appendix C. Non-convexity of certain functions	42
Appendix D. Concavity of log-Sobolev coefficients	45
Acknowledgement	46
References	46

1. INTRODUCTION

Log-Sobolev inequalities. Log-Sobolev inequalities (LSIs) are a class of inequalities bounding the rate of convergence of a Markov semigroup to its stationary distribution. They upper bound certain relative entropy (KL divergence) functions via a multiple of the Dirichlet form.

Let us introduce some standard notions. Let \mathcal{X} be a finite alphabet and $K : \mathcal{X} \times \mathcal{X} \rightarrow [0, 1]$ be a Markov kernel, i.e., for all $x \in \mathcal{X}$, we have $\sum_{y \in \mathcal{X}} K(x, y) = 1$. Let $L = K - I$. We consider the semigroup $(T_t)_{t \geq 0}$, where $T_t = \exp(tL)$. Let π be a stationary measure for the semigroup. For $f, g : \mathcal{X} \rightarrow \mathbb{R}$, the Dirichlet form is defined by

$$\mathcal{E}(f, g) := -\mathbb{E}_\pi[(Lf)g] = - \sum_{x, y \in \mathcal{X}} L(x, y) f(y) g(x) \pi(x).$$

For non-zero $f : \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}$, the relative entropy is defined by

$$\text{Ent}_\pi(f) := \mathbb{E}_\pi \left[f \log \frac{f}{\mathbb{E}_\pi[f]} \right] = \mathbb{E}_\pi[f] D(\pi^{(f)} || \pi)$$

where $\pi^{(f)}$ is a distribution defined as $\pi^{(f)}(x) = \frac{f(x)\pi(x)}{\mathbb{E}_\pi[f]}$, and $D(P||Q)$ is the Kullback-Leibler divergence¹

$$D(P||Q) := \int \log\left(\frac{dP}{dQ}\right) dP.$$

For $p > 1$, we say the semigroup $(T_t)_{t \geq 0}$ admits p -log-Sobolev inequality (p -LSI), if for some constant α_p , for all non-zero non-negative real functions f on \mathcal{X} , we have

$$\text{Ent}_\pi(f) \leq \frac{1}{\alpha_p} \mathcal{E}(f^{\frac{1}{p}}, f^{1-\frac{1}{p}}).$$

¹Throughout this paper, log means natural logarithm.

For $p = 1$, we define 1-LSI as

$$\text{Ent}_\pi(f) \leq \frac{1}{\alpha_1} \mathcal{E}(f, \log f).$$

The case $p = 2$ is the standard log-Sobolev inequality, originally studied in Gross [Gro75]. The case $p = 1$ is studied also under the name “modified log-Sobolev inequality” (e.g. [BT06]).

The relationship between 1-LSI and semigroup convergence can be seen from the following identity

$$(1) \quad \frac{d}{dt} \Big|_{t=0} \text{Ent}_\pi(T_t f) = -\mathcal{E}(f, \log f).$$

Therefore

$$(2) \quad \text{Ent}_\pi(T_t f) \leq \exp(-\alpha_1 t) \text{Ent}_\pi(f)$$

which corresponds to a property of T_t to exponentially fast relax to equilibrium (in the sense of relative entropy).

Polyanskiy and Samorodnitsky [PS19] introduced non-linear p -log-Sobolev inequalities (p -NLSI), a finer description of the relationship between relative entropy and Dirichlet forms. For $p \geq 1$, we say the semigroup satisfies p -LSI if for some non-negative function². $\Phi_p : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$, for all non-zero $f : \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}$, we have

$$(3) \quad \frac{\text{Ent}_\pi(f)}{\mathbb{E}_\pi[f]} \leq \Phi_p\left(\frac{\mathcal{E}(f^{\frac{1}{p}}, f^{1-\frac{1}{p}})}{\mathbb{E}_\pi[f]}\right),$$

where for $p = 1$, $\mathcal{E}(f^{\frac{1}{p}}, f^{1-\frac{1}{p}})$ should be replaced with $\mathcal{E}(f, \log f)$.

Non-linear p -log-Sobolev inequalities imply the ordinary p -log-Sobolev inequalities for

$$\alpha_p = \inf_{x>0} \frac{x}{\Phi_p(x)}.$$

When Φ_p is concave, this can be further simplified to $\alpha_p = (\Phi_p'(0))^{-1}$.

Mossel et al. [MOS13] proved that for reversible (K, π) ,

$$(4) \quad \frac{p^2(q-1)}{q^2(p-1)} \alpha_p \leq \alpha_q \leq \alpha_p$$

for $1 < q \leq p \leq 2$. We discuss some general facts about dependence of α_p and Φ_p on p in Appendix D.

Potts semigroup. In this paper, we focus on the simplest Markov semigroup, corresponding to the random walk on a complete graph. The Markov kernel is $K(x, y) = \frac{1}{k-1} \mathbb{1}\{x \neq y\}$, where $k = \#\mathcal{X}$.³ We call it the Potts semigroup, because every operator T_t in the semigroup is a ferromagnetic Potts channel. Its stationary distribution π is uniform on \mathcal{X} and its Dirichlet form is rescaled covariance:

$$\mathcal{E}(f, g) = \frac{k}{k-1} \text{Cov}_\pi(f, g)$$

A Potts channel PC_λ is a Markov kernel $[k] \times [k] \rightarrow [0, 1]$ defined by

$$\text{PC}_\lambda(x, y) = \frac{1-\lambda}{k} \mathbb{1}\{x \neq y\} + \left(\frac{1}{k} + \frac{k-1}{k} \lambda\right) \mathbb{1}\{x = y\}.$$

²[PS19] requires the function Φ_p to be concave. We do not make this assumption initially, however to extend these inequalities to product semigroups the concavification will be necessary – see Section 3.

³In the following, we always write $\mathcal{X} = [k]$.

We parametrize them by λ because λ is the second largest eigenvalue of PC_λ . The valid region of λ is $[-\frac{1}{k-1}, 1]$. When $\lambda > 0$, we call PC_λ a ferromagnetic Potts channel; when $\lambda < 0$, we call it an anti-ferromagnetic Potts channel. One can see that T_t in the Potts semigroup is exactly $\text{PC}_{\exp(-\frac{k}{k-1}t)}$.

Diaconis and Saloff-Coste [DSC96] computed the 2-log-Sobolev constant

$$(5) \quad \alpha_2 = \frac{k-2}{(k-1)\log(k-1)}.$$

They observed that the infimum of the ratio $\frac{\mathcal{E}(f,f)}{\text{Ent}_\pi[f^2]}$ is achieved at a two-valued function f , i.e., f takes exactly two values. In fact, the infimum is achieved at a function f where $f(1) = k-1$ and $f(i) = 1$ for $i \neq 1$. For $p \neq 2$, it seems hard to give a closed-form expression for α_p . Goel [Goe04] proved that

$$\frac{k}{k-1} \leq \alpha_1 \leq \left(1 + \frac{4}{\log(k-1)}\right) \frac{k}{k-1},$$

where the upper bound is by using a two-valued function f , where $f(1) = k+1$ and $f(i) = 1$ for $i \neq 1$. Bobkov and Tetali [BT06] also discussed bounds on α_1 and α_2 , proving that

$$\alpha_1 \geq \frac{k}{k-1} + \frac{2}{\sqrt{k-1}}.$$

These computations lead to the guess that for all p , the best possible p -LSI constant α_p for the Potts semigroup is achieved at a two-valued function. In Section 2, we prove that this is true, and in fact true for p -NLSIs for the Potts semigroup: For fixed $\frac{\text{Ent}_\pi(f)}{\mathbb{E}_\pi[f]}$, the unique function (up to scalar multiplication) of the form $f(1) \geq f(2) = \dots = f(k)$ minimizes $\frac{\mathcal{E}(f^{\frac{1}{p}}, f^{1-\frac{1}{p}})}{\mathbb{E}_\pi[f]}$. As a result we get the sharpest p -NLSIs for the Potts semigroup for all $p \geq 1$.

We define a useful function $\psi : [0, 1] \rightarrow \mathbb{R}$ as follows.

$$(6) \quad \psi(x) := \log k + x \log x + (1-x) \log \frac{1-x}{k-1}.$$

Note that $\psi(x)$ is the KL divergence between $(x, \frac{1-x}{k-1}, \dots, \frac{1-x}{k-1})$ and $\text{Unif}([k])$. Simple computation shows that ψ is non-negative, convex, $\psi(\frac{1}{k}) = 0$, strictly decreasing on $[0, \frac{1}{k}]$, strictly increasing on $[\frac{1}{k}, 1]$, and takes value in $[0, \log k]$.

Theorem 1 (p -NLSI for Potts semigroup). *For $p > 1$, define $\xi_p : [0, 1] \rightarrow \mathbb{R}$ as*

$$(7) \quad \xi_p(x) = \frac{k}{k-1} \left(1 - \frac{1}{k} \left(x^{\frac{1}{p}} + (k-1) \left(\frac{1-x}{k-1}\right)^{\frac{1}{p}}\right) \left(x^{1-\frac{1}{p}} + (k-1) \left(\frac{1-x}{k-1}\right)^{1-\frac{1}{p}}\right)\right).$$

Define $\xi_1 : [0, 1] \rightarrow \mathbb{R}$ as

$$(8) \quad \xi_1(x) = \frac{1}{k-1} \left(-\log x - (k-1) \log \frac{1-x}{k-1} + k \left(x \log x + (1-x) \log \frac{1-x}{k-1}\right)\right).$$

For $p \geq 1$, define $b_p : [0, \log k] \rightarrow \mathbb{R}$ as⁴

$$(9) \quad b_p(\psi(x)) = \xi_p(x)$$

for $x \in [\frac{1}{k}, 1]$, where ψ is defined in (6). Then the Potts semigroup satisfies p -NLSI with $\Phi_p = b_p^{-1}$.

⁴In the case $k = 2$, b_p differs from [PS19] by a constant factor due to a different parametrization of the semigroup.

In particular, we have

$$(10) \quad \alpha_p = \inf_{x \in (\frac{1}{k}, 1]} \frac{\xi_p(x)}{\psi(x)}.$$

Furthermore, this is the best possible p -NLSI for the Potts semigroup, in the sense that for any $y \in \text{dom } \Phi_p$, there exists function $f : [k] \rightarrow \mathbb{R}_{\geq 0}$ such that $\mathbb{E}_\pi f = 1$, $\mathcal{E}(f^{\frac{1}{p}}, f^{1-\frac{1}{p}}) = y$, and $\text{Ent}(f) = \Phi_p(y)$.

As a corollary of our 1-NLSI, we derive the second order behavior of α_1 as k goes to ∞ .

Proposition 2. For $k \geq 3$, we have

$$(11) \quad \frac{k}{k-1} \left(1 + \frac{1}{\log k}\right) \leq \alpha_1 \leq \frac{k}{k-1} \left(1 + \frac{1+o(1)}{\log k}\right).$$

Strong data processing inequalities. The data processing inequality⁵ states that $I(U; Y) \leq I(U; X)$ for any Markov chain $U \rightarrow X \rightarrow Y$, i.e., we cannot gain information by going through a channel. It is natural to think that when the channel $X \rightarrow Y$ is noisy, we should strictly lose information, i.e., $I(U; Y) < I(U; X)$. In fact, we have $I(U; Y) \leq \eta I(U; X)$, where the constant η depends only on the channel $P_{Y|X}$, but not on the distribution of U and X . Such inequalities are called strong data processing inequalities (SDPIs). We distinguish between the inequalities that depend on the distribution P_X and that are independent of it. Namely, fix a stochastic matrix (conditional distribution) W and input distribution (row-vector) Q_0 and define

$$\eta_{\text{KL}}(W) = \sup_{P, Q: 0 < D(P||Q) < \infty} \frac{D(PW||QW)}{D(P||Q)},$$

$$\eta_{\text{KL}}(W, Q_0) = \sup_{P: 0 < D(P||Q_0) < \infty} \frac{D(PW||Q_0W)}{D(P||Q_0)}$$

It can be shown, e.g. [PW17], that we also have alternative characterizations:

$$\eta_{\text{KL}}(W) = \sup_{U \rightarrow X \rightarrow Y} \frac{I(U; Y)}{I(U; X)},$$

$$\eta_{\text{KL}}(W, Q_0) = \sup_{U \rightarrow X \rightarrow Y, X \sim Q_0} \frac{I(U; Y)}{I(U; X)},$$

where $\mathbb{P}[Y = y|X = x] = W_{x,y}$. Even more generally, some channels can be shown to satisfy (for all Markov chains $U \rightarrow X \rightarrow Y$ with $P_{Y|X}$ as before, arbitrary U and fixed or arbitrary P_X)

$$(12) \quad I(U; Y) \leq s(I(U; X)),$$

for some non-linear function s .

From (2) and Proposition 2 we obtain

$$(13) \quad \eta_{\text{KL}}(\text{PC}_\lambda, \pi) \leq \lambda^{\frac{k-1}{k}\alpha_1} = \lambda^{1+\frac{1+o(1)}{\log k}}$$

for $\lambda \in [0, 1]$ and $o(1) \rightarrow 0$ as $k \rightarrow \infty$. It turns out that 1-NLSI can be seen as an infinitesimal version of the non-linear SDPIs (see [PS19, Theorem 2]). Thus, we can prove the best possible non-linear SDPI for Potts channels.

⁵We recall that for a pair of random variables X, Y we define $I(X; Y) = D(P_{X,Y}||P_X P_Y)$.

Theorem 3 (Non-linear SDPI for Potts channel). Fix $\lambda \in [-\frac{1}{k-1}, 1]$. Define $s_\lambda : [0, \log k] \rightarrow \mathbb{R}$ as

$$s_\lambda(\psi(x)) = \psi\left(\lambda x + \frac{1-\lambda}{k}\right),$$

for $x \in [\frac{1}{k}, 1]$, where ψ is defined in (6). Let \hat{s}_λ be the concave envelope of s_λ . For any Markov chain $U \rightarrow X \rightarrow Y$ where X has uniform distribution and $X \rightarrow Y$ is the Potts channel PC_λ , we have

$$I(U; Y) \leq \hat{s}_\lambda(I(U; X)).$$

In particular, we have

$$(14) \quad \eta_{\text{KL}}(\text{PC}_\lambda, \pi) = \sup_{x \in (\frac{1}{k}, 1]} \frac{\psi(\lambda x + \frac{1-\lambda}{k})}{\psi(x)}.$$

Furthermore, this is the best possible non-linear SDPI for Potts channels, in the sense that for any $c \in [0, \log k]$, there exists a Markov chain $U \rightarrow X \rightarrow Y$ where X has uniform distribution, $X \rightarrow Y$ is the Potts channel PC_λ , and $I(U; X) = c$, such that $I(U; Y) = \hat{s}_\lambda(c)$.

To compare the input-restricted η_{KL} with input-unrestricted one, in Appendix A we compute the exact value of $\eta_{\text{KL}}(\text{PC}_\lambda)$, and in Appendix B we prove that

$$(15) \quad \eta_{\text{KL}}(\text{PC}_\lambda, \pi) < \eta_{\text{KL}}(\text{PC}_\lambda)$$

for $k \geq 3$ and $\lambda \in [-\frac{1}{k-1}, 0) \cup (0, 1)$. See Section 2.4 for discussion on the tightness of the bound (13).

Remark 4 (Tensorization). In Section 3 we extend p -NLSI and SDPIs to product spaces/channels. In these results, the functions \check{b}_p (convexification of b_p) and \hat{s}_λ (concavification of s_λ) appear naturally. When $k = 2$, b_p is already convex, and s_λ is concave, leading to many good properties for the hypercube and for binary symmetric channels (e.g. Mrs. Gerber's Lemma [WZ73]). However, as shown in Proposition 25, these properties do not hold anymore for $k \geq 3$, implying a different structure of extremal distributions that are the slowest to relax to equilibrium as $t \rightarrow \infty$ in $T_t^{\otimes n}$, see Section 3.2.

Applications.

One of the implications of NLSIs are improved hypercontractivity inequalities for functions in $[k]^n$ supported on subsets of cardinality $k^{(1-\epsilon)n}$ – this was established generally (for any semigroup) in [PS19]. Here, we show how NLSIs can be used to close the gap (between functional-analytic proofs and explicit combinatorics of [Lin64]) in the edge-isoperimetric inequality for the $[k]^n$ – see Section 3.3.

Similarly, SDPIs have numerous applications. Originally introduced to study certain multi-user data-compression questions in information theory, they have been since adopted in many different scenarios. For example, Evans and Schulman [ES99] use SDPIs to investigate fundamental limits of fault tolerant computing. Polyanskiy and Wu [PW17, PW18] further developed the idea and related the amount of information transmitted in a directed or undirected graphical model in terms of the percolation probability (existence of an open path) on the same network. Other notable applications include distributed estimation [XR15, BGM⁺16] and communication complexity [HLPS19].

More directly related to our paper is the work of [EKPS00] which applies an SDPI (for a Potts channel with $k = 2$) to bound the threshold on for reconstruction

problem for the Ising model on a tree. We describe a more general class of such questions.

Consider a tree with a marked root. Each vertex of the tree has a random spin in $[k]$, generated in the following way: the root spin is generated according to some known distribution, and for each vertex, its spin is generated from its parent's spin, through some channel M . We say the problem has reconstruction if given spins of all nodes far away enough from the root, we can guess the root spin better than guessing from the initial distribution. Equivalently, the problem has reconstruction if the mutual information between the root spin and the spins of all nodes distance d away from the root goes to a non-zero limit, as d goes to ∞ . The problem has non-reconstruction otherwise.

Reconstruction problems trees have been studied for a long time. Kesten and Stigum [KS66] proved the so-called Kesten-Stigum bound, a reconstruction result based on the second eigenvalue of the channel. Physicists study the problem from a spin glass theoretic perspective (e.g., Mézard and Montanari [MM06]). Based on careful analysis of the evolution of magnetization, several authors have obtained very tight reconstruction thresholds for various models (e.g., Sly [Sly09a, Sly09b], Bhatnagar et al. [BST10], Liu and Ning [LN19]).

In this paper, we attack this problem using SDPIs. The method of [EKPS00] showed that reconstruction is impossible if $\text{br}(T)\eta_{\text{KL}}(M) < 1$. Here we improve their result by considering the input-restricted η_{KL} . We note that the idea of using input-restricted (but not for η_{KL}) has appeared in [FK09b] – see Remark 34 below.

Theorem 5. *Consider the broadcast model on a tree T with channel M . Let q^* be a stationary distribution, i.e., $q^*M = q^*$. Let M^\vee denote the reverse channel, i.e., any channel that satisfies $q_j^*M_{j,i}^\vee = q_i^*M_{i,j}$ for all $i, j \in [k]$. Then the model has non-reconstruction if*

$$\eta_{\text{KL}}(M^\vee, q^*) \text{br}(T) < 1,$$

where $\text{br}(T)$ is the branching number of T , whose definition is given in Definition 28.

Our method is very simple, is non-asymptotic, works for the branching number, and is often very tight. Previous results often impose additional restrictions on the tree (e.g., regular tree, or Galton-Watson with Poisson offspring distribution), and some only work as expected degree goes to infinity. We discuss in more detail in Section 4.

Applying Theorem 5 to the Potts model, we achieve improved non-reconstruction results for Potts models on a tree.

Theorem 6. *Consider the Potts model on a tree T . We have non-reconstruction whenever*

$$\eta_{\text{KL}}(\text{PC}_\lambda, \pi) \text{br}(T) < 1,$$

where $\eta_{\text{KL}}(\text{PC}_\lambda, \pi)$ is given by (14).

Theorem 6 strictly improves over the explicit bound of Mossel and Peres [MP03]. In the special case where the channel is the coloring channel ($\lambda = -\frac{1}{k-1}$), Theorem 6 recovers the reconstruction threshold up to the first order, which was previously obtained by Sly [Sly09b] and Bhatnagar et al. [BVVW11] using more complicated methods. (We note, however, that a major focus of those works was to obtain the lower-order terms.) More detailed analysis is done in Section 5.

Last but not least, we consider the problem of weak recovery for the stochastic block model with k communities (k -SBM). In k -SBM, n vertices each receive a random spin in $[k]$, and then a random graph is constructed, such that (1) for two vertices with the same spin, there exists an edge with probability $\frac{a}{n}$; (2) for two vertices with different spins, there exists an edge with probability $\frac{b}{n}$. The model is said to have weak recovery, if given the random graph, we can partition the vertices into k parts, such that the partition is correct (up to relabeling the parts) for at least ϵn vertices, for some absolute constant ϵ .

For $k = 2$, the weak recovery threshold is known: If $(a - b)^2 > 2(a + b)$, weak recovery is possible (Massoulié [Mas14], Mossel et al. [MNS18]); if $(a - b)^2 < 2(a + b)$, weak recovery is impossible (Mossel et al. [MNS15]). For $k \geq 3$, the weak recovery threshold is not completely determined. The stochastic block model is called assortative if $a > b$, and disassortative if $a < b$. The disassortative case was solved by Coja-Oghlan et al. [COKPZ18]. For the assortative case, the current best impossibility result for general k is by Banks et al. [BMNN16], which says weak recovery does not hold whenever

$$(16) \quad \frac{(a - b)^2}{a + (k - 1)b} < \frac{2k \log(k - 1)}{k - 1}.$$

By a standard reduction from the Potts model, we achieve improved impossibility results for weak recovery for the k -stochastic models.

Theorem 7. *Weak recovery of the stochastic block model is impossible if*

$$d\eta_{\text{KL}}(\text{PC}_\lambda, \pi) < 1,$$

where $d = \frac{a + (k-1)b}{k}$, $\lambda = \frac{a-b}{a + (k-1)b}$, and η_{KL} is given by (14).

We discuss in more detail, and show that this improves over previous results, in Section 6.

Organization. In Section 2 we prove the sharpest p -LSIs for the Potts semigroup (Theorem 1), and compute the input-restricted KL divergence contraction coefficients of all Potts channels (Theorem 3).

In Section 3 we discuss tensorization of p -NLSIs for the Potts semigroup, and non-linear SDPI for Potts channels.

In Section 4 we prove a non-reconstruction result for a general class of broadcast models on trees, based on strong data processing inequalities (Theorem 5).

In Section 5 we apply Theorem 5 to the Potts model on a tree (Theorem 6). We show that this improves previous non-reconstruction results. For a special case, the random coloring model on a tree, we obtain non-reconstruction results for arbitrary trees, generalizing previous results which work only for restricted classes of trees.

In Section 6, by a standard reduction from the Potts model, we prove impossibility results for weak recovery of the stochastic block model (Theorem 7). This results in improvements for the best known bounds for the k -SBM.

2. NON-LINEAR p -LOG-SOBOLEV INEQUALITIES FOR THE POTTS SEMIGROUP

In this section, we prove p -NLSIs for the Potts semigroup for $p \geq 1$. Because the form of the p -LSIs are slightly different for $p \neq 1$ and $p = 1$, we prove them separately.

Recall our setting. Alphabet $\mathcal{X} = [k]$ for some positive integer $k \geq 2$. The Potts semigroup $T_t = \exp(Lt)$ for generator $L = \frac{1}{k-1} \mathbb{1}\{x \neq y\} - \mathbb{1}\{x = y\}$. The stationary distribution is $\pi = \text{Unif}([k])$. The Dirichlet form is

$$\mathcal{E}(f, g) = -\mathbb{E}_\pi[(Lf)g] = -\frac{1}{k(k-1)} \left(\sum_x f(x) \right) \left(\sum_y g(y) \right) + \frac{1}{k-1} \sum_x f(x)g(x).$$

Relative entropy is

$$\text{Ent}_\pi(f) = \mathbb{E}_\pi \left[f \log \frac{f}{\mathbb{E}_\pi[f]} \right].$$

The non-linear p -log-Sobolev inequality says

$$\frac{\text{Ent}_\pi(f)}{\mathbb{E}_\pi[f]} \leq \Phi_p \left(\frac{\mathcal{E}(f^{\frac{1}{p}}, f^{1-\frac{1}{p}})}{\mathbb{E}_\pi[f]} \right)$$

for some concave Φ_p , where for $p = 1$, RHS is replaced with $\Phi_p \left(\frac{\mathcal{E}(f, \log f)}{\mathbb{E}_\pi[f]} \right)$. Because both sides of the inequality are fixed under scalar multiplication, we can wlog restrict f to be a distribution P . Then the relative entropy is

$$\text{Ent}_\pi(P) = \frac{1}{k} D(P|\pi) = \frac{1}{k} (\log k - H(P)).$$

2.1. Non-linear p -log-Sobolev inequality for $p > 1$. We prove Theorem 1 for $p > 1$. Before proving the theorem we show the following.

Proposition 8. *Fix $r \in (0, 1)$ and $c \in [0, \log k]$. Among all distributions $P = (p_1, \dots, p_k)$ with $H(P) = c$, the distribution of form $P = (x, \frac{1-x}{k-1}, \dots, \frac{1-x}{k-1})$ with $x \in [\frac{1}{k}, 1]$ achieves maximum $\sum_i p_i^r$. Furthermore, up to permutation of the alphabet this is the unique maximum-achieving distribution.*

Proof. The result for $c \in \{0, \log k\}$ is obvious. In the following, assume that $c \in (0, \log k)$. Write $F(P) := \sum_i p_i^r$. The set $\{P : H(P) = c\}$ is compact, so the maximum value of $F(P)$ is achieved at some point $P = (p_1, \dots, p_k)$.

Step 0.

Claim 9. *Fix $a, b > 0$ and $r \in (0, 1)$. Among all solutions $u, v, w \in [0, 1]$ with $u + v + w = a$ and $-u \log u - v \log v - w \log w = b$, the maximum of $u^r + v^r + w^r$ is not achieved at a point where $0 = u < v < w$.*

Proof. Suppose the maximum is achieved at such a point (u_0, v_0, w_0) where $0 = u_0 < v_0 < w_0$. Extend it to a curve $(u, v = v(u), w = w(u))$ on $u \in [0, \epsilon]$ for some $\epsilon > 0$, such that $u < v < w$ for all u , satisfying

$$(17) \quad u + v + w = a,$$

$$(18) \quad -u \log u - v \log v - w \log w = b,$$

and

$$v(0) = v_0, w(0) = w_0.$$

We prove that

$$f(u) := u^r + v^r + w^r$$

decreases as u approaches 0^+ , for small enough u .

By taking derivative of (17) and (18), one can compute that

$$\begin{aligned} v'(u) &= \frac{\log w - \log u}{\log v - \log w}, \\ w'(u) &= \frac{\log u - \log v}{\log v - \log w}. \end{aligned}$$

Therefore

$$\begin{aligned} f'(u) &= r(u^{r-1} + v^{r-1}v'(u) + w^{r-1}w'(u)) \\ &= r(u^{r-1} + v^{r-1}\frac{\log w - \log u}{\log v - \log w} + w^{r-1}\frac{\log u - \log v}{\log v - \log w}). \end{aligned}$$

Because $0 < v_0 < w_0$, the term u^{r-1} dominates the sum, and $f'(u) > 0$ for small enough $u > 0$. Therefore the maximum of f is not achieved at $u = 0$. \square

By Claim 9, if $p_i = 0$ for some i , then there can be at most two different values of p_i 's.

Step 1.

Claim 10. *If $u, v, w \in (0, 1)$ are all different, then*

$$\det \begin{pmatrix} 1 & \log u & u^{r-1} \\ 1 & \log v & v^{r-1} \\ 1 & \log w & w^{r-1} \end{pmatrix} \neq 0.$$

Proof of Claim. Suppose $\det = 0$. Then for some $a, b \in \mathbb{R}$, the equation $x^{r-1} + a \log x = b$ has at least three distinct solutions $x \in (0, 1)$. However

$$\frac{\partial}{\partial x}(x^{r-1} + a \log x) = (r-1)x^{r-2} + \frac{a}{x}$$

is smooth on $(0, 1)$, and takes zero at most once. So $x^{r-1} + a \log x$ takes each value at most once on $(0, 1)$. Contradiction. \square

By KKT conditions, the three vectors

$$\begin{aligned} \nabla F(P) &= (rp_i^{r-1})_{i \in [k]}, \\ \nabla H(P) &= (-1 - \log p_i)_{i \in [k]}, \\ \nabla \sum_{i \in [k]} p_i &= \mathbb{1} \end{aligned}$$

should be linear dependent. By Step 0 and Claim 10, there can be at most two different values of p_i 's.

So we can assume that $p_1 = \dots = p_m = x$, $p_{m+1} = \dots = p_k = \frac{1-mx}{k-m}$ for some $m \in [k-1]$, $x \in (\frac{1}{k}, \frac{1}{m}]$.

Step 2. For P of the above form, we have

$$\begin{aligned} -H(P) &= mx \log x + (1 - mx) \log \frac{1 - mx}{k - m}, \\ F(P) &= mx^r + (k - m) \left(\frac{1 - mx}{k - m} \right)^r. \end{aligned}$$

We smoothly continue both functions so that m can take any real value in $[1, k-1]$.

Claim 11. For $m \in (1, k-1]$ and $x \in (\frac{1}{k}, \frac{1}{m})$, we have

$$-\frac{\partial}{\partial x}H(P) > 0$$

and

$$\frac{\partial}{\partial x}H(P)\frac{\partial}{\partial m}F(P) - \frac{\partial}{\partial m}H(P)\frac{\partial}{\partial x}F(P) > 0.$$

Proof. We have

$$\begin{aligned} -\frac{\partial}{\partial x}H(P) &= m(\log x - \log \frac{1-mx}{k-m}) > 0, \\ -\frac{\partial}{\partial m}H(P) &= \frac{1-kx}{k-m} + x(\log x - \log \frac{1-mx}{k-m}), \\ \frac{\partial}{\partial x}F(P) &= rm(x^{r-1} - (\frac{1-mx}{k-m})^{r-1}), \\ \frac{\partial}{\partial m}F(P) &= x^r + (r\frac{1-kx}{1-mx} - 1)(\frac{1-mx}{k-m})^r. \end{aligned}$$

Let $a = \frac{kx-1}{1-mx}$. Then

$$\begin{aligned} G(P) &:= \frac{\partial}{\partial x}H(P)\frac{\partial}{\partial m}F(P) - \frac{\partial}{\partial m}H(P)\frac{\partial}{\partial x}F(P) \\ &= (r-1)m(x^r - (\frac{1-mx}{k-m})^r)(\log x - \log \frac{1-mx}{k-m}) \\ &\quad - rm\frac{1-kx}{k-m}(x^{r-1} - (\frac{1-mx}{k-m})^{r-1}) \\ &= x^{-r}m((r-1)(1-(a+1)^{-r})\log(a+1) - r\frac{a}{a+1}(1-(a+1)^{1-r})). \end{aligned}$$

The result then follows from Claim 12. \square

Claim 12. For all $r \in (0, 1)$ and $a > 0$ we have

$$(r-1)(1-(a+1)^{-r})\log(a+1) - r\frac{a}{a+1}(1-(a+1)^{1-r}) > 0.$$

Proof. Let

$$f(a) := (r-1)(1-(a+1)^{-r})\log(a+1) - r\frac{a}{a+1}(1-(a+1)^{1-r}).$$

Because $\lim_{a \rightarrow 0^+} f(a) = 0$, it suffices to prove that $f'(a) > 0$.

$$\begin{aligned} f'(a) &= (a+1)^{-r-1}(1-r - (a(1-r) + 1)((a+1)^{r-1} - r) - (1-r)r\log(a+1)) \\ &=: (a+1)^{-r-1}g(a). \end{aligned}$$

Because $\lim_{a \rightarrow 0^+} g(a) = 0$, it suffices to prove that $g'(a) > 0$.

$$g'(a) = \frac{ar(1-r)(1-(a+1)^{r-1})}{a+1} > 0.$$

\square

Now let us return to the proof of Proposition 8. The set of (m, x) where $m \in [1, k-1]$, $x \in (\frac{1}{k}, \frac{1}{m})$, and $H(P) = c$ can be parametrized as a curve $(m, x = x(m))$ for $m \in [1, m_c]$ for some constant m_c . Along the curve, $F(P)$ is continuous, and by Claim 11, is decreasing in m . Therefore $F(P)$ is maximized at $m = 1$. This finishes the proof. \square

Proof of Theorem 1 for $p > 1$. For a distribution $P = (p_1, \dots, p_k)$, we have

$$\mathcal{E}(P^{\frac{1}{p}}, P^{1-\frac{1}{p}}) = \frac{1}{k-1} \left(1 - \frac{1}{k} \left(\sum_i p_i^{\frac{1}{p}} \right) \left(\sum_i p_i^{1-\frac{1}{p}} \right) \right).$$

By Proposition 8, for fixed value of $\text{Ent}_\pi(P)$, the unique distribution of the form $(x, \frac{1-x}{k-1}, \dots, \frac{1-x}{k-1})$ with $x \in [\frac{1}{k}, 1]$ minimizes $\mathcal{E}(P^{\frac{1}{p}}, P^{1-\frac{1}{p}})$. Therefore for any non-zero non-negative f , we have

$$b_p \left(\frac{\text{Ent}_\pi(f)}{\mathbb{E}_\pi[f]} \right) \leq \frac{\mathcal{E}(f^{\frac{1}{p}}, f^{1-\frac{1}{p}})}{\mathbb{E}_\pi[f]}.$$

So p -NLSI holds with $\Phi_p = b_p^{-1}$. The statement about sharpness is obvious. \square

2.2. Non-linear 1-log-Sobolev inequality. We prove Theorem 1 for $p = 1$. Before proving the theorem we show the following.

Proposition 13. *Fix $0 \leq c \leq \log k$. Among all distributions P with $H(P) = c$, the distribution of form $P = (x, \frac{1-x}{k-1}, \dots, \frac{1-x}{k-1})$ with $x \in [\frac{1}{k}, 1]$ achieves maximum $\sum_i \log p_i$. Furthermore, up to permutation of the alphabet this is the unique minimum-achieving distribution.*

Proof. The result for $c \in \{0, \log k\}$ is obvious. In the following, assume that $0 < c < \log k$. Write $F(P) := \sum_i \log p_i$. The set $\{P : H(P) = c\}$ is compact, so the maximum value of $F(P)$ is achieved at some point $P = (p_1, \dots, p_k)$.

Step 0. If $p_i = 0$ for some i , then $F(P) = -\infty$. So $\min_{i \in [k]} p_i > 0$.

Step 1.

Claim 14. *If $u, v, w \in (0, 1)$ are all different, then*

$$\det \begin{pmatrix} 1 & \log u & \frac{1}{u} \\ 1 & \log v & \frac{1}{v} \\ 1 & \log w & \frac{1}{w} \end{pmatrix} \neq 0.$$

Proof of Claim. Suppose $\det = 0$. Then for some $a, b \in \mathbb{R}$, the equation $\frac{1}{x} + a \log x = b$ has at least three distinct solutions $x \in (0, 1)$. However, $\frac{\partial}{\partial x} (\frac{1}{x} + a \log x) = -\frac{1}{x^2} + \frac{a}{x}$ is smooth on $(0, 1)$, and takes zero at most once. So $\frac{1}{x} + a \log x$ takes each value at most once on $(0, 1)$. Contradiction. \square

By KKT conditions, the three vectors

$$\begin{aligned} \nabla F(P) &= \left(\frac{1}{p_i} \right)_{i \in [k]}, \\ \nabla H(P) &= (-1 - \log p_i)_{i \in [k]}, \\ \nabla \sum_{i \in [k]} p_i &= \mathbb{1} \end{aligned}$$

should be linear dependent. By Claim 14, there can be at most two different values of p_i 's.

So we can assume that $p_1 = \dots = p_m = x$, $p_{m+1} = \dots = p_k = \frac{1-mx}{k-m}$ for some $m \in [k-1]$, $x \in (\frac{1}{k}, \frac{1}{m})$.

Step 2. For P of the above form, we have

$$\begin{aligned} -H(P) &= mx \log x + (1 - mx) \log \frac{1 - mx}{k - m}, \\ F(P) &= m \log x + (k - m) \log \frac{1 - mx}{k - m}. \end{aligned}$$

We smoothly continue both functions so that m can take any real value in $[1, k - 1]$.

Claim 15. For $m \in (1, k - 1]$ and $x \in (\frac{1}{k}, \frac{1}{m})$, we have

$$-\frac{\partial}{\partial x} H(P) > 0$$

and

$$\frac{\partial}{\partial x} H(P) \frac{\partial}{\partial m} F(P) - \frac{\partial}{\partial m} H(P) \frac{\partial}{\partial x} F(P) > 0.$$

Proof of Claim. Let $f(x) = \log x - \log \frac{1 - mx}{k - m}$. Then we have

$$\begin{aligned} -\frac{\partial}{\partial x} H(P) &= mf(x) > 0, \\ -\frac{\partial}{\partial m} H(P) &= \frac{1 - kx}{k - m} + xf(x), \\ \frac{\partial}{\partial x} F(P) &= \frac{m(1 - kx)}{x(1 - mx)} \\ \frac{\partial}{\partial m} F(P) &= \frac{1 - kx}{1 - mx} + f(x). \end{aligned}$$

So

$$\begin{aligned} G(P) &:= \frac{\partial}{\partial x} H(P) \frac{\partial}{\partial m} F(P) - \frac{\partial}{\partial m} H(P) \frac{\partial}{\partial x} F(P) \\ &= m \left(\frac{(1 - kx)^2}{x(k - m)(1 - mx)} - f(x)^2 \right). \end{aligned}$$

Let $a = \frac{kx - 1}{1 - mx}$. Then $G(P) = m(\frac{a^2}{1 + a} - \log^2(a + 1))$. Because $a > 0$, we have $G(P) > 0$ by Lemma 16. \square

The set of (m, x) where $m \in [1, k - 1]$, $x \in (\frac{1}{k}, \frac{1}{m}]$, and $H(P) = c$ can be parametrized as a curve $(m, x = x(m))$ for $m \in [1, m_c]$ for some constant m_c . Along the curve, $F(P)$ is continuous, and by Claim 15, is decreasing in m . Therefore $F(P)$ is maximized at $m = 1$. This finishes the proof. \square

Lemma 16. For $a \in \mathbb{R}_{>-1}$, we have $\frac{a^2}{1 + a} \geq \log^2(a + 1)$. Equality holds only when $a = 0$.

Proof. We start from the well-known fact that $a \geq \log(a + 1)$ for $a \in \mathbb{R}_{>-1}$ (and equality holds only when $a = 0$). Let $f(a) = a(a + 2) - (2a + 2) \log(a + 1)$. We have $f(0) = 0$ and $f'(a) = 2(a - \log(a + 1)) \geq 0$ for $a \in \mathbb{R}_{>-1}$ (and equality holds only when $a = 0$). So f is negative on $(-1, 1)$ and positive on $(1, \infty)$.

Let $g(a) = \frac{a^2}{1 + a} - \log^2(a + 1)$. Clearly $g(0) = 0$. Because $g'(a) = \frac{f(a)}{(a + 1)^2}$, g is decreasing on $(-1, 1]$ and increasing on $[1, \infty)$. So $g(a) \geq 0$ for all $a \in \mathbb{R}_{>-1}$, and equality holds only when $a = 0$. \square

Proof of Theorem 1 for $p = 1$. For a distribution $P = (p_1, \dots, p_k)$, we have

$$\mathcal{E}(P, \log P) = \frac{1}{k-1} \sum_{i \in [k]} p_i \log p_i - \frac{1}{k(k-1)} \sum_{i \in [k]} \log p_i.$$

By Proposition 13, for fixed value of $\text{Ent}_\pi(P)$, the unique distribution of the form $(x, \frac{1-x}{k-1}, \dots, \frac{1-x}{k-1})$ with $x \in [\frac{1}{k}, 1]$ minimizes $\mathcal{E}(P, \log P)$. Therefore for any non-zero non-negative f , we have

$$b_1\left(\frac{\text{Ent}_\pi(f)}{\mathbb{E}_\pi[f]}\right) \leq \frac{\mathcal{E}(f, \log f)}{\mathbb{E}_\pi[f]}.$$

So 1-NLSI holds for $\Phi_1 = b_1^{-1}$. The statement about sharpness is obvious. \square

2.3. Input-restricted non-linear SDPI for Potts channels. In this section, we prove Theorem 3.

The subset of Potts channels corresponding to $\lambda \geq 0$ (i.e. ferromagnetic Potts channels) form a semigroup. For the semigroups, the optimal 1-NLSI is an “infinitesimal version” of the input-restricted non-linear SDPI. Consequently, by integrating the former we can get the latter (this is formalized in the first part of the proof below). Surprisingly, the result also extends beyond the semigroup to all of the Potts channels, namely we have the following.

Proposition 17. *Let $\lambda \in [-\frac{1}{k-1}, 1]$. Fix $0 \leq c \leq \log k$. Among all distributions P with $H(P) = c$, the distribution of form $P = (x, \frac{1-x}{k-1}, \dots, \frac{1-x}{k-1})$ with $x \in [\frac{1}{k}, 1]$ achieves minimum $H(\text{PC}_\lambda \circ P)$. Furthermore, when $\lambda \notin \{0, 1\}$, up to permutation of the alphabet this is the unique minimum-achieving distribution.*

Proof. The result for $\lambda \in \{0, 1\}$ is obvious. In the following assume that $\lambda \notin \{0, 1\}$. The result for $c \in \{0, \log k\}$ is obvious. In the following assume that $c \notin \{0, \log k\}$. The set $\{P : H(P) = c\}$ is compact, so the minimum value of $H(\text{PC}_\lambda \circ P)$ is achieved at some point $P = (p_1, \dots, p_k)$.

Step 0.

Claim 18. *Fix $a, b, d > 0$ and $c \in \mathbb{R}_{>-d} \setminus \{0\}$. Among all solutions $u, v, w \in [0, 1]$ with $u + v + w = a$ and $-u \log u - v \log v - w \log w = b$, the maximum of*

$$(cu + d) \log(cu + d) + (cv + d) \log(cv + d) + (cw + d) \log(cw + d)$$

is not achieved at a point where $0 = u < v < w$.

Proof. Suppose the maximum is achieved at such a point (u_0, v_0, w_0) where $0 = u_0 < v_0 < w_0$. Extend it to a curve $(u, v = v(u), w = w(u))$ on $u \in [0, \epsilon]$ for some $\epsilon > 0$, such that $u < v < w$ for all u , satisfying

$$(19) \quad u + v + w = a,$$

$$(20) \quad -u \log u - v \log v - w \log w = b,$$

and $v(0) = v_0, w(0) = w_0$.

We prove that

$$f(u) := (cu + d) \log(cu + d) + (cv + d) \log(cv + d) + (cw + d) \log(cw + d)$$

decreases as u approaches 0^+ for small enough u .

By taking derivative of (19) and (20), one can compute that

$$\begin{aligned} v'(u) &= \frac{\log w - \log u}{\log v - \log w}, \\ w'(u) &= \frac{\log u - \log v}{\log v - \log w}. \end{aligned}$$

Therefore

$$\begin{aligned} f'(u) &= c(\log(cu + d) + \log(cv + d)v'(u) + \log(cw + d)w'(u)) \\ &= c(\log(cu + d) + \log(cv + d)\frac{\log w - \log u}{\log v - \log w} + \log(cw + d)\frac{\log u - \log v}{\log v - \log w}). \end{aligned}$$

Because $0 < v_0 < w_0$, terms involving $\log u$ dominates the sum. The dominating term is

$$-c \log u \frac{\log(cv + d) - \log(cw + d)}{\log v - \log w} > 0.$$

Therefore the maximum of f is not achieved at $u = 0$. \square

By Claim 18, if $p_i = 0$ for some i , then there can be at most two different values of p_i 's.

Step 1.

Claim 19. *If $u, v, w \in (0, 1)$ are all different, then*

$$\det \begin{pmatrix} 1 & \log u & \log(\lambda u + \frac{1-\lambda}{k}) \\ 1 & \log v & \log(\lambda v + \frac{1-\lambda}{k}) \\ 1 & \log w & \log(\lambda w + \frac{1-\lambda}{k}) \end{pmatrix} \neq 0.$$

Proof of Claim. Suppose $\det = 0$. Then for some $a, b \in \mathbb{R}$, the equation

$$\log(\lambda x + \frac{1-\lambda}{k}) + a \log x = b$$

has at least three distinct solutions $x \in (0, 1)$. However,

$$\frac{\partial}{\partial x} (\log(\lambda x + \frac{1-\lambda}{k}) + a \log x) = \frac{\lambda}{\lambda x + \frac{1-\lambda}{k}} + \frac{a}{x}$$

is smooth on $(0, 1)$, and takes zero at most once. So

$$\log(\lambda x + \frac{1-\lambda}{k}) + a \log x$$

takes each value at most twice on $(0, 1)$. Contradiction. \square

By KKT conditions, the three vectors

$$\begin{aligned} \nabla H(\text{PC}_\lambda \circ P) &= (-\lambda \log(\lambda p_i + \frac{1-\lambda}{k}) - \lambda)_{i \in [k]}, \\ \nabla H(P) &= (-1 - \log p_i)_{i \in [k]}, \\ \nabla \sum_{i \in [k]} p_i &= \mathbb{1} \end{aligned}$$

should be linear dependent. By Claim 19, there can be at most two different values of p_i 's.

So we can assume that $p_1 = \dots = p_m = x$, $p_{m+1} = \dots = p_k = \frac{1-mx}{k-m}$ for some $m \in [k-1]$, $x \in (\frac{1}{k}, \frac{1}{m}]$.

Step 2. For P of the above form, we have

$$\begin{aligned} -H(P) &= mx \log x + (1 - mx) \log \frac{1 - mx}{k - m}, \\ -H(\text{PC}_\lambda \circ P) &= m\left(\lambda x + \frac{1 - \lambda}{k}\right) \log\left(\lambda x + \frac{1 - \lambda}{k}\right) \\ &\quad + (k - m)\left(\lambda \frac{1 - mx}{k - m} + \frac{1 - \lambda}{k}\right) \log\left(\lambda \frac{1 - mx}{k - m} + \frac{1 - \lambda}{k}\right). \end{aligned}$$

We smoothly continue both functions so that m can take any real value in $[1, k - 1]$.

Claim 20. For $m \in (1, k - 1]$ and $x \in (\frac{1}{k}, \frac{1}{m})$, we have

$$-\frac{\partial}{\partial x} H(P) > 0$$

and

$$\frac{\partial}{\partial m} H(P) \frac{\partial}{\partial x} H(\text{PC}_\lambda \circ P) - \frac{\partial}{\partial x} H(P) \frac{\partial}{\partial m} H(\text{PC}_\lambda \circ P) > 0.$$

Proof of Claim. Let

$$f(x) = \log x - \log \frac{1 - mx}{k - m}.$$

Then

$$\begin{aligned} -\frac{\partial}{\partial x} H(P) &= mf(x) > 0, \\ -\frac{\partial}{\partial m} H(P) &= \frac{1 - kx}{k - m} + xf(x), \\ -\frac{\partial}{\partial x} H(\text{PC}_\lambda \circ P) &= \lambda mf\left(\lambda x + \frac{1 - \lambda}{k}\right), \\ -\frac{\partial}{\partial m} H(\text{PC}_\lambda \circ P) &= \lambda \frac{1 - kx}{k - m} + \left(\lambda x + \frac{1 - \lambda}{k}\right) f\left(\lambda x + \frac{1 - \lambda}{k}\right), \end{aligned}$$

and

$$\begin{aligned} G(P) &:= \frac{\partial}{\partial m} H(P) \frac{\partial}{\partial x} H(\text{PC}_\lambda \circ P) - \frac{\partial}{\partial x} H(P) \frac{\partial}{\partial m} H(\text{PC}_\lambda \circ P) \\ &= \lambda m \frac{1 - kx}{k - m} \left(f\left(\lambda x + \frac{1 - \lambda}{k}\right) - f(x)\right) - mf(x) f\left(\lambda x + \frac{1 - \lambda}{k}\right) \frac{1 - \lambda}{k}. \end{aligned}$$

$$\frac{\partial}{\partial \lambda} \frac{G(P)}{m\lambda f(x) f\left(\lambda x + \frac{1 - \lambda}{k}\right)} = \frac{1}{k\lambda^2} + \frac{1 - kx}{k - m} \frac{\frac{\partial}{\partial \lambda} f\left(\lambda x + \frac{1 - \lambda}{k}\right)}{f\left(\lambda x + \frac{1 - \lambda}{k}\right)^2}.$$

Note that

(1)

$$\frac{G(P)}{m\lambda f(x) f\left(\lambda x + \frac{1 - \lambda}{k}\right)}$$

is continuous for $\lambda \in [-\frac{1}{k-1}, 1]$, and takes value 0 at $\lambda = 1$;

(2) $m\lambda f(x) f\left(\lambda x + \frac{1 - \lambda}{k}\right) \geq 0$ for $\lambda \in [-\frac{1}{k-1}, 1]$.

So we only need to prove that

$$\frac{\partial}{\partial \lambda} \frac{G(P)}{m\lambda f(x) f\left(\lambda x + \frac{1 - \lambda}{k}\right)} \leq 0,$$

i.e.,

$$f\left(\lambda x + \frac{1-\lambda}{k}\right)^2 \leq \frac{k\lambda^2(kx-1)}{k-m} \frac{\partial}{\partial \lambda} f\left(\lambda x + \frac{1-\lambda}{k}\right).$$

Let $y = \lambda x + \frac{1-\lambda}{k}$. Then the above inequality can be rewritten as

$$f(y)^2 \leq \frac{k\lambda^2(kx-1)}{k-m} \frac{\partial y}{\partial \lambda} \frac{\partial f(y)}{\partial y} = \frac{(ky-1)^2}{(k-m)y(1-my)}.$$

This is true by Lemma 16, applied to $a = \frac{ky-1}{1-my}$. Equality holds only when $y = \frac{1}{k}$, which cannot happen for $\lambda \neq 0$. \square

The set of (m, x) where $m \in [1, k-1]$, $x \in (\frac{1}{k}, \frac{1}{m}]$, and $H(P) = c$ can be parametrized as a curve $(m, x = x(m))$ for $m \in [1, m_c]$ for some constant m_c . Along the curve, $H(\text{PC}_\lambda \circ P)$ is continuous, and by Claim 20, is increasing in m . Therefore $H(\text{PC}_\lambda \circ P)$ is minimized at $m = 1$. This finishes the proof. \square

Proof of Theorem 3. Consider a Markov chain $U \rightarrow X \rightarrow Y$ where X has uniform distribution, and the channel $X \rightarrow Y$ is PC_λ . Because P_X and P_Y are both uniform, for any u , we have

$$\begin{aligned} D(P_{X|U=u}||P_X) &= \log k - H(P_{X|U=u}), \\ D(P_{Y|U=u}||P_Y) &= \log k - H(P_{Y|U=u}). \end{aligned}$$

So by Proposition 17 we get

$$D(P_{Y|U=u}||P_Y) \leq s_\lambda(D(P_{X|U=u}||P_X)).$$

Therefore

$$I(U; Y) = D(P_{Y|U}||P_Y|P_U) \leq \hat{s}_\lambda(D(P_{X|U}||P_X|P_U)) = \hat{s}_\lambda(I(U; X)).$$

Now we prove sharpness. Let $c \in [0, \log k]$. Choose $a, b \in [0, \log k]$ and $u \in [0, 1]$ such that $c = (1-u)a + ub$ and $\hat{s}_\lambda(c) = (1-u)s_\lambda(a) + us_\lambda(b)$. Choose $\rho, \tau \in [0, 1]$ such that $\text{Cap}(\text{PC}_\rho) = a$ and $\text{Cap}(\text{PC}_\tau) = b$, where Cap denotes channel capacity. Define random variable $U = (V, Z)$ such that $Z \sim \text{Ber}(u)$, and conditioned on $Z = 0$, $V \sim \text{PC}_\rho(X)$, and conditioned on $Z = 1$, $V \sim \text{PC}_\tau(X)$. One can check that

$$I(U; X) = (1-u)a + ub = c$$

and

$$I(U; Y) = (1-u)s_\lambda(a) + us_\lambda(b) = \hat{s}_\lambda(c).$$

\square

Let us discuss the relationship between Potts semigroup and ferromagnetic Potts channels. As discussed in the Introduction, ferromagnetic Potts channels are exactly the operators in the Potts semigroup, with $T_t = \text{PC}_{\exp(-\frac{k}{k-1}t)}$. Therefore 1-LSI for the Potts semigroup can be seen as infinitesimal SDPI for ferromagnetic Potts channels, and many results for the former can directly transfer to results for the latter.

We use Proposition 13 to give an alternative proof for Proposition 17 for ferromagnetic Potts channels.

Alternative proof of Proposition 17 for ferromagnetic Potts channels. Let P and Q be two distributions with $H(P) = H(Q) = c$, where P is of form $(x, \frac{1-x}{k-1}, \dots, \frac{1-x}{k-1})$ for some $x \in [\frac{1}{k}, 1]$, and Q is not of this form (up to permuting the alphabet). Define $P_t = T_t \circ P$ and $Q_t = T_t \circ Q$, where $(T_t)_{t \geq 0}$ is the Potts semigroup.

We prove that $H(P_t) < H(Q_t)$ for $t \in (0, \infty)$. Suppose this does not hold. Let $u = \inf\{t > 0 : H(P_t) \geq H(Q_t)\}$. Then we have $H(Q_u) = H(P_u)$ by continuity of semigroup. By Proposition 13, we have

$$\frac{\partial}{\partial t} \Big|_{t=u} H(Q_t) = \mathcal{E}(Q_u, \log Q_u) > \mathcal{E}(P_u, \log P_u) = \frac{\partial}{\partial t} \Big|_{t=u} H(P_t).$$

If $u = 0$, then for some $\epsilon > 0$, $H(Q_t) > H(P_t)$ for $t \in (0, \epsilon)$. If $u > 0$, then for some $\epsilon > 0$, $H(Q_t) < H(P_t)$ for $t \in (u - \epsilon, u)$. Both cases lead to contradiction with definition of u . So $H(P_t) < H(Q_t)$ for $t \in (0, \infty)$. This completes the proof of the result for $\lambda > 0$. \square

2.4. Behavior for $k \rightarrow \infty$. When should one use p -NLSI instead of p -LSI? To get some insights, we consider the case of $k \rightarrow \infty$. First, we prove Proposition 2 that $\alpha_1 = 1 + \frac{1+o(1)}{\log k}$.

Proof of Proposition 2. Lower bound. By Theorem 1, we need to show that for all $x \in (\frac{1}{k}, 1]$, we have

$$\frac{k}{k-1} \left(1 + \frac{1}{\log k}\right) \leq \frac{\xi_1(x)}{\psi(x)}.$$

Noting that

$$\xi_1(x) = \frac{k}{k-1} \left(\frac{1}{k} (-\log x - (k-1) \log \frac{1-x}{k-1}) - \log k + \psi(x) \right),$$

it suffices to prove that

$$f(x) := \frac{\log k}{k} (-\log x - (k-1) \log \frac{1-x}{k-1}) - \log^2 k - \psi(x) \geq 0.$$

We have $f(\frac{1}{k}) = 0$. So it suffices to prove that $f'(x) \geq 0$ for $x \in [\frac{1}{k}, 1]$.

$$f'(x) = \frac{\log k}{k} \left(-\frac{1}{x} + \frac{k-1}{1-x} \right) - \left(\log x - \log \frac{1-x}{k-1} \right).$$

We smoothly continue this function to $\{(k, x) \in \mathbb{R}^2 : k \geq 3, x \in [\frac{1}{k}, 1]\}$ and prove that it is non-negative in this region.

$$\begin{aligned} \frac{\partial}{\partial k} f'(x) &= \frac{1 - \log k}{k^2} \left(-\frac{1}{x} + \frac{k-1}{1-x} \right) + \frac{\log k}{k} \frac{1}{1-x} - \frac{1}{k-1} \\ &= \frac{(k-1) \log k + k(kx^2 - x - 1) + 1}{k^2(k-1)x(1-x)}. \end{aligned}$$

The numerator is a quadratic function in x , and for fixed k , it is minimized at $x = \frac{1}{k}$, leading to

$$\frac{\partial}{\partial k} f'(x) \geq \frac{(k-1) \log k - k + 1}{k^2(k-1)x(1-x)} = \frac{\log k - 1}{k^2 x(1-x)} \geq 0.$$

So we only need to prove $f'(x) \geq 0$ for minimum k , i.e., $k = \max\{3, \frac{1}{x}\}$. When $k = \frac{1}{x}$, one can verify that $f'(x) = 0$. So the only remaining case is $k = 3$. For

$k = 3$, we prove that f is convex in x , i.e., $f''(x) \geq 0$ for $x \in [0, 1]$.

$$\begin{aligned} f''(x) &= \frac{\log k}{k} \left(\frac{1}{x^2} + \frac{k-1}{(1-x)^2} \right) - \left(\frac{1}{x} + \frac{1}{1-x} \right) \\ &= \frac{\log k((1-x)^2 + (k-1)x^2) - kx(1-x)}{kx^2(1-x)^2} \\ &= \frac{(k \log k + k)x^2 - (k + 2 \log k)x + \log k}{kx^2(1-x)^2}. \end{aligned}$$

The numerator is a quadratic function in x , and its discriminant is

$$(k + 2 \log k)^2 - 4(k \log k + k) \log k = k^2 - 4(k-1) \log^2 k.$$

When $k = 3$, the above value is < 0 . So $f''(x) \geq 0$ for $k = 3$ and $x \in [0, 1]$. This finishes the proof of the lower bound.

Upper bound. For the upper bound, we need find $x \in (\frac{1}{k}, 1]$ such that $\frac{f(x)}{\psi(x)} = o(1)$. Because the upper bound to prove is asymptotic, we assume that k is large enough. Take $x = \frac{2}{\log k}$. Then we have

$$\begin{aligned} \psi(x) &= \log k + \frac{2}{\log k} \log \frac{2}{\log k} + \left(1 - \frac{2}{\log k}\right) \log \frac{1 - \frac{2}{\log k}}{k-1} \\ &= \log k - \left(1 - \frac{2}{\log k}\right) \log(k-1) + o(1) \\ &= 2 + o(1) \end{aligned}$$

and

$$\begin{aligned} f(x) &= \frac{\log k}{k} \left(-\log \frac{2}{\log k} - (k-1) \log \frac{1 - \frac{2}{\log k}}{k-1} \right) - \log^2 k - \psi(x) \\ &= \frac{\log k}{k} (k-1) (\log(k-1) - \log(1 - \frac{2}{\log k})) - \log^2 k - 2 + o(1) \\ &= \log k \cdot \left(1 + O\left(\frac{1}{k}\right)\right) \cdot (\log k + O\left(\frac{1}{k}\right) + \frac{2}{\log k} + O\left(\frac{1}{\log^2 k}\right)) - \log^2 k - 2 + o(1) \\ &= o(1). \end{aligned}$$

So $\frac{f(x)}{\psi(x)} = o(1)$. □

Remark 21. Numerical computation suggests $\frac{f(x)}{\psi(x)}$ is minimized at a point $x = \frac{2+o(1)}{\log k}$. This guides our proof of the upper bound in Proposition 2, but we have not attempted to prove this fact.

To understand the case $p > 1$, let us denote convexification of b_p as \check{b}_p . Then NLSI lower bound, assuming $\mathbb{E}[f] = 1$, gives

$$\mathcal{E}(f^{\frac{1}{p}}, f^{1-\frac{1}{p}}) \geq \check{b}_p(\text{Ent}(f)).$$

We see that this improves upon $\alpha_p \cdot \text{Ent}(f)$ the more the larger the entropy. In particular, the maximum improvement happens when $\text{Ent}(f) = \log k$. That is we have for $p > 1$

$$\alpha_p \leq \frac{\check{b}_p(x)}{x} \leq \frac{1}{\log k}.$$

Together with (4) and (5), we get $\alpha_p = \Theta(\frac{1}{\log k})$ as $k \rightarrow \infty$. Numerical computation suggests that $\alpha_p = \frac{1+o(1)}{\log k}$.

At the same time, the improvement given by the 1-NLSI (over 1-LSI) is much stronger, since $b_1(\log k) = \infty$. *To summarize*, the p -NLSI should be preferred for $p = 1$ or for cases where k is small and entropy is large (i.e. functions are highly spiky).

Next, we consider SDPIs and η_{KL} . First, we show that for a fixed $\lambda \geq 0$ we have

$$\eta_{\text{KL}}(\text{PC}_\lambda, \pi) = \lambda - \Theta\left(\frac{1}{\log k}\right).$$

Indeed, the upper bound is given by (13). For the lower bound we have

$$\begin{aligned} & \eta_{\text{KL}}(\text{PC}_\lambda, \pi) \\ & \geq \eta_{\min} := \frac{\psi(\lambda + \frac{1-\lambda}{k})}{\psi(1)} \\ & = \frac{\log k + (\lambda + \frac{1-\lambda}{k}) \log(\lambda + \frac{1-\lambda}{k}) + (1 - (\lambda + \frac{1-\lambda}{k})) \log \frac{1 - (\lambda + \frac{1-\lambda}{k})}{k-1}}{\log k} \\ & = \lambda + \frac{\lambda \log \lambda + (1 - \lambda) \log(1 - \lambda) + o(1)}{\log k}. \end{aligned}$$

On the other hand,

$$\eta_{\text{KL}}(\text{PC}_\lambda, \pi) \leq \eta_{\text{KL}}(\text{PC}_\lambda) \leq \eta_{\text{TV}}(\text{PC}_\lambda) = \lambda,$$

where η_{TV} is the contraction coefficient for the total variation distance (Dobrushin coefficient, see [CKZ98, PW17]).

Notice also that for \hat{s}_λ we have generally $\eta_{\min} \leq \frac{\hat{s}_\lambda(x)}{x} \leq \eta_{\text{KL}}(\text{PC}_\lambda, \pi)$. Therefore we have shown that

$$\lim_{k \rightarrow \infty} \frac{\hat{s}_\lambda(x)}{x} = \lim_{k \rightarrow \infty} \eta_{\text{KL}}(\text{PC}_\lambda, \pi) = \lim_{k \rightarrow \infty} \eta_{\text{KL}}(\text{PC}_\lambda) = \eta_{\text{TV}}(\text{PC}_\lambda) = \lambda.$$

The estimates of information quantities using the more sophisticated tools get improvement over simplistic coupling of at most multiplicative order $(1 + \Theta(\frac{1}{\log k}))$.

Note, however, if λ changes with k (e.g. $\lambda = -\frac{1}{k-1}$), then the improvement over η_{TV} can be as large as a multiplicative factor of $(1 + o(1)) \log k$, as shown in Proposition 32.

3. PRODUCT SPACES

In this section we study extensions of p -NLSIs and SDPIs to the product semi-group $(T_t^{\otimes n})_{t \geq 0}$ on the product space $[k]^n$ (and product channels $\text{PC}_\lambda^{\otimes n}$). The general property of tensorization of p -NLSI was established in [PS19, Theorem 1], and thus we only need to concavify functions Φ_p in (3). Similarly, we can show that (non-linear) strong data processing inequalities tensorize if one concavifies function $s(\cdot)$ in (12).

After showing these extensions to product spaces, we proceed to discussing implications of p -NLSI on speed of convergence to equilibrium in terms of $\text{Ent}(T_t^{\otimes n} \nu)$ and on edge-isoperimetric inequalities.

3.1. Tensorization.

Proposition 22. Fix $p \geq 1$. Recall b_p defined in Theorem 1. Let \check{b}_p be the convex envelope of b_p . Then p -LSI holds for the product semigroup $(T_t^{\otimes n})_{t \geq 0}$ with

$$\Phi_{n,p}(x) = n\check{b}_p^{-1}\left(\frac{x}{n}\right).$$

Proof. By Theorem 1 and [PS19, Theorem 1]. □

As we show below $\check{b}_p \neq b_p$ (Proposition 25).

For non-linear SDPI, we first prove a general tensorization result.

Proposition 23. Fix a probability kernel $P_{Y|X} : \mathcal{X} \rightarrow \mathcal{Y}$ and a distribution P_X on \mathcal{X} .

(1) Suppose for some non-decreasing function $s : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ we have

$$(21) \quad D(Q_Y || P_Y) \leq s(D(Q_X || P_X))$$

for all distribution Q_X on \mathcal{X} with $0 < D(Q_X || P_X) < \infty$. Then for all distribution Q_{X^n} on \mathcal{X}^n with $0 < D(Q_{X^n} || P_X^{\otimes n}) < \infty$, we have

$$(22) \quad D(Q_{Y^n} || P_Y^{\otimes n}) \leq n\hat{s}\left(\frac{1}{n}D(Q_{X^n} || P_X^{\otimes n})\right),$$

where \hat{s} is the concave envelope of s , and $Q_{Y^n} = P_{Y|X}^{\otimes n} \circ Q_{X^n}$.

(2) Suppose for some non-decreasing concave function $\hat{s} : [0, \log |\mathcal{X}|] \rightarrow \mathbb{R}_{\geq 0}$ we have

$$(23) \quad I(U; Y) \leq \hat{s}(I(U; X))$$

for all Markov chains

$$U \rightarrow X \rightarrow Y$$

where the distribution of X is P_X . Then for all Markov chains

$$U \rightarrow X^n \rightarrow Y^n$$

where the distribution of X is $P_X^{\otimes n}$, we have

$$(24) \quad I(U; Y^n) \leq n\hat{s}\left(\frac{1}{n}I(U; X^n)\right).$$

We have separate statements for non-linear SDPI defined via KL divergence and via mutual information, because they are not equivalent in general. It is not hard to show that if KL divergence type non-linear SDPI (Inequality (21)) holds for some function s , then mutual information type non-linear SDPI (Inequality (23)) holds for \hat{s} . However, it is not clear what is the best possible KL divergence type non-linear SDPI one can get starting from mutual information type non-linear SDPI. (Note the domain of function s would become larger during the translation.)

Proof of Proposition 23. Proof of (1). Perform induction on n . The base case $n = 1$ is trivial. Now consider $n \geq 2$. We have

$$\begin{aligned}
& D(Q_{Y^n} \| P_Y^{\otimes n}) \\
&= D(Q_{Y^{n-1}} \| P_Y^{\otimes(n-1)}) + D(Q_{Y_n | Y^{n-1}} \| P_Y | Q_{Y^{n-1}}) \\
&\leq D(Q_{Y^{n-1}} \| P_Y^{\otimes(n-1)}) + D(Q_{Y_n | X^{n-1}} \| P_Y | Q_{X^{n-1}}) \\
&\leq (n-1) \hat{s} \left(\frac{1}{n-1} D(Q_{X^{n-1}} \| P_X^{\otimes(n-1)}) \right) + s(D(Q_{X_n | X^{n-1}} \| P_X | Q_{X^{n-1}})) \\
&\leq n \hat{s} \left(\frac{1}{n} D(Q_{X^{n-1}} \| P_X^{\otimes(n-1)}) \right) + \frac{1}{n} D(Q_{X_n | X^{n-1}} \| P_X | Q_{X^{n-1}}) \\
&= n \hat{s} \left(\frac{1}{n} D(Q_{X^n} \| P_X^{\otimes n}) \right).
\end{aligned}$$

First step is by chain rule. Second step is because we have a Markov chain

$$Y^{n-1} - X^{n-1} - Y_n,$$

and conditioning on more information does not decrease conditioned divergence. Third step is by induction hypothesis. Fourth step is by concavity. Fifth step is by chain rule.

Proof of (2). Perform induction on n . The base case $n = 1$ is trivial. Now consider $n \geq 2$. We have

$$\begin{aligned}
I(U; Y^n) &= I(U; Y^{n-1}) + I(U; Y_n | Y^{n-1}) \\
&= I(U; Y^{n-1}) + I(U, Y^{n-1}; Y_n) \\
&\leq I(U; Y^{n-1}) + I(U, X^{n-1}; Y_n) \\
&= I(U; Y^{n-1}) + I(U; Y_n | X^{n-1}) \\
&\leq (n-1) \hat{s} \left(\frac{1}{n-1} I(U; X^{n-1}) \right) + \hat{s}(I(U; X_n | X^{n-1})) \\
&\leq n \hat{s} \left(\frac{1}{n} I(U; X^{n-1}) \right) + \frac{1}{n} I(U; X_n | X^{n-1}) \\
&= n \hat{s} \left(\frac{1}{n} I(U; X^n) \right).
\end{aligned}$$

First step is by chain rule. Second step is by chain rule, and that Y_n is independent with Y^{n-1} . Third step is by data processing inequality. Fourth step is by chain rule, and that Y_n is independent with X^{n-1} . Fifth step is by induction hypothesis. Sixth step is by concavity. Seventh step is by chain rule. \square

Corollary 24. Recall function s_λ defined in Theorem 3. Let Q_{X^n} be a distribution on $[k]^n$ and $Q_{Y^n} = \text{PC}_\lambda^{\otimes n} \circ Q_{X^n}$. Then we have

$$(25) \quad \frac{1}{n} H(Y^n) \geq \log k - \hat{s}_\lambda \left(\log k - \frac{1}{n} H(X^n) \right).$$

Furthermore, for every $c \in [0, \log k]$, there exist distributions X^n with $H(X^n) = (c + o(1))n$ such that $\frac{1}{n} H(Y^n) = \log k - \hat{s}_\lambda(\log k - c) + o(1)$.

Proof. Inequality (25) follows from Proposition 23 and that

$$D(Q_{X^n} \| \pi^{\otimes n}) = n \log k - H(Q_{X^n}).$$

For the second part, choose $a, b \in [0, \log k]$ and $u \in [0, 1]$ such that $c = (1 - u)a + ub$ and $\hat{s}_\lambda(\log k - c) = (1 - u)s_\lambda(\log k - a) + us_\lambda(\log k - b)$. Such a, b, u exist because \hat{s}_λ is the concave envelope of s_λ .

Let Q_A (resp. Q_B) be the unique distribution on $[k]$ of form $(x, \frac{1-x}{k-1}, \dots, \frac{1-x}{k-1})$ with $x \in [\frac{1}{k}, 1]$ and entropy a (resp. entropy b). Now let Q_{X^n} be the distribution $Q_A \times \dots \times Q_A \times Q_B \times \dots \times Q_B$, where Q_A appears $\lfloor (1 - u)n \rfloor$ times and Q_B appears $\lceil un \rceil$ times. It is easy to see that this distribution satisfies the required properties. \square

3.2. Linear piece. In Proposition 22 and Theorem 3, we make use of convexification of b_p and concavification of s_λ . When $k = 2$, we have $\check{b}_p = b_p$ and $\hat{s}_\lambda = s_\lambda$ (the latter fact is known as Mrs. Gerber's Lemma [WZ73]). However, for $k \geq 3$, the situation is vastly different.

Proposition 25. *Recall function $b_p : [0, \log k] \rightarrow \mathbb{R}$ defined in Theorem 1 and $s_\lambda : [0, \log k] \rightarrow \mathbb{R}$ defined in Theorem 3.*

- (1) *For all $k \geq 3$ and $p \geq 1$, b_p is not convex near 0.*
- (2) *For all $k \geq 3$, $\lambda \in [-\frac{1}{k-1}, 0) \cup (0, 1)$, s_λ not concave near 0.*

The proof is deferred to Appendix C. Proposition 25 implies that there is a linear piece near origin in the graph of \check{b}_p , $\hat{\Phi}_p$ and \hat{s}_λ .

This implies a curious new property distinguishing Potts semigroup with $k \geq 3$ from its binary cousin and from the Ornstein-Uhlenbeck semigroup. Both of the latter have their p -NLSI and SDPI strictly non-linear, which translates into the following fact: among all initial densities ν_0 with a given entropy $\text{Ent}(\nu_0)$ a simple product distributions simulateneously maximizes $\text{Ent}(T_t^{\otimes n} \nu_0)$ for all t . Stated differently we have (this is known as Mrs. Gerber's Lemma) when $k = 2$:

$$(26) \quad D(\text{PC}_\lambda \circ P_{X^n} || \pi^{\otimes n}) \leq D(\text{PC}_\lambda \circ \text{Ber}^{\otimes n}(p) || \pi^{\otimes n}),$$

where π is the uniform distribution on $[k]^n$, and $\text{Ber}^{\otimes n}(p)$ is an iid distribution on $[k]^n$ with $p \in [0, 1/2]$ solving $D(\text{Ber}(p) || \pi) = \frac{1}{n} D(\text{PC}_\lambda \circ P_{X^n} || \pi^{\otimes n})$. *That is, the slowest to relax to equilibrium is the product distribution.* For the Ornstein-Uhlenbeck a similar statement holds with $\text{Ber}(p)$ replaced by the $\mathcal{N}(0, \sigma^2 I_n)$.

This nice extremal property of product distributions is no longer true for $k \geq 3$ Potts semigroups, because s_λ is not concave, and the value of \hat{s}_λ at a point may be a mixture of two values of s_λ . More precisely, instead of (26), we have for every $\lambda \in [-\frac{1}{k-1}, 1]$ and every $c \in \mathbb{R}_{\geq 0}$, there exist two iid distributions P_1, P_2 on $[k]^n$ and $t \in [0, 1]$ satisfying

$$(1 - t)D(P_1 || \pi^{\otimes n}) + tD(P_2 || \pi^{\otimes n}) = c,$$

such that for every distribution P_{X^n} on $[k]^n$ with $D(P_{X^n} || \pi^{\otimes n}) = c$, we have

$$D(\text{PC}_\lambda \circ P_{X^n} || \pi^{\otimes n}) \leq (1 - t)D(\text{PC}_\lambda \circ P_1 || \pi^{\otimes n}) + tD(\text{PC}_\lambda \circ P_2 || \pi^{\otimes n}).$$

Note here P_1, P_2 and t all depend on c and λ , and thus there is no universal distribution that is the slowest to converge to equilibrium.

Let us discuss some general implications of non-convexity of b_p and non-concavity of s_λ near 0.

Let K be a Markov kernel with stationary distribution π . Consider the tightest possible non-linear p -NLSI given by

$$b_p(x) := \inf_{\substack{f: \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}, \\ \mathbb{E}_\pi f = 1, \text{Ent}_\pi(f) = x}} \mathcal{E}(f^{\frac{1}{p}}, f^{1-\frac{1}{p}}).$$

The p -log-Sobolev constant is

$$\alpha_p := \inf_{x > 0} \frac{b_p(x)}{x} = \inf_{f: \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}, \text{Ent}_\pi(f) > 0} \frac{\mathcal{E}(f^{\frac{1}{p}}, f^{1-\frac{1}{p}})}{\text{Ent}_\pi(f)}.$$

We also define the spectral gap

$$\lambda := \inf_{f: \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}, \text{Var}(f) > 0} \frac{\mathcal{E}(f, f)}{\text{Var}(f)},$$

where $\text{Var}(f) = \mathbb{E}_\pi(f - \mathbb{E}_\pi f)^2$. For any $p > 1$, we have

$$(27) \quad \frac{p^2}{2(p-1)} \alpha_p \leq \lambda.$$

The case $p = 2$ is proved in Diaconis and Saloff-Coste [DSC96], and the general case is proved in Mossel et al. [MOS13]. Their proof in fact implies a stronger inequality.

Lemma 26.

$$\limsup_{x \rightarrow 0^+} \frac{b_p(x)}{x} \leq \frac{2(p-1)}{p^2} \lambda.$$

In particular, when b_p is strictly concave near 0, we have

$$\alpha_p < \limsup_{x \rightarrow 0^+} \frac{b_p(x)}{x} \leq \frac{2(p-1)}{p^2} \lambda,$$

and (27) is strict.

Proof. Take any $g : \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}$ with $\text{Var}(g) > 0$. Define $f_\epsilon = 1 + \epsilon g$. As $\epsilon \rightarrow 0$, we have

$$\begin{aligned} \mathcal{E}(f_\epsilon^{\frac{1}{p}}, f_\epsilon^{1-\frac{1}{p}}) &= \epsilon^2 \frac{1}{p} \left(1 - \frac{1}{p}\right) \mathcal{E}(g, g) + o(\epsilon^2), \\ \text{Ent}_\pi(f_\epsilon) &= \frac{1}{2} \epsilon^2 \text{Var}(g) + o(\epsilon^2). \end{aligned}$$

Because $\text{Ent}_\pi(f_\epsilon) \rightarrow 0$ continuously as $\epsilon \rightarrow 0$, we have

$$\limsup_{x \rightarrow 0^+} \frac{b_p(x)}{x} \leq \lim_{\epsilon \rightarrow 0} \frac{\mathcal{E}(f_\epsilon^{\frac{1}{p}}, f_\epsilon^{1-\frac{1}{p}})}{\text{Ent}_\pi(f_\epsilon)} = \frac{2(p-1)}{p^2} \frac{\mathcal{E}(g, g)}{\text{Var}(g)}.$$

Lemma then follows because g is arbitrary. \square

Roughly speaking, existence of a ‘‘linear piece’’ near 0 in \check{b}_p implies that (27) is strict. For the Potts semigroup with $k \geq 3$, b_p is strictly concave near 0 by proof of Proposition 25. So (27) is strict for the Potts semigroup.

The story for non-linear SDPI is very similar. Let W be any channel and Q be any input distribution. Consider the tightest possible non-linear SDPI given by

$$s(x) := \sup_{P: D(P||Q) = x} D(PW||QW).$$

The input-restricted KL divergence contraction coefficient is

$$\eta_{\text{KL}}(W, Q) := \sup_{x>0} \frac{s(x)}{x} = \sup_{P:0<D(P||Q)<\infty} \frac{D(PW||QW)}{D(P||Q)}.$$

We also consider the input-restricted χ^2 -divergence contraction coefficient

$$\eta_{\chi^2}(W, Q) := \sup_{P:0<\chi^2(P||Q)<\infty} \frac{\chi^2(PW||QW)}{\chi^2(P||Q)}.$$

It is known (Ahlsvede and Gács [AG76]) that

$$(28) \quad \eta_{\text{KL}}(W, Q) \geq \eta_{\chi^2}(W, Q).$$

Similarly to the p -NLSI case, the proof of (28) implies a stronger inequality.

Lemma 27.

$$\liminf_{x \rightarrow 0^+} \frac{s(x)}{x} \geq \eta_{\chi^2}(W, Q).$$

In particular, when s is strictly convex near 0, we have

$$\eta_{\text{KL}}(W, Q) > \liminf_{x \rightarrow 0^+} \frac{s(x)}{x} \geq \eta_{\chi^2}(W, Q).$$

and (28) is strict.

Proof. Fix any distribution P with $0 < \chi^2(P||Q) < \infty$. Proof of [PW17, Theorem 2] constructs a sequence of distributions P_ϵ satisfying

$$\begin{aligned} D(P_\epsilon||Q) &= \epsilon^2 \chi^2(P||Q) + o(\epsilon^2), \\ D(P_\epsilon W||QW) &= \epsilon^2 \chi^2(PW||QW) + o(\epsilon^2), \end{aligned}$$

and $D(P_\epsilon||Q) \rightarrow 0$ continuously as $\epsilon \rightarrow 0$. Therefore

$$\liminf_{x \rightarrow 0^+} \frac{s(x)}{x} \geq \lim_{\epsilon \rightarrow 0} \frac{D(P_\epsilon W||QW)}{D(P_\epsilon||Q)} = \frac{\chi^2(PW||QW)}{\chi^2(P||Q)}.$$

Lemma follows because P is arbitrary. \square

Roughly speaking, existence of a “linear piece” near 0 in \hat{s} implies that (28) is strict. For Potts channels PC_λ with $\lambda \in [-\frac{1}{k-1}, 0) \cup (0, 1)$ and $k \geq 3$, s_λ is strictly convex near 0 by proof of Proposition 25. So (28) is strict for Potts channels.

3.3. Edge isoperimetric inequalities. As a toy application of the NLSIs for the product spaces, we derive an edge isoperimetric inequality for K_k^n , the graph whose vertex set is $[k]^n$, and edges connect vertex pairs with Hamming distance one. Given a graph $G = (V, E)$, edge isoperimetric inequalities solve the following combinatorial optimization problem:

$$\Psi_G(N) = \min\{|E(S, S^c)| : |S| = N\},$$

where $|E(S, S^c)| = \#\{e \in E : |e \cap S| = 1\}$. For K_k^n , the edge isoperimetric problem has been completely solved [Har64, Lin64, Ber67, Har76]. Specifically, Lindsey [Lin64] showed that the optimal S minimizing $|E(S, S^c)|$ for a fixed $|S|$ consists of largest elements in $[k]^n$ under a lexicographical order. In particular, we have

$$\Psi_{K_k^n}(k^m) = (n - m)(k - 1)k^m.$$

This was obtained by an explicit combinatorial argument (via a form of shifting/compression). What estimates obtained via LSIs and NLSIs?

Let $f = \mathbb{1}_S$ be the indicator function of a set S . Then for any $p > 1$ we have

$$\frac{\mathcal{E}(f^p, f^{1-p})}{\mathbb{E}_\pi[f]} = \frac{1}{k-1} \frac{|E(S, S^c)|}{|S|} \quad \text{and} \quad \frac{\text{Ent}(f)}{\mathbb{E}_\pi[f]} = \log \frac{k^n}{|S|}.$$

If we relate these two ratios via the 2-LSI (note that from (4), of all $p > 1$ the $p = 2$ gives the best result here) and by using the known value of α_2 from (5) we get

$$(29) \quad \Psi_{K_k^n}(k^m) \geq k^m(n-m)(k-2) \frac{\log k}{\log(k-1)}.$$

Clearly the coefficient in front of $(n-m)k^m$ here is not tight.

The p -NLSI allows us to perform a better comparison. First, again via (4) we get the best inequality for $p = 2$, which results in

$$(30) \quad \Psi_{K_k^n}(k^m) \geq (k-1)k^m n \check{b}_2\left(\frac{n-m}{n} \log k\right).$$

We know that the function \check{b}_2 is continuous with $\check{b}_2(\log k) = b_2(\log k) = 1$ (from (9)). Thus, for any $m = o(n)$ and $n \rightarrow \infty$ we get that (30) implies

$$\Psi_{K_k^n}(k^m) \geq (k-1)k^m(n-m)(1+o(1)),$$

which is tight in this regime. (However, from (9) we can also find that $\check{b}'_2(1) = \infty$ and thus, even when $m = o(n)$ the right-hand side of the above inequality is $(k-1)k^m(n-\omega(m))$, implying the behavior in terms of m is not optimal.)

4. NON-RECONSTRUCTION FOR BROADCAST MODELS ON TREES

In this section we prove non-reconstruction results for a general class of broadcast models on trees, using input-restricted KL divergence SDPI.

Fix a channel $M : [k] \rightarrow [k]$ with an invariant distribution q^* , i.e., $q^*M = q^*$. Let M^\vee denote the reverse channel, i.e., any channel that satisfies $q_j^*M_{j,i}^\vee = q_i^*M_{i,j}$ for all $i, j \in [k]$. Consider a (possibly infinite) tree T with a marked root ρ . For each vertex $v \in T$, we generate a spin $\sigma_v \in [k]$ according to the following rules:

- (1) $\mathbb{P}(\sigma_\rho = i) = q_i^*$.
- (2) If u is the parent of v , then $\mathbb{P}(\sigma_v = j | \sigma_u = i) = M_{i,j}$.

Let L_h denote the set of vertices of distance h to ρ . We say the model has non-reconstruction if and only if

$$\lim_{h \rightarrow \infty} I(\sigma_\rho; \sigma_{L_h}) = 0.$$

For any vertex u , let $c(u)$ denote the set of children of u .

Finally, recall the definition of the branching number $\text{br}(T)$ of a tree T by Lyons [Lyo90].

Definition 28 (Branching number). Define a flow to be a function $f : V(T) \rightarrow \mathbb{R}_{\geq 0}$ such that for every vertex u , we have

$$f_u = \sum_{v \in c(u)} f_v.$$

Define $\text{br}(T)$ to be the sup of all numbers λ such that there exists a flow f with $f_\rho > 0$, and $f_u \leq \lambda^{-d(u,\rho)}$ for all vertices u , where $d(u, \rho)$ is the distance between u and ρ .

Recall Theorem 5 states that the model has non-reconstruction when

$$\eta_{\text{KL}}(M^\vee, q^*) \text{br}(T) < 1.$$

Now we prove the theorem.

Proof of Theorem 5. For any vertex u , define $L_{u,h}$ to be the set of descendants of u that have distance h to ρ . Define

$$a_u = H(q^*)^{-1} \eta_{\text{KL}}(M^\vee, q^*)^{d(u,\rho)} \lim_{h \rightarrow \infty} I(\sigma_u; \sigma_{L_{u,h}}).$$

By data processing inequality, $I(\sigma_u; \sigma_{L_{u,h}})$ is non-increasing for $h \geq d(u, \rho)$, so the limit exists.

For any $v \in c(u)$, consider the reverse Markov chain

$$\sigma_{L_{v,h}} \rightarrow \sigma_v \rightarrow \sigma_u.$$

Because q^* is an invariant distribution, the distributions of σ_v and σ_u are both q^* . By SDPI, we have

$$I(\sigma_u; \sigma_{L_{v,h}}) \leq \eta_{\text{KL}}(M^\vee, q^*) I(\sigma_v; \sigma_{L_{v,h}}).$$

Because $(\sigma_{L_{v,h}})_{v \in c(u)}$ are independent conditioned on σ_u , we have

$$I(\sigma_u; \sigma_{L_{u,h}}) \leq \sum_{v \in c(u)} I(\sigma_u; \sigma_{L_{v,h}}).$$

Combine the two inequalities and let $h \rightarrow \infty$. We get that

$$a_u \leq \sum_{v \in c(u)} a_v.$$

Clearly,

$$a_u \leq \eta_{\text{KL}}(M^\vee, q^*)^{d(u,\rho)}$$

for all vertices u . However, a is not quite a flow yet. We define a flow b from a . For a vertex u , let $u_0 = \rho, \dots, u_d = u$ be the shortest path from ρ to u . Define

$$b_u = a_u \prod_{0 \leq j \leq d-1} \frac{a_{u_j}}{\sum_{v \in c(a_{u_j})} a_v}.$$

(If for some j , we have $\sum_{v \in c(a_{u_j})} a_v = 0$, then let $b_u = 0$.) It is not hard to check that

$$b_u = \sum_{v \in c(u)} b_v,$$

and that

$$b_u \leq a_u \leq \eta_{\text{KL}}(M^\vee, q^*)^{d(u,\rho)}.$$

By definition of branching number, we must have $b_\rho = 0$. This means

$$\lim_{h \rightarrow \infty} I(\sigma_\rho; \sigma_{L_h}) = 0,$$

and non-reconstruction holds. \square

Remark 29. In the definition of the model, it is not necessary to require σ_ρ to have distribution q^* . If we let $\sigma_{L_h}^i$ denote the leaf colors conditioned on $\sigma_\rho = i$, then Theorem 6 implies that when $\eta_{\text{KL}}(M^\vee, q^*) \text{br}(T) < 1$, we have

$$\lim_{h \rightarrow \infty} \text{TV}(\sigma_{L_h}^i, \sigma_{L_h}^j) = 0$$

for $i \neq j$ with $q_i^*, q_j^* > 0$.

Theorem 5 directly implies non-reconstruction results for Galton-Watson trees.

Corollary 30. *Let T be a Galton-Watson tree with expected offspring d . If*

$$\eta_{\text{KL}}(M^\vee, q^*)d < 1,$$

the model has non-reconstruction a.s.

Proof. If T extincts, then non-reconstruction obviously hold. Conditioned on non-extinction, we have $\text{br}(T) = d$ a.s. by [Lyo90], and then Theorem 5 applies. \square

Remark 31. As shown in Polyanskiy and Wu [PW17] reconstruction problems on arbitrary directed acyclic graphs (in particular trees) can be shown to be non-reconstructible by reducing to a percolation problem on the same graph. For example for trees, this results in non-reconstruction as long as

$$(31) \quad \eta_{\text{KL}}(M) \text{br}(T) < 1.$$

For any channel M , we have

$$\eta_{\text{KL}}(M, q^*) \leq \eta_{\text{KL}}(M),$$

and the inequality is often strict. So for reversible channels (i.e., $M = M^\vee$), Theorem 5 implies result (31). We do not know, however, how to extend Theorem 5 to general (non-tree) DAGs using input-restricted contraction coefficients.

Formentin and Külske [FK09b] proved a non-reconstruction result very similar to ours. They considered the symmetrized KL divergence

$$D_{\text{SKL}}(P||Q) = D(P||Q) + D(Q||P),$$

which is f -divergence with $f(x) = (x - 1) \log x$. They proved non-reconstruction holds for a Galton-Watson tree with expected offspring d if

$$\eta_{\text{SKL}}(M^\vee, q^*)d < 1.$$

With the method of proof as in Theorem 5 one can strengthen their result so that non-reconstruction holds for

$$\eta_{\text{SKL}}(M^\vee, q^*) \text{br}(T) < 1.$$

Proceeding to input-restricted contraction coefficients, we computed both numerically for several binary asymmetric channels and Potts channels. In Figure 1, we compare $\eta_{\text{SKL}}(M, q^*)$ and $\eta_{\text{KL}}(M, q^*)$ for the binary asymmetric channel

$$M = \begin{pmatrix} 1-a & a \\ b & 1-b \end{pmatrix},$$

for $a = 0.3$ and $b \in [0, 1]$. Simple computation shows that $q^* = (\frac{b}{a+b}, \frac{a}{a+b})$ and $M = M^\vee$.

In Figure 2, we compare the input-restricted SKL and KL contraction coefficients for Potts channels PC_λ for $k = 5$ and $\lambda \in [-\frac{1}{k-1}, 1]$. Because a simplified expression for $\eta_{\text{SKL}}(\text{PC}_\lambda, q^*)$ is not known, we use a lower bound $\bar{\eta}_{\text{SKL}}(\text{PC}_\lambda, q^*)$, which is defined as the sup of $\frac{D_{\text{SKL}}(\text{PC}_\lambda \circ P||q^*)}{D_{\text{SKL}}(P||q^*)}$ considering only distributions $P = (p_1, \dots, p_k)$ with $p_2 = \dots = p_k$. Clearly $\bar{\eta}_{\text{SKL}}(\text{PC}_\lambda, q^*) \leq \eta_{\text{SKL}}(\text{PC}_\lambda, q^*)$. It is conjectured in Formentin and Külske [FK09a] that $\bar{\eta}_{\text{SKL}}(\text{PC}_\lambda, q^*) = \eta_{\text{SKL}}(\text{PC}_\lambda, q^*)$ always.

As shown in Figure 1 and Figure 2, in all these cases, we observe

$$\eta_{\text{KL}}(M^\vee, q^*) \leq \eta_{\text{SKL}}(M^\vee, q^*),$$

which means Theorem 5 yields a stronger non-reconstruction result for these cases.




FIGURE 1. Contraction coefficient comparison for binary asymmetric channels with $a = 0.3$ and varying $b \in [0, 1]$. The figure shows $\eta_{\text{SKL}}(M, q^*) - \eta_{\text{KL}}(M, q^*)$ is always non-negative.




FIGURE 2. Contraction coefficient comparison for Potts channel with $k = 5$ and varying $\lambda \in [-\frac{1}{k-1}, 1]$. The figure shows $\bar{\eta}_{\text{SKL}}(\text{PC}_\lambda, q^*) - \eta_{\text{KL}}(\text{PC}_\lambda, q^*)$ is always non-negative.

We remark that the input-unrestricted KL and SKL contraction coefficients agree. Indeed, the function $x \mapsto (x - 1) \log x$ is operator convex (e.g., [Cha13, Example 3.6]), and thus by [CRS94, Theorem 1], we have

$$\eta_{\text{KL}}(M) = \eta_{\text{SKL}}(M)$$

for any channel M .

Suppose that for some function g , the g -mutual information satisfies the following subadditivity property: for any Markov chain $Y - X - Z$, we have

$$I_g(X; Y, Z) \leq I_g(X; Y) + I_g(X; Z).$$

Then non-reconstruction holds for a tree T with

$$\eta_g(M^\vee, q^*) \text{br}(T) < 1,$$

by the proof of Theorem 5. For mutual information the subadditivity is standard. For I_{SKL} , Formentin-Külske [FK09a] proved that

$$I_{\text{SKL}}(X; Y, Z) = I_{\text{SKL}}(X; Y) + I_{\text{SKL}}(X; Z).$$

It is an interesting question what is the best possible contraction coefficient one can achieve by varying g .

5. POTTS MODEL ON A TREE

In this section, we apply Theorem 5 to get non-reconstruction results for Potts models on a tree. In the Potts model, spins propagate through the Potts channel

PC_λ . Because Potts channels are reversible, $PC_\lambda^\vee = PC_\lambda$. Because Potts channels are symmetric, the invariant distribution is $\pi = \text{Unif}([k])$. Thus Theorem 6 directly follows from Theorem 5.

Let us briefly discuss previous non-reconstruction results for the Potts channel. Mossel and Peres [MP03] proved non-reconstruction for

$$\frac{k\lambda^2}{(k-2)\lambda+2} \text{br}(T) < 1.$$

By Proposition 38 we can see, thus, that this exactly corresponds to invoking a weaker (unrestricted-input) version of η_{KL} . Therefore, Theorem 6 is strictly stronger than [MP03] (see also discussion in Remark 31). Martinelli et al. [MSW07] proved non-reconstruction for regular trees for

$$d(1-\epsilon) \frac{k\lambda^2}{(k-2)\lambda+2} < 1$$

where $\epsilon > 0$ is a function of $k \geq 3$, d , λ in some involved way. Sly [Sly09a] obtained very sharp results for regular trees. In particular, he proved that Kesten-Stigum bound is tight for $k = 3$ and large enough d . It looks hard to extract what general bounds one can achieve by Sly's method. Formentin and Külske [FK09a] gave non-reconstruction results very similar to ours. As discussed in Remark 31, numerical computation suggests that Theorem 6 is stronger than their results.

For certain parameters, we can compute the contraction coefficient $\eta_{\text{KL}}(PC_\lambda, \pi)$ in closed form. In the following we show two examples.

5.1. Binary symmetric channel. For $k = 2$, PC_λ is the binary symmetric channel BSC_δ with $\delta = \frac{1-\lambda}{2}$, which is known (Ahlsvede and Gács [AG76]) to have SDPI coefficient $\eta_{\text{KL}}(\text{BSC}_\delta) = (1-2\delta)^2 = \lambda^2$. Theorem 6 implies non-reconstruction for $(1-2\delta)^2 \text{br}(T) < 1$, which was shown in Bleher et al. [BRZ95] (for regular trees) and Evans et al. [EKPS00] (for general trees).

5.2. Random coloring. The random coloring model with k -colors corresponds to the channel $\text{Col}_k := PC_{-\frac{1}{k-1}}$. This channel acts on input $x \in [k]$ by outputting $y \neq x$ uniformly among all $k-1$ alternatives.

Proposition 32.

$$\eta_{\text{KL}}(\text{Col}_k, \pi) = \frac{\log k - \log(k-1)}{\log k}.$$

Proof. By (14) we have

$$\begin{aligned} \eta_{\text{KL}}(\text{Col}_k, \pi) &= \sup_{x \in (\frac{1}{k}, 1]} \frac{\log k + \frac{1-x}{k-1} \log \frac{1-x}{k-1} + \frac{k+x-2}{k-1} \log \frac{k+x-2}{(k-1)^2}}{\log k + x \log x + (1-x) \log \frac{1-x}{k-1}} \\ &= \sup_{x \in (\frac{1}{k}, 1]} \frac{\log k - \log(k-1) + \frac{1-x}{k-1} \log(1-x) + \frac{k+x-2}{k-1} \log \frac{k+x-2}{k-1}}{\log k + x \log x + (1-x) \log \frac{1-x}{k-1}}. \end{aligned}$$

Taking $x = 1$, we get

$$\eta_{\text{KL}}(\text{Col}_k, \pi) \geq \frac{\log k - \log(k-1)}{\log k}.$$

To prove the proposition, we only need to prove that for $x \in (\frac{1}{k}, 1]$,

$$(32) \quad \frac{\frac{1-x}{k-1} \log(1-x) + \frac{k+x-2}{k-1} \log \frac{k+x-2}{k-1}}{x \log x + (1-x) \log \frac{1-x}{k-1}} \geq \frac{\log k - \log(k-1)}{\log k}.$$

(Note that both numerator and denominator in LHS are non-positive.) Define

$$\begin{aligned} g(x) &= (\log k - \log(k-1))x \log x - \frac{\log k}{k-1}(1-x) \log(1-x), \\ h(x) &= g(x) + (k-1)g\left(\frac{1-x}{k-1}\right). \end{aligned}$$

Rearranging (32), we only need to prove that $h(x) \geq 0$ for $x \in (\frac{1}{k}, 1]$.

We compute that

$$\begin{aligned} g'(x) &= (\log k - \log(k-1))(1 + \log x) + \frac{\log k}{k-1}(1 + \log(1-x)), \\ g''(x) &= (\log k - \log(k-1))\frac{1}{x} - \frac{\log k}{k-1} \frac{1}{1-x}, \\ g'''(x) &= -(\log k - \log(k-1))\frac{1}{x^2} - \frac{\log k}{k-1} \frac{1}{(1-x)^2} < 0. \end{aligned}$$

Claim 33. $h'''(x) < 0$ on $(0, 1)$.

Proof.

$$\begin{aligned} h'''(x) &= g'''(x) - \frac{1}{(k-1)^2} g'''\left(\frac{1-x}{k-1}\right) \\ &= -(\log k - \log(k-1))\frac{1}{x^2} - \frac{\log k}{k-1} \frac{1}{(1-x)^2} \\ &\quad + \frac{1}{(k-1)^2} \left((\log k - \log(k-1))\frac{1}{(\frac{1-x}{k-1})^2} + \frac{\log k}{k-1} \frac{1}{(1-\frac{1-x}{k-1})^2} \right) \\ &= \left(\log \frac{k}{k-1}\right) \left(\frac{1}{(1-x)^2} - \frac{1}{x^2} \right) + \frac{\log k}{k-1} \left(\frac{1}{(k-2+x)^2} - \frac{1}{(1-x)^2} \right) \\ &= \frac{1}{(1-x)^2} \left(\left(\log \frac{k}{k-1}\right) \left(1 - \frac{(1-x)^2}{x^2}\right) + \frac{\log k}{k-1} \left(\frac{(1-x)^2}{(k-2+x)^2} - 1 \right) \right) \\ &=: \frac{1}{(1-x)^2} (s(x) + t(x)). \end{aligned}$$

We have

- (1) $s(x) < 0$ for $x < \frac{1}{2}$, $s(x) > 0$ for $x > \frac{1}{2}$;
- (2) $t(x) < 0$ for $x \in (0, 1)$;
- (3) $s(x)$ is increasing for $x \in (0, 1)$;
- (4) $t(x)$ is decreasing for $x \in (0, 1)$.

So $h'''(x) < 0$ for $x \leq \frac{1}{2}$. For $x \geq \frac{1}{2}$, we have

$$s(x) + t(x) < s(1) + t\left(\frac{1}{2}\right) = \log \frac{k}{k-1} + \frac{\log k}{k-1} \left(\frac{1}{(2k-3)^2} - 1 \right).$$

It is not hard to verify that the last term is < 0 for $k \geq 3$. □

By Claim 33, $h'(x)$ is strictly concave. Because $h'(\frac{1}{k}) = 0$, $h(\frac{1}{k}) = h(1) = 0$, we get that $h(x) > 0$ for $x \in (1/k, 1)$. This finishes the proof. □

Theorem 6 and Proposition 32 together imply non-reconstruction for

$$\text{br}(T) < \frac{\log k}{\log k - \log(k-1)} = (1 - o(1))k \log k.$$

This result was proved for regular trees by Sly [Sly09b] and by Bhatnagar et al. [BVVW11]. Sly had more accurate lower order terms, and his proof works for Galton-Watson trees with Poisson offspring distribution. Efthymiou [Eft15] generalized the result to general Galton-Watson trees with weak assumptions on the offspring distribution. Our result does not assume any conditions on the degree distribution other than the expected offspring.

Remark 34. We show that previous methods based on information contraction do not give the threshold $(1 - o(1))k \log k$. The Evans-Schulman method is based on $\eta_{\text{KL}}(\text{Col}_k)$. For $P_\epsilon = (\frac{1}{2} - \epsilon, \frac{1}{2} + \epsilon, 0, \dots, 0)$ and $Q_\epsilon = (\frac{1}{2} + \epsilon, \frac{1}{2} - \epsilon, 0, \dots, 0)$, we can compute that

$$\eta_{\text{KL}}(\text{Col}_k) \geq \lim_{\epsilon \rightarrow 0} \frac{D(\text{Col}_k \circ P_\epsilon \| \text{Col}_k \circ Q_\epsilon)}{D(P_\epsilon \| Q_\epsilon)} = \frac{1}{k-1}.$$

On the other hand, by comparing with TV contraction coefficient, we have

$$\eta_{\text{KL}}(\text{Col}_k) \leq \eta_{\text{TV}}(\text{Col}_k) = \frac{1}{k-1}.$$

Therefore $\eta_{\text{KL}}(\text{Col}_k) = \frac{1}{k-1}$. The Evans-Schulman method gives non-reconstruction for $d < k - 1$.

The Formentin-Külske method is based on $\eta_{\text{SKL}}(\text{Col}_k, \pi)$. If we let $P_\epsilon = (1 - \epsilon, \frac{\epsilon}{k-1}, \dots, \frac{\epsilon}{k-1})$, then

$$\eta_{\text{SKL}}(\text{Col}_k, \pi) \geq \lim_{\epsilon \rightarrow 0} \frac{D_{\text{SKL}}(\text{Col}_k \circ P_\epsilon \| \pi)}{D_{\text{SKL}}(P_\epsilon \| \pi)} = \frac{1}{k-1}.$$

Therefore the Formentin-Külske method cannot give non-reconstruction results better than for $d < k - 1$.

6. STOCHASTIC BLOCK MODEL

In this section we study the problem of weak recovery of the stochastic block model. In this model, there are n vertices, each independently and uniformly randomly assigned one of k spins. Say vertex $v \in [n]$ has spin $\sigma_v \in [k]$. Generate a random graph, where two vertices u and v have an edge between them with probability

$$\begin{cases} \frac{a}{n} & \text{if } \sigma_u = \sigma_v, \\ \frac{b}{n} & \text{if } \sigma_u \neq \sigma_v, \end{cases}$$

for some absolute constants a and b . We say the model has weak recovery, if given the graph (without knowing the spins σ), we can construct an assignment $\hat{\sigma}$ of spins of the vertices, such that

$$\limsup_{n \rightarrow \infty} \mathbb{E}[d(\sigma, \hat{\sigma})] < 1 - \frac{1}{k},$$

where

$$d(\sigma, \hat{\sigma}) = \frac{1}{n} \min_{\tau \in \mathcal{S}_k} \sum_{i \in [n]} \mathbb{1}\{\sigma_i \neq \hat{\sigma}_{\tau(i)}\}.$$

This definition of distance d is meaningful because $\hat{\sigma}$ is in fact defined only up to a permutation of the spins. If $\hat{\sigma}$ is uniformly random, then the limit of $d(\sigma, \hat{\sigma})$ is $1 - \frac{1}{k}$. So the notion of weak recovery indicates whether we can recover the spin groups better than purely random guessing.

In the following, we show that SDPI-based non-reconstruction results for the Potts model lead to improved impossibility results for the stochastic block model.

6.1. Impossibility of weak recovery via information percolation. We first give an impossibility result via the information percolation method of Polyanskiy and Wu [PW18]. In op. cit., they proved the following statement.

Proposition 35 ([PW18, Proposition 8]). *Weak recovery for the stochastic block model is impossible, if the following tree model has non-reconstruction:*

Consider a Galton-Watson tree with offspring distribution Poisson with mean $d = (\sqrt{a} - \sqrt{b})^2$. For each vertex, we independently and uniformly randomly choose a spin. Say vertex v has spin σ_v . We observe $Y_{u,v} = \mathbb{1}\{\sigma_u = \sigma_v\}$ for each edge (u, v) .

Let ρ denote the root and L_h denote the set of vertices of distance h to ρ . Let Y denote the set of all observations. We say the model has reconstruction, if

$$\lim_{h \rightarrow \infty} \mathbb{E}I(\sigma_\rho; \sigma_{L_h} | Y) = 0.$$

In op. cit., it was proven that the tree model has non-reconstruction when $d < \frac{k}{2}$, using a coupling argument. Here we improve it to

$$d < \frac{1}{\frac{\log k - \log(k-1)}{\log k} \frac{k-1}{k} + \frac{1}{k}} = k - (1 + o(1))k / \log k.$$

Proposition 36. *The tree model in Proposition 35 has non-reconstruction if*

$$d < \frac{1}{\frac{\log k - \log(k-1)}{\log k} \frac{k-1}{k} + \frac{1}{k}}.$$

Proof. The tree model is equivalent to the following top-down process:

- (1) Choose σ_ρ uniformly randomly over $[k]$.
- (2) For an edge (u, v) where u is the parent of v , we randomly choose the transition matrix M , which is the identity I_k with probability $\frac{1}{k}$, and Col_k with probability $1 - \frac{1}{k}$. Then generate the spin of v according to $\mathbb{P}(\sigma_v = j | \sigma_u = i) = M_{i,j}$.

For any vertex u , define $L_{u,h}$ to be the set of descendants of u that have distance h to ρ . Let v be a child of u .

Note that $q^* = \text{Unif}([k])$ is an invariant distribution for both I_k and Col_k . We have $\eta_{\text{KL}}(I_k^\vee, q^*) = 1$, and by Proposition 32, $\eta_{\text{KL}}(\text{Col}_k^\vee, q^*) = \frac{\log k - \log(k-1)}{\log k}$. So if $Y_{u,v} = 1$, we have

$$I(\sigma_u; \sigma_{L_{v,h}} | Y) \leq \eta_{\text{KL}}(I_k^\vee, q^*) I(\sigma_v; \sigma_{L_{v,h}} | Y) = I(\sigma_v; \sigma_{L_{v,h}} | Y)$$

and if $Y_{u,v} = 0$, we have

$$\begin{aligned} I(\sigma_u; \sigma_{L_{v,h}} | Y) &\leq \eta_{\text{KL}}(\text{Col}_k^\vee, q^*) I(\sigma_v; \sigma_{L_{v,h}} | Y) \\ &= \frac{\log k - \log(k-1)}{\log k} I(\sigma_v; \sigma_{L_{v,h}} | Y). \end{aligned}$$

Taking expectation, we get

$$\mathbb{E}I(\sigma_u; \sigma_{L_{v,h}} | Y) \leq \left(\frac{\log k - \log(k-1)}{\log k} \frac{k-1}{k} + \frac{1}{k} \right) \mathbb{E}I(\sigma_v; \sigma_{L_{v,h}} | Y).$$

Rest of the proof is the same as Theorem 5. \square

The last two propositions together show that the stochastic block model does not have weak recovery when

$$(33) \quad (\sqrt{a} - \sqrt{b})^2 < \frac{1}{\frac{\log k - \log(k-1)}{\log k} \frac{k-1}{k} + \frac{1}{k}}.$$

As shown in Figure 3, for certain parameters, (33) leads to slight improvement over [BMNN16].

6.2. Impossibility of weak recovery via Potts channel. In the last section we have seen that the information percolation method together with our tree recursion gives a simple yet strong impossibility result for weak recovery of the stochastic block model. The information percolation method can be understood as comparison with the erasure channel. However, the stochastic block model is more closely related to the Potts channel. In this section we show an even better impossibility result via the Potts model on a tree.

Let $d = \frac{a+(k-1)b}{k}$ and $\lambda = \frac{a-b}{a+(k-1)b}$. We compare the stochastic block model to the Potts model (with Potts channel parameter λ) on a Galton-Watson tree with offspring distribution $\text{Po}(d)$.

Theorem 37 (Mossel et al. [MNS15]). *If the Potts model with parameter λ on a Galton-Watson tree with offspring distribution $\text{Po}(d)$ has non-reconstruction, the corresponding stochastic block model does not have weak recovery.*

Mossel et al. [MNS15] proved Theorem 37 for $k = 2$, and the proof works for general k with little change. Below we give a sketch of the proof.

Proof Sketch. The proof is in two parts. In the first part, we show that for some absolute constant $c > 0$, there exists a coupling between the $c \log n$ -neighborhood of a vertex in the SBM, and the $c \log n$ -neighborhood of the root in the Potts model on a Galton-Watson tree with offspring distribution $\text{Po}(d)$, such that the total variation distance between the two neighborhood (containing spins) is $o(1)$. Proof of this part is by observing that the $c \log n$ -neighborhood in SBM has no cycle with high probability, and can be constructed using a sequence of binomial random variables; on the other hand, the Potts model can be constructed using a sequence of Poisson variables. Then we compare the two sequences of random variables and find that they have very small total variation distance.

In the second part, we show that in the SBM, conditioned on the spins on the boundary of the $c \log n$ -neighborhood, the spins inside and the spins outside are approximately independent. More specifically, if A, B, C is a partition of V such that $\#(A \cup B) = o(n)$ and B separates A and C , then

$$\mathbb{P}(\sigma_A | \sigma_{B \cup C}, G) = (1 + o(1)) \mathbb{P}(\sigma_A | \sigma_B, G)$$

for G and σ with probability $1 - o(1)$. The proof is by writing out the partition function and removing exponents on $1 - \frac{a}{n}$ and $1 - \frac{b}{n}$ that have negligible effect.

Combining the two parts, one can prove that for any constant m and any vertices v_0, \dots, v_m ,

$$I(\sigma_{v_0}; \sigma_{v_1}, \dots, \sigma_{v_m} | G) = o(1).$$

This implies impossibility of weak recovery. \square

Theorem 37 and Theorem 6 imply Theorem 7 immediately. Figure 3 shows a comparison between the impossibility results for $k = 5$.




FIGURE 3. Impossibility of weak recovery results for SBM for $k = 5$. Horizontal axis is a , and vertical axis is b . In the assortative regime, (33) gives better results than [BMNN16] for certain parameters, and Theorem 7 gives the best results among the three.

Note that (16) is equivalent to $d \frac{\lambda^2(k-1)}{2 \log(k-1)} < 1$. In Appendix B, we prove that

$$(34) \quad \eta_{\text{KL}}(\text{PC}_\lambda, \pi) < \frac{\lambda^2(k-1)}{2 \log(k-1)}$$

holds for all $k \geq 3$ and $\lambda \in (0, 1]$. Therefore Theorem 7 strictly improves over (16) in the assortative regime.

APPENDIX A. INPUT-UNRESTRICTED CONTRACTION COEFFICIENT FOR POTTS CHANNELS

Computation of (input-restricted or input-unrestricted) contraction coefficients is often a daunting task. Previously, Makur and Polyanskiy [MP18] obtained lower and upper bounds of input-unrestricted KL divergence contraction coefficients for Potts channels. In this appendix we compute the exact value of these contraction coefficients.

Proposition 38.

$$\eta_{\text{KL}}(\text{PC}_\lambda) = \frac{k\lambda^2}{(k-2)\lambda + 2}.$$

Proof. The result is obvious for $\lambda \in \{0, 1\}$. In the following, assume that $\lambda \notin \{0, 1\}$.

We use the following characterization of contraction coefficient using Rényi correlation [Rén59] (see e.g. Sarmanov [Sar58]). For any channel M , we have

$$(35) \quad \eta_{\text{KL}}(M) = \left(\sup_P \sup_{f,g} \mathbb{E}[f(X)g(Y)] \right)^2$$

where P is a distribution on $[k]$, $X \sim P$, $Y \sim M \circ P$, $f : \mathcal{X} \rightarrow \mathbb{R}$ satisfies $\mathbb{E}_X[f] = 0$ and $\mathbb{E}_X[f^2] = 1$, and $g : \mathcal{Y} \rightarrow \mathbb{R}$ satisfies $\mathbb{E}_Y[g] = 0$ and $\mathbb{E}_Y[g^2] = 1$.

Specialize to $M = \text{PC}_\lambda$. Write $P = (p_1, \dots, p_k)$, $f = (f_1, \dots, f_k)$ and $g = (g_1, \dots, g_k)$. Then

$$(36) \quad \mathbb{E}[f(X)g(Y)] = \sum_{i,j} f_i p_i g_j \mathbb{P}[Y = j | X = i] = \lambda \sum f_i p_i g_i.$$

When $\lambda > 0$, we need to maximize $\sum f_i g_i p_i$. When $\lambda < 0$, we make the transform $f_i \leftarrow -f_i$, and still maximize $\sum f_i g_i p_i$. So we get the following optimization problem.

$$(37) \quad \max \sum f_i g_i p_i$$

$$(38) \quad \text{s.t.} \quad \sum f_i p_i = 0,$$

$$(39) \quad \sum f_i^2 p_i = 1,$$

$$(40) \quad \sum g_i \left(\lambda p_i + \frac{1-\lambda}{k} \right) = 0,$$

$$(41) \quad \sum g_i^2 \left(\lambda p_i + \frac{1-\lambda}{k} \right) = 1,$$

$$(42) \quad p_i \geq 0, \sum p_i = 1.$$

Lower bound. Take

$$\begin{aligned} P &= \left(\frac{1}{2}, \frac{1}{2}, 0, \dots, 0 \right), \\ f &= (1, -1, 0, \dots, 0), \\ g &= (u, -u, 0, \dots, 0) \end{aligned}$$

where

$$u = \sqrt{\frac{k}{(k-2)\lambda + 2}}.$$

Then

$$\sum f_i g_i p_i = u.$$

So

$$\eta_{\text{KL}}(\text{PC}_\lambda) \geq (\lambda u)^2 = \frac{k\lambda^2}{(k-2)\lambda + 2}.$$

Upper bound. Let us fix P and maximize over f and g . Assume for the sake of contrary that $\sum f_i g_i p_i > u$. The set of possible g is bounded; some coordinates of f may be unbounded, but their values do not affect the objective function. So the maximum value of $\sum f_i g_i p_i$ is achieved at some point f and g . Let us compute

the derivatives.

$$\begin{aligned}
\nabla_f \sum f_i g_i p_i &= (g_i p_i)_{i \in [k]}, \\
\nabla_f \sum f_i p_i &= (p_i)_{i \in [k]}, \\
\nabla_f \sum f_i^2 p_i &= (2f_i p_i)_{i \in [k]}, \\
\nabla_g \sum f_i g_i p_i &= (f_i p_i)_{i \in [k]}, \\
\nabla_g \sum g_i (\lambda p_i + \frac{1-\lambda}{k}) &= (\lambda p_i + \frac{1-\lambda}{k})_{i \in [k]}, \\
\nabla_g \sum g_i^2 (\lambda p_i + \frac{1-\lambda}{k}) &= (2g_i (\lambda p_i + \frac{1-\lambda}{k}))_{i \in [k]}.
\end{aligned}$$

By maximality in f , there exists some constants A and B such that

$$(42) \quad g_i p_i = A p_i + B f_i p_i$$

for all i . By maximality in g , there exists some constants C and D such that

$$(43) \quad f_i p_i = C (\lambda p_i + \frac{1-\lambda}{k}) + D g_i (\lambda p_i + \frac{1-\lambda}{k})$$

for all i .

By (42),

$$(44) \quad \sum f_i g_i p_i = \sum f_i (A p_i + B f_i p_i) = B.$$

By (43),

$$(45) \quad \sum f_i g_i p_i = \sum g_i (C (\lambda p_i + \frac{1-\lambda}{k}) + D g_i (\lambda p_i + \frac{1-\lambda}{k})) = D.$$

So $B = D > u > 0$.

For $p_i \neq 0$, we have $g_i = A + B f_i$ by (42).

If for some i , $p_i = 0$, then

$$\frac{1-\lambda}{k} (C + D g_i) = 0.$$

This means

$$\#\{g_i : p_i = 0\} = 1.$$

So we can choose f_i for such i such that

$$(46) \quad g_i = A + B f_i$$

for all i .

From (39), we get

$$\begin{aligned}
0 &= \sum g_i (\lambda p_i + \frac{1-\lambda}{k}) \\
&= \sum (A + B f_i) (\lambda p_i + \frac{1-\lambda}{k}) \\
&= A + B \frac{1-\lambda}{k} \sum f_i.
\end{aligned}$$

From (40), we get

$$\begin{aligned}
1 &= \sum g_i^2 \left(\lambda p_i + \frac{1-\lambda}{k} \right) \\
&= \sum (A^2 + 2ABf_i + B^2 f_i^2) \left(\lambda p_i + \frac{1-\lambda}{k} \right) \\
&= A^2 + 2AB \frac{1-\lambda}{k} \sum f_i + B^2 \lambda + B^2 \frac{1-\lambda}{k} \sum f_i^2 \\
&= B^2 \left(\lambda + \frac{1-\lambda}{k} \sum f_i^2 - \left(\frac{1-\lambda}{k} \sum f_i \right)^2 \right).
\end{aligned}$$

The result then follows from Claim 39 because we have

$$\begin{aligned}
B &= \frac{1}{\sqrt{\lambda + \frac{1-\lambda}{k} \left(\sum f_i^2 - \frac{1-\lambda}{k} \left(\sum f_i \right)^2 \right)}} \\
&\leq \frac{1}{\sqrt{\lambda + \frac{1-\lambda}{k} \left(\sum f_i^2 - \frac{1}{k-1} \left(\sum f_i \right)^2 \right)}} \\
&\leq \frac{1}{\sqrt{\lambda + \frac{1-\lambda}{k} \cdot 2}} = u.
\end{aligned}$$

□

Claim 39. For any distribution P and any f satisfying (37) and (38), we have

$$\sum f_i^2 - \frac{1}{k-1} \left(\sum f_i \right)^2 \geq 2.$$

Proof. Let us first prove the result for f with support size two. WLOG assume that $f_1 > 0$, $f_2 < 0$, $f_3 = \dots = f_k = 0$. One can compute that

$$f_1 = \sqrt{\frac{p_2}{p_1(p_1 + p_2)}}, \quad f_2 = -\sqrt{\frac{p_1}{p_1(p_1 + p_2)}}.$$

Then

$$\begin{aligned}
&f_1^2 + f_2^2 - \frac{1}{k-1} (f_1 + f_2)^2 \\
&\geq f_1^2 + f_2^2 - (f_1 + f_2)^2 \\
&= \frac{1}{p_1 + p_2} \left(\frac{p_2}{p_1} + \frac{p_1}{p_2} - \left(\sqrt{\frac{p_2}{p_1}} - \sqrt{\frac{p_1}{p_2}} \right)^2 \right) \\
&= \frac{2}{p_1 + p_2} \geq 2.
\end{aligned}$$

Let us define

$$\begin{aligned}
S(P) &:= \{f : \sum f_i p_i = 0, \sum f_i^2 p_i = 1\} \\
U(f) &:= \sum f_i^2 - \frac{1}{k-1} \left(\sum f_i \right)^2.
\end{aligned}$$

Now suppose that for some P and $f \in S(P)$ we have $U(f) < 2$. The set $S(P)/\{\pm\}$ is continuous, and there exists $f \in S(P)$ with $U(f) \geq 2$ (e.g., f with support size two), so for sufficiently small $\epsilon > 0$ there exists $f \in S(P)$ such that $U(f) \in (2 - \epsilon, 2)$.

Let $\lambda = -\frac{1}{k-1}$. Take ϵ small enough so that $\lambda + \frac{1-\lambda}{k}(2-\epsilon) > 0$ and choose $f \in S(P)$ with $U(f) \in (2-\epsilon, 2)$. Define

$$\begin{aligned} B &= \frac{1}{\sqrt{\lambda + \frac{1-\lambda}{k}U(f)}} > u, \\ A &= -B\frac{1-\lambda}{k} \sum f_i, \\ g_i &= A + Bf_i \forall i. \end{aligned}$$

One can check that g satisfies (39) and (40), and

$$\sum f_i g_i p_i = B > u.$$

By (35) and (36), this implies

$$\eta_{\text{KL}}(\text{PC}_{-\frac{1}{k-1}}) > \frac{1}{k-1}.$$

However, we have

$$\eta_{\text{KL}}(\text{PC}_{-\frac{1}{k-1}}) \leq \eta_{\text{TV}}(\text{PC}_{-\frac{1}{k-1}}) = \frac{1}{k-1}.$$

Contradiction. □

APPENDIX B. AN UPPER BOUND FOR INPUT-RESTRICTED CONTRACTION COEFFICIENT FOR POTTS CHANNELS

In this appendix we prove an upper bound for the input-restricted KL divergence contraction coefficient for ferromagnetic Potts channels.

Proposition 40. *Fix $k \geq 3$. For all $\lambda \in [0, 1]$, we have*

$$(47) \quad \eta_{\text{KL}}(\text{PC}_\lambda, \pi) \leq \frac{\lambda^2}{(1-\lambda)\frac{2(k-1)\log(k-1)}{k(k-2)} + \lambda}.$$

For all $\lambda \in [-\frac{1}{k-1}, 0]$, we have

$$(48) \quad \eta_{\text{KL}}(\text{PC}_\lambda, \pi) \leq \frac{\lambda^2}{(1+(k-1)\lambda)\frac{2(k-1)\log(k-1)}{k(k-2)} - \lambda\frac{\log k}{(k-1)(\log k - \log(k-1))}}.$$

We first prove a lemma.

Lemma 41. *$\frac{(kx-1)^2}{\psi(x)}$ is concave in $x \in [0, 1]$.*

Proof. Let $f(x) = \frac{(kx-1)^2}{\psi(x)}$.

$$\begin{aligned} f'(x) &= \frac{2k(kx-1)}{\psi(x)} - \frac{(kx-1)^2\psi'(x)}{\psi^2(x)}. \\ f''(x) &= \frac{2k^2}{\psi(x)} - \frac{4k(kx-1)\psi'(x)}{\psi^2(x)} - \frac{(kx-1)^2\psi''(x)}{\psi^2(x)} + \frac{2(kx-1)^2\psi'^2(x)}{\psi^3(x)} \\ &= \frac{2}{\psi^3(x)}(k\psi(x) - (kx-1)\psi'(x))^2 - \frac{(kx-1)^2\psi''(x)}{\psi^2(x)}. \end{aligned}$$

Therefore it suffices to prove that

$$g(x) := \psi^3(x)f''(x) = 2(k\psi(x) - (kx-1)\psi'(x))^2 - (kx-1)^2\psi(x)\psi''(x)$$

is non-positive for $x \in [0, 1]$. Note that $g(\frac{1}{k}) = 0$. So we only need to prove that $g'(x) \geq 0$ for $x \in [0, \frac{1}{k}]$ and $g'(x) \leq 0$ for $x \in [\frac{1}{k}, 1]$.

$$\begin{aligned} g'(x) &= -4(kx - 1)\psi''(x)(k\psi(x) - (kx - 1)\psi'(x)) - 2k(kx - 1)\psi(x)\psi''(x) \\ &\quad - (kx - 1)^2\psi'(x)\psi''(x) - (kx - 1)^2\psi(x)\psi'''(x) \\ &= (kx - 1)(-6k\psi(x)\psi''(x) + (kx - 1)(3\psi'(x)\psi''(x) - \psi(x)\psi'''(x))). \end{aligned}$$

Therefore we would like to prove that

$$u(k, x) := -6k\psi(x)\psi''(x) + (kx - 1)(3\psi'(x)\psi''(x) - \psi(x)\psi'''(x))$$

is non-positive. We enlarge the domain of u and prove that $u(k, x) \leq 0$ for real $k > 1$ and $x \in (0, 1)$.

We fix $x \in (0, 1)$ and consider $u_x(k) := u(k, x)$. We have $u_x(\frac{1}{x}) = 0$. So it suffices to prove that u_x is concave in k . We have

$$\begin{aligned} \psi'(x) &= \log x - \log \frac{1-x}{k-1}, \\ \psi''(x) &= \frac{1}{x} + \frac{1}{1-x}, \\ \psi'''(x) &= \frac{1}{(1-x)^2} - \frac{1}{x^2}, \\ \frac{\partial}{\partial k}\psi(x) &= \frac{1}{k} - \frac{1-x}{k-1}, \\ \frac{\partial}{\partial k}\psi'(x) &= \frac{1}{k-1}, \\ \frac{\partial}{\partial k}\psi''(x) &= \frac{\partial}{\partial k}\psi'''(x) = 0. \end{aligned}$$

So

$$\begin{aligned} u'_x(k) &= -6\psi(x)\psi''(x) - 6k\left(\frac{1}{k} - \frac{1-x}{k-1}\right)\psi''(x) + x(3\psi'(x)\psi''(x) - \psi(x)\psi'''(x)) \\ &\quad + (kx - 1)\left(3\frac{1}{k-1}\psi''(x) - \left(\frac{1}{k} - \frac{1-x}{k-1}\right)\psi'''(x)\right). \\ u''_x(k) &= -12\left(\frac{1}{k} - \frac{1-x}{k-1}\right)\psi''(x) - 6k\left(-\frac{1}{k^2} + \frac{1-x}{(k-1)^2}\right)\psi''(x) \\ &\quad + 6x\frac{1}{k-1}\psi''(x) - 2x\left(\frac{1}{k} - \frac{1-x}{k-1}\right)\psi'''(x) \\ &\quad + (kx - 1)\left(-3\frac{1}{(k-1)^2}\psi''(x) - \left(-\frac{1}{k^2} + \frac{1-x}{(k-1)^2}\right)\psi'''(x)\right) \\ &= \frac{(kx - 1)^2(1 - 2k + (k - 2)x)}{k^2(k - 1)^2x^2(1 - x)^2} \leq 0. \end{aligned}$$

We are done. □

Proof of Proposition 40. For fixed x , we would like to lower bound

$$f_x(\lambda) := \frac{\lambda^2\psi(x)}{\psi(\lambda x + \frac{1-\lambda}{k})}.$$

(Value of $f_x(0)$ is defined by continuity.) Because

$$f_x(\lambda) = \frac{\psi(x)}{(kx-1)^2} \cdot \frac{(k(\lambda x + \frac{1-\lambda}{k}) - 1)^2}{\psi(\lambda x + \frac{1-\lambda}{k})},$$

by Lemma 41, $f_x(\lambda)$ is concave for $\lambda \in [-\frac{1}{k-1}, 1]$.

Let us compute lower bounds of $f_x(\lambda)$ for $\lambda = -\frac{1}{k-1}, 0, 1$.

By Proposition 32, we have

$$(49) \quad f_x(-\frac{1}{k-1}) \geq \frac{\log k}{(k-1)^2(\log k - \log(k-1))}.$$

By L'Hôpital's rule,

$$\begin{aligned} f_x(0) &= \psi(x) \lim_{\lambda \rightarrow 0} \frac{2\lambda}{(x - \frac{1}{k})\psi'(\lambda x + \frac{1-\lambda}{k})} \\ &= \psi(x) \lim_{\lambda \rightarrow 0} \frac{2}{(x - \frac{1}{k})^2\psi''(\lambda x + \frac{1-\lambda}{k})} \\ &= \frac{2(k-1)\psi(x)}{(kx-1)^2}. \end{aligned}$$

By Lemma 41, $g(x) := \frac{(kx-1)^2}{\psi(x)}$ is concave in x . Also

$$g'(1 - \frac{1}{k}) = \frac{2k(k-2)}{\frac{1}{k}(k-2)\log(k-1)} - \frac{(k-2)^2 \cdot 2\log(k-1)}{(\frac{1}{k}(k-2)\log(k-1))^2} = 0.$$

So

$$g(x) \leq g(1 - \frac{1}{k}) = \frac{k(k-2)}{\log(k-1)}$$

and

$$(50) \quad f_x(0) \geq \frac{2(k-1)\log(k-1)}{k(k-2)}.$$

It is easy to see that

$$(51) \quad f_x(1) \geq 1.$$

Because $f_x(\lambda)$ is concave in λ , Inequality (47) follows from (50) and (51), and Inequality (48) follows from (49) and (50). \square

Remark 42. Proof of Proposition 40 implies the first order limit behavior of $\eta_{\text{KL}}(\text{PC}_\lambda, \pi)$ as $\lambda \rightarrow 0$.

$$\lim_{\lambda \rightarrow 0} \frac{\eta_{\text{KL}}(\text{PC}_\lambda, \pi)}{\lambda^2} = \frac{k(k-2)}{2(k-1)\log(k-1)}.$$

Note that for all $k \geq 3$ and $\lambda \in (0, 1]$,

$$\begin{aligned}\eta_{\text{KL}}(\text{PC}_\lambda, \pi) &\leq \frac{\lambda^2}{(1-\lambda)^{\frac{2(k-1)\log(k-1)}{k(k-2)}} + \lambda} \\ &\leq \lambda^2(1-\lambda) \frac{k(k-2)}{2(k-1)\log(k-1)} + \lambda^3 \\ &< \lambda^2 \frac{k(k-2)}{2(k-1)\log(k-1)} \\ &< \lambda^2 \frac{k-1}{2\log(k-1)}.\end{aligned}$$

(Second step is by Cauchy inequality.) So (47) implies (34).

For comparison with input-unrestricted contraction coefficient

$$\eta_{\text{KL}}(\text{PC}_\lambda) = \frac{k\lambda^2}{(k-2)\lambda + 2},$$

we note that $\frac{\lambda^2}{\eta_{\text{KL}}(\text{PC}_\lambda)}$ is linear in λ , and

$$\begin{aligned}\frac{1}{k-1} &< \frac{\log k}{(k-1)^2(\log k - \log(k-1))}, \\ \frac{2}{k} &< \frac{2(k-1)\log(k-1)}{k(k-2)}.\end{aligned}$$

So Proposition 40 implies (15).

APPENDIX C. NON-CONVEXITY OF CERTAIN FUNCTIONS

In this section we prove Proposition 25. Let us first prove a lemma.

Lemma 43. *Let g be a strictly increasing smooth function from $[x_0, x_1]$ to $[y_0, y_1]$, and f be a smooth function from $[x_0, x_1]$ to \mathbb{R} . Assume that $g'(x_0) = f'(x_0) = 0$ and $(g''f''' - f''g''')(x_0) > 0$. Then the function $h = f \circ g^{-1} : [y_0, y_1] \rightarrow \mathbb{R}$ is not concave near y_0 .*

Proof. Directives of h are

$$\begin{aligned}h'(x) &= \frac{f'(g^{-1}(x))}{g'(g^{-1}(x))}, \\ h''(x) &= \left(\frac{f''}{g'} - \frac{f'g''}{g'^2}\right)(g^{-1}(x)) \frac{1}{g'(g^{-1}(x))} \\ &= \left(\frac{f''}{g'^2} - \frac{f'g''}{g'^3}\right)(g^{-1}(x)).\end{aligned}$$

So it suffices to study the sign of $g'f''' - f'g''$ for x near x_0 . Let $u = g'f''' - f'g''$. We have $u(x_0) = 0$. Let us compute the derivatives.

$$\begin{aligned}u' &= g'f'''' - f'g''', \\ u'' &= g'f^{(4)} + g''f'''' - f''g'''' - g'g^{(4)}.\end{aligned}$$

So $u'(x_0) = 0$ and $u''(x_0) = (g''f'''' - f''g''''')(x_0) > 0$. So u is positive near x_0 . \square

Proof of Proposition 25. We apply Lemma 43 to $g = \psi$, $x_0 = \frac{1}{k}$, $x_1 = 1$, $y_0 = 0$, $y_1 = \log k$, and various f . We have

$$\begin{aligned}\psi'(\frac{1}{k}) &= 0, \\ \psi''(\frac{1}{k}) &= \frac{k^2}{k-1}, \\ \psi'''(\frac{1}{k}) &= -\frac{k^3(k-2)}{(k-1)^2}.\end{aligned}$$

Part 1. For b_1 , take

$$f(x) = -(k-1)\xi_1(x) = \log x + (k-1)\log \frac{1-x}{k-1} + k(\psi(x) - \log k).$$

Then

$$\begin{aligned}f'(x) &= \frac{1}{x} - \frac{k-1}{1-x} + k\psi'(x), \\ f''(x) &= -\frac{1}{x^2} - \frac{k-1}{(1-x)^2} + k\psi''(x), \\ f'''(x) &= \frac{2}{x^3} - \frac{2(k-1)}{(1-x)^3} + k\psi'''(x).\end{aligned}$$

So

$$\begin{aligned}f'(\frac{1}{k}) &= 0, \\ f''(\frac{1}{k}) &= -\frac{2k^3}{k-1}, \\ f'''(\frac{1}{k}) &= \frac{3(k-2)k^4}{(k-1)^2}.\end{aligned}$$

We have

$$\begin{aligned}(\psi'' f''' - f'' \psi''')(\frac{1}{k}) &= \frac{k^2}{k-1} \cdot \frac{3(k-2)k^4}{(k-1)^2} - (-\frac{2k^3}{k-1})(-\frac{k^3(k-2)}{(k-1)^2}) \\ &= \frac{k^6(k-2)}{(k-1)^3} > 0.\end{aligned}$$

So Lemma 43 applies.

Part 2. For b_p , $p > 1$, take

$$f(x) = k - (k-1)\xi_p(x) = (x^{\frac{1}{p}} + (k-1)(\frac{1-x}{k-1})^{\frac{1}{p}})(x^{1-\frac{1}{p}} + (k-1)(\frac{1-x}{k-1})^{1-\frac{1}{p}}).$$

For simplicity, write $r = \frac{1}{p}$ and let $u_r(x) = x^r + (k-1)(\frac{1-x}{k-1})^r$. Then $f(x) = u_r(x)u_{1-r}(x)$. Let us compute derivatives of u_r .

$$\begin{aligned}u_r'(x) &= r(x^{r-1} - (\frac{1-x}{k-1})^{r-1}), \\ u_r''(x) &= r(r-1)(x^{r-2} + \frac{1}{k-1}(\frac{1-x}{k-1})^{r-2}), \\ u_r'''(x) &= r(r-1)(r-2)(x^{r-3} - \frac{1}{(k-1)^2}(\frac{1-x}{k-1})^{r-3}).\end{aligned}$$

So

$$\begin{aligned}
u_r\left(\frac{1}{k}\right) &= k^{1-r}, \\
u'_r\left(\frac{1}{k}\right) &= 0, \\
u''_r\left(\frac{1}{k}\right) &= r(r-1)\frac{k}{k-1}\left(\frac{1}{k}\right)^{r-2}, \\
u'''_r\left(\frac{1}{k}\right) &= r(r-1)(r-2)\frac{k(k-2)}{(k-1)^2}\left(\frac{1}{k}\right)^{r-3}.
\end{aligned}$$

Now we compute derivatives of f .

$$\begin{aligned}
f'(x) &= u'_r(x)u_{1-r}(x) + u_r(x)u'_{1-r}(x), \\
f''(x) &= u''_r(x)u_{1-r}(x) + 2u'_r(x)u'_{1-r}(x) + u_r(x)u''_{1-r}(x), \\
f'''(x) &= u'''_r(x)u_{1-r}(x) + 3u''_r(x)u'_{1-r}(x) + 3u'_r(x)u''_{1-r}(x) + u_r(x)u'''_{1-r}(x).
\end{aligned}$$

So

$$\begin{aligned}
f'\left(\frac{1}{k}\right) &= 0, \\
f''\left(\frac{1}{k}\right) &= r(r-1)\frac{k}{k-1}\left(\frac{1}{k}\right)^{r-2} \cdot k^r + (1-r)(-r)\frac{k}{k-1}\left(\frac{1}{k}\right)^{-r-1} \cdot k^{1-r} \\
&= 2r(r-1)\frac{k^3}{(k-1)}, \\
f'''\left(\frac{1}{k}\right) &= r(r-1)(r-2)\frac{k(k-2)}{(k-1)^2}\left(\frac{1}{k}\right)^{r-3} \cdot k^r \\
&\quad + (1-r)(-r)(-r-1)\frac{k(k-2)}{(k-1)^2}\left(\frac{1}{k}\right)^{-r-2} \cdot k^{1-r}, \\
&= -3r(r-1)\frac{k^4(k-2)}{(k-1)^2}.
\end{aligned}$$

So

$$\begin{aligned}
(\psi'' f''' - f'' \psi''')\left(\frac{1}{k}\right) &= \frac{k^2}{k-1}(-3r(r-1)\frac{k^4(k-2)}{(k-1)^2}) \\
&\quad - 2r(r-1)\frac{k^3}{(k-1)}\left(-\frac{k^3(k-2)}{(k-1)^2}\right) \\
&= r(1-r)\frac{k^6(k-2)}{(k-1)^3} > 0.
\end{aligned}$$

So Lemma 43 applies.

Part 3. For s_λ , take

$$f(x) = \psi\left(\lambda x + \frac{1-\lambda}{k}\right).$$

Then

$$\begin{aligned}
f'(x) &= \lambda\psi'\left(\lambda x + \frac{1-\lambda}{k}\right), \\
f''(x) &= \lambda^2\psi''\left(\lambda x + \frac{1-\lambda}{k}\right), \\
f'''(x) &= \lambda^3\psi'''\left(\lambda x + \frac{1-\lambda}{k}\right).
\end{aligned}$$

So

$$\begin{aligned} f'\left(\frac{1}{k}\right) &= 0, \\ f''\left(\frac{1}{k}\right) &= \lambda^2 \psi''\left(\frac{1}{k}\right) = \lambda^2 \frac{k^2}{k-1}, \\ f'''\left(\frac{1}{k}\right) &= \lambda^3 \psi'''\left(\frac{1}{k}\right) = -\lambda^3 \frac{k^3(k-2)}{(k-1)^2}. \end{aligned}$$

We have

$$\begin{aligned} (\psi'' f''' - f'' \psi''')\left(\frac{1}{k}\right) &= \frac{k^2}{k-1} \left(-\lambda^3 \frac{k^3(k-2)}{(k-1)^2}\right) - \lambda^2 \frac{k^2}{k-1} \left(-\frac{k^3(k-2)}{(k-1)^2}\right) \\ &= \frac{k^5(k-2)}{(k-1)^3} (\lambda^2 - \lambda^3) > 0. \end{aligned}$$

So Lemma 43 applies. \square

APPENDIX D. CONCAVITY OF LOG-SOBOLEV COEFFICIENTS

Let K be a Markov kernel with stationary distribution π . Define Dirichlet form $\mathcal{E}(\cdot, \cdot)$ and entropy form $\text{Ent}(\cdot)$ as in the introduction.

For $r \in \mathbb{R}$, we consider the tightest $\frac{1}{r}$ -log-Sobolev inequality, corresponding to

$$\begin{aligned} b_{\frac{1}{r}}(x) &:= \inf_{\substack{f: \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}, \\ \mathbb{E}_\pi f = 1, \text{Ent}_\pi(f) = x}} \mathcal{E}(f^r, f^{1-r}), \\ \Phi_{\frac{1}{r}}(y) &:= \inf_{\substack{f: \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}, \\ \mathbb{E}_\pi f = 1, \mathcal{E}(f^r, f^{1-r}) = y}} \text{Ent}_\pi(f). \end{aligned}$$

The $\frac{1}{r}$ -log-Sobolev constant is

$$\alpha'_{\frac{1}{r}} := \inf_{x>0} \frac{b_{\frac{1}{r}}(x)}{x} = \inf_{y>0} \frac{y}{\Phi_{\frac{1}{r}}(y)}.$$

Remark 44. When $r = 0$, the fraction $\frac{1}{r}$ should be understood as a formal symbol. For $r \in (0, 1)$, $\alpha'_{\frac{1}{r}}$ is the same as $\alpha_{\frac{1}{r}}$ defined in the Introduction. However, in general $\alpha'_{\frac{1}{r}}$ is not equal to $\alpha_{\frac{1}{r}}$. We use the superscript $'$ to emphasize the difference.

Proposition 45. *We have*

- (1) For fixed x , $b_{\frac{1}{r}}(x)$ is concave in r .
- (2) For fixed y , $\Phi_{\frac{1}{r}}(y)$ is convex in r .
- (3) $\alpha'_{\frac{1}{r}}$ is concave in r .

Furthermore, if (K, π) is reversible, then

- (1) For fixed x , $b_{\frac{1}{r}}(x)$ is maximized at $r = \frac{1}{2}$.
- (2) For fixed y , $\Phi_{\frac{1}{r}}(y)$ is minimized at $r = \frac{1}{2}$.
- (3) $\alpha'_{\frac{1}{r}}$ is maximized at $r = \frac{1}{2}$.

Proof. Because $\Phi_{\frac{1}{r}}$ is the inverse function of $b_{\frac{1}{r}}$, it suffices to prove statements about $b_{\frac{1}{r}}$. Because inf of concave functions is still concave, it suffices to prove that for any $f: \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}$, $\mathbb{E}_\pi f = 1$, $\mathcal{E}(f^r, f^{1-r})$ is concave in r .

$$\begin{aligned}
\frac{d}{dr^2} \mathcal{E}(f^r, f^{1-r}) &= \frac{d}{dr^2} \sum_{x,y \in \mathcal{X}} (I - K)(x, y) f(y)^r f(x)^{1-r} \pi(x) \\
&= \sum_{x,y \in \mathcal{X}} (I - K)(x, y) f(y)^r f(x)^{1-r} \pi(x) (\log f(y) - \log f(x))^2 \\
&= \sum_{x \neq y \in \mathcal{X}} -K(x, y) f(y)^r f(x)^{1-r} \pi(x) (\log f(y) - \log f(x))^2 \\
&\leq 0.
\end{aligned}$$

When the Markov chain is reversible, we have $\mathcal{E}(f, g) = \mathcal{E}(g, f)$. So $b_{\frac{1}{r}}(x) = b_{\frac{1}{1-r}}(x)$ and by concavity, $b_{\frac{1}{r}}(x)$ is maximized at $r = \frac{1}{2}$. \square

ACKNOWLEDGEMENT

The authors are grateful to Jingbo Liu for helpful discussions on reconstruction problems on trees, and to anonymous reviewers for valuable comments.

REFERENCES

- [AG76] R. Ahlswede and P. Gács. Spreading of sets in product spaces and hypercontraction of the Markov operator. *The Annals of Probability*, pages 925–939, 1976.
- [Ber67] A. J. Bernstein. Maximally connected arrays on the n-cube. *SIAM Journal on Applied Mathematics*, 15(6):1485–1489, 1967.
- [BGM⁺16] M. Braverman, A. Garg, T. Ma, H. L. Nguyen, and D. P. Woodruff. Communication lower bounds for statistical estimation problems via a distributed data processing inequality. In *Proceedings of the forty-eighth annual ACM symposium on Theory of Computing*, pages 1011–1020. ACM, 2016.
- [BMNN16] J. Banks, C. Moore, J. Neeman, and P. Netrapalli. Information-theoretic thresholds for community detection in sparse networks. In *Conference on Learning Theory*, pages 383–416, 2016.
- [BRZ95] P. M. Bleher, J. Ruiz, and V. A. Zagrebnov. On the purity of the limiting Gibbs state for the Ising model on the Bethe lattice. *Journal of Statistical Physics*, 79(1-2):473–482, 1995.
- [BST10] B. Bhatnagar, A. Sly, and P. Tetali. Reconstruction threshold for the hardcore model. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*, pages 434–447. Springer, 2010.
- [BT06] S. G. Bobkov and P. Tetali. Modified logarithmic Sobolev inequalities in discrete settings. *Journal of Theoretical Probability*, 19(2):289–336, 2006.
- [BVVW11] N. Bhatnagar, J. Vera, E. Vigoda, and D. Weitz. Reconstruction for colorings on trees. *SIAM Journal on Discrete Mathematics*, 25(2):809–826, 2011.
- [Cha13] P. Chansangiam. Operator monotone functions: Characterizations and integral representations. *arXiv preprint arXiv:1305.2471*, 2013.
- [CKZ98] J. Cohen, J. H. B. Kempermann, and Gh. Zbăganu. *Comparisons of stochastic matrices with applications in information theory, statistics, economics and population*. Springer Science & Business Media, 1998.
- [COKPZ18] A. Coja-Oghlan, F. Krzakala, W. Perkins, and L. Zdeborova. Information-theoretic thresholds from the cavity method. *Advances in Mathematics*, 333:694–795, 2018.
- [CRS94] M. Choi, M.B. Ruskai, and E. Seneta. Equivalence of certain entropy contraction coefficients. *Linear algebra and its applications*, 208:29–36, 1994.
- [DSC96] P. Diaconis and L. Saloff-Coste. Logarithmic Sobolev inequalities for finite Markov chains. *The Annals of Applied Probability*, 6(3):695–750, 1996.
- [Eft15] C. Efthymiou. Reconstruction/non-reconstruction thresholds for colourings of general Galton-Watson trees. *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*, page 756, 2015.

- [EKPS00] W. Evans, C. Kenyon, Y. Peres, and L. J. Schulman. Broadcasting on trees and the Ising model. *The Annals of Applied Probability*, 10(2):410–433, 2000.
- [ES99] W. S. Evans and L. J. Schulman. Signal propagation and noisy circuits. *IEEE Transactions on Information Theory*, 45(7):2367–2373, 1999.
- [FK09a] M. Formentin and C. Külske. On the purity of the free boundary condition Potts measure on random trees. *Stochastic Processes and their Applications*, 119(9):2992–3005, 2009.
- [FK09b] M. Formentin and C. Külske. A symmetric entropy bound on the non-reconstruction regime of Markov chains on Galton-Watson trees. *Electronic Communications in Probability*, 14:587–596, 2009.
- [Goe04] S. Goel. Modified logarithmic Sobolev inequalities for some models of random walk. *Stochastic Processes and Their Applications*, 114(1):51–79, 2004.
- [Gro75] L. Gross. Logarithmic Sobolev inequalities. *American Journal of Mathematics*, 97(4):1061–1083, 1975.
- [Har64] L. H. Harper. Optimal assignments of numbers to vertices. *Journal of the Society for Industrial and Applied Mathematics*, 12(1):131–135, 1964.
- [Har76] S. Hart. A note on the edges of the n -cube. *Discrete Mathematics*, 14(2):157–163, 1976.
- [HLPS19] U. Hadar, J. Liu, Y. Polyanskiy, and O. Shayevitz. Communication complexity of estimating correlations. In *Proceedings of the fifty-first annual ACM symposium on Theory of computing*, pages 792–803. ACM, 2019.
- [KS66] H. Kesten and B. P. Stigum. Additional limit theorems for indecomposable multidimensional Galton-Watson processes. *The Annals of Mathematical Statistics*, 37(6):1463–1481, 1966.
- [Lin64] J. H. Lindsey. Assignment of numbers to vertices. *The American Mathematical Monthly*, 71(5):508–516, 1964.
- [LN19] W. Liu and N. Ning. Large degree asymptotics and the reconstruction threshold of the asymmetric binary channels. *Journal of Statistical Physics*, 174(6):1161–1188, 2019.
- [Lyo90] R. Lyons. Random walks and percolation on trees. *The Annals of Probability*, pages 931–958, 1990.
- [Mas14] L. Massoulié. Community detection thresholds and the weak Ramanujan property. In *Proceedings of the forty-sixth annual ACM symposium on Theory of computing*, pages 694–703. ACM, 2014.
- [MM06] M. Mézard and A. Montanari. Reconstruction on trees and spin glass transition. *Journal of statistical physics*, 124(6):1317–1350, 2006.
- [MNS15] E. Mossel, J. Neeman, and A. Sly. Reconstruction and estimation in the planted partition model. *Probability Theory and Related Fields*, 162(3-4):431–461, 2015.
- [MNS18] E. Mossel, J. Neeman, and A. Sly. A proof of the block model threshold conjecture. *Combinatorica*, 38(3):665–708, 2018.
- [MOS13] E. Mossel, K. Oleszkiewicz, and A. Sen. On reverse hypercontractivity. *Geometric and Functional Analysis*, 23(3):1062–1097, 2013.
- [MP03] E. Mossel and Y. Peres. Information flow on trees. *The Annals of Applied Probability*, 13(3):817–844, 2003.
- [MP18] A. Makur and Y. Polyanskiy. Comparison of channels: criteria for domination by a symmetric channel. *IEEE Transactions on Information Theory*, 64(8):5704–5725, 2018.
- [MSW07] F. Martinelli, A. Sinclair, and D. Weitz. Fast mixing for independent sets, colorings, and other models on trees. *Random Structures & Algorithms*, 31(2):134–172, 2007.
- [PS19] Y. Polyanskiy and A. Samorodnitsky. Improved log-Sobolev inequalities, hypercontractivity and uncertainty principle on the hypercube. *Journal of Functional Analysis*, page 108280, 2019.
- [PW17] Y. Polyanskiy and Y. Wu. Strong data-processing inequalities for channels and Bayesian networks. In *Convexity and Concentration*, pages 211–249. Springer, 2017.
- [PW18] Y. Polyanskiy and Y. Wu. Application of information-percolation method to reconstruction problems on graphs. *arXiv preprint arXiv:1806.04195*, 2018.
- [Rén59] A. Rényi. On measures of dependence. *Acta Mathematica Hungarica*, 10(3-4):441–451, 1959.

- [Sar58] O. V. Sarmanov. Maximal correlation coefficient (non-symmetric case). *Dokl. Akad. Nauk SSSR*, 121(1):52–55, 1958.
- [Sly09a] A. Sly. Reconstruction for the Potts model. In *Proceedings of the forty-first annual ACM symposium on Theory of computing*, pages 581–590. ACM, 2009.
- [Sly09b] A. Sly. Reconstruction of random colourings. *Communications in Mathematical Physics*, 288(3):943–961, 2009.
- [WZ73] A. Wyner and J. Ziv. A theorem on the entropy of certain binary sequences and applications–I. *IEEE Transactions on Information Theory*, 19(6):769–772, 1973.
- [XR15] A. Xu and M. Raginsky. Converses for distributed estimation via strong data processing inequalities. In *2015 IEEE International Symposium on Information Theory (ISIT)*, pages 2376–2380. IEEE, 2015.