

Exchangeability, Conformal Prediction, and Rank Tests

Arun Kumar Kuchibhotla

arunku@cmu.edu

Department of Statistics & Data Science, Carnegie Mellon University

Abstract

Conformal prediction has been a very popular method of distribution-free predictive inference in recent years in machine learning and statistics. Its popularity stems from the fact that it works as a wrapper around any prediction algorithm such as neural networks or random forests. Exchangeability is at the core of the validity of conformal prediction. The concept of exchangeability is also at the core of rank tests widely known in nonparametric statistics. In this paper, we review the concept of exchangeability and discuss the implications for conformal prediction and rank tests. We provide a low-level introduction to these topics, and discuss the similarities between conformal prediction and rank tests.

1 Introduction

Exchangeability of random variables is one of the fundamental concepts in statistics, probably right next to the concept of independent and identically distributed (i.i.d.) random variables. Although these two concepts are very closely related, the fact that exchangeability allows for a specific type of dependence between the random variables leads to numerous implications/applications of this concept. One of the most important implications of exchangeability is that the indexing of random variables is immaterial. In technical words, this means that the ranks of real-valued exchangeable random variables are uniform over the set of all permutations. Just this one implication has led to the development of two very different fields in statistics and machine learning, namely, non-parametric rank tests and conformal prediction.

Conformal prediction fills an important gap in machine learning (prediction) algorithms and forecasting. For example, in the context of regression, classical algorithms only provide the point prediction for the response without any uncertainty quantification. Conformal prediction intervals centered at the point prediction algorithm act as such uncertainty measure. Further, classical prediction intervals such as those in linear regression are based on well-specified linear model assumptions. Conformal methods yield prediction regions without any such distributional assumptions.

On the other hand, rank tests concern the statistical problem of testing hypotheses. Most commonly used hypothesis testing procedures asymptotically control the type I error and depend on certain distributional

assumptions so as to ensure “good” asymptotic properties of the test statistic. Rank tests, when available, are finite sample valid and are distribution-free.

The main purpose of this article is to define exchangeability, discuss its implications, and then exhibit the uses of this concept for conformal prediction and rank tests. Both conformal prediction and rank tests make significant use of exchangeability to yield finite sample guarantees for prediction regions and type I error control, respectively, without any distributional assumptions. Conformal prediction regions for data in arbitrary dimensions has been well discussed in the literature, while rank tests for arbitrary dimensions is not as widely discussed. This is not to say that rank tests for multivariate or high-dimensional cases are unknown; see [Friedman \(2003\)](#), [Vayatis et al. \(2009\)](#) for some works. The popularity of conformal prediction stems from the fact that it can be wrapped around any arbitrary algorithm that provides point predictions and leads to a finite sample valid prediction regions for future observations.

In this article, we show how non-parametric rank tests (usually defined for real-valued cases) can also be thought of as wrappers and be applied to arbitrary spaces. The idea of (data independent) dimension reduction for rank tests in arbitrary spaces trivially leads to a finite sample type I error control as mentioned in [Matthews and Taylor \(1996\)](#) and [Lh eritier \(2015\)](#). Note that this also includes the case of transformations based on sample splitting as noted in [Friedman \(2003\)](#) and [Vayatis et al. \(2009\)](#). In this article, we show that the dimension reduction algorithm need not be independent of the data.

Conformal prediction pioneered by [Vovk et al. \(2005\)](#) was introduced to the statistics community by [Lei et al. \(2013\)](#) and further explored in several works ([Lei and Wasserman, 2014](#); [Lei et al., 2018](#); [Chernozhukov et al., 2018](#); [Romano et al., 2019](#); [Foygel Barber et al., 2019](#); [Barber et al., 2021](#); [Chernozhukov et al., 2019](#); [Gupta et al., 2021](#)), among others. For a general overview of the topic, we refer the reader to [Balasubramanian et al. \(2014\)](#). The discussion in all of these papers starts with a “conformity” score. In this article, we do not formally define a “conformity” score but show the application of exchangeability and it is done also because similar thinking helps when we discuss rank tests.

The organization of the article is as follows. In [Section 2](#), we introduce the concept of exchangeability and discuss its implications for ranks of real-valued random variables. Exchangeability is a very intuitive concept that can make it hard to verify rigorously, in some cases. For this reason, in [Section 2.5](#), we discuss the issue of preserving exchangeability via transformations. In [Section 3](#), we discuss the applications of the implication of exchangeability for the construction of distribution-free finite sample valid prediction regions. In [Section 4](#), we discuss the applications of the implication of exchangeability for the construction of distribution-free finite-sample valid rank tests for testing equality of distributions as well as testing independence of two random variables. In both these tests of hypotheses, we allow the random variables to take values in an arbitrary space, thus showing the full strength of exchangeability for this application. In [Section 5](#), we summarize the article and discuss a few open questions.

Most of the results in the article are either known or standard. All the results follow from the definition of exchangeability.

Notation. We use the following notation throughout the article. The notation $\stackrel{d}{=}$ represents the equality in distribution of two random variables. We abbreviate the set $\{1, 2, \dots, n\}$ by $[n]$ for any $n \geq 1$. We write i.i.d. for independent and identically distributed.

2 Exchangeability and Implications

2.1 Definition of Exchangeability

Random variables W_1, \dots, W_n for $n \geq 1$ are said to be exchangeable if

$$(W_1, \dots, W_n) \stackrel{d}{=} (W_{\pi(1)}, \dots, W_{\pi(n)}), \quad (1)$$

for any permutation $\pi : [n] \rightarrow [n]$. Intuitively, exchangeability means that the index of the random variables is immaterial. If W_1, \dots, W_n are real-valued random variables, then the definition (1) is equivalent to the condition that (W_1, \dots, W_n) has the same (joint) cumulative distribution function as that of $(W_{\pi(1)}, \dots, W_{\pi(n)})$, that is, for any $a_1, \dots, a_n \in \mathbb{R}$, and any permutation $\pi : [n] \rightarrow [n]$,

$$\mathbb{P}(W_1 \leq a_1, \dots, W_n \leq a_n) = \mathbb{P}(W_{\pi(1)} \leq a_1, \dots, W_{\pi(n)} \leq a_n). \quad (2)$$

Here $\mathbb{P}(\cdot)$ represents the probability of the event with respect to the probability measure of (W_1, \dots, W_n) . If (W_1, \dots, W_n) has a density $p(\cdot, \dots, \cdot)$ with respect to the Lebesgue measure, then this condition is further equivalent to

$$p(a_1, \dots, a_n) = p(a_{\pi(1)}, \dots, a_{\pi(n)}), \quad (3)$$

for any permutation $\pi : [n] \rightarrow [n]$ and any $a_1, \dots, a_n \in \mathbb{R}$. For random variable in an arbitrary measurable space \mathcal{X} , definition (1) is equivalent to

$$\mathbb{P}(W_1 \in A_1, \dots, W_n \in A_n) = \mathbb{P}(W_{\pi(1)} \in A_1, \dots, W_{\pi(n)} \in A_{\pi(n)}), \quad (4)$$

for any permutation $\pi : [n] \rightarrow [n]$ and any Borel measurable sets A_1, \dots, A_n . A simple consequence of definition (4) is that exchangeable random variables must be identically distributed. To see this, fix a $j \in [n]$ and take the $\pi : [n] \rightarrow [n]$ such that $\pi(1) = j$. Choosing $A_2 = A_3 = \dots = A_n = \mathcal{X}$ in (4) yields $\mathbb{P}(W_1 \in A_1) = \mathbb{P}(W_j \in A_1)$ and because $j \in [n]$ is arbitrary, the result follows. Hence, *identical distributions is a necessary (but not a sufficient) condition for exchangeability*.

Further, it is not hard to verify using (4) that if W_1, \dots, W_n are independent and identically distributed (i.i.d.), then they are exchangeable.

2.2 Examples and Counter-examples

In the following, we provide a few examples of exchangeable random variables.

1. Suppose W_1, W_2 have the joint distribution

$$\begin{pmatrix} W_1 \\ W_2 \end{pmatrix} \sim N \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix} \right). \quad (5)$$

For any $\rho \in [-1, 1]$, exchangeability of W_1 and W_2 can be verified readily using (3). Suppose the mean of the bivariate normal distribution in (5) is changed to $(\mu_1, \mu_2)^\top$. In this case, W_1 and W_2 are not exchangeable unless $\mu_1 = \mu_2$. This follows by noting that W_1 and W_2 do not have the same marginal distribution if $\mu_1 \neq \mu_2$.

2. Suppose $W_i = X_i + Z$ for i.i.d. random variables X_i and another random variable Z independent of X_1, \dots, X_n . Then W_1, \dots, W_n are exchangeable. To prove this, note that

$$\begin{aligned}
\mathbb{P}(W_1 \leq w_1, \dots, W_n \leq w_n) &= \mathbb{E}[\mathbb{P}(W_1 \leq w_1, \dots, W_n \leq w_n | Z)] \\
&= \mathbb{E}[\mathbb{P}(X_1 \leq w_1 - Z, \dots, X_n \leq w_n - Z | Z)] \\
&\stackrel{(a)}{=} \mathbb{E}[\mathbb{P}(X_1 \leq w_1 - Z | Z) \cdots \mathbb{P}(X_n \leq w_n - Z | Z)] \\
&\stackrel{(b)}{=} \mathbb{E}[P_X(w_1 - Z) \cdots P_X(w_n - Z)] \\
&= \mathbb{P}(W_{\pi(1)} \leq w_1, \dots, W_{\pi(n)} \leq w_n).
\end{aligned}$$

Here (a) follows from the assumption that X_1, \dots, X_n are independent and (b) follows from the assumption that X_i 's are identically distributed with distribution function $P_X(\cdot)$. Finally, the last equality follows by retracing the steps with a permutation. Observe that the random variables W_1, \dots, W_n are not independent and their dependence stems from the common random variable Z . This shows that exchangeability in general does not imply independence.

3. Suppose $W_i = f(X_i, Z)$ for a function f , i.i.d. random variables X_1, \dots, X_n and another random variable Z independent of X_1, \dots, X_n . Then W_1, \dots, W_n are exchangeable. The proof is almost verbatim as in the previous case.

2.3 De Finetti's Theorem

A commonality of the last two examples in Section 2.2 is that there exists a random variable Z conditional on which W_1, \dots, W_n are i.i.d.. This represents one of the most general ways of constructing exchangeable random variables. One of the most important results in Bayesian statistics states that if $n = \infty$, then there is no other way of constructing exchangeable random variables. Formally, we have the following De Finetti's representation theorem for infinite sequence of exchangeable random variables. The following statement is taken from [Schervish \(2012, Theorem 1.49\)](#).

Theorem 1 (De Finetti's Representation Theorem). *Let (S, \mathcal{A}, μ) be a probability space, and let $(\mathcal{W}, \mathcal{B})$ be a Borel space. For each n , let $W_n : S \rightarrow \mathcal{X}$ be measurable. The sequence $\{W_i\}_{i=1}^\infty$ is exchangeable if and only if there is a random probability measure \mathbf{P} on $(\mathcal{W}, \mathcal{B})$ such that, conditional on $\mathbf{P} = P$, $\{W_i\}_{i=1}^\infty$ are independent and identically distributed with distribution P . Furthermore, if the sequence is exchangeable, then the distribution of \mathbf{P} is unique, and $n^{-1} \sum_{i=1}^n \mathbb{1}\{W_i \in B\}$ converges to $\mathbf{P}(B)$ almost surely for each $B \in \mathcal{B}$.*

[De Finetti \(1929\)](#) proved the result only for random variables W_i taking values in $\{0, 1\}$. It was extended to arbitrary compact Hausdorff spaces by [Hewitt and Savage \(1955\)](#). Interested readers can refer to [Ressel \(1985\)](#) and [Aldous \(1985\)](#) for a review and a detailed discussion of probabilistic aspects of exchangeability. The hypothesis that there are infinite number of elements in the sequence is crucial and it can be shown that the result is false for a finite sequence of exchangeable random variables, in general. See [Schervish \(2012, Chapter 1, Section 1.2\)](#) for a counterexample. Also, see Theorem 1.48 of [Schervish \(2012\)](#) for a representation theorem for a finite sequence of exchangeable random variables.

2.4 Implication for Ranks

One of the most important implications of exchangeability of real-valued random variables is that the ranks of W_1, \dots, W_n (among this sequence) are uniformly distributed on $\{1, 2, \dots, n\}$. If $\mathcal{X} := \{x_1, \dots, x_n\}$ is a set with n elements (meaning that all the elements in \mathcal{X} are distinct), then the rank of x_i among \mathcal{X} can be defined as

$$\text{rank}^*(x_i; \mathcal{X}) := |\{j \in [n] : x_j \leq x_i\}|. \quad (6)$$

In other words, rank of x_i is the number of elements in \mathcal{X} (including itself) that are smaller than or equal to x_i . If the set \mathcal{X} has fewer than n elements (meaning that some elements of \mathcal{X} are equal), then ranks as defined in (6) will lead to ties, that is, the elements of \mathcal{X} that are equal get the same rank.

Ranking with ties would, in general, lead to a distribution of $(\text{rank}(X_i; \mathcal{X}), i \in [n])$ that depends on the distribution of X_i . For example, if we have a sequence of $n = 100$ i.i.d. Bernoulli(p) random variables, then the number of 1's and 2's in the sequence of ranks defined by (6) depends on p . This causes a hindrance to the distribution-free prediction and non-parametric ranks. There are different ways of breaking ties in ranks obtained from (6). For simplicity, we consider the following definition (from Vorlickova (1972)) of ranks that works for all sequences alike.

Definition 1 (Rank). *For a set of real numbers $\mathcal{X} := \{x_1, \dots, x_n\}$, define the rank of x_i among \mathcal{X} as*

$$\begin{aligned} \text{rank}(x_i; \mathcal{X}) &:= |\{j \in [n] : x_j + \xi U_j \leq x_i + \xi U_i\}| \\ &= \text{rank}^*(x_i + \xi U_i; \mathcal{X} + \xi \mathcal{U}), \end{aligned} \quad (7)$$

where $\xi > 0$ is arbitrary and U_1, \dots, U_n are iid $\text{Unif}[-1, 1]$ random variables. Here $\mathcal{X} + \xi \mathcal{U} = \{x_i + \xi U_i : 1 \leq i \leq n\}$.

Because U_1, \dots, U_n are almost surely distinct, we get that $x_i + \xi U_i, i \in [n]$ are also distinct with probability one, irrespective of whether x_1, \dots, x_n have ties. Essentially, jittering the original sequence with some noise breaks the ties and brings the situation back to the case of no ties. This is important to obtain the distribution-free nature of the results to be described and with ties this would not be possible. For further convenience, we formally define the jittered sequence.

Definition 2 (Jittered Sequence). *For any $n \geq 1$ and sequence x_1, \dots, x_n of real numbers, the jittered sequence with parameter $\xi > 0$ is defined as x_1^*, \dots, x_n^* with*

$$x_i^* := x_i + \xi U_i.$$

Here U_1, \dots, U_n are iid $\text{Unif}[-1, 1]$ random variables. We suppress ξ in the notation of x_i^* for convenience.

In general, the definition of rank above depends on $\xi > 0$. If $x_i, i \in [n]$ do not have ties, then $\text{rank}(x_i; \mathcal{X})$ in (7) matches the one in (6) as ξ tends to zero. For a general sequence with ties, the rank in (7) breaks the ties for ranking randomly as ξ tends to zero. For the purposes of exchangeability, the size of ξ is immaterial but in practice, fixing ξ to be a small constant such as 10^{-8} relative to the spacings in the data works as expected for all sets.

Definition 1 of ranks coupled with the definition of exchangeability implies the following result proved in Appendix B.

Theorem 2. If W_1, \dots, W_n are exchangeable random variables, then for any $\xi > 0$,

$$(\text{rank}(W_i; \{W_1, \dots, W_n\}) : i \in [n]) \sim \text{Unif}(\{\pi : [n] \rightarrow [n]\}).$$

Here $\text{Unif}(\{\pi : [n] \rightarrow [n]\})$ represents the uniform distribution over all permutations of $[n]$, that is, each permutation has an equal probability of $1/n!$.

Theorem 2 shows that the ranks of $W_i, i \in [n]$ are exchangeable, and further that their distribution does not depend on the distribution of W_i . It should be mentioned here that the distribution of the ranks is computed including the randomness of U_1, \dots, U_n ; they are not conditioned on. This theorem also represents one of the most useful implications of exchangeability and is crucial in proving the validity of rank tests as well as conformal prediction.

For the validity guarantees of rank tests, Theorem 2 in its form is enough. For the validity guarantees of conformal prediction, we need the following corollary (proved in Appendix B) of Theorem 2.

Corollary 1. Under the assumptions of Theorem 2, for any $\xi > 0$, we have

$$\mathbb{P}\left(\text{rank}(W_n; \{W_1, \dots, W_n\}) \leq t\right) = \frac{\lfloor t \rfloor}{n},$$

where, for $t \in \mathbb{R}$, $\lfloor t \rfloor$ represents the largest integer smaller than or equal to t . Moreover, the random variable $P := \text{rank}(W_n; \{W_1, \dots, W_n\})/n$ is a valid p -value, i.e.,

$$\mathbb{P}(P \leq \alpha) \leq \alpha \quad \text{for all } \alpha \in [0, 1].$$

2.5 Transformations Preserving Exchangeability

Theorem 2 holds for real-valued random variables¹ and to explore the full strength of exchangeability in arbitrary spaces, we transform random variables from arbitrary spaces to the real line. If the transformation to the real line does not depend on the data (or is constructed from an independent data), then it is relatively easy to verify that the transformed variables also form an exchangeable sequence. In many cases, one might not have access to independent data or might want to use the full data for a more “powerful” transformation. For such purposes, we need a result to verify exchangeability of random variables after transformation stated below. As a motivation, consider the following examples:

- Suppose W_1, \dots, W_n are exchangeable. Consider the transformed variables $W_1 - \overline{W}_n, \dots, W_n - \overline{W}_n$, where \overline{W}_n is the average of the n variables. In this case the transformation takes n variables to n variables. Intuitively, these are exchangeable but how does one prove it rigorously. One could use (2).
- In the same setting as above, consider the transformed variables to be $W_1 - \overline{W}_{-3}, W_2 - \overline{W}_{-3}, W_3 - \overline{W}_{-3}$, where \overline{W}_{-3} is the average of W_4, \dots, W_n (the sequence without the first three elements).
- In the same setting as above, consider the transformed variables to be $W_1 - \overline{W}_{n-1}, \dots, W_{n-1} - \overline{W}_{n-1}, W_n - \overline{W}_{n-1}$. In this case, the transformation depends only on first $n - 1$ variables and takes a sequence of n variables to n variables.

¹For random variables in a metric space, the definition of ranks can be extended and a result similar to Theorem 2 can be proved; see [Deb and Sen \(2021\)](#) for details.

The following is an important result about transformations preserving exchangeability taken from [Dean and Verducci \(1990\)](#) and [Commenges \(2003, Section 2.5\)](#). The setting is as follows: W_1, \dots, W_n are random variables taking values in a space \mathcal{W} and G is a transformation taking a vector of n elements in \mathcal{W} to a vector of m elements in another space \mathcal{W}' . Here \mathcal{W} and \mathcal{W}' are arbitrary sets. (Usually \mathcal{W} would be an arbitrary space and \mathcal{W}' is the real line.) For any $y = (y_1, \dots, y_m)$ and a permutation $\pi_1 : [m] \rightarrow [m]$, set $\pi_1 y = (y_{\pi_1(1)}, \dots, y_{\pi_1(m)})$.

Theorem 3 ([Dean and Verducci \(1990\)](#)). *Suppose $W = (W_1, \dots, W_n) \in \mathcal{W}^n$ is a vector of exchangeable random variables. Fix a transformation $G : \mathcal{W}^n \rightarrow (\mathcal{W}')^m$. If for each permutation $\pi_1 : [m] \rightarrow [m]$ there exists a permutation $\pi_2 : [n] \rightarrow [n]$ such that*

$$\pi_1 G(w) = G(\pi_2 w), \quad \text{for all } w \in \mathcal{W}^n, \quad (8)$$

then $G(\cdot)$ preserves exchangeability of W . Conversely, if $G(\cdot)$ preserves exchangeability of W whatever the distribution of W , then for each permutation $\pi_1 : [m] \rightarrow [m]$ and $w \in \mathcal{W}^n$, there exists a permutation $\pi_2 : [n] \rightarrow [n]$ (possibly depending on w) such that $\pi_1 G(w) = G(\pi_2 w)$. Furthermore, if $G(\cdot)$ is a linear transformation, then G is exchangeability preserving if and only if (8) holds true.

Theorem 3 follows from the proof of Theorem 4 of [Dean and Verducci \(1990\)](#). [Commenges \(2003, Section 2.5\)](#) states (without proof) that (8) is a necessary and sufficient condition for $G(\cdot)$ to be exchangeability preserving. At present, we could only prove Theorem 3; see Appendix B.3. The only difference between the necessary and sufficient conditions in Theorem 3 is that the permutation π_2 can depend on w in the necessary condition. This theorem, in words, states that a transformation of exchangeable random variables is exchangeable if a permutation of the transformed random variables is equal to the transformation applied to a permutation of the original exchangeable random variables.

As an application, we will revisit the examples discussed above.

- In the first example above, the transformation is

$$G : (W_1, \dots, W_n) \mapsto (W_1 - \bar{W}_n, \dots, W_n - \bar{W}_n).$$

A permutation of the right hand side is

$$(W_{\pi(1)} - \bar{W}_n, \dots, W_{\pi(n)} - \bar{W}_n),$$

and this is equal to $G(W_{\pi(1)}, \dots, W_{\pi(n)})$, because the average of n variables is a symmetric function and does not change with a permutation. Hence, $G(W_1, \dots, W_n)$ is a vector of exchangeable random variables whenever W_1, \dots, W_n are exchangeable by Theorem 3.

- In the second example, the transformation is

$$G : (W_1, \dots, W_n) \mapsto (W_1 - \bar{W}_{-3}, W_2 - \bar{W}_{-3}, W_3 - \bar{W}_{-3}).$$

In this case if we permute the right hand side to get $(W_{\pi(1)} - \bar{W}_{-3}, W_{\pi(2)} - \bar{W}_{-3}, W_{\pi(3)} - \bar{W}_{-3})$, then it corresponds to applying the same permutation on the first three elements of W_1, \dots, W_n and leaving the remaining elements as is. Formally, take $\pi_1 : [n] \rightarrow [n]$ such that $\pi_1(1) = \pi(1), \pi_1(2) = \pi(2), \pi_1(3) = \pi(3)$, and $\pi_1(i) = i$ for $i \geq 4$. This implies that $W_1 - \bar{W}_{-3}, W_2 - \bar{W}_{-3}, W_3 - \bar{W}_{-3}$ are exchangeable if W_1, \dots, W_n are exchangeable, again by Theorem 3. This application is related to the split conformal method discussed in Section 3.3.

- For the third example, the transformation is

$$G : (W_1, \dots, W_n) \mapsto (W_1 - \bar{W}_{n-1}, \dots, W_{n-1} - \bar{W}_{n-1}, W_n - \bar{W}_{n-1}).$$

Note that this is a linear transformation and Theorem 3 provides a necessary and sufficient condition. If we apply a permutation π on $G(W_1, \dots, W_n)$, we get

$$(W_{\pi(1)} - \bar{W}_{n-1}, \dots, W_{\pi(n-1)} - \bar{W}_{n-1}, W_{\pi(n)} - \bar{W}_{n-1}). \quad (9)$$

Because \bar{W}_{n-1} is an asymmetric function of (W_1, \dots, W_n) , the vector in (9) is not equal to the transformation G applied to $(W_{\pi(1)}, \dots, W_{\pi(n)})$. This implies that, in general, $G(W_1, \dots, W_n)$ is not a vector of exchangeable random variables.

Having described in details the implications of exchangeability, we now proceed to explore the applications for conformal prediction and rank tests. All the results that follow are corollaries of the results in the current section. It might be worth mentioning here that none of the results in the paper are new or difficult to prove. They are all standard. This is the main intent of the article: to show that most of conformal prediction and non-parametric rank tests follow from some basic facts about exchangeability.

3 Conformal Prediction

Conformal prediction is a generic tool for finite sample, distribution-free valid predictive inference introduced by Vovk et al. (2005) and Shafer and Vovk (2008). This method of predictive inference was reintroduced to the statistics community by Lei et al. (2013).

3.1 Formulation of the Problem

The general formulation of the prediction problem is as follows. Given realizations of n exchangeable random variables W_1, \dots, W_n , construct a prediction region for a future random variable, W_{n+1} , that is exchangeable with the first n random variables, i.e., W_1, \dots, W_{n+1} is a sequence of exchangeable random variables. Mathematically, for $\alpha \in [0, 1]$, construct a prediction region $\hat{\mathcal{R}}_{n,\alpha}$ depending on W_1, \dots, W_n , that is,

$$\hat{\mathcal{R}}_{n,\alpha} = \hat{\mathcal{R}}_{n,\alpha}(W_1, \dots, W_n),$$

such that the $(n+1)$ -st random variable W_{n+1} belongs in this region with a probability of at least $1 - \alpha$:

$$\mathbb{P}(W_{n+1} \in \hat{\mathcal{R}}_{n,\alpha}) \geq 1 - \alpha, \quad (10)$$

whenever W_1, \dots, W_{n+1} form a sequence of exchangeable random variables. In (10), the probability $\mathbb{P}(\cdot)$ is the probability of the event with respect to the joint distribution of (W_1, \dots, W_{n+1}) . In the general formulation here, there is no restriction on the random variables W_1, \dots, W_{n+1} to be real-valued; in fact, they may be elements of an arbitrary sample space \mathcal{W} . Prediction problems in general spaces occur in applications such as functional data analysis (Lei et al., 2015) and image prediction (Bates et al., 2021).

The idea of conformal prediction for real-valued random variables would be that the rank of the future observation W_{n+1} among the collection $\{W_1, \dots, W_{n+1}\}$ is equally likely to be any of $1, 2, \dots, n+1$. We will deal with prediction in arbitrary spaces by using transformations that map these spaces to the real line, so that the rank transformation can be applied and the uniform distribution of the ranks can be leveraged (Section 2.4). To this end, Theorem 3 would play an important role.

3.2 Full Conformal Prediction

3.2.1 Real-valued Random Variables.

If W_1, \dots, W_{n+1} are real valued and exchangeable, then with ranks defined as in Definition 1, Corollary 1 implies that

$$\mathbb{P}\left(\text{rank}(W_{n+1}; \{W_1, \dots, W_{n+1}\}) \leq \lceil (n+1)(1-\alpha) \rceil\right) = \frac{\lceil (n+1)(1-\alpha) \rceil}{n+1}.$$

It is easy to verify that the right hand side is at least $1-\alpha$ and at most $1-\alpha+1/(n+1)$. Hence, a one-sided prediction region can be constructed as follows:

$$\widehat{\mathcal{R}}_{n,\alpha} := \left\{ w \in \mathbb{R} : \text{rank}(w; \{W_1, \dots, W_n, w\}) \leq \lceil (n+1)(1-\alpha) \rceil \right\},$$

This is documented in the following result.

Proposition 1. *If $W_1, \dots, W_{n+1} \in \mathbb{R}$ form a sequence of exchangeable random variables, then*

$$1-\alpha \leq \mathbb{P}\left(W_{n+1} \in \widehat{\mathcal{R}}_{n,\alpha}\right) \leq 1-\alpha + \frac{1}{n+1}, \quad \text{for all } n \geq 1, \alpha \in [0, 1],$$

where the probability extends over all variables W_1, \dots, W_{n+1} .

This result is essentially proved above and follows from the basic corollary 1 of the definition of exchangeability. Although the result is a restatement of Corollary 1, formulating the result in terms of prediction regions provides a form of finite sample distribution-free valid inference. Furthermore, the interval is not overly conservative in that the coverage is at most $1/(n+1)$ away from the required coverage of $(1-\alpha)$. The set $\widehat{\mathcal{R}}_{n,\alpha}$ is defined implicitly and we now describe the computation of this prediction set.

Pseudocode 1: The set $\widehat{\mathcal{R}}_{n,\alpha}$ can be computed as follows.

1. Take $\xi = 10^{-8}$ and generate U_1, \dots, U_n, U_{n+1} from the $\text{Unif}[-1, 1]$ distribution. Define the jittered sequence $W_i^* := W_i + \xi U_i$, $i \in [n]$.
2. Sort the jittered random variables and let the sorted vector be

$$W_{(1)}^* \leq \dots \leq W_{(n)}^*.$$

3. Compute $I := \lceil (n+1)(1-\alpha) \rceil$ and report the interval

$$\left(-\infty, W_{(I)}^* - \xi U_{n+1}\right].$$

This is the simplest example of the full conformal method where exchangeability is invoked on the original set of real-valued random variables W_1, \dots, W_n, W_{n+1} . Sometimes it might be useful to apply Proposition 1 to a transformed data. For example, note that the prediction region $\widehat{\mathcal{R}}_{n,\alpha}$ is a one-sided interval and a two-sided interval might be preferable in practice.

3.2.2 Arbitrary Spaces.

For notational convenience, let $Z_1, Z_2, \dots, Z_n \in \mathcal{Z}$ be exchangeable random variables and we want to construct a prediction region for Z_{n+1} , when Z_1, \dots, Z_n, Z_{n+1} is a sequence of exchangeable random variables in \mathcal{Z} . Here \mathcal{Z} can be \mathbb{R}, \mathbb{R}^d , or any other arbitrary space.

For any (non-random) transformation $g : \mathcal{Z} \rightarrow \mathbb{R}$,

$$g(Z_1), g(Z_2), \dots, g(Z_n), g(Z_{n+1}),$$

are real-valued exchangeable random variables. This fact can be verified based on Theorem 3. Hence, Proposition 1 applies and we obtain

$$\mathbb{P}\left(g(Z_{n+1}) \leq (g(Z))_{([(n+1)(1-\alpha)])}^* - \xi U_{n+1}\right) \geq 1 - \alpha, \quad (11)$$

where $(g(Z))_{([(n+1)(1-\alpha)])}^*$ is the $[(n+1)(1-\alpha)]$ -th largest value of $g(Z_i) + \xi U_i, 1 \leq i \leq n$. Inequality (11) yields a valid finite sample prediction region, irrespective of what $g : \mathcal{Z} \rightarrow \mathbb{R}$ is.

For concrete examples, one can consider the following transformations. If $\mathcal{Z} = \mathbb{R}$, taking $g(z) = |z|$ yields the prediction region

$$\{z \in \mathbb{R} : |z| \leq |Z|_{([(n+1)(1-\alpha)])}^* - \xi U_{n+1}\}. \quad (12)$$

This is a two-sided interval centered at 0. If $\mathcal{Z} = \mathbb{R}^d$ or a general normed linear space with norm $\|\cdot\|$, then taking $g(z) = \|z\|$ yields the prediction region

$$\{z \in \mathcal{Z} : \|z\| \leq \|Z\|_{([(n+1)(1-\alpha)])}^* - \xi U_{n+1}\}. \quad (13)$$

This is a symmetric bounded set centered at $0 \in \mathcal{Z}$. Prediction sets (12) and (13) both suffer from the same disadvantage: they are symmetric around zero. If the true distribution of Z_i 's has a mode at some $z_0 \neq 0$, then these prediction regions are unnecessarily large. For instance, for the normal distribution with mean 1, the shortest prediction interval is centered at 1. However, a valid symmetric prediction interval centered at 0 is about 1.34 times larger than the shortest interval centered at 1.

This disadvantage can be rectified by considering a data-dependent transformation. Now we need to consider transformations that retain exchangeability and hence, Theorem 3 plays a crucial role. A data-dependent transformation $g : \mathcal{Z} \rightarrow \mathbb{R}$ denoted by $g(z; Z_1, \dots, Z_n, Z_{n+1})$ is said to be permutation invariant if for any permutation $\pi : [n+1] \rightarrow [n+1]$,

$$g(z; Z_1, \dots, Z_n, Z_{n+1}) = g(z; Z_{\pi(1)}, Z_{\pi(2)}, \dots, Z_{\pi(n)}, Z_{\pi(n+1)}). \quad (14)$$

For intuition, one may consider the following examples of permutation invariant transformations.

- **Location Centering:** If $\mathcal{Z} = \mathbb{R}$, then an example transformation is

$$g(z; Z_1, \dots, Z_n, Z_{n+1}) := \left| z - \frac{1}{n+1} \sum_{j=1}^{n+1} Z_j \right|.$$

Because the mean is permutation invariant, this is a permutation invariant transformation. Clearly, one can replace the mean by the median of Z_1, \dots, Z_{n+1} , which is also a symmetric function. If \mathcal{Z} is a normed linear space with the norm $\|\cdot\|$, then define

$$g(z; Z_1, \dots, Z_n, Z_{n+1}) := \left\| z - \frac{1}{n+1} \sum_{j=1}^{n+1} Z_j \right\|.$$

Once again, this is also permutation invariant.

- **Density Transformation:** If $\mathcal{Z} = \mathbb{R}$, then define

$$g(z; Z_1, \dots, Z_n, Z_{n+1}) := \frac{1}{\hat{p}_{n+1}(z)}, \quad (15)$$

where $\hat{p}_{n+1}(\cdot)$ is a density estimator that depends permutation invariantly on Z_1, \dots, Z_{n+1} . For example, one can take $\hat{p}_{n+1}(\cdot)$ to be the kernel density estimator

$$\hat{p}_{n+1}(s) := \frac{1}{(n+1)h} \sum_{i=1}^{n+1} k\left(\frac{s - Z_j}{h}\right),$$

for a kernel function $k(\cdot)$ and bandwidth $h > 0$. A similar density estimator can be constructed in normed spaces. The transformation (15) was considered in [Lei et al. \(2013\)](#) to construct asymptotically optimal prediction sets in \mathbb{R}^d . Here optimality is in terms of smallest volume or Lebesgue measure.

- **Regression Residual:** If $\mathcal{Z} = \mathbb{R} \times \mathcal{X}$ and $Z_i = (Y_i, X_i), 1 \leq i \leq n+1$ for a regression data, then an example transformation targeting the response is

$$g(z; Z_1, \dots, Z_n, Z_{n+1}) := |y - \hat{\mu}(x; Z_1, \dots, Z_{n+1})|, \quad (16)$$

for $z = (y, x)$. Here $\hat{\mu}(\cdot; Z_1, \dots, Z_{n+1})$ represents a non-parametric regression mean estimator that depends permutation invariantly on Z_1, \dots, Z_{n+1} . For example, it can be the kernel regression estimator

$$\hat{\mu}(x; Z_1, \dots, Z_{n+1}) := \frac{\sum_{j=1}^{n+1} Y_j k((x - X_j)/h)}{\sum_{j=1}^{n+1} k((x - X_j)/h)},$$

for a kernel function $k(\cdot)$ and bandwidth $h > 0$. The transformation (16) leads to a non-trivial prediction region for the response but a trivial one for the predictors. This feature will be discussed later in [Section 3.4](#).

Under the permutation invariance condition (14), [Theorem 3](#) implies that when $z = Z_{n+1}$,

$$W_i(z) := g(Z_i; Z_1, \dots, Z_n, z), 1 \leq i \leq n, W_{n+1}(z) := g(z; Z_1, \dots, Z_n, z),$$

is a sequence of exchangeable real-valued random variables; see [Proposition 4](#) (of [Appendix A](#)) for a formal result. Hence applying [Proposition 1](#) to $W_i(z)$'s, we obtain the prediction region

$$\{z \in \mathcal{Z} : \text{rank}(W_{n+1}(z); \{W_1(z), \dots, W_n(z), W_{n+1}(z)\}) \leq [(n+1)(1-\alpha)]\}. \quad (17)$$

A distinguishing feature of this prediction region compared to the one from [Pseudocode 1](#), and in (12), (13), is that there is no closed form expression. Note that all the W_i 's depend on the unknown $z = Z_{n+1}$. Computing the region (17), in general, requires computing $g(\cdot; Z_1, \dots, Z_n, z)$ for all $z \in \mathcal{Z}$ and then verifying the rank condition in (17). Hence, the prediction region (17) is, in general, computationally inefficient. It should be mentioned that there do exist cases where the full conformal prediction set (17) can be computed efficiently; see [Burnaev and Vovk \(2014\)](#), [Chen et al. \(2018\)](#), [Lei \(2019\)](#), and [Ndiaye and Takeuchi \(2019\)](#) for some examples.

Because of the heavy computational cost of the full conformal method, we now focus on the split conformal method that provides the same finite sample validity guarantees and is computationally efficient.

3.3 Split Conformal Prediction

3.3.1 Real-valued Random Variables.

Following Papadopoulos et al. (2002) and Lei et al. (2018), we now discuss a split conformal prediction method which can be used to construct efficient prediction regions without the computational burden of the full conformal method. The procedure, in words, is as follows. We split the exchangeable sequence into two parts, and from the first part, we compute the average. Then the variables in the second part (along with the future variable) centered at the average of the first part are exchangeable, which leads us to a two-sided prediction interval. Formally, given random variables $Z_1, \dots, Z_n \in \mathbb{R}$ and $n_1 \in [n]$, construct the split

$$\mathcal{T} := \{Z_1, \dots, Z_{n_1}\}, \quad \text{and} \quad \mathcal{C} := \{Z_{n_1+1}, \dots, Z_n\}. \quad (18)$$

(The notations \mathcal{T} and \mathcal{C} stand for training and calibration sets, respectively.) From the training set, compute $\bar{Z}_{\mathcal{T}}$, the average of the observations in \mathcal{T} . From the calibration set, compute the random variables $|Z_{n_1+1} - \bar{Z}_{\mathcal{T}}|, \dots, |Z_n - \bar{Z}_{\mathcal{T}}|$. Proposition 3 (of Appendix A) implies that these random variables are exchangeable with $|Z_{n_1+1} - \bar{Z}_{\mathcal{T}}|$. Now applying Proposition 1 with

$$W_1 := |Z_{n_1+1} - \bar{Z}_{\mathcal{T}}|, \dots, W_{n-n_1} := |Z_n - \bar{Z}_{\mathcal{T}}|$$

yields the prediction region:

$$\hat{\mathcal{R}}_{n,\alpha}^{\text{split}} := \left\{ z \in \mathbb{R} : |z - \bar{Z}_{\mathcal{T}}| + \xi U_{n+1} \leq W_{i_{[(n-n_1+1)(1-\alpha)]}}^* \right\}. \quad (19)$$

Here $U_{n+1} \sim U[0, 1]$, W^* represents the jittered sequence in Definition 2, and $W_{i_{[(n-n_1+1)(1-\alpha)]}}^*$ represents the $[(n-n_1+1)(1-\alpha)]$ -th largest value among $W_1^*, \dots, W_{n-n_1}^*$.

Proposition 2. *If $Z_1, \dots, Z_{n+1} \in \mathbb{R}$ form a sequence of exchangeable random variables, then for all $n \geq n_1 \geq 1$ and $\alpha \in [0, 1]$,*

$$1 - \alpha \leq \mathbb{P}\left(Z_{n+1} \in \hat{\mathcal{R}}_{n,\alpha}^{\text{split}}\right) \leq 1 - \alpha + \frac{1}{n - n_1 + 1},$$

where the probability extends over all variables, including Z_1, \dots, Z_n used to construct $\hat{\mathcal{R}}_{n,\alpha}^{\text{split}}$.

Although the mean $\bar{Z}_{\mathcal{T}}$ is a natural choice in (19) given the fact that the average is in some sense the best point predictor of a random variable,² Proposition 2 continues to hold true if $\bar{Z}_{\mathcal{T}}$ is generalized by replacing it with $h(Z_1, \dots, Z_{n_1})$ for any function h of Z_1, \dots, Z_{n_1} . This again is because of Proposition 3.

3.3.2 Arbitrary Spaces.

Suppose $Z_1, Z_2, \dots, Z_{n+1} \in \mathcal{Z}$ are exchangeable random variables. Here \mathcal{Z} can be a space of functions or a space of images or a space of documents. For any (non-random) transformation $g : \mathcal{Z} \rightarrow \mathbb{R}$,

$$g(Z_1), g(Z_2), \dots, g(Z_n), g(Z_{n+1}),$$

are real valued exchangeable random variables. Hence Proposition 2 applies to $W_i = g(Z_{n_1+i}), 1 \leq i \leq n - n_1$ and leads to prediction regions

$$\hat{\mathcal{R}}_{n,\alpha}^{\text{split}} := \left\{ z \in \mathcal{Z} : |g(z) - \overline{g(Z)}_{\mathcal{T}}| \leq W_{i_{[(n-n_1+1)(1-\alpha)]}}^* - \xi U_{n+1} \right\}.$$

²The value $a \in \mathbb{R}$ that minimizes $\mathbb{E}[(Z - a)^2]$ (the prediction risk of a) is given by $\mathbb{E}[Z]$.

Here again $W_{\lfloor (n-n_1+1)(1-\alpha) \rfloor}^*$ is the $\lfloor (n-n_1+1)(1-\alpha) \rfloor$ -th largest value of W_1, \dots, W_{n-n_1} . It is noteworthy that for the prediction coverage validity of $\widehat{\mathcal{R}}_{n,\alpha}^{\text{split}}$, the only requirement is that the random variables $g(Z_1), \dots, g(Z_{n+1})$ are exchangeable and this can hold for some data-driven transformations. More precisely, the analyst is allowed to use the training part of the data to construct a transformation $\widehat{f}_{\mathcal{T}}(\cdot)$. Then exchangeability of Z_1, \dots, Z_{n+1} implies that $\widehat{f}_{\mathcal{T}}(Z_{n_1+1}), \dots, \widehat{f}_{\mathcal{T}}(Z_{n+1})$ are exchangeable and can be used for constructing a prediction interval for Z_{n+1} . This leads to the following result proved in Appendix ?? (This result is given the status of a theorem because it shows the generality of conformal prediction.) Recall here that \mathcal{Z} denotes the space in which the random variables Z_1, \dots, Z_n lie.

Theorem 4. For any $\widehat{f}_{\mathcal{T}} : \mathcal{Z} \rightarrow \mathbb{R}$, an arbitrary function depending only on \mathcal{T} , define

$$\widehat{\mathcal{R}}_{n,\alpha}^{\text{split}} := \left\{ z \in \mathcal{Z} : \widehat{f}_{\mathcal{T}}(z) \leq (\widehat{f}_{\mathcal{T}}(Z))_{\lfloor (n-n_1+1)(1-\alpha) \rfloor}^* - \xi U_{n+1} \right\},$$

where $(\widehat{f}_{\mathcal{T}}(Z))_{\lfloor (n-n_1+1)(1-\alpha) \rfloor}^*$ is the $\lfloor (n-n_1+1)(1-\alpha) \rfloor$ -th largest value among the jittered sequence $(\widehat{f}_{\mathcal{T}}(Z_i))^*, i \in \{n_1+1, \dots, n\}$. Then

$$1 - \alpha \leq \mathbb{P}\left(Z_{n+1} \in \widehat{\mathcal{R}}_{n,\alpha}^{\text{split}}\right) \leq 1 - \alpha + \frac{1}{n - n_1 + 1}.$$

In words, Theorem 4 implies that if we find a real-valued transformation based on the training set and construct a prediction region based on the transformed calibration set, then it is also a valid $(1-\alpha)$ prediction, under exchangeability of the set of random variables Z_1, \dots, Z_{n+1} .

Theorem 3.1 is not new and is well-known in the conformal prediction literature. See, for example, Theorem 2 of [Lei et al. \(2018\)](#). In the language of conformal scores, the transformation $\widehat{f}_{\mathcal{T}}(Z_i)$ would be a conformal score corresponding to Z_i . As can be seen from Theorem 3.1, no specific properties of this transformation are required for validity. Because we are constructing a prediction set that contains smaller values of $\widehat{f}_{\mathcal{T}}(\cdot)$, the set $\widehat{\mathcal{R}}_{n,\alpha}^{\text{split}}$ would be sensible if a smaller value of $\widehat{f}_{\mathcal{T}}(z)$ corresponds to z “conforming” with the training data. For instance, the split conformal prediction set (19) is constructed based on the conformal score $\widehat{f}_{\mathcal{T}}(z) = |z - \bar{Z}_{\mathcal{T}}|$. A smaller value here means z is close to the “center” of training data — conforming. A larger value means z is away from the training data — not conforming.

Pseudocode 2: Computationally, Theorem 4 works in practice as follows:

1. Split the data Z_1, \dots, Z_n into two parts: \mathcal{T} and \mathcal{C} as in (18).
2. Based on the training data \mathcal{T} , find a transformation $\widehat{f}_{\mathcal{T}} : \mathcal{Z} \rightarrow \mathbb{R}$. See Section 3.3.3 for some examples of $\widehat{f}_{\mathcal{T}}$.
3. Take $\xi = 10^{-8}$ and generate U_1, \dots, U_{n+1} from $\text{Unif}[-1, 1]$ distribution. Define the jittered sequence (based on the elements in the calibration set) as $W_i^* := \widehat{f}_{\mathcal{T}}(Z_i) + \xi U_i$, $n_1 + 1 \leq i \leq n$.
4. Sort the observations W_i as $W_{(1)}^* \leq W_{(2)}^* \leq \dots \leq W_{(n-n_1+1)}^*$.
5. Report the prediction region

$$\left\{ z \in \mathcal{Z} : \widehat{f}_{\mathcal{T}}(z) \leq W_{\lfloor (n-n_1+1)(1-\alpha) \rfloor}^* - \xi U_{n+1} \right\}.$$

3.3.3 Some Concrete Examples.

In the following, we present a few concrete examples/applications of Theorem 4. Note that the problem is still prediction: given Z_1, \dots, Z_n , we want to predict Z_{n+1} . In the context of Theorem 3.1, we are essentially doing this based on $\hat{f}_{\mathcal{T}}(Z_{n_1+1}), \dots, \hat{f}_{\mathcal{T}}(Z_n)$. In the following examples, we describe the construction of some useful $\hat{f}_{\mathcal{T}}(\cdot)$. The construction of the prediction region is based on Theorem 4. Recall here that \mathcal{Z} denotes the space in which the random variables Z_1, \dots, Z_n lie.

1. **Norm-ball around the Mean:** Suppose $\mathcal{Z} = \mathbb{R}^d$ and let $\bar{Z}_{\mathcal{T}}$ be the average of observations in \mathcal{T} , the training set. Take $\hat{f}_{\mathcal{T}}(Z_i) = \|Z_i - \bar{Z}_{\mathcal{T}}\|$ for $i \in \{n_1 + 1, \dots, n\}$. Here $\|\cdot\|$ can be any semi-norm in \mathbb{R}^d ; for instance the Euclidean norm or the Manhattan norm or the ℓ_p -norm or even the absolute value of a single coordinate. In the multivariate case, calculating the norm may not be meaningful if different coordinates of Z have different units. If this is the case, then one can compute the norm of “whitened” vectors. More precisely, let $\hat{\Sigma}_{\mathcal{T}}$ represent the sample covariance based on \mathcal{T} and take $\hat{f}_{\mathcal{T}}(Z_i) := \|\hat{\Sigma}_{\mathcal{T}}^{-1/2}(Z_i - \bar{Z}_{\mathcal{T}})\|$. If $\hat{\Sigma}_{\mathcal{T}}$ is not invertible (which can happen if dimension is larger than n_1), then take

$$\hat{f}_{\mathcal{T}}(Z_i) := \|\text{diag}(\hat{\Sigma}_{\mathcal{T}})^{-1/2}(Z_i - \bar{Z}_{\mathcal{T}})\|.$$

Here $\text{diag}(\hat{\Sigma}_{\mathcal{T}})$ is the diagonal matrix corresponding to $\hat{\Sigma}_{\mathcal{T}}$.

2. **Principal Component Analysis (PCA):** In the previous example, the function $\hat{f}_{\mathcal{T}}$ uses all the coordinates of Z_i with no regard to the coordinates that matter more. For cases where the distribution of Z_i 's is supported on a low-dimensional manifold, this might be wasteful. One way to account for low-dimensionality is by using a dimension reduction technique on \mathcal{T} . For one concrete example, when $\mathcal{Z} = \mathbb{R}^d$, apply PCA on $\mathcal{T} = \{Z_1, \dots, Z_{n_1}\}$ and fix $1 \leq \hat{k} \leq d$ to be the number of principal components (PCs) to be used. (This choice of \hat{k} can be based on any rule as long as it depends only on \mathcal{T} .) Let the first \hat{k} PCs be written into a matrix $\Pi \in \mathbb{R}^{\hat{k} \times d}$ and consider

$$\hat{f}_{\mathcal{T}}(Z_i) := \|\text{diag}(\hat{\Sigma}_{\Pi, \mathcal{T}})^{-1/2}(\Pi Z_i - (\overline{\Pi Z})_{\mathcal{T}})\|.$$

Here $\hat{\Sigma}_{\Pi, \mathcal{T}}$ and $(\overline{\Pi Z})_{\mathcal{T}}$ represent the sample covariance matrix and sample average of $\Pi Z_1, \dots, \Pi Z_{n_1}$. The semi-norm $\|\cdot\|$ above is arbitrary as in the previous example. Similar to the previous example, PCA is not special here and any of the many existing dimension reduction (linear or non-linear) techniques (Cunningham, 2008; Xie et al., 2017; Sorzano et al., 2014; Nguyen and Holmes, 2019; Hinton and Salakhutdinov, 2006; Wang et al., 2014; Tenenbaum et al., 2000; Silva and Tenenbaum, 2003) can be used to get ΠZ_i . In the context of functional data, Lei et al. (2015) propose a few examples of Π .

3. **Level Sets:** The prediction regions mentioned in the discussions before are all convex sets; in \mathbb{R} , these are intervals. These may, however, not be the optimal ones. For example, if the true distribution of Z_i is a mixture of $N(3, 1)$ and $N(7, 1)$, then the optimal prediction region is a union of two intervals centered at the two modes; see Figure 1. This is similar to the definition of high density regions, popular in Bayesian statistics; see Hyndman (1996) for details. If the density of Z_i is $p(\cdot)$, then the (oracle) optimal prediction region is given by

$$\mathcal{R}^{\text{opt}} := \{z \in \mathcal{Z} : p(z) \geq t_{\alpha}\}, \quad (20)$$

where t_{α} is the largest t solving $\int_{z:p(z) \geq t} p(z) dz \geq 1 - \alpha$. Sets of this type are called level sets. Of course, in practice we do not know the density p and it might not even exist (with respect to the Lebesgue




Figure 1: Illustration of level sets with the density of mixture of Gaussians centered at 3 and 7.

measure). One way to imitate this optimal prediction region is by taking

$$\hat{f}_{\mathcal{T}}(Z_i) := 1/\hat{p}_{\mathcal{T}}(Z_i), \quad (21)$$

where $\hat{p}_{\mathcal{T}}(\cdot)$ is an estimate of the density based on Z_1, \dots, Z_{n_1} . Combining with the previous example, the density estimator here can be performed after an initial dimension reduction step. Furthermore, any of the existing density estimation methodologies (along with tuning parameter selection methods) can be used. The validity guarantees of conformal prediction does not depend on the estimation accuracy of the dimension reduction or density estimation methods. It can be proved that (21) along with Theorem 4 leads to prediction regions that “converge” to the optimal one (20); see [Lei et al. \(2013\)](#) for details.

In some of the examples above, we have made an assumption that \mathcal{Z} (the space in which Z_1, \dots, Z_n lie) is the Euclidean space \mathbb{R}^d . Some real data examples, such as image classification or topic modeling or text mining, do not satisfy this assumption readily. In all these examples, however, classical machine learning algorithms first convert the image or text data into a high-dimensional real-valued vector. Dimension is not an issue for conformal prediction because the validity is finite sample.

3.4 Conformal Prediction for Regression

In previous subsections, we have discussed the problem of prediction with no side information, i.e., we do not have any information at the future random variable. Most classical prediction algorithms in machine learning and statistics have covariate information for future random variable and the response is to be predicted. This forms one of the most interesting applications of conformal prediction. Here the information in each observation is in two parts: covariates or predictors or features (X) and the response or class (Y). Let $Z_1 = (X_1, Y_1), \dots, Z_n = (X_n, Y_n)$ be n observations from a space $\mathcal{X} \times \mathcal{Y}$ and we want to predict Y_{n+1} for $X_{n+1} \in \mathcal{X}$, whenever (X_{n+1}, Y_{n+1}) is exchangeable with $(X_i, Y_i), 1 \leq i \leq n$. Formally, the goal is to construct $\hat{\mathcal{R}}_{n,\alpha}(X_{n+1})$ such that

$$\mathbb{P}\left(Y_{n+1} \in \hat{\mathcal{R}}_{n,\alpha}(X_{n+1})\right) \geq 1 - \alpha. \quad (22)$$

The probability on the left hand side is with respect to (X_{n+1}, Y_{n+1}) and also with respect to $(X_i, Y_i), i \in [n]$.

Marginal versus Conditional Coverage. Mathematically, there is nothing wrong with the formulation (22), however, it is notationally misleading because of the alternative goal

$$\mathbb{P}\left(Y_{n+1} \in \widehat{\mathcal{R}}_{n,\alpha}(X_{n+1}) \mid X_{n+1} = x\right) \geq 1 - \alpha, \quad (23)$$

for all $x \in \mathcal{X}$. Formulation (22) provides a prediction region that covers Y_{n+1} whenever (X_{n+1}, Y_{n+1}) comes from the same distribution. Formulation (23), however, requires a prediction region that covers Y_{n+1} whatever the value of X_{n+1} is. To understand the philosophical difference between these two goals, consider the following scenario. Suppose we have bivariate classification data with body mass index (BMI) as the covariate and indicator for the presence of cancer cells as the response. The goal (23) provides a region such that whoever the next patient is, his/her response would lie in the constructed region with probability of at least $1 - \alpha$. If we get 100 new patients, then (23) implies that among the patients with BMI= 15 (say), the proportion of patients for which the true response lies in the constructed prediction set is about $1 - \alpha$ and the same is true among the patients with BMI= 15.7, 16, and so on (any real number). On the other hand, the goal (22) cannot guarantee this but only implies that if we have 100 new patients and we give a region for each patient, then out of these 100 patients, for about $(1 - \alpha)$ proportion of them the true response lies in their corresponding region. Importantly, among the patients with BMI= 15 (say), there is no specific control on the proportion of patients for which the true response lies in the constructed prediction set.

It is easy to show that (23) implies (22). It turns out that (23) is too ambitious a goal in that it cannot be attained, non-trivially, in finite samples in a distribution-free setting; see Balasubramanian et al. (2014, Section 2.6.1) and Foygel Barber et al. (2019). The latter reference discusses alternative conditional goals that are attainable sensibly. In this section, we will restrict attention to marginal validity as in (22) and refer to Foygel Barber et al. (2019) for details on conditional validity (23).

Cross-sectional Conformal Method. One simple way to attain the guarantee (22) is as follows. Based on $(X_1, Y_1), \dots, (X_n, Y_n) \in \mathcal{X} \times \mathcal{Y}$, construct a prediction region $\widehat{\mathcal{R}}_{n,\alpha} \subseteq \mathcal{X} \times \mathcal{Y}$. Any of the constructions mentioned in Subsection 3.3 can be used. These regions satisfy

$$\mathbb{P}\left((X_{n+1}, Y_{n+1}) \in \widehat{\mathcal{R}}_{n,\alpha}\right) \geq 1 - \alpha, \quad (24)$$

and hence

$$\mathbb{P}\left(Y_{n+1} \in \widehat{\mathcal{R}}_{n,\alpha}(X_{n+1})\right) \geq 1 - \alpha, \quad (25)$$

where

$$\widehat{\mathcal{R}}_{n,\alpha}(X_{n+1}) := \{y \in \mathcal{Y} : (X_{n+1}, y) \in \widehat{\mathcal{R}}_{n,\alpha}\}.$$

This set is a cross-section of $\widehat{\mathcal{R}}_{n,\alpha}$ at X_{n+1} . See Figure 2 for an illustration of this method of attaining (22). In light of the discussion regarding conditional and marginal validity, we prefer writing (24) instead of (25) for clarity. The notation (24) makes it clear that the construction of the region has nothing to do with the realized value of X_{n+1} . This also shows that there is no formal difference between conformal prediction for regression and conformal prediction for general spaces. The only difference is in the construction of the transformation; now the transformation focuses on how well the response value conforms with the training data for the realized value of X_{n+1} .




Figure 2: Illustration of Cross-sectional Conformal Prediction for Regression and Classification.

Figure 2 shows a commonly mentioned disadvantage of this cross-sectional method. For some values of X_{n+1} , $\widehat{\mathcal{R}}_{n,\alpha}(X_{n+1})$ can be empty. For example, in Figure 2, $\widehat{\mathcal{R}}_{n,\alpha}(x) = \emptyset$ and $\widehat{\mathcal{R}}_{n,\alpha}(x') \neq \emptyset$. From a different view point, this is more of an advantage than a disadvantage because the validity guarantee (22) requires X_{n+1} from the same distribution as the X 's in the training data. Hence, an empty cross-section at x as in Figure 2 actually informs the practitioner that the value x is not likely, given the training values of X 's. This might be useful in raising a red flag when the practitioner is about to do extrapolation. For determining extrapolation, it might be better to first perform a dimension reduction that highlights the information in X pertaining to predicting Y ; this dimension reduction map can be based on the training split of the data.

Conformal Regions with Focus on Response. The cross-sectional conformal prediction gives equal weight to the covariates and the response in that the prediction region $\widehat{\mathcal{R}}_{n,\alpha}$ does not focus more on either. In the regression or classification context, one might want to focus more on the response and ignore prediction for the covariates. Ignoring covariates for prediction means that the coverage guarantee for covariates can be 1, or in other words, the covariate cross-section of the region $\widehat{\mathcal{R}}_{n,\alpha}$ is the whole space of covariates. We will now provide ways to apply Theorem 4 for prediction in regression and classification settings with the motivation above. This means that we will provide some concrete ways of designing $\widehat{f}_{\mathcal{T}}$ for the regression and classification framework.

- *Conditional Mean Estimation:* Based on the training data $\{(X_i, Y_i) : 1 \leq i \leq n_1\} \subseteq \mathcal{X} \times \mathbb{R}$, estimate the conditional mean $\mathbb{E}[Y|X = x]$. Let the estimate be $\widehat{\mu}_{\mathcal{T}}(\cdot)$ based on the training data \mathcal{T} . Define, for $Z_i = (X_i, Y_i)$,

$$\widehat{f}_{\mathcal{T}}(Z_i) := |Y_i - \widehat{\mu}_{\mathcal{T}}(X_i)|, \quad n_1 + 1 \leq i \leq n + 1.$$

This yields a prediction region of the form $[\widehat{\mu}_{\mathcal{T}}(X_{n+1}) - t, \widehat{\mu}_{\mathcal{T}}(X_{n+1}) + t]$ for some t . The validity guarantee holds true irrespective of what $\widehat{\mu}_{\mathcal{T}}(\cdot)$ is. Any algorithm can be used to get an estimator; it may not even be consistent for $\mathbb{E}[Y|X = \cdot]$. Unlike the cross-sectional method, this function leads to a non-empty prediction region for all realized values of X_{n+1} and as mentioned above, this can be a disadvantage. There exist

many variations of these regions including those based on conditional variance normalization (Lei and Wasserman, 2014; Lei et al., 2018), conditional quantile estimator (Romano et al., 2019; Kivaranovic et al., 2020; Sesia and Candès, 2020), and conditional distributions (Chernozhukov et al., 2019; Izbicki et al., 2019; Gupta et al., 2021). These variations aim to get approximate conditional coverage and have width adapting to the conditional heteroscedasticity. See Corollary 1 of Sesia and Candès (2020) for details.

- *Conditional Probability Density Estimation:* Although conditional validity (23) is non-trivially impossible in a distribution-free setting, one can still construct a marginally valid prediction region that can asymptotically attain the conditional validity guarantee. Similar to the optimal prediction region (20) for the whole vector $Z_{n+1} = (X_{n+1}, Y_{n+1})$, the optimal conditional prediction region for Y_{n+1} given X_{n+1} is given by

$$\mathcal{R}_\alpha^{\text{opt}}(x) := \{y \in \mathcal{Y} : p(y|x) \geq t_\alpha(x)\}, \quad (26)$$

where $p(y|x)$ is the conditional probability density function of Y given $X = x$ and $t_\alpha(x)$ is the largest $t \geq 0$ such that

$$\int_{y:p(y|x) \geq t} p(y|x) dy \geq 1 - \alpha.$$

An imitation of this region is given by replacing $p(y|x)$ by an estimator $\hat{p}_\mathcal{T}(y|x)$ based on the training data $(X_i, Y_i), 1 \leq i \leq n_1$. This replacement does not guarantee any validity and any such guarantees depend on the accuracy of $\hat{p}_\mathcal{T}(y|x)$ for $p(y|x)$. To guarantee validity, define for $z = (x, y)$,

$$\hat{f}_\mathcal{T}(z) := 1 - \inf\{\alpha \in [0, 1] : y \in \hat{\mathcal{R}}_{\mathcal{T}, \alpha}^*(x)\}, \quad (27)$$

where

$$\hat{\mathcal{R}}_{\mathcal{T}, \alpha}^*(x) := \{y \in \mathcal{Y} : \hat{p}_\mathcal{T}(y|x) \geq \hat{t}_{\mathcal{T}, \alpha}(x)\}.$$

Here $\hat{t}_{\mathcal{T}, \alpha}(x)$ is the largest $t > 0$ such that $\int_{y:\hat{p}_\mathcal{T}(y|x) \geq t} \hat{p}_\mathcal{T}(y|x) dy \geq 1 - \alpha$. Now Theorem 4 with $\hat{f}_\mathcal{T}(\cdot)$ in (27) leads to a prediction region that is guaranteed to satisfy (22). We stress once again that although this region is imitating the optimal conditional prediction region (26), it does *not* have a finite sample conditional guarantee. Because of the imitation, it is expected that the region from Theorem 4 based on (27) will asymptotically satisfy the conditional guarantee (23). Similar construction of conformal prediction sets for regression and classification has also been discussed in Izbicki et al. (2019) and Gupta et al. (2021). Unlike the conditional mean estimation based prediction region, the region from (27) is sensible for classification and regression alike. For classification, the prediction set using $\hat{f}_\mathcal{T}(\cdot)$ in (27) gathers those classes with higher estimated probabilities from the classifier. See Romano et al. (2020) and Kuchibhotla and Berk (2021) for a similar approach for classification.

3.5 Testing Interpretation of Conformal Prediction and other Variants

The full conformal method, as mentioned before, is computationally prohibitive, in general. But this method makes the full use of the data for prediction purposes. The split conformal method although computationally efficient uses some part of the data for training and only some part of the data for prediction calibration purposes. For this reason, many authors (Balasubramanian et al., 2014; Lei et al., 2018; Barber et al., 2021) have argued that split conformal method could incur statistical inefficiency due to this splitting. Several methods have been proposed to make better use of the data; see, for example, Carlsson et al. (2014); Vovk

(2015); Linusson et al. (2017); Vovk and Wang (2019); Lei et al. (2018); Barber et al. (2021); Gupta et al. (2021); Kim et al. (2020). In this section, we describe these variants briefly using the hypothesis testing interpretation of conformal prediction regions.

Recall that the goal of conformal prediction is to construct a set $\widehat{\mathcal{R}}_{n,\alpha}$ based on W_1, \dots, W_n such that

$$\mathbb{P}\left(W_{n+1} \in \widehat{\mathcal{R}}_{n,\alpha}\right) \geq 1 - \alpha.$$

Following the duality of testing and confidence regions, we can formally think of testing the hypothesis $H_0 : W_{n+1} = w$ for some value w . (This is not a traditional hypothesis because it relates to a random variable W_{n+1} .) Based on a test for H_0 , a valid prediction set can be constructed by collecting all the w 's for which $H_0 : W_{n+1} = w$ is not rejected. We will now define a p -value and the corresponding test when W_1, \dots, W_n, W_{n+1} are real-valued exchangeable random variables. The case of arbitrary spaces can be dealt with similarly using transformations. Define the p -value

$$P_w := 1 - \frac{\text{rank}(w; \{W_1, W_2, \dots, W_n, w\}) + 1}{n + 1}. \quad (28)$$

From Corollary 1, it follows that $\mathbb{P}(P_{W_{n+1}} \leq \alpha) \leq \alpha$ for all $\alpha \in [0, 1]$. In other words, P_w is a valid p -value under $H_0 : W_{n+1} = w$. Hence, the region $\{w : P_w > \alpha\}$ is a valid $(1 - \alpha)$ prediction region. This p -value interpretation of conformal prediction method was mentioned in Shafer and Vovk (2008, Section 4.2) and Lei et al. (2013, Section 2), among others. For an interesting modification of these p -values in relation to conformal prediction, see Carlsson et al. (2015).

We are now ready to discuss the conformal prediction methods that lie in between the split and full conformal methods.

- Jackknife and CV methods (Vovk, 2015; Barber et al., 2021) are based on the idea of splitting the data into multiple disjoint folds (instead of just 2) and then combine the ranks or transformed variables in some way. To elaborate, we briefly describe the jackknife+ method from Barber et al. (2021). Suppose $Z_1, \dots, Z_n, Z_{n+1} \in \mathcal{Z}$ are exchangeable random variables with the goal of predicting Z_{n+1} . Let $\widehat{f}_{-(i,j)} : \mathcal{Z} \rightarrow \mathbb{R}$ be a permutation invariant transformation computed based on $\{Z_1, \dots, Z_{n+1}\} \setminus \{Z_i, Z_j\}$. In regression data with $Z_i = (X_i, Y_i)$, for example, $\widehat{f}_{-(i,j)}(z)$ can be $|y - \widehat{\mu}_{-(i,j)}(x)|$, where $\widehat{\mu}_{-(i,j)}(\cdot)$ is a regression function computed based on $\{Z_1, \dots, Z_{n+1}\} \setminus \{Z_i, Z_j\}$. Define the prediction set for Z_{n+1} as

$$\widehat{\mathcal{R}}_{n,\alpha}^{\text{Jack}} := \left\{ z \in \mathcal{Z} : \sum_{j=1}^n \mathbb{1} \left\{ \widehat{f}_{-(n+1,i)}(z) > \widehat{f}_{-(i,n+1)}(Z_i) \right\} < (1 - \alpha)(n + 1) \right\}.$$

Note that $\widehat{f}_{-(i,n+1)}(\cdot) \equiv \widehat{f}_{-(n+1,i)}(\cdot)$ and that these transformations can be computed without the knowledge of Z_{n+1} . Further, $\widehat{f}_{-(i,n+1)}(\cdot)$ is a leave-one-out transformation on the data Z_1, \dots, Z_n . Theorem 1 of Barber et al. (2021) can be used to prove that $\mathbb{P}(Z_{n+1} \in \widehat{\mathcal{R}}_{n,\alpha}^{\text{Jack}}) \geq 1 - 2\alpha$, although the theorem is only stated for regression data with absolute residual. The proof of Theorem 1 of Barber et al. (2021) hinges on the fact that

$$G \begin{pmatrix} Z_1 \\ \vdots \\ Z_{n+1} \end{pmatrix} := \begin{pmatrix} W_1 \\ \vdots \\ W_{n+1} \end{pmatrix} = \begin{pmatrix} \sum_{j=2}^{n+1} \mathbb{1} \{ \widehat{f}_{-(1,j)}(Z_1) > \widehat{f}_{-(j,1)}(Z_j) \} \\ \vdots \\ \sum_{j=1}^n \mathbb{1} \{ \widehat{f}_{-(n+1,j)}(Z_{n+1}) > \widehat{f}_{-(j,n+1)}(Z_j) \} \end{pmatrix}, \quad (29)$$

is an exchangeability preserving transformation; this can be verified using Theorem 3. Step 1 in Section 6 of Barber et al. (2021) shows that the number of coordinates in the right hand side of (29) that are

larger than or equal to $(1 - \alpha)(n + 1)$ is bounded by $2\alpha(n + 1)$; this is a deterministic inequality and does not require exchangeability of Z_1, \dots, Z_{n+1} . Using the fact that $G(\cdot)$ is exchangeability preserving, we get that

$$\begin{aligned} \mathbb{P}(W_{n+1} \geq (1 - \alpha)(n + 1)) &= \frac{1}{n + 1} \sum_{j=1}^{n+1} \mathbb{P}(W_j \geq (1 - \alpha)(n + 1)) \\ &= \mathbb{E} \left[\frac{\sum_{j=1}^n \mathbb{1}\{W_j \geq (1 - \alpha)(n + 1)\}}{n + 1} \right] \leq 2\alpha. \end{aligned}$$

Here the inequality above follows from Step 1 in Section 6 of [Barber et al. \(2021\)](#). The CV+ method is defined similarly where instead of leave-one-out, one uses a leave-a-fold-out; see Section 3 of [Barber et al. \(2021\)](#) for details. Also, see [Solari and Djordjilović \(2021\)](#) for a different argument.

- Subsampling or repeated split methods ([Carlsson et al., 2014](#); [Lei et al., 2018](#); [Gupta et al., 2021](#)) repeat the split conformal method several times on the data and combine the resulting prediction sets in some way. To elaborate, we briefly discuss the subsampling or Bonferroni method discussed in [Lei et al. \(2018\)](#) and [Gupta et al. \(2021\)](#). Recall that the split conformal method can be interpreted in terms of a retention region from a p -value (28). If we repeat the splitting process on the data K times, then we get K p -values $P_{Z_{n+1}}^{(1)}, \dots, P_{Z_{n+1}}^{(K)}$. It is very important to observe that these are dependent p -values, dependent through Z_{n+1} . This implies that $K \min_{1 \leq k \leq K} P_{Z_{n+1}}^{(k)}$ is also a valid p -value:

$$\mathbb{P} \left(K \min_{1 \leq k \leq K} P_{Z_{n+1}}^{(k)} \leq \alpha \right) \leq \mathbb{P} \left(\bigcup_{k=1}^K \left\{ P_{Z_{n+1}}^{(k)} \leq \frac{\alpha}{K} \right\} \right) \leq \sum_{k=1}^K \frac{\alpha}{K} \leq \alpha.$$

Hence, we get that $\{z : P_z^{(k)} > \alpha/K \text{ for all } 1 \leq k \leq K\}$ is a valid $(1 - \alpha)$ prediction region for Z_{n+1} . The combination of p -values above is the Bonferroni correction from the multiple testing literature. One can use other combinations of p -values such as twice the arithmetic or geometric mean, and so on; see [Vovk and Wang \(2019\)](#) for more examples. The use of multiple testing for other conformal methods can be seen in [Lei et al. \(2018, Section 2.3\)](#), [Vovk and Wang \(2019, Section 4\)](#), [Gupta et al. \(2021, Appendix B & C\)](#).

Although these variants make better use of the full data, there is a clear lack of great advantage of these complicated methods in performance per computational cost in comparison to the split conformal method. Firstly, these variants require computing the transformation multiple times. Secondly, they can be both conservative and anti-conservative in practice. Finally, the volume of the resulting prediction regions can also be larger than that of the split method. See [Barber et al. \(2021, Figure 2\)](#) and [Gupta et al. \(2021, Table 2\)](#) for some comparisons. This is not to say that split conformal is always the best. Conformal methods that make use of (close to) full data perform best when the dimension of the training algorithm is close to the sample size; for example, fitting linear regression with n observations and d ($\approx n$) covariates ([Barber et al., 2021, Figure 2](#)). With machine learning algorithms such as random forests that automatically yield several training and calibration sets, the out-of-bag or aggregated conformal methods ([Kim et al., 2020](#); [Gupta et al., 2021](#)) can yield better performance in comparison to split conformal without increasing the computational cost.

4 Nonparametric Rank Tests

Testing equality of distributions and independence of random vectors are two of the most fundamental problems in statistics. In the following two sections, we discuss each of these problems and detail the implications of exchangeability. We have seen in the previous section that a basic prediction interval for real-valued random variables can be used to construct prediction sets for random variables in arbitrary spaces by a data-driven transformation. In this section, we show that the classical non-parametric rank tests, defined for real-valued random variables, can also be used for tests for random variables in arbitrary spaces by a data-driven transformation. All this is made possible by exchangeability and its consequences. It should be mentioned that this dimension reduction idea in rank tests is not new and has been discussed in some works such as [Matthews and Taylor \(1996\)](#) and [Friedman \(2003\)](#).

A brief description of the tests is as follows:

1. **Equality of Distributions:** Given two datasets, one might want to test if the datasets are obtained from the same distribution. More formally, given n i.i.d. observations X_1, \dots, X_n from P_X and m i.i.d. observations Y_1, \dots, Y_m from P_Y with both sets of observations independent, we want to test

$$H_0 : P_X = P_Y \quad \text{versus} \quad H_1 : P_X \neq P_Y. \quad (30)$$

This is also known as a two-sample testing problem and has numerous applications in pharmaceutical studies ([Farris and Schopflocher, 1999](#)), causal inference ([Folkes et al., 1987](#)), remote sensing ([Conradsen et al., 2003](#)), and econometrics ([Mayer, 1975](#)).

2. **Independence:** Given n i.i.d. paired observations $(X_i, Y_i), 1 \leq i \leq n$, one might want to test if X_i is independent of Y_i . Formally, suppose the distribution of (X_i, Y_i) is P , the marginal distribution of X_i is P_X , and the marginal distribution of Y_i is P_Y . Then we want to test

$$H_0 : P = P_X \otimes P_Y \quad \text{versus} \quad H_1 : P \neq P_X \otimes P_Y.$$

Independence testing has found applications in statistical genetics ([Liu et al., 2010](#)), marketing and finance ([Grover and Dillon, 1985](#)), survival analysis ([Martin and Betensky, 2005](#)), and ecological risk assessment ([Dishion et al., 1999](#)). Furthermore, the test for equality of distributions can be formulated as a test for independence by defining Y as a binary random variable labeling the sample to which X belongs to; see, e.g., [Heller et al. \(2016, Section 1\)](#).

Unlike in the case of conformal prediction, in this section, we will assume that the underlying observations are independent and identically distributed. The reason for this change in assumption can be understood as follows. Consider the testing problem (30) of equality of distributions. Under the null hypothesis H_0 , we want the data obtained by combining $\{X_1, \dots, X_n\}$ and $\{Y_1, \dots, Y_m\}$ to form an exchangeable sequence. For this exchangeability, under the independence of $\{X_1, \dots, X_n\}$ and $\{Y_1, \dots, Y_m\}$, the i.i.d. assumption seems the most sensible.

4.1 Testing Equality of Distributions

Suppose X_1, \dots, X_n are independent and identically distributed random variables from a probability measure P_X and Y_1, \dots, Y_m are independent and identically distributed random variables from a probability measure P_Y . Random variables X_1, \dots, X_n are independent of Y_1, \dots, Y_m . The hypothesis to test is

$$H_0 : P_X = P_Y \quad \text{versus} \quad H_1 : P_X \neq P_Y.$$

There exist numerous tests for this hypothesis. We refer to reader to [Bhattacharya \(2020, 2019\)](#) and [Deb and Sen \(2021\)](#) for an overview of the existing literature. Wilcoxon rank sum test ([Hollander et al., 2013](#), Section 4.1) is one of the classical rank tests for this hypothesis when the observations are real-valued. Let $\mathcal{C}_{n+m} := \{Z_1, \dots, Z_{n+m}\}$ represent the random variables X_1, \dots, X_n and Y_1, \dots, Y_m put together. Under H_0 , the collection of random variables in \mathcal{C}_{n+m} are independent and identically distributed and hence, by [Theorem 2](#), $\text{rank}(Z_1; \mathcal{C}_{n+m}), \dots, \text{rank}(Z_{n+m}; \mathcal{C}_{n+m})$ are distributed uniformly over all permutations of $[n+m]$; recall $[n+m] = \{1, 2, \dots, n+m\}$. The Wilcoxon rank sum test statistic is given by

$$T_n := \sum_{i=1}^n \text{rank}(Z_i; \mathcal{C}_{n+m}),$$

the sum of ranks of X_1, \dots, X_n among \mathcal{C}_{n+m} . Recall the definition of rank from [Definition 1](#). From this definition, it follows that the distribution of T_n does not depend on the true distributions P_X and P_Y under H_0 .

Using this, we now extend the Wilcoxon test to random variables taking values in arbitrary space \mathcal{Z} . Let $\hat{f}_{n+m} : \mathcal{Z} \rightarrow \mathbb{R}$ be any transformation that depends *permutation invariantly* on $Z_1, \dots, Z_{n+m} \in \mathcal{C}_{n+m}$. This means that if we write

$$\hat{f}_{n+m}(z) = \hat{f}_{n+m}(z; Z_1, \dots, Z_{n+m}),$$

then permutation invariance means for any $z \in \mathcal{Z}$ and any $\pi : [n+m] \rightarrow [n+m]$,

$$\hat{f}_{n+m}(z; Z_1, \dots, Z_{n+m}) = \hat{f}_{n+m}(z; Z_{\pi(1)}, \dots, Z_{\pi(n+m)}).$$

[Proposition 4](#) (in [Appendix A](#)) shows that $W_1 := \hat{f}_{n+m}(Z_1), \dots, W_{n+m} := \hat{f}_{n+m}(Z_{n+m})$ are exchangeable and hence

$$T_n^{\text{exch}} := \sum_{i=1}^n \text{rank}(W_i; \{W_1, \dots, W_{n+m}\}),$$

the sum of ranks of W_1, \dots, W_n among the collection $\{W_1, \dots, W_{n+m}\}$, is also distribution-free and has the same distribution as T_n under H_0 . This proves the following result.

Theorem 5. *Suppose X_1, \dots, X_n are independent and identically distributed random variables from a probability measure P_X supported on \mathcal{Z} and Y_1, \dots, Y_m are independent and identically distributed random variables from a probability measure P_Y supported also on \mathcal{Z} . Then for any transformation \hat{f}_{n+m} depending permutation invariantly on $X_1, \dots, X_n, Y_1, \dots, Y_m$,*

$$T_n^{\text{exch}} := \sum_{i=1}^n \text{rank}(\hat{f}_{n+m}(X_i); \{\hat{f}_{n+m}(X_1), \dots, \hat{f}_{n+m}(Y_m)\}),$$

is distribution-free under H_0 and matches the null distribution of the Wilcoxon rank sum statistic.

The main conclusion of the theorem is that any permutation invariant transformation \hat{f}_{n+m} obtained from the data will lead to a distribution-free test with a finite sample control of the type I error, irrespective of the domain of the data. This is the analogy with the full conformal prediction method where for any permutation invariant transformation [\(14\)](#), the prediction region has a finite sample control of the coverage. [Theorem 5](#) is obvious if \hat{f}_{n+m} does not depend on the data Z_1, \dots, Z_{n+m} . Such application of rank tests after dimension reduction is well-known. The main novelty of [Theorem 5](#) is that the transformation \hat{f}_{n+m} can depend on the data. Although relatively easy to derive based on [Theorem 3](#), [Theorem 5](#) is new to the best of our knowledge.

Pseudocode 3: Computationally, Theorem 5 works as follows:

1. Rename the data as

$$Z_1 = X_1, \dots, Z_n = X_n, Z_{n+1} = Y_1, \dots, Z_{n+m} = Y_m.$$

2. Find a permutation invariant transformation \hat{f}_{n+m} based on Z_1, \dots, Z_{n+m} . This can be any cluster revealing transformation; see Section 4.1.1 below.
3. Take $\xi = 10^{-8}$ and compute the ranks of the first n elements $\hat{f}_{n+m}(Z_1), \dots, \hat{f}_{n+m}(Z_n)$ among $\hat{f}_{n+m}(Z_1), \dots, \hat{f}_{n+m}(Z_{n+m})$.
4. Sum the ranks and compute T_n^{exch} . Reject the null hypothesis H_0 if T_n^{exch} deviates from its expected value $n(n+m+1)/2$. The critical value here is the same as that of Wilcoxon rank sum test; see [Hollander et al. \(2013, Section 4.1\)](#) and [Wilcoxon \(1945\)](#) for details.

4.1.1 Some Concrete Examples.

In the following, we discuss a few examples of \hat{f}_{n+m} in Theorem 5. The most generic application is based on applying any of the many unsupervised clustering algorithms.

1. **k -means Clustering:** Suppose Z_1, \dots, Z_{n+m} are elements of a space \mathcal{Z} with a quasi-norm $\|\cdot\|$.³ Fix $k \geq 1$, the number of clusters and apply the k -means clustering algorithm on the data Z_1, \dots, Z_{n+m} , that is, find $\hat{c}_1, \dots, \hat{c}_k \in \mathcal{Z}$ such that

$$(\hat{c}_1, \dots, \hat{c}_k) := \arg \min_{c_1, \dots, c_k \in \mathcal{Z}} \sum_{i=1}^{n+m} \min_{1 \leq j \leq k} \|Z_i - c_j\|^2. \quad (31)$$

Note that the minimizer can at best be unique up to permutations, i.e., if $(\hat{c}_1, \dots, \hat{c}_k)$ is a minimizer, then $(\hat{c}_{\pi(1)}, \dots, \hat{c}_{\pi(k)})$ is also a minimizer for arbitrary permutation $\pi : [k] \rightarrow [k]$. For our purposes, we choose (arbitrarily) and fix a minimizer. It is easy to verify that $(\hat{c}_1, \dots, \hat{c}_k)$ is a permutation invariant function of $\{Z_1, \dots, Z_{n+m}\}$. One transformation \hat{f}_{n+m} based on this clustering method is given by

$$\hat{f}_{n+m}(Z_i) := \|Z_i - \hat{c}_1\| - \min_{1 \leq j \leq k} \|Z_i - \hat{c}_j\|.$$

To gain intuition for this particular permutation invariant transformation, suppose H_1 is true (that is, $P_X \neq P_Y$) and $P_X = N(\mu_X, \sigma^2)$ and $P_Y = N(\mu_Y, \sigma^2)$. If we perform k -means clustering with $k = 2$, then asymptotically $\hat{c}_1 = \mu_X$ and $\hat{c}_2 = \mu_Y$ (up to labels) and hence, asymptotically,

$$\begin{aligned} \min_{1 \leq j \leq k} \|Z_i - \hat{c}_j\| &= \|Z_i - \hat{c}_1\|, \quad \Rightarrow \quad \hat{f}_{n+m}(Z_i) = 0, \quad \text{for } 1 \leq i \leq m, \\ \min_{1 \leq j \leq k} \|Z_i - \hat{c}_j\| &= \|Z_i - \hat{c}_2\|, \quad \text{for } m+1 \leq i \leq m+n, \\ \Rightarrow \hat{f}_{n+m}(Z_i) &= \|Z_i - \hat{c}_1\| - \|Z_i - \hat{c}_2\| > 0, \quad \text{for } i > m. \end{aligned}$$

This implies that under H_1 (asymptotically) $T_n^{\text{exch}} = \sum_{i=1}^n i = n(n+1)/2$ which is significantly smaller than the mean leading to a rejection.

³Quasi-norm means that $\|\cdot\|$ does not need to satisfy the triangle inequality.

For the purposes of Theorem 5, it does not matter if one is able to obtain the global minimum in (31). Any method for obtaining $\hat{c}_1, \dots, \hat{c}_k$ is permissible, as long as the procedure does not depend on the X, Y labels. For instance, Lloyd’s k -means clustering algorithm, k -means++ algorithm (Arthur and Vassilvitskii, 2007) and many other variants (Celebi et al., 2013; Hamerly, 2010; Ding et al., 2015; Hamerly and Drake, 2015; Shen et al., 2017; Newling and Fleuret, 2016) will work. In addition to the variants of the algorithms, one can also use the data to choose k , the number of clusters based on the data. In particular, the well-known elbow rule can be used for this purpose.

2. **Dimension Reduction and k -means:** In the same setting as above, due to the high computational complexity (Grönlund et al., 2017; Aloise et al., 2009) of the k -means clustering, one might opt to perform a preliminary dimension reduction and then use k -means clustering. Another reason for dimension reduction could be a possibility that distributions differ along a low-dimensional projection. Many of the dimension reduction techniques discussed in Section 3.3.3 can be used. The final transformation \hat{f}_{n+m} is given by

$$\hat{f}_{n+m}(Z_i) := \|\Pi(Z_i) - \hat{c}_1\| - \min_{1 \leq j \leq k} \|\Pi(Z_i) - \hat{c}_j\|,$$

where $\Pi(\cdot)$ denotes the preliminary dimension reduction map with $\Pi(Z_1), \dots, \Pi(Z_{n+m})$ representing the dimension reduced data and $\hat{c}_1, \dots, \hat{c}_k$ now represent the k -cluster centers obtained from the dimension reduced data.

3. **Clustering and Density Estimation:** In this case, we provide a generic method of converting any unsupervised clustering method into a valid transformation for Theorem 5. Suppose we have an unsupervised clustering procedure \mathcal{P} that partitions the collection of random variables Z_1, \dots, Z_{n+m} into disjoint sets B_1, \dots, B_k (where k could itself be a part of \mathcal{P}). This clustering procedure should treat the data Z_1, \dots, Z_{n+m} in a permutation invariant way. See Wasserman and Tibshirani (2017) for some examples. From all the random variables in $B_j, 1 \leq j \leq k$, estimate the density in a permutation invariant way; see Wang and Scott (2019) and Kim et al. (2019) for some examples. Then the transformation $\hat{f}_{n+m}(\cdot)$ is given by

$$\hat{f}_{n+m}(Z_i) := \frac{\hat{p}_{B_1}(Z_i)}{\max_{1 \leq j \leq k} \hat{p}_{B_j}(Z_i)}.$$

To gain intuition for this transformation, observe that if we obtain $\hat{c}_1, \dots, \hat{c}_k$ from a nearest neighbor clustering and we take the density estimator $\hat{p}_{B_j}(x) = \phi(x; \hat{c}_j, I)$, the multivariate normal density with location \hat{c}_j and variance identity, then $\hat{f}_{n+m}(Z_i)$ is a simple transformation of $\|Z_i - \hat{c}_1\| - \min_{1 \leq j \leq k} \|Z_i - \hat{c}_j\|$ matching the one in first example above.

As in the previous example, one can first apply a dimension reduction procedure on the data and then apply the unsupervised clustering method. In this case, the density estimator could be based on the dimension reduced data.

4.2 Testing Independence

Suppose $(X_1, Y_1), \dots, (X_n, Y_n) \in \mathcal{X} \times \mathcal{Y}$ are independent and identically distributed random variables from a probability measure P on $\mathcal{X} \times \mathcal{Y}$. If the marginal distribution of X_i ’s is P_X and the marginal distribution of Y_i ’s is P_Y , then the hypothesis to test is

$$H_0 : P = P_X \otimes P_Y \quad \text{versus} \quad H_1 : P \neq P_X \otimes P_Y. \quad (32)$$

Here $P_X \otimes P_Y$ represents the joint probability distribution with independent marginals of P_X and P_Y . Similar to the problem of testing equality of distributions, there exist numerous tests for (32) and we refer to Han et al. (2017); Deb and Sen (2021); Shi et al. (2020) for an overview. One of the classical tests for independence hypothesis is based on the Spearman's rank correlation (Hollander et al., 2013, Section 8.5). This test applies when P_X and P_Y are supported on the real line. Suppose the observations are $(X_1, Y_1), \dots, (X_n, Y_n) \in \mathbb{R}^2$. Let R_1^X, \dots, R_n^X denote the ranks of X_1, \dots, X_n and R_1^Y, \dots, R_n^Y denote the ranks of Y_1, \dots, Y_n . Under the null hypothesis H_0 , (R_1^X, \dots, R_n^X) and (R_1^Y, \dots, R_n^Y) are independent random vectors. Further each of these vectors is distributed as uniform on all permutations of $(1, 2, \dots, n)$ because of Theorem 2. The Spearman's rank correlation is given by

$$\rho_n := 1 - \frac{6}{n} \sum_{i=1}^n \frac{(R_i^X - R_i^Y)^2}{(n^2 - 1)}. \quad (33)$$

Because the distributions of (R_1^X, \dots, R_n^X) and (R_1^Y, \dots, R_n^Y) do not depend on P_X and P_Y , ρ_n in (33) has the same distribution (under H_0) irrespective of what P_X and P_Y are.

Noting that the distribution-free nature of the test depends only on the fact that ranks are distribution-free, we get that we can transform the data in each coordinate almost arbitrarily. Let \hat{f}_X be a transformation that depends on X_1, \dots, X_n permutation invariantly and let \hat{f}_Y be a transformation that depends on Y_1, \dots, Y_n permutation invariantly. Then by Theorem 3, $\hat{f}_X(X_1), \dots, \hat{f}_X(X_n)$ are exchangeable and $\hat{f}_Y(Y_1), \dots, \hat{f}_Y(Y_n)$ are also exchangeable. Further under the null hypothesis H_0 , the vectors $(\hat{f}_X(X_1), \dots, \hat{f}_X(X_n))$ and $(\hat{f}_Y(Y_1), \dots, \hat{f}_Y(Y_n))$ are independent. This leads to the following result.

Theorem 6. *Suppose $(X_1, Y_1), \dots, (X_n, Y_n) \in \mathcal{X} \times \mathcal{Y}$ are independent observations. Suppose \hat{f}_X and \hat{f}_Y are transformations depending permutation invariantly only on (X_1, \dots, X_n) and (Y_1, \dots, Y_n) , respectively. Then the statistic*

$$\rho_n(\hat{f}_X, \hat{f}_Y) := 1 - \frac{6}{n} \sum_{i=1}^n \frac{(R_i^X - R_i^Y)^2}{n^2 - 1},$$

has the same distribution as Spearman's rank correlation (33) under H_0 . Here

$$\begin{aligned} R_i^X &:= \text{rank}(\hat{f}_X(X_i); \{\hat{f}_X(X_1), \dots, \hat{f}_X(X_n)\}), \\ R_i^Y &:= \text{rank}(\hat{f}_Y(Y_i); \{\hat{f}_Y(Y_1), \dots, \hat{f}_Y(Y_n)\}). \end{aligned}$$

The main conclusion of Theorem 6 is that any permutation invariant data-driven transformations \hat{f}_X and \hat{f}_Y will lead to a distribution-free finite sample valid test for the independence hypothesis (32), irrespective of the domain of the data. Theorem 6 is, however, lacking in one important way. If allowed, one might want to use transformations \hat{f}_X and \hat{f}_Y depending on $(X_i, Y_i), i \in [n]$ that leads to the maximal correlation between $\hat{f}_X(X_i), i \in [n]$ and $\hat{f}_Y(Y_i), i \in [n]$ (Rényi, 1959). These transformations would depend on both coordinates. This, however, does not lead to validity at least through Theorem 6. Once again, Theorem 6 can be seen as an analogy of the full conformal prediction method.

4.3 Rank Tests based on Sample Splitting

In the previous sections, we have restricted the nature of data-driven transformations; in case of equality of distributions, the transformations should not depend on the X, Y labels and in case of independence, the transformations have to depend on X_1, \dots, X_n and Y_1, \dots, Y_n only marginally not jointly. Although the tests control the type I error, these restrictions can drastically effect the power. We can avoid these

restrictions based on sample splitting, which will be described below. The following procedures based on sample splitting can be thought of as analogues to the split conformal prediction method.

Although the full sample based transformation is lacking in the literature, the sample splitting transformation for rank tests has been described in the literature albeit seems not widely known. See [Friedman \(2003\)](#) and [Vayatis et al. \(2009\)](#).

Equality of Distributions: Recall that the hypotheses are $H_0 : P_X = P_Y$ and $H_1 : P_X \neq P_Y$. The observations are X_1, \dots, X_n i.i.d from P_X and Y_1, \dots, Y_m i.i.d. from P_Y . Under the null hypothesis H_0 , Z_1, \dots, Z_{n+m} defined by $Z_i = X_i$ for $1 \leq i \leq n$ and $Z_i = Y_{i-n}$ for $n+1 \leq i \leq n+m$ are independent and identically distributed. Randomly split this collection into two parts:

$$S_1 := \{Z_i : i \in \mathcal{I}_1\} \quad \text{and} \quad S_2 := \{Z_i : i \in \mathcal{I}_2\},$$

where \mathcal{I}_1 contains a random subset of $[n+m]$ and $\mathcal{I}_2 = [n+m] \setminus \mathcal{I}_1$. For any transformation \hat{f}_{S_1} based on S_1 , Proposition 3 proves that $\hat{f}_{S_1}(Z_i), i \in \mathcal{I}_2 = \mathcal{I}_1^c$ are exchangeable. Hence,

$$T^{\text{split}} := \sum_{i=1, i \in \mathcal{I}_2}^n \text{rank}(\hat{f}_{S_1}(Z_i); \{\hat{f}_{S_1}(Z_j) : j \in \mathcal{I}_2\}),$$

the sum of ranks of the X_i random variables in the second split S_2 is a valid test statistic. It follows from the discussion in Section 4.1 that under H_0 , T^{split} has a distribution independent of $P_X = P_Y$ and the critical values can be obtained from the Wilcoxon rank sum test based on the total sample size of $|\mathcal{I}_2|$.

In comparison to the test statistic T_n^{exch} in Section 4.1, the transformation $\hat{f}_{S_1}(\cdot)$ can now depend arbitrarily on the X, Y labels. In particular, we describe a specific example below. Based on the first split S_1 of the data, obtain $\hat{p}_{X, S_1}(\cdot)$ and $\hat{p}_{Y, S_1}(\cdot)$, the estimators of density of P_X and P_Y ; the density estimator can be arbitrary ([Kim et al., 2019](#); [Wang and Scott, 2019](#)). The final transformation $\hat{f}_{S_1}(\cdot)$ is given by

$$\hat{f}_{S_1}(Z_i) := \frac{\hat{p}_{X, S_1}(Z_i)}{\hat{p}_{Y, S_1}(Z_i)}, \quad \text{for all } i \in \mathcal{I}_2.$$

Further examples can be derived by writing the data in S_1 as $(Z_i, W_i), i \in \mathcal{I}_1$, where $W_i = 0$ if $Z_i \sim P_X$ and $W_i = 1$ if $Z_i \sim P_Y$ and finding an estimator $\hat{\eta}_{S_1}(\cdot)$ of the conditional probability $\eta(z) := \mathbb{P}(W = 1 | Z = z)$ based only on S_1 . The final transformation then would be $\hat{f}_{S_1}(Z_i) = \hat{\eta}_{S_1}(Z_i)$, which would naturally be higher for cases where $W_i = 1$ for $i \in \mathcal{I}_2$ (under H_1). This can also be combined with the methods of central subspace estimation ([Ma and Zhu, 2013](#)), which is fruitful when $\eta(\cdot)$ depends on a few coordinates or directions. For instance, if P_X and P_Y only differ in the distribution of the first coordinate, then one can at first apply a subspace estimation algorithm for W on Z with the data in S_1 and then find a classifier based on the reduced subspace.

Independence: Recall that the hypotheses are $H_0 : P = P_X \otimes P_Y$ and $H_1 : P \neq P_X \otimes P_Y$. The observations are $(X_1, Y_1), \dots, (X_n, Y_n)$ which are i.i.d. from P . As before, randomly split the data into two parts:

$$S_1 := \{(X_i, Y_i) : i \in \mathcal{I}_1\} \quad \text{and} \quad S_2 := \{(X_i, Y_i) : i \in \mathcal{I}_2\},$$

where \mathcal{I}_1 contains a random subset of $\{1, 2, \dots, n\}$ and $\mathcal{I}_2 = \{1, 2, \dots, n\} \setminus \mathcal{I}_1$. For any transformations $\hat{f}_{X, S_1}(\cdot)$ and $\hat{f}_{Y, S_1}(\cdot)$ based on S_1 , Proposition 3 yields that the bivariate random vectors $(\hat{f}_{X, S_1}(X_i), \hat{f}_{Y, S_1}(Y_i)), i \in \mathcal{I}_2$ are exchangeable and hence

$$\rho^{\text{split}}(\hat{f}_{X, S_1}, \hat{f}_{Y, S_1}) := 1 - \frac{6}{|\mathcal{I}_2|} \sum_{i \in \mathcal{I}_2} \frac{(R_i^X - R_i^Y)^2}{|\mathcal{I}_2|^2 - 1},$$

is a valid test statistic, where

$$\begin{aligned} R_i^X &:= \text{rank}(\widehat{f}_{X,S_1}(X_i); \{\widehat{f}_{X,S_1}(X_j) : j \in \mathcal{I}_2\}), \\ R_i^Y &:= \text{rank}(\widehat{f}_{Y,S_1}(Y_i); \{\widehat{f}_{Y,S_1}(Y_j) : j \in \mathcal{I}_2\}). \end{aligned}$$

Following the discussion in Section 4.2, we conclude that under the null hypothesis $H_0 : P = P_X \otimes P_Y$, $\rho^{\text{split}}(\widehat{f}_{X,S_1}, \widehat{f}_{Y,S_1})$ has the same distribution as the Spearman’s rank correlation based on sample size $|\mathcal{I}_2|$.

In comparison to the test statistic in Section 4.2, we can now use transformations \widehat{f}_{X,S_1} and \widehat{f}_{Y,S_1} that can depend on $(X_i, Y_i), i \in S_1$ jointly not just marginally. We describe a specific example below. Based on the first split S_1 of the data, obtain transformations $\widehat{f}_{X,S_1}(\cdot)$ and $\widehat{f}_{Y,S_1}(\cdot)$ that maximize the “correlation” between X_i, Y_i for $i \in S_1$; any technique can be used here and the “correlation” measure is also arbitrary. See [Breiman and Friedman \(1985\)](#) for an example and one can also mix this methodology with dimension reduction techniques ([Ma and Zhu, 2013](#)). These transformations can be used in the statistic $\rho^{\text{split}}(\widehat{f}_{X,S_1}, \widehat{f}_{Y,S_1})$ above. The critical values for this statistic can be obtained as before.

Summarizing the discussion in Section 4, we have shown that the distribution-free nature of the rank tests continues to hold under a large class of data-driven transformations. In all these sections, we have described the procedures only through two classical tests: Wilcoxon rank-sum test and Spearman’s rank correlation test. Because most rank tests only depend on the fact that ranks are distributed uniformly over all permutations, the procedures can also be used with other rank tests ([Hollander et al., 2013](#)).

5 Summary and Concluding Remarks

We have described the fundamental concept of exchangeability and its implications for prediction regions as well as rank tests. By describing the basic components, the intention is to bring the conformal prediction more into practice and also to show the wide range of flexibility hiding within the rank tests. In both these topics, we have (intentionally) not done an in-depth survey of the existing literature. We encourage the reader to refer to the cited literature to explore these topics further.

Of course, in both cases (prediction and testing), it is also of interest to understand the “power”. For a prediction region, this could be the length/volume of the region and for a test, it is the usual power ($1 - \text{type II error}$). In the case of conformal prediction, we discussed the imitation of the optimal volume prediction region but it should be stressed that, in general, optimality is hard to attain in finite samples in a distribution-free way because it requires the transformation used in practice to be the optimal transformation and in general, one can only consistently estimate that optimal transformation under “smoothness” assumptions. See [Lei et al. \(2013\)](#), [Györfi and Walk \(2020\)](#), and [Yang and Kuchibhotla \(2021\)](#) for some optimality results.

In the case of testing, the optimal transformation for the equality of distribution testing would require estimation of the optimal distribution separating transformation. For instance, suppose P_X and P_Y are two distributions on \mathbb{R}^d and in truth, they differ in their distributions only in the first coordinate. Then the optimal transformation to use is $x \in \mathbb{R}^d \mapsto x_1 \in \mathbb{R}$. Among all the tests of the form suggested in Pseudocode 3, the optimal transformation should converge to $x \mapsto x_1$ asymptotically for optimality in this class of tests. As with conformal prediction, this can be hard because of distributional assumptions and the curse of dimensionality in estimating the optimal transformation and also because the full-data transformation \widehat{f}_{n+m} is not allowed to use the true X, Y labels of Z_1, \dots, Z_{n+m} which makes it an unsupervised problem. The second issue can be alleviated by sample splitting.

In light of the discussion here, we now briefly mention a few open questions. Firstly, regarding conformal prediction, we have focused on the split conformal method for computational efficiency. This method uses one part of the data for training and the other part for calibrating the prediction region. As mentioned, it has been argued in the literature that the split conformal method could incur statistical inefficiency due to this splitting. Because prediction regions (unlike confidence regions) do not shrink to a singleton, it is not clear how to characterize this statistical inefficiency. The results of [Lei et al. \(2018\)](#) and [Sesia and Candès \(2020\)](#) already prove that, under certain assumptions, the split conformal regions can “converge” to the optimal prediction region. In this sense, asymptotic volume optimality holds in general but to understand the sub-optimality stemming from splitting, we need refined results. We believe it to be an open question on how these refined results look. Secondly, related to rank tests, we introduced sample splitting as a way of avoiding restrictions on the data-driven transformations. There is, however, a trade-off in that sample splitting tests are only based on a fraction of the total sample size and hence can also sacrifice power. It would be interesting to understand if there is a way to improve power and make use of data more cleverly; this could be done based in p-value combination techniques ([Vovk and Wang, 2019](#)) or the leave-one-out analogues ([Barber et al., 2021](#)). Study of power gains of such procedures requires further exploration. Furthermore, it would be interesting to study the (asymptotic) power properties of the tests discussed in Section 4 when the data-driven transformations are assumed to be consistent (in suitable metric) to their targets.

Acknowledgments

The author thanks Richard Berk, Andreas Buja, Rohit Patra for reading the earlier versions of the article and providing constructive comments that led to an improvement in the exposition. The author is also grateful to the reviewers, the associate editor, and the editor for their comments that led to the improved presentation.

References

- Aldous, D. J. (1985). Exchangeability and related topics. In *École d’Été de Probabilités de Saint-Flour XIII—1983*, pages 1–198. Springer.
- Aloise, D., Deshpande, A., Hansen, P., and Popat, P. (2009). NP-hardness of euclidean sum-of-squares clustering. *Machine learning*, 75(2):245–248.
- Arthur, D. and Vassilvitskii, S. (2007). k-means++: the advantages of careful seeding, p 1027–1035. In *SODA’07: proceedings of the eighteenth annual ACM-SIAM symposium on discrete algorithms. Society for Industrial and Applied Mathematics, Philadelphia, PA*.
- Balasubramanian, V., Ho, S.-S., and Vovk, V. (2014). *Conformal prediction for reliable machine learning: theory, adaptations and applications*. Morgan Kaufmann Publishers Inc.
- Barber, R. F., Candès, E. J., Ramdas, A., and Tibshirani, R. J. (2021). Predictive inference with the jackknife+. *The Annals of Statistics*, 49(1):486–507.

- Bates, S., Angelopoulos, A., Lei, L., Malik, J., and Jordan, M. I. (2021). Distribution-free, risk-controlling prediction sets. *arXiv preprint arXiv:2101.02703*.
- Bhattacharya, B. B. (2019). A general asymptotic framework for distribution-free graph-based two-sample tests. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 81(3):575–602.
- Bhattacharya, B. B. (2020). Asymptotic distribution and detection thresholds for two-sample tests based on geometric graphs. *Annals of Statistics*, 48(5):2879–2903.
- Breiman, L. and Friedman, J. H. (1985). Estimating optimal transformations for multiple regression and correlation. *Journal of the American statistical Association*, 80(391):580–598.
- Burnaev, E. and Vovk, V. (2014). Efficiency of conformalized ridge regression. In *Conference on Learning Theory*, pages 605–622.
- Carlsson, L., Ahlberg, E., Boström, H., Johansson, U., and Linusson, H. (2015). Modifications to p-values of conformal predictors. In *International Symposium on Statistical Learning and Data Sciences*, pages 251–259. Springer.
- Carlsson, L., Eklund, M., and Norinder, U. (2014). Aggregated conformal prediction. In *IFIP International Conference on Artificial Intelligence Applications and Innovations*, pages 231–240. Springer.
- Celebi, M. E., Kingravi, H. A., and Vela, P. A. (2013). A comparative study of efficient initialization methods for the k-means clustering algorithm. *Expert systems with applications*, 40(1):200–210.
- Chen, W., Chun, K.-J., and Barber, R. F. (2018). Discretized conformal prediction for efficient distribution-free inference. *Stat*, 7(1):e173.
- Chernozhukov, V., Wüthrich, K., and Yinchu, Z. (2018). Exact and robust conformal inference methods for predictive machine learning with dependent data. pages 732–749.
- Chernozhukov, V., Wüthrich, K., and Zhu, Y. (2019). Distributional conformal prediction. *arXiv:1909.07889*.
- Commenges, D. (2003). Transformations which preserve exchangeability and application to permutation tests. *Journal of nonparametric statistics*, 15(2):171–185.
- Conradsen, K., Nielsen, A. A., Schou, J., and Skriver, H. (2003). A test statistic in the complex wishart distribution and its application to change detection in polarimetric sar data. *IEEE Transactions on Geoscience and Remote Sensing*, 41(1):4–19.
- Cunningham, P. (2008). Dimension reduction. In *Machine learning techniques for multimedia*, pages 91–112. Springer.
- De Finetti, B. (1929). Funzione caratteristica di un fenomeno aleatorio. In *Atti del Congresso Internazionale dei Matematici: Bologna del 3 al 10 de settembre di 1928*, pages 179–190.
- Dean, A. and Verducci, J. (1990). Linear transformations that preserve majorization, schur concavity, and exchangeability. *Linear algebra and its applications*, 127:121–138.
- Deb, N. and Sen, B. (2021). Multivariate rank-based distribution-free nonparametric testing using measure transportation. *Journal of the American Statistical Association*, (just-accepted):1–45.

- Ding, Y., Zhao, Y., Shen, X., Musuvathi, M., and Mytkowicz, T. (2015). Yinyang k-means: A drop-in replacement of the classic k-means with consistent speedup. In *International Conference on Machine Learning*, pages 579–587.
- Dishion, T. J., Capaldi, D. M., and Yoerger, K. (1999). Middle childhood antecedents to progressions in male adolescent substance use: An ecological analysis of risk and protection. *Journal of Adolescent Research*, 14(2):175–205.
- Farris, K. B. and Schopflocher, D. P. (1999). Between intention and behavior: an application of community pharmacists’ assessment of pharmaceutical care. *Social science & medicine*, 49(1):55–66.
- Folkes, V. S., Koletsky, S., and Graham, J. L. (1987). A field study of causal inferences and consumer reaction: the view from the airport. *Journal of consumer research*, 13(4):534–539.
- Foygel Barber, R., Candès, E. J., Ramdas, A., and Tibshirani, R. J. (2019). The limits of distribution-free conditional predictive inference. *Information and Inference: A Journal of the IMA*.
- Friedman, J. H. (2003). On multivariate goodness-of-fit and two-sample testing. *Statistical problems in particle physics, astrophysics, and cosmology*, page 311.
- Grønlund, A., Larsen, K. G., Mathiasen, A., Nielsen, J. S., Schneider, S., and Song, M. (2017). Fast exact k-means, k-medians and bregman divergence clustering in 1d. *arXiv:1701.07204*.
- Grover, R. and Dillon, W. R. (1985). A probabilistic model for testing hypothesized hierarchical market structures. *Marketing Science*, 4(4):312–335.
- Gupta, C., Kuchibhotla, A. K., and Ramdas, A. K. (2021). Nested conformal prediction and quantile out-of-bag ensemble methods. *Accepted at Pattern Recognition. Preprint at arXiv:1910.10562*.
- Györfi, L. and Walk, H. (2020). Nearest neighbor based conformal prediction. *Pub. Inst. Stat. Univ. Paris*, Special issue in honour of Denis Bosq’s 80th birthday(63):173–190.
- Hamerly, G. (2010). Making k-means even faster. In *Proceedings of the 2010 SIAM international conference on data mining*, pages 130–140. SIAM.
- Hamerly, G. and Drake, J. (2015). Accelerating Lloyd’s algorithm for k-means clustering. In *Partitional clustering algorithms*, pages 41–78. Springer.
- Han, F., Chen, S., and Liu, H. (2017). Distribution-free tests of independence in high dimensions. *Biometrika*, 104(4):813–828.
- Heller, R., Heller, Y., Kaufman, S., Brill, B., and Gorfine, M. (2016). Consistent distribution-free k-sample and independence tests for univariate random variables. *The Journal of Machine Learning Research*, 17(1):978–1031.
- Hewitt, E. and Savage, L. J. (1955). Symmetric measures on cartesian products. *Transactions of the American Mathematical Society*, 80(2):470–501.
- Hinton, G. E. and Salakhutdinov, R. R. (2006). Reducing the dimensionality of data with neural networks. *science*, 313(5786):504–507.

- Hollander, M., Wolfe, D. A., and Chicken, E. (2013). *Nonparametric statistical methods*. John Wiley & Sons.
- Hyndman, R. J. (1996). Computing and graphing highest density regions. *The American Statistician*, 50(2):120–126.
- Izbicki, R., Shimizu, G. T., and Stern, R. B. (2019). Flexible distribution-free conditional predictive bands using density estimators. *arXiv preprint arXiv:1910.05575*.
- Kim, B., Xu, C., and Foygel Barber, R. (2020). Predictive inference is free with the jackknife+-after-bootstrap. *Advances in Neural Information Processing Systems*, 33.
- Kim, J., Shin, J., Rinaldo, A., and Wasserman, L. (2019). Uniform convergence rate of the kernel density estimator adaptive to intrinsic volume dimension. pages 3398–3407.
- Kivaranovic, D., Johnson, K. D., and Leeb, H. (2020). Adaptive, distribution-free prediction intervals for deep networks. pages 4346–4356.
- Kuchibhotla, A. K. and Berk, R. A. (2021). Nested conformal prediction sets for classification with applications to probation data. *arXiv preprint arXiv:2104.09358*.
- Lei, J. (2019). Fast exact conformalization of the lasso using piecewise linear homotopy. *Biometrika*, 106(4):749–764.
- Lei, J., G’Sell, M., Rinaldo, A., Tibshirani, R. J., and Wasserman, L. (2018). Distribution-free predictive inference for regression. *Journal of the American Statistical Association*, 113(523):1094–1111.
- Lei, J., Rinaldo, A., and Wasserman, L. (2015). A conformal prediction approach to explore functional data. *Annals of Mathematics and Artificial Intelligence*, 74(1-2):29–43.
- Lei, J., Robins, J., and Wasserman, L. (2013). Distribution-free prediction sets. *J. Amer. Statist. Assoc.*, 108(501):278–287.
- Lei, J. and Wasserman, L. (2014). Distribution-free prediction bands for non-parametric regression. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76(1):71–96.
- Lh eritier, A. (2015). *Nonparametric methods for learning and detecting multivariate statistical dissimilarity*. PhD thesis, Universit e Nice Sophia Antipolis.
- Linusson, H., Norinder, U., Bostr om, H., Johansson, U., and L ofstr om, T. (2017). On the calibration of aggregated conformal predictors. In *The 6th Symposium on Conformal and Probabilistic Prediction with Applications, (COPA 2017), 13-16 June, 2017, Stockholm, Sweden*, pages 154–173.
- Liu, J. Z., Mcrae, A. F., Nyholt, D. R., Medland, S. E., Wray, N. R., Brown, K. M., Hayward, N. K., Montgomery, G. W., Visscher, P. M., and Martin, N. G. (2010). A versatile gene-based test for genome-wide association studies. *The American Journal of Human Genetics*, 87(1):139–145.
- Ma, Y. and Zhu, L. (2013). Efficient estimation in sufficient dimension reduction. *Annals of statistics*, 41(1):250.
- Martin, E. C. and Betensky, R. A. (2005). Testing quasi-independence of failure and truncation times via conditional Kendall’s tau. *Journal of the American Statistical Association*, 100(470):484–492.

- Matthews, K. E. and Taylor, M. S. (1996). Nonparametric methods for multivariate analysis using statistically equivalent blocks. Technical report, Army research lab aberdeen proving ground md.
- Mayer, T. (1975). Selecting economic hypotheses by goodness of fit. *The Economic Journal*, 85(340):877–883.
- Ndiaye, E. and Takeuchi, I. (2019). Computing full conformal prediction set with approximate homotopy. In *Advances in Neural Information Processing Systems*, pages 1386–1395.
- Newling, J. and Fleuret, F. (2016). Fast k-means with accurate bounds. In *International Conference on Machine Learning*, pages 936–944.
- Nguyen, L. H. and Holmes, S. (2019). Ten quick tips for effective dimensionality reduction. *PLoS computational biology*, 15(6).
- Papadopoulos, H., Proedrou, K., Vovk, V., and Gammerman, A. (2002). Inductive confidence machines for regression. In *European Conference on Machine Learning*, pages 345–356. Springer.
- Rényi, A. (1959). On measures of dependence. *Acta mathematica hungarica*, 10(3-4):441–451.
- Ressel, P. (1985). De Finetti-type theorems: an analytical approach. *The Annals of Probability*, 13(3):898–922.
- Romano, Y., Patterson, E., and Candes, E. (2019). Conformalized quantile regression. In *Advances in Neural Information Processing Systems*, pages 3538–3548.
- Romano, Y., Sesia, M., and Candès, E. J. (2020). Classification with valid and adaptive coverage. *arXiv preprint arXiv:2006.02544*.
- Schervish, M. J. (2012). *Theory of statistics*. Springer Science & Business Media.
- Sesia, M. and Candès, E. J. (2020). A comparison of some conformal quantile regression methods. *Stat*, 9(1):e261.
- Shafer, G. and Vovk, V. (2008). A tutorial on conformal prediction. *Journal of Machine Learning Research*, 9(Mar):371–421.
- Shen, X., Liu, W., Tsang, I., Shen, F., and Sun, Q.-S. (2017). Compressed k-means for large-scale clustering. In *Thirty-First AAAI Conference on Artificial Intelligence*.
- Shi, H., Drton, M., and Han, F. (2020). Distribution-free consistent independence tests via center-outward ranks and signs. *Journal of the American Statistical Association*, pages 1–16.
- Silva, V. D. and Tenenbaum, J. B. (2003). Global versus local methods in nonlinear dimensionality reduction. In *Advances in neural information processing systems*, pages 721–728.
- Solari, A. and Djordjilović, V. (2021). Multi split conformal prediction. *arXiv preprint arXiv:2103.00627*.
- Sorzano, C. O. S., Vargas, J., and Montano, A. P. (2014). A survey of dimensionality reduction techniques. *arXiv:1403.2877*.
- Tenenbaum, J. B., De Silva, V., and Langford, J. C. (2000). A global geometric framework for nonlinear dimensionality reduction. *science*, 290(5500):2319–2323.

- Vayatis, N., Depecker, M., and Cléménçon, S. (2009). AUC optimization and the two-sample problem. *Advances in Neural Information Processing Systems*, 22:360–368.
- Vorlickova, D. (1972). Asymptotic properties of rank tests of symmetry under discrete distributions. *The Annals of Mathematical Statistics*, 43(6):2013–2018.
- Vovk, V. (2015). Cross-conformal predictors. *Annals of Mathematics and Artificial Intelligence*, 74(1-2):9–28.
- Vovk, V. (2020). Testing randomness online. *Statistical Science (forthcoming)*. *arXiv:1906.09256*.
- Vovk, V., Gammerman, A., and Shafer, G. (2005). *Algorithmic learning in a random world*. Springer Science & Business Media.
- Vovk, V. and Wang, R. (2019). Combining p-values via averaging. *arXiv:1212.4966v3*.
- Wang, W., Huang, Y., Wang, Y., and Wang, L. (2014). Generalized autoencoder: A neural network framework for dimensionality reduction. In *Proceedings of the IEEE conference on computer vision and pattern recognition workshops*, pages 490–497.
- Wang, Z. and Scott, D. W. (2019). Nonparametric density estimation for high-dimensional data—algorithms and applications. *Wiley Interdisciplinary Reviews: Computational Statistics*, 11(4):e1461.
- Wasserman, L. and Tibshirani, R. J. (2017). Clustering. <https://www.stat.cmu.edu/~ryantibs/statml/lectures/clustering.pdf>.
- Wilcoxon, F. (1945). Individual comparisons by ranking methods. *Biometrics*, 1(6):80–83.
- Xie, H., Li, J., and Xue, H. (2017). A survey of dimensionality reduction techniques based on random projection. *arXiv:1706.04371*.
- Yang, Y. and Kuchibhotla, A. K. (2021). Finite-sample efficient conformal prediction. *arXiv preprint arXiv:2104.13871*.

A Auxiliary Results

The following two results follow from Theorem 3 and will play an important role for both conformal prediction and rank tests.

Proposition 3. *Suppose Z_1, \dots, Z_n, Z_{n+1} are exchangeable random variables. If $1 \leq n_1 \leq n$, and $\hat{T}_{n_1} := g(Z_1, \dots, Z_{n_1})$ is any statistic computed based only on Z_1, \dots, Z_{n_1} , then for any function \hat{f}_{n_1} depending arbitrarily on Z_1, \dots, Z_{n_1} ,*

$$\hat{f}_{n_1}(Z_{n_1+1}, \hat{T}_{n_1}), \hat{f}_{n_1}(Z_{n_1+2}, \hat{T}_{n_1}), \dots, \hat{f}_{n_1}(Z_n, \hat{T}_{n_1}), \hat{f}_{n_1}(Z_{n+1}, \hat{T}_{n_1}),$$

are exchangeable random variables.

Proof. Define the function

$$G \begin{pmatrix} Z_1 \\ Z_2 \\ \vdots \\ Z_{n+1} \end{pmatrix} := \begin{pmatrix} \hat{f}_{n_1}(Z_{n_1+1}, \hat{T}_{n_1}) \\ \vdots \\ \hat{f}_{n_1}(Z_{n+1}, \hat{T}_{n_1}) \end{pmatrix}.$$

For any permutation $\pi : \{n_1 + 1, \dots, n + 1\} \rightarrow \{n_1 + 1, \dots, n + 1\}$,

$$\pi G \begin{pmatrix} Z_1 \\ Z_2 \\ \vdots \\ Z_{n+1} \end{pmatrix} = \begin{pmatrix} \hat{f}_{n_1}(Z_{\pi(n_1+1)}, \hat{T}_{n_1}) \\ \dots \\ \hat{f}_{n_1}(Z_{\pi(n+1)}, \hat{T}_{n_1}) \end{pmatrix} = G \begin{pmatrix} Z_1 \\ \vdots \\ Z_{n_1} \\ \pi \begin{pmatrix} Z_{n_1+1} \\ \vdots \\ Z_{n+1} \end{pmatrix} \end{pmatrix} \stackrel{(a)}{=} G \left(\pi_1 \begin{pmatrix} Z_1 \\ Z_2 \\ \vdots \\ Z_{n+1} \end{pmatrix} \right),$$

for a permutation $\pi_1 : [n + 1] \rightarrow [n + 1]$, where

$$\pi_1(i) = i \quad \text{for all } 1 \leq i \leq n_1 \quad \text{and} \quad \pi_1(i) = \pi(i) \quad \text{for } i > n_1.$$

Equality (a) above follows from the fact that \hat{f}_{n_1} depends only on Z_1, \dots, Z_{n_1} . Hence Theorem 3 implies the result. \square

Proposition 4. *Suppose Z_1, \dots, Z_n are exchangeable, and \hat{f} is a function depending on Z_1, \dots, Z_n permutation invariantly.⁴ Then $\hat{f}(Z_1), \dots, \hat{f}(Z_n)$ are exchangeable.*

This result is same as Proposition 2.1 of Vovk (2020).

Proof of Proposition 4. Like in Proposition 3, we apply Theorem 3. Define the function

$$G \begin{pmatrix} Z_1 \\ \vdots \\ Z_n \end{pmatrix} := \begin{pmatrix} \hat{f}(Z_1) \\ \vdots \\ \hat{f}(Z_n) \end{pmatrix}.$$

⁴This means that the algorithm outputting \hat{f} does not use the indexing of Z_1, \dots, Z_n .

For any permutation $\pi : [n] \rightarrow [n]$,

$$\pi G \begin{pmatrix} Z_1 \\ \vdots \\ Z_n \end{pmatrix} = \begin{pmatrix} \hat{f}(Z_{\pi(1)}) \\ \vdots \\ \hat{f}(Z_{\pi(n)}) \end{pmatrix} = G \begin{pmatrix} Z_{\pi(1)} \\ \vdots \\ Z_{\pi(n)} \end{pmatrix}.$$

The first equality above holds because \hat{f} depends on Z_1, \dots, Z_n permutation invariantly. The result now follows from Theorem 3. \square

B Proofs of Results in Sections 2 and 3

B.1 Proof of Theorem 2

Fix $\xi > 0$ and set $Z_i = W_i + \xi U_i$. Because U_1, \dots, U_n continuously distributed and are distinct with probability 1, we have that the events

$$\{Z_{\pi(1)} \leq Z_{\pi(2)} \leq \dots \leq Z_{\pi(n)}\},$$

over all permutations $\pi : [n] \rightarrow [n]$, are disjoint and further, one of them has to have occurred. Hence

$$\sum_{\pi: [n] \rightarrow [n]} \mathbb{P}(Z_{\pi(1)} \leq \dots \leq Z_{\pi(n)}) = 1. \quad (34)$$

Because U_1, \dots, U_n are iid and W_1, \dots, W_n are exchangeable, Z_1, \dots, Z_n are exchangeable. This, by definition, implies that

$$(Z_1, \dots, Z_n) \sim (Z_{\pi(1)}, \dots, Z_{\pi(n)}),$$

for any permutation $\pi : [n] \rightarrow [n]$ and taking $A := \{(x_1, \dots, x_n) : x_1 \leq \dots \leq x_n\}$ yields

$$\begin{aligned} \mathbb{P}(Z_1 \leq \dots \leq Z_n) &= \mathbb{P}((Z_1, \dots, Z_n) \in A) \\ &= \mathbb{P}((Z_{\pi(1)}, \dots, Z_{\pi(n)}) \in A) \\ &= \mathbb{P}(Z_{\pi(1)} \leq \dots \leq Z_{\pi(n)}). \end{aligned}$$

This combined with (34) proves that for every permutation π ,

$$\mathbb{P}(Z_{\pi(1)} \leq \dots \leq Z_{\pi(n)}) = \frac{1}{n!}.$$

This proves the result because $\{Z_{\pi(1)} \leq \dots \leq Z_{\pi(n)}\}$ is equivalent to the event that $(\text{rank}(Z_i) : i \in [n])$ is a particular permutation of $[n]$.

B.2 Proof of Corollary 1

Because $\text{rank}(\cdot; \cdot)$ takes values in $\{1, 2, \dots, n\}$, we get that

$$\begin{aligned} \mathbb{P}\left(\text{rank}(W_n; \{W_1, \dots, W_n\}) \leq t\right) &= \mathbb{P}\left(\text{rank}(W_n; \{W_1, \dots, W_n\}) \leq [t]\right) \\ &= \sum_{i=1}^{\lfloor t \rfloor} \mathbb{P}\left(\text{rank}(W_n; \{W_1, \dots, W_n\}) = i\right) \\ &= \sum_{i=1}^{\lfloor t \rfloor} \frac{(n-1)!}{n!}, \end{aligned}$$

which proves the result.

B.3 Proof of Theorem 3

Fix the transformation $G : \mathcal{W}^n \rightarrow (\mathcal{W}')^m$ and a vector W of exchangeable random variables. Suppose for each permutation $\pi_1 : [m] \rightarrow [m]$, there exists a permutation $\pi_2 : [n] \rightarrow [n]$ such that

$$\pi_1 G(w) = G(\pi_2 w), \quad \text{for all } w \in \mathcal{W}^n.$$

Then for any permutation $\pi_1 : [m] \rightarrow [m]$, the distribution of $\pi_1 G(W)$ is same as the distribution of $G(\pi_2 W)$. Because W is a vector of exchangeable random variables, $\pi_2 W$ has the same distribution as W . Therefore, $G(\pi_2 W)$ (and hence, $\pi_1 G(W)$) has the same distribution as $G(W)$. This completes the proof of exchangeability of the vector $G(W)$ of random variables. Hence the transformation G preserves exchangeability.

To prove the second part, suppose $G : \mathcal{W}^n \rightarrow (\mathcal{W}')^m$ is a transformation preserving exchangeability. This means that whenever $W \in \mathcal{W}^n$ is a vector of exchangeable random variables, $G(W)$ is also a vector of exchangeable random variables in $(\mathcal{W}')^m$. Fix a (non-random) vector $w \in \mathcal{W}^n$ and define a random vector $W \in \mathcal{W}^n$ via the distribution

$$\mathbb{P}(W = \pi'_2 w) = \frac{1}{n!}, \quad \text{for all permutations } \pi'_2 : [n] \rightarrow [n].$$

The idea for this distribution comes from the proof of Theorem 4 in [Dean and Verducci \(1990\)](#). Random vector W has a uniform distribution on the set of all permutations of $w \in \mathcal{W}^n$. It is easy to verify that W is a vector of exchangeable random variables. Because, by assumption, $G : \mathcal{W}^n \rightarrow (\mathcal{W}')^m$ is an exchangeability preserving transformation, we get that $G(W)$ is also a vector of exchangeable random variables. In particular, the support of $\pi_1 G(W)$ remains constant over all permutations $\pi_1 : [m] \rightarrow [m]$. Note that the support of $G(W)$ is

$$\{G(\pi'_2 w) : \pi'_2 \text{ a permutation on } [n]\}.$$

Therefore, for all permutations $\pi_1 : [m] \rightarrow [m]$,

$$\begin{aligned} & \pi_1 \{G(\pi'_2 w) : \pi'_2 \text{ a permutation on } [n]\} \\ &= \{G(\pi_2 w) : \pi_2 \text{ a permutation on } [n]\}. \end{aligned} \tag{35}$$

Note that $\pi_1 G(w)$ is an element in the set on the left hand side of (35). This implies that $\pi_1 G(w)$ is equal to $G(\pi_2 w)$ for some permutation $\pi_2 : [n] \rightarrow [n]$. In other words, for all $w \in \mathcal{W}^n$ and all permutations $\pi_1 : [m] \rightarrow [m]$, there exists a permutation $\pi_2 : [n] \rightarrow [n]$ (possibly depending on w) such that $\pi_1 G(w) = G(\pi_2 w)$. The final part about necessary and sufficient condition for linear transformations follows from Theorem 4 of [Dean and Verducci \(1990\)](#). This completes the proof.

B.4 Proof of Theorem 4

Proposition 3 proves that $\hat{f}_{\mathcal{T}}(Z_{n_1+1}), \dots, \hat{f}_{\mathcal{T}}(Z_{n+1})$ are exchangeable. The result now follows from Proposition 1.