

Two equalities expressing the determinant of a matrix in terms of expectations over matrix-vector products

Jascha Sohl-Dickstein
jaschasd@google.com

Abstract—We introduce two equations expressing the inverse determinant of a full rank matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$ in terms of expectations over matrix-vector products. The first relationship is $|\det(\mathbf{A})|^{-1} = \mathbb{E}_{\mathbf{s} \sim S^{n-1}} [|\mathbf{A}\mathbf{s}|^{-n}]$, where expectations are over vectors drawn uniformly on the surface of an n -dimensional radius one hypersphere. The second relationship is $|\det(\mathbf{A})|^{-1} = \mathbb{E}_{\mathbf{x} \sim q} [p(\mathbf{A}\mathbf{x}) / q(\mathbf{x})]$, where p and q are smooth distributions, and q has full support.

I. DERIVATION

A. General relationship: $|\mathbf{A}|^{-1} = \mathbb{E}_{\mathbf{x} \sim q} [p(\mathbf{A}\mathbf{x}) / q(\mathbf{x})]$

Let $\mathbf{A} \in \mathbb{R}^{n \times n}$ be a full rank matrix, $|\mathbf{A}|$ be the absolute determinant of \mathbf{A} , and p and q be smooth distributions over \mathbb{R}^n , where q has full support. The inverse determinant can be related to an expectation over functions of matrix-vector products:

$$\begin{aligned} |\mathbf{A}|^{-1} &= |\mathbf{A}|^{-1} \int d^n \mathbf{x} p(\mathbf{x}) \\ &= \int d^n \mathbf{x} p(\mathbf{A}\mathbf{x}) \\ &= \int d^n \mathbf{x} q(\mathbf{x}) \frac{p(\mathbf{A}\mathbf{x})}{q(\mathbf{x})} \\ &= \mathbb{E}_{\mathbf{x} \sim q} \left[\frac{p(\mathbf{A}\mathbf{x})}{q(\mathbf{x})} \right] \end{aligned} \quad (1)$$

B. Refined relationship: $|\mathbf{A}|^{-1} = \mathbb{E}_{\mathbf{s} \sim S^{n-1}} [|\mathbf{A}\mathbf{s}|^{-n}]$

Set $q = p = \mathcal{N}$, where $\mathcal{N}(\mathbf{x})$ is the probability density at \mathbf{x} of a Gaussian with mean 0 and identity covariance. Additionally, let $\mathbb{E}_{\mathbf{s} \sim S^{n-1}} [\cdot]$ be an expectation over vectors drawn uniformly on the surface of an n -dimensional radius one hypersphere, and $\mathbb{E}_{r \sim \chi(n)} [\cdot]$ be an expectation over a chi distribution with n degrees of freedom. The following more refined relationship then follows:

$$\begin{aligned} |\mathbf{A}|^{-1} &= \mathbb{E}_{\mathbf{x} \sim \mathcal{N}} \left[\frac{\mathcal{N}(\mathbf{A}\mathbf{x})}{\mathcal{N}(\mathbf{x})} \right] \\ &= \mathbb{E}_{\mathbf{x} \sim \mathcal{N}} \left[\exp \left(\frac{1}{2} \left[\|\mathbf{x}\|^2 - \|\mathbf{A}\mathbf{x}\|^2 \right] \right) \right] \\ &= \mathbb{E}_{\mathbf{s} \sim S^{n-1}} \left[\mathbb{E}_{r \sim \chi(n)} \left[\exp \left(\frac{1}{2} \left[\|r\mathbf{s}\|^2 - \|\mathbf{A}r\mathbf{s}\|^2 \right] \right) \right] \right] \\ &= \mathbb{E}_{\mathbf{s} \sim S^{n-1}} \left[\mathbb{E}_{r \sim \chi(n)} \left[\exp \left(\frac{r^2}{2} \left[1 - \|\mathbf{A}\mathbf{s}\|^2 \right] \right) \right] \right] \\ &= \mathbb{E}_{\mathbf{s} \sim S^{n-1}} [|\mathbf{A}\mathbf{s}|^{-n}] \end{aligned} \quad (2)$$

Because $|\mathbf{A}|^{-1} = |\mathbf{A}^{-1}|$, Equation 2 also provides an unbiased stochastic estimator for the determinant of a matrix,

Convergence of stochastic estimator




Fig. 1. An experimental validation of the relationship in Equation 3. We stochastically estimate the determinant of a matrix \mathbf{A} , by averaging $|\mathbf{A}^{-1}\mathbf{s}|^{-n}$ over random unit-norm vectors \mathbf{s} . Here, \mathbf{A} is a 10×10 matrix, with iid, variance one, Gaussian entries.

in terms of matrix-vector products with its inverse:

$$|\mathbf{A}| = \mathbb{E}_{\mathbf{s} \sim S^{n-1}} [|\mathbf{A}^{-1}\mathbf{s}|^{-n}] \quad (3)$$

Experimental validation of this relationship is presented in Figure 1.

II. RELATED WORK

The equality in Section I-A has been used in physics [14], but appears not to have been previously published as an explicit identity. Related expressions appear in work on ratios of moments of quadratic forms [11], and in techniques for rewriting certain determinants in terms of integrals which can be evaluated by Monte Carlo [13, 5, 4]. In the special case of positive symmetric definite \mathbf{A} , Gaussian quadrature techniques have been used to stochastically estimate determinants [1]. Other work derives stochastic estimators of classes of log determinants [7, 12, 2]. Hadamard’s inequality can be reinterpreted as a stochastic upper bound on $|\mathbf{A}|$ in terms of the norms of row or column vectors [6].

III. DISCUSSION

We hope that the stochastic estimators presented in this note will enable new Monte Carlo techniques for estimating, or stochastically bounding, functions of matrix determinants. These relationships may be especially useful in machine learning for training and evaluating both normalizing flow models [3, 10, 8] and Gaussian process kernels [9].

ACKNOWLEDGMENTS

Thank you to Alex Alemi, Anudhyan Boral, Ricky Chen, Arnaud Doucet, Guy Gur-Ari, Albin Jones, Abhishek Kumar, Peyman Milanfar, Jeffrey Pennington, Christian Szegedy, Srinivas Vasudevan,

and Max Vladymyrov for helpful discussion and links to related work.

REFERENCES

- [1] Zhaojun Bai, Gark Fahey, and Gene Golub. Some large-scale matrix computation problems. *Journal of Computational and Applied Mathematics*, 74(1-2):71–89, 1996.
- [2] Tian Qi Chen, Jens Behrmann, David K Duvenaud, and Jörn-Henrik Jacobsen. Residual flows for invertible generative modeling. In *Advances in Neural Information Processing Systems*, pages 9913–9923, 2019.
- [3] Laurent Dinh, Jascha Sohl-Dickstein, and Samy Bengio. Density estimation using real nvp. *arXiv preprint arXiv:1605.08803*, 2016.
- [4] Jacob Friedrich Finkenrath. *Stochastic Methods for the Fermion Determinant in Lattice Quantum Chromodynamics*. PhD thesis, Universität Wuppertal, Fakultät für Mathematik und Naturwissenschaften, 2018.
- [5] F Fucito, G Parisi, E Marinari, and C Rebbi. A proposal for monte carlo simulations of fermionic systems. *Nucl. Phys. B*, 180(CERN-TH-2960):369–377, 1980.
- [6] Jacques Hadamard. Resolution d’une question relative aux determinants. *Bull. des sciences math.*, 2:240–246, 1893.
- [7] Insu Han, Dmitry Malioutov, and Jinwoo Shin. Large-scale log-determinant computation through stochastic chebyshev expansions. In *International Conference on Machine Learning*, pages 908–917, 2015.
- [8] Mahdi Karami, Dale Schuurmans, Jascha Sohl-Dickstein, Laurent Dinh, and Daniel Duckworth. Invertible convolutional flow. In *Advances in Neural Information Processing Systems*, pages 5636–5646, 2019.
- [9] Carl Edward Rasmussen. Gaussian processes in machine learning. In *Summer School on Machine Learning*, pages 63–71. Springer, 2003.
- [10] Danilo Jimenez Rezende and Shakir Mohamed. Variational inference with normalizing flows. *arXiv preprint arXiv:1505.05770*, 2015.
- [11] Andrew L Rukhin. Identities for negative moments of quadratic forms in normal variables. *Statistics & probability letters*, 79(8):1004–1007, 2009.
- [12] Arvind K Saibaba, Alen Alexanderian, and Ilse CF Ipsen. Randomized matrix-free trace and log-determinant estimators. *Numerische Mathematik*, 137(2):353–395, 2017.
- [13] DH Weingarten and DN Petcher. Monte carlo integration for lattice gauge theories with fermions. *Physics Letters B*, 99(4):333–338, 1981.
- [14] Robert W Zwanzig. High-temperature equation of state by a perturbation method. i. nonpolar gases. *The Journal of Chemical Physics*, 22(8):1420–1426, 1954.