

ON THE HOPF ALGEBRA STRUCTURE OF THE LUSZTIG QUANTUM DIVIDED POWER ALGEBRAS

NICOLÁS ANDRUSKIEWITSCH, IVÁN ANGIOÑO AND CRISTIAN VAY

ABSTRACT. We study the Hopf algebra structure of Lusztig's quantum groups. First we show that the zero part is the tensor product of the group algebra of a finite abelian group with the enveloping algebra of an abelian Lie algebra. Second we build them from the plus, minus and zero parts by means of suitable actions and coactions within the formalism presented by Sommerhauser to describe triangular decompositions.

1. INTRODUCTION

There are two versions of quantum groups at roots of 1: the one introduced and studied by De Concini, Kac and Procesi [DK, DP] and the quantum divided power algebra of Lusztig [Lu1, Lu2, Lu3, Lu4]. The small quantum groups (aka Frobenius-Lusztig kernels) appear as quotients of the first and Hopf subalgebras of the second; in both cases they fit into suitable exact sequences of Hopf algebras.

The key actor in all these constructions is what we now call a Nichols algebra of diagonal type. Indeed all the Hopf algebras involved have triangular decompositions compatible with the mentioned exact sequences; the positive part of the small quantum group is a Nichols algebra. The celebrated classification of the finite-dimensional Nichols algebras of diagonal type was achieved in [H]. The positive parts of the small quantum groups correspond to braidings of Cartan type, but there are also braidings of super and modular types in the list, see the survey [AA].

The question of defining the versions of the quantum groups of De Concini, Kac and Procesi on one hand, and of Lusztig on the other, for every Nichols algebra in the classification arises unsurprisingly. The first was solved in [Ang2] introducing Hopf algebras also with triangular decompositions and whose positive parts are now the distinguished pre-Nichols algebras of diagonal type. These were introduced earlier in [Ang1], instrumental to the description of the defining relations of the Nichols algebras. A distinguished pre-Nichols algebra projects onto the corresponding Nichols algebra and the

2010 *Mathematics Subject Classification.* 16T05.

The work of N. A., I. A. and C. V. was partially supported by CONICET and Secyt (UNC). The work of C.V. was partially supported by Foncyt PICT 2016-3927.

kernel is a normal Hopf subalgebra that is even central under a mild technical hypothesis, see [Ang2, AAR2]. The geometry behind these new Hopf algebras is studied in [AAY] for Nichols algebras coming in families.

Towards the second goal, the graded duals of those distinguished pre-Nichols algebras were studied in [AAR1] under the name of Lusztig algebras; when the braiding is of Cartan type one recovers in this way the positive (and the negative) parts of Lusztig's quantum groups. A Lusztig algebra contains the corresponding Nichols algebra as a normal Hopf subalgebra and the cokernel is an enveloping algebra $U(\mathfrak{n})$ under the same mild technical hypothesis mentioned above, see [AAR2]. In [AAR2, AAR3] it was shown that \mathfrak{n} is either 0 or the positive part of a semisimple Lie algebra that was determined explicitly in each case.

In order to construct the analogues of Lusztig's quantum groups at roots of one for Nichols algebras of diagonal type, we still need to define the 0-part and the interactions with the positive and negative parts. This leads us to understand the Hopf algebra structure of a Lusztig's quantum group which is the objective of this Note.

Let V be the $\mathbb{Z}[v, v^{-1}]$ -Hopf algebra as in [Lu3, 2.3]; the quantum group is defined by specialization of V . Our first goal is to describe the specialization of the 0-part V^0 , a commutative and cocommutative Hopf subalgebra of V . We show in Theorem 3.10 that it splits as the tensor product of the group algebra of a finite group and the enveloping algebra of the Cartan subalgebra of the corresponding Lie algebra. For this we use some skew-primitive elements $h_{i,n} \in V^0$, cf. Definition 3.4, defined from the elements $[K_i; 0]_t$ and K_i^n of the original presentation of [Lu3]. The elements $h_{i,n}$, or rather multiples of them, were already introduced in [Le2] towards defining unrolled versions of quantum groups; see Remark 3.5. We point out that the definition in *loc. cit.* is by a limit procedure, while ours is explicit in terms of polynomials $p_{n,s} \in \mathbb{Z}[v, v^{-1}]$ that we define recursively in Lemma 3.3. Theorem 3.10 appears in [Le1, Le2].

In [S] it is explained that Hopf algebras U with a triangular decomposition $U \simeq A \otimes H \otimes B$, where H is a Hopf subalgebra, A is a Hopf algebra in the category of left Yetter-Drinfeld modules and B is the same but right, plus natural compatibilities, can be described by some specific structure that we call a TD-datum. Our second goal is to spell out the TD-datum corresponding to the quantum group, see Theorem 4.4.

The paper is organized as follows. In Section 2 we set up some notation and recall the formalism of [S]. Section 3 contains the analysis of the Hopf algebra V^0 from [Lu3]. In Section 4 we recall the definition of Lusztig's version of quantum groups at roots of 1, show that it fits into the setting of [S] and prove Theorem 4.4. For simplicity of the exposition we assume that the underlying Dynkin diagram is simply-laced; in the last Subsection we discuss how one would extend the material to the general case.

The Lusztig's quantum groups enter into a cleft short exact sequence of Hopf algebras [A, 3.4.1, 3.4.4] and contain an unrolled version of the finite

quantum groups [Le2] but as is apparent from the description here, they are not unrolled Hopf algebras.

We are not aware of other papers containing information on the matter of our interest. Other versions of triangular decompositions similar to [S] appear in [Ma, L].

2. PRELIMINARIES

2.1. Conventions. We adhere to the notation in [Lu3, Lu4] as much as possible. If $t \in \mathbb{N}_0$, $\theta \in \mathbb{N}$ and $t < \theta$, then $\mathbb{I}_{t,\theta} := \{t, t+1, \dots, \theta\}$, $\mathbb{I}_\theta := \mathbb{I}_{1,\theta}$.

Let $\mathcal{A} = \mathbb{Z}[v, v^{-1}]$, the ring of Laurent polynomials in the indeterminate v , $\mathcal{A}' = \mathbb{Q}(v)$, its field of fractions; later we also need $\mathcal{A}'' := \mathbb{Z}[v, v^{-1}, (1-v^2)^{-1}]$. The v -numbers are the polynomials

$$[s]_v = \frac{v^s - v^{-s}}{v - v^{-1}}, \quad [N]_v! = \prod_{s=1}^N [s]_v, \quad \begin{bmatrix} N \\ i \end{bmatrix}_v = \frac{[N]_v!}{[N-i]_v! [i]_v!} \in \mathcal{A},$$

$s, N \in \mathbb{N}$, $i \in \mathbb{I}_{0,N}$. We denote $\begin{bmatrix} N \\ i \end{bmatrix}_v = 0$ when $i > N$, $i, N \in \mathbb{N}$.

If B is a commutative ring and $\xi \in B$ is a unit, then B is an \mathcal{A} -algebra via $v \mapsto \xi$; the elements $[s]_v$, $[N]_v!$, $\begin{bmatrix} N \\ i \end{bmatrix}_v$ of \mathcal{A} specialize to $[s]_\xi$, $[N]_\xi!$, $\begin{bmatrix} N \\ i \end{bmatrix}_\xi$ of B . As in [Lu5, 35.1.3], we fix $\ell \in \mathbb{N}$ and set

$$\ell' = \begin{cases} \ell & \text{if } \ell \text{ is odd,} \\ 2\ell & \text{if } \ell \text{ is even.} \end{cases}$$

This convention is slightly different from the one in [Lu3, 5.1, pp. 287 ff].

Let $\phi_{\ell'} \in \mathbb{Z}[v]$ be the ℓ' -th cyclotomic polynomial; let \mathcal{B} be the field of fractions of $\mathcal{A}/\langle \phi_{\ell'} \rangle$ and let ξ be the image of v in \mathcal{B} . We have in \mathcal{B}

$$(2.1) \quad \phi_{\ell'}(\xi^2) = 0, \quad \xi^\ell = (-1)^{\ell'+1}, \quad \xi^{2\ell} = \xi^{\ell^2} = 1,$$

$$(2.2) \quad \begin{bmatrix} N+M \\ M \end{bmatrix}_\xi = 0, \quad N, M \in \mathbb{I}_{0,\ell-1}, \quad N+M \geq \ell.$$

We also have that

$$(2.3) \quad \frac{[m\ell]_\xi}{[n\ell]_\xi} = \xi^{(m-n)\ell} \frac{m}{n}, \quad \frac{[m\ell+j]_\xi}{[n\ell+j]_\xi} = \xi^{(m-n)\ell}, \quad m, n \in \mathbb{N}_0, j \in \mathbb{I}_{\ell-1}.$$

Hence for all $m \geq n \in \mathbb{N}_0$ and $j \in \mathbb{I}_{\ell-1}$, we have

$$(2.4) \quad \begin{bmatrix} m\ell \\ n\ell \end{bmatrix}_\xi = \binom{m}{n}, \quad \begin{bmatrix} m\ell+j \\ j \end{bmatrix}_\xi = \xi^{mj\ell}, \quad \begin{bmatrix} m\ell+j-1 \\ j \end{bmatrix}_\xi = 0.$$

Let \mathbb{k} be a field; all algebras, coalgebras, etc. below are over \mathbb{k} unless explicitly stated otherwise. If A is an associative unital \mathbb{k} -algebra, then we identify \mathbb{k} with a subalgebra of A .

2.2. Hopf algebras with triangular decomposition. Let H be a Hopf algebra with multiplication m , comultiplication Δ (with Sweedler notation $\Delta(h) = h_{(1)} \otimes h_{(2)}$), counit ε and bijective antipode \mathcal{S} ; we add a subscript H when precision is desired. We denote by ${}^H_H\mathcal{YD}$, respectively \mathcal{YD}_H^H , the category of left-left, respectively right-right Yetter-Drinfeld modules over H . If $M \in {}^H_H\mathcal{YD}$, then the action of H on M is denoted by \triangleright while the coaction is $m \mapsto m_{(-1)} \otimes m_{(0)}$, whereas if $N \in \mathcal{YD}_H^H$, the action is denoted by \triangleleft and the coaction is $n \mapsto n_{(0)} \otimes n_{(1)}$. For Hopf algebras either in ${}^H_H\mathcal{YD}$ or \mathcal{YD}_H^H , we use notations as above but with the variation $\Delta(r) = r^{(1)} \otimes r^{(2)}$. Given Hopf algebras R in ${}^H_H\mathcal{YD}$ and S in \mathcal{YD}_H^H , their bosonizations are denoted $R\#H$, $H\#S$. If $A \xrightleftharpoons[\pi]{} H$ are morphisms of Hopf algebras with $\pi\iota = \text{id}_H$, then $R\#H \simeq A \simeq H\#S$ where R and S are the subalgebras of right and left coinvariants of π ; see [R, §11.6, §11.7].

A *TD-datum* over H [S, Definition 3.2] is a collection $(A, B, \rightharpoonup, \leftharpoonup, \sharp)$ where

- (i) A is a Hopf algebra in ${}^H_H\mathcal{YD}$;
- (ii) B is a Hopf algebra in \mathcal{YD}_H^H ;
- (iii) $\rightharpoonup: B \otimes A \rightarrow A$ is a left action so that A is a left B -module, and the following identities hold for all $a \in A$, $b \in B$ and $h \in H$:

$$(2.5) \quad \begin{aligned} b \rightharpoonup (h \triangleright a) &= h_{(1)} \triangleright ((b \triangleleft h_{(2)}) \rightharpoonup a), & b \rightharpoonup 1 &= \varepsilon_B(b), \\ \Delta_A(b \rightharpoonup a) &= \left(b^{(1)}_{(0)} \rightharpoonup a^{(1)} \right) \otimes \left(b^{(1)}_{(1)} \triangleright \left(b^{(2)} \rightharpoonup a^{(2)} \right) \right); \end{aligned}$$

- (iv) $\leftharpoonup: B \otimes A \rightarrow B$ is a right action so that B is a right A -module, and the following identities hold for all $a \in A$, $b \in B$ and $h \in H$:

$$(2.6) \quad \begin{aligned} (b \triangleleft h) \leftharpoonup a &= (b \leftharpoonup (h_{(1)} \triangleright a)) \triangleleft h_{(2)}, & 1 \leftharpoonup a &= \varepsilon_A(a), \\ \Delta_B(b \leftharpoonup a) &= \left((b^{(1)} \leftharpoonup a^{(1)}) \triangleleft a^{(2)}_{(-1)} \right) \otimes \left(b^{(2)} \leftharpoonup a^{(2)}_{(0)} \right); \end{aligned}$$

both actions also satisfy for all $a \in A$ and $b \in B$:

$$(2.7) \quad \begin{aligned} & \left(b^{(1)} \rightharpoonup a^{(1)} \right) \otimes \left(b^{(2)} \leftharpoonup a^{(2)} \right) \\ &= \left(b^{(1)}_{(1)} \triangleright \left(b^{(2)} \rightharpoonup a^{(2)}_{(0)} \right) \right) \otimes \left(\left(b^{(1)}_{(0)} \leftharpoonup a^{(1)} \right) \triangleleft a^{(2)}_{(-1)} \right) \end{aligned}$$

- (v) $\sharp: B \otimes A \rightarrow H$ is a linear map, $b \otimes a \mapsto b\sharp a$, satisfying the following identities for all $a, c \in A$, $b, d \in B$, $h \in H$.

Compatibility of \sharp with the structure of H :

$$(2.8) \quad \begin{aligned} (b\sharp (h_{(1)} \triangleright a)) h_{(2)} &= h_{(1)} \left((b \triangleleft h_{(2)}) \sharp a \right), \\ \Delta_H(b\sharp a) &= \left(b^{(1)}_{(0)} \sharp a^{(1)} \right) a^{(2)}_{(-1)} \otimes b^{(1)}_{(1)} \left(b^{(2)} \sharp a^{(2)}_{(0)} \right), \\ \varepsilon_H(b\sharp a) &= \varepsilon_B(b) \varepsilon_A(a), \end{aligned}$$

Compatibility of \sharp with the products of A and B :

$$(2.9) \quad \begin{aligned} b\sharp(ac) &= \left(b^{(1)}\sharp a^{(1)}\right) a^{(2)}{}_{(-1)} \left(\left(b^{(2)} \leftarrow a^{(2)}{}_{(0)}\right)\sharp c\right), \\ (bd)\sharp a &= \left(b\sharp\left(d^{(1)}{}_{(0)} \rightarrow a^{(1)}\right)\right) d^{(1)}{}_{(1)} \left(d^{(2)}\sharp a^{(2)}\right), \\ b\sharp 1 &= \varepsilon_B(b), \quad 1\sharp a = \varepsilon_A(a), \end{aligned}$$

Compatibility of the actions with the multiplications via \sharp :

$$(2.10) \quad \begin{aligned} b \rightarrow (ac) &= \left(b^{(1)}{}_{(0)} \rightarrow a^{(1)}\right) \times \\ &\quad \times \left(b^{(1)}{}_{(1)}(b^{(2)}\sharp a^{(2)})a^{(3)}{}_{(-1)} \triangleright \left[\left(b^{(3)} \leftarrow a^{(3)}{}_{(0)}\right) \rightarrow c\right]\right), \\ (bd) \leftarrow a &= \left(\left[b \leftarrow \left(d^{(1)}{}_{(0)} \rightarrow a^{(1)}\right)\right] \triangleleft d^{(1)}{}_{(1)}(d^{(2)}\sharp a^{(2)})a^{(3)}{}_{(-1)}\right) \times \\ &\quad \times \left(d^{(3)} \leftarrow a^{(3)}{}_{(0)}\right); \end{aligned}$$

Compatibility of the coactions with the comultiplications via \sharp :

$$(2.11) \quad \begin{aligned} &\left(b^{(1)}{}_{(0)} \rightarrow a^{(1)}\right)_{(-1)} b^{(1)}{}_{(1)} \left(b^{(2)}\sharp a^{(2)}\right) \otimes \left(b^{(1)}{}_{(0)} \rightarrow a^{(1)}\right)_{(0)} \\ &= \left(b^{(1)}{}_{(0)}\sharp a^{(1)}\right) a^{(2)}{}_{(-1)} \otimes \left(b^{(1)}{}_{(1)} \triangleright \left(b^{(2)} \rightarrow a^{(2)}{}_{(0)}\right)\right); \\ &\left(b^{(2)} \leftarrow a^{(2)}{}_{(0)}\right)_{(0)} \otimes \left(b^{(1)}\sharp a^{(1)}\right) a^{(2)}{}_{(-1)} \left(b^{(2)} \leftarrow a^{(2)}{}_{(0)}\right)_{(1)} \\ &= \left(\left(b^{(1)}{}_{(0)} \leftarrow a^{(1)}\right) \triangleleft a^{(2)}{}_{(-1)}\right) \otimes b^{(1)}{}_{(1)} \left(b^{(2)}\sharp a^{(2)}\right)_{(0)}. \end{aligned}$$

Proposition 2.1. (a) [S, 3.3, 3.4] *Let $(A, B, \rightarrow, \leftarrow, \sharp)$ be a TD-datum over H . Then $U := A \otimes H \otimes B$ is a Hopf algebra with multiplication, comultiplication and antipode:*

$$\begin{aligned} (a \otimes h \otimes b)(c \otimes k \otimes d) &= a \left(h_{(1)} \triangleright \left(b^{(1)}{}_{(0)} \rightarrow c^{(1)}\right)\right) \\ &\quad \otimes h_{(2)} b^{(1)}{}_{(1)} \left(b^{(2)}\sharp c^{(2)}\right) c^{(3)}{}_{(1)} k_{(1)} \otimes \left(\left(b^{(3)} \leftarrow c^{(3)}{}_{(2)}\right) \triangleleft k_{(2)}\right) d, \\ \Delta(a \otimes h \otimes b) &= \left(a^{(1)} \otimes a^{(2)}{}_{(-1)} h_{(1)} \otimes b^{(1)}{}_{(0)}\right) \otimes \left(a^{(2)}{}_{(0)} \otimes h_{(2)} b^{(1)}{}_{(1)} \otimes b^{(2)}\right), \\ \mathcal{S}(a \otimes h \otimes b) &= (1 \otimes 1 \otimes \mathcal{S}_B(b_{(0)}))(1 \otimes \mathcal{S}_H(a_{(-1)} h b_{(1)}) \otimes 1)(\mathcal{S}_A(a_{(0)}) \otimes 1 \otimes 1). \end{aligned}$$

(b) [S, 3.5] *Let \mathcal{U} be a Hopf algebra. Let A and B be Hopf algebras in ${}^H_H\mathcal{YD}$ and \mathcal{YD}_H^H respectively, provided with injective algebra maps*

$$\iota_A : A \hookrightarrow \mathcal{U}, \quad \iota_H : H \hookrightarrow \mathcal{U}, \quad \iota_B : B \hookrightarrow \mathcal{U}.$$

Assume that

- (i) *The map $A \otimes H \otimes B \xrightarrow{m_{\mathcal{U}}(\iota_A \otimes \iota_H \otimes \iota_B)} \mathcal{U}$ is a linear isomorphism.*
- (ii) *The induced maps $A\sharp H \rightarrow \mathcal{U}$, $H\sharp B \rightarrow \mathcal{U}$ are Hopf algebra maps.*

Then there exists a TD-datum $(A, B, \rightarrow, \leftarrow, \sharp)$ over H such that $\mathcal{U} \simeq U$. \square

Clearly these constructions are mutually inverse. In the setting of the Proposition, we say that $U \simeq A \otimes H \otimes B$ is a *triangular decomposition*.

As observed in [S], the verification of the conditions in the definition of TD-datum is easier when H is commutative and cocommutative.

3. THE ALGEBRA V^0

3.1. Basic definitions. We fix $\theta \in \mathbb{N}$. For simplicity, set $\mathbb{I} := \mathbb{I}_\theta$. Following [Lu3, 2.3, pp. 268 ff] we consider the \mathcal{A} -algebra V^0 presented by generators

$$(3.1) \quad K_i, \quad K_i^{-1}, \quad \begin{bmatrix} K_i; c \\ t \end{bmatrix}, \quad i \in \mathbb{I}, c \in \mathbb{Z}, t \in \mathbb{N}_0$$

and relations for all $i \in \mathbb{I}$, tagged as in *loc. cit.*,

$$(g5) \quad (v - v^{-1}) \begin{bmatrix} K_i; 0 \\ 1 \end{bmatrix} = K_i - K_i^{-1},$$

$$(g6) \quad \text{the generators (3.1) commute with each other,}$$

$$(g7) \quad K_i K_i^{-1} = 1, \quad \begin{bmatrix} K_i; 0 \\ 0 \end{bmatrix} = 1,$$

$$(g8) \quad \begin{bmatrix} t + t' \\ t \end{bmatrix}_v \begin{bmatrix} K_i; 0 \\ t + t' \end{bmatrix} = \sum_{0 \leq j \leq t'} (-1)^j v^{t(t'-j)} \begin{bmatrix} t + j - 1 \\ j \end{bmatrix}_v K_i^j \begin{bmatrix} K_i; 0 \\ t \end{bmatrix} \begin{bmatrix} K_i; 0 \\ t' - j \end{bmatrix},$$

$$t \geq 1, t' \geq 0,$$

$$(g9) \quad \begin{bmatrix} K_i; -c \\ t \end{bmatrix} = \sum_{0 \leq j \leq t} (-1)^j v^{c(t-j)} \begin{bmatrix} c + j - 1 \\ j \end{bmatrix}_v K_i^j \begin{bmatrix} K_i; 0 \\ t - j \end{bmatrix}, \quad t \geq 0, c \geq 1,$$

$$(g10) \quad \begin{bmatrix} K_i; c \\ t \end{bmatrix} = \sum_{0 \leq j \leq t} v^{c(t-j)} \begin{bmatrix} c \\ j \end{bmatrix}_v K_i^{-j} \begin{bmatrix} K_i; 0 \\ t - j \end{bmatrix}, \quad t \geq 0, c \geq 0.$$

Observe that (g9) and (g10) actually define the elements $\begin{bmatrix} K_i; c \\ t \end{bmatrix}$, $c \in \mathbb{Z} - 0$, in terms of $K_i^{\pm 1}$ and

$$(3.2) \quad k_{i,t} := \begin{bmatrix} K_i; 0 \\ t \end{bmatrix}, \quad t \in \mathbb{N}, i \in \mathbb{I}.$$

See also §4.4 for an equivalent formulation. Set

$$(3.3) \quad a_{i,t} = \frac{v^{-t} K_i - v^t K_i^{-1}}{v - v^{-1}} \quad i \in \mathbb{I}, t \in \mathbb{Z}.$$

Thus $\mathcal{S}(a_{i,t}) = -a_{i,-t}$. Taking $t' = 1$ in (g8) we have

$$(3.4) \quad [t+1]_v k_{i,t+1} = k_{i,t} (v^t k_{i,1} - [t]_v K_i) = k_{i,t} a_{i,t},$$

hence

$$(3.5) \quad [t]_v^! k_{i,t} = \prod_{0 \leq s < t} a_{i,s}.$$

Multiplying (3.4) by $v - v^{-1}$, we get

$$(3.6) \quad K_i^2 k_{i,t} = v^t (v^{t+1} - v^{-t-1}) K_i k_{i,t+1} + v^{2t} k_{i,t}.$$

Proposition 3.1. (a) [Lu3, Lemma 2.21] *The \mathcal{A} -module V^0 is free with basis*

$$(3.7) \quad K_1^{\delta_1} \cdots K_\theta^{\delta_\theta} k_{1,t_1} \cdots k_{\theta,t_\theta}, \quad \delta_i \in \{0, 1\}, t_i \in \mathbb{N}_0, i \in \mathbb{I}.$$

(b) [Lu3, 2.22] $V^0 \otimes_{\mathcal{A}} \mathcal{A}' \simeq \mathcal{A}'[\mathbb{Z}^{\mathbb{I}}]$ as \mathcal{A}' -algebras. \square

Thus V^0 is an \mathcal{A} -form of the group algebra $\mathcal{A}'[\mathbb{Z}^{\mathbb{I}}]$; actually it is a form of the Hopf algebra structure as we see next.

Lemma 3.2. *The \mathcal{A} -algebra V^0 is a Hopf algebra with comultiplication determined by*

$$(3.8) \quad \Delta(K_i^{\pm 1}) = K_i^{\pm 1} \otimes K_i^{\pm 1} \quad i \in \mathbb{I}.$$

Proof. Since V^0 is a subalgebra of the Hopf algebra $\mathcal{A}'[\mathbb{Z}]$, we need to see that $\Delta(V^0) \subset V^0 \otimes_{\mathcal{A}} V^0$. By (g9) and (g10), it is enough to show that

$$(3.9) \quad \Delta(k_{i,t}) = \sum_{0 \leq s \leq t} k_{i,t-s} K_i^{-s} \otimes k_{i,s} K_i^{t-s}$$

for all $i \in \mathbb{I}$ and $t \in \mathbb{N}$. We proceed by induction on t . If $t = 1$, then

$$\Delta(k_{i,1}) = \frac{1}{v - v^{-1}} (K_i \otimes K_i - K_i^{-1} \otimes K_i^{-1}) = k_{i,1} \otimes K_i + K_i^{-1} \otimes k_{i,1}.$$

If (3.9) is valid for t , then

$$\begin{aligned} \Delta(k_{i,t+1}) &= \frac{1}{[t+1]_v} \Delta(k_{i,t}) \Delta(a_{i,t}) = \frac{1}{[t+1]_v} \left(\sum_{0 \leq s \leq t} k_{i,t-s} K_i^{-s} \otimes k_{i,s} K_i^{t-s} \right) \times \\ &\quad \left(\frac{v^{-t} K_i \otimes K_i - v^t K_i^{-1} \otimes K_i^{-1}}{v - v^{-1}} \right) \\ &= \sum_{0 \leq s \leq t} \frac{v^{-t} k_{i,t-s} K_i^{1-s} \otimes k_{i,s} K_i^{t+1-s} - v^t k_{i,t-s} K_i^{-1-s} \otimes k_{i,s} K_i^{t-1-s}}{[t+1]_v (v - v^{-1})} \\ &= \sum_{0 \leq s \leq t} \frac{v^{-s}}{[t+1]_v} \left([t-s+1]_v k_{i,t+1-s} + \frac{v^{t-s} k_{i,t-s} K_i^{-1}}{v - v^{-1}} \right) K_i^{-s} \otimes k_{i,s} K_i^{t+1-s} \\ &\quad + \frac{1}{[t+1]_v} \sum_{0 \leq s \leq t} v^{t-s} k_{i,t-s} K_i^{-1-s} \otimes \left([s+1]_v k_{i,s+1} - \frac{v^{-s} k_{i,s} K_i}{v - v^{-1}} \right) K_i^{t-s} \\ &= k_{i,t+1} \otimes K_i^{t+1} + \frac{1}{[t+1]_v} \sum_{1 \leq s \leq t} v^{-s} [t-s+1]_v k_{i,t+1-s} K_i^{-s} \otimes k_{i,s} K_i^{t+1-s} \\ &\quad + \sum_{0 \leq s \leq t-1} v^{t-s} \frac{[s+1]_v}{[t+1]_v} k_{i,t-s} K_i^{-1-s} \otimes k_{i,s+1} K_i^{t-s} + K_i^{-1-t} \otimes k_{i,t+1} \\ &= k_{i,t+1} \otimes K_i^{t+1} + K_i^{-1-t} \otimes k_{i,t+1} \end{aligned}$$

$$\begin{aligned}
& + \sum_{1 \leq j \leq t} \frac{v^{-j}[t-j+1]_v + v^{t+1-j}[j]_v}{[t+1]_v} k_{i,t+1-j} K_i^{-j} \otimes k_{i,j} K_i^{t+1-j} \\
& = \sum_{0 \leq s \leq t+1} k_{i,t+1-s} K_i^{-s} \otimes k_{i,s} K_i^{t+1-s},
\end{aligned}$$

which completes the inductive step. \square

3.2. Some skew-primitive elements. We introduce some notation:

- For $n \in \mathbb{N}$, $\phi_n : \mathbb{N} \rightarrow \{0, 1\}$ is the map given by $\phi_n(j) = 0$ if $n - j$ is even and $\phi_n(j) = 1$ if $n - j$ is odd.
- $\Phi : V^0 \rightarrow V^0$ is the algebra automorphism determined by

$$(3.10) \quad K_i \mapsto -K_i, \quad K_i^{-1} \mapsto -K_i^{-1}, \quad \begin{bmatrix} K_i; c \\ t \end{bmatrix} \mapsto (-1)^t \begin{bmatrix} K_i; c \\ t \end{bmatrix}.$$

It is easy to see that (3.10) defines an algebra map. Notice that

$$(3.11) \quad \Phi(k_{i,n}) = (-1)^n k_{i,n}, \quad \Phi(K_i^{\pm n}) = (-1)^n K_i^{\pm n}, \quad n \in \mathbb{N}, i \in \mathbb{I}.$$

Lemma 3.3. *Let $n \in \mathbb{N}$. We define $p_{n,s} \in \mathbb{Z}[v, v^{-1}]$, $s \in \mathbb{I}_n$, recursively on s by $p_{n,1} = v^{-\phi_n(1)}$,*

$$p_{n,s} = \frac{v^{ns} - v^{-ns}}{v^{\phi_n(s)s}(v^n - v^{-n})} - \sum_{t \in \mathbb{I}_{s-1}} p_{n,t} \begin{bmatrix} s \\ t \end{bmatrix}_v v^{(\phi_n(t) - \phi_n(s))s}, \quad s > 1.$$

Then

$$(3.12) \quad K_i^n - K_i^{-n} = (v^n - v^{-n}) \sum_{s \in \mathbb{I}_n} p_{n,s} k_{i,s} K_i^{\phi_n(s)}.$$

Proof. Fix $i \in \mathbb{I}$. By Proposition 3.1 (a), $K_i^n - K_i^{-n}$ is a linear combination of $k_{i,t}$, $k_{i,t} K_i$, $t \in \mathbb{N}_0$. Indeed, it can be shown by induction on n that $K_i^{\pm n}$ belongs to the \mathcal{A} -submodule spanned by $k_{i,t}$, $k_{i,t} K_i$, $t \leq n$. Using the involution Φ , we see by (3.11) that there are $a_{n,t} \in \mathcal{A}$, $t \in \mathbb{I}_n$ such that

$$(3.13) \quad K_i^n - K_i^{-n} = \sum_{t \in \mathbb{I}_{0,n}} a_{n,t} k_{i,t} K_i^{\phi_n(t)}.$$

We extend scalars as in Proposition 3.1 (b) and consider the algebra maps

$$(3.14) \quad \Xi_{i,j} : \mathcal{A}'[\mathbb{Z}^{\mathbb{I}}] \rightarrow \mathcal{A}', \quad K_i \mapsto v^j, \quad K_p \mapsto 1, \quad p \neq i,$$

$j \in \mathbb{N}_0$. Notice that, with the convention $\begin{bmatrix} N \\ n \end{bmatrix}_v = 0$ when $n > N$,

$$(3.15) \quad \Xi_{i,j}(k_{i,t}) = \frac{1}{[t]_v!} \prod_{0 \leq s < t} \Xi_{i,j}(a_{i,s}) = \frac{1}{[t]_v!} \prod_{0 \leq s < t} \frac{v^{j-s} - v^{s-j}}{v - v^{-1}} = \begin{bmatrix} j \\ t \end{bmatrix}_v.$$

Applying $\Xi_{i,0}$ and $\Xi_{i,1}$ to (3.13), we see that $0 = a_{n,0}$, $v^n - v^{-n} = a_{n,1} v^{\phi_n(1)}$. Now we apply $\Xi_{i,s}$, $s > 1$, to (3.13):

$$v^{ns} - v^{-ns} = \sum_{t \in \mathbb{I}_s} a_{n,t} \begin{bmatrix} s \\ t \end{bmatrix}_v v^{\phi_n(t)s};$$

this implies the recursive formula holds since

$$a_{n,s} = v^{-\phi_n(s)s}(v^{ns} - v^{-ns}) - \sum_{t \in \mathbb{I}_{s-1}} a_{n,t} \begin{bmatrix} s \\ t \end{bmatrix}_v v^{(\phi_n(t) - \phi_n(s))s}. \quad \square$$

Definition 3.4. Let $n \in \mathbb{N}$. We set

$$(3.16) \quad h_{i,n} := \frac{K_i^n - K_i^{-n}}{n(v^n - v^{-n})} K_i^n = \frac{1}{n} \left(\sum_{s \in \mathbb{I}_n} p_{n,s} k_{i,s} K_i^{\phi_n(s)} \right) K_i^n \in V^0.$$

Then $h_{i,n} = \frac{K_i^{2n} - 1}{n(v^n - v^{-n})}$ is $(1, K_i^{2n})$ -skew primitive.

Remark 3.5. The elements H'_α defined in [Le2, Theorem 3.1] are multiples of the above elements in the particular case $n = \ell$. Explicitly, $H'_\alpha = \frac{\ell(v^\ell - v^{-\ell})}{\phi'_\ell(v^2)} \mathbf{h}_{i,\ell}$. Notice that H'_α are defined by taking a limit while (3.16) is an explicit expression in terms of the polynomials $p_{n,s}$ that are defined recursively. We discuss now these polynomials.

Lemma 3.6. Let $n \in \mathbb{N}$. Then

$$(3.17) \quad p_{n,n} = v^{-\binom{n}{2}} (-1)^{n-1} (v - v^{-1})^{n-1} [n-1]_v!.$$

Proof. We compute $k_{i,t}$ in $V^0 \otimes_{\mathcal{A}} \mathcal{A}' \simeq \mathcal{A}'[\mathbb{Z}^1]$:

$$k_{i,t} = \frac{1}{[t]_v!} \prod_{j=0}^{t-1} \frac{v^{-j} K_i - v^j K_i^{-1}}{v - v^{-1}} = \sum_{s=-t}^t f_{t,s} K_i^s, \quad \text{for some } f_{t,s} \in \mathcal{A}'.$$

In particular, $f_{t,-t} = \frac{(-1)^t v^{\binom{t}{2}}}{[t]_v! (v - v^{-1})^t}$. Looking at the equality (3.12), K_i^{-n} appears only in one summand, $p_{n,n} k_{i,n}$, on the right hand side. Hence

$$-1 = (v^n - v^{-n}) f_{n,-n} p_{n,n} = [n]_v (v - v^{-1}) \frac{(-1)^n v^{\binom{n}{2}}}{[n]_v! (v - v^{-1})^n} p_{n,n},$$

and the claimed equality follows. \square

Given $\ell \in \mathbb{N}$ we consider the lower triangular matrix

$$\mathbf{P}_{i,\ell} = \begin{pmatrix} p_{11} & 0 & 0 & \dots & \dots & 0 \\ p_{21} K_i & p_{22} & 0 & \dots & \dots & 0 \\ p_{31} & p_{32} K_i & p_{32} & \dots & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ p_{\ell 1} K_i^{\phi_\ell(1)} & p_{\ell 2} K_i^{\phi_\ell(2)} & p_{\ell 3} K_i^{\phi_\ell(3)} & \dots & p_{\ell \ell-1} K_i & p_{\ell \ell} \end{pmatrix},$$

and the column vectors

$$\mathbf{k}_{i,\ell} = \begin{pmatrix} k_{i,1} \\ \vdots \\ k_{i,\ell} \end{pmatrix}, \quad \tilde{\mathbf{h}}_{i,\ell} = \begin{pmatrix} \tilde{h}_{i,1} \\ \vdots \\ \tilde{h}_{i,\ell} \end{pmatrix}, \quad \text{where } \tilde{h}_{i,n} := n h_{i,n} K_i^{-n}.$$

Then (3.16) says that $\mathbf{P}_{i,\ell} \mathbf{k}_{i,\ell} = \tilde{\mathbf{h}}_{i,\ell}$. Recall $\mathcal{A}'' = \mathbb{Z}[v, v^{-1}, (1-v)^{-1}]$ so that the matrix $\mathbf{P}_{i,\ell}$ becomes invertible in $V^0 \otimes_{\mathcal{A}} \mathcal{A}''$ by (3.17). Let

$$\mathbf{P}_{i,\ell}^{-1} = \begin{pmatrix} q_{11} & 0 & 0 & \cdots & \cdots & 0 \\ q_{21} & q_{22} & 0 & \cdots & \cdots & 0 \\ q_{31} & q_{32} & q_{32} & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ q_{\ell 1} & q_{\ell 2} & q_{\ell 3} & \cdots & q_{\ell \ell-1} & q_{\ell \ell} \end{pmatrix}.$$

Then

$$k_{i,n} = \sum_{s \in \mathbb{I}_n} q_{n,s} \tilde{h}_{i,s} = \sum_{s \in \mathbb{I}_n} q_{n,s} h_{i,s} K_i^{-s}, \quad n \in \mathbb{N}, i \in \mathbb{I}.$$

Example 3.7. We compute $p_{n,s}$ for small values of n . For $n = 2$, $p_{2,1} = v^{-1}$,

$$p_{2,2} = \frac{v^4 - v^{-4}}{v^2 - v^{-2}} - p_{2,1} \begin{bmatrix} 2 \\ 1 \end{bmatrix}_v v^2 = v^2 + v^{-2} - v(v + v^{-1}) = v^{-2} - 1.$$

This agrees with (3.17). Thus,

$$h_{i,2} = \left(\frac{v^{-2} - 1}{2} k_{i,2} + \frac{v^{-1}}{2} k_{i,1} K_i \right) K_i^2.$$

For $n = 3$, we have that $p_{3,1} = 1$,

$$\begin{aligned} p_{3,2} &= \frac{v^6 - v^{-6}}{v^2(v^3 - v^{-3})} - p_{3,1} \begin{bmatrix} 2 \\ 1 \end{bmatrix}_v v^{-2} = \frac{(v - v^{-1})^2 [2]_v}{v^2}, \\ p_{3,3} &= \frac{v^9 - v^{-9}}{v^3 - v^{-3}} - \sum_{t \in \mathbb{I}_2} p_{3,t} \begin{bmatrix} 3 \\ t \end{bmatrix}_v v^{3\phi_3(t)} = \frac{v^9 - v^{-9}}{v^3 - v^{-3}} - p_{3,1} \begin{bmatrix} 3 \\ 1 \end{bmatrix}_v - p_{3,2} \begin{bmatrix} 3 \\ 2 \end{bmatrix}_v v^3 \\ &= 1 - v^{-2} - v^{-4} + v^{-6} = \frac{(v - v^{-1})^2 [2]_v}{v^3}. \end{aligned}$$

Again this agrees with (3.17). Thus,

$$h_{i,3} = \left(\frac{(v - v^{-1})^2 [2]_v}{3v^3} k_{i,3} + \frac{(v - v^{-1})^2 [2]_v}{3v^2} k_{i,2} K_i + k_{i,1} \right) K_i^3.$$

The element H' computed in [Le2, Example 3.2] (assuming $\ell' = 4$ in our notation) is a multiple of $h_{i,2}$ above.

Remark 3.8. For instance, from the preceding formulas we conclude:

$$(3.18) \quad k_{i,1} = h_{i,1} K_i^{-1},$$

$$(3.19) \quad k_{i,2} = \frac{2}{v^{-2} - 1} h_{i,2} K_i^{-2} - \frac{v^{-1}}{v^{-2} - 1} h_{i,1},$$

$$(3.20) \quad \begin{aligned} k_{i,3} &= \frac{3v^3}{(v - v^{-1})^2 [2]_v} h_{i,3} K_i^{-3} - \frac{2v}{v^{-2} - 1} h_{i,2} K_i^{-1} + \frac{1}{v^{-2} - 1} h_{i,1} K_i \\ &\quad - \frac{3v^3}{(v - v^{-1})^2 [2]_v} h_{i,1} K_i^{-1}. \end{aligned}$$

3.3. Specializations of V^0 . Recall that $\ell' \in \mathbb{N}$ is defined in Section 2.1 and that \mathcal{B} is the field of fractions of $\mathcal{A}/\langle \phi_{\ell'} \rangle$. We study now $V_{\mathcal{B}}^0 := V^0 \otimes_{\mathcal{A}} \mathcal{B}$. Thus the map $\mathcal{A} \rightarrow \mathcal{B}$ factorizes through \mathcal{A}'' .

Lemma 3.9. [Lu2, Lemma 4.4], [Lu3, Lemma 2.21] *The algebra $V_{\mathcal{B}}^0$ is generated by K_i and $k_{i,\ell} = \begin{bmatrix} K_i; 0 \\ \ell \end{bmatrix}$, $i \in \mathbb{I}$. Furthermore,*

$$(3.21) \quad K_i^{2\ell} = 1.$$

Proof. We first prove (3.21). Taking $t = \ell - 1$ and $t' = 1$ in (g8) we have:

$$\begin{aligned} 0 &\stackrel{(2.2)}{=} \begin{bmatrix} \ell \\ \ell - 1 \end{bmatrix}_{\xi} \begin{bmatrix} K_i; 0 \\ \ell \end{bmatrix} = \xi^{\ell-1} \begin{bmatrix} K_i; 0 \\ \ell - 1 \end{bmatrix} \begin{bmatrix} K_i; 0 \\ 1 \end{bmatrix} - \begin{bmatrix} \ell - 1 \\ 1 \end{bmatrix}_{\xi} K_i \begin{bmatrix} K_i; 0 \\ \ell - 1 \end{bmatrix} \\ &= \begin{bmatrix} K_i; 0 \\ \ell - 1 \end{bmatrix} \frac{\xi K_i - \xi^{-1} K_i^{-1}}{\xi - \xi^{-1}} \stackrel{(3.5)}{=} \frac{\prod_{0 \leq s < \ell} \xi^{-s} K_i - \xi^s K_i^{-1}}{[\ell - 1]_{\xi} (\xi - \xi^{-1})^{\ell}} \\ &= \xi^{-\binom{\ell}{2}} \frac{K_i^{\ell} - K_i^{-\ell}}{[\ell - 1]_{\xi} (\xi - \xi^{-1})^{\ell}}. \end{aligned}$$

Hence (3.21) holds.

Let $t \in \mathbb{I}_{0,\ell-1}$. Then $[t]_{\xi}^! \neq 0$, so $k_{i,t} = \frac{1}{[t]_{\xi}^!} \prod_{0 \leq s < t} a_{i,s}$ belongs to the subalgebra generated by K_i . Now we claim that

$$(3.22) \quad k_{i,n\ell} = \frac{1}{n!} \prod_{0 \leq s < n} \left(k_{i,\ell} - s K_i^{\ell} \right), \quad \text{for all } n \in \mathbb{N}.$$

We take $t = n\ell$ and $t' = \ell$ in (g8). By (2.4) the left-hand side is

$$\begin{bmatrix} (n+1)\ell \\ n\ell \end{bmatrix}_{\xi} k_{i,(n+1)\ell} = (n+1) k_{i,(n+1)\ell},$$

while the right-hand side is, by (2.3),

$$\begin{aligned} &\sum_{0 \leq j \leq \ell} (-1)^j \xi^{n\ell(\ell-j)} \begin{bmatrix} n\ell + j - 1 \\ j \end{bmatrix}_{\xi} K_i^j k_{i,n\ell} k_{i,\ell-j} \\ &= k_{i,n\ell} k_{i,\ell} + (-1)^{\ell} \xi^{n\ell(\ell-1)} \frac{[n\ell]_{\xi}}{[\ell]_{\xi}} K_i^{\ell} k_{i,n\ell} = k_{i,n\ell} \left(k_{i,\ell} - n K_i^{\ell} \right). \end{aligned}$$

Hence $k_{i,(n+1)\ell} = \frac{1}{n+1} k_{i,n\ell} \left(k_{i,\ell} - n K_i^{\ell} \right)$, so we obtain (3.22) recursively.

Finally we take $t = m\ell$, $t' \in \mathbb{I}_{\ell-1}$ in (g8). Using (2.4),

$$k_{i,m\ell+t'} = \sum_{0 \leq j \leq t'} (-1)^j \begin{bmatrix} m\ell + j - 1 \\ j \end{bmatrix}_{\xi} K_i^j k_{i,m\ell} k_{i,t'-j} = k_{i,m\ell} k_{i,t'}.$$

Hence the claim follows from Proposition 3.1 (a). \square

Let $\Gamma = (\mathbb{Z}/2\ell)^{\mathbb{I}}$, with $g_i \in \Gamma$ being generators of the corresponding copies of the cyclic group $\mathbb{Z}/2\ell$. Let \mathfrak{h} be the abelian Lie algebra with basis $(t_i)_{i \in \mathbb{I}}$, so that $U(\mathfrak{h}) \simeq \mathcal{B}[t_i : i \in \mathbb{I}]$.

Theorem 3.10. [Le1, Theorem 4.1] *The assignment*

$$(3.23) \quad \Psi(g_i) = K_i, \quad \Psi(t_i) = h_{i,\ell}, \quad i \in \mathbb{I},$$

determines an isomorphism of Hopf algebras $\Psi : \mathcal{B}\Gamma \otimes U(\mathfrak{h}) \rightarrow V_{\mathcal{B}}^0$.

We present a different proof involving the polynomials $p_{i,t}$.

Proof. That (3.23) defines an algebra map follows by (g6) and (3.21); that is a surjective Hopf algebra map, by (3.8), Definition 3.4 and Lemma 3.9. It remains to prove that Ψ is injective. By [M, 5.3.1] it reduces to prove that Ψ is injective on the first term of the coradical filtration, i.e. that the set $\{K_i^p h_{r,\ell}^j : i, r \in \mathbb{I}_\theta, p \in \mathbb{I}_{0,2\ell-1}, j \in \mathbb{I}_{0,1}\}$ is linearly independent. By the assumption $\xi^2 \neq 1$, we have $p_{\ell,\ell}(\xi) = n$, so (3.16) implies that

$$(3.24) \quad h_{i,\ell} \in k_{i,\ell} K_i^\ell + \mathcal{B}\langle K_i \rangle,$$

see the line before (3.22). We need then to prove that the set

$$\{K_i^p k_{r,\ell}^j : i, r \in \mathbb{I}_\theta, p \in \mathbb{I}_{0,2\ell-1}, j \in \mathbb{I}_{0,1}\}$$

is linearly independent. Indeed, suppose that

$$(3.25) \quad 0 = \sum_{i,p} e_{i,p} K_i^p + b_{i,p} K_i^p k_{i,\ell},$$

where $e_{i,p}, b_{i,p} \in \mathcal{B}$. Fix $i \in \mathbb{I}$. The \mathcal{A}' -algebra maps $\Xi_{i,j} : \mathcal{A}'[\mathbb{Z}] \rightarrow \mathcal{A}'$ as in (3.14) satisfy $\Xi_{i,j}(V^0) \subseteq \mathcal{A}$ by (3.15). We restrict to \mathcal{A} -algebra maps $\Xi_{i,j} : V^0 \rightarrow \mathcal{A}$ and tensorize to get \mathcal{B} -algebra maps $\Xi_{i,j} : V_{\mathcal{B}}^0 \rightarrow \mathcal{B}$ such that

$$\Xi_{i,j}(K_i) = \xi^j, \quad \Xi_{i,j}(k_{i,\ell}) = \begin{bmatrix} j \\ \ell \end{bmatrix}_\xi, \quad \Xi_{i,j}(K_r) = 1, \quad \Xi_{i,j}(k_{r,\ell}) = 0 \text{ if } r \neq i.$$

Applying $\Xi_{i,j}$ to (3.25), we get

$$(3.26) \quad 0 = \sum_{p \in \mathbb{I}_{0,2\ell-1}} e_{i,p} \xi^{pj}, \quad 0 \leq j < \ell;$$

$$(3.27) \quad 0 = \sum_{p \in \mathbb{I}_{0,2\ell-1}} e_{i,p} \xi^{pj} + b_{i,p} \xi^{pj}, \quad \ell \leq j < 2\ell.$$

If ℓ' is even, then $\ell' = 2\ell$ and from (3.26) we deduce that $e_{i,p} = 0$ for all $p \in \mathbb{I}_{0,2\ell-1}$. Hence $0 = \sum_{p \in \mathbb{I}_{0,2\ell-1}} b_{i,p} \xi^{pj}$ for all $0 \leq j < \ell$ by (3.27), and the same argument shows that $b_{i,p} = 0$ for all $p \in \mathbb{I}_{0,2\ell-1}$.

If $\ell' = \ell$ is odd, then $e_{i,p} + e_{i,p+\ell} \stackrel{*}{=} 0$ for all $p \in \mathbb{I}_{0,\ell-1}$ by (3.26). Similarly as above, we consider the algebra maps $\tilde{\Xi}_{i,j} : \mathcal{A}'[\mathbb{Z}^j] \rightarrow \mathcal{A}'$ such that $K_i \mapsto -v^j$ and $K_r \mapsto 1$ for $r \neq i$; we get algebra maps $\tilde{\Xi}_{i,j} : V_{\mathcal{B}}^0 \rightarrow \mathcal{B}$ such that

$$\tilde{\Xi}_{i,j}(K_i) = -\xi^j, \quad \tilde{\Xi}_{i,j}(k_{i,\ell}) = -\begin{bmatrix} j \\ \ell \end{bmatrix}_\xi, \quad \tilde{\Xi}_{i,j}(K_r) = 1, \quad \tilde{\Xi}_{i,j}(k_{r,\ell}) = 0, \quad r \neq i.$$

Applying $\tilde{\Xi}_{i,j}$ to the previous equality (3.25), we see that

$$0 = \sum_{p \in \mathbb{I}_{0,\ell-1}} (-1)^i (e_{i,p} - e_{i,p+\ell}) \xi^{pj}, \quad 0 \leq j < \ell.$$

Hence $e_{i,p} - e_{i,p+\ell} = 0$ for all $p \in \mathbb{I}_{0,\ell-1}$, so $e_{i,p} = 0$ for all $p \in \mathbb{I}_{0,2\ell-1}$. Analogously $b_{i,p} = 0$ for all $p \in \mathbb{I}_{0,2\ell-1}$. \square

4. THE ALGEBRA V , SIMPLY-LACED DIAGRAM

4.1. Definitions and first properties. As in [Lu3], we fix a finite Cartan matrix $A = (a_{ij})_{i,j \in \mathbb{I}}$ whose Dynkin diagram is connected and simply-laced, that is, of type A, D or E.

Following [Lu3, 2.3, pp. 268 ff] we consider the \mathcal{A} -algebra V presented by generators (3.1), $E_i^{(N)}$, $F_i^{(N)}$, $i \in \mathbb{I}$, $N \in \mathbb{N}_0$ with relations (g5), ... (g10), together with the following, tagged again as in *loc. cit.*,

$$(d1) \quad E_i^{(N)} E_i^{(M)} = \begin{bmatrix} N+M \\ M \end{bmatrix}_v E_i^{(N+M)}, \quad E_i^{(0)} = 1;$$

$$(f1) \quad F_i^{(N)} F_i^{(M)} = \begin{bmatrix} N+M \\ M \end{bmatrix}_v F_i^{(N+M)}, \quad F_i^{(0)} = 1;$$

if $i \neq j \in \mathbb{I}$, $a_{ij} = 0$:

$$(d2) \quad E_i^{(N)} E_j^{(M)} = E_j^{(M)} E_i^{(N)},$$

$$(f2) \quad F_i^{(N)} F_j^{(M)} = F_j^{(M)} F_i^{(N)},$$

if $i \neq j \in \mathbb{I}$, $a_{ij} = -1$, $i < j$:

$$(d3) \quad E_i^{(N)} E_j^{(M)} = \sum_{t=0}^{\min\{M,N\}} v^{t+(N-t)(M-t)} E_j^{(M-t)} E_{ij}^{(t)} E_i^{(N-t)},$$

$$(d4) \quad v^{NM} E_i^{(N)} E_{ij}^{(M)} = E_{ij}^{(M)} E_i^{(N)},$$

$$(d5) \quad v^{NM} E_{ij}^{(M)} E_j^{(N)} = E_j^{(N)} E_{ij}^{(M)},$$

$$(f3) \quad F_i^{(N)} F_j^{(M)} = \sum_{t=0}^{\min\{M,N\}} v^{-t-(N-t)(M-t)} F_j^{(M-t)} F_{ij}^{(t)} F_i^{(N-t)},$$

$$(f4) \quad v^{NM} F_i^{(N)} F_{ij}^{(M)} = F_{ij}^{(M)} F_i^{(N)},$$

$$(f5) \quad v^{NM} F_{ij}^{(M)} F_j^{(N)} = F_j^{(N)} F_{ij}^{(M)},$$

where $E_{ij}^{(N)} = \sum_{k=0}^N (-1)^{N-k} v^{-k} E_i^{(k)} E_j^{(N)} E_i^{(N-k)}$ (cf. [Lu3, Lemma 2.5 (d)]),

$$F_{ij}^{(N)} = \sum_{k=0}^N (-1)^{N-k} v^{-k} F_i^{(k)} F_j^{(N)} F_i^{(N-k)};$$

$$(h1) \quad E_i^{(N)} F_j^{(M)} = F_j^{(M)} E_i^{(N)}, \quad i \neq j,$$

$$(h2) \quad E_i^{(N)} F_i^{(M)} = \sum_{0 \leq t \leq \min\{N, M\}} F_i^{(M-t)} \begin{bmatrix} K_i; 2t - N - M \\ t \end{bmatrix} E_i^{(N-t)},$$

$$(h3) \quad K_i^{\pm 1} E_j^{(N)} = v^{\pm N a_{ij}} E_j^{(N)} K_i^{\pm 1},$$

$$(h4) \quad K_i^{\pm 1} F_j^{(N)} = v^{\mp N a_{ij}} F_j^{(N)} K_i^{\pm 1},$$

$$(h5) \quad \begin{bmatrix} K_i; c \\ t \end{bmatrix} E_j^{(N)} = E_j^{(N)} \begin{bmatrix} K_i; c + N a_{ij} \\ t \end{bmatrix},$$

$$(h6) \quad \begin{bmatrix} K_i; c \\ t \end{bmatrix} F_j^{(N)} = F_j^{(N)} \begin{bmatrix} K_i; c - N a_{ij} \\ t \end{bmatrix}.$$

Let V^+ , respectively V^- , be the subalgebra of V generated by $E_i^{(N)}$, respectively $F_i^{(N)}$, $i \in \mathbb{I}$, $N \in \mathbb{N}_0$. Let

$$(4.1) \quad \begin{bmatrix} K_i^{-1}; c \\ t \end{bmatrix} = \mathcal{S} \left(\begin{bmatrix} K_i; c \\ t \end{bmatrix} \right).$$

The following formula is analogous to (h2), cf. [Lu5, Corollary 3.19]:

$$(4.2) \quad F_i^{(N)} E_i^{(M)} = \sum_{0 \leq t \leq \min\{N, M\}} E_i^{(M-t)} \begin{bmatrix} K_i^{-1}; 2t - N - M \\ t \end{bmatrix} F_i^{(N-t)}.$$

By [Lu3, Proposition 4.8, p. 287], we know that V has a unique Hopf algebra structure determined by (3.8) and

$$(4.3) \quad \begin{aligned} \Delta(E_i^{(N)}) &= \sum_{0 \leq b \leq N} v^{b(N-b)} E_i^{(N-b)} K_i^b \otimes E_i^{(b)}, \\ \Delta(F_i^{(N)}) &= \sum_{0 \leq a \leq N} v^{-a(N-a)} F_i^{(a)} \otimes K_i^{-a} F_i^{(N-a)}, \end{aligned} \quad i \in \mathbb{I}, N \in \mathbb{N}_0.$$

4.2. Specializations of V . We define next

$$V_{\mathcal{B}}^+ = V^+ \otimes_{\mathcal{A}} \mathcal{B}, \quad V_{\mathcal{B}}^- = V^- \otimes_{\mathcal{A}} \mathcal{B}, \quad V_{\mathcal{B}} = V \otimes_{\mathcal{A}} \mathcal{B}.$$

By [Lu2, Proposition 3.2 (b)], $V_{\mathcal{B}}^+$ is generated by E_i and $E_i^{(\ell)}$, $i \in \mathbb{I}$; $V_{\mathcal{B}}^-$ is generated by F_i and $F_i^{(\ell)}$, $i \in \mathbb{I}$. From now on, we abbreviate

$$k_{i,N} = \begin{bmatrix} K_i; 0 \\ N \end{bmatrix}, \quad N \in \mathbb{N}_0, \quad E_i = E_i^{(1)}, \quad F_i = F_i^{(1)}.$$

Lemma 4.1. *Let $i, j \in \mathbb{I}$. We have in $V_{\mathcal{B}}$:*

$$(4.4) \quad k_{i,\ell} E_i = E_i (k_{i,\ell} + \xi^{-2} [2]_{\xi} K_i^{-1} k_{i,\ell-1} + \xi^{-4} K_i^{-2} k_{i,\ell-2}),$$

$$(4.5) \quad k_{i,\ell} F_i = F_i \left(k_{i,\ell} + \sum_{j \in \mathbb{I}_{\ell-2}} (-1)^j \xi^{-2j} [j+1]_{\xi} K_i^j k_{i,\ell-j} - K_i^{\ell} \right),$$

$$(4.6) \quad k_{i,\ell} E_j = E_j k_{i,\ell}, \quad i \neq j, a_{ij} = 0,$$

$$(4.7) \quad k_{i,\ell} F_j = F_j k_{i,\ell}, \quad i \neq j, a_{ij} = 0,$$

$$(4.8) \quad k_{i,\ell}E_j = \xi^\ell E_j \left(\sum_{s=0}^{\ell} (-\xi)^{-s} K_i^s k_{i,\ell-s} \right), \quad i \neq j, a_{ij} = -1,$$

$$(4.9) \quad k_{i,\ell}F_j = \xi^\ell F_j (k_{i,\ell} + \xi^{-1} K_i^{-1} k_{i,\ell-1}), \quad i \neq j, a_{ij} = -1,$$

$$(4.10) \quad k_{i,\ell}E_j^{(\ell)} = E_j^{(\ell)} (k_{i,\ell} + a_{ij} K_i^\ell),$$

$$(4.11) \quad k_{i,\ell}F_j^{(\ell)} = F_j^{(\ell)} (k_{i,\ell} - a_{ij} K_i^\ell).$$

Proof. We consider first the case $j = i$. We take $t = \ell$, $c = 0$ and $N = 1$ in (h6) and use (g9) to obtain (4.5):

$$\begin{aligned} k_{i,\ell}F_i &= F_i \left[\begin{matrix} K_i; -2 \\ \ell \end{matrix} \right] = F_i \left(\sum_{0 \leq j \leq \ell} (-1)^j \xi^{2(\ell-j)} \begin{bmatrix} j+1 \\ j \end{bmatrix}_\xi K_i^j k_{i,\ell-j} \right) \\ &= F_i \left(k_{i,\ell} + \sum_{j \in \mathbb{I}_{\ell-2}} (-1)^j \xi^{-2j} [j+1]_\xi K_i^j k_{i,\ell-j} + (-\xi)^\ell K_i^\ell \right). \end{aligned}$$

For (4.11), we take $t = \ell$, $c = 0$ and $N = \ell$ in (h6) and use (g9):

$$\begin{aligned} k_{i,\ell}F_i^{(\ell)} &= F_i^{(\ell)} \left[\begin{matrix} K_i; -2\ell \\ \ell \end{matrix} \right] = F_i^{(\ell)} \left(\sum_{0 \leq j \leq \ell} (-1)^j \xi^{2\ell(\ell-j)} \begin{bmatrix} 2\ell+j-1 \\ j \end{bmatrix}_\xi K_i^j k_{i,\ell-j} \right) \\ &= F_i^{(\ell)} \left(k_{i,\ell} + (-1)^\ell \xi^{2\ell(\ell-1)} 2K_i^\ell \right) = F_i^{(\ell)} \left(k_{i,\ell} - 2K_i^\ell \right). \end{aligned}$$

Now we take $j \neq i$ and $a_{ij} = 0$. From (h6), $k_{i,\ell}F_j^{(N)} = F_j^{(N)} k_{i,\ell}$ for all $N \in \mathbb{N}$, hence we obtain (4.7) when $N = 1$, and (4.11) when $N = \ell$.

Next we take $j \neq i$ and $a_{ij} = -1$. From (h6) and (g10) we derive (4.9) when $N = 1$, and (4.11) when $N = \ell$:

$$\begin{aligned} k_{i,\ell}F_j &= F_j \left[\begin{matrix} K_i; 1 \\ \ell \end{matrix} \right] = F_j \left(\sum_{s=0}^{\ell} \xi^{\ell-s} \begin{bmatrix} 1 \\ s \end{bmatrix}_\xi K_i^{-s} \begin{bmatrix} K_i; 0 \\ \ell-s \end{bmatrix} \right) \\ &= \xi^\ell F_j (k_{i,\ell} + \xi^{-1} K_i^{-1} k_{i,\ell-1}), \\ k_{i,\ell}F_j^{(\ell)} &= F_j^{(\ell)} \left[\begin{matrix} K_i; \ell \\ \ell \end{matrix} \right] = F_j^{(\ell)} \left(\sum_{s=0}^{\ell} \xi^{\ell(\ell-s)} \begin{bmatrix} \ell \\ s \end{bmatrix}_\xi K_i^{-s} \begin{bmatrix} K_i; 0 \\ \ell-s \end{bmatrix} \right) = F_j^{(\ell)} (k_{i,\ell} + K_i^{-\ell}). \end{aligned}$$

Finally we get (4.4), (4.6), (4.8) and (4.10) similarly but from (h5). \square

Now we compute relations involving $h_{i,\ell}$. The formulas (4.12) and (4.14) appear in [Le2, Theorem 3.1].

Lemma 4.2. *Let $i, j \in \mathbb{I}$. We have in $V_{\mathcal{B}}$:*

$$(4.12) \quad h_{i,\ell}E_j = E_j h_{i,\ell} + a_{ij} \ell^{-1} E_j,$$

$$(4.13) \quad h_{i,\ell}E_j^{(\ell)} = E_j^{(\ell)} h_{i,\ell} + a_{ij} E_j^{(\ell)},$$

$$(4.14) \quad h_{i,\ell}F_j = F_j h_{i,\ell} - a_{ij} \ell^{-1},$$

$$(4.15) \quad h_{i,\ell}F_j^{(\ell)} = F_j^{(\ell)} h_{i,\ell} - a_{ij} F_j^{(\ell)}.$$

Proof. By (3.16), there exist $\mathbf{b}_t \in \mathcal{B}$ such that

$$(4.16) \quad h_{i,\ell} = \left(\sum_{s \in \mathbb{I}_\ell} \frac{p_{\ell,s}}{\ell} k_{i,s} K_i^{\phi_\ell(s)} \right) K_i^\ell = k_{i,\ell} K_i^\ell + \sum_{t=0}^{\ell-1} \mathbf{b}_t K_i^{2t}.$$

Indeed, for each $s \in \mathbb{I}_{\ell-1}$,

$$k_{i,s} = \frac{1}{[s]_\xi!} \prod_{j=0}^{s-1} \frac{\xi^{-j} K_i - \xi^j K_i^{-1}}{\xi - \xi^{-1}} \in \sum_{p=0}^s \mathcal{B} K_i^{2p-s}.$$

Using (h3), (4.10) and (3.21),

$$\begin{aligned} h_{i,\ell} E_j^{(\ell)} &= (k_{i,\ell} K_i^\ell + \sum_{t=0}^{\ell-1} \mathbf{b}_t K_i^{2t}) E_j^{(\ell)} \\ &= \xi^{a_{ij} \ell^2} E_j^{(\ell)} \left(k_{i,\ell} + a_{ij} K_i^\ell \right) K_i^\ell + \sum_{t=0}^{\ell-1} \mathbf{b}_t \xi^{2ta_{ij} \ell} E_j^{(\ell)} K_i^t = E_j^{(\ell)} h_{i,\ell} + a_{ij} E_j^{(\ell)}. \end{aligned}$$

The proof of (4.15) is similar. Next we check (4.12). By a direct computation,

$$\Delta([h_{i,\ell}, E_j]) = [h_{i,\ell}, E_j] \otimes 1 + K_j \otimes [h_{i,\ell}, E_j].$$

Thus $[h_{i,\ell}, E_j]$ is $(1, K_j)$ -primitive and belongs to the subalgebra generated by K_j , $h_{i,\ell}$ and E_j , so $[h_{i,\ell}, E_j] = c_{ij} E_j + d_{ij} (1 - K_j)$ for some $c_{ij}, d_{ij} \in \mathcal{B}$. As $E_j^\ell = 0$, we have that

$$\begin{aligned} 0 &= [h_{i,\ell}, E_j^\ell] = \sum_{k \in \mathbb{I}_\ell} E_j^{k-1} [h_{i,\ell}, E_j] E_j^{\ell-k} \\ &= \sum_{k \in \mathbb{I}_\ell} E_j^{k-1} (c_{ij} E_j + d_{ij} (1 - K_j)) E_j^{\ell-k} = d_{ij} \sum_{k \in \mathbb{I}_\ell} E_j^{k-1} (1 - K_j) E_j^{\ell-k} \\ &= d_{ij} \left(\ell E_j^{\ell-1} - \left(\sum_{k \in \mathbb{I}_\ell} \xi^{-2k} \right) E_j^{\ell-1} K_j \right) = d_{ij} \ell E_j^{\ell-1}. \end{aligned}$$

Hence $d_{ij} = 0$. Analogously $[h_{i,\ell}, F_j] = c'_{ij} F_j$ for some $c'_{ij} \in \mathcal{B}$. Now

$$0 = [h_{i,\ell}, K_j - K_j^{-1}] = (\xi - \xi^{-1}) [h_{i,\ell}, [E_j, F_j]] = (c_{ij} + c'_{ij}) (K_j - K_j^{-1}),$$

so $c'_{ij} = -c_{ij}$. We consider three cases:

- (1) $j \neq i$, $a_{ij} = 0$. Then $[h_{i,\ell}, E_j] = 0$ by (4.6) and (h3). Thus $c_{ij} = 0$.
- (2) $j \neq i$, $a_{ij} = -1$. From (4.9), (4.16) and (h4):

$$\begin{aligned} -c_{ij} F_j &= [h_{i,\ell}, F_j] = [k_{i,\ell} K_i^\ell, F_j] + \sum_{t=0}^{\ell-1} \mathbf{b}_t [K_i^{2t}, F_j] \\ &= \xi^{-1} F_j K_i^{\ell-1} k_{i,\ell-1} + \sum_{t=0}^{\ell-1} \mathbf{b}_t (\xi^{2t} - 1) F_j K_i^{2t}. \end{aligned}$$

As the set $\{F_j K_i^{2t} : t \in \mathbb{I}_{0, \ell-1}\}$ is linearly independent,

$$c_{ij} = -\xi^{-1} b_0$$

where $b_t \in \mathcal{B}$ denote the elements satisfying

$$k_{i, \ell-1} = \frac{1}{[\ell-1]_\xi!} \prod_{j=0}^{\ell-2} \frac{\xi^{-j} K_i - \xi^j K_i^{-1}}{\xi - \xi^{-1}} = \sum_{t=0}^{\ell-1} b_t K_i^{2t-\ell+1}.$$

Since

$$b_0 = \frac{1}{[\ell-1]_\xi!} \frac{\prod_{j=0}^{\ell-2} -\xi^j}{(\xi - \xi^{-1})^{\ell-1}} = \frac{(-1)^{\ell-1} \xi^{\binom{\ell-1}{2}}}{\xi^{\binom{\ell}{2}} \prod_{j \in \mathbb{I}_{\ell-1}} 1 - \xi^{-2j}} = \frac{\xi}{\ell},$$

we have that $c_{ij} = -\ell^{-1}$.

(3) $j = i$. The proof is analogous to the previous case, using (4.4).

Hence $c_{ij} = \frac{a_{ij}}{\ell}$ in all the cases, so (4.12) and (4.14) follow. \square

4.3. The Hopf algebra structure of $V_{\mathcal{B}}$. Recall that by Theorem 3.10, $V_{\mathcal{B}}^0 = \mathcal{B}[K_i, h_{i, \ell} : i \in \mathbb{I}] \simeq \mathcal{B}\Gamma \otimes U(\mathfrak{h})$.

Remark 4.3. The counit on the elements $\begin{bmatrix} K_i^{\pm 1}; c \\ t \end{bmatrix}$ takes the following values.

$$(4.17) \quad \varepsilon \left(\begin{bmatrix} K_i^{\pm 1}; c \\ t \end{bmatrix} \right) = \begin{cases} 1 & \text{if } c = t = 0, \\ 0 & \text{if } c = 0 \text{ and } t \neq 0, \\ \begin{bmatrix} c \\ t \end{bmatrix}_\xi & \text{if } c > 0, \\ (-1)^t \begin{bmatrix} -c+t-1 \\ t \end{bmatrix}_\xi & \text{if } c < 0. \end{cases}$$

In fact, we first note that $\varepsilon \left(\begin{bmatrix} K_i^{-1}; c \\ t \end{bmatrix} \right) = \varepsilon \mathcal{S} \left(\begin{bmatrix} K_i; c \\ t \end{bmatrix} \right) = \varepsilon \left(\begin{bmatrix} K_i; c \\ t \end{bmatrix} \right)$ by (4.1). The formula for $c = 0$ holds by (3.5). Then, for $c > 0$, we use (g10):

$$\varepsilon \left(\begin{bmatrix} K_i; c \\ t \end{bmatrix} \right) = \sum_{0 \leq j \leq t} v^{c(t-j)} \begin{bmatrix} c \\ j \end{bmatrix}_v \varepsilon(K_i^{-j}) \varepsilon \left(\begin{bmatrix} K_i; 0 \\ t-j \end{bmatrix} \right) = \begin{bmatrix} c \\ t \end{bmatrix}_v.$$

While for $c < 0$, we use (g9):

$$\begin{aligned} \varepsilon \left(\begin{bmatrix} K_i; c \\ t \end{bmatrix} \right) &= \sum_{0 \leq j \leq t} (-1)^j v^{-c(t-j)} \begin{bmatrix} -c+j-1 \\ j \end{bmatrix}_v \varepsilon(K_i^j) \varepsilon \left(\begin{bmatrix} K_i; 0 \\ t-j \end{bmatrix} \right) \\ &= (-1)^t \begin{bmatrix} -c+t-1 \\ t \end{bmatrix}_v \end{aligned}$$

Theorem 4.4. *The Hopf algebra $V_{\mathcal{B}}$ has a triangular decomposition given by a TD-datum $(V_{\mathcal{B}}^+, V_{\mathcal{B}}^-, \rightarrow, \leftarrow, \#)$ over $V_{\mathcal{B}}^0$. The left action \rightarrow of $V_{\mathcal{B}}^-$ on*

$V_{\mathcal{B}}^+$, the right action \leftarrow of $V_{\mathcal{B}}^+$ on $V_{\mathcal{B}}^-$ and the map $\sharp : V_{\mathcal{B}}^- \otimes V_{\mathcal{B}}^+ \rightarrow V_{\mathcal{B}}^0$ are determined as follows:

$$(4.18) \quad \begin{aligned} F_i^{(N)} \rightharpoonup E_j^{(M)} &= \delta_{ij}(-1)^N \begin{bmatrix} M-1 \\ N \end{bmatrix}_{\xi} E_i^{(M-N)} \\ F_i^{(N)} \leftarrow E_j^{(M)} &= \delta_{ij}(-1)^M \begin{bmatrix} N-1 \\ M \end{bmatrix}_{\xi} F_i^{(N-M)} \\ F_i^{(N)} \sharp E_j^{(M)} &= \delta_{M,N} \delta_{ij} \begin{bmatrix} K_i^{-1}; 0 \\ N \end{bmatrix}, \end{aligned}$$

cf. (4.1), where $E_i^{(n)} = 0 = F_i^{(n)}$ if $n < 0$.

Proof. For the first claim, we just need to verify that the conditions of Proposition 2.1 (b) hold.

Let $V_{\mathcal{B}}^{\geq 0} := V_{\mathcal{B}}^+ V_{\mathcal{B}}^0$ and $V_{\mathcal{B}}^{\leq 0} := V_{\mathcal{B}}^0 V_{\mathcal{B}}^-$; these are Hopf subalgebras of $V_{\mathcal{B}}$ by definition. It is easy to see that the inclusions $V_{\mathcal{B}}^0 \hookrightarrow V_{\mathcal{B}}^{\geq 0}$ and $V_{\mathcal{B}}^0 \hookrightarrow V_{\mathcal{B}}^{\leq 0}$ admit Hopf algebra sections π^+ and π^- respectively and that

$$V_{\mathcal{B}}^+ = \left(V_{\mathcal{B}}^{\geq 0} \right)^{\text{co } \pi^+}, \quad V_{\mathcal{B}}^- = \text{co } \pi^- \left(V_{\mathcal{B}}^{\geq 0} \right).$$

Thus $V_{\mathcal{B}}^+$ is a Hopf algebra in $\frac{V_{\mathcal{B}}^0}{V_{\mathcal{B}}^0} \mathcal{YD}$ and $V_{\mathcal{B}}^{\geq 0} \simeq V_{\mathcal{B}}^+ \# V_{\mathcal{B}}^0$, respectively $V_{\mathcal{B}}^-$ is a Hopf algebra in $\mathcal{YD}_{\frac{V_{\mathcal{B}}^0}{V_{\mathcal{B}}^0}}$ and $V_{\mathcal{B}}^{\leq 0} \simeq V_{\mathcal{B}}^0 \# V_{\mathcal{B}}^-$. Also, by [Lu3, Theorem 4.5 (a)], the multiplication induces a linear isomorphism $V_{\mathcal{B}}^+ \otimes V_{\mathcal{B}}^0 \otimes V_{\mathcal{B}}^- \simeq V_{\mathcal{B}}$. Thus we may apply Proposition 2.1 (b).

The verification of (4.18) is direct using the formulas in the proof of [S, Theorem 3.5] and the natural projections $\varpi^{\star} : V_{\mathcal{B}} \rightarrow V_{\mathcal{B}}^{\star}$, for $\star \in \{+, 0, -\}$. In fact, $F_i^{(N)} \rightharpoonup E_j^{(M)} = \varpi^+(F_i^{(N)} E_j^{(M)})$. If $i \neq j$, this is zero by (h1). Otherwise, we use (4.2):

$$\begin{aligned} F_i^{(N)} \rightharpoonup E_i^{(M)} &= \varpi^+(F_i^{(N)} E_i^{(M)}) \\ &= \sum_{0 \leq t \leq \min\{N, M\}} E_i^{(M-t)} \varepsilon \left(\begin{bmatrix} K_i^{-1}; 2t - N - M \\ t \end{bmatrix} \right) \varepsilon \left(F_i^{(N-t)} \right) \end{aligned}$$

which is zero for $N \geq M$ by Remark 4.3. If $N < M$, then

$$F_i^{(N)} \rightharpoonup E_i^{(M)} = E_i^{(M-N)} \varepsilon \left(\begin{bmatrix} K_i^{-1}; N - M \\ N \end{bmatrix} \right) = (-1)^N \begin{bmatrix} M-1 \\ N \end{bmatrix} E_i^{(M-N)}.$$

We can verify the formulas for \leftarrow and \sharp in a similar way.

We next show by induction that (4.18) completely determines \rightharpoonup , \leftarrow and \sharp . We will use that the comultiplication of $V_{\mathcal{B}}^{\pm}$ in the respective Yetter-Drinfeld

category is given by

$$\begin{aligned}\Delta(E_i^{(N)}) &= \sum_{0 \leq b \leq N} v^{b(N-b)} E_i^{(N-b)} \otimes E_i^{(b)}, \\ \Delta(F_i^{(N)}) &= \sum_{0 \leq a \leq N} v^{-a(N-a)} F_i^{(a)} \otimes F_i^{(N-a)},\end{aligned}\quad i \in \mathbb{I}, N \in \mathbb{N}_0.$$

This follows from (4.3).

Let $E = E_{j_1}^{(M_1)} \cdots E_{j_r}^{(M_r)}$ and $F = F_{i_1}^{(N_1)} \cdots F_{i_s}^{(N_s)}$. First, we assume that

$$F_i^{(N)} \rightharpoonup E, \quad F \leftarrow E_j^{(M)}, \quad F_i^{(N)} \sharp E \quad \text{and} \quad F \sharp E_j^{(M)}$$

are determined by (4.18) for all $r, s \leq n$ and prove the same claim for $s = r = n + 1$. By (2.10), we have that

$$(4.19) \quad \begin{aligned}F_i^{(N)} \rightharpoonup (E_j^{(M)} E) &= \left((F_i^{(N)})_{(0)}^{(1)} \rightharpoonup (E_j^{(M)})_{(0)}^{(1)} \right) \times \\ &\times \left((F_i^{(N)})_{(1)}^{(1)} \left((F_i^{(N)})_{(1)}^{(2)} \sharp (E_j^{(M)})_{(1)}^{(2)} \right) (E_j^{(M)})_{(-1)}^{(3)} \right) \triangleright \\ &\left(\left((F_i^{(N)})_{(0)}^{(3)} \leftarrow (E_j^{(M)})_{(0)}^{(3)} \right) \rightharpoonup E \right).\end{aligned}$$

Hence, (4.19) is determined by (4.18) because of the inductive hypothesis.

The same holds for $F_i^{(N)} \sharp (E_j^{(M)} E)$ since

$$\begin{aligned}F_i^{(N)} \sharp (E_j^{(M)} E) &= \left((F_i^{(N)})_{(0)}^{(1)} \sharp (E_j^{(M)})_{(0)}^{(1)} \right) \times \\ &\left((E_j^{(M)})_{(-1)}^{(2)} \left(\left((F_i^{(N)})_{(0)}^{(2)} \leftarrow (E_j^{(M)})_{(0)}^{(2)} \right) \sharp E \right) \right)\end{aligned}$$

by (2.9). A similar argument works for $(FF_i^{(N)}) \leftarrow E_j^{(M)}$ and $(FF_i^{(N)}) \sharp E_j^{(M)}$.

Second, we prove that

$$F \rightharpoonup E, \quad F \leftarrow E \quad \text{and} \quad F \sharp E$$

are determined by (4.18) for all $r, s \geq 0$. This is true for $F \rightharpoonup E$ and $F \leftarrow E$ because \rightharpoonup and \leftarrow are actions. For the others we proceed again by induction on r (or on s) using (2.9); notice that the initial inductive step $r = 1$ was proved above. \square

Remark 4.5. Here are some particular instances of the first line in (4.18):

$$\begin{aligned}F_i \rightharpoonup E_j^{(M)} &= F_i^{(\ell)} \rightharpoonup E_j^{(M)} = 0, & \text{if } i \neq j, \\ F_i \rightharpoonup E_i^{(M)} &= (-1)^{M-1} [M-1]_{\xi} E_i^{(M-1)}, \\ F_i^{(\ell)} \rightharpoonup E_i^{(M)} &= 0 & \text{if } \ell \text{ does not divide } M, \\ F_i^{(\ell)} \rightharpoonup E_i^{(\ell n)} &= (-1)^{n-1} (n-1) E_i^{(\ell n - \ell)}.\end{aligned}$$

Remark 4.6. The structure of $V_{\mathcal{B}}^+$ as an object in ${}_{V_{\mathcal{B}}^0}^{V_{\mathcal{B}}^0}\mathcal{YD}$ is as follows: the (left) action of $V_{\mathcal{B}}^0$ on $V_{\mathcal{B}}^+$ is given by (h3), (4.12) and (4.13), while the coaction $\lambda : V_{\mathcal{B}}^+ \rightarrow V_{\mathcal{B}}^0 \otimes V_{\mathcal{B}}^+$ is determined by

$$\lambda(E_i^{(N)}) = K_i^N \otimes E_i^{(N)}, \quad i \in \mathbb{I}, N \in \mathbb{N}.$$

Analogously, the structure of $V_{\mathcal{B}}^-$ as an object in $\mathcal{YD}_{V_{\mathcal{B}}^0}^{V_{\mathcal{B}}^0}$ is as follows: the (right) action of $V_{\mathcal{B}}^0$ on $V_{\mathcal{B}}^-$ is given by (h4), (4.14) and (4.15); meanwhile the coaction $\rho : V_{\mathcal{B}}^- \rightarrow V_{\mathcal{B}}^- \otimes V_{\mathcal{B}}^0$ is determined by

$$\rho(F_i^{(N)}) = F_i^{(N)} \otimes K_i^{-N}, \quad i \in \mathbb{I}, N \in \mathbb{N}.$$

4.4. The multiply-laced diagrams. The arguments above can be extended to the diagrams of types B, C, F, G. We just discuss the torus part here.

Let $\mathbf{d} = (d_i)_{i \in \mathbb{I}} \in \mathbb{N}^{\mathbb{I}}$. Following [Lu4, 6.4] we consider the \mathcal{A} -algebra \mathbb{V}^0 that is a (multiply-laced!) variation of the V^0 studied so far. For the agility of the exposition we do not stress \mathbf{d} in the notation. This \mathbb{V}^0 is presented by the generators analogous to those (3.1) of V^0 :

$$(4.20) \quad \mathbb{K}_i, \quad \mathbb{K}_i^{-1}, \quad \begin{bmatrix} \mathbb{K}_i; c \\ t \end{bmatrix}, \quad i \in \mathbb{I}, c \in \mathbb{Z}, t \in \mathbb{N}_0$$

with slightly modified relations. Tagging them as in [Lu4], these are:

(b1) the generators (4.20) commute with each other,

$$(b2) \quad \mathbb{K}_i \mathbb{K}_i^{-1} = 1, \quad \begin{bmatrix} \mathbb{K}_i; c \\ 0 \end{bmatrix} = 1,$$

$$(b3) \quad \begin{bmatrix} \mathbb{K}_i; 0 \\ t \end{bmatrix} \begin{bmatrix} \mathbb{K}_i; -t \\ t' \end{bmatrix} = \begin{bmatrix} t + t' \\ t \end{bmatrix}_{v^{d_i}} \begin{bmatrix} \mathbb{K}_i; 0 \\ t + t' \end{bmatrix}, \quad t, t' \geq 0,$$

$$(b4) \quad \begin{bmatrix} \mathbb{K}_i; c \\ t \end{bmatrix} - v^{-d_i t} \begin{bmatrix} \mathbb{K}_i; c + 1 \\ t \end{bmatrix} = -v^{-d_i(c+1)} \mathbb{K}_i^{-1} \begin{bmatrix} \mathbb{K}_i; c \\ t - 1 \end{bmatrix}, \quad t \geq 1,$$

$$(b5) \quad (v^{d_i} - v^{-d_i}) \begin{bmatrix} \mathbb{K}_i; 0 \\ 1 \end{bmatrix} = \mathbb{K}_i - \mathbb{K}_i^{-1}.$$

The algebra \mathbb{V}^0 is related to V^0 in the following way. For $i \in \mathbb{I}$, let V_i^0 , respectively \mathbb{V}_i^0 , be the subalgebra of V^0 , respectively \mathbb{V}^0 , generated by $K_i^{\pm 1}$ and $\begin{bmatrix} K_i; c \\ t \end{bmatrix}$, respectively $\mathbb{K}_i^{\pm 1}$ and $\begin{bmatrix} \mathbb{K}_i; c \\ t \end{bmatrix}$, $c \in \mathbb{Z}$, $t \in \mathbb{N}_0$. Then (g6) and Proposition 3.1, respectively (b1) and [Lu4, Theorem 6.7] imply that there are algebra isomorphisms

$$(4.21) \quad V^0 \simeq V_1^0 \otimes V_2^0 \otimes \dots \otimes V_{\theta}^0, \quad \mathbb{V}^0 \simeq \mathbb{V}_1^0 \otimes \mathbb{V}_2^0 \otimes \dots \otimes \mathbb{V}_{\theta}^0.$$

Lemma 4.7. *Let $\tilde{\mathcal{A}} = \mathcal{A}$ regarded as \mathcal{A} -algebra via $v \mapsto v^{d_i}$. Then $\mathbb{V}_i^0 \simeq V_i^0 \otimes_{\mathcal{A}} \tilde{\mathcal{A}}$ as algebras.*

Proof. First we claim that there is an algebra map $\psi_i : \mathbb{V}_i^0 \rightarrow V_i^0 \otimes_{\mathcal{A}} \tilde{\mathcal{A}}$ given by $\mathbb{K}_i^{\pm 1} \mapsto K_i^{\pm 1} \otimes 1$ and $\begin{bmatrix} \mathbb{K}_i; c \\ t \end{bmatrix} \mapsto \begin{bmatrix} K_i; c \\ t \end{bmatrix} \otimes 1$. Indeed, the images satisfy (b2) by (g7) and (b5) by (g5). Taking $c = t$ and $y = t'$ in (g9) and inserting the right hand side in (g9) we get (b3), while (b4) follows applying (g9) to both sides. The claim is proved and implies in turn that ψ_i is an isomorphism, as it sends a basis to a basis by Proposition 3.1 and [Lu4, Theorem 6.7]. \square

From Lemma 3.2, (4.21) and Lemma 4.7 we see that \mathbb{V}^0 is a Hopf algebra over \mathcal{A} with comultiplication determined by the K_i 's being group-likes. Let

$$\mathbb{k}_{i,t} := \begin{bmatrix} \mathbb{K}_i; 0 \\ t \end{bmatrix},$$

$$\mathbf{h}_{i,n} := \frac{\mathbb{K}_i^n - \mathbb{K}_i^{-n}}{n(v^{d_i n} - v^{-d_i n})} \mathbb{K}_i^n = \frac{1}{n} \left(\sum_{s \in \mathbb{I}_n} p_{n,s}(v^{d_i}) \mathbb{k}_{i,s} \mathbb{K}_i^{\phi_n(s)} \right) \mathbb{K}_i^n \in \mathbb{V}^0,$$

$t, n \in \mathbb{N}$, $i \in \mathbb{I}$. Let $U(\mathfrak{h}) \simeq \mathcal{B}[t_i : i \in \mathbb{I}]$ as above and let $\Gamma = (\mathbb{Z}/2\ell)^{\mathbb{I}}$ with generators $(g_i)_{i \in \mathbb{I}}$. From the previous considerations we conclude:

Proposition 4.8. *Assume that $\xi^{2d_i} \neq 1$ for all $i \in \mathbb{I}$. Then $V_{\mathcal{B}}^0 \simeq \mathcal{B}\Gamma \otimes U(\mathfrak{h})$ as Hopf algebras via $g_i \mapsto \mathbb{K}_i$, $t_i \mapsto \mathbf{h}_{i,\ell}$. \square*

REFERENCES

- [A] N. Andruskiewitsch, *Notes on extensions of Hopf algebras*. Canad. J. Math. **48** (1996), 3–42.
- [AA] N. Andruskiewitsch, I. Angiono, *On Finite dimensional Nichols algebras of diagonal type*. Bull. Math. Sci. **7** 353–573 (2017).
- [AAR1] N. Andruskiewitsch, I. Angiono, F. Rossi Bertone. *The quantum divided power algebra of a finite-dimensional Nichols algebra of diagonal type*, Math. Res. Lett. **24** (2017), 619–643.
- [AAR2] ——— *A finite-dimensional Lie algebra arising from a Nichols algebra of diagonal type (rank 2)*, Bull. Belg. Math. Soc. Simon Stevin **24** (1) (2017), 15–34.
- [AAR3] ——— *Lie algebras arising from Nichols algebras of diagonal type* [arXiv:1911.06586](https://arxiv.org/abs/1911.06586).
- [AAY] N. Andruskiewitsch, I. Angiono, M. Yakimov. *Poisson orders on large quantum groups*, in preparation.
- [Ang1] I. Angiono, *On Nichols algebras of diagonal type*, J. Reine Angew. Math. **683** (2013), 189–251.
- [Ang2] ——— *Distinguished pre-Nichols algebras*, Transform. Groups **21** (2016), 1–33.
- [DK] C. De Concini, V. Kac, *Representations of quantum groups at roots of 1*. Operator algebras, unitary representations, enveloping algebras, and invariant theory (Paris, 1989), 471–506, Progr. Math., **92** (1990), Birkhauser, Boston, MA.
- [DP] C. De Concini, C. Procesi, *Quantum groups*. D-modules, representation theory, and quantum groups, 31–140, Lecture Notes in Math. **1565**, Springer, 1993.
- [H] I. Heckenberger, *Classification of arithmetic root systems*, Adv. Math. **220** (2009) 59–124.
- [L] R. Laugwitz, *Pointed Hopf algebras with triangular decomposition. A characterization of multiparameter quantum groups*, Algebr. Represent. Theory **19** 547–578 (2016).
- [Le1] S. Lentner, *A Frobenius homomorphism for Lusztig’s quantum groups for arbitrary roots of unity*. Commun. Contemp. Math. **18** (2016), no. 3, 1550040, 42 pp.

- [Le2] ———, *The unrolled quantum group inside Lusztig's quantum group of divided powers*. Lett. Math. Phys. **109** (2019), 1665–1682.
- [Lu1] G. Lusztig, *Quantum deformations of certain simple modules over enveloping algebras*, Adv. Math. **70** (1988), 237–249.
- [Lu2] ———, *Modular representations and quantum groups*. Contemp. Math. **82** (1989), 59–77.
- [Lu3] ———, *Finite dimensional Hopf algebras arising from quantized universal enveloping algebras*. J. Amer. Math. Soc. **3** (1990), 257–296.
- [Lu4] ———, *Quantum groups at roots of 1*. Geom. Dedicata **35** (1990), 89–114.
- [Lu5] ———, *Introduction to quantum groups*. Birkhäuser (1993).
- [Ma] S. Majid, *Double-bosonization of braided groups and the construction of $U_q(g)$* . Math. Proc. Camb. Philos. Soc. **125**, 151–192 (1999).
- [M] S. Montgomery. *Hopf algebras and their actions on rings*, CMBS **82**, Amer. Math. Soc. (1993).
- [R] Radford, D. E., *Hopf algebras*, Series on Knots and Everything 49. Hackensack, NJ: World Scientific. xxii, 559 p. (2012).
- [S] Y. Sommerhauser, *Deformed Enveloping Algebras*, New York J. Math. **2** (1996), 35–58.

FACULTAD DE MATEMÁTICA, ASTRONOMÍA Y FÍSICA, UNIVERSIDAD NACIONAL DE CÓRDOBA. CIEM – CONICET. MEDINA ALLENDE S/N (5000) CIUDAD UNIVERSITARIA, CÓRDOBA, ARGENTINA

E-mail address: (andrus|angiono|vay)@famaf.unc.edu.ar