

ON A FAMILY OF DISCRETE LOG-SYMMETRIC DISTRIBUTIONS

H. SAULO, R. VILA, L. PAIVA AND N. BALAKRISHNAN

ABSTRACT. The use of continuous probability distributions has been widespread in problems with purely discrete nature. In general, such distributions are not appropriate in this scenario. In this paper, we introduce a class of discrete and asymmetric distributions based on the family of continuous log-symmetric distributions. Some properties are discussed as well as estimation by the maximum likelihood method. A Monte Carlo simulation study is carried out to evaluate the performance of the estimators, and censored and uncensored data sets are used to illustrate the proposed methodology.

1. INTRODUCTION

Continuous log-symmetric distributions are of particular interest for describing strictly positive and asymmetric data with the possibility of outlier observations; see, for example, Jones (2008), Vanegas and Paula (2016a,b), Saulo and Leão (2017), Balakrishnan et al. (2017), Medeiros and Ferrari (2017) and Ventura et al. (2019), for some discussions and applications of log-symmetric models. A continuous random variable Y follows a log-symmetric distribution if its probability density function (PDF) is given by

$$(1) \quad f_Y(y|\boldsymbol{\theta}) = \frac{(Z_g)^{-1}}{\sqrt{\phi}y} g[a_{\boldsymbol{\theta}}^2(y)], \quad a_{\boldsymbol{\theta}}(y) = \log\left(\frac{y}{\lambda}\right)^{1/\sqrt{\phi}}, \quad y > 0;$$

where $\boldsymbol{\theta} = (\lambda, \phi)$, $\lambda > 0$ is a scale parameter and also the median of Y , $\phi > 0$ is a shape parameter associated with the skewness or relative dispersion, $Z_g = \int_{-\infty}^{\infty} g(w^2) dw$ is the partition function, and the function g is a density generating kernel such that $g(u) > 0$ for $u > 0$. The function g is associated with an additional parameter ξ (or vector $\boldsymbol{\xi}$). We use the notation $Y \sim \text{LS}(\boldsymbol{\theta}, g)$. Note that if $g(u)$ in (1) is $\exp(-u/2)$; $[1 + (u/\xi)]^{-(\xi+1)/2}$, $\xi > 0$; $\exp[-u^{1/(1+\xi)}/2]$, $-1 < \xi \leq 1$; $\sqrt{\xi_2} \exp(-\xi_2 u/2) + [(1 - \xi_1)/\xi_1] \exp(-u/2)$, $0 < \xi_1, \xi_2 < 1$; $\cosh(u^{1/2}) \exp[-(2/\xi^2) \sinh^2(u^{1/2})]$, $\xi > 0$; or $\cosh(u^{1/2}) [\xi_2 \xi_1^2 + 4 \sinh^2(u^{1/2})]^{-(\xi_2+1)/2}$, $\xi_1, \xi_2 > 0$; we have the log-normal, log-Student- t , log-power-exponential, log-contaminated-normal, extended Birnbaum-Saunders or extended Birnbaum-Saunders- t distributions, respectively; see Vanegas and Paula (2016a) and Ventura et al. (2019). If $Y \sim \text{LS}(\boldsymbol{\theta}, g)$, then the associated cumulative distribution function (CDF) is given by $F_Y(y|\boldsymbol{\theta}) = G[a_{\boldsymbol{\theta}}(y)]$, where the function $G : (-\infty, +\infty) \rightarrow [0, 1]$ is defined as

$$(2) \quad G(r) = (Z_g)^{-1} \int_{-\infty}^r g(z^2) dz, \quad -\infty < r < +\infty.$$

This mapping is easily seen to have the following properties:

$$(a) \quad G(0) = 0.5, \quad G(+\infty) = \lim_{r \rightarrow +\infty} G(r) = 1, \quad G(-\infty) = \lim_{r \rightarrow -\infty} G(r) = 0;$$

Date: March 2, 2022.

Key words and phrases. Discrete distributions; Maximum likelihood methods; Monte Carlo simulation; R software.

- (b) $G(\cdot)$ is a continuous function and that $G(\cdot)$ is strictly monotonically increasing. Hence $G(\cdot)$ has an inverse function, denoted by $G^{-1}(\cdot)$;
- (c) From Items (a) and (b), $G(\cdot)$ is a CDF; and
- (d) $G^{-1}(1 - p) = -G^{-1}(p)$ for $p \in (0, 1)$ given.

Despite the huge use of log-symmetric distributions – its most famous member is the log-normal model – they are not appropriate in purely discrete contexts. For example, to model the number of cycles before failure of a equipment or the number of weeks to cure a patient, among others; see Vila et al. (2019). Moreover, despite useful, continuous log-symmetric models do not include the zero. In this paper, we define a discrete random variable associated to Y in (1) as $X = \lfloor Y \rfloor$, where $\lfloor y \rfloor$ denotes the largest integer less than or equal to y . In other words, we propose a class of discrete log-symmetric distributions. The proposed class incorporates every distribution belonging to the log-symmetric family, and it is useful for asymmetric and non-negative discrete data.

The rest of the paper proceeds as follows. In Section 2, we introduce the class of discrete log-symmetric models. In Section 3, we discuss some mathematical properties. In Section 4, estimation of the model parameters are approached via the maximum likelihood method for the censored and uncensored cases. In Section 5, we carry out a simulation study to evaluate the performance of the estimators taking into account different censoring proportions. In Section 6, we illustrate the proposed methodology with two real data sets. Finally, in Section 6, we make some concluding remarks and discuss future work.

2. DISCRETE LOG-SYMMETRIC DISTRIBUTIONS

We say that a discrete random variable X , taking values in the set $\{0, 1, \dots\}$, follows a discrete log-symmetric distribution with parameter vector $\boldsymbol{\theta} = (\lambda, \phi)$, where $\lambda > 0, \phi > 0$, denoted by $X \sim \text{LS}_d(\boldsymbol{\theta}, g)$, if its probability mass function (PMF) is given by

$$(3) \quad p(x|\boldsymbol{\theta}) = G[a_{\boldsymbol{\theta}}(x+1)] - G[a_{\boldsymbol{\theta}}(x)], \quad x = 0, 1, \dots,$$

where $a_{\boldsymbol{\theta}}(\cdot)$ and $G(\cdot)$ are as in (1) and (2), respectively. Note that $G[a_{\boldsymbol{\theta}}(0)] = G(-\infty) = 0$ and that $G[a_{\boldsymbol{\theta}}(+\infty)] = G(+\infty) = 1$. Given the density generating kernel g , defined below Item (1), the parameters λ and ϕ completely determine the PMF (3) at $x = 0$. Since $G(\cdot)$ and $a_{\boldsymbol{\theta}}(\cdot)$ are strictly increasing functions, and

$$\lim_{n \rightarrow \infty} \sum_{x=0}^n p(x|\boldsymbol{\theta}) = \lim_{n \rightarrow \infty} G[a_{\boldsymbol{\theta}}(n+1)] = G(+\infty) = 1,$$

it is clear that $p(x|\boldsymbol{\theta})$ is a PDF.

The CDF, reliability function (RF) and hazard rate (HR) of the LS_d distribution, respectively, are given by

$$F(x|\boldsymbol{\theta}) = 1 - R(x|\boldsymbol{\theta}) = G[a_{\boldsymbol{\theta}}(\lfloor x \rfloor + 1)], \quad x \geq 0;$$

$$H(x|\boldsymbol{\theta}) = \frac{p(x|\boldsymbol{\theta})}{p(x|\boldsymbol{\theta}) + R(x|\boldsymbol{\theta})} = \frac{G[a_{\boldsymbol{\theta}}(x+1)] - G[a_{\boldsymbol{\theta}}(x)]}{1 - G[a_{\boldsymbol{\theta}}(x)]}, \quad x = 0, 1, \dots$$

3. MATHEMATICAL PROPERTIES

This section, if not explicitly mentioned otherwise, consists of mathematical properties valid for any discrete random variable X with support $\{0, 1, \dots\}$.

Let (b_n) be a sequence of real numbers. For technical reasons in the next result we use the convention $\prod_{y=0}^{-1} b_y = 1$. The next result provides a characterization of the PMF and RF of a discrete distribution in terms of the HR.

Proposition 3.1. *If X is a discrete random variable then, for each $x = 0, 1, 2, \dots$,*

$$(a) \quad p(x|\boldsymbol{\theta}) = \frac{H(x|\boldsymbol{\theta})}{1 - H(x|\boldsymbol{\theta})} \prod_{y=0}^{x-1} [1 - H(y|\boldsymbol{\theta})];$$

$$(b) \quad R(x|\boldsymbol{\theta}) = \prod_{y=0}^{x-1} [1 - H(y|\boldsymbol{\theta})];$$

where $H(\cdot|\boldsymbol{\theta})$ is the HR.

Proof. By using the identity $p(x|\boldsymbol{\theta}) = R(x|\boldsymbol{\theta}) - R(x+1|\boldsymbol{\theta}) = [p(x|\boldsymbol{\theta}) + R(x|\boldsymbol{\theta})]H(x|\boldsymbol{\theta})$, $x = 0, 1, \dots$, we have

$$1 = H(x|\boldsymbol{\theta}) + \frac{R(x|\boldsymbol{\theta})H(x|\boldsymbol{\theta})}{p(x|\boldsymbol{\theta})}, \quad x = 0, 1, 2, \dots$$

Since $p(x|\boldsymbol{\theta})/H(x|\boldsymbol{\theta}) = p(x|\boldsymbol{\theta}) + R(x|\boldsymbol{\theta}) = R(x-1|\boldsymbol{\theta})$, it follows that

$$\frac{R(x|\boldsymbol{\theta})}{R(x-1|\boldsymbol{\theta})} = 1 - H(x|\boldsymbol{\theta}), \quad x = 0, 1, 2, \dots$$

Exchanging x for y in the above identity and then multiplying from $y = 0$ to $y = x - 1$, we get

$$R(x|\boldsymbol{\theta}) = \prod_{y=0}^{x-1} \frac{R(y|\boldsymbol{\theta})}{R(y-1|\boldsymbol{\theta})} = \prod_{y=0}^{x-1} [1 - H(y|\boldsymbol{\theta})], \quad x = 0, 1, 2, \dots,$$

verifying the identity for $R(x|\boldsymbol{\theta})$. On the other hand, combining the above identity with the definition of HR, the identity for $p(x|\boldsymbol{\theta})$ follows. \square

3.1. Moments and variance.

Theorem 3.2. *If X is a discrete random variable possessing all the higher-order moments, then*

$$(a) \quad \mathbb{E}(X^r) = \sum_{x=0}^{\infty} [(x+1)^r - x^r] R(x|\boldsymbol{\theta});$$

$$(b) \quad \mathbb{E}(X^r) = \sum_{x=0}^{\infty} \sum_{k=0}^r \sum_{i=0}^{r-k} \binom{r-k}{i} x^{k+i} R(x|\boldsymbol{\theta});$$

$$(c) \quad \text{Var}(X) = 2 \sum_{x=0}^{\infty} x R(x|\boldsymbol{\theta}) + \sum_{x=0}^{\infty} R(x|\boldsymbol{\theta}) \left[1 - \sum_{x=0}^{\infty} R(x|\boldsymbol{\theta}) \right];$$

where $R(\cdot|\boldsymbol{\theta})$ is the RF.

Proof. In order to prove Item (a), using the telescopic series $\sum_{x=0}^{i-1} [(x+1)^r - x^r] = i^r$, it follows that

$$\mathbb{E}(X^r) = \sum_{i=0}^{\infty} \sum_{x=0}^{\infty} \mathbb{1}_{\{x < i\}} [(x+1)^r - x^r] p(i|\boldsymbol{\theta}) = \sum_{x=0}^{\infty} [(x+1)^r - x^r] \sum_{i=0}^{\infty} \mathbb{1}_{\{i > x\}} p(i|\boldsymbol{\theta}),$$

where in the second equality we exchange the orders of the summations because the following series

$$\sum_{x=0}^{\infty} \mathbf{1}_{\{x < i\}} |(x+1)^r - x^r| p(i|\boldsymbol{\theta}) = \sum_{x=0}^{\infty} \mathbf{1}_{\{x < i\}} \cdot [(x+1)^r - x^r] p(i|\boldsymbol{\theta}) = i^r p(i|\boldsymbol{\theta}) =: M_i$$

is finite for each $i = 0, 1, \dots$, and, by hypothesis, the expectation $\mathbb{E}(X^r) = \sum_{i=0}^{\infty} M_i$ always exists. This proves the first item. The second item follows by combining the expression for $\mathbb{E}X^r$ given in the first item with the polynomial identity $a^n - b^n = (a-b) \sum_{k=0}^{n-1} a^{n-k} b^k$ and with the binomial expansion. The proof of the third item immediately follows from Item (a). Thus, the proof is complete. \square

3.2. The p -quantile.

Theorem 3.3. *Let $X = \lfloor Y \rfloor$ be a discrete random variable obtained from a positive continuous random variable Y with CDF $F_Y(\cdot|\boldsymbol{\theta})$. Given $p \in (0, 1)$, let $Q_p = F_Y^{-1}(p|\boldsymbol{\theta})$ be the p -quantile for Y . The following statements are valid:*

- (a) *If $Q_p > 0$ is a natural number, then $Q_p - 1$ is the p -quantile for X ;*
- (b) *If $Q_p > 0$ is not a natural number, then all $y \in [\lfloor Q_p \rfloor, \lfloor Q_p \rfloor + 1)$ is a p -quantile for X .*

Proof. Given $p \in (0, 1)$, assume that $Q_p = F_Y^{-1}(p|\boldsymbol{\theta})$. By using the relations, for all $x > 0$,

$$(4) \quad F_X(x^-|\boldsymbol{\theta}) = F_Y(\lfloor x \rfloor|\boldsymbol{\theta});$$

$$(5) \quad \lfloor x \rfloor \leq x < \lfloor x \rfloor + 1;$$

where $F_X(x^-|\boldsymbol{\theta}) = \lim_{\delta \rightarrow 0} F_X(x - \delta|\boldsymbol{\theta})$ for all $\delta > 0$, we have

$$F_X[(Q_p - 1)^-|\boldsymbol{\theta}] \leq F_X(Q_p - 1|\boldsymbol{\theta}) = F_X(Q_p^-|\boldsymbol{\theta}) \stackrel{(4)}{=} F_Y(Q_p|\boldsymbol{\theta}) = p,$$

whenever $Q_p > 0$ is a natural number. Then, by definition of p -quantile for a discrete random variable, the statement in Item (a) follows.

Already, when $Q_p > 0$ is not a natural number, from (4) and (5) we have

$$F_X(Q_p^-|\boldsymbol{\theta}) \stackrel{(4)}{=} F_Y(\lfloor Q_p \rfloor|\boldsymbol{\theta}) \leq F_Y(Q_p|\boldsymbol{\theta}) = p;$$

$$F_X(Q_p|\boldsymbol{\theta}) = F_X[(Q_p + 1)^-|\boldsymbol{\theta}] \stackrel{(4)}{=} F_Y(\lfloor Q_p \rfloor + 1|\boldsymbol{\theta}) \stackrel{(5)}{\geq} F_Y(Q_p|\boldsymbol{\theta}) = p.$$

Therefore, $F_X(Q_p^-|\boldsymbol{\theta}) \leq p \leq F_X(Q_p|\boldsymbol{\theta})$. Hence, the p -quantile for X can be represented by any value in the interval $[\lfloor Q_p \rfloor, \lfloor Q_p \rfloor + 1)$, and the proof of Item (b) follows. This completes the proof. \square

The following two results are applied exclusively to random variables with discrete log-symmetric distribution.

Proposition 3.4. *Let X be a random variable with $LS_{\mathbf{d}}$ distribution. The following statements hold:*

- (a) *If λ is a natural number, then $\lambda - 1$ is the median for X ;*
- (b) *If λ is not a natural number, then the median of the distribution of X can be represented by any value in the set $[\lfloor \lambda \rfloor, \lfloor \lambda \rfloor + 1)$.*

Proof. Let $X \sim \text{LS}_d(\boldsymbol{\theta}, g)$. Since $G(\cdot)$ and $a_{\boldsymbol{\theta}}(\cdot)$ are strictly increasing functions, the function $G[a_{\boldsymbol{\theta}}(\cdot)]$ is a strictly increasing CDF corresponding to some continuous random variable Y with log-symmetric distribution $\text{LS}(\boldsymbol{\theta}, g)$. Furthermore, note that the median $Q_{0.5}$ for Y can be written as

$$Q_{0.5} = (G \circ a_{\boldsymbol{\theta}})^{-1}(0.5) = a_{\boldsymbol{\theta}}^{-1}[G^{-1}(0.5)] = \lambda \exp \left[\sqrt{\phi} G^{-1}(0.5) \right] = \lambda,$$

where in the last equality we use that $G^{-1}(0.5) = 0$; see Item (a) below Item (2) in Section 1. Then, by Theorem 3.3, the proof of Items (a) and (b) follows. \square

Let $X \sim \text{LS}_d(\boldsymbol{\theta}, g)$. For given $p \in (0, 1)$, let $Q_{d;p}$ be a p -quantile for X . Let us define

$$\text{Dispersion: } \zeta = Q_{d;0.75} - Q_{d;0.25}, \quad 0 < \zeta < \infty;$$

$$\text{Relative dispersion: } \varpi = \frac{\zeta}{\zeta + 2Q_{d;0.25}}, \quad 0 < \varpi < 1;$$

$$\text{Skewness: } \varkappa(p) = \frac{Q_{d;p} + Q_{d;1-p} - 2Q_{d;0.5}}{Q_{d;1-p} + Q_{d;p}}, \quad 0 < \varkappa(p) < 1, \quad 0 < p < 0.5;$$

$$\text{Kurtosis: } \varsigma = \frac{Q_{d;7/8} - Q_{d;5/8} + Q_{d;3/8} - Q_{d;1/8}}{Q_{d;6/8} - Q_{d;2/8}}, \quad 0 \leq \varsigma < \infty.$$

The relative dispersion, skewness and kurtosis have appeared in Zwillinger and Kokoska (2000), Hinkley (1975) and Moors (1988), respectively.

Proposition 3.5. *Given $p \in (0, 1)$, let $X \sim \text{LS}_d(\boldsymbol{\theta}, g)$ and let Q_p be the p -quantile of the corresponding continuous log-symmetric distribution. If Q_p is a natural number, then*

- (a) $\zeta = 2\lambda \sinh [\sqrt{\phi} G^{-1}(0.75)]$;
- (b) $\varpi = \left\{ \coth [\sqrt{\phi} G^{-1}(0.75)] - \operatorname{cosech} [\sqrt{\phi} G^{-1}(0.75)] \right\}^{-1}$;
- (c) $\varkappa(p) = \lambda$;
- (d) $\varsigma = \frac{\sinh [\sqrt{\phi} G^{-1}(7/8)] - \sinh [\sqrt{\phi} G^{-1}(5/8)]}{\sinh [\sqrt{\phi} G^{-1}(6/8)]}$;

where $G(\cdot)$ was defined in (2).

Proof. Since Q_p is a natural number, by Theorem 3.3, $Q_{d;p} = Q_p - 1$ is a p -quantile for X , where $Q_p = \lambda \exp [\sqrt{\phi} G^{-1}(p)]$. By using the identity $G^{-1}(1-p) = -G^{-1}(p)$ (see Item (d) below Item (2) in Section 1), from Proposition 3.4 and a simple algebraic computation, the proof of statements in Items (a)-(d) follows. \square

3.3. Shape properties. The next result shows that the shape of a discrete log-symmetric distribution depends on choice of density generating kernel and on the distance between the modes of the corresponding continuous log-symmetric distribution.

Theorem 3.6. *Let g be a density generating kernel so that the corresponding continuous log-symmetric distribution, of a random variable Y , is bimodal. Then the discrete log-symmetric distribution of $X = \lfloor Y \rfloor$ has the following shapes:*

- (a) *It is bimodal, whenever the distance between the modes is big enough;*
- (b) *It is unimodal, whenever the distance between the modes is small enough.*

Proof. Since the proof of Item (b) follows the same analysis and steps as the first item, we are concerned with proving only Item (a).

Let $f_Y(y|\boldsymbol{\theta})$, $t > 0$, be the bimodal PDF of the continuous random variable $Y \sim \text{LS}(\boldsymbol{\theta}, g)$, where the distance between their modes, denoted by $y_0 > 0$ and $y_\epsilon = y_0 + \epsilon$, is big enough ($\epsilon > 6$). From bimodality property, there is $y_* \in (y_0, y_\epsilon)$ such that the following inequalities hold:

$$(6) \quad f_Y(y|\boldsymbol{\theta}) \geq f_Y(y-1|\boldsymbol{\theta}) \quad \text{for all } y \leq y_0 \text{ and } y_* \leq y \leq y_\epsilon;$$

$$(7) \quad f_Y(y|\boldsymbol{\theta}) \geq f_Y(y+1|\boldsymbol{\theta}) \quad \text{for all } y_* \geq y \geq y_0 \text{ and } y \geq y_\epsilon.$$

If x is a natural number such that $x \leq \lfloor y_0 \rfloor - 1$ and $\lfloor y_* \rfloor + 1 \leq x \leq \lfloor y_\epsilon \rfloor - 1$, from above inequalities, we have

$$p(x|\boldsymbol{\theta}) = \int_x^{x+1} f_Y(y|\boldsymbol{\theta}) dy \stackrel{(6)}{\geq} \int_x^{x+1} f_Y(y-1|\boldsymbol{\theta}) dy = p(x-1|\boldsymbol{\theta}).$$

Already, if x is a natural number such that $\lfloor y_* \rfloor - 1 \geq x \geq \lfloor y_0 \rfloor + 1$ and $x \geq \lfloor y_\epsilon \rfloor + 1$, we have

$$p(x|\boldsymbol{\theta}) = \int_x^{x+1} f_Y(y|\boldsymbol{\theta}) dy \stackrel{(7)}{\geq} \int_x^{x+1} f_Y(y+1|\boldsymbol{\theta}) dy = p(x+1|\boldsymbol{\theta}).$$

In other words, we have the following

$$(8) \quad \begin{aligned} p(0|\boldsymbol{\theta}) &\leq p(1|\boldsymbol{\theta}) \leq \dots \leq p(\lfloor y_0 \rfloor - 1|\boldsymbol{\theta}); \\ p(\lfloor y_0 \rfloor + 1|\boldsymbol{\theta}) &\geq \dots \geq p(\lfloor y_* \rfloor - 2|\boldsymbol{\theta}) \geq p(\lfloor y_* \rfloor - 1|\boldsymbol{\theta}); \end{aligned}$$

$$(9) \quad \begin{aligned} p(\lfloor y_* \rfloor + 1|\boldsymbol{\theta}) &\leq p(\lfloor y_* \rfloor + 2|\boldsymbol{\theta}) \leq \dots \leq p(\lfloor y_\epsilon \rfloor - 1|\boldsymbol{\theta}); \\ p(\lfloor y_\epsilon \rfloor + 1|\boldsymbol{\theta}) &\geq p(\lfloor y_\epsilon \rfloor + 2|\boldsymbol{\theta}) \geq \dots. \end{aligned}$$

From monotonicities (8) and (9), one can guarantee the bimodality property of the discrete log-symmetric distribution. By using (8), we show how to obtain only one of the modes, since the other one can be obtained following a similar path. Indeed, by (8) it remains to relate the probabilities $p(\lfloor y_0 \rfloor - 1|\boldsymbol{\theta})$, $p(\lfloor y_0 \rfloor|\boldsymbol{\theta})$ and $p(\lfloor y_0 \rfloor + 1|\boldsymbol{\theta})$ to find the first mode, denoted by x_0 , of the discrete log-symmetric distribution. A simple observation shows that x_0 is given by

$$x_0 = \begin{cases} \lfloor y_0 \rfloor + 1, & \text{if } p(\lfloor y_0 \rfloor - 1|\boldsymbol{\theta}) \leq p(\lfloor y_0 \rfloor|\boldsymbol{\theta}) < p(\lfloor y_0 \rfloor + 1|\boldsymbol{\theta}), \\ \lfloor y_0 \rfloor, & \text{if } p(\lfloor y_0 \rfloor - 1|\boldsymbol{\theta}) < p(\lfloor y_0 \rfloor|\boldsymbol{\theta}) > p(\lfloor y_0 \rfloor + 1|\boldsymbol{\theta}), \\ \lfloor y_0 \rfloor - 1, & \text{if } p(\lfloor y_0 \rfloor - 1|\boldsymbol{\theta}) > p(\lfloor y_0 \rfloor|\boldsymbol{\theta}) \geq p(\lfloor y_0 \rfloor + 1|\boldsymbol{\theta}), \\ \lfloor y_0 \rfloor - 1 \text{ and } \lfloor y_0 \rfloor, & \text{if } p(\lfloor y_0 \rfloor - 1|\boldsymbol{\theta}) = p(\lfloor y_0 \rfloor|\boldsymbol{\theta}) > p(\lfloor y_0 \rfloor + 1|\boldsymbol{\theta}), \\ \lfloor y_0 \rfloor \text{ and } \lfloor y_0 \rfloor + 1, & \text{if } p(\lfloor y_0 \rfloor - 1|\boldsymbol{\theta}) < p(\lfloor y_0 \rfloor|\boldsymbol{\theta}) = p(\lfloor y_0 \rfloor + 1|\boldsymbol{\theta}), \\ \lfloor y_0 \rfloor, & \text{if } p(\lfloor y_0 \rfloor - 1|\boldsymbol{\theta}) = p(\lfloor y_0 \rfloor|\boldsymbol{\theta}) < p(\lfloor y_0 \rfloor + 1|\boldsymbol{\theta}). \end{cases}$$

Notice that, any other possible relation between $p(\lfloor y_0 \rfloor - 1|\boldsymbol{\theta})$, $p(\lfloor y_0 \rfloor|\boldsymbol{\theta})$ and $p(\lfloor y_0 \rfloor + 1|\boldsymbol{\theta})$ contradicts the fact that y_0 is a mode of the continuous log-symmetric distribution. As mentioned above, using (9), the second mode of the discrete log-symmetric distribution is obtained in an analogous way. So we completed the proof. \square

As an immediate consequence of Theorem 3.6, the following result follows.

Corollary 3.7. *Let g be a density generating kernel so that the corresponding continuous log-symmetric distribution, of a random variable Y , is unimodal. Then the discrete log-symmetric distribution of $X = \lfloor Y \rfloor$ is also unimodal.*

4. MAXIMUM LIKELIHOOD ESTIMATION

4.1. **Uncensored data.** Let (X_1, \dots, X_n) be a random sample of size n from a random variable X with PMF given by (3) and $\mathbf{x} = (x_1, \dots, x_n)$ their observations (data). Then, the log-likelihood function for a parameter vector $\boldsymbol{\theta} = (\lambda, \phi)$ is given by

$$(10) \quad \ell(\boldsymbol{\theta}) = \ell(\boldsymbol{\theta}|\mathbf{x}) = \sum_{i=1}^n \log p(x_i|\boldsymbol{\theta}) = \sum_{i=1}^n \log \{G[a_{\boldsymbol{\theta}}(x_i + 1)] - G[a_{\boldsymbol{\theta}}(x_i)]\}.$$

The roots of the system formed by the partial derivatives of the log-likelihood function $\ell(\boldsymbol{\theta})$ with respect to λ and ϕ are the estimates of these parameters, respectively. Thus, we must solve the following system of equations:

$$\frac{\partial \ell(\boldsymbol{\theta})}{\partial \theta} = (Z_g)^{-1} \sum_{i=1}^n \sum_{j=0}^1 (-1)^{j+1} \frac{\partial a_{\boldsymbol{\theta}}(x_i + j)}{\partial \theta} \frac{g[a_{\boldsymbol{\theta}}^2(x_i + j)]}{p(x_i|\boldsymbol{\theta})} = 0, \quad \theta \in \{\lambda, \phi\},$$

where $Z_g = \int_{-\infty}^{\infty} g(w^2)dw$ and,

$$(11) \quad \frac{\partial a_{\boldsymbol{\theta}}(x_i)}{\partial \lambda} = -(\lambda \phi^{1/2})^{-1}, \quad \frac{\partial a_{\boldsymbol{\theta}}(x_i)}{\partial \phi} = \log \left(\frac{x_i}{\lambda} \right)^{1/(2\phi^{3/2})}.$$

Note that they must be solved by an iterative procedure for non-linear optimization, such as the Broyden-Fletcher-Goldfarb-Shanno (BFGS) quasi-Newton method; see Mittelhammer et al. (2000, p. 199).

Inference for $\boldsymbol{\theta}$ of the LS_d model can be based on the asymptotic distribution of the maximum likelihood estimator $\hat{\boldsymbol{\theta}}$. Under classic regularity conditions, this estimator is bivariate normal distributed with mean $\boldsymbol{\theta}$ and covariance matrix $\boldsymbol{\Sigma}_{\hat{\boldsymbol{\theta}}}$, namely,

$$\sqrt{n}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}) \xrightarrow{D} N_2(\mathbf{0}, \boldsymbol{\Sigma}_{\hat{\boldsymbol{\theta}}} = [\mathcal{I}(\boldsymbol{\theta})]^{-1}),$$

as $n \rightarrow \infty$, where \xrightarrow{D} means ‘‘convergence in distribution’’, $\mathcal{I}(\boldsymbol{\theta})$ is the expected Fisher information matrix, and $\mathcal{J}(\boldsymbol{\theta}) = \lim_{n \rightarrow \infty} (1/n)[\mathcal{I}(\boldsymbol{\theta})]$. Observe that $[\hat{\mathcal{I}}(\boldsymbol{\theta})]^{-1}$ is a consistent estimator of the asymptotic covariance matrix of $\hat{\boldsymbol{\theta}}$. Observe also that one may use the Hessian matrix to obtain the observed version of the expected Fisher information matrix. The Hessian matrix of $\ell(\boldsymbol{\theta})$ is given by

$$[\ddot{\ell}_{\theta\theta'}(\boldsymbol{\theta})]_{2 \times 2} = \begin{bmatrix} \frac{\partial^2 \ell(\boldsymbol{\theta})}{\partial \lambda^2} & \frac{\partial^2 \ell(\boldsymbol{\theta})}{\partial \lambda \partial \phi} \\ \frac{\partial^2 \ell(\boldsymbol{\theta})}{\partial \phi \partial \lambda} & \frac{\partial^2 \ell(\boldsymbol{\theta})}{\partial \phi^2} \end{bmatrix},$$

where its elements, for each $\theta, \theta' \in \{\lambda, \phi\}$, are

$$\ddot{\ell}_{\theta\theta'}(\boldsymbol{\theta}) = (Z_g)^{-1} \sum_{i=1}^n \sum_{j=0}^1 (-1)^{j+1} \left[\frac{\partial^2 a_{\boldsymbol{\theta}}(x_i + j)}{\partial \theta \partial \theta'} + \Theta_j(x_i) - \Omega_j(x_i) \right] \frac{g[a^2(x_i + j)]}{p(x_i|\boldsymbol{\theta})}.$$

Here we adopt the following notation:

$$(12) \quad \Theta_j(x_i) = 2a_{\boldsymbol{\theta}}(x_i + j) g'[a^2(x_i + j)] \frac{\partial a_{\boldsymbol{\theta}}(x_i + j)}{\partial \theta} \frac{\partial a_{\boldsymbol{\theta}}(x_i + j)}{\partial \theta'};$$

$$(13) \quad \Omega_j(x_i) = (Z_g)^{-1} \frac{\partial a_{\boldsymbol{\theta}}(x_i + j)}{\partial \theta} \sum_{k=0}^1 (-1)^{k+1} \frac{\partial a_{\boldsymbol{\theta}}(x_i + k)}{\partial \theta'} \frac{g[a^2(x_i + k)]}{p(x_i|\boldsymbol{\theta})};$$

whenever the density generating kernel g be differentiable. The above second-order partial derivatives of $a_{\boldsymbol{\theta}}(\cdot)$, with respect to the parameters, are given by

$$(14) \quad \begin{aligned} \frac{\partial^2 a_{\boldsymbol{\theta}}(x_i)}{\partial \lambda^2} &= (\lambda^2 \phi^{1/2})^{-1}; & \frac{\partial^2 a_{\boldsymbol{\theta}}(x_i)}{\partial \phi^2} &= \log \left(\frac{x_i}{\lambda} \right)^{-3/(4\phi^{5/2})}; \\ \frac{\partial a_{\boldsymbol{\theta}}(x_i)}{\partial \lambda \partial \phi} &= (2\lambda \phi^{3/2})^{-1}; & \frac{\partial a_{\boldsymbol{\theta}}(x_i)}{\partial \phi \partial \lambda} &= (2\lambda \phi^{3/2})^{-1}. \end{aligned}$$

Under certain regularity conditions, the Fisher information matrix

$$[\mathcal{I}_{\theta\theta'}(\boldsymbol{\theta})]_{2 \times 2} = - \begin{bmatrix} \mathbb{E} \frac{\partial^2 p(X|\boldsymbol{\theta})}{\partial \lambda^2} & \mathbb{E} \frac{\partial^2 p(X|\boldsymbol{\theta})}{\partial \lambda \partial \phi} \\ \mathbb{E} \frac{\partial^2 p(X|\boldsymbol{\theta})}{\partial \phi \partial \lambda} & \mathbb{E} \frac{\partial^2 p(X|\boldsymbol{\theta})}{\partial \phi^2} \end{bmatrix}, \quad X \sim \text{LS}_d(\boldsymbol{\theta}, g),$$

has elements of the following form

$$\mathcal{I}_{\theta\theta'}(\boldsymbol{\theta}) = (Z_g)^{-1} \sum_{x=0}^{\infty} \sum_{j=0}^1 (-1)^j \left[\frac{\partial^2 a_{\boldsymbol{\theta}}(x+j)}{\partial \theta \partial \theta'} + \Theta_j(x) - \Omega_j(x) \right] g[a^2(x+j)],$$

for each $\theta, \theta' \in \{\lambda, \phi\}$, where $\Theta_j(\cdot)$ and $\Omega_j(\cdot)$ are given in (12) and (13), respectively, and whenever the above series converges absolutely.

The extra parameter ξ (or parameter vector $\boldsymbol{\xi}$) associated with g is selected by using the profile log-likelihood function. For instance, in the case of the discrete log-Student- t distribution, two steps are require:

- i) Let $\xi_k = k$ and for each $k = 1, \dots, 100$ compute the k -th maximum likelihood estimate of $\boldsymbol{\theta}_k = (\lambda_k, \phi_k)^\top$, $\hat{\boldsymbol{\theta}}_k = (\hat{\lambda}_k, \hat{\phi}_k)^\top$ say. Compute also the k -th log-likelihood function value $\ell_k(\hat{\boldsymbol{\theta}}_k)$;
- ii) The final estimate of ξ , $\hat{\xi} = \xi_k$ say, is the one which maximizes the log-likelihood function, that is, $\hat{\xi} \in \{\text{argmax}_{\xi_k} \ell_k(\hat{\boldsymbol{\theta}}_k)\}$, and the estimate of $\boldsymbol{\theta}$ is $\hat{\boldsymbol{\theta}}_k = (\hat{\lambda}_k, \hat{\phi}_k)^\top$.

4.2. Censored data. Let $Y_i \sim \text{LS}(\boldsymbol{\theta}, g)$ be the failure time of the i -th individual and let δ_i indicate whether the i -th individual is censored or not. Let us define $d_k =$ “number of failures at time t_k ”, $q_k =$ “number censored at time t_k ” and $N_k = \sum_{i=k}^{\infty} (d_i + q_i)$. Note that $N_k - d_k$ represents the number survived just before time $t_k + 1$. That is, in each given time t_k , there are d_k failures and $N_k - d_k$ survivals.

Since the data are discrete observing $\{(Y_i, \delta_i)\}$ is equivalent to observing $\{(d_k, q_k)\}$, the likelihood function for the random censoring is given by

$$L^R(\boldsymbol{\theta}) = \prod_{i=1}^n [f_Y(y_i|\boldsymbol{\theta})]^{\delta_i} [1 - F_Y(y_i|\boldsymbol{\theta})]^{1-\delta_i} = \prod_{k=1}^{\infty} [p(x_k|\boldsymbol{\theta})]^{d_k} [p(x_k|\boldsymbol{\theta}) + R(x_k|\boldsymbol{\theta})]^{q_k}.$$

This type of censoring has as special case type I and II censoring. The corresponding log-likelihood is

$$(15) \quad \begin{aligned} \ell^R(\boldsymbol{\theta}) &= \sum_{k=1}^{\infty} \left\{ d_k \log p(x_k|\boldsymbol{\theta}) + q_k \log [p(x_k|\boldsymbol{\theta}) + R(x_k|\boldsymbol{\theta})] \right\} \\ &= \sum_{k=1}^{\infty} d_k \log \{ G[a_{\boldsymbol{\theta}}(x_k + 1)] - G[a_{\boldsymbol{\theta}}(x_k)] \} + \sum_{k=1}^{\infty} q_k \log \{ 1 - G[a_{\boldsymbol{\theta}}(x_k)] \}. \end{aligned}$$

Remark 4.1. *By Proposition 3.1, the log-likelihood (15) can be rewritten in terms of HR as*

$$\ell^R(\boldsymbol{\theta}) = \sum_{k=1}^{\infty} (d_k + q_k) \left\{ \frac{d_k}{d_k + q_k} \log [H(x_k|\boldsymbol{\theta})] - \log [1 - H(x_k|\boldsymbol{\theta})] + \sum_{y=0}^{x_k-1} \log [1 - H(y|\boldsymbol{\theta})] \right\},$$

whenever the above series converges absolutely.

Differentiating in (15), a straightforward computation shows that

$$\begin{aligned} \frac{\partial \ell^R(\boldsymbol{\theta})}{\partial \theta} &= (Z_g)^{-1} \sum_{k=1}^{\infty} d_k \sum_{j=0}^1 (-1)^{j+1} \frac{\partial a_{\boldsymbol{\theta}}(x_k + j)}{\partial \theta} \frac{g[a_{\boldsymbol{\theta}}^2(x_k + j)]}{p(x_k|\boldsymbol{\theta})} \\ &\quad - (Z_g)^{-1} \sum_{k=1}^{\infty} q_k \frac{\partial a_{\boldsymbol{\theta}}(x_k)}{\partial \theta} \frac{g[a_{\boldsymbol{\theta}}^2(x_k)]}{p(x_k|\boldsymbol{\theta}) + R(x_k|\boldsymbol{\theta})}, \quad \theta \in \{\lambda, \phi\}, \end{aligned}$$

where $Z_g = \int_{-\infty}^{\infty} g(w^2) dw$. The mixed partial derivatives of $\ell^R(\boldsymbol{\theta})$ are given by

$$\begin{aligned} \frac{\partial^2 \ell^R(\boldsymbol{\theta})}{\partial \theta \partial \theta'} &= (Z_g)^{-1} \sum_{k=1}^{\infty} d_k \sum_{j=0}^1 (-1)^{j+1} \left[\frac{\partial^2 a_{\boldsymbol{\theta}}(x_k + j)}{\partial \theta \partial \theta'} + \Theta_j(x_k) - \Omega_j(x_k) \right] \frac{g[a_{\boldsymbol{\theta}}^2(x_k + j)]}{p(x_k|\boldsymbol{\theta})} \\ &\quad - (Z_g)^{-1} \sum_{k=1}^{\infty} q_k \left[\frac{\partial^2 a_{\boldsymbol{\theta}}(x_k)}{\partial \theta \partial \theta'} + \Theta_0(x_k) + \hat{\Omega}(x_k) \right] \frac{g[a_{\boldsymbol{\theta}}^2(x_k)]}{p(x_k|\boldsymbol{\theta}) + R(x_k|\boldsymbol{\theta})}, \quad \theta, \theta' \in \{\lambda, \phi\}, \end{aligned}$$

where

$$\hat{\Omega}(x_k) = (Z_g)^{-1} \frac{g[a^2(x_k)]}{p(x_k|\boldsymbol{\theta}) + R(x_k|\boldsymbol{\theta})} \frac{\partial a_{\boldsymbol{\theta}}(x_k)}{\partial \theta} \frac{\partial a_{\boldsymbol{\theta}}(x_k)}{\partial \theta'},$$

and $\Theta_j(\cdot), \Omega_j(\cdot)$ are as in (12) and (13), respectively. The first and second derivatives of function $a_{\boldsymbol{\theta}}(\cdot)$ are given in (11) and (14), respectively.

5. MONTE CARLO SIMULATION STUDY

A Monte Carlo simulation study was carried out to evaluate the performance of the maximum likelihood estimators for the LS_d models, particularly the log-normal, log-Student-*t*, log-contaminated-normal, log-power-exponential, extended Birnbaum-Saunders and extended Birnbaum-Saunders-*t* cases. Note that when $\phi = 4$ (fixed) the Birnbaum-Saunders and Birnbaum-Saunders-*t* are obtained. All numerical evaluations were done in the R software; see R Core Team (2016).

The simulation scenario considers: sample size $n \in \{40, 120, 400\}$, values of true parameters $\phi \in \{1, 4, 8\}$, $\lambda \in \{2.00\}$, censoring proportions $\{0\%, 10\%, 30\%\}$, and 1,000 Monte Carlo replications for each sample size. The values of the true extra parameters are presented in the caption of each table.

The maximum likelihood estimation results are presented in Tables 1–6. The following sample statistics for the maximum likelihood estimates are reported: empirical mean, bias, and mean squared error (MSE). A look at the results in Tables 1–6 allows us to conclude that, as the sample size increases, the bias and MSE of all the estimators decrease, indicating that they are asymptotically unbiased, as expected. Moreover, as the censoring proportion increases, the performances of the estimators of ϕ and λ , deteriorate. Generally, all of these results show the good performance of the proposed model.

TABLE 1. Empirical values of mean, bias and MSE from simulated discrete log-normal data for the indicated maximum likelihood estimators.

n	Cen.		$\phi = 1$			$\phi = 4$			$\phi = 8$		
			Mean	Bias	MSE	Mean	Bias	MSE	Mean	Bias	MSE
40		$\hat{\phi}$	1.0119	0.0119	0.0851	4.0752	0.0752	1.7247	8.1388	0.1388	7.5728
		$\hat{\lambda}$	2.0262	0.0262	0.1166	2.1126	0.1126	0.5629	2.2586	0.2586	1.4043
120	0%	$\hat{\phi}$	1.0059	0.0059	0.0260	4.0413	0.0413	0.5237	8.0768	0.0768	2.1683
		$\hat{\lambda}$	2.0086	0.0086	0.0394	2.0317	0.0317	0.1759	2.0750	0.0750	0.3897
400		$\hat{\phi}$	1.0041	0.0041	0.0077	4.0184	0.0184	0.1522	8.0404	0.0404	0.6641
		$\hat{\lambda}$	2.0021	0.0021	0.0110	2.0086	0.0086	0.0464	2.0198	0.0198	0.1020
40		$\hat{\phi}$	1.0189	0.0189	0.0998	4.1507	0.1507	2.2755	8.3002	0.3002	10.0324
		$\hat{\lambda}$	2.0301	0.0301	0.1198	2.1154	0.1154	0.5850	2.2528	0.2528	1.4323
120	10%	$\hat{\phi}$	1.0114	0.0114	0.0290	4.0519	0.0519	0.5978	8.1162	0.1162	2.5822
		$\hat{\lambda}$	2.0099	0.0099	0.0396	2.0416	0.0416	0.1795	2.0842	0.0842	0.4037
400		$\hat{\phi}$	1.0015	0.0015	0.0090	4.0262	0.0262	0.1817	8.0675	0.0675	0.8152
		$\hat{\lambda}$	2.0012	0.0012	0.0109	2.0014	0.0014	0.0466	2.0073	0.0073	0.0984
40		$\hat{\phi}$	1.0562	0.0562	0.1839	4.4113	0.4113	5.7506	8.9965	0.9965	28.9825
		$\hat{\lambda}$	2.0435	0.0435	0.1407	2.1396	0.1396	0.6858	2.2925	0.2925	1.7228
120	30%	$\hat{\phi}$	1.0218	0.0218	0.0458	4.1116	0.1116	1.1065	8.2539	0.2539	5.0762
		$\hat{\lambda}$	2.0124	0.0124	0.0438	2.0450	0.0450	0.1915	2.0868	0.0868	0.4270
400		$\hat{\phi}$	1.0093	0.0093	0.0133	4.0840	0.0840	0.3158	8.1961	0.1961	1.5102
		$\hat{\lambda}$	2.0046	0.0046	0.0119	2.0076	0.0076	0.0501	2.0131	0.0131	0.1043

TABLE 2. Empirical values of mean, bias and MSE from simulated discrete log-Student- t data for the indicated maximum likelihood estimators with $\xi = 4$.

n	Cen.		$\phi = 1$			$\phi = 4$			$\phi = 8$		
			Mean	Bias	MSE	Mean	Bias	MSE	Mean	Bias	MSE
40		$\hat{\phi}$	1.0250	0.0250	0.1313	4.1432	0.1432	2.6039	8.3035	0.3035	14.7992
		$\hat{\lambda}$	2.0351	0.0351	0.1589	2.1416	0.1416	0.7463	2.3098	0.3098	1.9332
120	0%	$\hat{\phi}$	1.0093	0.0093	0.0408	4.0316	0.0316	0.8793	7.9743	-0.0257	4.0502
		$\hat{\lambda}$	2.0179	0.0179	0.0497	2.0657	0.0657	0.2195	2.1694	0.1694	0.7087
400		$\hat{\phi}$	0.9974	-0.0026	0.0123	3.9778	-0.0222	0.4006	7.9731	-0.0269	2.8371
		$\hat{\lambda}$	1.9986	-0.0014	0.0137	2.0148	0.0148	0.0684	2.0657	0.0657	0.2455
40		$\hat{\phi}$	1.0332	0.0332	0.1499	4.2117	0.2117	3.2204	8.5131	0.5131	14.0700
		$\hat{\lambda}$	2.0318	0.0318	0.1692	2.1338	0.1338	0.8105	2.2965	0.2965	2.1397
120	10%	$\hat{\phi}$	1.0149	0.0149	0.0466	4.0564	0.0564	0.9162	8.1134	0.1134	4.0340
		$\hat{\lambda}$	2.0068	0.0068	0.0510	2.0434	0.0434	0.2086	2.0916	0.0916	0.4621
400		$\hat{\phi}$	1.0026	0.0026	0.0126	4.0285	0.0285	0.2729	8.0335	0.0335	1.1661
		$\hat{\lambda}$	1.9987	-0.0013	0.0146	2.0000	0.0000	0.0598	2.0137	0.0137	0.1208
40		$\hat{\phi}$	1.0772	0.0772	0.2681	4.5456	0.5456	8.5155	9.3122	1.3122	41.0324
		$\hat{\lambda}$	2.0430	0.0430	0.2007	2.1605	0.1605	1.0059	2.3570	0.3570	3.1707
120	30%	$\hat{\phi}$	1.0148	0.0148	0.0666	4.0801	0.0801	1.4871	8.1426	0.1426	6.8379
		$\hat{\lambda}$	2.0032	0.0032	0.0534	2.0429	0.0429	0.2210	2.0866	0.0866	0.4900
400		$\hat{\phi}$	1.0129	0.0129	0.0168	4.0890	0.0890	0.4097	8.1679	0.1679	1.8873
		$\hat{\lambda}$	2.0021	0.0021	0.0160	2.0040	0.0040	0.0645	2.0179	0.0179	0.1297

TABLE 3. Empirical values of mean, bias and MSE from simulated discrete log-contaminated-normal data for the indicated maximum likelihood estimators with $\xi = (0.5, 0.5)^\top$.

n	Cen.		$\phi = 1$			$\phi = 4$			$\phi = 8$		
			Mean	Bias	MSE	Mean	Bias	MSE	Mean	Bias	MSE
40		$\hat{\phi}$	1.0057	0.0057	0.0947	4.0459	0.0459	1.8954	8.0944	0.0944	7.9832
		$\hat{\lambda}$	2.0437	0.0437	0.1706	2.1750	0.1750	0.8474	2.3687	0.3687	2.1616
120	0%	$\hat{\phi}$	1.0024	0.0024	0.0300	4.0207	0.0207	0.6042	8.0165	0.0165	2.5295
		$\hat{\lambda}$	2.0161	0.0161	0.0545	2.0589	0.0589	0.2503	2.1297	0.1297	0.5661
400		$\hat{\phi}$	1.0020	0.0020	0.0085	4.0061	0.0061	0.1779	7.9912	-0.0088	0.8225
		$\hat{\lambda}$	2.0024	0.0024	0.0165	2.0161	0.0161	0.0745	2.0410	0.0410	0.1751
40		$\hat{\phi}$	1.0336	0.0336	0.1484	4.1971	0.1971	3.1525	8.4153	0.4153	12.8221
		$\hat{\lambda}$	2.0562	0.0562	0.1819	2.1952	0.1952	0.9257	2.3944	0.3944	2.3858
120	10%	$\hat{\phi}$	1.0154	0.0154	0.0396	4.0676	0.0676	0.7806	8.1552	0.1552	3.3793
		$\hat{\lambda}$	2.0182	0.0182	0.0588	2.0688	0.0688	0.2649	2.1338	0.1338	0.5975
400		$\hat{\phi}$	1.0067	0.0067	0.0103	4.0278	0.0278	0.2113	8.0581	0.0581	0.9135
		$\hat{\lambda}$	1.9988	-0.0012	0.0163	2.0077	0.0077	0.0735	2.0222	0.0222	0.1586
40		$\hat{\phi}$	1.0644	0.0644	0.2448	4.4774	0.4774	7.4371	9.1280	1.1280	33.9336
		$\hat{\lambda}$	2.0635	0.0635	0.2088	2.2106	0.2106	1.0845	2.4422	0.4422	3.1094
120	30%	$\hat{\phi}$	1.0310	0.0310	0.0591	4.1681	0.1681	1.3522	8.4147	0.4147	6.3718
		$\hat{\lambda}$	2.0237	0.0237	0.0635	2.0767	0.0767	0.2895	2.1436	0.1436	0.6536
400		$\hat{\phi}$	1.0126	0.0126	0.0161	4.0609	0.0609	0.3648	8.1301	0.1301	1.6252
		$\hat{\lambda}$	2.0005	0.0005	0.0176	2.0102	0.0102	0.0771	2.0237	0.0237	0.1628

TABLE 4. Empirical values of mean, bias and MSE from simulated discrete log-power-exponential data for the indicated maximum likelihood estimators with $\xi = -0.5$.

n	Cen.		$\phi = 1$			$\phi = 4$			$\phi = 8$		
			Mean	Bias	MSE	Mean	Bias	MSE	Mean	Bias	MSE
40		$\hat{\phi}$	0.9783	-0.0217	0.0470	3.9497	-0.0503	1.1520	7.9186	-0.0814	5.6648
		$\hat{\lambda}$	2.0072	0.0072	0.0481	2.0493	0.0493	0.2856	2.1332	0.1332	0.6806
120	0%	$\hat{\phi}$	0.9970	-0.0030	0.0134	4.0014	0.0014	0.3399	7.9966	-0.0034	1.5695
		$\hat{\lambda}$	2.0032	0.0032	0.0153	2.0185	0.0185	0.0863	2.0482	0.0482	0.1996
400		$\hat{\phi}$	0.9969	-0.0031	0.0042	3.9906	-0.0094	0.1019	7.9864	-0.0136	0.4742
		$\hat{\lambda}$	1.9998	-0.0002	0.0047	2.0036	0.0036	0.0282	2.0098	0.0098	0.0640
40		$\hat{\phi}$	0.9941	-0.0059	0.0612	4.0123	0.0123	1.5967	8.0169	0.0169	8.0172
		$\hat{\lambda}$	2.0046	0.0046	0.0510	2.0472	0.0472	0.2900	2.1398	0.1398	0.6989
120	10%	$\hat{\phi}$	0.9999	-0.0001	0.0167	4.0151	0.0151	0.4585	8.0260	0.0260	2.0633
		$\hat{\lambda}$	2.0010	0.0010	0.0165	2.0099	0.0099	0.0866	2.0349	0.0349	0.1968
400		$\hat{\phi}$	1.0020	0.0020	0.0053	4.0178	0.0178	0.1283	8.0441	0.0441	0.6083
		$\hat{\lambda}$	2.0010	0.0010	0.0050	2.0044	0.0044	0.0298	2.0084	0.0084	0.0661
40		$\hat{\phi}$	1.0093	0.0093	0.0890	4.1602	0.1602	3.3402	8.2982	0.2982	15.8037
		$\hat{\lambda}$	2.0111	0.0111	0.0646	2.0512	0.0512	0.3185	2.1408	0.1408	0.7509
120	30%	$\hat{\phi}$	1.0040	0.0040	0.0250	4.0485	0.0485	0.7736	8.1109	0.1109	3.7337
		$\hat{\lambda}$	2.0019	0.0019	0.0191	2.0104	0.0104	0.0927	2.0350	0.0350	0.2090
400		$\hat{\phi}$	1.0026	0.0026	0.0079	4.0215	0.0215	0.1934	8.0665	0.0665	0.9711
		$\hat{\lambda}$	2.0010	0.0010	0.0056	2.0045	0.0045	0.0318	2.0093	0.0093	0.0694

TABLE 5. Empirical values of mean, bias and MSE from simulated discrete extended Birnbaum-Saunders data for the indicated maximum likelihood estimators with $\zeta = 0.5$.

n	Cen.		$\phi = 1$			$\phi = 4$			$\phi = 8$		
			Mean	Bias	MSE	Mean	Bias	MSE	Mean	Bias	MSE
40		$\hat{\phi}$	0.8741	-0.1259	0.1721	3.9719	-0.0281	0.9531	7.9871	-0.0129	4.3427
		$\hat{\lambda}$	2.0126	0.0126	0.0075	2.0117	0.0117	0.0275	2.0152	0.0152	0.0542
120	0%	$\hat{\phi}$	0.9747	-0.0253	0.0353	4.0090	0.0090	0.3178	8.0289	0.0289	1.3258
		$\hat{\lambda}$	2.0049	0.0049	0.0029	2.0032	0.0032	0.0090	2.0037	0.0037	0.0185
400		$\hat{\phi}$	0.9913	-0.0087	0.0117	4.0042	0.0042	0.0978	8.0266	0.0266	0.4048
		$\hat{\lambda}$	2.0020	0.0020	0.0009	2.0011	0.0011	0.0026	2.0004	0.0004	0.0051
40		$\hat{\phi}$	0.8825	-0.1175	0.1945	4.0107	0.0107	1.2217	8.1117	0.1117	5.3717
		$\hat{\lambda}$	2.0128	0.0128	0.0076	2.0120	0.0120	0.0289	2.0172	0.0172	0.0565
120	10%	$\hat{\phi}$	0.9699	-0.0301	0.0460	4.0048	0.0048	0.3449	8.0872	0.0872	1.6128
		$\hat{\lambda}$	2.0047	0.0047	0.0028	2.0043	0.0043	0.0096	2.0031	0.0031	0.0187
400		$\hat{\phi}$	0.9907	-0.0093	0.0134	4.0000	0.0000	0.1167	8.0092	0.0092	0.4932
		$\hat{\lambda}$	2.0005	0.0005	0.0009	1.9997	-0.0003	0.0026	2.0005	0.0005	0.0050
40		$\hat{\phi}$	0.7674	-0.2326	0.3577	4.0841	0.0841	2.0748	8.3517	0.3517	8.9625
		$\hat{\lambda}$	2.0152	0.0152	0.0067	2.0178	0.0178	0.0343	2.0280	0.0280	0.0692
120	30%	$\hat{\phi}$	0.9263	-0.0737	0.0967	4.0323	0.0323	0.5129	8.1723	0.1723	2.5286
		$\hat{\lambda}$	2.0052	0.0052	0.0028	2.0057	0.0057	0.0106	2.0055	0.0055	0.0210
400		$\hat{\phi}$	0.9852	-0.0148	0.0212	4.0188	0.0188	0.1660	8.0638	0.0638	0.7290
		$\hat{\lambda}$	2.0009	0.0009	0.0009	2.0011	0.0011	0.0028	2.0028	0.0028	0.0056

TABLE 6. Empirical values of mean, bias and MSE from simulated discrete extended Birnbaum-Saunders- t data for the indicated maximum likelihood estimators with $\xi = (0.5, 4)^\top$.

n	Cen.		$\phi = 1$			$\phi = 4$			$\phi = 8$		
			Mean	Bias	MSE	Mean	Bias	MSE	Mean	Bias	MSE
40		$\hat{\phi}$	0.9871	-0.0129	0.1511	3.9863	-0.0137	1.4404	8.0258	0.0258	6.0994
		$\hat{\lambda}$	2.0028	0.0028	0.0114	2.0100	0.0100	0.0387	2.0167	0.0167	0.0756
120	0%	$\hat{\phi}$	0.9998	-0.0002	0.0457	4.0319	0.0319	0.4972	8.0689	0.0689	2.1333
		$\hat{\lambda}$	2.0026	0.0026	0.0037	2.0061	0.0061	0.0128	2.0086	0.0086	0.0248
400		$\hat{\phi}$	0.9984	-0.0016	0.0141	3.9996	-0.0004	0.1569	8.0282	0.0282	0.6348
		$\hat{\lambda}$	2.0018	0.0018	0.0010	2.0031	0.0031	0.0038	2.0035	0.0035	0.0074
40		$\hat{\phi}$	1.0002	0.0002	0.1663	4.1093	0.1093	1.8263	8.2395	0.2395	7.9333
		$\hat{\lambda}$	2.0009	0.0009	0.0109	2.0029	0.0029	0.0387	2.0095	0.0095	0.0760
120	10%	$\hat{\phi}$	1.0055	0.0055	0.0512	4.0492	0.0492	0.5522	8.1287	0.1287	2.3725
		$\hat{\lambda}$	2.0027	0.0027	0.0037	2.0044	0.0044	0.0129	2.0080	0.0080	0.0245
400		$\hat{\phi}$	1.0003	0.0003	0.0157	3.9944	-0.0056	0.1649	8.0085	0.0085	0.6848
		$\hat{\lambda}$	2.0011	0.0011	0.0012	2.0020	0.0020	0.0042	2.0028	0.0028	0.0078
40		$\hat{\phi}$	0.9829	-0.0171	0.2728	4.1695	0.1695	2.7474	8.4158	0.4158	12.0941
		$\hat{\lambda}$	2.0027	0.0027	0.0114	2.0055	0.0055	0.0438	2.0151	0.0151	0.0882
120	30%	$\hat{\phi}$	1.0115	0.0115	0.0708	4.1024	0.1024	0.7985	8.2566	0.2566	3.5320
		$\hat{\lambda}$	2.0029	0.0029	0.0038	2.0065	0.0065	0.0140	2.0119	0.0119	0.0273
400		$\hat{\phi}$	1.0009	0.0009	0.0238	4.0094	0.0094	0.2337	8.0494	0.0494	0.9990
		$\hat{\lambda}$	2.0014	0.0014	0.0012	2.0037	0.0037	0.0047	2.0049	0.0049	0.0087

6. ILLUSTRATIVE EXAMPLES

The LS_d models are now used to analyze two real-world data sets. It is considered the following discrete LS_d models: log-normal (LN), log-Student- t (Lt), log-contaminated-normal (LCN), log-power-exponential (LPE), Birnbaum-Saunders (BS), extended Birnbaum-Saunders (EBS), Birnbaum-Saunders- t (BSt), and extended Birnbaum-Saunders- t (EBSt).

Example 6.1. *The first data set corresponds to the number of times that a DEC-20 computer broke down in each of 128 consecutive weeks of operation. This computer has operated at the Open University during the 1980s; see Table 9 and Trenkler (1995). Descriptive statistics for the computer breaks data set are the following: 128(sample size), 0(minimum), 22(maximum), 3(median), 4.016(mean), 3.808(standard deviation), 94.839(coefficent of variation), 1.732(coefficent of skewness) and 3.995(coefficent of kurtosis). From these results, we observe the positive skewness and a high degree of kurtosis. Figure 1(left) shows the histogram for the computer breaks data, from where it is confirmed the positive skewness. Moreover, Figure 1(right) shows the usual and adjusted boxplots, and we note that some potential outliers are not in fact outliers when the adjusted boxplot is observed.*

TABLE 7. Computer breaks data.

x	0	1	2	3	4	5	6	7	8	9	10	11	12	13	16	17	22
frequency	15	19	23	14	15	10	8	4	6	2	3	3	2	1	1	1	1




FIGURE 1. Histogram (left) and boxplots (right) for the computer breaks data.

Table 8 presents the maximum likelihood estimates, computed by the BFGS method, and standard errors (SEs) for the LS_d models parameters. Moreover, the p -values of the χ^2 and Cramer-Von Mises (CVM) statistics, and the Akaike (AIC) and Bayesian information (BIC) criteria, are also reported. The results of Table 8 reveal that the discrete BS model

provides the best adjustment compared to other models based on the values of AIC and BIC.

TABLE 8. Maximum likelihood estimates (with SE in parentheses) and model selection measures for fit to the computer breaks data.

Model	Estimates			p-value		AIC	BIC
	$\hat{\lambda}$ (SE)	$\hat{\phi}$ (SE)	$\hat{\xi}$ (ξ)	χ^2	CMV		
LN	3.2280 (0.2526)	0.7541 (0.1048)	—	0.7841	0.6959	643.5141	652.0702
Lt	3.2653 (0.2574)	0.7065 (0.1026)	20.0	0.8306	0.7762	644.2248	652.7809
LCN	3.2283 (0.2526)	0.6858 (0.0953)	(0.9, 0.9)	0.7996	0.6967	645.5205	656.9287
LPE	3.1624 (0.2770)	1.0176 (0.0555)	-0.2	0.7096	0.5151	642.8785	651.4346
BS	3.1704 (0.0589)	*	0.9	0.6007	0.5283	640.6061	646.3102
EBS	3.1436 (0.2090)	2.9392 (0.0438)	1.1	0.6517	0.4700	642.4045	650.9605
BSt	3.1803 (0.3008)	*	(0.9, 20.0)	0.7966	0.5799	642.9026	651.4587
EBSt	3.1406 (0.2193)	2.0820 (0.0818)	(1.3, 20.0)	0.6492	0.4364	644.5534	655.9615

* indicates that $\phi = 4$ (fixed).

Example 6.2. The second data set refers to the number of physiotherapy sessions until a patient's chronic back pain is reduced or alleviated; see Table 9. The patients were submitted to electric currents and the study was developed by the School of Physiotherapy Clinics of City University of Sao Paulo (UNICID), Sao Paulo, Brazil; see Silva et al. (2017). Observations were considered censored to the right when patients did not report pain reduction or relief after 12 treatment sessions, or if they had been lost to follow-up. Such as in Vila et al. (2019), the variable of interest is defined as $T = X - 1$, $t = 0, 1, 2, 3, \dots$, where $t = 0$ denotes a patient who presented pain relief in the first session performed. Descriptive statistics for the pain relief data are the following: 100 (sample size), 0 (minimum), 11 (maximum), 0 (median), 0.98 (mean), 1.933 (standard deviation), 197.258 (coefficient of variation), 2.802 (coefficient of skewness) and 9.015 (coefficient of kurtosis). These statistics values indicate the positive skewness and a high degree of kurtosis. Figure 2 shows the histogram and the fitted survival function by the Kaplan-Meier (KM) method.

TABLE 9. Number of sessions until a patient's chronic back pain is reduced or alleviated.

Sessions	T	# at risk	# of events	censoring indicator
1	0	100	64	0
2	1	36	16	0
3	2	20	5	1
4	3	14	4	0
5	4	10	4	0
6	5	6	3	0
7	6	3	0	0
8	7	3	1	0
9	8	2	0	0
10	9	2	1	0
11	10	1	0	0
12	11	1	0	1

The maximum likelihood estimates of the discrete log-symmetric distribution parameters, along with AIC and BIC criteria are reported in Table 10. We note that the log-Student-t




FIGURE 2. Histogram (left) and KM (right) for the pain relief data.

model provides better adjustment compared to the other models based on the values of AIC and BIC. Table 10 and 11 present the fitted survival functions obtained by the KM and the discrete log-symmetric models. These results suggest that (extended) Birnbaum-Saunders and log-normal models yield the best fits to the pain relief data.

TABLE 10. Maximum likelihood estimates and model selection measures for fit to the pain relief data.

Discrete distribution	Estimates (SE)	AIC	BIC
Log-normal	$\hat{\lambda}=2.3229 (0.1329)$ $\hat{\phi}=0.462 (0.0571)$	540.0872	549.1191
Log-Student-t	$\hat{\lambda}=1.8745 (0.1250)$ $\hat{\phi}=0.122 (0.0538)$ $\hat{\zeta}=2$	513.1153	522.1472
Log-Power-Exponential	$\hat{\lambda}=2.0046 (0.0999)$ $\hat{\phi}=0.1713 (0.0259)$ $\hat{\zeta}=0.5$	528.1035	537.1354
Log-Contaminated-Normal	$\hat{\lambda}=1.8654 (0.1329)$ $\hat{\phi}=0.1018 (0.0519)$ $\hat{\zeta}=(0.37;0.10)$	513.3146	525.3571
Birnbaum-Saunders	$\hat{\lambda}=2.4767 (0.8043)$ $\hat{\zeta}=0.7$	543.6507	549.6719
Extended Birnbaum-Saunders	$\hat{\lambda}=2.3263 (0.156)$ $\hat{\phi}=184.8547 (0.0701)$ $\hat{\zeta}=0.1$	540.2164	549.2483
Birnbaum-Saunders-t	$\hat{\lambda}=1.8966 (0.2047)$ $\hat{\zeta}=(0.4;2.0)$	515.0695	524.1014
Extended Birnbaum-Saunders-t	$\hat{\lambda}=1.8751 (0.1477)$ $\hat{\phi}=49.0515 (0.1995)$ $\hat{\zeta}=(0.1;2.0)$	515.1634	527.2060

TABLE 11. Estimates of the survival function via KM and discrete log-symmetric distributions.

x	KM	LN	$L-t$	LPE	LCN	BS	EBS	$BS-t$	$EBS-t$
0	1	0.8925	0.8931	0.8698	0.8848	0.9100	0.8930	0.8774	0.8930
1	0.4333	0.5871	0.4350	0.5018	0.4354	0.6202	0.5880	0.4533	0.4354
2	0.2400	0.3533	0.1552	0.2412	0.1610	0.3919	0.3541	0.1835	0.1557
3	0.1933	0.2120	0.0811	0.1315	0.0885	0.2447	0.2126	0.0982	0.0812
4	0.1588	0.1297	0.0534	0.0791	0.0614	0.1528	0.1301	0.0640	0.0534
5	0.1299	0.0813	0.0398	0.0511	0.0458	0.0958	0.0815	0.0466	0.0396
6	0.0866	0.0523	0.0318	0.0348	0.0352	0.0603	0.0524	0.0364	0.0316
7	0.0794	0.0344	0.0267	0.0247	0.0276	0.0381	0.0344	0.0297	0.0265
8	0.0577	0.0232	0.0231	0.0182	0.0220	0.0242	0.0231	0.0250	0.0228
10	0.0505	0.0111	0.0184	0.0106	0.0145	0.0098	0.0110	0.0190	0.0181
12	0.0361	0.0056	0.0155	0.0067	0.0101	0.0040	0.0056	0.0153	0.0152




FIGURE 3. Estimation of the survival function using the KM (solid) and discrete log-symmetric distributions (dashed) with the pain relief data.

7. CONCLUDING REMARKS

We have proposed a new class of distributions to deal with cases where the data are discrete, asymmetric and nonnegative. The proposed approach is a discrete version of the family of continuous log-symmetric distributions. We have considered estimation about the model parameters based on the maximum likelihood method with censored and uncensored data. A Monte Carlo simulation study was carried out to evaluate the behavior of the maximum likelihood estimators. We have applied the proposed models to two real-world data sets. In general, the results have shown that the proposed discrete family proved to be an useful model for discrete data. As part of future research, it is of interest to discuss regression models as well as multivariate extensions. Moreover, time series models based on the proposed class may be of interest. Work on these issues is currently in progress and we hope to report some findings in future papers.

REFERENCES

- Balakrishnan, N., Saulo, H., Bourguignon, M., and Zhu, X. (2017). On moment-type estimators for a class of log-symmetric distributions. *Computational Statistics*, 32(4):1339–1355.
- Hinkley, D. V. (1975). On power transformations to symmetry. *Biometrika*, 62:101–111.
- Jones, M. C. (2008). On reciprocal symmetry. *Journal of Statistical Planning and Inference*, 138:3039–3043.
- Medeiros, F. M. C. and Ferrari, S. L. P. (2017). Small-sample testing inference in symmetric and log-symmetric linear regression models. *Statistica Neerlandica*, 71:200–224.
- Mittelhammer, R. C., Judge, G. G., and Miller, D. J. (2000). *Econometric Foundations*. Cambridge University Press, New York, US.
- Moors, J. J. A. (1988). A quantile alternative for kurtosis. *The Statistician*, 37:25–32.
- R Core Team (2016). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Saulo, H. and Leão, J. (2017). On log-symmetric duration models applied to high frequency financial data. *Economics Bulletin*, 37:1089–1097.
- Silva, J. F., Liebano, R. E., Corrêa, J. B., Matsushita, R. Y., and Nakano, E. Y. (2017). Analysis of the time to relieving pain in patients with chronic non-specific low back pain via Cox proportional hazard model. *Ciência e Natura*, 39:233–243.
- Trenkler, D. (1995). A handbook of small data sets: Hand, d.j., daly, f., lunn, a.d., mcconway, k.j. & ostrowski, e. (1994): Chapman & hall, london. *Computational Statistics & Data Analysis*, 19(1):101–101.
- Vanegas, L. H. and Paula, G. A. (2016a). An extension of log-symmetric regression models: R codes and applications. *Journal of Statistical Simulation and Computation*, 86:1709–1735.
- Vanegas, L. H. and Paula, G. A. (2016b). *ssym: Fitting Semi-Parametric log-Symmetric Regression Models*. R package version 1.5.7.

- Ventura, M., Saulo, H., Leiva, V., and Monsueto, S. (2019). Log-symmetric regression models: information criteria and application to movie business and industry data with economic implications. *Applied Stochastic Models in Business and Industry*, 35(4):963–977.
- Vila, R., Nakano, E. Y., and Saulo, H. (2019). Theoretical results on the discrete Weibull distribution of Nakagawa and Osaki. *Statistics*, 53(2):339–363.
- Zwillinger, D. and Kokoska, S. (2000). *Standard Probability and Statistical Tables and Formula*. Chapman & Hall, Boca Raton.

DEPARTAMENTO DE ESTATÍSTICA - UNIVERSIDADE DE BRASÍLIA, 70910-900, DF, BRAZIL,
Helton Saulo , EMAIL: heltonsaulo@gmail.com
Roberto Vila, EMAIL: rovig161@gmail.com
Leonardo Paiva, EMAIL: leopaiva23@hotmail.com

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY, HAMILTON, ON, CANADA
Narayanaswamy Balakrishnan, EMAIL: bala@mcmaster.ca