

Robust Sure Independence Screening for Non-polynomial dimensional Generalized Linear Models

Abhik Ghosh¹ and Magne Thoresen²

¹ Indian Statistical Institute, Kolkata, India

² University of Oslo, Oslo, Norway

February 28, 2022

Abstract

We consider the problem of variable screening in ultra-high dimensional (of non-polynomial order) generalized linear models (GLMs). Since the popular SIS approach is extremely unstable in the presence of contamination and noises, which may frequently arise in the large scale sample data (e.g., Omics data), we discuss a new robust screening procedure based on the minimum density power divergence estimator (MDPDE) of the marginal regression coefficients. Our proposed screening procedure performs extremely well both under pure and contaminated data scenarios. We also theoretically justify the use of this marginal MDPDEs for variable screening from the population as well as sample aspects; in particular, we prove that these marginal MDPDEs are uniformly consistent leading to the sure screening property of our proposed algorithm. We have also proposed an appropriate MDPDE based extension for robust conditional screening in the GLMs along with the derivation of its sure screening property.

Keywords: Sure Independence Screening; Density power Divergence; DPD-SIS; High-dimensional Statistics; Conditional Screening; Robustness.

1 Introduction

The class of generalized linear models (GLMs) is a rich class of parametric regression models that allows to study a wide range of relationship structures for different types of response data; it makes the GLMs to be arguably the most popular statistical tool for real-life applications across several applied disciplines. Let us consider the GLM in its canonical form where, given a set of p predictor variables X_1, X_2, \dots, X_p , the scalar response variables Y follows a distribution from the exponential family having density

$$f(y; \theta) = \exp \{y\theta - b(\theta) + c(y)\}, \quad (1)$$

for some appropriate (known) functions $b(\cdot)$ and $c(\cdot)$ and the unknown canonical parameter θ . For simplicity, we do not consider a dispersion parameter in the model (e.g., logistic or

Poisson regression) although it can easily be incorporated in all our methodological discussions and the theories derived throughout the paper with slight modifications. We concentrate on mean regression model for θ given by

$$E[Y|\mathbf{X} = \mathbf{x}] = b'(\theta) = g^{-1}(\mathbf{x}^T \boldsymbol{\beta}), \quad (2)$$

where $\mathbf{X} = (X_0 = 1, X_1, \dots, X_p)^T$, $\boldsymbol{\beta} = (\beta_0, \beta_1, \dots, \beta_p)^T \in \mathbb{R}^{p+1}$ is the vector of unknown regression coefficients and g is a monotone differentiable link function. Given independent and identically distributed (IID) data (y_i, \mathbf{x}_i) , $i = 1, \dots, n$, our objective is to fit a GLM by efficiently estimating $\boldsymbol{\beta}$ and use it for subsequent inference.

Commonly, the regression coefficient $\boldsymbol{\beta}$ is estimated through the likelihood based approaches (or suitable extensions) under the classical low dimensional set-up ($n < p$). However, recent advancement of technologies across disciplines generates data on a large number of possible covariates with limited observations leading to $p \gg n$, known as the high-dimensional set-up. In this paper, we would consider the ultra-high dimensional GLMs with the number of covariates being of non-polynomial (NP) order of n , i.e., $\log(p) = O(n^l)$ for some $0 < l < 1$. However, to perform meaningful inference under such set-up, we need to assume sparsity of the model — only $s \ll n$ covariates (out of the vast pool of p covariates) are actually important to explain the variability in the response although we do not know which. There are several statistical procedures like LASSO or other regularized approaches (Fan and Li, 2001; Buhlmann and Van De Geer, 2011; Hastie et al., 2015; Giraud, 2014; Ghosh and Majumdar, 2019) to simultaneously select these important variables and estimate the corresponding (non-zero) regression coefficients. Although they often work reasonably well in moderately high dimensions, their computation becomes highly extensive in ultra-high dimensional set-ups. Then it is more efficient to first reduce the set of all covariates to a sufficiently small size (maybe $< n$) through some initial screening procedure. And in this respect, the most popular one is the sure independence screening (SIS) proposed in the pioneer paper by Fan and Lv (2008) for the linear regression model and later extended to GLMs by Fan and Song (2010). The SIS has become extremely popular for its simplicity, elegance, computational speed as well as the theoretical guarantees for sure screening of the true model asymptotically with probability tending to one. Subsequently SIS has been extended to different types of data and associated statistical problems; see, e.g., Barut et al. (2016); Zhao and Li (2012); Luo et al. (2014); Saldana and Feng (2018) among many others.

However, the SIS procedure and its extensions are mainly based on the Pearson correlation or the maximum likelihood estimator (MLE) of the marginal regression coefficients, both of which are extremely non-robust against possible outliers in the data. This non-robustness of SIS was, in fact, first noted in the discussion of the original paper itself by Gather and Guddat (2008); they proposed an alternative robust SIS using the Gnanadesikan-Kettenring correlation in place of the usual correlation while ranking the covariates in a linear regression model. Subsequently, several other robust versions of SIS, mostly non-parametric in nature, are proposed in the literature, for the high-dimensional linear regression model only (Hall and Miller, 2009; Li et al., 2012a,b; Mu and Xiong, 2014; Wang et al., 2017). Although these non-parametric versions of SIS can potentially be applied to the GLMs as well (possibly with appropriate modulation), they are never theoretically studied in the literature. Thus there is a need for a robust variable screening procedure for the ultra-high dimensional

GLM with proper theoretical guarantees of its sure screening property. We aim to fill this gap in the literature by developing a robust sure screening procedure for the general class of GLMs.

Compared to any non-parametric robust procedure, a parametric robust approach is known to provide significantly better efficiency when the assumed model is valid for a majority of the data except only for the noise/contamination part (e.g., Hampel et al., 1986). Recently, a robust parametric version of SIS, namely the DPD-SIS, has been proposed for ultra-high dimensional linear regression models by Ghosh and Thoresen (2020). This DPD-SIS is empirically studied and found to have significantly improved performance compared to the other existing non-parametric SIS procedures under data contamination, although no theoretical guarantees are provided in Ghosh and Thoresen (2020). They have indeed proposed to use the marginal regression approach as in Fan and Song (2010) but to estimate the marginal regression slopes by the robust minimum density power divergence estimator (MDPDE) instead of the MLE. These MDPDEs were first proposed by Basu et al. (1998) as a robust generalization of the MLE for the simple IID problems. Due to the high robustness of these MDPDEs along with high efficiency and simple computation, they are subsequently extended to more complex statistical models. For linear regression models, the MDPDEs are studied by Ghosh and Basu (2013) and subsequently used by Ghosh and Thoresen (2020). For different GLMs as well, the MDPDEs are seen to provide highly efficient and robust parameter estimates under the classical low dimensional set-ups (Basu et al., 2011; Ghosh and Basu, 2016; Basu et al., 2017, 2018; Ghosh, 2019). In this paper, we will utilize the MDPDEs under the marginal regression approach to develop a robust variable screening procedure for the ultra-high dimensional GLMs. Our proposal is thus an extension of the robust DPD-SIS of Ghosh and Thoresen (2020) to the class of GLMs, but we will additionally prove that the proposed procedure indeed satisfy the sure screening property and can also control the selection of false positive under appropriate assumptions on the covariates. The required assumptions are verified to hold under some mild conditions for common examples of GLMs.

A natural extension of the usual SIS has been the conditional SIS (CSIS) proposed by Barut et al. (2016) which takes care of additional information (whenever available) about some previously chosen important variables. Such conditioning variables may be available from similar previously conducted studies or may be estimated via some penalization methods. Among several advantages of such an approach (see Barut et al., 2016), most importantly, it helps to select the hidden important variables (having big impact on the response despite being weakly correlated with it). However, the CSIS is also extremely non-robust under data contamination just as SIS, and there is no literature available on its (parametric) robust version. So, in this paper, we also extend our proposed DPD-SIS further to develop a robust conditional screening procedure under the NP-dimensional GLMs, which we will refer to as the conditional DPD-SIS. The population-level justifications as well as the sample-level sure screening property of this conditional DPD-SIS are also derived rigorously under practically reasonable assumptions.

For simplicity in presentation, all proofs are deferred to the (Online) Supplementary material.

2 The Proposed DPD-SIS for NP-dimensional GLMs

Let us consider the GLM described in (1)-(2) with ultra-high dimensional covariates; for simplicity in presentation, throughout the rest of the paper we will assume canonical link function so that $b' = g^{-1}$ and hence $\theta = \mathbf{x}^T \boldsymbol{\beta}$ in (2). Suppose that the true value of the regression coefficient $\boldsymbol{\beta}$ is denoted by $\boldsymbol{\beta}_0 = (\beta_{00}, \beta_{01}, \dots, \beta_{0p})^T$. We assume that the true model, denoted as $\mathcal{M}_0 = \{1 \leq j \leq p : \beta_{0j} \neq 0\}$, is sparse with model size $s = |\mathcal{M}_0| < n$. Our aim is to perform an initial screening of the covariates in a robust manner such that it indeed cover all the truly important variables corresponding to \mathcal{M}_0 ; this later property is referred to as the sure screening property in the literature.

We follow the marginal regression approach of Fan and Song (2010) to consider the GLM for Y based on X_j (plus an intercept term) separately for each $j = 1, \dots, p$; let us denote the associated regression coefficients for these marginal models by $\boldsymbol{\beta}_j^M = (\beta_{j0}^M, \beta_j^M)$, respectively. However, instead of using the MLE of $\boldsymbol{\beta}_j^M$ as in Fan and Song (2010), we propose to use the MDPDE of these parameters. Note that, each marginal GLM is of low-dimension having only two parameters in $\boldsymbol{\beta}_j^M$. Hence we can follow Ghosh and Basu (2016) and define their MDPDE as the minimizer of appropriately defined DPD measure between the observed data (y_i, \mathbf{x}_i) , $i = 1, \dots, n$, and the assumed GLM density (1). After simplifications, the MDPDE of $\boldsymbol{\beta}_j^M$ with (given) tuning parameter $\alpha > 0$ is defined as

$$\widehat{\boldsymbol{\beta}}_j^{M\alpha} = \left(\widehat{\beta}_{j0}^{M\alpha}, \widehat{\beta}_j^{M\alpha} \right) = \arg \min_{\beta_{j0}, \beta_j} \frac{1}{n} \sum_{i=1}^n l_\alpha(y_i, \beta_{j0} + \beta_j x_{ij}), \quad (3)$$

where $\mathbf{x}_i = (1, x_{i1}, \dots, x_{ip})^T$ for each $i = 1, \dots, n$, and

$$l_\alpha(y, \theta) = \int f(s; \theta)^{1+\alpha} ds - \left(1 + \frac{1}{\alpha} \right) f(y; \theta)^\alpha + \frac{1}{\alpha}. \quad (4)$$

The tuning parameter α in the definition of the MDPDE is known to control the trade-off between the efficiency under pure data and the robustness against data contamination. In fact, $l_0(y, \theta) := \lim_{\alpha \rightarrow 0} l_\alpha(y, \theta) = -\log f(y, \theta)$ so that the MDPDE at $\alpha = 0$ (in a limiting sense) is nothing but the most efficient and highly non-robust MLE. At $\alpha > 0$, the MDPDE provide a robust extension of the MLE having increasing robustness with a slight loss in efficiency as α increases (Ghosh and Basu, 2016). For any given $\alpha \geq 0$, each MDPDE $\widehat{\boldsymbol{\beta}}_j^{M\alpha}$ can also be obtained by solving the corresponding estimating equation $\sum_{i=1}^n \psi_\alpha(y_i, \beta_0 + \beta_j x_{ij}) [1, x_{ij}]^T = \mathbf{0}_2$, where $\xi_\alpha(\theta) = \int (y - b'(\theta)) f(y; \theta)^{1+\alpha} dy$ and

$$\psi_\alpha(y, \theta) = (y - b'(\theta)) f(y; \theta)^\alpha - \xi_\alpha(\theta). \quad (5)$$

Note that $\xi_0(\theta) = 0$ and hence $\psi_0(y, \theta) = (y - b'(\theta))$, the usual score function, which again leads to the MLE. We propose to use an appropriate $\alpha > 0$, to be discussed later, to construct a robust variable screening procedure for GLMs.

Based on the MDPDEs $\widehat{\boldsymbol{\beta}}_j^{M\alpha}$ for the marginal regression coefficients, for each $j = 1, \dots, p$, and a given $\alpha > 0$, we choose a suitable pre-defined threshold γ_n and select the variables in

the set

$$\widehat{M}_\alpha(\gamma_n) = \left\{ 1 \leq j \leq p : \left| \widehat{\beta}_j^{M\alpha} \right| \geq \gamma_n \right\}. \quad (6)$$

By choosing γ_n appropriately, we can reduce the number of covariates from a large p to any smaller target, say $d < n$ so that the subsequent computation becomes feasible. With these d variables selected in $\widehat{M}_\alpha(\gamma_n)$, we can then fit any appropriate low-dimensional estimation procedure or regularized approach to get the estimated coefficient vector, say $\widehat{\beta}_d = (\widehat{\beta}_{d0}, \widehat{\beta}_{d1}, \dots, \widehat{\beta}_{dd})^T$ and subsequently the final model $\widehat{M} = \left\{ 1 \leq j \leq p : \widehat{\beta}_{dj} \neq 0 \right\}$.

It is important to note the similarity with the DPD-SIS in Ghosh and Thoresen (2020), which was proposed for the linear regression model only. In our DPD-SIS, we suggest to choose γ_n from the target of attaining a fixed model size. However, in practice, it may be chosen in several other ways, e.g., controlling the false positives, or prediction error, etc.; see Section 3 for an optimal rate of γ_n . Note that, the case $\alpha = 0$ reduces to the ordinary SIS, which properties are already studied by Fan and Song (2010). Along the same lines, we will show the sure screening property of our proposed DPD-SIS, so that asymptotically $\widehat{M}_\alpha(\gamma_n)$ contains the true model \mathcal{M}_0 with probability tending to one, under appropriate conditions for any given $\alpha > 0$.

Remark 2.1. *The robustness of the proposed DPD-SIS procedure under data contamination follows directly from those of the marginal MDPDEs $\widehat{\beta}_j^{M\alpha}$. Since the existing theory of MDPDE (Basu et al., 2011; Ghosh and Basu, 2016) yields their increasing robustness for the increasing value of $\alpha > 0$, the same is also expected for the proposed DPD-SIS with any $\alpha > 0$. See also similar discussion in Ghosh and Thoresen (2020) for the linear regression model.*

3 Sure Screening property of the DPD-SIS

We are considering the GLM in (1)-(2) with the canonical link function and the true sparse parameter value β_0 having support \mathcal{M}_0 of size $s = |\mathcal{M}_0| < n$, as described in previous sections. Recall that, under the ultra-high dimensional set-up considered here, the number of covariates $p = p_n$ is assumed to grow exponentially with the sample size n ; we also allow the true model size $s = s_n$ to depend on n . Further we assume that the data (y_i, \mathbf{x}_i) , for $i = 1, \dots, n$, are IID from a true joint distribution $\Pi(dy, d\mathbf{x}) = F_{\beta_0}(dy|\mathbf{x})Q(d\mathbf{x})$, where F_{β_0} is the (conditional) distribution corresponding to the GLM in (1)-(2) and Q is the marginal distribution of the covariates (for which no model is assumed). Then, it is straightforward from the definition of ψ_α in (5) that, for any $\alpha \geq 0$,

$$E[\psi_\alpha(Y, \mathbf{x}^T \beta_0) | \mathbf{X} = \mathbf{x}] = E[\psi_\alpha(Y, \mathbf{x}_1^T \beta_{01}) | \mathbf{X} = \mathbf{x}] = \mathbf{0}, \quad (7)$$

where we have used the notation $\mathbf{x}_1 = (x_j : j \in \mathcal{M}_0)^T$ and $\beta_{01} = (\beta_{0j} : j \in \mathcal{M}_0)^T$. Without loss of generality, let us assume $\mathcal{M}_0 = \{1, 2, \dots, s\}$ and consider the partitions $\mathbf{x}^T = (\mathbf{x}_1, \mathbf{x}_2)$, $\beta_0^T = (\beta_{01}^T, \beta_{02}^T)^T$ and similarly $\mathbf{x}_i^T = (\mathbf{x}_{i1}, \mathbf{x}_{i2})$, $\beta^T = (\beta_1, \beta_2)$ and so on for any p -vectors. We also assume that, for simplicity, the covariates are standardized so that $E(X_j) = 0$ and $E(X_j^2) = 1$ for all $j = 1, \dots, p$.

3.1 Population level results

We first investigate the proposed DPD-SIS at the population level. The population version (functional) of the marginal MDPDE $\widehat{\beta}_j^{M\alpha}$, defined in (3), is given by

$$\beta_j^{M\alpha} = (\beta_{j0}^{M\alpha}, \beta_j^{M\alpha}) = \arg \min_{\beta_{j0}, \beta_j} E[l_\alpha(Y, \beta_{j0} + \beta_j X_j)]. \quad (8)$$

This marginal MDPDE functional $\beta_j^{M\alpha}$ then satisfies the estimating equations

$$E[\psi_\alpha(Y, \beta_0 + \beta_j X_j)] = 0, \quad E[\psi_\alpha(Y, \beta_0 + \beta_j X_j) X_j] = 0. \quad (9)$$

Let us denote

$$B_\alpha(v(\mathbf{x})) = b'(\mathbf{x}^T \beta_0) - E[\psi_\alpha(Y, v(\mathbf{x})) | \mathbf{X} = \mathbf{x}]. \quad (10)$$

Clearly $B_\alpha(\mathbf{x}^T \beta_0) = b'(\mathbf{x}^T \beta_0)$ but $B_\alpha(\beta_{j0}^{M\alpha} + \beta_j^{M\alpha} x_j)$ do not necessarily equal $b'(\beta_{j0}^{M\alpha} + \beta_j^{M\alpha} x_j)$. However, at $\alpha = 0$ we always have $B_0(v(\mathbf{x})) = b'(v(\mathbf{x}))$ for any $v(\mathbf{x})$. Using equations (7) and (9), we have proved the following two theorems. They show, at the population level, why the proposed DPD-SIS is expected to have the targeted sure screening property.

Theorem 3.1. *For a given $\alpha \geq 0$, suppose that the function $B_\alpha(t)$ is strictly monotone in t and $B_\alpha(v)$ is constant for any constant v (independent of \mathbf{x}). Then for any $j = 1, \dots, p$, the marginal MDPDE functional $\beta_j^{M\alpha} = 0$ if and only if $\text{Cov}(b'(\mathbf{X}^T \beta_0), X_j) = \text{Cov}(Y, X_j) = 0$.*

Theorem 3.2. *Given any $\alpha \geq 0$, along with the assumptions of Theorem 3.1 on $B_\alpha(\cdot)$, let us additionally assume that either $B'_\alpha(\cdot)$ is bounded or $G_\alpha(|x|) = \sup_{|u| \leq |x|} |B_\alpha(u)|$ satisfies the relation*

$$E[G(a|X_j)|X_j|I(|X_j| \geq n^\eta)] \leq dn^{-\kappa}, \quad (11)$$

for all $j \in \mathcal{M}_0$ and constants $a, d > 0$, $\eta \in (0, \kappa)$. If there exists a constant $c_1 > 0$ such that $|\text{Cov}(b'(\mathbf{X}^T \beta_0), X_j)| \geq c_1 n^{-\kappa}$ for all $j \in \mathcal{M}_0$, then we have

$$\min_{j \in \mathcal{M}_0} |\beta_j^{M\alpha}| \geq c_2 n^{-\kappa}, \quad \text{for some constant } c_2 > 0.$$

The above two theorems are exactly similar to Theorems 1 and 3 of Fan and Song (2010) from the context of SIS, although they require slightly stronger assumptions depending on the value of $\alpha > 0$. These assumptions, however, automatically hold at $\alpha = 0$.

In the same spirit of Fan and Song (2010), Theorem 3.1 implies that if the set of unimportant covariates $\{X_j : j \notin \mathcal{M}_0\}$ is independent of the set of important covariates $\{X_j : j \in \mathcal{M}\}$ then $\beta_j^{M\alpha} = 0$ for all $j \notin \mathcal{M}_0$ and all $\alpha \geq 0$. Further, note that, an important covariate X_j having non-zero correlation with the response has a marginal regression coefficient $\beta_j^{M\alpha} \neq 0$. These together indicate the existence of a threshold γ_n satisfying $\min_{j \in \mathcal{M}_0} |\beta_j^{M\alpha}| \geq \gamma_n$ and $\max_{j \notin \mathcal{M}_0} |\beta_j^{M\alpha}| = 0$. This forms the theoretical basis for the model selection consistency of the proposed DPD-SIS with any $\alpha \geq 0$ and justifies our proposal as a authenticate screening criterion.

On the other hand, Theorem 3.2 provides the conditions to yield $\min_{j \in \mathcal{M}_0} |\beta_j^{M\alpha}| \geq O(n^{-\kappa})$ for some $\kappa < 1/2$, which can be interpreted as the marginal signals being stronger than the maximum stochastic noise level. It is an intuitive necessity for the proposed DPD-SIS, the sample version (6), to have the sure screening property. Other than the α -dependent assumption, one crucial condition in Theorem 3.2 is $|\text{Cov}(b'(\mathbf{X}^T \boldsymbol{\beta}_0), X_j)| \geq c_1 n^{-\kappa}$ which is the same as required by the usual SIS in Fan and Song (2010); it can be further simplified for jointly Gaussian covariates following Proposition 1 of Fan and Song (2010) and the discussion thereafter. This theorem also provides the necessary framework to achieve the sparsity in the final selected model (6).

3.2 Sample level results

We first show that the marginal MDPDEs $\widehat{\boldsymbol{\beta}}_j^{M\alpha}$, $j = 1, \dots, p$, are uniformly consistent at an exponential rate which would then lead to the sure screening property (sample level) of our proposed DPD-SIS. In this regard, let us note that the marginal MDPDE functional $\boldsymbol{\beta}_j^{M\alpha}$ is unique and is an interior point of the parameter space by convexity of the DPD loss function $l_\alpha(Y, \beta_0 + \beta_j X_j)$ in $\boldsymbol{\beta}_j = (\beta_{0j}, \beta_j)$ for each j . So, we can restrict the minimization of the marginal DPD loss function over the compact set $\mathcal{B} = \{|\beta_{j0}| \leq B, |\beta_j| \leq B\}$ for some large enough constant $B > 0$ such that $\boldsymbol{\beta}_j^{M\alpha}$ is also an interior point of \mathcal{B} . For each $j = 1, \dots, p$, let $\mathbf{X}_j = (1, X_j)^T$ and define the matrices

$$\mathbf{J}_{j,\alpha}(\boldsymbol{\beta}_j) = E [\nabla^2 l_\alpha(Y, \beta_0 + \beta_j X_j)] = (1 + \alpha) E [\Gamma_\alpha(\mathbf{X}_j^T \boldsymbol{\beta}_j) \mathbf{X}_j \mathbf{X}_j^T], \quad (12)$$

$$\begin{aligned} \mathbf{K}_{j,\alpha}(\boldsymbol{\beta}_j) &= E \left[(\nabla l_\alpha(Y, \beta_0 + \beta_j X_j)) (\nabla l_\alpha(Y, \beta_0 + \beta_j X_j))^T \right] \\ &= (1 + \alpha)^2 E [\{\Gamma_{2\alpha}(\mathbf{X}_j^T \boldsymbol{\beta}_j) - \xi_\alpha^2(\mathbf{X}_j^T \boldsymbol{\beta}_j)\} \mathbf{X}_j \mathbf{X}_j^T], \end{aligned} \quad (13)$$

$$\text{where } \Gamma_\alpha(\theta) = \int (y - b'(\theta))^2 f(y; \theta)^{1+\alpha} dy. \quad (14)$$

Then, the following assumptions are needed for our subsequent theoretical investigation of the DPD-SIS; here $\alpha \geq 0$ is a fixed given tuning parameter and $\Lambda_{\min}[\cdot]$ and $\Lambda_{\max}[\cdot]$, respectively, denote the minimum and maximum eigenvalues of its argument matrix.

(A1) The GLM is such that the density f^α in (1) is bounded (say, by $L_\alpha > 0$) and the function $b''(\cdot)$ is continuous and positive. Also, $|\xi_\alpha(\cdot)|$ is non-decreasing.

(A2) For all $\boldsymbol{\beta}_j \in \mathcal{B}$, there exists some constant $V > 0$ such that $\Lambda_{\min}[\mathbf{J}_{j,\alpha}(\boldsymbol{\beta}_j)] \geq V$ uniformly over $j = 1, \dots, p$.

(A3) $\mathbf{K}_{j,\alpha}(\boldsymbol{\beta}_j^{M\alpha})$ is finite and positive definite for each $j = 1, \dots, p$. Also, the norm $\|\mathbf{K}_{j,\alpha}(\boldsymbol{\beta}_j)\|_{\mathcal{B}}$ is bounded from above for each j , where $\|\mathbf{K}_{j,\alpha}(\boldsymbol{\beta}_j)\|_{\mathcal{B}} = \sup_{\boldsymbol{\beta}_j \in \mathcal{B}, \|\mathbf{u}\|=1} \|\mathbf{K}_{j,\alpha}(\boldsymbol{\beta}_j)^{1/2} \mathbf{u}\|$.

(A4) There exist an $\epsilon_1 > 0$ and a large constant $K_n > 0$, such that

$$\sup_{\boldsymbol{\beta}_j \in \mathcal{B}: \|\boldsymbol{\beta}_j - \boldsymbol{\beta}_j^{M\alpha}\| \leq \epsilon_1} E [B_\alpha(\mathbf{X}_j^T \boldsymbol{\beta}_j) \|\mathbf{X}_j\|_2 I(|X_j| > K_n)] \leq o\left(\frac{1}{n}\right), \quad \text{for all } j = 1, 2, \dots, p.$$

(A5) The distribution of the covariate X_j is such that, for sufficiently large $t > 0$ and some positive constants m_0, m_1, m_2, m_3 and τ , we have

$$P(|X_j| > t) = (m_1 - m_2)e^{-m_0 t^\tau}, \quad j = 1, 2, \dots, p, \quad \text{and}$$

$$E [\exp (b(\mathbf{X}^T \boldsymbol{\beta}_0 + m_3) - b(\mathbf{X}^T \boldsymbol{\beta}_0))] + E [\exp (b(\mathbf{X}^T \boldsymbol{\beta}_0 - m_3) - b(\mathbf{X}^T \boldsymbol{\beta}_0))] \leq m_2.$$

(A6) $\text{Var}(\mathbf{X}^T \boldsymbol{\beta}_0)$ is bounded both from below and above by finite positive constants.

(A7) Either $b''(\cdot)$ is bounded or $\widetilde{\mathbf{X}} = (X_1, \dots, X_p)^T$ follows an elliptically contoured distribution with variance Σ_1 and $|E [b'(\mathbf{X}^T \boldsymbol{\beta}_0)(\mathbf{X}^T \boldsymbol{\beta}_0 - \beta_{00})]|$ is bounded.

Note that Assumptions (A1)–(A6) are appropriate extensions, in the same spirit, of the assumptions made by Fan and Song (2010) to prove the sure screening property of the usual SIS; they coincide at $\alpha = 0$ since $L_0 = 1$, $\xi_0 \equiv 0$, $\mathbf{J}_{j,0} = \mathbf{K}_{j,0} = E [b''(\mathbf{X}_j^T \boldsymbol{\beta}_j) \mathbf{X}_j \mathbf{X}_j^T]$ and $B_0(v(\mathbf{x})) = b'(v(\mathbf{x}))$ for any $v(\mathbf{x})$. For general $\alpha > 0$ as well, Assumption (A1) clearly holds for most common examples of GLM including the normal, Poisson and logistic regression models; other assumptions are also valid for these GLMs under mild sufficient conditions. Interestingly, Assumptions (A5)–(A7) are independent of the choice of α and are exactly the same as Assumption (D), (F) and (G) of Fan and Song (2010), respectively. In particular, Assumption (A5) ensures that the covariates and the response variable have light tails; it implies, via Lemma 1 of Fan and Song (2010), that

$$P\left(|Y| \geq \frac{m_0}{m_3} t^\tau\right) \leq m_2 e^{-m_0 t^\tau}, \quad \text{for any } t > 0. \quad (15)$$

Assumption (A6), on the other hand, implies that the variance of the response Y is bounded. In fact, denoting the variance of \mathbf{X} by $\Sigma = \text{Diag}\{0, \Sigma_1\}$, Assumption (A6) states that

$$\text{Var}(\mathbf{X}^T \boldsymbol{\beta}_0) = \boldsymbol{\beta}_0^T \Sigma \boldsymbol{\beta}_0 = O(1).$$

Note that the maximum eigenvalue of Σ_1 in Assumption (A7) is the same as $\Lambda_{\max}(\Sigma)$, which is a positive finite number by Assumption (A6). Further, Assumptions (A6)–(A7) along with the positiveness of $b''(\cdot)$ from Assumption (A1) imply that for any $\boldsymbol{\beta}_j$ in the interior of \mathcal{B} (and hence in particular for $\boldsymbol{\beta}_j = \boldsymbol{\beta}_j^{M\alpha}$), we have

$$\|\boldsymbol{\beta}_j\|_2^2 = O(\|\Sigma \boldsymbol{\beta}_0\|_2^2) = O(\Lambda_{\max}(\Sigma)) = O(\Lambda_{\max}(\Sigma_1)). \quad (16)$$

Here, the first equality is as shown in the proof of Theorem 5 of Fan and Song (2010) while the remaining equalities are argued above.

Now, under Assumptions (A1)–(A5), we have the exponential convergence result for the marginal MDPDE as presented in the following Lemma.

Lemma 3.3. *Suppose Assumptions (A1)–(A5) hold for a given $\alpha \geq 0$. Then, for any $t > 0$,*

$$P\left(\sqrt{n} \left| \widehat{\boldsymbol{\beta}}_j^{M\alpha} - \boldsymbol{\beta}_j^{M\alpha} \right| \geq \frac{16k_n^{(\alpha)}}{V}(1+t)\right) \leq e^{-\frac{2t^2}{K_n^2}} + nm_1 e^{-m_0 K_n^\tau}, \quad j = 1, \dots, p, \quad (17)$$

where $k_n^{(\alpha)} = (1 + \alpha) \left[\frac{m_0}{m_3} K_n^2 L_\alpha + |b'(K_n B + B)| L_\alpha + \xi_\alpha(K_n B + B) \right]$.

Note that the constant bounds involved in the above Lemma are independent of the index j leading to the uniform convergence of all the marginal regression models through union bound. We will utilize this fact to derive the sure screening property of the proposed DPD-SIS along with its rate of false positive control (based on (16)), which is presented in the following theorem.

Theorem 3.4. *Suppose Assumptions (A1)–(A5) hold for a given $\alpha \geq 0$ and $\frac{n^{1-2\kappa}}{(k_n K_n)^2} \rightarrow \infty$ as $n \rightarrow \infty$, where $k_n = k_n^{(\alpha)}$ is as defined in Lemma 3.3. Then, we have the following results.*

(a) *For any given $c_3 > 0$, there exists $C > 0$ such that*

$$P\left(\max_{1 \leq j \leq p} |\widehat{\beta}_j^{M\alpha} - \beta_j^{M\alpha}| \geq c_3 n^{-\kappa}\right) \leq pR_n, \quad (18)$$

where

$$R_n = \left[e^{-\frac{n^{1-2\kappa}C}{(k_n K_n)^2}} + nm_1 e^{-m_0 K_n^\tau} \right].$$

(b) *If additionally the assumptions of Theorem 3.2 hold, then taking $\gamma_n = c_4 n^{-\kappa}$ with $c_4 \leq c_2/2$, we have $P\left(\widehat{\mathcal{M}}(\gamma_n) \supset \mathcal{M}_0\right) \geq 1 - sR_n$.*

(c) *If additionally Assumptions (A6)–(A7) hold, taking $\gamma_n = c_4 n^{-2\kappa}$ with $c_4 > 0$, we get*

$$P\left(|\widehat{\mathcal{M}}(\gamma_n)| \leq O(n^{2\kappa} \Lambda_{\max}(\Sigma))\right) \geq 1 - pR_n.$$

It is important to note that the bound R_n in the above theorem is exactly the same (except for the value of $k_n = k_n^{(\alpha)}$) as obtained by Fan and Song (2010) for usual SIS and it would be exponentially small for standard GLMs with appropriate choices of K_n ; see discussions in Fan and Song (2010). Thus, along with the additional robustness property, our proposed DPD-SIS at any $\alpha > 0$ also enjoys the same optimal rate of convergence and false discovery control as well as the similar sure screening property as the usual SIS under slightly modified assumptions. This is the most striking benefit of our proposal in the context of robust variable screening under high-dimensionality. Additionally, the sure screening property of the DPD-SIS, as stated in Theorem 3.4(b), does not depend on the variance and the correlation structure of the covariates for any choices of $\alpha \geq 0$. However, greater correlation among covariates may surely increase the false positive selection which can be seen by the dependence of the size of $\widehat{\mathcal{M}}(\gamma_n)$ selected via the DPD-SIS on Σ or more precisely on $\Lambda_{\max}(\Sigma)$ [Theorem 3.4(c)]. As we have less correlation among covariates and hence smaller values of $\Lambda_{\max}(\Sigma)$, the number of variables selected via our DPD-SIS reduces leading to less false positives due to its sure independence property. As in the usual SIS, we can also achieve model selection consistency for DPD-SIS at any $\alpha \geq 0$, i.e.,

$$P\left(\widehat{\mathcal{M}}(\gamma_n) = \mathcal{M}_0\right) = 1 - o(1),$$

under appropriate assumptions on $\Lambda_{\max}(\Sigma)$ along with proper control of K_n . As a particular (extreme) example, it holds with the choice of γ_n as in Theorem 3.4(b) if we have $|\text{Cov}(b'(\mathbf{X}^T \boldsymbol{\beta}_0), X_j)| = o(n^{-\kappa})$ for all $j \notin \mathcal{M}_0$, along with the other necessary conditions of the theorem depending on $\alpha \geq 0$.

4 Robust Conditional Screening: The DPD-CSIS

Let us now extend the DPD-SIS approach to conditional screening problems in GLMs. Consider the set-up and notation of Section 2. Additionally, suppose that the information is available to always include a set of q covariates, say \mathbf{X}_C and we need to robustly select additional variables from the remaining $d := p - q$ variables (say, \mathbf{X}_D). For simplicity, in this section, we will assume no intercept terms, since that can be easily incorporated within the given \mathbf{X}_C . Further, without loss of generality, we may assume that $\mathbf{X}_C = (X_1, \dots, X_q)$ so that $\mathbf{X}_D = (X_{q+1}, \dots, X_p)$; denote $\mathcal{C} = \{1, \dots, q\}$ and $\mathcal{D} = \{q + 1, \dots, p\}$ and hence $\boldsymbol{\beta}_C = (\beta_1, \dots, \beta_q)^T \in \mathbb{R}^q$ and $\boldsymbol{\beta}_D = (\beta_{q+1}, \dots, \beta_p)^T \in \mathbb{R}^d$. Now, we can easily extend the DPD-SIS, for a given $\alpha \geq 0$, in this case to choose the variables from \mathbf{X}_D based on the marginal estimators defined as

$$\widehat{\boldsymbol{\beta}}_{C_j}^{M\alpha} = \left(\widehat{\boldsymbol{\beta}}_{C_j 1}^{M\alpha}, \widehat{\beta}_j^{M\alpha} \right) = \arg \min_{\boldsymbol{\beta}_C, \beta_j} \frac{1}{n} \sum_{i=1}^n l_\alpha(y_i, \mathbf{x}_{iC}^T \boldsymbol{\beta}_C + \beta_j x_{ij}), \quad j = q + 1, \dots, p, \quad (19)$$

where $l_\alpha(y, \theta)$ is as defined in (4) and \mathbf{x}_{iC} is the i -th observation on \mathbf{X}_C . Then, as in (6), given a suitable pre-defined threshold γ_n , we may select the variables in the set

$$\widehat{M}_\alpha(\gamma_n | \mathcal{D}) = \left\{ q + 1 \leq j \leq p : \left| \widehat{\beta}_j^{M\alpha} \right| \geq \gamma_n \right\}. \quad (20)$$

We refer to this extension as the conditional DPD-SIS, or the DPD-CSIS in short. Clearly, the DPD-CSIS again coincides with the usual CSIS of Barut et al. (2016) at $\alpha = 0$ and provides a robust generalization at $\alpha > 0$. Further, when the conditioning variable set \mathbf{X}_C is empty (or contains only the intercept), we are back to our DPD-SIS. We here study the properties of the DPD-CSIS in line with the results derived in Section 3.

4.1 Population-level results: Justifications behind the DPD-CSIS

Let us continue with the notation of Section 3.1 and additionally assume that $E(X_j | \mathbf{X}_C) = 0$ for all $j \in \mathcal{D}$. We define the population version (functional) of $\widehat{\boldsymbol{\beta}}_{C_j}^{M\alpha}$ from (19) as

$$\boldsymbol{\beta}_{C_j}^{M\alpha} = (\boldsymbol{\beta}_{C_j 1}^{M\alpha}, \beta_j^{M\alpha}) = \arg \min_{\boldsymbol{\beta}_C, \beta_j} E [l_\alpha(Y, \mathbf{X}_C^T \boldsymbol{\beta}_C + \beta_j X_j)], \quad j = q + 1, \dots, p. \quad (21)$$

Additionally, let us define the functional for the baseline parameter given only \mathbf{X}_C , without any additional variable, as

$$\boldsymbol{\beta}_C^{M\alpha} = \arg \min_{\boldsymbol{\beta}_C} E [l_\alpha(Y, \mathbf{X}_C^T \boldsymbol{\beta}_C)]. \quad (22)$$

Then, throughout all theoretical discussions of DPD-CSIS, as in Barut et al. (2016), we need to assume that the functionals $\boldsymbol{\beta}_{C_j}^{M\alpha}$ and $\boldsymbol{\beta}_C^{M\alpha}$ are unique, i.e., the associated marginal problems are fully identifiable. Now, for DPD-CSIS at any given $\alpha \geq 0$, we consider the random variables $m_{\alpha, j}$, for each $j = q + 1, \dots, p$, defined as

$$m_{\alpha, j} = \frac{B_\alpha(\mathbf{X}_{C_j}^T \boldsymbol{\beta}_{C_j}^{M\alpha}) - B_\alpha(\mathbf{X}_C^T \boldsymbol{\beta}_C^{M\alpha})}{\mathbf{X}_{C_j}^T \boldsymbol{\beta}_{C_j}^{M\alpha} - \mathbf{X}_C^T \boldsymbol{\beta}_C^{M\alpha}}, \quad (23)$$

where $\mathbf{X}_{Cj} = (\mathbf{X}_C^T, X_j)^T$ for each j and B_α is as defined in (10). Denote by $\mathcal{M}_{0\mathcal{D}} = \mathcal{M}_0 \cap \mathcal{D}$ the indices of the truly important variables in $\mathbf{X}_{\mathcal{D}}$. Then, we have the following results, in analogue of Theorems 3.1 and 3.2, that justify our DPD-CSIS algorithm as a reasonable procedure for conditional screening. Here, in analogue of Barut et al. (2016), we define

$$\text{Cov}_L(Y, X_j | \mathbf{X}_C) := E[(Y - E_L[Y | \mathbf{X}_C])(X_j - E_L[X_j | \mathbf{X}_C])], \quad \text{for any } j \in \mathcal{D}, \quad (24)$$

where $E_L[\cdot | \mathbf{X}_C]$ denote the best linear regression fit given \mathbf{X}_C ; clearly $E_L[Y | \mathbf{X}_C] = b'(\mathbf{X}_C^T \boldsymbol{\beta}_C^{M\alpha})$.

Theorem 4.1. *For a given $\alpha \geq 0$ and any $j \in \mathcal{D}$, the (conditional) marginal MDPDE functional $\beta_j^{M\alpha}$ in (21) is zero if and only if $\text{Cov}_L(Y, X_j | \mathbf{X}_C) = 0$.*

Theorem 4.2. *Given any $\alpha \geq 0$, suppose that $E[m_{\alpha,j} X_j^2] \leq c_2$ uniformly in $j \in \mathcal{D}$, for some constant c_2 . If there exist constants $c_1 > 0, \kappa < -1/2$ such that $|\text{Cov}_L(Y, X_j | \mathbf{X}_C)| \geq c_1 n^{-\kappa}$ for all $j \in \mathcal{M}_{0\mathcal{D}}$, then we have*

$$\min_{j \in \mathcal{M}_{0\mathcal{D}}} |\beta_j^{M\alpha}| \geq c_3 n^{-\kappa}, \quad \text{for another constant } c_3 > 0.$$

Note that, the conditioning on some covariates in DPD-CSIS helps us to avoid a few necessary assumptions from the corresponding (population level) results (Theorems 3.1, 3.2) on the unconditional DPD-SIS.

4.2 Sample-level properties: Sure Screening via the DPD-CSIS

We now extend the results of Section 3.2 for the unconditional DPD-SIS to the case of conditional screening to show the uniform convergence of the associated (conditional) MDPDEs and the resulting sure screening property of the DPD-CSIS. We continue with the notation of Section 3.2 and assume that Assumptions (A1)–(A7) hold with $\boldsymbol{\beta}_j$ and $\boldsymbol{\beta}_j^{M\alpha}$ replaced by $\boldsymbol{\beta}_{Cj} \in \mathbb{R}^{q+1}$ and $\boldsymbol{\beta}_{Cj}^{M\alpha}$, respectively, in (A2)–(A4). We also assume the following additional condition.

(A8) There exists $C > 0$ such that $\Lambda_{\min}(E[m_{\alpha,j} \mathbf{X}_{Cj} \mathbf{X}_{Cj}^T]) > C$, uniformly over $j \in \mathcal{D}$.

Note that Assumption (A8) is mild (and regular) if B_α is strictly convex implying $m_{\alpha,j} > 0$ almost surely. Further, we define

$$\mathbf{Z} = E(E_L[\mathbf{X}_{\mathcal{D}} | \mathbf{X}_C] [\mathbf{X}^T \boldsymbol{\beta}_0 - \mathbf{X}_C^T \boldsymbol{\beta}_C^{M\alpha}]), \quad (25)$$

$$\Sigma_{\mathcal{D}|C} = E(\mathbf{X}_{\mathcal{D}} - E_L[\mathbf{X}_{\mathcal{D}} | \mathbf{X}_C])(\mathbf{X}_{\mathcal{D}} - E_L[\mathbf{X}_{\mathcal{D}} | \mathbf{X}_C])^T. \quad (26)$$

We can show that Assumptions (A6)–(A8) imply the following analogue of (16) for this conditional case, given by

$$\|\boldsymbol{\beta}_{\mathcal{D}}\|_2^2 = O(\Sigma_{\mathcal{D}|C} + \mathbf{Z}\mathbf{Z}^T). \quad (27)$$

Then, we have the desired results in analogue to Theorem 3.4 for the present conditional case of the DPD-CSIS which is presented in the following theorem. The proof is similar to that of Theorem 3.4, but using (27) instead of (16), and is hence omitted for brevity.

Theorem 4.3. *Suppose that, for a given $\alpha \geq 0$, Assumptions (A1)–(A5) hold with β_j and $\beta_j^{M\alpha}$ replaced by $\beta_{Cj} \in \mathbb{R}^{q+1}$ and $\beta_{Cj}^{M\alpha}$, respectively, in (A2)–(A4). Also, let $\frac{n^{1-2\kappa}}{(k_n K_n)^2} \rightarrow \infty$ as $n \rightarrow \infty$, where $k_n = k_n^{(\alpha)}$ is as defined in Lemma 3.3. Then, we have the following results.*

(a) *For any given $c_3 > 0$, there exists $C > 0$ such that*

$$P\left(\max_{q+1 \leq j \leq p} |\widehat{\beta}_j^{M\alpha} - \beta_j^{M\alpha}| \geq c_3 n^{-\kappa}\right) \leq dR_n, \quad (28)$$

where R_n is as defined in Theorem 3.4.

(b) *If additionally the assumptions of Theorem 4.2 hold, then taking $\gamma_n = c_4 n^{-\kappa}$ with $c_4 \leq c_2/2$, we have $P\left(\widehat{\mathcal{M}}(\gamma_n) \supset \mathcal{M}_0\right) \geq 1 - sR_n$.*

(c) *If additionally Assumptions (A6)–(A8) hold, taking $\gamma_n = c_4 n^{-2\kappa}$ with $c_4 > 0$, we get*

$$P\left(|\widehat{\mathcal{M}}(\gamma_n)| \leq O\left(n^{2\kappa} \Lambda_{\max}(\Sigma_{\mathcal{D}|C} + \mathbf{Z}\mathbf{Z}^T)\right)\right) \geq 1 - dR_n. \quad (29)$$

Note that the rate of convergences in the above theorem is exactly the same as in the unconditional case (Theorem 3.4) and are in line with the existing literature on variable screening. In the particular case of the linear regression model, we have $\mathbf{Z} = \mathbf{0}$, and hence the result (29) in Theorem 4.3 reduces to

$$P\left(|\widehat{\mathcal{M}}(\gamma_n)| \leq O\left(n^{2\kappa} \Lambda_{\max}(\Sigma_{\mathcal{D}|C})\right)\right) \geq 1 - dR_n. \quad (30)$$

In general, if we additionally assume $\|\mathbf{Z}\|_2^2 = o(\Lambda_{\min}(\Sigma_{\mathcal{D}|C}))$, as in Condition 3(iii) of Barut et al. (2016), we can also have (30) instead of (29) in Theorem 4.3.

A Supplementary Material: Proof of the Results

A.1 Proof of Theorem 3.1

The case $\alpha = 0$ is the same as Theorem 1 of Fan and Song (2010), so here we will follow similar arguments to prove the result for a given $\alpha > 0$. Note that $E[l_\alpha(Y, \beta_0 + \beta_j X_j)]$ is still convex in β_j for any $\alpha > 0$.

Fix a $j \in \{1, 2, \dots, p\}$ and assume that $\text{Cov}(b'(\mathbf{X}^T \boldsymbol{\beta}_0), X_j) = \text{Cov}(Y, X_j) = 0$. Then, using the estimating equations (9), we have

$$E[B_\alpha(\beta_{j0}^{M\alpha} + \beta_j^{M\alpha} X_j)X_j] = E[b'(\mathbf{X}^T \boldsymbol{\beta}_0)X_j].$$

But, $E(X_j) = 0$ and hence, this leads to

$$\text{Cov}(B_\alpha(\beta_{j0}^{M\alpha} + \beta_j^{M\alpha} X_j), X_j) = \text{Cov}(b'(\mathbf{X}^T \boldsymbol{\beta}_0), X_j) = 0. \quad (31)$$

By assumption the functions $g(t) = B_\alpha(\beta_{j0}^{M\alpha} + t)$ and $h(t)$ are strictly monotone in t and hence, for $t \neq 0$, we have $[g(t) - g(0)](t - 0) > 0$.

Now, if $\beta_j^{M\alpha} \neq 0$, we take $t = \beta_j^{M\alpha} X_j$ in the above and get

$$\beta_j^{M\alpha} \text{Cov}(g(\beta_j^{M\alpha} X_j), X_j) = E[E[g(t) - g(0)](t - 0)|X_j \neq 0] > 0,$$

contradicting (31). Hence, we must have $\beta_j^{M\alpha} = 0$.

Next, to prove the converse, again fix a $j \in \{1, 2, \dots, p\}$ and assume $\beta_j^{M\alpha} = 0$. Then, from the estimating equations (9) and (7), we have

$$E[B_\alpha(\beta_{j0}^{M\alpha})X_j] = E[b'(\mathbf{X}^T \boldsymbol{\beta}_0)X_j].$$

By our second assumption in the theorem, we have that $B_\alpha(\beta_{j0}^{M\alpha})$ is a constant and hence

$$\text{Cov}(b'(\mathbf{X}^T \boldsymbol{\beta}_0), X_j) = E[b'(\mathbf{X}^T \boldsymbol{\beta}_0)X_j] = B_\alpha(\beta_{j0}^{M\alpha})E[X_j] = 0.$$

□

A.2 Proof of Theorem 3.2

For $\alpha = 0$ this theorem is identical to Theorem 3 of Fan and Song (2010). We now extend their argument to prove it for any given $\alpha > 0$. Let us fix an $\alpha > 0$ and $j \in \mathcal{M}_0$.

First consider the cases where $B'_\alpha(\cdot)$ is bounded and let D be its upper bound. Then $B_\alpha(\cdot)$ is Lipschitz continuous and hence

$$|\{B_\alpha(\beta_{j0}^{M\alpha} + \beta_j^{M\alpha} X_j) - B_\alpha(\beta_{j0}^{M\alpha})\} X_j| \leq D|\beta_j^{M\alpha}|X_j^2.$$

Taking expectation, we get

$$\begin{aligned} D|\beta_j^{M\alpha}| &\geq |E[\{B_\alpha(\beta_{j0}^{M\alpha} + \beta_j^{M\alpha} X_j) - B_\alpha(\beta_{j0}^{M\alpha})\} X_j]| \\ &= |E[B_\alpha(\beta_{j0}^{M\alpha} + \beta_j^{M\alpha} X_j)X_j]| = |\text{Cov}(B_\alpha(\beta_{j0}^{M\alpha} + \beta_j^{M\alpha} X_j), X_j)|. \end{aligned}$$

But, from the estimating equation in (9), we get

$$E[B_\alpha(\beta_{j_0}^{M\alpha} + \beta_j^{M\alpha} X_j) X_j] = E[b'(\mathbf{X}^T \boldsymbol{\beta}_0) X_j],$$

and hence, using $E(X_j) = 0$,

$$\text{Cov}(B_\alpha(\beta_{j_0}^{M\alpha} + \beta_j^{M\alpha} X_j), X_j) = \text{Cov}(b'(\mathbf{X}^T \boldsymbol{\beta}_0), X_j). \quad (32)$$

Then, in view of the condition of the theorem, we get $|\beta_j^{M\alpha}| \geq D_1^{-1} c_1 n^{-\kappa}$ completing the proof of the theorem.

Next, we consider the cases where the second assumption, namely Condition (11), holds. Clearly if $|\beta_j^{M\alpha}| \geq cn^{-\kappa}$ for a sufficiently large universal constant $c > 0$, the result holds and we are done. So, assume $|\beta_j^{M\alpha}| \leq \tilde{c}_1 n^{-\kappa}$ for some $\tilde{c} > 0$ and let $\beta_0^{M\alpha}$ be a constant such that $B_\alpha(\beta_0^{M\alpha}) = E[Y]$. We first prove the following claim.

Claim 1: $|\beta_{j_0}^{M\alpha} - \beta_0^{M\alpha}| \leq \tilde{c}_2$ for all $j \in \mathcal{M}_0$ and some constant $\tilde{c}_2 > 0$.

To prove the claim, we fix a $j \in \mathcal{M}_0$ and consider the marginal MDPDE objective function (population version) as a function of β_0 only as $Q(\beta_0) = E[l_\alpha(Y, \beta_0 + \beta_j^{M\alpha} X_j)]$ so that we get

$$Q'(\beta_0) = E[Y - B_\alpha(\beta_0 + \beta_j^{M\alpha} X_j)] = B_\alpha(\beta_0^{M\alpha}) - E[B_\alpha(\beta_0 + \beta_j^{M\alpha} X_j)].$$

But,

$$\begin{aligned} & |E[B_\alpha(\beta_0 + \beta_j^{M\alpha} X_j)] - B_\alpha(\beta_0)| \\ & \leq \sup_{|x| \leq \tilde{c}_1 n^{\eta-\kappa}} |B_\alpha(\beta_0 + x) - B_\alpha(\beta_0)| + 2E[G(a|X_j)|X_j|I(|X_j| > n^\eta)] \\ & = o(1) + o(1), \end{aligned}$$

by the continuity of $B_\alpha(\cdot)$ and Condition (11). Therefore, we get $Q'(\beta_0) = B_\alpha(\beta_0^{M\alpha}) - B_\alpha(\beta_0) + o(1)$ and hence, for a $\tilde{c}_2 > 0$, we have $Q'(\beta_0^{M\alpha} - \tilde{c}_2) < 0$ and $Q'(\beta_0^{M\alpha} + \tilde{c}_2) > 0$ since $B_\alpha(\cdot)$ is strictly increasing. Hence $|\beta_{j_0}^{M\alpha} - \beta_0^{M\alpha}| \leq \tilde{c}_2$ proving our Claim 1.

Finally, to prove the theorem, we note that if $|X_j| \leq n^\kappa$, then Claim 1 ensures that the points $\beta_{j_0}^{M\alpha}$ and $(\beta_{j_0}^{M\alpha} + \beta_j^{M\alpha} X_j)$, for all $j \in \mathcal{M}_0$, belong to the interval $I = (\beta_0^{M\alpha} - h, \beta_0^{M\alpha} + h)$ independent of j , where $h = \tilde{c} - 1 + \tilde{c}_2$. Let $\tilde{D} = \max_{x \in I} B'_\alpha(x)$, which is finite by Lipschitz continuity of $B_\alpha(\cdot)$ in a neighborhood of $\beta_0^{M\alpha}$ and hence, for $|X_j| \leq n^\kappa$, we have

$$|\{B_\alpha(\beta_{j_0}^{M\alpha} + \beta_j^{M\alpha} X_j) - B_\alpha(\beta_{j_0}^{M\alpha})\}| \leq \tilde{D} |\beta_j^{M\alpha}| X_j^2.$$

Taking expectation over the region $\{|X_j| \leq n^\kappa\}$, we get

$$\begin{aligned} \tilde{D} |\beta_j^{M\alpha}| & \geq |E[\{B_\alpha(\beta_{j_0}^{M\alpha} + \beta_j^{M\alpha} X_j) - B_\alpha(\beta_{j_0}^{M\alpha})\} X_j I(|X_j| \leq n^\kappa)]| \\ & = |\text{Cov}(b'(\mathbf{X}^T \boldsymbol{\beta}_0), X_j)| - A_0 - A_1, \end{aligned} \quad (33)$$

by a similar calculation leading to (32), where $A_m = E[B_\alpha(\beta_{j_0}^{M\alpha} + \beta_j^{M\alpha} X_j^m) X_j I(|X_j| > n^\kappa)]$ for $m = 0, 1$. But, $|\beta_{j_0}^{M\alpha} + \beta_j^{M\alpha} X_j^m| \leq a|X_j|$ for $|X_j| > n^\kappa$ with a sufficiently large n independent of j and m , we get from Condition (11) that

$$A_m \leq E[G(a|X_j)|X_j|I(|X_j| \geq n^\kappa)] \leq dn^{-\kappa},$$

for both $m = 0, 1$. Then, the theorem follows from (33) using the given condition that $|\text{Cov}(b'(\mathbf{X}^T \boldsymbol{\beta}_0), X_j)| \geq c_1 n^{-\kappa}$. \square

A.3 Proof of Lemma 3.3

The result in the lemma holds directly by Theorem 1 of Fan and Song (2010), provided we can show that their Conditions (A), (B) and (C) are implied by our Assumptions (A1)–(A5). In this regard, note that Assumption (A3) is indeed a reformulation of Condition (A) of Fan and Song (2010). Further, Assumption (A2) implies Condition (C) of Fan and Song (2010) via a second order Taylor series expansion of $l_\alpha(Y, \mathbf{X}_j^T \boldsymbol{\beta}_j)$ with respect to $\boldsymbol{\beta}_j$ around $\boldsymbol{\beta}_j = \boldsymbol{\beta}_j^{M\alpha}$. Finally it remains to show that Condition (B) of Fan and Song (2010) holds under Assumptions (A1), (A4) and (A5).

Let us define $\Omega_n = \{(X_j, Y) : |X_j| \leq K_n, |Y| \leq K_n^*\}$, where K_n is as in Assumption (A4) and $K_n^* = \frac{m_0}{m_3} K_n^\tau$ with m_0, m_3 and τ being as in Assumption (A5). Then, for our present case, Condition (B) of Fan and Song (2010) becomes equivalent to

$$|l_\alpha(Y, \mathbf{X}_j^T \boldsymbol{\beta}_j) - l_\alpha(Y, \mathbf{X}_j^T \boldsymbol{\beta}'_j)| I((X_j, Y) \in \Omega_n) \leq k_n^{(\alpha)} |\mathbf{X}_j^T \boldsymbol{\beta}_j - \mathbf{X}_j^T \boldsymbol{\beta}'_j| I((X_j, Y) \in \Omega_n),$$

$$\boldsymbol{\beta}_j, \boldsymbol{\beta}'_j \in \mathcal{B}, \quad (34)$$

$$\text{and} \quad \sup_{\boldsymbol{\beta}_j \in \mathcal{B}: \|\boldsymbol{\beta}_j - \boldsymbol{\beta}_j^{M\alpha}\| \leq \epsilon_1} |E[l_\alpha(Y, \mathbf{X}_j^T \boldsymbol{\beta}_j) - l_\alpha(Y, \mathbf{X}_j^T \boldsymbol{\beta}_j^{M\alpha})] I((X_j, Y) \notin \Omega_n)| \leq o(n^{-1}), \quad (35)$$

where $k_n^{(\alpha)}$ is as defined in the statement of the Lemma and ϵ_1 as in Assumption (A4). First, to show (34), we use a first order Taylor series expansion to get

$$l_\alpha(Y, \mathbf{X}_j^T \boldsymbol{\beta}_j) - l_\alpha(Y, \mathbf{X}_j^T \boldsymbol{\beta}'_j) = D(\tilde{\boldsymbol{\beta}}_j) [\mathbf{X}_j^T \boldsymbol{\beta}_j - \mathbf{X}_j^T \boldsymbol{\beta}'_j], \quad (36)$$

where $\tilde{\boldsymbol{\beta}}_j \in \mathcal{B}$ lies on the line segment joining $\boldsymbol{\beta}_j$ and $\boldsymbol{\beta}'_j$ and

$$D(\tilde{\boldsymbol{\beta}}_j) = (1 + \alpha) \left[\xi_\alpha(\mathbf{X}_j^T \tilde{\boldsymbol{\beta}}_j) - (Y - b'(\mathbf{X}_j^T \tilde{\boldsymbol{\beta}}_j)) f^\alpha(Y; \mathbf{X}_j^T \tilde{\boldsymbol{\beta}}_j) \right].$$

But, on Ω_n , we have

$$\begin{aligned} |D(\tilde{\boldsymbol{\beta}}_j)| &\leq (1 + \alpha) \left[\left| \xi_\alpha(\mathbf{X}_j^T \tilde{\boldsymbol{\beta}}_j) \right| + (|Y| + |b'(\mathbf{X}_j^T \tilde{\boldsymbol{\beta}}_j)|) \left| f^\alpha(Y; \mathbf{X}_j^T \tilde{\boldsymbol{\beta}}_j) \right| \right] \\ &\leq (1 + \alpha) \left[\left| \xi_\alpha(K_n B + B) \right| + \left(\frac{m_0}{m_3} K_n^2 + |b'(K_n B + B)| \right) L_\alpha \right] = k_n^{(\alpha)}, \end{aligned}$$

by Assumption (A1), (A5) and the subsequent result (15). Substituting it in (36), we get Condition (34).

Next, to prove (35), we again consider the expansion (36) with $\boldsymbol{\beta}'_j = \boldsymbol{\beta}_j^{M\alpha}$ and by taking expectation we get

$$\begin{aligned} |E[l_\alpha(Y, \mathbf{X}_j^T \boldsymbol{\beta}_j) - l_\alpha(Y, \mathbf{X}_j^T \boldsymbol{\beta}_j^{M\alpha})]| &= E \left| D(\tilde{\boldsymbol{\beta}}_j) [\mathbf{X}_j^T \boldsymbol{\beta}_j - \mathbf{X}_j^T \boldsymbol{\beta}_j^{M\alpha}] \right| \\ &\leq (1 + \alpha) \|\boldsymbol{\beta}_j - \boldsymbol{\beta}_j^{M\alpha}\|_2 E \left[\left| B_\alpha(\mathbf{X}_j^T \tilde{\boldsymbol{\beta}}_j) - B_\alpha(\mathbf{X}_j^T \boldsymbol{\beta}_j^{M\alpha}) \right| \|\mathbf{X}_j\|_2 \right], \end{aligned}$$

by an application of the Cauchy-Schwartz inequality. Therefore, we get

$$\begin{aligned}
& \sup_{\beta_j \in \mathcal{B}: \|\beta_j - \beta_j^{M\alpha}\| \leq \epsilon_1} |E [l_\alpha(Y, \mathbf{X}_j^T \beta_j) - l_\alpha(Y, \mathbf{X}_j^T \beta_j^{M\alpha})] I((X_j, Y) \notin \Omega_n)| \\
& \leq (1 + \alpha)\epsilon_1 \sup_{\beta_j \in \mathcal{B}: \|\beta_j - \beta_j^{M\alpha}\| \leq \epsilon_1} E \left[|B_\alpha(\mathbf{X}_j^T \tilde{\beta}_j)| \|\mathbf{X}_j\|_2 + |B_\alpha(\mathbf{X}_j^T \beta_j^{M\alpha})| \|\mathbf{X}_j\|_2 \right] I(|X_j| > K_n), \\
& \leq o(n^{-1}),
\end{aligned}$$

by Assumption (A4), and this completes the proof. \square

A.4 Proof of Theorem 3.4

Part (a):

We start with Lemma 3.3 and take $(1 + t) = c_3 V n^{\frac{1}{2} - \kappa} (16k_n^{(\alpha)})^{-1} > 0$ to get

$$P \left(\left| \widehat{\beta}_j^{M\alpha} - \beta_j^{M\alpha} \right| \geq c_3 n^{-\kappa} \right) \leq e^{-\frac{n^{1-2\kappa}}{\kappa_n^2 \kappa_n^2} C} + n m_1 e^{-m_0 K_n^\tau} = R_n, \quad j = 1, \dots, p, \quad (37)$$

Then, the uniform convergence result in Part (a) of the theorem holds from the relation (37) via union bound of probabilities.

Part (b):

Let us consider the event $\mathcal{E}_n = \left\{ \max_{j \in \mathcal{M}_0} \left| \widehat{\beta}_j^{M\alpha} - \beta_j^{M\alpha} \right| \leq c_2 n^{-\kappa} / 2 \right\}$.

By Theorem 3.2, on \mathcal{E}_n , we then have $\left| \widehat{\beta}_j^{M\alpha} \right| \geq c_2 n^{-\kappa} / 2$ for all $j \in \mathcal{M}_0$. Therefore, for the choice of γ_n as given in the statement of the theorem, we have $\mathcal{M}_0 \subset \widehat{\mathcal{M}}_\alpha(\gamma_n)$ on \mathcal{E}_n , and hence

$$P \left(\widehat{\mathcal{M}}(\gamma_n) \supset \mathcal{M}_0 \right) \geq P(\mathcal{E}_n) = 1 - P(\mathcal{E}_n^c).$$

But, since \mathcal{M}_0 has s elements, by a union bound of probability, we get from (37) that $P(\mathcal{E}_n^c) \leq s R_n$ completing the proof of Part (b).

Part (c):

The proof is based on the result (16) evaluated at $\beta_j = \beta_j^{M\alpha}$, which implies that the number of variables having $|\beta_j^{M\alpha}| > \epsilon n^{-\kappa}$ cannot exceed $O(n^{2\kappa} \Lambda_{\max}(\Sigma))$ for any given $\epsilon > 0$. Now, let us consider the event

$$\widetilde{\mathcal{E}}_n = \left\{ \max_{1 \leq j \leq p} \left| \widehat{\beta}_j^{M\alpha} - \beta_j^{M\alpha} \right| \leq \epsilon n^{-\kappa} \right\}.$$

Then, on the event $\widetilde{\mathcal{E}}_n$, we have

$$\left| \left\{ j : \left| \widehat{\beta}_j^{M\alpha} \right| > 2\epsilon n^{-\kappa} \right\} \right| \leq \left| \left\{ j : \left| \beta_j^{M\alpha} \right| > 2\epsilon n^{-\kappa} \right\} \right| \leq O(n^{2\kappa} \Lambda_{\max}(\Sigma)).$$

Hence, taking $\epsilon = c_5 / 2$ for the choice of γ_n as given in the statement of the theorem, we get

$$P \left(\left| \widehat{\mathcal{M}}(\gamma_n) \right| \leq O(n^{2\kappa} \Lambda_{\max}(\Sigma)) \right) \geq P(\widetilde{\mathcal{E}}_n) = 1 - P(\widetilde{\mathcal{E}}_n^c).$$

But, by Part (a) of the theorem, we have $P(\widetilde{\mathcal{E}}_n^c) \leq p R_n$ completing the proof. \square

A.5 Proofs of Theorems 4.1 and 4.2

We note that, for each $j \in \mathcal{D}$, the quantity $\beta_{\mathcal{C}_j}^{M\alpha}$, defined in (21), satisfies the estimating equations given by

$$E [\psi_\alpha (Y, \mathbf{X}_{\mathcal{C}_j}^T \beta_{\mathcal{C}_j}^{M\alpha}) \mathbf{X}_{\mathcal{C}}] = 0, \quad E [\psi_\alpha (Y, \mathbf{X}_{\mathcal{C}_j}^T \beta_{\mathcal{C}_j}^{M\alpha}) X_j] = 0. \quad (38)$$

On the other hand, the baseline quantity $\beta_{\mathcal{C}}^{M\alpha}$, defined in (22), satisfies

$$E [\psi_\alpha (Y, \mathbf{X}_{\mathcal{C}}^T \beta_{\mathcal{C}}^{M\alpha}) \mathbf{X}_{\mathcal{C}}] = 0. \quad (39)$$

Further, for any $j \in \mathcal{D}$, using $E[X_j | \mathbf{X}_{\mathcal{C}}] = 0$, we have

$$\begin{aligned} \text{Cov}_L(Y, X_j | \mathbf{X}_{\mathcal{C}}) &= E [(Y - E[Y | \mathbf{X}_{\mathcal{C}}]) X_j] = E [Y X_j] \\ &= E [E(Y | \mathbf{X}) X_j] = E [b'(\mathbf{X}^T \beta_0) X_j], \end{aligned} \quad (40)$$

and hence, invoking the definitions of ψ_α and B_α , we get

$$\begin{aligned} E [\psi_\alpha (Y, \mathbf{X}_{\mathcal{C}}^T \beta_{\mathcal{C}}^{M\alpha}) X_j] &= E [(B_\alpha(\mathbf{X}_{\mathcal{C}}^T \beta_{\mathcal{C}}^{M\alpha}) - b'(\mathbf{X}^T \beta_0)) X_j] \\ &= EE [(B_\alpha(\mathbf{X}_{\mathcal{C}}^T \beta_{\mathcal{C}}^{M\alpha}) - b'(\mathbf{X}^T \beta_0)) X_j | \mathbf{X}_{\mathcal{C}}] \\ &= -E [b'(\mathbf{X}^T \beta_0) X_j] \\ &= -\text{Cov}_L(Y, X_j | \mathbf{X}_{\mathcal{C}}). \end{aligned} \quad (41)$$

A.5.1 Proof of Theorem 4.1

Firstly, if $\beta_j^{M\alpha} = 0$ for some $j \in \mathcal{D}$, from (38) we get

$$E [\psi_\alpha (Y, \mathbf{X}_{\mathcal{C}}^T \beta_{\mathcal{C}_{j1}}^{M\alpha}) \mathbf{X}_{\mathcal{C}}] = 0, \quad E [\psi_\alpha (Y, \mathbf{X}_{\mathcal{C}}^T \beta_{\mathcal{C}_{j1}}^{M\alpha}) X_j] = 0. \quad (42)$$

Combining the first equation with (39) and the uniqueness of its solution we have $\beta_{\mathcal{C}_{j1}}^{M\alpha} = \beta_{\mathcal{C}}^{M\alpha}$ and hence the second equation in (42) becomes

$$E [\psi_\alpha (Y, \mathbf{X}_{\mathcal{C}}^T \beta_{\mathcal{C}}^{M\alpha}) X_j] = 0. \quad (43)$$

This leads to the desired condition $\text{Cov}_L(Y, X_j | \mathbf{X}_{\mathcal{C}}) = 0$ by (41).

On the other hand, if $\text{Cov}_L(Y, X_j | \mathbf{X}_{\mathcal{C}}) = 0$ for some $j \in \mathcal{D}$, then by (41), the equation in (43) hold. Combining (43) with (39), we see that $\beta_{\mathcal{C}_j}^{M\alpha} = (\beta_{\mathcal{C}}^{M\alpha}, 0)^T$ is a solution of the estimating equations in (38), leading to $\beta_j^{M\alpha} = 0$. \square

A.5.2 Proof of Theorem 4.2

Fix any $j \in \mathcal{M}_{0\mathcal{D}}$ and define $\boldsymbol{\Omega}_j = E [m_{\alpha,j} \mathbf{X}_{\mathcal{C}_j} \mathbf{X}_{\mathcal{C}_j}^T]$ and $\beta_{\Delta,j} = (\beta_{\mathcal{C}_{j1}}^{M\alpha} - \beta_{\mathcal{C}}^{M\alpha})$. Consider a partition of $\boldsymbol{\Omega}_j$ as given by

$$\boldsymbol{\Omega}_j = \begin{bmatrix} \boldsymbol{\Omega}_{11,j} & \boldsymbol{\Omega}_{12,j} \\ \boldsymbol{\Omega}_{21,j}^T & \boldsymbol{\Omega}_{22,j} \end{bmatrix} = \begin{pmatrix} E [m_{\alpha,j} \mathbf{X}_{\mathcal{C}} \mathbf{X}_{\mathcal{C}}^T] & E [m_{\alpha,j} \mathbf{X}_{\mathcal{C}} X_j] \\ E [m_{\alpha,j} X_j \mathbf{X}_{\mathcal{C}}^T] & E [m_{\alpha,j} X_j^2] \end{pmatrix}.$$

Now, from the estimating equations (38) and (39), along with the definitions of B_α and $m_{\alpha,j}$, we get

$$\begin{aligned}
0 &= E \left[(B_\alpha(\mathbf{X}_{\mathcal{C}j}^T \boldsymbol{\beta}_{\mathcal{C}j}^{M\alpha}) - B_\alpha(\mathbf{X}_{\mathcal{C}}^T \boldsymbol{\beta}_{\mathcal{C}}^{M\alpha})) \mathbf{X}_{\mathcal{C}} \right] \\
&= E \left[m_{\alpha,j} (\mathbf{X}_{\mathcal{C}j}^T \boldsymbol{\beta}_{\mathcal{C}j}^{M\alpha} - \mathbf{X}_{\mathcal{C}}^T \boldsymbol{\beta}_{\mathcal{C}}^{M\alpha}) \mathbf{X}_{\mathcal{C}} \right] \\
&= E \left[m_{\alpha,j} (\mathbf{X}_{\mathcal{C}}^T \boldsymbol{\beta}_{\Delta,j} + X_j \boldsymbol{\beta}_j^{M\alpha}) \mathbf{X}_{\mathcal{C}} \right].
\end{aligned}$$

Therefore, by solving, we get $\boldsymbol{\beta}_{\Delta,j} = -\Omega_{11,j}^{-1} \Omega_{12,j} \boldsymbol{\beta}_j^{M\alpha}$. Further, note that, for any integrable function $h(\mathbf{X}_{\mathcal{C}})$, we have

$$E[h(\mathbf{X}_{\mathcal{C}})X_j] = EE[h(\mathbf{X}_{\mathcal{C}})X_j|\mathbf{X}_{\mathcal{C}}] = E[h(\mathbf{X}_{\mathcal{C}})E(X_j|\mathbf{X}_{\mathcal{C}})] = 0.$$

Hence, by (40), (38), and the definition of $m_{\alpha,j}$, we get

$$\begin{aligned}
\text{Cov}_L(Y, X_j | \mathbf{X}_{\mathcal{C}}) &= E[b'(\mathbf{X}^T \boldsymbol{\beta}_0) X_j] = E[(b'(\mathbf{X}^T \boldsymbol{\beta}_0) - B_\alpha(\mathbf{X}_{\mathcal{C}}^T \boldsymbol{\beta}_{\mathcal{C}}^{M\alpha})) X_j] \\
&= E[(B_\alpha(\mathbf{X}_{\mathcal{C}j}^T \boldsymbol{\beta}_{\mathcal{C}j}^{M\alpha}) - B_\alpha(\mathbf{X}_{\mathcal{C}}^T \boldsymbol{\beta}_{\mathcal{C}}^{M\alpha})) X_j] \\
&= E[m_{\alpha,j} (\mathbf{X}_{\mathcal{C}j}^T \boldsymbol{\beta}_{\mathcal{C}j}^{M\alpha} - \mathbf{X}_{\mathcal{C}}^T \boldsymbol{\beta}_{\mathcal{C}}^{M\alpha}) X_j] \\
&= \Omega_{12,j}^T \boldsymbol{\beta}_{\Delta,j} + \Omega_{22,j} \boldsymbol{\beta}_j^{M\alpha} \\
&= \left[\Omega_{22,j} - \Omega_{12,j}^T \Omega_{11,j}^{-1} \Omega_{12,j} \right] \boldsymbol{\beta}_j^{M\alpha}.
\end{aligned}$$

Now, taking absolute value in the above and using the assumptions of the theorem, we get

$$c_1 n^{-\kappa} \leq c_2 |\boldsymbol{\beta}_j^{M\alpha}|.$$

Since this holds for all $j \in \mathcal{M}_{0\mathcal{D}}$, taking minimum over all such j we get the desired conclusion of the theorem with $c_3 = c_1/c_2$. \square

Acknowledgment: A major part of this research work has been done while the first author (AG) was visiting University of Oslo, Norway.

References

- Barut, E., Fan, J., and Verhasselt, A. (2016). Conditional sure independence screening. *J Amer Stat Assoc*, 111(515), 1266-1277.
- Basu, A., Harris, I. R., Hjort, N. L., and Jones, M. C. (1998). Robust and efficient estimation by minimising a density power divergence. *Biometrika*, **85**, 549–559.
- Basu, A., Shioya, H. and Park, C. (2011). *Statistical Inference: The Minimum Distance Approach*. Chapman & Hall/CRC, Boca de Raton.
- Basu, A., Ghosh, A., Mandal, A., Martin, N. and Pardo, L. (2017) A Wald-type test statistic for testing linear Hypothesis in logistic regression models based on minimum density power divergence estimator. *Electron. J. Stat.*, **11**, 2741–2772.
- Basu, A., Ghosh, A., Mandal, A., Martin, N. and Pardo, L. (2018) Robust Wald-type tests in GLM with random design based on minimum density power divergence estimators. *ArXiv Pre-print*, arXiv:1804.00160v3.
- Buhlmann, P., and Van De Geer, S. (2011). *Statistics for high-dimensional data: methods, theory and applications*. Springer Science & Business Media.
- Fan, J. and Li, R. (2001). Variable Selection via Nonconcave Penalized Likelihood and its Oracle Properties. *J Amer Statist Assoc*, 96:1348-1360.
- Fan, J., and Lv, J. (2008). Sure independence screening for ultrahigh dimensional feature space. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 70(5), 849-911.
- Fan, J., and Song, R. (2010). Sure independence screening in generalized linear models with NP-dimensionality. *The Annals of Statistics*, 38(6), 3567-3604.
- Fu, L., and Wang, Y. G. (2018). Variable selection in rank regression for analyzing longitudinal data. *Stat Methods Med Res*, 27(8), 2447-2458.
- Gather, U., and Guddat, C. (2008). Comment on Sure Independence Screening for Ultrahigh Dimensional Feature Space by Fan, JQ and Lv, J. *J Royal Stat Soc B*, 70, 893-895.
- Ghosh, A. (2019). Robust inference under the beta regression model with application to health care studies. *Stat. Method. Med Res*, 28(3), 871–888.
- Ghosh, A., and Basu, A. (2013). Robust estimation for independent non-homogeneous observations using density power divergence with applications to linear regression. *Electron. J. Stat.*, 7, 2420–2456.

- Ghosh, A., and Basu, A. (2016). Robust Estimation in Generalized Linear Models : The Density Power Divergence Approach. *Test*, 25(2), 269–290.
- Ghosh, A. and Majumdar, S. (2019). Ultrahigh-dimensional Robust and Efficient Sparse Regression using Non-Concave Penalized Density Power Divergence. *ArXiv pre-print*, arXiv:1802.04906v4.
- Ghosh, A. and Thoresen, M. (2020). A Robust Variable Screening procedure for Ultra-high dimensional data. *ArXiv pre-print*, .
- Giraud, C. (2014). *Introduction to high-dimensional statistics*. Chapman and Hall/CRC.
- Hall, P., and Miller, H. (2009). Using generalized correlation to effect variable selection in very high dimensional problems. *J Comput Graphical Stat*, 18(3), 533-550.
- Hampel, F. R., Ronchetti, E., Rousseeuw, P. J., and Stahel W.(1986). *Robust Statistics: The Approach Based on Influence Functions*. New York, USA: John Wiley & Sons.
- Hastie, T., Tibshirani, R., and Wainwright, M. (2015). *Statistical learning with sparsity: the lasso and generalizations*. CRC press.
- Li, G., Peng, H., Zhang, J., and Zhu, L. (2012a). Robust rank correlation based screening. *Ann Stat*, 40(3), 1846-1877.
- Li, R., Zhong, W., and Zhu, L. (2012b). Feature screening via distance correlation learning. *J Amer Statist Assoc*, 107(499), 1129-1139.
- Luo, S., Song, R., and Witten, D. (2014). Sure Screening for Gaussian Graphical Models. *Stat*, 1050, 29.4
- Mu, W., and Xiong, S. (2014). Some notes on robust sure independence screening. *J App Stat*, 41(10), 2092–2102.
- Saldana, D. F., and Feng, Y. (2018). SIS: An R package for sure independence screening in ultrahigh-dimensional statistical models. *J Stat Software*, 83(2), 1–25.
- van der Vaart, A. W. (1998). *Asymptotic statistics*. Cambridge University Press.
- Wainwright, M. J. (2019). *High-dimensional statistics: A non-asymptotic viewpoint* (Vol. 48). Cambridge University Press.
- Wang, T., Zheng, L., Li, Z., and Liu, H. (2017). A robust variable screening method for high-dimensional data. *J App Stat*, 44(10), 1839-1855.
- Zhao, S. D., and Li, Y. (2012). Principled sure independence screening for Cox models with ultra-high-dimensional covariates. *J Mult Anal*, 105(1), 397-411.
- Zhong, W. (2014). Robust sure independence screening for ultrahigh dimensional non-normal data. *Acta Mathematica Sinica, English Series*, 30(11), 1885–1896.