

Representations of the fixed point subalgebra of the vertex algebra associated to a non-degenerate even lattice by an automorphism of order 2

Kenichiro Tanabe*

Department of Mathematics
Hokkaido University
Kita 10, Nishi 8, Kita-Ku, Sapporo, Hokkaido, 060-0810
Japan

ktanabe@math.sci.hokudai.ac.jp

Abstract

Let V_L be the vertex algebra associated to a non-degenerate even lattice L , θ the automorphism of V_L induced from the -1 -isometry of L , and V_L^+ the fixed point subalgebra of V_L under the action of θ . We show that every weak V_L^+ -module is completely reducible.

1 Introduction

The vertex (operator) algebras V such that every (\mathbb{N} -graded) weak V -module is completely reducible are a fundamental class of vertex (operator) algebras as with groups, rings, and Lie algebras. In the case of weak modules, a typical example of such a vertex algebra is the vertex algebra V_L associated to

*Research was partially supported by the Grant-in-aid (No. 18K03198) for Scientific Research, JSPS.

a non-degenerate even lattice L [7, 13, 16]. A vertex operator algebra V is called *rational* (resp. *regular*) if every \mathbb{N} -graded weak (resp. weak) V -module is completely reducible. Thus, the vertex operator algebra V_L associated to a positive definite lattice L is an example of regular vertex operator algebra, where we note that the vertex algebra V_L is a vertex operator algebra only in the case when L is positive definite. Examples of other regular, and hence rational vertex operator algebras include the moonshine vertex operator algebra V^\natural [8, 16], vertex operator algebras associated to the irreducible vacuum representations for affine Kac–Moody algebras with positive integer levels [13, 17], vertex operator algebras of irreducible highest weight modules for the Virasoro algebra of highest weight zero and minimal central charges [13, 17, 29], the vertex operator algebras associated to minimal representations of W -algebras [4]. In most cases above, the Zhu algebras, which are associative \mathbb{C} -algebras introduced in [31], play an essential role in showing the rationality since [31, Theorem 2.2.1] shows that for a vertex operator algebra V there is a one to one correspondence between the set of all isomorphism classes of irreducible \mathbb{N} -graded weak V -modules and that of irreducible modules for the Zhu algebra associated to V . We remark that if a vertex operator algebra $V = \bigoplus_{n=0}^{\infty} V_n$ with $V_0 = \mathbb{C}\mathbf{1}$ is regular, then every weak V -module is \mathbb{N} -graded by [3, Theorem 4.5 and Proposition 5.7].

It is expected that the property of a simple vertex operator algebra of being rational is preserved by taking the fixed point subalgebra under the action of any finite automorphism group. Namely, let V be a vertex algebra, G a finite automorphism group of V , and V^G the fixed point subalgebra of V under the action of G : $V^G = \{u \in V \mid gu = u \text{ for all } g \in G\}$. It is conjectured that if a simple vertex operator algebra V is rational, then so is V^G [6]. This conjecture is an analogue of a result due to Levitzki ([20], see also [22, Theorem 1.15]) in ring theory. Let L be a non-degenerate even lattice and θ the automorphism of V_L induced from the -1 -isometry of L . The conjecture has been confirmed in the vertex operator algebra $V_L^+ := V_L^{(\theta)}$ [2, 10], where L is a positive definite even lattice. It is worth mentioning that the moonshine vertex operator algebra V^\natural is constructed as a direct sum of V_Λ^+ and an irreducible V_Λ^+ -module in [16] where Λ is the Leech lattice. This example demonstrates the importance of the fixed point vertex subalgebras, in particular V_L^+ . When L is not positive definite, the vertex algebra V_L is not a vertex operator algebra as mentioned above, however, it is established in [30, Theorems 3.8 and 3.16] that every \mathbb{N} -graded weak V_L^+ -module is completely reducible by using the Zhu algebra. There are some

other examples in which the conjecture is confirmed (cf. [11, 27, 28]).

Contrary to the case of \mathbb{N} -graded weak modules, the study of non- \mathbb{N} -graded weak modules for vertex (operator) algebras has not progressed and is widely known to be difficult because of the absence of useful tool like the Zhu algebras for non- \mathbb{N} -graded weak modules. So far, the vertex algebras V_L such that L are not positive definite are only one class of vertex algebras which are not vertex operator algebras and whose weak modules are well understood. For some vertex operator algebras, the irreducible weak modules with Whittaker vectors, which are non- \mathbb{N} -graded, are classified in [25, 26].

The aim of this paper is the study of weak V_L^+ -modules for any non-degenerate even lattice L of finite rank. More precisely, as a continuation of [24, Theorem 1.1] which classifies the irreducible weak V_L^+ -modules, we shall show the following result, which means that an analogue of the conjecture for the fixed point vertex operator subalgebras above holds for the vertex algebra V_L^+ :

Theorem 1.1. *Every weak V_L^+ -module is completely reducible for any non-degenerate even lattice L of finite rank.*

When L is positive definite, this result is shown in [2, Theorem 7.7] in the case that $\text{rank } L = 1$ and in [10, Theorem 6.5] in the general case.

Let us explain the basic idea briefly. Let L be a non-degenerate even lattice. Since [24, Lemma 7.3] shows that every weak V_L^+ -module has an irreducible weak V_L^+ -module, it is enough to show that $\text{Ext}_{V_L^+}^1(M, W) = 0$ for any pair of irreducible weak V_L^+ -modules M and W . Let $M(1)$ be the Heisenberg vertex operator algebra constructed from $\mathfrak{h} := \mathbb{C} \otimes_{\mathbb{Z}} L$. Then, $M(1)$ is a subalgebra of V_L and the fixed point subspace $M(1)^+ = \{a \in M(1) \mid \theta(a) = a\}$ is a subalgebra of V_L^+ . The irreducible $M(1)^+$ -modules are classified in [14, Theorem 4.5] in the case that $\dim_{\mathbb{C}} \mathfrak{h} = 1$ and in [15, Theorem 6.2.2] in the general case. We have classified the irreducible weak V_L^+ -modules in [24, Theorem 1.1] and we remark that each irreducible weak V_L^+ -module is a direct sum of irreducible $M(1)^+$ -modules. We follow the notation in these results. We first show that for any irreducible $M(1)^+$ -submodule K of a weak V_L^+ -module which is not isomorphic to $M(1, \lambda)$ for any $\lambda \in L$, the weak V_L^+ -submodule generated by K is completely reducible (Lemma 3.1). We note that each irreducible weak V_L^+ -module includes such an irreducible $M(1)^+$ -submodule K . Thus, it is enough to show that for any pair of irreducible weak V_L^+ -modules M and W , any exact sequence of weak

V_L^+ -modules

$$0 \rightarrow W \rightarrow N \xrightarrow{\pi} M \rightarrow 0, \quad (1.1)$$

and the irreducible $M(1)^+$ -submodule K of M in Lemma 3.1, there exists an $M(1)^+$ -submodule of N which is isomorphic to K under the restriction of π in (1.1). When (M, W) is not isomorphic to (V_L^\pm, V_L^\mp) , $(V_{\lambda+L}, V_{\lambda+L})$ with $\lambda \in L^\perp$ and $2\lambda \notin L$, $(V_{\lambda+L}^\rho, V_{\lambda+L}^\sigma)$ with $\lambda \in L^\perp \setminus L$, $2\lambda \in L$, and $\rho, \sigma \in \{+, -\}$, we can carry out this procedure using results in [24] and [30] (Corollary 3.3). For the other (M, W) , we take a certain preimage U of the lowest space of K under the map $\pi : N \rightarrow M$ in (1.1). If $(M, W) \not\cong (V_L^+, V_L^-)$, then a similar argument as in the proof of [24, Lemmas 3.10 and 5.5] shows that U is a subspace of

$$\Omega_{M(1)^+}(N) = \left\{ u \in N \mid \begin{array}{l} a_i u = 0 \text{ for all homogeneous } a \in V \\ \text{and } i > \text{wt } a - 1. \end{array} \right\} \quad (1.2)$$

and is an irreducible module for the Zhu algebra $A(M(1)^+)$. It follows from [24, Corollary 5.9] that the $M(1)^+$ -submodule of N generated by U is isomorphic to K (Lemmas 3.6 and 3.8). If $(M, W) \cong (V_L^+, V_L^-)$, then U is spanned by a preimage u of the vacuum element $\mathbf{1} \in M$. We show that $\omega_0 u = 0$ and hence the V_L^+ -submodule of N generated by u is isomorphic to V_L^+ (Lemma 3.7).

Throughout this paper, complicated computation has been done by a computer algebra system Risa/Asir[23].

The organization of the paper is as follows. In Section 2 we recall some basic properties of the Heisenberg vertex operator algebra $M(1)$, the vertex algebra V_L associated to a non-degenerate even lattice L , and the fixed point subalgebras $M(1)^+$ and V_L^+ . In Section 3 we give a proof of Theorem 1.1.

2 Preliminary

We assume that the reader is familiar with the basic knowledge on vertex algebras as presented in [5, 16, 19, 21].

Throughout this paper, p is a non-zero integer, \mathbb{N} denotes the set of all non-negative integers, \mathbb{Z} denotes the set of all integers, L is a non-degenerate even lattice of finite rank d with a bilinear form $\langle \cdot, \cdot \rangle$, L^\perp is the dual of L : $L^\perp = \{\alpha \in \mathbb{Q} \otimes_{\mathbb{Z}} L \mid \langle \alpha, \beta \rangle \in \mathbb{Z} \text{ for all } \beta \in L\}$, and $(V, Y, \mathbf{1})$ is a vertex

algebra. Recall that V is the underlying vector space, $Y(\cdot, x)$ is the linear map from $V \otimes_{\mathbb{C}} V$ to $V((x))$, and $\mathbf{1}$ is the vacuum vector. Throughout this paper, we always assume that V has an element ω such that $\omega_0 a = a_{-2} \mathbf{1}$ for all $a \in V$. For a vertex operator algebra V , this condition automatically holds since V has the conformal vector (Virasoro element). We write down the definition of a weak V -module:

Definition 2.1. A *weak V -module* M is a vector space over \mathbb{C} equipped with a linear map

$$\begin{aligned} Y_M(\cdot, x) : V \otimes_{\mathbb{C}} M &\rightarrow M((x)) \\ a \otimes u &\mapsto Y_M(a, x)u = \sum_{n \in \mathbb{Z}} a_n u x^{-n-1} \end{aligned} \quad (2.1)$$

such that the following conditions are satisfied:

- (1) $Y_M(\mathbf{1}, x) = \text{id}_M$.
- (2) For $a, b \in V$ and $u \in M$,

$$\begin{aligned} &x_0^{-1} \delta\left(\frac{x_1 - x_2}{x_0}\right) Y_M(a, x_1) Y_M(b, x_2) u - x_0^{-1} \delta\left(\frac{x_2 - x_1}{-x_0}\right) Y_M(b, x_2) Y_M(a, x_1) u \\ &= x_1^{-1} \delta\left(\frac{x_2 + x_0}{x_1}\right) Y_M(Y(a, x_0) b, x_2) u. \end{aligned} \quad (2.2)$$

For $i \in \mathbb{Z}$, we define

$$\mathbb{Z}_{<i} = \{j \in \mathbb{Z} \mid j < i\} \text{ and } \mathbb{Z}_{>i} = \{j \in \mathbb{Z} \mid j > i\}. \quad (2.3)$$

For $n \in \mathbb{C}$ and a weak V -module M , we define $M_n = \{u \in V \mid \omega_1 u = nu\}$. For $a \in V_n$ ($n \in \mathbb{C}$), $\text{wt } a$ denotes n . For a vertex algebra V which admits a decomposition $V = \bigoplus_{n \in \mathbb{Z}} V_n$ and a subset U of a weak V -module, we define

$$\Omega_V(U) = \left\{ u \in U \mid \begin{array}{l} a_i u = 0 \text{ for all homogeneous } a \in V \\ \text{and } i > \text{wt } a - 1. \end{array} \right\}. \quad (2.4)$$

For a vertex algebra V which admits a decomposition $V = \bigoplus_{n \in \mathbb{Z}} V_n$, a weak V -module N is called \mathbb{N} -graded if N admits a decomposition $N = \bigoplus_{n=0}^{\infty} N(n)$ such that $a_i N(n) \subset N(\text{wt } a - i - 1 + n)$ for all homogeneous $a \in V$, $i \in \mathbb{Z}$, and $n \in \mathbb{Z}_{\geq 0}$, where we define $N(n) = 0$ for all $n < 0$. For a triple of weak

V -modules M, N, W , $u \in M, v \in W$, and an intertwining operator $I(\cdot, x)$ from $M \times W$ to N , we write the expansion of $I(u, x)v$ by

$$I(u, x)v = \sum_{i \in \mathbb{C}} u_i v x^{-i-1} \in N\{x\}. \quad (2.5)$$

In this paper, we consider only the case that the image of $I(\cdot, x)$ is contained in $N((x))$, namely $I(\cdot, x) : M \times W \rightarrow N((x))$. For a subset X of W ,

$$M \cdot X \text{ denotes } \text{Span}_{\mathbb{C}}\{a_i u \mid a \in M, i \in \mathbb{Z}, u \in X\} \subset N. \quad (2.6)$$

For an intertwining operator $I(\cdot, x) : M \times W \rightarrow N((x))$, $u \in M$, and $v \in W$, we define $\epsilon_I(u, v) = \epsilon(u, v) \in \mathbb{Z} \cup \{-\infty\}$ by

$$u_{\epsilon_I(u, v)} v \neq 0 \text{ and } u_i v = 0 \text{ for all } i > \epsilon_I(u, v) \quad (2.7)$$

if $I(u, x)v \neq 0$ and $\epsilon_I(u, v) = -\infty$ if $I(u, x)v = 0$. If M is irreducible, then $\epsilon_I(u, v) \in \mathbb{Z}$ by [12, Proposition 11.9]. For a weak module (M, Y_M) , we write $\epsilon_M = \epsilon_{Y_M}$ for simplicity.

We recall the *Zhu algebra* $A(V)$ of a vertex operator algebra V from [31, Section 2]. For homogeneous $a \in V$ and $b \in V$, we define

$$a \circ b = \sum_{i=0}^{\infty} \binom{\text{wt } a}{i} a_{i-2} b \in V \quad (2.8)$$

and

$$a * b = \sum_{i=0}^{\infty} \binom{\text{wt } a}{i} a_{i-1} b \in V. \quad (2.9)$$

We extend (2.8) and (2.9) for any $a \in V$ by linearity. We also define $O(V) = \text{Span}_{\mathbb{C}}\{a \circ b \mid a, b \in V\}$. Then, the quotient space

$$A(V) = M/O(V) \quad (2.10)$$

called the *Zhu algebra* associated to V , is an associative \mathbb{C} -algebra with multiplication (2.9) by [31, Theorem 2.1.1]. In [31, Theorem 2.2.1], Zhu shows that for a vertex operator algebra V there is a one to one correspondence between the set of all isomorphism classes of irreducible \mathbb{N} -graded weak V -modules and that of irreducible modules for the Zhu algebra associated to V .

For a finite dimensional vector space \mathfrak{h} equipped with a non-degenerate symmetric bilinear form $\langle \cdot, \cdot \rangle$, $M(1)$ denotes *the Heisenberg vertex operator algebra* constructed from \mathfrak{h} (cf. [19, Section 6.3]). As a vector space, $M(1)$ is the symmetric algebra of $\mathfrak{h} \otimes \mathbb{C}[t^{-1}]$. For $\alpha \in \mathfrak{h}$ and $n \in \mathbb{Z}_{<0}$, $\alpha(n)$ denotes $\alpha \otimes t^n \in \mathfrak{h} \otimes \mathbb{C}[t^{-1}]$. The conformal vector of $M(1)$ is given by

$$\omega = \frac{1}{2} \sum_{i=1}^{\dim \mathfrak{h}} h_i(-1) h'_i(-1) \mathbf{1} \quad (2.11)$$

where $\{h_1, \dots, h_{\dim \mathfrak{h}}\}$ is a basis of \mathfrak{h} and $\{h'_1, \dots, h'_{\dim \mathfrak{h}}\}$ is its dual basis. For $\beta \in \mathfrak{h}$, $M(1, \beta)$ denotes the irreducible $M(1)$ -module generated by the vector e^β such that $(\alpha(-1)\mathbf{1})_0 e^\beta = \langle \alpha, \beta \rangle e^\beta$ and $(\alpha(-1)\mathbf{1})_n e^\beta = 0$ for all $\alpha \in \mathfrak{h}$ and $n \in \mathbb{Z}_{>0}$.

Let \hat{L} be the canonical central extension of L by the cyclic group $\langle \kappa \rangle$ of order 2 with the commutator map $c(\alpha, \beta) = \kappa^{\langle \alpha, \beta \rangle}$ for $\alpha, \beta \in L$:

$$0 \rightarrow \langle \kappa \rangle \rightarrow \hat{L} \xrightarrow{\pi} L \rightarrow 0. \quad (2.12)$$

Taking $M(1)$ for $\mathfrak{h} := \mathbb{C} \otimes_{\mathbb{Z}} L$, we define $V_{\lambda+L} := \bigoplus_{\beta \in \lambda+L} M(1, \beta)$ for $\lambda + L \in L^\perp/L$. Then, V_L admits a unique vertex algebra structure compatible with the action of $M(1)$ and is called *the lattice vertex algebra* (cf. [19, Section 6.4]). For each $\lambda + L \in L^\perp/L$ the $M(1)$ -module $V_{\lambda+L}$ is an irreducible weak V_L -module which admits the following decomposition:

$$V_{\lambda+L} = \bigoplus_{n \in \langle \lambda, \lambda \rangle / 2 + \mathbb{Z}} (V_{\lambda+L})_n \text{ where } (V_{\lambda+L})_n = \{a \in V_{\lambda+L} \mid \omega_1 a = na\}. \quad (2.13)$$

It is shown in [7, Theorem 3.1] that $\{V_{\lambda+L} \mid \lambda + L \in L^\perp/L\}$ is a complete set of representatives of equivalence classes of the irreducible weak V_L -modules where L^\perp is the dual lattice of L , and in [13, Theorem 3.16] that every weak V_L -module is completely reducible. Note that if L is positive definite, then $\dim_{\mathbb{C}}(V_{\lambda+L})_n < +\infty$ for all $n \in \lambda + L$ and $(V_{\lambda+L})_{\langle \lambda, \lambda \rangle / 2 + i} = 0$ for sufficiently small $i \in \mathbb{Z}$. If L is not positive definite, then

$$\dim_{\mathbb{C}}(V_{\lambda+L})_n = +\infty \quad (2.14)$$

for all $n \in \langle \lambda, \lambda \rangle / 2 + \mathbb{Z}$, which implies that $V_{\lambda+L}$ is not a V_L -module. The -1 -isometry of L induces an automorphism of \hat{L} of order 2 and an automorphism of V_L of order 2. By abuse of notation we denote these automorphisms by

the same symbol θ . For a weak V_L -module M , we define a weak V_L -module $(M \circ \theta, Y_{M \circ \theta})$ by $M \circ \theta = M$ and

$$Y_{M \circ \theta}(a, x) = Y_M(\theta(a), x) \quad (2.15)$$

for $a \in V_L$. Then $V_{\lambda+L} \circ \theta \cong V_{-\lambda+L}$ for $\lambda \in L^\perp$. Thus, for $\lambda \in L^\perp$ with $2\lambda \in L$ we define

$$V_{\lambda+L}^\pm = \{u \in V_{\lambda+L} \mid \theta(u) = \pm u\}. \quad (2.16)$$

For $\alpha \in \mathfrak{h}$, we define

$$E(\alpha) = e^\alpha + \theta(e^\alpha). \quad (2.17)$$

Let $\alpha \in L \setminus \{0\}$. By [19, (6.4.62)], we have

$$\epsilon_{V_{\lambda+L}}(E(\alpha), e^\lambda) = |\langle \alpha, \lambda \rangle| - 1 \quad (2.18)$$

for $\lambda \in L^\perp \setminus \{0, \pm\alpha\}$ and

$$\epsilon_{V_{\lambda+L}^\pm}(E(\alpha), e^\lambda + \theta(e^\lambda)) = |\langle \alpha, \lambda \rangle| - 1. \quad (2.19)$$

for $\lambda \in L^\perp \setminus \{0, \pm\alpha\}$ with $2\lambda \in L$.

Set a submodule $K = \{\theta(a)a^{-1} \mid a \in \hat{L}\}$ of \hat{L} . The vector space $M(1)(\theta)$ denotes the θ -twisted $M(1)$ -module, T_χ denotes the irreducible \hat{L}/K -module associated to a central character χ such that $\chi(\kappa) = -1$, and $V_L^{T_\chi}$ denotes the irreducible θ -twisted V_L -module associated to χ (cf. [16, Section 9] and [9]). We define

$$\begin{aligned} M(1)(\theta)^\pm &= \{u \in M(1)(\theta) \mid \theta(u) = \pm u\}, \\ (V_L^{T_\chi})^\pm &= \{u \in V_L^{T_\chi} \mid \theta(u) = \pm u\}. \end{aligned} \quad (2.20)$$

It is shown in [14, Theorem 4.5] and [15, Theorem 6.2.2] that the following is a complete set of representatives of equivalence classes of the irreducible irreducible $M(1)^+$ -module:

$$M(1)^\pm, M(1, \lambda) \cong M(1, -\lambda) \text{ with } \lambda \in \mathfrak{h} \setminus \{0\}, M(1)(\theta)^\pm. \quad (2.21)$$

It is also shown in [24, Theorem 1.1] that for a non-degenerate even lattice L of finite rank, the following is a complete set of representatives of equivalence classes of the irreducible weak V_L^+ -modules:

- (1) $V_{\lambda+L}^\pm$, $\lambda + L \in L^\perp/L$ with $2\lambda \in L$,
- (2) $V_{\lambda+L} \cong V_{-\lambda+L}$, $\lambda + L \in L^\perp/L$ with $2\lambda \notin L$,
- (3) $V_L^{T_\chi, \pm}$ for any irreducible \hat{L}/K -module T_χ with central character χ .

Let $h^{[1]}, \dots, h^{[d]}$ be an orthonormal basis of \mathfrak{h} . For $i = 1, \dots, d$, we define

$$\begin{aligned}
\omega^{[i]} &= \frac{1}{2}h^{[i]}(-1)^2, \\
\omega &= \omega^{[1]} + \dots + \omega^{[d]}, \\
H^{[i]} &= \frac{1}{3}h^{[i]}(-3)h^{[i]}(-1)\mathbf{1} - \frac{1}{3}h^{[i]}(-2)^2\mathbf{1}, \\
H^{(6),[i]} &= \frac{1}{5}h^{[i]}(-5)h^{[i]}(-1)\mathbf{1} - \frac{13}{10}h^{[i]}(-4)h^{[i]}(-2)\mathbf{1} + \frac{11}{10}(h(-3)^{[i]})^2\mathbf{1}. \quad (2.22)
\end{aligned}$$

We recall the following notation and some results from [15, Sections 4 and 5]: for any pair of distinct elements $i, j \in \{1, \dots, d\}$ and $l, m \in \mathbb{Z}_{>0}$, we define

$$\begin{aligned}
S_{ij}(l, m) &= h^{[i]}(-l)h^{[j]}(-m), \\
E_{ij}^u &= 5S_{ij}(1, 2) + 25S_{ij}(1, 3) + 36S_{ij}(1, 4) + 16S_{ij}(1, 5), \\
E_{ij}^t &= -16S_{ij}(1, 2) + 145S_{ij}(1, 3) + 19S_{ij}(1, 4) + 8S_{ij}(1, 5), \\
\Lambda_{ij} &= 45S_{ij}(1, 2) + 190S_{ij}(1, 3) + 240S_{ij}(1, 4) + 96S_{ij}(1, 5). \quad (2.23)
\end{aligned}$$

It follows from [15, Proposition 5.3.14] that in $A(M(1)^+)$, $A^u = \oplus_{i,j} \mathbb{C}E_{ij}^u$ and $A^t = \oplus_{i,j} \mathbb{C}E_{ij}^t$ are two-sided ideals, each of which is isomorphic to the $d \times d$ matrix algebra and $A^u A^t = A^t A^u = 0$. By [15, Proposition 5.3.15], $A(M(1)^+)/(A^u + A^t)$ is a commutative algebra generated by the images of $\omega^{[i]}$, $H^{[i]}$ and Λ_{jk} where $i = 1, \dots, d$ and $j, k \in \{1, \dots, d\}$ with $j \neq k$.

It is well known that $M(1)^+$ is strongly generated by $\omega^{[i]}$, $H^{[i]}$, and $S_{lm}(1, r)$ ($1 \leq i \leq d, 1 \leq m < l \leq d, r = 1, 2, 3$) in the sense of [18, p.111]. For a coset $\Gamma \in L^\perp/L$ with $\Gamma \neq L$ and $2\Gamma \subset L$, we take a subset S_Γ of Γ so that $\Gamma = S_\Gamma \cup (-S_\Gamma)$ and $S_\Gamma \cap (-S_\Gamma) = \emptyset$ where $-S_\Gamma = \{-\mu \mid \mu \in S_\Gamma\}$. We also take a subset S_L of L so that $L \setminus \{0\} = S_L \cup (-S_L)$ and $S_L \cap (-S_L) = \emptyset$.

Then,

$$\begin{aligned}
V_L^+ &\cong M(1)^+ \oplus \bigoplus_{\alpha \in S_L} M(1, \alpha), \\
V_L^- &\cong M(1)^- \oplus \bigoplus_{\alpha \in S_L} M(1, \alpha), \\
V_{\lambda+L} &\cong \bigoplus_{\mu \in \lambda+L} M(1, \mu) \text{ for } \lambda \in L^\perp \setminus L \text{ with } 2\lambda \notin L, \\
V_{\lambda+L}^\pm &\cong \bigoplus_{\mu \in S_{\lambda+L}} M(1, \mu) \text{ for } \lambda \in L^\perp \setminus L \text{ with } 2\lambda \in L, \\
V_L^{T_X, \pm} &\cong M(1)(\theta)^\pm \otimes_{\mathbb{C}} T_X
\end{aligned} \tag{2.24}$$

as $M(1)^+$ -modules.

The rest of this section we assume $\text{rank } L = \dim_{\mathbb{C}} \mathfrak{h} = 1$. Let $h = h^{[1]}$ be an orthonormal basis of \mathfrak{h} . We write $\omega = \omega^{[1]}$, $H = H^{[1]}$, $H^{(6)} = H^{(6), [1]}$ for simplicity. In [2, (4.4)], H and $H^{(6)}$ are denoted by H^4 and H^6 respectively. We recall the following formulas from [2, (4.6)–(4.9)]: for $m \in \mathbb{Z}$,

$$[H_3, \omega_{m+1}] = -3mH_{m+3} - \frac{m^2(m+1)}{2}\omega_{m+1}, \tag{2.25}$$

$$\begin{aligned}
[H_3, [H_3, \omega_{m+1}]] &= 9m^2H_{m+5}^{(6)} + \frac{3m^3(5m+9)}{4}H_{m+3} \\
&\quad + \frac{m^3(3m^3+7m^2+3m-3)}{10}\omega_{m+1},
\end{aligned} \tag{2.26}$$

$$\begin{aligned}
[H_5^{(6)}, \omega_{m+1}] &= -5mH_{m+5}^{(6)} - \frac{15m^2(m+1)}{4}H_{m+3} \\
&\quad - \frac{m^2(m+1)(2m^2+2m-1)}{6}\omega_{m+1},
\end{aligned} \tag{2.27}$$

$$[H_4, \omega_{m+1}] = -(3m-1)H_{m+4} - \frac{m(m+1)(3m+1)}{6}\omega_{m+2}. \tag{2.28}$$

By (2.25)–(2.27),

$$\begin{aligned}
\omega_{m+1} &= \frac{-5}{m^3}[H_3, \omega_{m+1}] + \frac{5}{m^6}[H_3, [H_3, \omega_{m+1}]] + \frac{9}{m^5}[H_5^{(6)}, \omega_{m+1}] \\
&= \frac{-5}{m^3}(H_3\omega_{m+1} - \omega_{m+1}H_3) \\
&\quad + \frac{5}{m^6}((H_3)^2\omega_{m+1} - 2H_3\omega_{m+1}H_3 + \omega_{m+1}(H_3)^2) \\
&\quad + \frac{9}{m^5}(H_5^{(6)}\omega_{m+1} - \omega_{m+1}H_5^{(6)}). \tag{2.29}
\end{aligned}$$

Let $L = \mathbb{Z}\alpha$. We write $p = \langle \alpha, \alpha \rangle \in 2\mathbb{Z} \setminus \{0\}$ and $E = E(\alpha)$ for simplicity. As stated in [24, [(3.27), (3.32), (3.33), and (3.35)], the following elements in V_L^+ are zero:

$$\begin{aligned}
P^{(8),H} &= -2376\omega_{-2}\omega_{-2}\omega_{-1}\mathbf{1} + 3168\omega_{-3}\omega_{-1}\omega_{-1}\mathbf{1} - 6256\omega_{-3}\omega_{-3}\mathbf{1} - 11799\omega_{-4}\omega_{-2}\mathbf{1} \\
&\quad + 30456\omega_{-5}\omega_{-1}\mathbf{1} + 2310\omega_{-7}\mathbf{1} - 9504\omega_{-1}\omega_{-1}H_{-1}\mathbf{1} - 6024\omega_{-3}H_{-1}\mathbf{1} \\
&\quad - 13419\omega_{-2}H_{-2}\mathbf{1} - 6516\omega_{-1}H_{-3}\mathbf{1} + 11868H_{-5}\mathbf{1} + 5040H_{-1}^2\mathbf{1}, \tag{2.30}
\end{aligned}$$

$$\begin{aligned}
Q^{(4)} &= 2(p-2)(-27 + 54p - 44p^2 + 40p^3)\omega_{-3}E \\
&\quad - 12p(p-2)(-3 + 4p)\omega_{-1}^2E \\
&\quad - 6p(p-2)(-9 + 2p)(-1 + 2p)H_{-1}E \\
&\quad + (-72p^3 - 96p^2 + 210p - 90)\omega_0\omega_{-2}E \\
&\quad + (120p^2 - 48p + 36)\omega_0^2\omega_{-1}E \\
&\quad + (-48p - 9)\omega_0^4E, \tag{2.31}
\end{aligned}$$

$$\begin{aligned}
Q^{(5,1)} &= 3(p-2)(10p^2 - 29p + 32)(10p^2 - 4p + 3)\omega_{-4}E \\
&\quad - 12p(3p-4)(10p^2 - 4p + 3)\omega_{-2}\omega_{-1}E \\
&\quad - 3(p-8)(p-2)(2p-1)(10p^2 - 4p + 3)H_{-2}E \\
&\quad + 8(2p-7)(15p^3 - 22p^2 + 8p - 6)\omega_0\omega_{-3}E \\
&\quad + 24p^2(8p-9)\omega_0\omega_{-1}\omega_{-1}E \\
&\quad - 12(p-2)(2p-1)(6p^2 - 5p + 6)\omega_0H_{-1}E \\
&\quad - 6(2p^3 - 32p^2 + 29p + 12)\omega_0\omega_0\omega_{-2}E \\
&\quad - 6(8p-9)\omega_0\omega_0\omega_0\omega_0E, \tag{2.32}
\end{aligned}$$

$$\begin{aligned}
Q^{(6)} = & 2(3696p^8 - 22564p^7 + 66284p^6 - 84937p^5 + 56207p^4 \\
& - 91528p^3 + 11774p^2 + 29190p - 13500)\omega_{-5}E \\
& - 4p(352p^6 + 2152p^5 - 8282p^4 + 7951p^3 - 11696p^2 \\
& + 6304p - 1542)\omega_{-3}\omega_{-1}E \\
& - 3p(1584p^6 - 5572p^5 + 6456p^4 - 6877p^3 + 5214p^2 \\
& - 3040p + 642)\omega_{-2}\omega_{-2}E \\
& + 720p^3(p-2)(4p-1)\omega_{-1}\omega_{-1}\omega_{-1}E \\
& - 24p(p-2)(2p-1)(44p^4 - 98p^3 + 157p^2 - 88p + 48)\omega_{-1}H_{-1}E \\
& - 3(p-2)(2p-25)(2p-1)^2(44p^4 - 13p^3 + 62p^2 - 48p + 18)H_{-3}E \\
& + 3(1760p^7 - 9382p^6 + 1391p^5 + 28130p^4 - 14380p^3 \\
& + 29762p^2 - 25851p + 7650)\omega_0\omega_{-4}E \\
& + 12p(352p^5 - 1459p^4 + 2396p^3 - 2894p^2 + 1254p - 225)\omega_0\omega_{-2}\omega_{-1}E \\
& - 3(p-2)(2p-1)(352p^5 + 101p^4 + 86p^3 - 614p^2 + 804p - 225)\omega_0H_{-2}E \\
& + 12(88p^6 + 1104p^5 - 4136p^4 + 3714p^3 - 3944p^2 + 2670p - 675)\omega_0\omega_0\omega_{-3}E \\
& - 6(352p^5 - 1099p^4 + 686p^3 - 689p^2 + 804p - 225)\omega_0\omega_0\omega_0\omega_{-2}E \\
& - 90(p-2)(4p-1)\omega_0\omega_0\omega_0\omega_0\omega_0E. \tag{2.33}
\end{aligned}$$

The following result will be used in Lemma 3.8.

Lemma 2.2. *Let $L = \mathbb{Z}\alpha$ be an even lattice of rank one such that $\langle \alpha, \alpha \rangle \notin \{0, 2\}$ and M a weak V_L^+ -module such that V_L^+ is a submodule of M . Let u be a non-zero element of M such that $H_4u, \omega_2u \in \mathbb{C}\mathbf{1} \subset V_L^+$, and $H_{4+i}u = \omega_{2+i}u = 0$ for all $i \in \mathbb{Z}_{>0}$.*

- (1) *If $\epsilon_M(E(\alpha), u) \geq 0$, $\omega_1u = H_3u = u$, and $H_4u = 0$, then $\epsilon_M(\omega, u) = 1$ and $\epsilon_M(E(\alpha), u) = 0$.*
- (2) *If $\epsilon_M(E(\alpha), u) \geq -1$, $\omega_1u = H_3u = 0$, and $H_4u = (-1/3)\omega_2u$, then $\epsilon_M(\omega, u) \leq 0$ and $\epsilon_M(E(\alpha), u) = -1$.*

Proof. We write $p = \langle \alpha, \alpha \rangle$ and $E = E(\alpha)$ for simplicity. Let t be an integer such that $t \geq \epsilon_M(E(\alpha), u)$. Using [24, Lemma 2.2], we expand $Q_{t+4}^{(4)}u, Q_{t+5}^{(5,1)}u$, and $Q_{t+6}^{(6)}u$ so that the resulting expressions are linear combinations of elements of the form

$$a_{i_1}^{(1)} \cdots a_{i_l}^{(l)} E_m b_{j_1}^{(1)} \cdots b_{j_n}^{(n)} u \tag{2.34}$$

where $l, n \in \mathbb{Z}_{\geq 0}$, $m \in \mathbb{Z}$, and

$$\begin{aligned} (a^{(1)}, i_1), \dots, (a^{(l)}, i_l) &\in \{(\omega, k) \mid k \leq 0\} \cup \{(H, k) \mid k \leq 2\}, \\ (b^{(1)}, j_1), \dots, (b^{(n)}, j_n) &\in \{(\omega, k) \mid k \geq 1\} \cup \{(H, k) \mid k \geq 3\}. \end{aligned} \quad (2.35)$$

Then, we have the following results:

$$\begin{aligned} 0 &= -(t+1)^2((16p+3)t^2 + (-16p^2 + 36p + 12)t \\ &\quad + 4p^3 - 18p^2 + 14p + 12)E_t u \\ &\quad + 2((20p^2 - 8p + 6)t^2 + (-16p^3 + 44p^2 - 10p + 12)t \\ &\quad + 4p^4 - 18p^3 + 18p^2 + p + 6)E_t \omega_1 u \\ &\quad + 2((20p^2 - 8p + 6)t^2 + (-28p^3 + 28p^2 + 25p - 3)t \\ &\quad + 20p^4 - 70p^3 + 42p^2 + 6p)E_{t-1} \omega_2 u \\ &\quad - 4p(p-2)(4p-3)E_t \omega_1 \omega_1 u \\ &\quad - 8p(p-2)(4p-3)E_{t-1} \omega_1 \omega_2 u \\ &\quad - 2p(p-2)(2p-9)(2p-1)E_t H_3 u \\ &\quad - 2p(p-2)(2p-9)(2p-1)E_{t-1} H_4 u, \end{aligned} \quad (2.36)$$

$$\begin{aligned} 0 &= 2(t+1)^2((8p-9)t^3 + (72p^2 + 44p - 45)t^2 \\ &\quad + (-78p^3 + 166p^2 + 115p - 78)t + 20p^4 - 88p^3 + 52p^2 + 112p - 48)E_t u \\ &\quad - 4((102p^3 - 64p^2 + 43p - 12)t^2 + (-78p^4 + 226p^3 - 77p^2 + 71p - 24)t \\ &\quad + 20p^5 - 88p^4 + 82p^3 + 20p^2 + 22p - 12)E_t \omega_1 u \\ &\quad - 2((202p^3 - 96p^2 + 57p - 36)t^2 + (-240p^4 + 134p^3 + 560p^2 - 295p + 12)t \\ &\quad + 220p^5 - 818p^4 + 558p^3 + 117p^2 - 102p)E_{t-1} \omega_2 u \\ &\quad - 8p((8p^2 - 9p)t - 30p^3 + 92p^2 - 70p + 12)E_t \omega_1 \omega_1 u \\ &\quad - 4p((32p^2 - 36p)t - 150p^3 + 420p^2 - 305p + 60)E_{t-1} \omega_1 \omega_2 u \\ &\quad + 4(p-2)(2p-1)((6p^2 - 5p + 6)t + 10p^3 - 54p^2 + 10p + 6)E_t H_3 u \\ &\quad + (p-2)(2p-1)((24p^2 - 20p + 24)t + 50p^3 - 300p^2 + 75p)E_{t-1} H_4 u, \end{aligned} \quad (2.37)$$

$$\begin{aligned}
0 = & -30(t+1)^2((12p^2 - 27p + 6)t^4 + (228p^2 - 513p + 114)t^3 \\
& + (704p^5 - 569p^4 + 994p^3 + 335p^2 - 2316p + 501)t^2 \\
& + (-704p^6 + 2246p^5 - 2098p^4 + 2602p^3 + 1314p^2 - 4422p + 864)t \\
& + 176p^7 - 932p^6 + 1496p^5 - 1175p^4 + 410p^3 + 2656p^2 - 3198p + 540)E_t u \\
& - 60((176p^5 - 455p^4 + 514p^3 - 349p^2 + 276p - 45)t^3 \\
& + (-1056p^6 + 2355p^5 - 4527p^4 + 4854p^3 - 2877p^2 + 1791p - 270)t^2 \\
& + (704p^7 - 2950p^6 + 5142p^5 - 8303p^4 + 6886p^3 - 3486p^2 + 2574p - 405)t \\
& - 176p^8 + 932p^7 - 1848p^6 + 2609p^5 - 3033p^4 + 1085p^3 \\
& - 352p^2 + 990p - 180)E_t \omega_1 u \\
& - 6((2112p^5 - 5649p^4 + 5826p^3 - 4179p^2 + 3564p - 675)t^3 \\
& + (-11792p^6 + 21956p^5 - 34675p^4 + 31030p^3 - 11613p^2 + 10944p - 2025)t^2 \\
& + (10384p^7 - 18258p^6 + 8401p^5 + 8859p^4 - 76550p^3 + 86952p^2 - 19887p + 900)t \\
& - 10560p^8 + 50640p^7 - 91020p^6 + 126690p^5 - 114150p^4 \\
& + 22110p^3 + 21600p^2 - 4770p)E_{t-1} \omega_2 u \\
& + 120p((176p^5 - 455p^4 + 694p^3 - 754p^2 + 366p - 45)t \\
& - 352p^6 + 1434p^5 - 2623p^4 + 3729p^3 - 3523p^2 + 1524p - 180)E_t \omega_1 \omega_1 u \\
& + 12p((3872p^5 - 10559p^4 + 16276p^3 - 17974p^2 + 8574p - 1125)t \\
& - 10032p^6 + 36036p^5 - 61980p^4 + 88110p^3 - 81438p^2 + 34164p - 4050)E_{t-1} \omega_1 \omega_2 u \\
& + 720p^3(p-2)(4p-1)E_t \omega_1 \omega_1 \omega_1 u \\
& + 2160p^3(p-2)(4p-1)E_{t-1} \omega_1 \omega_1 \omega_2 u \\
& - 6(p-2)(2p-1)((880p^5 - 190p^4 + 800p^3 - 1580p^2 + 1800p - 450)t \\
& + 880p^6 - 6596p^5 + 9372p^4 - 14533p^3 + 11462p^2 - 72p - 450)E_t H_3 u \\
& - 24p(p-2)(2p-1)(44p^4 - 98p^3 + 157p^2 - 88p + 48)E_t H_3 \omega_1 u \\
& - 24p(p-2)(2p-1)(44p^4 - 98p^3 + 157p^2 - 88p + 48)E_{t-1} H_3 \omega_2 u \\
& - 3(p-2)(2p-1)((2112p^5 - 279p^4 + 1686p^3 - 3774p^2 + 4404p - 1125)t \\
& + 2640p^6 - 22780p^5 + 29470p^4 - 47255p^3 + 39830p^2 - 6000p)E_{t-1} H_4 u \\
& - 24p(p-2)(2p-1)(44p^4 - 98p^3 + 157p^2 - 88p + 48)E_{t-1} H_4 \omega_1 u.
\end{aligned} \tag{2.38}$$

We note that $\omega_i \omega_2 u = \omega_i H_4 u = H_i \omega_2 u = H_i H_4 u = E_i \omega_2 u = E_i H_4 u = 0$ for all $i \geq 0$ in (2.36)–(2.38) since $H_4 u, \omega_2 u \in \mathbb{C}\mathbf{1}$.

- (1) Substituting $H_3 u = \omega_1 u = u$ and $H_4 u = 0$ into (2.36)–(2.38) and deleting the terms including $E_{t-1} \omega_2 u$ from the obtained relations, we

have the following results:

$$\begin{aligned}
0 &= t^2(10p^2 - 4p + 3) \\
&\quad \times ((32p - 36)t^5 + (-80p^2 + 281p - 198)t^4 \\
&\quad + (24p^3 - 376p^2 + 783p - 378)t^3 \\
&\quad + (-12p^4 + 176p^3 - 673p^2 + 887p - 294)t^2 \\
&\quad + (24p^5 + 16p^4 - 370p^3 + 660p^2 - 367p + 78)t \\
&\quad - 8p^6 - 40p^5 + 498p^4 - 1424p^3 + 1697p^2 - 864p + 156)E_t u,
\end{aligned} \tag{2.39}$$

$$\begin{aligned}
0 &= t^2((2400p^4 - 6360p^3 + 4080p^2 - 2100p + 360)t^6 \\
&\quad + (33792p^6 - 87408p^5 + 137589p^4 - 189186p^3 + 124737p^2 - 41898p + 5355)t^5 \\
&\quad + (-81664p^7 + 382200p^6 - 726064p^5 + 1055205p^4 - 1153306p^3 \\
&\quad + 669765p^2 - 193386p + 22095)t^4 \\
&\quad + (25344p^8 - 412844p^7 + 1292964p^6 - 2146889p^5 + 2965056p^4 - 2780396p^3 \\
&\quad + 1427091p^2 - 369111p + 39375)t^3 \\
&\quad + (-9152p^9 + 180504p^8 - 874824p^7 + 1952142p^6 - 2951142p^5 \\
&\quad + 3684897p^4 - 2957890p^3 + 1327089p^2 - 306300p + 30465)t^2 \\
&\quad + (20416p^{10} + 4648p^9 - 377744p^8 + 1165462p^7 - 2048798p^6 + 2665009p^5 \\
&\quad - 2403075p^4 + 1390194p^3 - 493932p^2 + 103029p - 9630)t \\
&\quad - 7040p^{11} - 36640p^{10} + 546080p^9 - 2091064p^8 + 4414496p^7 \\
&\quad - 6655612p^6 + 7729798p^5 - 6378462p^4 + 3419178p^3 - 1123344p^2 \\
&\quad + 210006p - 17820)E_t u.
\end{aligned} \tag{2.40}$$

Deleting t^i ($i \geq 3$) from (2.39) and (2.40) (cf. (1) in the proof of [24, Lemma 3.10]), we have

$$0 = t^2 p(p-2)(2p-1)(8p-9)g_1 g_2 g_3 g_4 g_5 E_t u \tag{2.41}$$

where

$$\begin{aligned}
g_1 &= 720896p^8 - 6533120p^7 + 12732160p^6 - 11571376p^5 + 8753247p^4 \\
&\quad - 6117402p^3 + 2934828p^2 - 718146p + 40095, \\
g_2 &= 30976p^{10} - 93632p^9 - 274896p^8 + 676496p^7 + 70580p^6 \\
&\quad - 1376964p^5 + 1569114p^4 - 766098p^3 + 138753p^2 + 14580p - 6561, \\
g_3 &= 32480690176p^{17} - 686708228096p^{16} + 4100113563648p^{15} \\
&\quad - 9261356843008p^{14} + 4721613180928p^{13} + 27252555600512p^{12} \\
&\quad - 89074476796752p^{11} + 153500461862476p^{10} - 188119215355208p^9 \\
&\quad + 182750873232189p^8 - 146333976903441p^7 + 95994434529360p^6 \\
&\quad - 50207935079160p^5 + 20230310418021p^4 - 6034639211379p^3 \\
&\quad + 1264660375698p^2 - 169751870910p + 11394160650, \\
g_4 &= 108295298266169344p^{28} - 1578632220535422976p^{27} \\
&\quad + 9311723577440993280p^{26} - 30020826118265765888p^{25} \\
&\quad + 56480859583533809664p^{24} - 34201181968036986880p^{23} \\
&\quad - 166134072751850102784p^{22} + 653988138655346800640p^{21} \\
&\quad - 1269321655065859079168p^{20} + 1420723402232025004160p^{19} \\
&\quad - 291714635577902883936p^{18} - 2582683778801669449520p^{17} \\
&\quad + 6770894754722207547920p^{16} - 10944187113221235636592p^{15} \\
&\quad + 13555805386036866935018p^{14} - 13737333902075174510823p^{13} \\
&\quad + 11720517307272109891506p^{12} - 8535382247957070808665p^{11} \\
&\quad + 5336352269983480520232p^{10} - 2867703024491554846995p^9 \\
&\quad + 1322914878412254530550p^8 - 522431105033231973729p^7 \\
&\quad + 175773803489479604430p^6 - 49918422180208562604p^5 \\
&\quad + 11741259740661392544p^4 - 2205270148985139708p^3 \\
&\quad + 309462836760861120p^2 - 28630525595816700p \\
&\quad + 1296455338665000
\end{aligned}$$

$$\begin{aligned}
g_5 = & 125419598061634584576p^{38} - 4058775327682313846784p^{37} \\
& + 61364532083457777467392p^{36} - 582936719229084397731840p^{35} \\
& + 3943379872911033271844864p^{34} - 20371782223780152285790208p^{33} \\
& + 83996702697827014071894016p^{32} - 284517625089600742756231168p^{31} \\
& + 805663391079536900830277632p^{30} - 1920640734900916387237049344p^{29} \\
& + 3836954501857572677445864064p^{28} - 6271979192897114647393521856p^{27} \\
& + 7760578552733394021465806368p^{26} - 4960446924702123953479190320p^{25} \\
& - 7465071488664393767860076264p^{24} + 35708801005589262962946399572p^{23} \\
& - 84222727411601389378170057962p^{22} + 152434039009084653430665673211p^{21} \\
& - 232437295292831894456601157082p^{20} + 309491421641544381910932790025p^{19} \\
& - 366054612786171099332876352444p^{18} + 388035250642103571764606350307p^{17} \\
& - 370337448344968087579269096586p^{16} + 318820688670520334196524737033p^{15} \\
& - 247602958017955809443453495754p^{14} + 173233782350652213882420588060p^{13} \\
& - 108885985952016051266803373628p^{12} + 61222945466874004618327931304p^{11} \\
& - 30611663342976087375389167464p^{10} + 13506540937664381785106978544p^9 \\
& - 5208033211875215795239060032p^8 + 1733872407863047422573508704p^7 \\
& - 490817666374303669993349712p^6 + 115795112960087840103771888p^5 \\
& - 22152274459759464382843776p^4 + 3301481575103941278100080p^3 \\
& - 359562883864819986252000p^2 + 25437319335136373184000p \\
& - 875923032140153280000. \tag{2.42}
\end{aligned}$$

Substituting $t = \epsilon_M(E(\alpha), u)$ into (2.41), we have $\epsilon_M(E(\alpha), u) = 0$ since $g_i \neq 0$ for all $p \in \mathbb{Z}$ and $i = 1, \dots, 5$. Substituting $t = 0$ into (2.36), we have

$$0 = p(10p^3 - 35p^2 + 21p + 3)E(\alpha)_{-1}\omega_2u. \tag{2.43}$$

Since $\omega_2u \in \mathbb{C}\mathbf{1}$, we have $\omega_2u = 0$ and hence $\epsilon_M(\omega, u) = 1$.

- (2) The same argument as above shows that if $\omega_1u = H_3u = 0$ and $H_4u = (-1/3)\omega_2u$, then $\epsilon_M(E(\alpha), u) = -1$. Substituting $t = 0$, $H_3u = \omega_1u = 0$, and $H_4u = (-1/3)\omega_2u$ into (2.36) and deleting the terms including E_0u from the obtained relation, we have $E(\alpha)_{-1}\omega_2u = 0$. Since $\omega_2u \in \mathbb{C}\mathbf{1}$, we have $\omega_2u = 0$ and hence $\epsilon_M(\omega, u) \leq 0$.

□

3 Proof of Theorem 1.1

In this section we shall show Theorem 1.1. Let L be a non-degenerate even lattice of finite rank d . Since [24, Lemma 7.3] shows that every weak V_L^+ -module has an irreducible weak V_L^+ -module, it is enough to show that $\text{Ext}_{V_L^+}^1(M, W) = 0$ for any pair of irreducible weak V_L^+ -modules M and W (cf. the last half of the proof of [2, Theorem 2.8]).

Lemma 3.1. *Let M be a weak V_L^+ -modules and K an $M(1)^+$ -submodule of M which is isomorphic to $M(1)^\pm$, $M(1)(\theta)^\pm$, or $M(1, \lambda)$ for some $\lambda \in L^\perp \setminus L$. Then, $V_L^+ \cdot K$ is a completely reducible weak V_L^+ -module.*

Proof. If $K \cong M(1)(\theta)^\pm$, then $V_L^+ \cdot K$ is a V_L^+ -module by [24, Lemma 7.2 (1)] and hence is completely reducible by [30, Theorem 3.16]. If $K \cong M(1)^+$, then $V_L^+ \cdot K \cong V_L^+$ as V_L^+ -modules by [19, Proposition 4.7.7]. If $K \cong M(1)^-$, then $V_L^+ \cdot K \cong V_L^-$ as V_L^+ -modules by [24, Lemma 7.1]. If $K \cong M(1, \lambda)$ for some $\lambda \in L^\perp \setminus L$, then the proof of [24, Lemma 7.3] shows that $V_L^+ \cdot K \cong V_{\lambda+L}$ if $2\lambda \notin L$ and $V_L^+ \cdot K \cong V_{\lambda+L}^\pm$ or $V_{\lambda+L}^+ \oplus V_{\lambda+L}^-$ if $2\lambda \in L$. Thus, $V_L^+ \cdot K$ is a completely reducible weak V_L^+ -module in each case. The proof is complete. \square

Lemma 3.2. *Let M and W be two irreducible weak V_L^+ -modules. We set a $M(1)^+$ -submodule K of M by*

$$K = \begin{cases} M(1)^+, & \text{if } M \cong V_L^+, \\ M(1)^-, & \text{if } M \cong V_L^-, \\ \text{any irreducible } M(1)^+\text{-submodule of } M, & \text{otherwise.} \end{cases} \quad (3.1)$$

If $\text{Ext}_{M(1)^+}^1(K, U) = 0$ for any $M(1)^+$ -submodule U of W , then $\text{Ext}_{V_L^+}^1(M, W) = 0$.

Proof. Let $0 \rightarrow W \rightarrow N \xrightarrow{\pi} M \rightarrow 0$ be an exact sequence of weak V_L^+ -modules. Since any weak V_L^+ -module is a direct sum of irreducible $M(1)^+$ -modules, it follows from [2, Lemma 2.6] that $\text{Ext}_{M(1)^+}^1(K, W) = 0$ and hence we can take an $M(1)^+$ -submodule of N which is isomorphic to K and intersects with W trivially. We denote again by K this $M(1)^+$ -submodule of N . Since $K \cap W = 0$ and M is an irreducible weak V_L^+ -module, we have $N = W + (V_L^+ \cdot K)$. By Lemma 3.1, $V_L^+ \cdot K$ is a completely reducible weak V_L^+ -module and hence $N \cong M \oplus W$ as V_L^+ -modules. \square

Corollary 3.3. *If a pair (M, W) of irreducible V_L^+ -modules satisfies one of the following conditions, then $\text{Ext}_{V_L^+}^1(M, W) = 0$.*

- (1) $M \cong V_L^+$ and $W \not\cong V_L^-$.
- (2) $M \cong V_L^-$ and $W \not\cong V_L^+$.
- (3) $M \cong V_{\lambda+L}$ with $\lambda \in L^\perp$ and $2\lambda \notin L$ and $W \not\cong V_{\lambda+L}$.
- (4) $M \cong V_{\lambda+L}^+$ with $\lambda \in L^\perp$ and $2\lambda \in L$ and $W \not\cong V_{\lambda+L}^\pm$.
- (5) $M \cong V_{\lambda+L}^-$ with $\lambda \in L^\perp$ and $2\lambda \in L$ and $W \not\cong V_{\lambda+L}^\pm$.
- (6) $M \cong V_L^{T_\chi, \pm}$ where χ is a central character for \hat{L}/K with $\chi(\kappa) = -1$.

Proof. The result follows from (2.24), Lemma 3.2, [24, Proposition 5.8], and [30, Theorem 3.16] \square

Let $h^{[1]}, \dots, h^{[d]}$ be an orthonormal basis of \mathfrak{h} . For a weak V_L^+ -module N and $\zeta = (\zeta^{[i]})_{i=1}^d, \xi = (\xi^{[i]})_{i=1}^d, \rho = (\rho^{[i]})_{i=1}^d \in \mathbb{C}^d$, we define a generalized eigenspace

$$N_{\zeta, \xi, \rho} = \left\{ u \in N \mid \begin{array}{l} \text{For } i = 1, \dots, d \text{ there exists } j \in \mathbb{Z}_{>0} \text{ such that} \\ (\omega_1^{[i]} - \zeta^{[i]})^j u = (H_3^{[i]} - \xi^{[i]})^j u = (H_5^{(6), [i]} - \rho^{[i]})^j u = 0 \end{array} \right\}. \quad (3.2)$$

Let $0 \rightarrow W \rightarrow N \xrightarrow{\pi} M \rightarrow 0$ be an exact sequence of weak V_L^+ -modules where W and M are irreducible weak V_L^+ -modules. For all $u \in N_{\zeta, \xi, \rho}$ and $a, b \in \{\omega_1^{[i]} - \zeta^{[i]}, H_3^{[i]} - \xi^{[i]}, H_5^{(6), [i]} - \rho^{[i]} \mid i = 1, \dots, d\}$, we have

$$au \in M \text{ and } abu = 0 \quad (3.3)$$

since an arbitrary irreducible weak V_L^+ -module is a completely reducible module for the commutative \mathbb{C} -algebra generated by $\{\omega_1^{[i]}, H_3^{[i]}, H_5^{(6), [i]} \mid i = 1, \dots, d\}$. For $\alpha \in \mathfrak{h}$ with $\alpha \in \mathbb{C}h^{[1]}$ and $u \in N_{\zeta, \xi, \rho}$, we have

$$\begin{aligned} & (\omega_1^{[1]} - \zeta^{[1]} - \frac{\langle \alpha, \alpha \rangle}{2} + \epsilon(E(\alpha), u) + 1)^2 E(\alpha)_{\epsilon(E(\alpha), u)} u \\ & = E(\alpha)_{\epsilon(E(\alpha), u)} (\omega_1^{[1]} - \zeta^{[1]})^2 u = 0. \end{aligned} \quad (3.4)$$

Lemma 3.4. *Let L be a non-degenerate integral lattice of finite rank d . and λ a non-zero element of $\mathbb{Q} \otimes_{\mathbb{Z}} L$. Then, there exists a sublattice $\oplus_{i=1}^d \mathbb{Z}\alpha^{[i]}$ of L such that $\langle \alpha_i, \alpha_i \rangle \notin \{0, 2\}$ for all $i = 1, \dots, d$, $\langle \alpha_i, \alpha_j \rangle = 0$ for all pairs of distinct elements $i, j \in \{1, \dots, d\}$, and $\langle \alpha_i, \lambda \rangle \neq 0$ for all $i = 1, \dots, d$.*

Proof. We take a sublattice $\Gamma = \oplus_{i=1}^d \mathbb{Z}\alpha^{[i]}$ of L so that $\langle \alpha^{[i]}, \alpha^{[i]} \rangle \neq 0$ for all $i = 1, \dots, d$ and $\langle \alpha^{[i]}, \alpha^{[j]} \rangle = 0$ for any distinct pair of elements $i, j \in \{1, \dots, d\}$. Since L is non-degenerate, we may assume $\langle \alpha^{[1]}, \lambda \rangle \neq 0$. If $\langle \alpha^{[2]}, \lambda \rangle = 0$, then writing $p = \langle \alpha^{[1]}, \alpha^{[1]} \rangle$ and $q = \langle \alpha^{[2]}, \alpha^{[2]} \rangle$ for simplicity, we take a pair of non-zero integers x and y so that $px^2 + qy^2 \notin \{0, \pm 1, \pm 2\}$ and set

$$\beta^{[1]} := x\alpha^{[1]} + y\alpha^{[2]}, \quad \beta^{[2]} := -yq\alpha^{[1]} + px\alpha^{[2]} \in L. \quad (3.5)$$

Then,

$$\begin{aligned} \langle \beta^{[1]}, \beta^{[1]} \rangle &= px^2 + qy^2 \neq 0, & \langle \beta^{[2]}, \beta^{[2]} \rangle &= pq(px^2 + qy^2) \neq 0, \\ \langle \beta^{[1]}, \beta^{[2]} \rangle &= 0, \\ \langle \beta^{[i]}, \alpha^{[j]} \rangle &= 0 \quad (i = 1, 2, j = 3, \dots, d), \\ \langle \beta^{[1]}, \lambda \rangle &= x\langle \alpha^{[1]}, \lambda \rangle \neq 0, & \langle \beta^{[2]}, \lambda \rangle &= -yq\langle \alpha^{[1]}, \lambda \rangle \neq 0, \end{aligned} \quad (3.6)$$

and

$$(\alpha^{[1]}, \alpha^{[2]}) = \frac{1}{(px^2 + qy^2)} (\beta^{[1]}, \beta^{[2]}) \begin{pmatrix} px & yq \\ -y & x \end{pmatrix}. \quad (3.7)$$

We replace $\alpha^{[1]}$ and $\alpha^{[2]}$ by $\beta^{[1]}$ and $\beta^{[2]}$ respectively. Repeating this procedure, we have the result. \square

Remark 3.5. The reason why we avoid the case that $\langle \alpha^{[i]}, \alpha^{[i]} \rangle = 2$ in Lemma 3.4 is that when $\langle \alpha^{[i]}, \alpha^{[i]} \rangle = 2$, we need to change the generators of $V_{\mathbb{Z}\alpha^{[i]}}^+$ and hence the commutation relations in the proof of Lemmas 3.6 and 3.8 below.

Lemma 3.6. *For $\lambda \in L^\perp$ with $2\lambda \notin L$, we have $\text{Ext}_{V_L^+}^1(V_{\lambda+L}, V_{\lambda+L}) = 0$. For $\lambda \in L^\perp \setminus L$ with $2\lambda \in L$, we have $\text{Ext}_{V_L^+}^1(V_{\lambda+L}^+, V_{\lambda+L}^+) = \text{Ext}_{V_L^+}^1(V_{\lambda+L}^-, V_{\lambda+L}^-) = \text{Ext}_{V_L^+}^1(V_{\lambda+L}^+, V_{\lambda+L}^-) = \text{Ext}_{V_L^+}^1(V_{\lambda+L}^-, V_{\lambda+L}^+) = 0$.*

Proof. Let $\lambda \in L^\perp \setminus L$. By Lemma 3.4, we can take a sublattice $\oplus_{i=1}^d \mathbb{Z}\alpha_i$ of L of rank d such that $\langle \alpha_i, \alpha_i \rangle \notin \{0, 2\}$ for all $i = 1, \dots, d$, $\langle \alpha_i, \alpha_j \rangle = 0$ for any

distinct pair of elements $i, j \in \{1, \dots, d\}$, and $\langle \alpha_i, \lambda \rangle \neq 0$ for all $i = 1, \dots, d$. We take an orthonormal basis $h^{[1]}, \dots, h^{[d]}$ of $\mathfrak{h} = \mathbb{C} \otimes_{\mathbb{Z}} L$ defined by

$$h^{[i]} = \frac{1}{\sqrt{\langle \alpha^{[i]}, \alpha^{[i]} \rangle}} \alpha^{[i]} \quad (i = 1, \dots, d). \quad (3.8)$$

We set $\zeta = (\langle \lambda, h^{[1]} \rangle^2/2, \dots, \langle \lambda, h^{[d]} \rangle^2/2)$ and $\mathbf{0} = (0, \dots, 0)$. When $2\lambda \notin L$, we set $(M, W) = (V_{\lambda+L}, V_{\lambda+L})$, and $v = e^\lambda \in M, w = e^\lambda \in W$. When $2\lambda \in L$, we set $(M, W) = (V_{\lambda+L}^\rho, V_{\lambda+L}^\sigma)$ with $\rho, \sigma \in \{+, -\}$, and $v = e^\lambda \pm \theta(e^\lambda) \in M = V_{\lambda+L}^\pm, w = e^\lambda \pm \theta(e^\lambda) \in W = V_{\lambda+L}^\pm$.

Let $0 \rightarrow W \rightarrow N \xrightarrow{\pi} M \rightarrow 0$ be an exact sequence of weak V_L^+ -modules. We take $u \in N_{\zeta, \mathbf{0}, \mathbf{0}}$ such that $\pi(u) = v$. We fix $i \in \{1, \dots, d\}$. Using a slight modification of the proof of [2, Lemma 4.8], we shall first show that $\omega_{2+j}^{[i]} u = H_{4+j}^{[i]} u = 0$ for all $j \geq 0$. It follows from [2, Proposition 4.3] that $H_3^{[i]} u, H_5^{6, [i]} u \in \mathbb{C}w \subset W$ and hence

$$\omega_j^{[i]} H_3^{[i]} u = \omega_j^{[i]} H_5^{(6), [i]} u = 0 \quad (3.9)$$

for all $j \in \mathbb{Z}_{\geq 2}$. For $m \in \mathbb{Z}_{\geq 1}$, by (2.29) and (3.9),

$$0 = (-1 + \frac{-5}{m^3} H_3^{[i]} + \frac{5}{m^6} (H_3^{[i]})^2 + \frac{9}{m^5} H_5^{(6), [i]}) \omega_{m+1}^{[i]} u. \quad (3.10)$$

It follows from [2, Proposition 4.3] that $H_3^{[i]}$ acts diagonally on W and the eigenvalues of $H_3^{[i]}$ on W are non-negative integers. Since $0 \neq -1 - 5a + 5a^2 + 9b$ for any pair of $a, b \in \mathbb{Z}_{\geq 0}$ and $\omega_2^{[i]} u \in W$, it follows from (3.10) with $m = 1$ that $\omega_2^{[i]} u = 0$. We have $H_4^{[i]} u = 0$ by (2.25), $H_6^{(6), [i]} u = 0$ by (2.27), and $H_5^{[i]} u = (-2/3)\omega_3 u$ by (2.28). By (2.25) with $m = 2$ and (3.9),

$$H_3^{[i]} \omega_3^{[i]} u = -6H_5^{[i]} u - 6\omega_3^{[i]} u = -2\omega_3^{[i]} u. \quad (3.11)$$

Since any eigenvalue of $H_3^{[i]}$ on W is non-negative, we have $\omega_3^{[i]} u = 0$. It follows from (2.25) and

$$[\omega_j^{[i]}, \omega_k^{[i]}] = (i - j)\omega_{j+k-1}^{[i]} + \delta_{j+k-2, 0} \frac{j(j-1)(j-2)}{12} \quad (3.12)$$

for all $j, k \in \mathbb{Z}$ that $\omega_{2+j}^{[i]} u = H_{4+j}^{[i]} u = 0$ for all $j \geq 0$.

We shall show that u is a simultaneous eigenvector for $\{\omega_1^{[i]}, H_3^{[i]}\}_{i=1}^d$ and is an element of $\Omega_{M(1)+(N_{\zeta, \mathbf{0}, \mathbf{0}})}$. We fix $i = 1, \dots, d$. After renumbering $\alpha_1, \dots, \alpha_d$, we may assume $i = 1$ and we write $\alpha = \alpha^{[1]}$, $p = \langle \alpha^{[1]}, \alpha^{[1]} \rangle$, $E = E(\alpha^{[1]})$, and $t = \epsilon(E(\alpha^{[1]}), u)$ for simplicity. We may assume

$$\langle \lambda, \alpha \rangle \geq 0. \quad (3.13)$$

Since $N/W \cong M$, we have

$$t \geq \langle \lambda, \alpha \rangle - 1 \quad (3.14)$$

by (2.18) and (2.19). In [24, (3.68)–(3.71)] we have computed $Q_{t+4}^{(4)} u$, $Q_{t+5}^{(5,1)} u$, $Q_{t+5}^{(5,2)} u$, and $Q_{t+6}^{(6)} u$ so that the resulting expressions are linear combinations of elements of the form

$$a_{i_1}^{(1)} \cdots a_{i_l}^{(l)} E_m b_{j_1}^{(1)} \cdots b_{j_n}^{(n)} u \quad (3.15)$$

where $l, n \in \mathbb{Z}_{\geq 0}$, $m \in \mathbb{Z}$, and

$$\begin{aligned} (a^{(1)}, i_1), \dots, (a^{(l)}, i_l) &\in \{(\omega, k) \mid k \leq 1\} \cup \{(H, k) \mid k \leq 2\}, \\ (b^{(1)}, j_1), \dots, (b^{(n)}, j_n) &\in \{(\omega, k) \mid k \geq 2\} \cup \{(H, k) \mid k \geq 3\}. \end{aligned} \quad (3.16)$$

We shall compute $Q_{t+4}^{(4)} u$, $Q_{t+5}^{(5,1)} u$, $Q_{t+5}^{(5,2)} u$, and $Q_{t+6}^{(6)} u$ so that the resulting expressions are linear combinations of elements of the form

$$a_{i_1}^{(1)} \cdots a_{i_l}^{(l)} E_m b_{j_1}^{(1)} \cdots b_{j_n}^{(n)} u \quad (3.17)$$

where $l, n \in \mathbb{Z}_{\geq 0}$, $m \in \mathbb{Z}$, and

$$\begin{aligned} (a^{(1)}, i_1), \dots, (a^{(l)}, i_l) &\in \{(\omega, k) \mid k \leq 1\} \cup \{(H, k) \mid k \leq 3\}, \\ (b^{(1)}, j_1), \dots, (b^{(n)}, j_n) &\in \{(\omega, k) \mid k \geq 2\} \cup \{(H, k) \mid k \geq 4\}. \end{aligned} \quad (3.18)$$

The results are

$$\begin{aligned}
0 &= ((t+1-p)^2 - 2p\omega_1^{[1]}) \\
&\quad \times \left(-(p-2)(p-2\omega_1^{[1]})(4p-3) + 2p(8p-11)t - (16p+3)t^2 \right) E_t u \\
&\quad - 2p(p-2)(2p-9)(2p-1)H_3^{[1]} E_t u, \tag{3.19}
\end{aligned}$$

$$\begin{aligned}
0 &= ((t+1-p)^2 - 2p\omega_1^{[1]}) \\
&\quad \times \left(2(p-2)(p-2\omega_1^{[1]})(15p^2 - 16p + 3) \right. \\
&\quad \quad - (118p^3 + (-16\omega_1^{[1]} - 193)p^2 + (18\omega_1^{[1]} + 35)p + 6)t \\
&\quad \quad \left. + (112p^2 + 6p - 21)t^2 + (8p - 9)t^3 \right) E_t u \\
&\quad + 2(p-2)(2p-1)((6p^2 - 5p + 6)t + 10p^3 - 54p^2 + 10p + 6)H_3^{[1]} E_t u, \tag{3.20}
\end{aligned}$$

$$\begin{aligned}
0 &= ((t+1-p)^2 - 2p\omega_1^{[1]}) \\
&\quad \times \left((p-2)(p-2\omega_1^{[1]})(72p^3 + 44p^2 - 235p + 120) \right. \\
&\quad \quad - (284p^4 + (-40\omega_1^{[1]} - 3)p^3 + (6\omega_1^{[1]} - 1120)p^2 + (88\omega_1^{[1]} + 754)p - 48\omega_1^{[1]} + 60)t \\
&\quad \quad \left. + (272p^3 + 410p^2 - 363p - 270)t^2 \right. \\
&\quad \quad \left. + (16p^2 + 61p - 102)t^3 \right) E_t u \\
&\quad + 2(p-2)(2p-1)((14p^3 + 21p^2 - 74p + 60)t + 24p^4 \\
&\quad \quad - 90p^3 - 221p^2 + 220p + 60)H_3^{[1]} E_t u, \tag{3.21}
\end{aligned}$$

$$\begin{aligned}
0 &= ((t+1-p)^2 - 2p\omega_1^{[1]}) \\
&\quad \times \left(-3(p-2)(p-2\omega_1^{[1]})(616p^5 - 1262p^4 + (-40\omega_1^{[1]} + 1958)p^3 \right. \\
&\quad \quad \left. + (10\omega_1^{[1]} - 2397)p^2 + 1232p - 150) \right. \\
&\quad \quad + 3(2464p^6 + (-704\omega_1^{[1]} - 6537)p^5 + (1618\omega_1^{[1]} + 9700)p^4 \\
&\quad \quad \left. + (-2452\omega_1^{[1]} - 12590)p^3 + (2848\omega_1^{[1]} + 7496)p^2 \right. \\
&\quad \quad \left. + (-1388\omega_1^{[1]} - 623)p + 150\omega_1^{[1]} - 90)t \right. \\
&\quad \quad - 3(2464p^5 - 2793p^4 + (40\omega_1^{[1]} + 4907)p^3 + (-90\omega_1^{[1]} - 3483)p^2 \\
&\quad \quad \left. + (20\omega_1^{[1]} - 1252)p + 385)t^2 \right. \\
&\quad \quad \left. - 15(p-2)(2p+19)(4p-1)t^3 - 15(p-2)(4p-1)t^4 \right) E_t u \\
&\quad - (p-2)(2p-1)((1056p^5 - 582p^4 + 1428p^3 - 1932p^2 + 1992p - 450)t \\
&\quad \quad + 792p^6 + (176\omega_1^{[1]} - 6224)p^5 + (-392\omega_1^{[1]} + 8666)p^4 \\
&\quad \quad + (628\omega_1^{[1]} - 13729)p^3 + (-352\omega_1^{[1]} + 11014)p^2 \\
&\quad \quad \left. + (192\omega_1^{[1]} + 120)p - 450)H_3^{[1]} E_t u. \tag{3.22}
\end{aligned}$$

Deleting the terms including $E_t H_3^{[1]} u$ and $((t+1-p)^2 - 2p\omega_1^{[1]})^2 E_t u$ from [24, (3.68)–(3.71)], we have

$$\begin{aligned} 0 &= p^2(2p-9)(2p-1)(4p^2-12p+15)(10p^2-4p+3) \\ &\quad \times (44p^4-13p^3+62p^2-48p+18) \\ &\quad \times t(t-p+2)(2t-p+1)(2t-p+2)(2t-p+3) \\ &\quad \times ((t+1-p)^2 - 2p\omega_1^{[1]}) E_t u. \end{aligned} \quad (3.23)$$

We set

$$\tilde{u} := ((t+1-p)^2 - 2p\omega_1^{[1]}) E_t u = E_t((t+1)^2 - 2p\omega_1^{[1]}) u \quad (3.24)$$

and assume $\tilde{u} \neq 0$. By (3.23), we have $t = 0, p/2 - 1$, or $p - 2$.

- (1) If $t = p - 2$, then deleting the terms including $E_t H_3^{[1]} u$ from [24, (3.68) and (3.69)], we have $(\omega_1^{[1]} - 1)\tilde{u} = 0$. Since $\langle \lambda, \alpha \rangle^2 / (2p) + p/2 - t - 1 = 1$ by (3.4), we have $\langle \lambda, \alpha \rangle^2 = p^2$ and hence $t = p - 2 < \langle \lambda, \alpha \rangle - 1$ by (3.13), which contradicts (3.14).
- (2) If $t = p/2 - 1$, then deleting the terms including $H_3^{[1]} E_t u$ from (3.19)–(3.22), we have

$$0 = (16\omega_1^{[1]} - 1)(16\omega_1^{[1]} - 9)\tilde{u} \quad (3.25)$$

and hence there exists a non-zero $w \in \langle \omega_1^{[1]} \rangle u$ such that $(\omega_1^{[1]} w, H_3^{[1]} w) = ((1/16)w, (-1/128)w)$ or $((9/16)w, (15/128)w)$ by (3.19). This contradicts to [2, Proposition 4.3] which shows that the eigenvalues of $H_3^{[1]}$ on $V_{\lambda+L}$ for any $\lambda \in L^\perp$ are non-negative integers.

- (3) If $t = 0$, then deleting the terms including $H_3^{[1]} E_t u$ from (3.19)–(3.22), we have $\omega_1^{[1]}\tilde{u} = (p/2)\tilde{u}$. Since $\langle \lambda, \alpha \rangle / (2p) + p/2 - t - 1 = p/2$ by (3.4), we have $\langle \lambda, \alpha \rangle = 2p$, which contradicts (3.13) and (3.14).

Thus, $\tilde{u} = (\omega_1^{[1]} - (t+1-p)^2 / (2p)) E_t u = E_t(\omega_1^{[1]} - (t+1)^2 / (2p)) u = 0$ and hence $(t+1)^2 = \langle \lambda, \alpha \rangle^2$ by (3.4), which leads

$$t = \langle \lambda, \alpha \rangle - 1 \quad (3.26)$$

by (3.14). Since $(\omega_1^{[1]} - (t+1)^2 / (2p)) u \in W_{\zeta, \mathbf{0}, \mathbf{0}} = \mathbb{C}w$, it follows from (2.18) and (2.19) that $(\omega_1^{[1]} - (t+1)^2 / (2p)) u = 0$. By [24, (3.68)], $E_t H_3^{[1]} u = 0$. Since

$H_3^{[1]}u \in W_{\zeta, \mathbf{0}, \mathbf{0}} = \mathbb{C}w$, it follows from (2.18) and (2.19) that $H_3^{[1]}u = 0$. We conclude that u is a simultaneous eigenvector for $\{\omega_1^{[i]}, H_3^{[i]}\}_{i=1}^d$. The same argument as in the proof (1) of [24, Lemma 5.5] shows that $\epsilon(S_{ij}(1, 1), u) \leq 1$ for any pair of distinct elements $i, j \in \{1, \dots, d\}$ and hence $u \in \Omega_{M(1)^+}(N_{\zeta, \mathbf{0}, \mathbf{0}})$.

Since $M(1)^-$ and $M(1)(\theta)^-$ are not $M(1)^+$ -submodules of N , $A^u u = A^t u = 0$ and hence $A(M(1)^+) \cdot u = \langle \Lambda_{ij} \mid i, j = 1, \dots, d \text{ with } i \neq j \rangle u$. By [15, (6.1.15)], $\Lambda_{ij}^2 u = 4\omega_1^{[i]}\omega_1^{[j]}u = \langle h^{[i]}, \lambda \rangle^2 \langle h^{[j]}, \lambda \rangle^2 u \neq 0$. Since $(\Lambda_{ij} - \langle h^{[i]}, \lambda \rangle \langle h^{[j]}, \lambda \rangle)^2 u = 0$, we have $\Lambda_{ij} u = \langle h^{[i]}, \lambda \rangle \langle h^{[j]}, \lambda \rangle u$ and hence $\mathbb{C}u \cong M(1, \lambda)(0)$ as $A(M(1)^+)$ -modules. By [24, Corollary 5.9], $M(1)^+ \cdot u \cong M(1, \lambda)$ as $M(1)^+$ -modules. Now, the result follows from Lemma 3.1. \square

Lemma 3.7. $\text{Ext}_{V_L^+}^1(V_L^+, V_L^-) = 0$.

Proof. Let $0 \rightarrow V_L^- \rightarrow N \xrightarrow{\pi} V_L^+ \rightarrow 0$ be an exact sequence of weak V_L^+ -modules. Since $(V_L^+)_{\mathbf{0}, \mathbf{0}, \mathbf{0}} = \mathbb{C}\mathbf{1}$ and $(V_L^-)_{\mathbf{0}, \mathbf{0}, \mathbf{0}} = 0$, we can take a non-zero $u \in N_{\mathbf{0}, \mathbf{0}, \mathbf{0}}$ such that $\pi(u) = \mathbf{1}$ and $\omega_1^{[i]}u = H_3^{[i]}u = H_5^{(6), [i]}u = 0$ for all $i = 1, \dots, d$. The same argument as in the proof of Lemma 3.6 (cf. (3.9)–(3.12)) shows that $\omega_{2+j}^{[i]}u = H_{4+j}^{[i]}u = 0$ for all $i = 1, \dots, d$ and $j \geq 0$. Since $P_6^{(8), H}u = 0$, we have

$$\omega_0^{[i]}u = 3H_2^{[i]}u. \quad (3.27)$$

For $i, j = 1, \dots, d$,

$$\omega_1^{[j]}\omega_0^{[i]}u = -[\omega_0^{[i]}, \omega_1^{[j]}]u = \delta_{ij}\omega_0^{[i]}u \quad (3.28)$$

and

$$H_3^{[j]}\omega_0^{[i]}u = -[\omega_0^{[i]}, H_3^{[j]}]u = \delta_{ij}3H_2^{[i]}u = \delta_{ij}\omega_0^{[i]}u. \quad (3.29)$$

It follows from [2, Proposition 4.3] that for any $i = 1, \dots, d$

$$\begin{aligned} \{v \in V_L^+ \mid \omega_1^{[j]}v = H_3^{[j]}v = \delta_{ij}v \text{ for all } j = 1, \dots, d\} &= \{0\} \text{ and} \\ \{v \in V_L^- \mid \omega_1^{[j]}v = H_3^{[j]}v = \delta_{ij}v \text{ for all } j = 1, \dots, d\} &\subset \mathbb{C}h^{[i]}(-1)\mathbf{1} \end{aligned} \quad (3.30)$$

and hence that $\omega_0^{[i]}u \in \mathbb{C}h^{[i]}(-1)\mathbf{1} \subset V_L^-$. We take $\beta \in \mathfrak{h}$ so that $\beta(-1)\mathbf{1} = \omega_0 u = \sum_{i=1}^d \omega_0^{[i]}u \in V_L^-$. The following argument is a slight modification of a

part of the proof of [1, Proposition 4.6]. Since for all $\alpha \in L$ and $n \in \mathbb{Z}$,

$$\begin{aligned} E(\alpha)_n \beta(-1)\mathbf{1} &= -[\beta(-1), E(\alpha)_n]\mathbf{1} + \beta(-1)E(\alpha)_n\mathbf{1} \\ &= -(\beta(0)E(\alpha))_{n-1}\mathbf{1} + \beta(-1)E(\alpha)_n\mathbf{1} \\ &= -\langle \alpha, \beta \rangle (e^\alpha - \theta(e^\alpha))_{n-1}\mathbf{1} + \beta(-1)E(\alpha)_n\mathbf{1}, \end{aligned} \quad (3.31)$$

we have $E(\alpha)_n \omega_0 u = 0$ for all $n \geq 1$. Since $E(\alpha)_n = (-1/(n+1))[\omega_0, E(\alpha)_{n+1}]$ for all $n \geq 0$, we have $\epsilon(E(\alpha), u) \leq 0$. Moreover, since $E(\alpha)_0 u = -[\omega_0, E(\alpha)_1]u = 0$, we have $E(\alpha)_0 \omega_0 u = \omega_0 E(\alpha)_0 u = 0$. Since L is non-degenerate, it follows from (3.31) with $n = 0$ that $\omega_0 u = 0$ and hence $V_L^+ \cdot u \cong V_L^+$. Now, the result follows from Lemma 3.1. \square

Lemma 3.8. $\text{Ext}_{V_L^+}^1(V_L^-, V_L^+) = 0$.

Proof. Let $0 \rightarrow V_L^+ \rightarrow N \xrightarrow{\pi} V_L^- \rightarrow 0$ be an exact sequence of weak V_L^+ -modules. By Lemma 3.4, we can take a sublattice $\bigoplus_{i=1}^d \mathbb{Z}\alpha^{[i]}$ of L so that $\langle \alpha^{[i]}, \alpha^{[i]} \rangle \notin \{0, 2\}$ for all $i = 1, \dots, d$ and $\langle \alpha^{[i]}, \alpha^{[j]} \rangle = 0$ for all pairs of distinct elements $i, j \in \{1, \dots, d\}$. We define $p_i := \langle \alpha^{[i]}, \alpha^{[i]} \rangle$ and $h^{[i]} := \alpha^{[i]}/\sqrt{p_i}$. For $i = 1, \dots, d$ and $\zeta^{(i)} = (\delta_{ij})_{j=1}^d$, let $u^{[i]} \in N_{\zeta^{(i)}, \zeta^{(i)}, \zeta^{(i)}}$ such that

$$\pi(u^{[i]}) = \alpha^{[i]}(-1)\mathbf{1} \in V_L^-. \quad (3.32)$$

Since $(V_L^+)_{\zeta^{(i)}, \zeta^{(i)}, \zeta^{(i)}} = \mathbf{0}$, we have

$$(\omega_1^{[j]} - \delta_{ij})u^{[i]} = (H_3^{[j]} - \delta_{ij})u^{[i]} = (H_5^{(6), [j]} - \delta_{ij})u^{[i]} = 0 \quad (3.33)$$

for $j = 1, \dots, d$. We define

$$U = \sum_{i=1}^d \mathbb{C}u^{[i]}. \quad (3.34)$$

Since $\text{Ext}_{M(1)^+}^1(M(1)^-, M(1, \alpha)) = 0$ for all $\alpha \in \mathfrak{h} \setminus \{0\}$ by [24, Proposition 5.8] and $M(1)_1^+ = 0$, the vector space U is an $A(M(1)^+)$ -module which is isomorphic to $M(1)_1^- = M(1)^-(0)$. For any pair of distinct element $i, j \in \{1, \dots, d\}$ and $l, m \in \mathbb{Z}_{>0}$, we set

$$\tilde{S}_{ij}(l, m) = \alpha^{[i]}(-l)\alpha^{[j]}(-m), \quad (3.35)$$

which is a non-zero scalar multiple of $S_{ij}(l, m)$. In order to show $U \subset \Omega_{M(1)^+}(N)$, it is convenient to use $\tilde{S}_{ij}(l, m)$ instead of $S_{ij}(l, m)$. We fix

$i \in \{1, \dots, d\}$. For $k = 1, \dots, d$, since $\omega_2^{[k]} u^{[i]} \in \mathbb{C}\mathbf{1}$, we have $H_3^{[k]} \omega_2^{[k]} u^{[i]} = 0$. For $j \in \{1, \dots, d\}$ with $j \neq i$, by (3.33) and (2.25) with $m = 1$,

$$\begin{aligned} -\omega_2^{[i]} u^{[i]} &= [H_3^{[i]}, \omega_2^{[i]}] u^{[i]} = -3H_4^{[i]} u^{[i]} - \omega_2^{[i]} u^{[i]}, \\ 0 &= [H_3^{[j]}, \omega_2^{[j]}] u^{[i]} = -3H_4^{[j]} u^{[i]} - \omega_2^{[j]} u^{[i]} \end{aligned} \quad (3.36)$$

and hence $H_4^{[i]} u^{[i]} = 0$ and $3H_4^{[j]} u^{[i]} = -\omega_2^{[j]} u^{[i]}$. By Lemma 2.2, we have $\epsilon_M(\omega^{[i]}, u^{[i]}) = 1$, $\epsilon_M(E(\alpha^{[i]}), u^{[i]}) = 0$ and $\epsilon_M(\omega^{[j]}, u^{[i]}) \leq 0$, $\epsilon_M(E(\alpha^{[j]}), u^{[i]}) = -1$ for $j \neq i$.

We shall show that $\epsilon_M(\tilde{S}_{ij}(1, 1), u) \leq 1$ for all non-zero $u \in U$ and all pairs of distinct elements $i, j \in \{1, \dots, d\}$. We may assume $(i, j) = (2, 1)$. We take a pair of non-zero integers x and y so that $p_1 x^2 + p_2 y^2 \notin \{0, \pm 1, \pm 2\}$ and define the sequence $\tilde{\alpha}^{[1]}, \tilde{\alpha}^{[2]}, \dots, \tilde{\alpha}^{[d]} \in L$ by

$$\begin{aligned} \tilde{\alpha}^{[1]} &= x\alpha^{[1]} + y\alpha^{[2]}, \\ \tilde{\alpha}^{[2]} &= -yp_2\alpha^{[1]} + xp_1\alpha^{[2]}, \text{ and} \\ \tilde{\alpha}^{[i]} &= \alpha^{[i]}, \quad i = 3, \dots, d. \end{aligned} \quad (3.37)$$

For $i = 1, \dots, d$ and $j = 3, \dots, d$, we also define

$$\begin{aligned} \tilde{h}^{[i]} &= \frac{\tilde{\alpha}^{[i]}}{\sqrt{\langle \tilde{\alpha}^{[i]}, \tilde{\alpha}^{[i]} \rangle}}, \\ \tilde{\omega}^{[i]} &= \frac{1}{2} \tilde{h}^{[i]} (-1)^2 \mathbf{1}, \\ \tilde{H}^{[i]} &= \frac{1}{3} \tilde{h}^{[i]} (-3) h^{[i]} (-1) \mathbf{1} - \frac{1}{3} \tilde{h}^{[i]} (-2)^2 \mathbf{1}, \\ \tilde{u}^{[1]} &= xu^{[1]} + yu^{[2]}, \\ \tilde{u}^{[2]} &= -yp_2u^{[1]} + xp_1u^{[2]}, \\ \tilde{u}^{[j]} &= u^{[j]}. \end{aligned} \quad (3.38)$$

Then, $\langle \tilde{\alpha}^{[i]}, \tilde{\alpha}^{[i]} \rangle \notin \{0, 2\}$ for all $i = 1, \dots, d$, $\langle \tilde{\alpha}^{[i]}, \tilde{\alpha}^{[j]} \rangle = 0$ for all pairs of distinct element $i, j \in \{1, \dots, d\}$, $\tilde{\omega}_1^{[i]} \tilde{u}^{[j]} = \tilde{H}_3^{[i]} \tilde{u}^{[j]} = \delta_{ij} \tilde{u}^{[j]}$, and $U = \sum_{i=1}^d \mathbb{C} \tilde{u}^{[i]}$. The same argument as above shows that $\epsilon(\tilde{\omega}^{[i]}, u) \leq 1$ for all non-zero $u \in U$ and $i = 1, \dots, d$. For a non-zero $u \in U$, since $\epsilon(\tilde{\omega}^{[1]}, u), \epsilon(\omega^{[k]}, u) \leq 1$ for all $k = 1, \dots, d$, and

$$\tilde{S}_{21}(1, 1) = \frac{1}{xy} (\langle \tilde{\alpha}^{[1]}, \tilde{\alpha}^{[1]} \rangle \tilde{\omega}^{[1]} - p_1 x^2 \omega^{[1]} - p_2 y^2 \omega^{[2]}), \quad (3.39)$$

we have $\epsilon(\widetilde{S}_{21}(1,1), u) \leq 1$ and hence $U \subset \Omega_{M(1)^+}(N)$. Now, the result follows from Lemma 3.1. \square

(*Proof of Theorem 1.1*). Since [24, Lemma 7.3] shows that every weak V_L^+ -module has an irreducible weak V_L^+ -module, it is enough to show that $\text{Ext}_{V_L^+}^1(M, W) = 0$ for any pair of irreducible weak V_L^+ -modules M and W (cf. the last half of the proof of [2, Theorem 2.8]). The result follows from Corollary 3.3, Lemma 3.6, Lemma 3.7, and Lemma 3.8. \square

References

- [1] T. Abe, *The charge conjugation orbifold $V_{\mathbb{Z}\alpha}^+$ is rational when $\langle \alpha, \alpha \rangle / 2$ is prime*, Int. Math. Res. Not. **2002** (2002), 647–665.
- [2] T. Abe, *Rationality of the vertex operator algebra V_L^+ for a positive definite even lattice L* , Math. Z. **249** (2005), 455–484.
- [3] T. Abe, G. Buhl and C. Dong, *Rationality, regularity, and C_2 -cofiniteness*, Trans. Amer. Math. Soc. **356** (2004), 3391–3402.
- [4] T. Arakawa, *Rationality of W -algebras: principal nilpotent cases*, Ann. of Math. (2) **182** (2015), 565–604.
- [5] R. Borcherds, *Vertex algebras, Kac-Moody algebras, and the Monster*, Proc. Nat. Acad. Sci. U.S.A. **83** (1986), 3068–3071.
- [6] R. Dijkgraaf, C. Vafa, E. Verlinde, and H. Verlinde, *The operator algebra of orbifold models*, Comm. Math. Phys. **123** (1989), 485–526.
- [7] C. Dong, *Vertex algebras associated with even lattices*, J. Algebra **160** (1993), 245–265.
- [8] C. Dong, *Representations of the moonshine module vertex operator algebra*, Mathematical aspects of conformal and topological field theories and quantum groups (South Hadley, MA, 1992), Contemp. Math. **175**, 27–36 (1994).
- [9] C. Dong, *Twisted modules for vertex algebras associated with even lattices*, J. Algebra **165** (1994), 91–112.

- [10] C. Dong, C. Jiang and X. Lin, *Rationality of vertex operator algebra V_L^+ : higher rank*, Proc. Lond. Math. Soc. **104** (2012), 799–826.
- [11] C. Dong, C. H. Lam, K. Tanabe, H. Yamada, and K. Yokoyama, *\mathbb{Z}_3 symmetry and W_3 algebra in lattice vertex operator algebras*, Pacific J. Math. **215** (2004), 245–296.
- [12] C. Dong and J. Lepowsky, *Generalized vertex algebras and relative vertex operators*, Progress in Mathematics **112**, Birkhauser Boston, Inc., Boston, MA, 1993.
- [13] C. Dong, H. S. Li and G. Mason, *Regularity of rational vertex operator algebras*, Adv. Math. **132** (1997), 148–166.
- [14] C. Dong and K. Nagatomo, *Classification of irreducible modules for the vertex operator algebra $M(1)^+$* , J. Algebra **216** (1999), 384–404.
- [15] C. Dong and K. Nagatomo, *Classification of irreducible modules for the vertex operator algebra $M(1)^+$:II. higher rank*, J. Algebra **240** (2001), 289–325.
- [16] I. B. Frenkel, J. Lepowsky and A. Meurman, *Vertex operator algebras and the monster*, Pure and Applied Math. **134**, Academic Press, 1988.
- [17] I. B. Frenkel and Y. Zhu, *Vertex operator algebras associated to representations of affine and virasoro algebras*, Duke Math. J. **66** (1992), 123–168.
- [18] V. Kac, *Vertex algebras for beginners, second edition*, University Lecture Series **10**, American Mathematical Society, 1998.
- [19] J. Lepowsky and H. S. Li, *Introduction to vertex operator algebras and their representations*, Progress in Mathematics **227**, Birkhauser Boston, Inc., Boston, MA, 2004.
- [20] J. Levitzki, *On automorphisms of certain rings*, Ann. of Math. (2) **36** (1935), 984–992.
- [21] H. S. Li, *Local systems of vertex operators, vertex superalgebras and modules*, J. Pure Appl. Algebra **109** (1996), 143–195.

- [22] S. Montgomery, *Fixed rings of finite automorphism groups of associative rings*, Lecture Notes in Mathematics **818**, Springer, Berlin, 1980.
- [23] Risa/Asir, <http://www.math.kobe-u.ac.jp/Asir/asir.html>.
- [24] K. Tanabe, *The irreducible weak modules for the fixed point subalgebra of the vertex algebra associated to a non-degenerate even lattice by an automorphism of order 2*, <https://arxiv.org/abs/1910.07126>.
- [25] K. Tanabe, *Simple weak modules for the fixed point subalgebra of the heisenberg vertex operator algebra of rank 1 by an automorphism of order 2 and whittaker vectors*, Proc. Amer. Math. Soc. **145** (2017), 4127–4140.
- [26] K. Tanabe, *Simple weak modules for some subalgebras of the Heisenberg vertex algebra and Whittaker vectors*, Algebr. Represent. Theory **23** (2020), 53–66.
- [27] K. Tanabe and H. Yamada, *The fixed point subalgebra of a lattice vertex operator algebra by an automorphism of order three*, Pacific J. Math. **230** (2007), 469–510.
- [28] K. Tanabe and H. Yamada, *Fixed point subalgebras of lattice vertex operator algebras by an automorphism of order three*, J. Math. Soc. Japan **65** (2013), 1169–1242.
- [29] W. Wang, *Rationality of virasoro vertex operator algebras*, International Mathematics Research Notices **1993** (1993), 197–211.
- [30] G. Yamskulna, *Rationality of the vertex algebra V_L^+ when L is a non-degenerate even lattice of arbitrary rank*, J. Algebra **321** (2009), 1005–1015.
- [31] Y. Zhu, *Modular invariance of characters of vertex operator algebras*, J. Amer. Math. Soc. **9** (1996), 237–302.