

Portfolio optimization with mixture vector autoregressive models

Davide Ravagli

davide.ravagli@manchester.ac.uk

Georgi N. Boshnakov

georgi.boshnakov@manchester.ac.uk

Department of Mathematics
The University of Manchester
Manchester, United Kingdom

Abstract

Obtaining reliable estimates of conditional covariance matrices is an important task of heteroskedastic multivariate time series. In portfolio optimization and financial risk management, it is crucial to provide measures of uncertainty and risk as accurately as possible. We propose using mixture vector autoregressive (MVAR) models for portfolio optimization. Combining a mixture of distributions that depend on the recent history of the process, MVAR models can accommodate asymmetry, multimodality, heteroskedasticity and cross-correlation in multivariate time series data. For mixtures of Normal components, we exploit a property of the multivariate Normal distribution to obtain explicit formulas of conditional predictive distributions of returns on a portfolio of assets. After showing how the method works, we perform a comparison with other relevant multivariate time series models on real stock return data.

Keywords: Forecasting; Heteroskedasticity; Mixture vector autoregressive model; MVAR model; Portfolio.

1 Introduction

Financial and econometric data often presents the feature of heteroskedasticity. For multivariate time series, this implies that the covariance matrix of an observation at a given time point depends upon the recent history of the process. This may be due changes in the volatility of a single series, as well as in the cross-correlations between any two series of interest. As a result, one cannot trust sample estimates of the (unconditional) covariance matrix or linear time series models to build reliable predictions about the future. Therefore, obtaining reliable estimates of covariance matrices remains an important challenge in portfolio optimization and financial risk management which use, for instance, modern portfolio theory (Markowitz 1952). Bollerslev et al. (1988) and Engle and Kroner (1995) pioneered in the attempt to model conditional covariance matrices of predictors for multivariate time series with multivariate GARCH models, using different parametrizations known respectively as VEC and BEKK. Engle (2002) extended the idea of multivariate GARCH to the so

called Dynamic Conditional Correlation models, in which each element of the time-dependent covariance matrix of the data is modelled to follow a GARCH process. Such models have computational advantages over multivariate GARCH models in that the number of parameters to be estimated in the correlation process is independent of the number of series to be correlated, by use of common parameters across all correlations to be estimated.

Since then, much work has been done to develop multivariate GARCH models for portfolio optimization. In particular, many attempts have been made in combining GARCH and factor models, with the aim of dimensionality reduction. These models rely on the assumption that financial returns are described by a small number of underlying common variables, or factors, which can be used to model the data more parsimoniously. Although all equal in concept, different approaches used different assumptions on such factors, so that different techniques are used to derive them. For instance, Alexander (2000) uses a principal components analysis in which factors are assumed to follow independent GARCH processes, whereas Van der Weide (2002) considers the case in which factors are not orthogonal. Finally, Santos and Moura (2014) introduced the dynamic factor GARCH model with time-varying factor loadings.

We propose using a mixture vector autoregressive (MVAR) model (Fong et al. 2007) for portfolio optimization. MVAR models are the multivariate extension of the mixture autoregressive model by Wong and Li (2000). Combining predictive distributions which depend on the recent history of the process, MVAR models can accommodate asymmetry, multimodality, heteroskedasticity and cross-correlation in multivariate time series data. Theoretical properties of MVAR were explored for the case of a multivariate Gaussian mixture in Fong et al. (2007) and Kalliovirta et al. (2016).

Using the Gaussian MVAR model assumption, we are able to fully specify conditional predictive distributions for future observations. We will show how it is possible to combine modern portfolio theory (Markowitz 1952) and the assumption of Gaussian mixture vector autoregressive model for portfolio optimization. Under this model assumption, we will also estimate the risk associated with the forecast. Finally, we will compare the performance of our method with that of the dynamic conditional correlation model by Engle (2002) and the vector autoregressive model (VAR).

2 The mixture vector autoregressive model

Mixture vector autoregressive models or MVAR (Fong et al. 2007) are a multivariate extension of Mixture Autoregressive Models (Wong and Li 2000).

The MVAR model with g Gaussian components, and an m dimensional vector Y_t is defined as

$$F(Y_t | \mathcal{F}_{t-1}) = \sum_{k=1}^g \pi_k \Phi \left(\Omega_k^{-1/2} \left(Y_t - \Theta_{k0} - \sum_{i=1}^{p_k} \Theta_{ki} Y_{t-i} \right) \right) \quad (1)$$

where

- Y_t is a $m \times 1$ data vector at time t .
- $\boldsymbol{\pi} = (\pi_1, \dots, \pi_g)$ are the mixing weights.

- Ω_k is the covariance matrix of component k .
- p_k , $k = 1, \dots, g$ is the autoregressive order of component k . We denote $p = \max(p_k)$.
- Θ_{k0} is a $m \times 1$ intercept vector for component k , and $\Theta_{k1}, \dots, \Theta_{kp_k}$ are $m \times m$ matrices of autoregressive parameters. If $p_k < p$, then $\Theta_{kl} = 0_m$ for $p_k < l \leq p$, where 0_m is the zero-matrix of size $m \times m$.
- $\Phi(\cdot)$ is the CDF of the standard multivariate normal distribution, and $\phi(\cdot)$ is the corresponding pdf.
- Assuming start at $t = 1$, (1) holds for $t > p$.

Regularity conditions and parameter estimation by EM algorithm are discussed in Fong et al. (2007) and Kalliovirta et al. (2016).

MVAR may be seen as an alternative to multivariate GARCH when the data presents heteroskedasticity and time-dependent cross correlations, while also accounting for possible multimodality and asymmetry in the distribution.

Suppose that a m -variate time series $\{Y_t\}$ of length n follows a MVAR process. Let $\mathbf{Z} = (\mathbf{Z}_1, \dots, \mathbf{Z}_n)$ be an unobserved allocation random variable, where \mathbf{Z}_t is a g -dimensional vector with component k equal to 1 if Y_t comes from the k^{th} component, and 0 otherwise.

Following notation from Fong et al. (2007), let $\tilde{\Theta}_k = [\Theta_{k0}, \Theta_{k1}, \dots, \Theta_{kp_k}]$ and $X_{tk} = (1, Y_{t-1}^T, \dots, Y_{t-p_k}^T)^T$. In addition, let ϑ denote the complete set of parameters. Parameter estimates are then obtained by EM-algorithm with the following steps:

• **E-step**

$$\tau_{tk} = \mathbb{E}[Z_{tk} | Y_t, \vartheta] = \frac{\pi_k \phi \left(\Omega_k^{-1/2} \left(Y_t - \Theta_{k0} - \sum_{i=1}^{p_k} \Theta_{ki} Y_{t-i} \right) \right)}{\sum_{l=1}^g \pi_l \phi \left(\Omega_l^{-1/2} \left(Y_t - \Theta_{l0} - \sum_{i=1}^{p_l} \Theta_{li} Y_{t-i} \right) \right)} \quad (2)$$

• **M- step**

$$\begin{aligned} \hat{\pi}_k &= \frac{1}{n-p} \sum_{t=p+1}^n \tau_{tk} \\ \hat{\Theta}_k &= \left(\sum_{t=p+1}^n \tau_{tk} X_{tk} X_{tk}^T \right)^{-1} \left(\sum_{t=p+1}^n \tau_{tk} X_{tk} Y_t^T \right) \\ \hat{\Omega}_k &= \frac{\sum_{t=p+1}^n \tau_{tk} e_{tk} e_{tk}^T}{\sum_{t=p+1}^n \tau_{tk}} \end{aligned} \quad (3)$$

where $e_{tk} = Y_t - \Theta_{k0} - \sum_{i=1}^{p_k} \Theta_{ki} Y_{t-i}$.

E-step and M-step are repeated recursively until convergence to maximum likelihood estimates of the parameters.

First and second order stationarity conditions are discussed by Saikkonen (2007), see also Boshnakov (2011) for the univariate case. Let

$$A_k = \begin{bmatrix} \Theta_{k1} & \Theta_{k2} & \dots & \Theta_{kp-1} & \Theta_{kp} \\ I_m & 0_m & \dots & 0_m & 0_m \\ 0_m & I_m & \dots & 0_m & 0_m \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0_m & 0_m & \dots & I_m & 0_m \end{bmatrix}, \quad k = 1, \dots, g \quad (4)$$

where I_m and 0_m are respectively the identity matrix and the zero matrix of size $m \times m$. A necessary and sufficient condition for the MVAR model to be stationary is that the eigenvalues of $\sum_{k=1}^g \pi_k A_k \otimes A_k$ are smaller than 1 in modulus. A MVAR model that satisfies this condition is said to be *Stable*. In practice, to assess stability of the fitted model parameters are replaced by their estimates.

2.1 Prediction with mixture vector autoregressive models

By model assumption, the one step ahead conditional predictive distribution at time t is fully specified, and it is that of (1) where, for notational convenience, we replace t with $t + 1$, i.e.

$$F(Y_{t+1} | \mathcal{F}_t) = \sum_{k=1}^g \pi_k \Phi \left(\Omega_k^{-1/2} \left(Y_{t+1} - \Theta_{k0} - \sum_{i=1}^{p_k} \Theta_{ki} Y_{t+1-i} \right) \right)$$

Hence, the conditional distribution of the one step ahead predictor is a mixture of g Gaussian components and it depends on previous values of the process. In particular, the conditional covariance matrix depends on previous values of the process, a defining property of heteroskedasticity. To obtain the conditional mean and the covariance matrix let $\mu_{t+1,k} = \Theta_{k0} + \sum_{i=1}^{p_k} \Theta_{ki} Y_{t+1-i}$, for $k = 1, \dots, g$. Then

$$\begin{aligned} \mathbb{E}[Y_{t+1} | \mathcal{F}_t] &= \sum_{k=1}^g \pi_k \mu_{t+1,k} = \mu_{t+1} \\ \text{Cov}(Y_{t+1} | \mathcal{F}_t) &= \sum_{k=1}^g \pi_k \Omega_k + \sum_{k=1}^g \pi_k (\mu_{t+1,k} - \mu_{t+1})(\mu_{t+1,k} - \mu_{t+1})^T \\ &= \sum_{k=1}^g \pi_k \Omega_k + \sum_{k=1}^g \pi_k \mu_{t+1,k} \mu_{t+1,k}^T - \mu_{t+1} \mu_{t+1}^T \end{aligned} \quad (5)$$

Using a method analogous to that of Boshnakov (2009), we can derive the conditional distribution for the two steps ahead predictor as a mixture of g^2 Gaussian components:

$$F(Y_{t+2} | \mathcal{F}_t) = \sum_{k=1}^g \sum_{l=1}^g \pi_k \pi_l \Phi \left(\Psi_{kl}^{-1/2} (Y_{t+2} - \mu_{kl}) \right) \quad (6)$$

where, for each $k, l = 1, \dots, g$,

$$\begin{aligned}\mu_{kl} &= \Theta_{k0} + \Theta_{k1}\Theta_{l0} + \sum_{i=1}^{p-1} (\Theta_{k,i+1} + \Theta_{k1}\Theta_{li}) Y_{t-1-i} + \Theta_{k1}\Theta_{lp} Y_{t-1-p} \\ \Psi_{kl} &= \Omega_k + \Theta_{k1}\Omega_l\Theta_{k1}^T\end{aligned}$$

Note that, in general, $\mu_{kl} \neq \mu_{lk}$ and $\Psi_{kl} \neq \Psi_{lk}$. Expectation and covariance matrix of this predictor are:

$$\begin{aligned}\mathbb{E}[Y_{t+2} | \mathcal{F}_t] &= \sum_{k=1}^g \sum_{l=1}^g \pi_k \pi_l \mu_{kl} = \mu_{t+2} \\ \text{Cov}(Y_{t+2} | \mathcal{F}_t) &= \sum_{k=1}^g \pi_k \pi_l \Psi_{kl} + \sum_{k=1}^g \sum_{l=1}^g \pi_k \pi_l \mu_{kl} \mu_{kl}^T - \mu_{t+2} \mu_{t+2}^T\end{aligned}\tag{7}$$

Full derivation of (7), as well as proof of the conditional distribution of Y_{t+2} , is available in Appendix B. By recursing this procedure, we could derive a full distribution for any horizon h . However, the number of components in the mixture increases as g^h when h increases and therefore simulation methods may be preferred for approximate computation of predictive densities for larger horizons.

3 Portfolio optimization with MVAR models

Suppose that a multivariate time series $\{Y_t\}$ of asset returns is observed, and it is believed that the underlying generating process is MVAR. From Section 2.1, conditional distributions of the 1 and 2 step predictors are fully specified, and can be estimated by plugging parameter estimates into the relevant equations.

Now, let w denote the weights of a portfolio built with assets Y_t (allowing short selling), and let $R_{t+1} = w^T Y_{t+1}$ be the portfolio return at time t . Intuitively, because our model consists of a mixture of multivariate normal components, we can apply the property in (16) to conclude that the conditional distribution of R_{t+1} is also (univariate) mixture normal, with corresponding mixing weights $\boldsymbol{\pi}$ from the fitted multivariate model. By model assumption in fact, at each time $t+1$ an observation Y_{t+1} is assumed to be generated from one of g components of the mixture. Consequently, R_{t+1} is obtained by applying (16) to the selected component. Recursing this for all g components the result is itself a mixture distribution for R_{t+1} .

In terms of model parameters we write:

$$F(R_{t+1} | \mathcal{F}_t) = \sum_{k=1}^g \pi_k \Phi\left(\frac{R_{t+1} - w^T \mu_{t+1,k}}{\sqrt{w^T \Omega_k w}}\right)\tag{8}$$

Conditional mean and variance of R_{t+1} are:

$$\begin{aligned}\mathbb{E}[R_{t+1} | \mathcal{F}_t] &= \sum_{k=1}^g \pi_k (w^T \mu_{t+1,k}) = \sum_{k=1}^g \pi_k \mu_{t+1,k}^* = \mu^* \\ \text{Var}(R_{t+1} | \mathcal{F}_t) &= \sum_{k=1}^g \pi_k (w^T \Omega_k w) + \sum_{k=1}^g \pi_k (\mu_{t+1,k}^*)^2 - (\mu^*)^2\end{aligned}\tag{9}$$

Modern portfolio theory (Markowitz 1952) gives us a way to calculate weights w^* to construct the most efficient portfolio for a given return, and to calculate the efficient portfolio of assets with the minimum possible variance. A portfolio with target return μ is said to be an efficient portfolio when the variance associated with it is the lowest amongst all portfolios of the same assets having the same target return. The minimum variance portfolio is the efficient portfolio with the lowest possible variance of all efficient portfolios of the same assets. Throughout this paper we will denote efficient portfolios with the subscript EFF, and minimum variance portfolios with the subscript MVP. We now see how modern portfolio theory can be used to predict future observations assuming a MVAR model.

For the MVAR case, let $E[Y_{t+1} | \mathcal{F}_t] = \mu_{t+1}$ and $\text{Cov}(Y_{t+1} | \mathcal{F}_t) = \Omega_{t+1}$. In addition, let

$$A = \mathbb{1}\Omega_{t+1}^{-1}\mu_{t+1} \quad , \quad B = \mu_{t+1}\Omega_{t+1}^{-1}\mu_{t+1} \quad , \quad C = \mathbb{1}\Omega_{t+1}^{-1}\mathbb{1} \quad , \quad D = CB - A^2 \quad (10)$$

where $\mathbb{1}$ is a vector of 1s of the same length as μ_{t+1} .

It can be proved that optimal weights for an efficient portfolio of these assets and target return μ_{EFF} are

$$w_{\text{EFF}} = \frac{1}{D} (B\Omega_{t+1}^{-1}\mathbb{1} - A\Omega_{t+1}^{-1}\mu_{t+1} + \mu^* (C\Omega_{t+1}^{-1}\mu_{t+1} - A\Omega_{t+1}^{-1}\mathbb{1})) \quad (11)$$

and the variance of such portfolio can be calculated equivalently as $\text{Var}(R_t | \mathcal{F}_{t-1})$ (MVAR model assumption) or $w^T\Omega_t w$ (the variance of an efficient portfolio of assets) since

$$\begin{aligned} w_{\text{EFF}}^T\Omega_{t+1}w_{\text{EFF}} &= \sum_{k=1}^g \pi_k (w_{\text{EFF}}^T\Omega_k w_{\text{EFF}}) + \sum_{k=1}^g \pi_k (w_{\text{EFF}}^T\mu_{t+1,k})^2 \\ &\quad - \left[\sum_{k=1}^g \pi_k (w_{\text{EFF}}^T\mu_{t+1,k}) \right]^2 = \text{Var}(R_{t+1} | \mathcal{F}_t) \end{aligned} \quad (12)$$

In practice, $\mu_{t+1,k}$ and Ω_{t+1} are replaced with their estimates $\hat{\mu}_{t+1,k}$ and $\hat{\Omega}_{t+1}$.

Weights of the minimum variance portfolio of same assets $\{Y_t\}$, and corresponding return, are:

$$w_{\text{MVP}} = \frac{\Omega_{t+1}^{-1}\mathbb{1}}{C} \quad \mu_{\text{MVP}} = \frac{A}{C} \quad (13)$$

Conditional predictive distributions can also be calculated analytically for any $h \geq 2$. However, one must keep in mind that such predictive distribution would be a mixture of g^h components, so that simulation methods may be preferred in some cases as h increases.

Consider the case $h = 2$. The conditional predictive distribution $F(Y_{t+2} | \mathcal{F}_t)$ for the MVAR model is a mixture of g^2 Gaussian components given in (7). Similarly to the case $h = 1$, we can derive the full conditional distribution of R_{t+2} , which is again a mixture of g^2 Gaussian components:

$$F(R_{t+2} | \mathcal{F}_t) = \sum_{k,l=1}^g \pi_k \pi_l \Phi \left(\frac{R_{t+2} - w^{(2)}\mu_{kl}}{w^{(2)T}\Psi_{kl}w^{(2)}} \right) \quad (14)$$

where $w^{(2)}$ is the vector of optimal weights for this portfolio. Similarly to the case $h = 1$, one can now calculate $E[Y_{t+2} | \mathcal{F}_t] = \mu_{t+2}$ and $\text{Cov}(Y_{t+2} | \mathcal{F}_t) = \Omega_{t+2}$ and adapt (10), (11), (12) and (13) to obtain an efficient or minimum variance portfolio.

4 Simulation example

We simulate a series of size $n = 500$ of hypothetical stock returns from the 3-variate MVAR(2; 1, 1) process

$$F(Y_t | \mathcal{F}_{t-1}) = 0.75\Phi\left(\frac{Y_t - v_1}{\Omega_1}\right) + 0.25\Phi\left(\frac{Y_t - v_2}{\Omega_2}\right)$$

where

$$v_1 = \Theta_{10} + \Theta_{11}Y_{t-1} \quad v_2 = \Theta_{20} + \Theta_{21}Y_{t-1}$$

and

$$\Theta_{10} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \quad \Theta_{11} = \begin{bmatrix} 0.5 & 0 & 0.4 \\ -0.3 & 0 & 0.5 \\ -0.6 & 0.5 & -0.3 \end{bmatrix} \quad \Omega_1 = \begin{bmatrix} 1 & 0.5 & -0.40 \\ 0.5 & 2 & 0.8 \\ -0.4 & 0.8 & 4 \end{bmatrix}$$

$$\Theta_{20} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \quad \Theta_{21} = \begin{bmatrix} -0.5 & 1 & -0.4 \\ 0.3 & 0 & -0.2 \\ 0 & -0.5 & 0.5 \end{bmatrix} \quad \Omega_2 = \begin{bmatrix} 1 & 0.2 & 0 \\ 0.2 & 2 & -0.55 \\ 0 & -0.55 & 4 \end{bmatrix}$$

The three series can be seen in Figure 1, with their autocorrelation and cross-correlation plots in Figure 2. The data is very representative of what we should be looking for, in a real case scenario, to assume an underlying MVAR process. We notice in fact signs of heteroskedasticity in the series, and autocorrelations and cross-correlations significantly different from 0 at lags larger than 0. The latter is what separates MVAR from multivariate GARCH models, which assume the original series to be uncorrelated.

Parameter estimates were calculated using the EM Algorithm approach for MVAR model by Fong et al. (2007). In order to perform out of sample prediction, data from Y_1 to Y_{498} were used for estimation, with Y_{499} and Y_{500} being left out as observations 1 and 2 time points in the future:

$$\hat{\pi} = (0.7242, 0.2758)$$

$$\Theta_{10} = \begin{bmatrix} -0.0022 \\ -0.0303 \\ 0.1276 \end{bmatrix} \quad \hat{\Theta}_{11} = \begin{bmatrix} 0.4931 & -0.0339 & 0.4169 \\ -0.3156 & -0.0012 & 0.5078 \\ -0.6141 & 0.6007 & -0.3844 \end{bmatrix} \quad \hat{\Omega}_1 = \begin{bmatrix} 0.9551 & 0.4783 & -0.2776 \\ 0.4783 & 1.9123 & 0.9736 \\ -0.2776 & 0.9736 & 3.9455 \end{bmatrix}$$

$$\hat{\Theta}_{20} = \begin{bmatrix} 0.0338 \\ 0.5499 \\ -0.7580 \end{bmatrix} \quad \hat{\Theta}_{21} = \begin{bmatrix} -0.4595 & 1.0124 & -0.4004 \\ 0.3343 & -0.1423 & -0.1551 \\ -0.1273 & -0.2336 & 0.6509 \end{bmatrix} \quad \hat{\Omega}_2 = \begin{bmatrix} 0.8767 & 0.4794 & -0.3627 \\ 0.4794 & 2.9148 & -0.6576 \\ -0.3627 & -0.6576 & 9.8135 \end{bmatrix}$$

We then calculated the one step ahead conditional mean and variance based




Figure 1: Simulated time series of stock returns Asset 1 (top left), Asset 2 (top right) and Asset 3 (bottom).

on parameter estimates:

$$\begin{aligned}
 E[Y_{499} | \mathcal{F}_{498}] &= \hat{\mu}_{499} = \begin{bmatrix} -0.1750 \\ -0.9655 \\ -1.4361 \end{bmatrix} \\
 \text{Cov}(Y_{499} | \mathcal{F}_{498}) &= \hat{\Omega}_{499} = \begin{bmatrix} 1.3109 & -0.6080 & -0.0768 \\ -0.6080 & 5.3174 & -0.5642 \\ -0.0768 & -0.5642 & 5.9420 \end{bmatrix}
 \end{aligned}$$

Given $\hat{\Omega}_{499}$, we can calculate the minimum variance portfolio, which is obtained for weights $w_{\text{MVP}} = (0.6434, 0.2228, 0.1338)^T$. The corresponding expected return on this portfolio at $(t + 1) = 499$ is $\mu_{\text{MVP}} = -0.5198$, with standard deviation $\sigma_{\text{MVP}} = 0.8475$.

Suppose now that we wish to increase our return to $\mu^* = 0$, i.e. no expected loss, at the cost of a larger variance. We can calculate weights to construct an efficient portfolio of these assets as seen in Section 3. We obtain:

$$w_{\text{EFF}} = \begin{bmatrix} 1.1097 \\ 0.0781 \\ -0.1878 \end{bmatrix}$$

The interpretation of w_{EFF} is that the optimal portfolio yielding expected return of 0 is constructed by short-selling a small amount of Asset 3, and investing 110.97% and 7.81% of the initial capital (meanwhile increased by short selling) into Asset 1 and Asset 2 respectively. Notice that the target return is $w_{\text{EFF}}\mu_t = \mu_{\text{EFF}} = 0$ as desired.

We can now calculate the quantities we need for the conditional predictive




Figure 2: Autocorrelation and cross-correlation plots of the simulated time series data.

distribution of R_{499} :

$$\begin{aligned} \mu_1^* &= w^* \mu_{499,1} = 0.2642 & \sigma_1^{*2} &= 1.4968 \approx (1.2235)^2 \\ \mu_2^* &= w^* \mu_{499,2} = -0.6939 & \sigma_2^{*2} &= 1.6062 \approx (1.3025)^2 \end{aligned}$$

Therefore, the conditional distribution of $R_{499} = w^{*T} Y_{499}$ is

$$F(R_{499} | \mathcal{F}_{498}) = 0.7242 \times \Phi\left(\frac{R_{499} - 0.2642}{1.2235}\right) + 0.2758 \times \Phi\left(\frac{R_{499} - 0.6939}{1.3025}\right)$$

The standard deviation associated to this portfolio is $\sigma_{\text{EFF}} = 1.3173$ which as expected is larger than σ_{MVP} . More importantly, we can use the distribution assumption on R_{499} to estimate risk measures. Figure 3 shows the conditional distribution of R_{499} . The black dot on the left hand side, highlighted with a dashed line, is the value at risk at 95% level. It was found that the value at risk at such level is -2.2039 , with expected shortfall of -2.7912 . This means that an investor could expect a loss on this portfolio higher than 2.2039 with probability 0.05, and when this threshold is exceeded, the expected loss is of 2.7912. The observed return is also shown in Figure 3 as a red dot with red dashed line. We notice that it lies on a region of high density of the predictive distribution.

We can also estimate the conditional distribution of the 2 step ahead predictor at $t = 498$, $F(R_{500} | \mathcal{F}_{498})$ (shown in Figure 4).

The minimum variance portfolio for a two steps ahead portfolio of assets is calculated with weights $w_{\text{MVP}}^{(2)} = (0.4367, 0.2822, 0.2811)$, with an expected return $\mu_{\text{MVP}}^{(2)} = -0.3918$, with $\sigma_{\text{MVP}}^{(2)} = 1.1784$, showing the increasing uncertainty as we attempt to predict further into the future.

Once again we consider building a portfolio of assets yielding expected return




Figure 3: Conditional density of the one-step conditional distribution of R_{499} for the simulated data, with VaR at 95% (black) and observed return (red) highlighted.

$\mu^* = 0$. Optimal weights for this portfolio are

$$w_{\text{MVP}}^{(2)} = \begin{bmatrix} -0.9404 \\ 1.5193 \\ 0.4211 \end{bmatrix}$$

From Figure 4, we notice how the density is now flatter, which is sign of a larger variability. In fact, the estimated standard deviation of R_{500} is 3.5056, which is a significant increase. This also results in much larger estimated VaR = -5.0207 (in absolute value) at the same 95% level, with expected shortfall equal to -7.4505 . Once again, the observed return (red dot and dashed line) is in a high density region of the predictive distribution.

Overall, we can be satisfied with the performance of our method in predicting portfolio returns.

5 Application to the US stock market

We consider a multivariate dataset of $m = 4$ stocks on the US stock market: Dell Technologies Inc. (DELL), Microsoft Corporation (MSFT), Intel Corporation (INTC), and International Business Machine Corporation (IBM). The data were obtained from Yahoo! Finance (<https://finance.yahoo.com>). The original time series include daily Adjusted Close Prices between January 2nd 2016 and January 29th 2020 (867 observations). For each series and $t = 2, \dots, 867$, we calculated daily returns as $(\text{Price}_t - \text{Price}_{t-1})/\text{Price}_{t-1}$. The resulting series, displayed in Figure 5, includes 866 observations.

All four series in Figure 5 appear to be heteroskedastic. Their histograms also show signs of heavy tails, which was confirmed by calculation of sample excess kurtosis (all significantly larger than 0). In addition, from a preliminary analysis, it was noticed that the data present autocorrelation at least at lags 1 and 2, and cross-correlations at lags 0, 1 and 2 (see Figure 6). Therefore, it is reasonable to consider a MVAR generating process for the data.




Figure 4: Conditional density of the two-step conditional distribution of R_{500} for the simulated data, with VaR at 95% (black) and observed return (red) highlighted.

Several models were fitted. In terms of diagnostics, a $MVAR(3; 3, 2, 1)$ was chosen as best fit. Estimation was carried out on the first 864 observations, omitting the last two for out-of-sample prediction.

Given parameter estimates and the one step ahead predictive distribution at $t = 864$, we calculated weights for the minimum variance portfolio made of these assets, which yields a mean return of approximately 0.0024 (0.24%). The standard deviation associated with this portfolio is $\sigma_{MVP} = 0.0092$.

Now, assume we would like to increase our mean return to $0.007 = 0.7\%$. We can calculate optimal weights

$$w_{EFF} = (-0.5832, 0.9538, 0.1085, 0.5209)^T$$

Weights are interpreted as follows: an investor shall short-sell an amount of around 0.58 times their initial capital in DELL stocks, and reinvest the new total in the remaining three assets, with a major bet on MFST and IBM. The idea behind this is that it is believed that DELL stocks will decrease in value between the present and the nearest future, and therefore one could short-sell to make a profit. On the other hand, it is believed that the remaining three assets will increase their value in the same time span, and in particular MSFT stocks. However, the standard deviation associated with this portfolio is $\sigma_{EFF} = 0.0139$, a slight increase compared to σ_{MVP} , considering the scale of the data.

For the latter portfolio, we calculated the one step ahead conditional distribution of $R_{865} = \sum_{m=1}^4 w_m^* y_{m,865}$ using parameter estimates from the MVAR model fitting:

$$\begin{aligned} F(R_{865} | \mathcal{F}_{864}) &= 0.1316\Phi\left(\frac{R_{865} + 0.00052}{0.0266}\right) + 0.5627\Phi\left(\frac{R_{865} + 0.00178}{0.0093}\right) \\ &+ 0.3057\Phi\left(\frac{R_{865} + 0.01932}{0.0169}\right) \end{aligned}$$

The corresponding predictive density can be seen in Figure 7




Figure 5: Time series of returns of DELL (top left), MSFT (top right), INTC (bottom left) and IBM (bottom right).

Value at risk at $\alpha = 95\%$ is estimated at -0.0174 , with expected shortfall of -0.0299 . Observed return $R_{865} = -0.0062$, we can see that it lies on a region of high density, and therefore is somewhat plausible.

We can also look at building a portfolio of the same assets looking at 2 steps in the future, at $t = 866$. The minimum variance portfolio in this case yielded an expected return $\mu_{\text{MVP}}^{(2)} = -0.011$, with associated standard deviation $\sigma_{\text{MVP}}^{(2)} = 0.0101$. We used the distribution assumptions for Y_{866} , its expected value and covariance matrix to estimate optimal weights to look once again to increasing our return by bulding an efficient portfolio with same target return $\mu^* = 0.007$ as before:

$$w_{\text{EFF}}^{(2)} = (0.0402, 0.6174, 0.7554, -0.4130)^T$$

Using parameter estimates and Equation 14, the conditional distribution of R_{866} is a mixture of $3^2 = 9$ components:

$$\begin{aligned} F(R_{866} | \mathcal{F}_{864}) = & 0.0173 \Phi\left(\frac{R_{866} - 0.0170}{0.0285}\right) + 0.0741 \Phi\left(\frac{R_{866} - 0.0190}{0.0268}\right) \\ & + 0.0402 \Phi\left(\frac{R_{866} - 0.0252}{0.0276}\right) + 0.0741 \Phi\left(\frac{R_{866} - 0.0086}{0.0101}\right) \\ & + 0.3166 \Phi\left(\frac{R_{866} - 0.0069}{0.0096}\right) + 0.1720 \Phi\left(\frac{R_{866} - 0.0051}{0.0096}\right) \\ & + 0.0402 \Phi\left(\frac{R_{866} - 0.0098}{0.0206}\right) + 0.1720 \Phi\left(\frac{R_{866} - 0.0040}{0.0187}\right) \\ & + 0.0935 \Phi\left(\frac{R_{866} + 0.0077}{0.0196}\right) \end{aligned}$$

The conditional distribution of R_{866} can be seen in Figure 8. We notice,




Figure 6: Autocorrelation and cross-correlation plots for the multivariate time series. Notice the presence of correlation and cross-correlation in the data.




Figure 7: Conditional density of the one-step conditional distribution of R_{865} for the US stock data, with VaR at 95% (black) and observed return (red) highlighted.




Figure 8: Conditional density of the two-step conditional distribution of R_{866} for the US stock data, with VaR at 95% (black) and observed return (red) highlighted.

as expected, an increase in the standard deviation of the distribution, $\sigma_{\text{EFF}}^{(2)} = 0.0177$ with respect to $\sigma_{\text{EFF}} = 0.0177$. Overall, the two shapes in Figure 7 and 8 look similar, however the observed return R_{866} is not in a high density region of its predictive distribution, as it actually exceeds the expectations. VaR was now estimated at -0.021 , with expected shortfall equal to -0.0315 .

6 Comparing VAR, MVAR and DCC

Dynamic Conditional Correlation models (DCC, Engle 2002) are a class of multivariate GARCH models in which conditional correlations between elements of a vector series are time dependent. In particular, given the conditional covariance matrix of the model at time t , H_t , each entry h_{ij} of the matrix is modelled as a univariate GARCH model. DCC models are used in finance to predict behavior of vector time series in which the assets are correlated and heteroskedastic, thanks to the fact that the conditional covariance matrix of a predictor is always fully specified.

We compare here the performance of modeling the data in Section 5 with an MVAR model and a DCC model. We also add a comparison with a fitted vector autoregressive model of order 3 (VAR(3)). We compare the three models in terms of expected return, variance, risk measures and Continuous Ranked Probability Score (CRPS) associated to their respective minimum variance portfolios for 1 and 2 days density forecasts.

CRPS (see for instance Gneiting and Raftery 2007) was chosen as a strictly proper scoring rule to directly compare forecasting performance of the three methods: the lower the score, the better the forecast. Given an observation x and the associated forecast distribution F , CRPS is defined mathematically as:

$$CRPS(F, x) = \int_{\mathbb{R}} (F(y) - \mathcal{I}(y \geq x))^2 dy \quad (15)$$

	MVAR(2;3,2,1)	DCC-GARCH(1,1)	VAR(3)
$\hat{\mu}_{t+1}$	0.0024	0.0006	-0.0009
$\hat{\sigma}_{t+1}$	0.0092	0.0091	0.0053
VaR _{0.95}	-0.0134	-0.0144	-0.0096
ES _{0.95}	-0.0190	-0.0182	-0.0189
CRPS	0.0141	0.0159	0.0158

Table 1: Summary statistics for MVAR, DCC-GARCH and VAR for the one step ahead predictive distribution at $t = 864$.

	MVAR(2;3,2,1)	DCC-GARCH(1,1)	VAR(3)
$\hat{\mu}_{t+2}$	-0.0011	0.0010	-0.0018
$\hat{\sigma}_{t+2}$	0.0101	0.0094	0.0053
VaR _{0.95}	-0.0172	-0.0145	-0.0105
ES _{0.95}	-0.0240	-0.0184	-0.0127
CRPS	0.0118	0.0162	0.0120

Table 2: Summary statistics for MVAR and DCC-GARCH for the two step ahead predictive distribution at $t = 864$.

where $\mathcal{I}(\cdot)$ is the indicator function assuming value 1 when the argument $y > x$ is true, and 0 otherwise. CRPS is a measure of discrepancy between the forecast CDF, F , and the empirical CDF of the observation x . Notice that it was purposely chosen not to involve factor models due to the small number of assets considered.

A DCC-GARCH(1,1) was found to be the best model for the return series assuming multivariate normal innovations. Weights for minimum variance portfolios for both 1 and 2 days future observations were calculated using the respective conditional covariance matrices, and results were compared to our fitted MVAR(2;3,2,1) model.

Tables 1 and 2, and Figure 9 compare some summary statistics, risk measures and predictive densities obtained with MVAR and DCC-GARCH. For the DCC-GARCH, the distribution of returns is normal by model assumption by application of the property in Equation (16). MVAR is showing slightly heavy tails and a skewed distribution for the predictive distribution of R_{865} (which is a sign of more flexibility!). The two methods seem to perform similarly in terms of forecasting, with MVAR having a slightly lower CRPS score for both forecasts.

The fitted VAR(3) model seems to fail in capturing the heavy-tailed nature of the data, as the predictive distribution is very narrow around the predictor. While its performance in terms of CRPS is comparable to that of the DCC-GARCH model in 1-step prediction, the VAR model is clearly outperformed by both methods in the 2-step predictor, a sign that this linear time series model is not reliable as we try to predict further in the future.

We can conclude that our method can be a useful alternative to existing methods. At least for the examples discussed in the paper, our method performs at least as well as the DCC-GARCH method, while it outperforms the VAR model.




Figure 9: Comparison of conditional predictive densities at time $t = 864$ with MVAR(black line) and DCC-GARCH(red line).

7 Conclusions

The paper presents an innovative way of using Mixture vector autoregressive models for portfolio optimization. The method consists in deriving analytically predictive distributions of future observations, and use the covariance matrix, together with modern portfolio theory, to build an efficient portfolio and obtain a distribution for future returns. We have seen in fact that, assuming multivariate normal distributions for mixture components, the conditional predictive distribution of the portfolio return at a future horizon h itself follows a (univariate) mixture of g^h normal components, depending on observation up to the present.

The methodology was tested both on a simulated and a real dataset and compared with the widely used dynamic conditional correlation model, which uses multivariate GARCH to estimate conditional correlations. The conclusion was that the two methods have similar performance on the analyzed datasets, suggesting the proposed method can be considered a valid alternative.

A possible extension of this method is to incorporate factor models, in order to be able to model a large number of possibly correlated assets at a time. In addition, distribution assumptions other than normal can be made on the innovation terms. For example, considering a distribution with heavy tails might need a smaller number of components to fit the data. However, estimation would become much more complicated, and numerical algorithms would be required.

A A property of the multivariate normal distribution

At several places we use the standard result that any linear combination of the elements of a random vector from the multivariate normal distribution is univariate normal. More specifically, let X be a random vector of length m following a multivariate normal distribution with mean μ and covariance matrix Σ . Let a be a constant vector of the same length as X . Then

$$a^T X = \sum_{j=1}^m a_j X_j \sim N(a^T \mu, a^T \Sigma a) \quad (16)$$

B Derivation of the representation of y_{t+2} and its characteristic function

We here derive the analytic expression for the predictor y_{t+2} when information up to time t is available. We also derive its characteristic function, to show that the distribution of such predictor is a mixture of g^2 normal distributions. Proof is simply the multivariate version of the proof in Boshnakov (2009). Let $Z_t \in \{1, \dots, g\}$ be the allocation random variable defined in Section 2, and assume $z_{t+2} = k, z_{t+1} = l$ at times $t+2$ and $t+1$. We have that

$$\begin{aligned} y_{t+2} &= \mu_{t+2,k} + \Omega^{1/2} \varepsilon_{t+2,k} \\ &= \mu_{t+2,k} - \Theta_{k,1} Y_{t+1} + \Theta_{k,1} Y_{t+1} + \Omega^{1/2} \varepsilon_{t+2,k} \\ &= (\mu_{t+2,k} - \Theta_{k,1} Y_{t+1} + \Theta_{k,1} \mu_{t+1,l}) + \Theta_{k,1} \Omega_l^{1/2} \varepsilon_{t+1,l} + \Omega^{1/2} \varepsilon_{t+2,k} \\ &= \mu_{t+2;k,l} + \Theta_{k,1} \Omega_l^{1/2} \varepsilon_{t+1,l} + \Omega^{1/2} \varepsilon_{t+2,k} \end{aligned} \quad (17)$$

where $\varepsilon_{t+h,k}$ is the innovation term associated with the k^{th} component.

We want an expression that does not contain Y_{t+1} . Hence, we rewrite $\mu_{t+2;k,l}$ as

$$\begin{aligned} \mu_{t+2;k,l} &= \mu_{t+2,k} - \Theta_{k,1} Y_{t+1} + \Theta_{k,1} \mu_{t+1,l} \\ &= \Theta_{k,0} + \sum_{i=1}^p \Theta_{k,i} Y_{t+2-i} - \Theta_{k,1} Y_{t+1} + \Theta_{k,1} \left(\Theta_{l,0} + \sum_{i=1}^p \Theta_{l,i} Y_{t+1-i} \right) \\ &= \Theta_{k,0} + \Theta_{k,1} \Theta_{l,0} - \Theta_{k,1} Y_{t-1} + \Theta_{k,1} Y_{t+1} + \sum_{i=2}^p \Theta_{k,i} Y_{t+2-i} \\ &\quad + \Theta_{k,1} \sum_{i=1}^p \Theta_{l,i} Y_{t+1-i} \\ &= \Theta_{k,0} + \Theta_{k,1} \Theta_{l,0} + \sum_{i=1}^{p-1} \Theta_{k,i+1} Y_{t+1-i} + \Theta_{k,1} \sum_{i=1}^p \Theta_{l,i} Y_{t+1-i} \\ &= \Theta_{k,0} + \Theta_{k,1} \Theta_{l,0} + \sum_{i=1}^{p-1} (\Theta_{k,i+1} + \Theta_{k,1} \Theta_{l,i}) Y_{t+1-i} + \Theta_{k,1} \Theta_{l,p} Y_{t+1-p} \end{aligned}$$

And therefore we have the expression for Y_{t+2}

$$Y_{t+2} = \Theta_{k,0} + \Theta_{k,1}\Theta_{l,0} + \sum_{i=1}^{p-1} (\Theta_{k,i+1} + \Theta_{k,1}\Theta_{l,i})Y_{t+1-i} + \Theta_{k,1}\Theta_{l,p}Y_{t+1-p} \\ + \Theta_{k,1}\Omega_l^{1/2}\varepsilon_{t+1,l} + \Omega_k^{1/2}\varepsilon_{t+2,k}$$

We deduce that, given observed z_{t+2}, z_{t+1} :

$$\begin{aligned} \mathbb{E}[Y_{t+2} \mid Z_{t+2}, Z_{t+1}, \mathcal{F}_t] \\ = \Theta_{k,0} + \Theta_{k,1}\Theta_{l,0} + \sum_{i=1}^{p-1} (\Theta_{k,i+1} + \Theta_{k,1}\Theta_{l,i})Y_{t+1-i} + \Theta_{k,1}\Theta_{l,p}Y_{t+1-p} \\ \text{Cov}(Y_{t+2} \mid Z_{t+2}, Z_{t+1}, \mathcal{F}_t) = \Theta_{k,1}\Omega_l\Theta_{k,1} + \Omega_k \end{aligned}$$

We now need to derive the characteristic function for the predictor. Recall the characteristic function for the multivariate normal distribution and Y_{t+1} can be written as

$$\varphi_{t+1|t} \equiv \mathbb{E} \left[e^{is^T y_{t+1}} \mid \mathcal{F}_{t-1} \right] = \mathbb{E} \left[\sum_{k=1}^g \pi_k e^{is^T \mu_{t+1;k}} \varphi_k(\Omega_k^{1/2} s) \right]$$

It follows that, for Y_{t+2} , we have

$$\begin{aligned} \varphi_{t+2|t}(s) &\equiv \mathbb{E} \left[e^{is^T Y_{t+2}} \mid \mathcal{F}_t \right] = \mathbb{E} \left[\mathbb{E} \left(e^{is^T Y_{t+2}} \mid z_{t+2}, z_{t+1}, \mathcal{F}_t \right) \mid \mathcal{F}_t \right] \\ &= \mathbb{E} \left[i s^T \mu_{t+2;k,l} \mathbb{E} \left(e^{\Theta_{k,1}\Omega_l^{1/2}\varepsilon_{t+1,l} + \Omega_k^{1/2}\varepsilon_{t+2,k}} \mid z_{t+2}, z_{t+1}, \mathcal{F}_t \right) \mid \mathcal{F}_t \right] \\ &= \sum_{k,l=1}^g \pi_k \pi_l e^{is^T \mu_{t+2;k,l}} \varphi_1(\Theta_{k,1}\Omega_l^{1/2} s) \varphi_2(\Omega_k^{1/2} s) \end{aligned}$$

Thus, the conditional distribution of Y_{t+2} given \mathcal{F}_t is a mixture of g^2 components with mixing weights $\pi_k \pi_l$. For a normal mixture, we also have that:

$$\varphi_1(\Theta_{k,1}\Omega_l^{1/2} s) \varphi_2(\Omega_k^{1/2} s) = e^{\Theta_{k,1}\Omega_l\Theta_{k,1}^T} e^{\Omega_k} = e^{\Theta_{k,1}\Omega_l\Theta_{k,1}^T + \Omega_k}$$

which shows that the conditional distribution of the two-step predictor is a mixture of Normals with means $\mu_{t+2;k,l}$ and covariance matrices $\Theta_{k,1}\Omega_l\Theta_{k,1}^T + \Omega_k$.

References

- Alexander, C.: 2000, A primer on the orthogonal garch model, manuscript ISMA Centre, University of Reading, UK 2.
- Bollerslev, T., Engle, R. F. and Wooldridge, J. M.: 1988, A capital asset pricing model with time-varying covariances, Journal of Political Economy **96**(1), 116–131.
URL: <http://www.jstor.org/stable/1830713>
- Boshnakov, G. N.: 2009, Analytic expressions for predictive distributions in mixture autoregressive models., Stat. Probab. Lett. **79**(15), 1704–1709.

- Boshnakov, G. N.: 2011, On first and second order stationarity of random coefficient models, Linear Algebra and its Applications **434**(2), 415 – 423.
URL: <http://www.sciencedirect.com/science/article/pii/S0024379510004866>
- Engle, R.: 2002, Dynamic conditional correlation, Journal of Business & Economic Statistics **20**(3), 339–350.
URL: <https://doi.org/10.1198/073500102288618487>
- Engle, R. F. and Kroner, K. F.: 1995, Multivariate simultaneous generalized arch, Econometric Theory **11**(1), 122–150.
URL: <http://www.jstor.org/stable/3532933>
- Fong, P. W., Li, W. K., Yau, C. W. and Wong, C. S.: 2007, On a mixture vector autoregressive model, The Canadian Journal of Statistics / La Revue Canadienne de Statistique **35**(1), 135–150.
URL: <http://www.jstor.org/stable/20445243>
- Gneiting, T. and Raftery, A. E.: 2007, Strictly proper scoring rules, prediction, and estimation, Journal of the American Statistical Association **102**(477), 359–378.
URL: <https://doi.org/10.1198/016214506000001437>
- Kalliovirta, L., Meitz, M. and Saikkonen, P.: 2016, Gaussian mixture vector autoregression, Journal of Econometrics **192**(2), 485 – 498. Innovations in Multiple Time Series Analysis.
URL: <http://www.sciencedirect.com/science/article/pii/S030440761630015X>
- Markowitz, H.: 1952, Portfolio selection, The Journal of Finance **7**(1), 77–91.
URL: <http://www.jstor.org/stable/2975974>
- Saikkonen, P.: 2007, Stability of mixtures of vector autoregressions with autoregressive conditional heteroskedasticity, Statistica Sinica **17**(1), 221–239.
- Santos, A. A. and Moura, G. V.: 2014, Dynamic factor multivariate garch model, Computational Statistics & Data Analysis **76**, 606 – 617. CFEnetwork: The Annals of Computational and Financial Econometrics.
URL: <http://www.sciencedirect.com/science/article/pii/S0167947312003398>
- Van der Weide, R.: 2002, Go-garch: a multivariate generalized orthogonal garch model, Journal of Applied Econometrics **17**(5), 549–564.
- Wong, C. S. and Li, W. K.: 2000, On a mixture autoregressive model., J. R. Stat. Soc., Ser. B, Stat. Methodol. **62**(1), 95–115.