

L^p -Kato class measures and their relations with Sobolev embedding theorems for Dirichlet spaces

Takahiro Mori*

Abstract

In this paper, we discuss relationships between the continuous embeddings of Dirichlet spaces $(\mathcal{F}, \mathcal{E}_1)$ into Lebesgue spaces and the integrability of the associated resolvent kernel $r_\alpha(x, y)$. For a positive measure μ , we consider the following two properties; the first one is that the Dirichlet space $(\mathcal{F}, \mathcal{E}_1)$ is continuously embedded into $L^{2p}(E; \mu)$ (which we write as $(\text{Sob})_p$), and the second one is that the family of 1-order resolvent kernels $\{r_1(x, y)\}_{x \in E}$ is uniformly p -th integrable in y with respect to the measure μ (which we write as $(\text{Dyn})_p$).

Under some assumptions, for a measure μ satisfying $(\text{Dyn})_1$, we prove $(\text{Dyn})_{p'}$ implies $(\text{Sob})_p$ for $1 \leq p \leq p' < \infty$, and prove $(\text{Sob})_{p'}$ implies $(\text{Dyn})_p$ for $1 \leq p < p' < \infty$. To prove these results we introduce L^p -Kato class, an L^p -version of the set of Kato class measures, and discuss its properties. We also give variants of such relations corresponding to the Gagliardo-Nirenberg type interpolation inequalities. As an application, we discuss the continuity of intersection measures in time.

Keywords: Dirichlet form; Sobolev embedding theorem; Kato class; Resolvent kernel

Mathematics Subject Classification (2010): 31C25 (primary); 46E35; 60J35; 60J45

1 Introduction

In this paper, we discuss relationships between the continuous embeddings of Dirichlet spaces into Lebesgue spaces and the integrability of the associated resolvent kernel.

The prototype of the relationships we are focusing on is the classical Dirichlet integral $(\frac{1}{2}\mathbf{D}, H^1(\mathbb{R}^d))$ on \mathbb{R}^d and the associated resolvent kernel $r_\alpha(x, y)$, $x, y \in \mathbb{R}^d$, $\alpha > 0$, that is, $H^1(\mathbb{R}^d)$ is the Sobolev space on \mathbb{R}^d ,

$$\mathbf{D}(u, v) = \sum_{i=1}^d \int_{\mathbb{R}^d} \frac{\partial u}{\partial x_i} \frac{\partial v}{\partial x_i} dx \quad \text{for } u, v \in H^1(\mathbb{R}^d)$$

and

$$r_\alpha(x, y) = \frac{1}{(2\pi)^{d/2}} \int_0^\infty \frac{1}{t^{d/2}} \exp\left\{-\left(\alpha t + \frac{|x-y|^2}{2t}\right)\right\} dt \quad \text{for } x, y \in \mathbb{R}^d, \alpha > 0.$$

The classical Sobolev embedding theorem on \mathbb{R}^d is well known:

$$H^1(\mathbb{R}^d) \text{ is continuously embedded into } L^{2p}(\mathbb{R}^d) \text{ only for } p \in [1, \infty) \text{ with} \quad (1.1)$$

$$d - p(d-2) \geq 0 \text{ (hence, } 2p \leq 2d/(d-2) \text{ when } d \geq 2).$$

By an elementary calculation, it holds that

$$\sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d} r_1(x, y)^p dy < \infty \text{ if and only if } d - p(d-2) > 0. \quad (1.2)$$

Note that $d - p(d-2)$ appears in both conditions (1.1) and (1.2). This means that there is a relation between the Sobolev embedding theorem and the integrability of the resolvent.

*Research Institute for Mathematical Sciences, Kyoto University, Kyoto, 606-8502, JAPAN. tmori@kurims.kyoto-u.ac.jp

The main purpose of this paper is to generalize such relations from the perspective of the Dirichlet form theory. Let E be a locally compact separable metric space, m be a Radon measure on E with $\text{supp}[m] = E$, and let $(\mathcal{E}, \mathcal{F})$ be a Dirichlet form on $L^2(E; m)$ with the associated resolvent kernel $r_\alpha(x, y)$ with respect to m . We simply write the inner product $\mathcal{E}_1(u, v) = \mathcal{E}(u, v) + \int_E uv dm$ for $u, v \in \mathcal{F}$. Suppose μ is a Radon measure on E . For $p \in [1, \infty)$, we consider two properties;

(Sob) $_p$ the Hilbert space $(\mathcal{F}, \mathcal{E}_1)$ is continuously embedded into $L^{2p}(E; \mu)$, that is, there exists a positive constant $C > 0$ such that $\|u\|_{L^{2p}(E; \mu)}^2 \leq C\mathcal{E}_1(u, u)$ for all $u \in \mathcal{F}$,

(Dyn) $_p$ it holds that $\sup_{x \in E} \int_E r_1(x, y)^p \mu(dy) < \infty$.

The property (Dyn) $_p$ is named after Dynkin, which is equivalent to $\mathcal{D}^p(X)$ defined later. The aim of this paper is to show the following: under some conditions, for any measure μ satisfying (Dyn) $_1$,

• if (Dyn) $_{p'}$ holds for some $p' \in [1, \infty)$, then (Sob) $_p$ holds for all $1 \leq p \leq p'$, (1.3)

• if (Sob) $_{p'}$ holds for some $p' \in [1, \infty)$, then (Dyn) $_p$ holds for all $1 \leq p < p'$. (1.4)

See Sections 4.1 and 4.2 for precise statements and proofs. We will prove (1.3) in Theorem 4.1 (i) and the following Corollary 4.2. We will also prove (1.4) in Theorem 4.7 (i). Regarding (1.4), Lemma 4.6 gives a stronger result when μ is the reference measure m , that is, we obtain the order of decay for the quantity $\sup_{x \in E} \int_E r_\alpha(x, y)^p m(dy)$ as $\alpha \uparrow \infty$. By (1.3) and (1.4), (1.2) follows from (1.1) and (1.1) with $d - p(d - 2) > 0$ follows from (1.2). In Theorem 4.1 and Theorem 4.7, we also give some variants of such relationships.

When μ is the reference measure m , the Sobolev inequality has been studied for various settings; Euclidean space, Riemannian manifolds, Lie groups, and so on (see [22, 3] for example). It is known that the Sobolev inequality is equivalent to the ultra-contractivity of the associated transition semigroup [30], the Nash type inequality [6], and the capacity isoperimetric inequality [15, 13].

When $p = p' = 1$, our result (1.3) is related to the theory of the Kato class of measures. Kato class is introduced to analyse the Schrödinger semigroups and analyse integral kernels of semigroups given by Feynman-Kac functionals (see [1, 2] for example). The set of measures satisfying (Dyn) $_1$ is the so-called the Dynkin class (1-order version of Green-bounded measures). The embedding result (1.3) for $p = p' = 1$ is proved by Stollmann and Voigt [25] via operator theory, and later Shiozawa and Takeda [24] proved it in terms of Dirichlet forms.

The organization of the paper is as follows. In Section 2.1, we give the framework. In Section 2.2, we introduce an L^p -Dynkin class, which is equivalent to (Dyn) $_p$, and an L^p -Kato class, which is an L^p -analog of the classical Kato class. We will give equivalent conditions of these classes in terms of heat kernels, so we can check that a measure is in the classes once an upper bound of the heat kernel such as the (sub-)Gaussian estimate (see (2.13)) or the jump type estimate (see (2.14)) holds for a short time. These estimates are established for many processes. Regarding the (sub-)Gaussian estimate, it is obtained for Brownian motion on a manifold [20], Brownian motion on a metric measure space with Riemannian curvature dimension condition [26], Brownian motion on the Sierpiński gasket [5] and other diffusions on fractals [4], and so on. Regarding the jump-type estimate, stable-like processes on d sets [8] are studied for example.

In Section 2.3, we introduce a subclass of the L^p -Kato class (denoted by $\mathcal{K}^{p, \delta}(X)$), which has additional information on the order of decay for the quantity $\sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy)$ as $\alpha \uparrow \infty$. Similarly to the L^p -Dynkin and the L^p -Kato classes, this subclass can be characterized via the heat kernel estimate, and the above example satisfies the estimate.

In Section 3, a relation between L^p -Dynkin classes for a process and for its time changed process is discussed. This section plays a key role later to prove (1.4). Sections 4.1 and 4.2 are devoted to proving (1.3) and (1.4) as mentioned above. In these sections, we also prove variants of (1.3) and

(1.4) corresponding to the L^p -Kato class and the subclass $\mathcal{K}^{p,\delta}(X)$, respectively, and the later gives relations with the Gagliardo-Nirenberg type interpolation inequality.

Section 5 is an application to the intersection of the paths of independent stochastic processes. Analysis of the intersection of Brownian paths was initiated by Dvoretzky, Erdős, Kakutani [9, 10] and Dvoretzky, Erdős, Kakutani and Taylor [11]. They gave the following dichotomy: for p independent Brownian motions $B^{(1)}, \dots, B^{(p)}$ on \mathbb{R}^d ,

$$\begin{aligned} & \text{the paths intersect, i.e., } B^{(1)}(0, \infty) \cap \dots \cap B^{(p)}(0, \infty) \neq \emptyset \text{ almost surely if} \\ & d - p(d - 2) > 0, \text{ and do not intersect almost surely if } d - p(d - 2) \leq 0. \end{aligned} \quad (1.5)$$

Note that the same number $d - p(d - 2)$ appears as in (1.1) and (1.2).

Motivated by problems in statistical physics such as the configurations of interacting polymers, a random measure called the intersection local time has been introduced; see [19] for example. In this paper, we consider the occupation measure of the set of intersections for independent processes $X^{(1)}, \dots, X^{(p)}$ with the same distribution X which is formally written as

$$\ell_t^{\text{IS}}(A) = \int_A \left[\prod_{i=1}^p \int_0^{t_i} \delta_x(X^{(i)}(s_i)) ds_i \right] m(dx) \quad \text{for } A \in \mathcal{B}(E)$$

and for $\mathbf{t} = (t_1, \dots, t_p) \in [0, \infty)^p$, where δ_x is the Dirac measure at x , m is the reference measure of the processes and $\mathcal{B}(E)$ is the family of Borel sets in E . We call the measure as *the (mutual) intersection measure* named after König and Mukherjee [17]. Here and in the following, the superscript “IS” means “InterSection”.

In Theorem 5.1 we prove the following; if the reference measure m belongs to the subclass $\mathcal{K}^{p,\delta}(X)$ introduced in Section 2.3, then the measure-valued process $\mathbf{t} \mapsto \ell_{\mathbf{t}}^{\text{IS}}(dx)$ has a continuous modification, and the real-valued process $\mathbf{t} \mapsto \langle f, \ell_{\mathbf{t}}^{\text{IS}} \rangle$ has a Hölder continuous modification for each bounded Borel function f . This is a generalization of [7, Section 2.2], in which the results are obtained for independent Brownian motions.

2 L^p -Kato class and its variant

In this section, we give the framework and introduce the L^p -Kato class and its variants.

2.1 Framework

Let E be a locally compact, separable metric space and let m be a Radon measure on E with $\text{supp}[m] = E$. Let ∂ be a point added to E so that $E_{\partial} := E \cup \{\partial\}$ is the one-point compactification of E . The point ∂ serves as the cemetery point for E . Suppose $(\mathcal{E}, \mathcal{F})$ is a regular Dirichlet form on $L^2(E; m)$ and $X = (\Omega, X_t, \zeta, \mathbb{P}_x)$ is an associated m -symmetric Hunt process. For $\alpha > 0$ and $u \in \mathcal{F}$, we simply write $\mathcal{E}_{\alpha}(u, u) = \|u\|_{\mathcal{E}_{\alpha}}^2 := \mathcal{E}(u, u) + \alpha \int_E u^2 dm$. In this paper, we always take the quasi-continuous version of the element u of \mathcal{F} (see [12, Section 2] for example).

Throughout this paper, we assume that the transition kernel $(P_t)_{t>0}$ of X satisfies the absolute continuity condition:

$$\begin{aligned} & P_t(x, dy) \text{ is absolutely continuous with respect to } m(dy) \text{ for each} \\ & t > 0 \text{ and } x \in E. \end{aligned} \quad (2.1)$$

Note that the condition (2.1) implies the measurability of the heat kernel (see [31, Theorem 2] for example):

$$\begin{aligned} & (P_t)_{t>0} \text{ admits a heat kernel } p_t(x, y) \text{ which is jointly measurable on} \\ & (0, \infty) \times E \times E \text{ such that } p_t(x, y) = p_t(y, x) \text{ and} \\ & p_{t+s}(x, y) = \int_E p_s(x, z) p_t(z, y) m(dz) \text{ for all } s, t > 0, x, y \in E. \end{aligned} \quad (2.2)$$

Remark 2.1. We may consider a slightly weaker condition than (2.1):

There exists a Borel properly exceptional set N such that $P_t(x, dy)$ is absolutely continuous with respect to $m(dy)$ for each $t > 0$ and $x \in E \setminus N$. (2.3)

It is known that the Sobolev inequality implies (2.3) (see [12, Theorem 4.27] for example). Under the condition, we can obtain the corresponding results of this paper by replacing $\sup_{x \in E}$ by $\inf_{\text{Cap}(N)=0} \sup_{x \in E \setminus N}$ as in [2, (3.3)], where Cap is the (1-)capacity of the Dirichlet form $(\mathcal{E}, \mathcal{F})$. In this paper, we do not give detailed calculations under the assumption (2.3) and we always impose (2.1) for simplicity.

For each $\alpha > 0$, write the α -order resolvent kernel of X by $r_\alpha(x, y) = \int_0^\infty e^{-\alpha t} p_t(x, y) dt$. We denote by $S_{00}(X)$ the set of positive Borel measures μ such that $\mu(E) < \infty$ and $R_1\mu(x) := \int_E r_1(x, y) \mu(dy)$ is uniformly bounded in $x \in E$. A positive Borel measure μ on E is said to be *smooth in the strict sense* if there exists a sequence $\{E_n\}_{n=1}^\infty$ of Borel sets increasing to E such that $1_{E_n} \cdot \mu \in S_{00}(X)$ for each n and

$$\mathbb{P}_x \left(\lim_{n \rightarrow \infty} \sigma_{E \setminus E_n} \geq \zeta \right) = 1, \quad \text{for all } x \in E,$$

where $\sigma_{E \setminus E_n}$ is the first hitting time of $E \setminus E_n$. The totality of smooth measures in the strict sense is denoted by $S_1(X)$.

2.2 The class $\mathcal{K}^p(X)$

In this section, we introduce the L^p -version of the Kato class measures.

Definition 2.2. Let $p \in [1, \infty)$. For a positive Radon measure μ on E , μ is said to be in the *L^p -Kato class with respect to X* (in symbols $\mu \in \mathcal{K}^p(X)$) if

$$\limsup_{\alpha \uparrow \infty} \sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy) = 0 \tag{2.4}$$

and μ is said to be in the *L^p -Dynkin class with respect to X* (in symbols $\mu \in \mathcal{D}^p(X)$) if

$$\sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy) < \infty \quad \text{for some } \alpha > 0. \tag{2.5}$$

Clearly $\mathcal{K}^p(X) \subset \mathcal{D}^p(X)$. The condition $(\text{Dyn})_p$ which we introduced in Section 1 means nothing else than the L^p -Dynkin class.

Remark 2.3.

1. $\mathcal{K}^1(X)$ and $\mathcal{D}^1(X)$ are so-called the set of Kato and Dynkin class measures, respectively. The reference measure m always belongs to $\mathcal{K}^1(X)$.
2. If $\mu(E) < \infty$, Hölder's inequality gives that $\mu \in \mathcal{K}^{p'}(X)$ implies $\mu \in \mathcal{K}^p(X)$ for $1 \leq p < p'$.
3. The L^p -Kato and the L^p -Dynkin classes are well-defined up to the quasi-everywhere equivalence of the processes. Indeed, it holds that

$$\sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy) = \inf_{N \subset E, \text{Cap}(N)=0} \sup_{x \in E \setminus N} \int_E r_\alpha(x, y)^p \mu(dy). \tag{2.6}$$

Obviously the left-hand side is equal or greater than the right-hand side. To prove the converse inequality, let μ be a Radon measure on E , $\alpha > 0$, and $N \subset E$ be a zero-capacity set. For fixed $x \in E$, $M > 0$ and a compact set $K \subset E$, we write $\nu(dy) = 1_K(y) (r_\alpha(x, y) \wedge M)^{p-1} \mu(dy)$ and $R_\alpha \nu = \int_E r_\alpha(\cdot, y) \nu(dy)$. For $z \in E \setminus N$, we have by Hölder's inequality

$$\begin{aligned} R_\alpha \nu(z) &= \int_K r_\alpha(z, y) (r_\alpha(x, y) \wedge M)^{p-1} \mu(dy) \\ &\leq \left\{ \sup_{z \in E \setminus N} \int_E r_\alpha(z, y)^p \mu(dy) \right\}^{\frac{1}{p}} \left\{ \int_K (r_\alpha(x, y) \wedge M)^p \mu(dy) \right\}^{\frac{p-1}{p}}. \end{aligned}$$

Since $R_\alpha \nu$ is α -excessive and the absolute continuity condition (2.1) holds, we have

$$\begin{aligned} R_\alpha \nu(x) &= \lim_{\varepsilon \downarrow 0} e^{-\alpha\varepsilon} \mathbb{E}_x[R_\alpha \nu(X_\varepsilon)] = \lim_{\varepsilon \downarrow 0} e^{-\alpha\varepsilon} \int_{E \setminus N} p_\varepsilon(x, z) R_\alpha \nu(z) m(dz) \\ &\leq \left\{ \sup_{z \in E \setminus N} \int_E r_\alpha(z, y)^p \mu(dy) \right\}^{\frac{1}{p}} \left\{ \int_K (r_\alpha(x, y) \wedge M)^p \mu(dy) \right\}^{\frac{p-1}{p}}, \end{aligned}$$

which means that

$$\int_K (r_\alpha(x, y) \wedge M)^p \mu(dy) \leq \sup_{z \in E \setminus N} \int_E r_\alpha(z, y)^p \mu(dy).$$

Taking $M \uparrow \infty$ and $K \uparrow E$, we have the desired inequality by the dominated convergence theorem.

Hence one can define the L^p -Kato and the L^p -Dynkin classes by the right-hand side of (2.6) without probabilistic notations and denote them as $\mathcal{K}^p(\mathcal{E})$ and $\mathcal{D}^p(\mathcal{E})$ in symbols. Throughout this paper, we always use the sup-notation for simplicity because one can find such replacements of $\sup_{x \in E}$ with $\inf_{N \subset E, \text{Cap}(N)=0} \sup_{x \in E \setminus N}$ by similar calculations.

Example 2.4 (Brownian motion on \mathbb{R}^d). Suppose $E = \mathbb{R}^d$, m is the Lebesgue measure on \mathbb{R}^d and X is a Brownian motion on \mathbb{R}^d . Let $p \in [1, \infty)$ with $d - p(d - 2) > 0$ and μ be a positive Radon measure on \mathbb{R}^d . By the proof of [1, Theorem 4.5], $\mu \in \mathcal{K}^p(X)$ if and only if

$$\begin{aligned} \limsup_{r \downarrow 0} \sup_{x \in \mathbb{R}^d} \int_{|x-y| < r} \frac{\mu(dy)}{|x-y|^{p(d-2)}} &= 0, \quad d \geq 3, \\ \limsup_{r \downarrow 0} \sup_{x \in \mathbb{R}^d} \int_{|x-y| < r} (-\log|x-y|)^p \mu(dy) &= 0, \quad d = 2, \\ \sup_{x \in \mathbb{R}^d} \int_{|x-y| \leq 1} \mu(dy) &< \infty, \quad d = 1. \end{aligned}$$

In particular, when $d = 1$, $\mathcal{K}^1(X) = \mathcal{K}^p(X)$ for any $p > 1$.

By the above characterization, we may give a sufficient condition for $\mathcal{K}^p(X)$. If a Borel function f on \mathbb{R}^d satisfies $\sup_{x \in \mathbb{R}^d} \int_{|x-y| \leq 1} |f(y)|^r dy < \infty$ for some $r > d/(d - p(d - 2))$ for $d \geq 2$, or $r \geq 1$ for $d = 1$, then the measure $|f(x)|dx$ is in the class $\mathcal{K}^p(X)$. This gives an extension of [1, Theorem 1.4 (iii)], in which the result is obtained for $p = 1$. (See also [18], in which such results are obtained under more general heat kernel estimates.) In particular, $|x|^{-\beta} dx \in \mathcal{K}^p(X)$ if $\beta < d - p(d - 2)$ for $d \geq 2$, and $\beta < 1$ for $d = 1$.

When $d \geq 3$, Schechter [23] introduced related classes $M_{\alpha, r}$ ($\alpha > 0$, $r > 1$) of functions V given by

$$\sup_{x \in \mathbb{R}^d} \int_{|x-y| \leq 1} \frac{|V(y)|^r}{|x-y|^{d-\alpha}} dy < \infty,$$

and [1] studied relations between $M_{\alpha, r}$ and the classical Kato class $\mathcal{K}^1(X)$. By Hölder's inequality, we have $M_{\alpha, r} \subset \mathcal{K}^p(X)$ if $r > \alpha/(d - p(d - 2))$. This is an extension of [1, Proposition 4.1, 4.2], in which the result are obtained for $p = 1$. (Note that there are typos in [1]; $\beta > 2$ in Proposition 4.1 (resp. $\alpha > 2p$ in Proposition 4.2) should be $\beta < 2$ (resp. $\alpha < 2p$).)

Remark 2.5. Güneysu [14] gives another generalization of the Kato class to show the Hölder continuity of the Schrödinger semigroups. For $\alpha \in [0, 1]$, a Radon measure μ on E is said to be in the α -Kato class in the sense of Güneysu (in symbols $\mu \in \tilde{\mathcal{K}}^\alpha(X)$) if

$$\limsup_{t \downarrow 0} \sup_{x \in E} \int_E \left(\int_0^t s^{-\frac{\alpha}{2}} p_s(x, y) ds \right) \mu(dy) = 0.$$

Obviously $\tilde{\mathcal{K}}^0(X) = \mathcal{K}^1(X)$, which is the classical Kato class. When X is a Brownian motion on \mathbb{R}^d , by the proof of [1, Theorem 4.5] $\mu \in \tilde{\mathcal{K}}^\alpha(X)$ if and only if

$$\begin{aligned} \limsup_{r \downarrow 0} \sup_{x \in \mathbb{R}^d} \int_{|x-y| < r} \frac{\mu(dy)}{|x-y|^{d+\alpha-2}} &= 0, \quad d + \alpha > 2, \\ \limsup_{r \downarrow 0} \sup_{x \in \mathbb{R}^d} \int_{|x-y| < r} (-\log|x-y|) \mu(dy) &= 0, \quad d + \alpha = 2, \\ \sup_{x \in \mathbb{R}^d} \int_{|x-y| \leq 1} \mu(dy) &< \infty, \quad d + \alpha < 2. \end{aligned}$$

A comparison with Example 2.4 gives that the coincidence $\tilde{\mathcal{K}}^\alpha(X) = \mathcal{K}^p(X)$ also holds when $p \geq 1$, $d + \alpha - 2 = p(d - 2) > 0$, or when $p \geq 1$, $d = 1$, $0 \leq \alpha < 1$.

The following proposition is the L^p -version of [2, Proposition 3.8] in some sense.

Proposition 2.6. *Let $p \in [1, \infty)$. It holds that*

$$\mathcal{D}^p(X) \subset S_1(X).$$

Proof. Suppose $\mu \in \mathcal{D}^p(X)$ and assume first $\mu(E) < \infty$. As in Remark 2.3, we have $\mu \in \mathcal{D}^1(X)$, that is, $\sup_{x \in E} R_1 \mu(x) < \infty$. This means μ is in $S_{00}(X)$ and hence is in $S_1(X)$.

When $\mu \in \mathcal{D}^p(X)$ may not be a finite measure, take a sequence $\{E_n\}_{n=1}^\infty$ of relatively compact open sets that go to E as $n \uparrow \infty$. By the above, we have $1_{E_n} \cdot \mu \in S_{00}(X)$. Set $\sigma = \lim_{n \rightarrow \infty} \sigma_{E \setminus E_n}$. For each $x \in E$, the quasi-left-continuity of the Hunt process X (see for example, [12, Appendix A.2]) implies that

$$\lim_{n \rightarrow \infty} X_{\sigma_{E \setminus E_n}} = X_\sigma, \quad \mathbb{P}_x\text{-a.s. on } \{\sigma < \infty\}$$

and then $\mathbb{P}_x(\sigma \geq \zeta) = 1$, which concludes $\mu \in S_1(X)$. \square

The next two propositions characterize the L^p -Dynkin class and the L^p -Kato class in terms of the heat kernel.

Proposition 2.7. *Let $p \in [1, \infty)$. For a Radon measure μ on E , the following are equivalent:*

- (i) $\mu \in \mathcal{D}^p(X)$,
- (ii) $\sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy) < \infty$ for all $\alpha > 0$,
- (iii) $\sup_{x \in E} \int_E \left(\int_0^t p_s(x, y) ds \right)^p \mu(dy) < \infty$ for some $t > 0$,
- (iv) $\sup_{x \in E} \int_E \left(\int_0^t p_s(x, y) ds \right)^p \mu(dy) < \infty$ for all $t > 0$.

Proof. Trivially (ii) implies (i) and (iv) implies (iii).

Assume $\mu \in \mathcal{D}^p(X)$ and take $\beta > 0$ such that $\sup_{x \in E} \int_E r_\beta(x, y)^p \mu(dy) < \infty$. For $\alpha > \beta$, the monotonicity of the resolvent clearly implies that $\sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy) < \infty$. For $0 < \alpha < \beta$, fix $x \in E$ and set $F(\cdot) := \int_E r_\alpha(x, z) r_\beta(z, \cdot) m(dz)$. Then we have by Hölder's inequality,

$$\begin{aligned} \int_E F(y)^p \mu(dy) &= \int_E \left(\int_E F(y)^{p-1} r_\beta(z, y) \mu(dy) \right) r_\alpha(x, z) m(dz) \\ &\leq \int_E \left(\int_E F(y)^p \mu(dy) \right)^{\frac{p-1}{p}} \left(\int_E r_\beta(z, y)^p \mu(dy) \right)^{\frac{1}{p}} r_\alpha(x, z) m(dz) \\ &\leq \frac{1}{\alpha} \left(\int_E F(y)^p \mu(dy) \right)^{\frac{p-1}{p}} \left(\sup_{z \in E} \int_E r_\beta(z, y)^p \mu(dy) \right)^{\frac{1}{p}}, \end{aligned}$$

which implies that

$$\left(\int_E F(y)^p \mu(dy) \right)^{\frac{1}{p}} \leq \frac{1}{\alpha} \left(\sup_{z \in E} \int_E r_\beta(z, y)^p \mu(dy) \right)^{\frac{1}{p}}.$$

Hence the resolvent equation $r_\alpha(x, y) = r_\beta(x, y) + (\beta - \alpha) \int_E r_\alpha(x, z) r_\beta(z, y) m(dz)$ yields that

$$\begin{aligned} \left(\int_E r_\alpha(x, y)^p \mu(dy) \right)^{\frac{1}{p}} &\leq \left(\int_E r_\beta(x, y)^p \mu(dy) \right)^{\frac{1}{p}} + (\beta - \alpha) \left(\int_E F(y)^p \mu(dy) \right)^{\frac{1}{p}} \\ &\leq \frac{\beta}{\alpha} \left(\sup_{z \in E} \int_E r_\beta(z, y)^p \mu(dy) \right)^{\frac{1}{p}} < \infty, \end{aligned} \quad (2.7)$$

which concludes (ii). Moreover, we have for any $t > 0$, $x \in E$ and $\alpha > 0$,

$$\int_E \left(\int_0^t p_s(x, y) ds \right)^p \mu(dy) \leq e^{p\alpha t} \sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy), \quad (2.8)$$

which concludes (iv).

Next, assume (iii). Take $t_0 > 0$ such that $\sup_{x \in E} \int_E \left(\int_0^{t_0} p_s(x, y) ds \right)^p \mu(dx) < \infty$. For any $t \leq t_0$ and $a > 0$, we have

$$\int_E \left(\int_a^{a+t} p_s(x, y) ds \right)^p \mu(dy) \leq \sup_{z \in E} \int_E \left(\int_0^t p_s(x, y) ds \right)^p \mu(dy). \quad (2.9)$$

Indeed, the Chapman-Kolmogorov equation gives that the left-hand side equals

$$\begin{aligned} &\int_E \left\{ \int_0^t \int_E p_a(x, z) p_s(z, y) m(dz) ds \right\}^p \mu(dy) \\ &= \int_E \left\{ \int_E \left(\int_0^t p_s(z, y) ds \right) p_a(x, z) m(dz) \right\}^p \mu(dy). \end{aligned}$$

Applying Hölder's inequality with the measure $p_a(x, z) m(dz)$, the above equation is bounded from above by

$$\begin{aligned} &\int_E \left\{ \int_E \left(\int_0^t p_s(z, y) ds \right)^p p_a(x, z) m(dz) \right\}^{\frac{p}{p-1}} \left\{ \int_E 1^{\frac{p}{p-1}} p_a(x, z) m(dz) \right\}^{\frac{p(p-1)}{p}} \mu(dy) \\ &\leq \int_E \int_E \left(\int_0^t p_s(z, y) ds \right)^p \mu(dy) p_a(x, z) m(dz) \\ &\leq \sup_{z \in E} \int_E \left(\int_0^t p_s(z, y) ds \right)^p \mu(dy), \end{aligned}$$

where we used $P_a 1 \leq 1$ in the last two lines. This proves (2.9).

Now, suppose $t > 0$. By taking large N such that $Nt_0 \geq t$, we have from (2.9)

$$\begin{aligned} \left\{ \int_E \left(\int_0^t p_s(x, y) ds \right)^p \mu(dy) \right\}^{\frac{1}{p}} &\leq \sum_{n=0}^{N-1} \left\{ \int_E \left(\int_{nt_0}^{(n+1)t_0} p_s(x, y) ds \right)^p \mu(dy) \right\}^{\frac{1}{p}} \\ &\leq N \left\{ \sup_{z \in E} \int_E \left(\int_0^{t_0} p_s(x, y) ds \right)^p \mu(dy) \right\}^{\frac{1}{p}}, \end{aligned}$$

which concludes (iv).

Further, for any $\alpha > 0$, $x \in E$ and $t \leq t_0$, we have from the triangle inequality,

$$\begin{aligned} \left(\int_E r_\alpha(x, y)^p \mu(dy) \right)^{\frac{1}{p}} &= \left\| \int_0^\infty e^{-\alpha s} p_s(x, \cdot) ds \right\|_{L^p(E; \mu)} \\ &\leq \sum_{n=0}^\infty \left\| \int_{nt}^{(n+1)t} e^{-\alpha s} p_s(x, \cdot) ds \right\|_{L^p(E; \mu)} \end{aligned}$$

and by (2.9), the right-hand side is bounded from above by

$$\sum_{n=0}^\infty e^{-\alpha n t} \left(\sup_{x \in E} \int_E \left(\int_0^t p_s(x, y) ds \right)^p \mu(dy) \right)^{\frac{1}{p}}$$

$$= \frac{1}{1 - e^{-\alpha t}} \left(\sup_{x \in E} \int_E \left(\int_0^t p_s(x, y) ds \right)^p \mu(dy) \right)^{\frac{1}{p}}, \quad (2.10)$$

which concludes (ii). \square

Corollary 2.8. *Let $p \in [1, \infty)$. Then, for a Radon measure μ on E , the following are equivalent:*

(i) $\mu \in \mathcal{K}^p(X)$,

(ii) $\limsup_{t \downarrow 0} \sup_{x \in E} \int_E \left(\int_0^t p_s(x, y) ds \right)^p \mu(dy) = 0$.

Proof. (i) implies (ii) by letting $t \downarrow 0$ and then $\alpha \uparrow \infty$ in (2.8). Conversely, (ii) implies (i) by letting $\alpha \uparrow \infty$ and then $t \downarrow 0$ in (2.10). \square

Remark 2.9. Let $X^{(1)}$ be the 1-subprocess of X , that is, the m -symmetric Markov process with transition probability $e^{-t} p_t(x, y) m(dy)$. Clearly $X^{(1)}$ satisfies the absolute continuity condition (2.1). We claim that $\mathcal{K}^p(X) = \mathcal{K}^p(X^{(1)})$. Indeed, we have the inclusion $\mathcal{K}^p(X) \subset \mathcal{K}^p(X^{(1)})$ since the α -order resolvent kernel of $X^{(1)}$ is $r_{1+\alpha}(x, y)$ and the inequality $r_{1+\alpha}(x, y) \leq r_\alpha(x, y)$ holds. By applying (2.7) with $\beta = \alpha + 1$, we have the converse inclusion $\mathcal{K}^p(X^{(1)}) \subset \mathcal{K}^p(X)$. In the same way, we also have $\mathcal{D}^p(X) = \mathcal{D}^p(X^{(1)})$.

2.3 The class $\mathcal{K}^{p,\delta}(X)$

In this section, we introduce a subclass of L^p -Kato class, which has additional information on the order of decay of the quantities $\sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy)$ (they have been introduced in Definition 2.2) as $\alpha \uparrow \infty$.

Definition 2.10. Let $p \in [1, \infty)$ and $\delta \in (0, 1]$. For a positive Radon measure μ on E , μ is said to be in the L^p -Kato class with order δ (in symbols $\mu \in \mathcal{K}^{p,\delta}(X)$) if

$$\sup_{x \in E} \left(\int_E r_\alpha(x, y)^p \mu(dy) \right)^{\frac{1}{p}} = O(\alpha^{-\delta}) \quad \text{as } \alpha \rightarrow \infty.$$

That is, there exist constants $C > 0$ and $\alpha_0 > 0$ such that the left-hand side is bounded from above by $C\alpha^{-\delta}$ for all $\alpha > \alpha_0$. Clearly $\mathcal{K}^{p,\delta}(X) \subset \mathcal{K}^p(X)$.

Similarly to Proposition 2.7 and Corollary 2.8, we can characterize the set of L^p -Kato class measures with order δ in terms of the heat kernel.

Proposition 2.11. *Let $p \in [1, \infty)$ and $\delta \in (0, 1]$. For a Radon measure μ on E , the following are equivalent:*

(i) $\mu \in \mathcal{K}^{p,\delta}(X)$,

(ii) $\sup_{x \in E} \left(\int_E \left(\int_0^t p_s(x, y) ds \right)^p \mu(dy) \right)^{\frac{1}{p}} = O(t^\delta)$ as $t \rightarrow 0$.

Proof. By setting $\alpha t = 1$, (i) implies (ii) from (2.8) and (ii) implies (i) from (2.10). \square

In the following, we write

$$\gamma(\alpha, \mu, p) := \sup_{x \in E} \left(\int_E r_\alpha(x, y)^p \mu(dy) \right)^{\frac{1}{p}} \quad (2.11)$$

and write

$$\eta(t, \mu, p) := \sup_{x \in E} \left(\int_E \left(\int_0^t p_s(x, y) ds \right)^p \mu(dy) \right)^{\frac{1}{p}}. \quad (2.12)$$

In the sequel, we write as $\gamma(\alpha)$ and $\eta(t)$ in short if there is no danger of confusion.

Remark 2.12.

1. As we see in Corollary 2.8, if $\mu \in \mathcal{K}^p(X)$ then $\eta(t) < \infty$ for all $t > 0$. Hence, under (2.2), $\mu \in \mathcal{K}^{p,\delta}(X)$ if and only if $\sup_{0 < t \leq T} \{t^{-\delta} \eta(t)\} < \infty$ for some (also for all) $T > 0$.
2. Let $X^{(1)}$ be the 1-subprocess of X . By the same way as in Remark 2.9, one can show that $\mathcal{K}^{p,\delta}(X) = \mathcal{K}^{p,\delta}(X^{(1)})$.

Example 2.13 (Brownian motion). We continue with Example 2.4. The heat kernel of the Brownian motion on \mathbb{R}^d is $p_t(x - y)$, where

$$p_t(x) = (2\pi t)^{-\frac{d}{2}} e^{-\frac{|x|^2}{2t}} \quad t > 0, x \in \mathbb{R}^d.$$

Then, as in the proof of [7, Lemma 2.2.4], we have for $t > 0$,

$$\int_{\mathbb{R}^d} \left(\int_0^t p_s(x) ds \right)^p dx \leq (2\pi)^{-\frac{d(p-1)}{2}} p^{-\frac{d}{2}} \left(\frac{2p}{2p - d(p-1)} \right)^p t^{\frac{2p-d(p-1)}{2}},$$

whenever $2p - d(p-1) = d - p(d-2) > 0$. Hence, the Lebesgue measure on \mathbb{R}^d is in $\mathcal{K}^{p, \frac{d-p(d-2)}{2p}}(X)$ if $d - p(d-2) > 0$.

Example 2.14 ((Sub-)Gaussian heat kernel estimate). Let ρ denote the metric on E , let $\text{diam}(E) := \sup\{\rho(x, y) : x, y \in E\}$ denote the diameter of E , and denote with $B(x, r)$ the open ball with center $x \in E$ and radius $r > 0$. Assume that $\text{diam}(E) = 1$, $m(E) < \infty$ and there exist constants $c_1, c_2 > 0$ and $d_f \geq 1$ such that $c_1 r^{d_f} \leq m(B(x, r)) \leq c_2 r^{d_f}$ for all $x \in E$, $r \in (0, 1]$. We also assume that $p_t(x, y)$ enjoys the (sub-)Gaussian heat kernel upper estimate: there exist constants $c_3, c_4 > 0$ and $d_w \geq 2$ such that

$$p_t(x, y) \leq c_3 t^{-\frac{d_f}{d_w}} \exp \left\{ -c_4 \left(\frac{\rho(x, y)^{d_w}}{t} \right)^{\frac{1}{d_w-1}} \right\} \quad \text{for all } x, y \in E, t \in (0, 1]. \quad (2.13)$$

Then, a straightforward calculation gives that $m \in \mathcal{K}^{p,\delta}(X)$ if $pd_w\delta < d_f - p(d_f - d_w)$, i.e., $\delta < (d_s - p(d_s - 2))/2p$ by setting $d_s := 2d_f/d_w$. d_f , d_w and d_s are the so-called fractal dimension of E and walk dimension and spectrum dimension of the process X , respectively.

Example 2.15 (Jump-type heat kernel estimate). Under the setting of Example 2.14 we assume that $p_t(x, y)$ enjoys the jump-type heat kernel upper estimate: there exist constants $c_3 > 0$ and $d_w \geq 2$ such that

$$p_t(x, y) \leq c_3 \left\{ t^{-\frac{d_f}{d_w}} \wedge \frac{t}{\rho(x, y)^{d_f+d_w}} \right\} \quad \text{for all } x, y \in E, t \in (0, 1]. \quad (2.14)$$

Then, a straightforward calculation gives that $m \in \mathcal{K}^{p,\delta}(X)$ if $pd_w\delta < d_f - p(d_f - d_w)$, i.e., $\delta < (d_s - p(d_s - 2))/2p$ by setting $d_s := 2d_f/d_w$.

In [21, Section 1.3], these computations are made in the context of the intersection measure for such processes.

3 L^p -Dynkin class with respect to the time changed process

In this section, we discuss a relation between L^p -Dynkin classes with respect to X and with respect to its time changed process. The goal of this section is to prove Proposition 3.2.

We first introduce the notation about the time changed processes of X (for detail, see [12, Section 6] for example). Let $\{\mathcal{F}_t\}_{t \geq 0}$ is the minimum completed admissible filtration of X , that is, $\mathcal{F}_t := \bigcap_{\mu \in \mathcal{P}(E_\partial)} \mathcal{F}_t^\mu$ for $0 \leq t < \infty$, where $\mathcal{P}(E_\partial)$ is the set of probability measures on E_∂ , and \mathcal{F}_t^μ is the σ -algebra generated by $\{X_s : s \leq t\}$ and the null sets of the \mathbb{P}_μ -completion of $\sigma(X_t : 0 \leq t < \infty)$. We also let θ_t be the translation operator on Ω , that is, θ_t is a map from Ω to Ω such that $X_s \circ \theta_t = X_{s+t}$

for all $s \geq 0$. A stochastic process $\{A_t\}_{t \geq 0}$ is said to be a *positive continuous additive functional in the strict sense* (PCAF in abbreviation) if the following conditions hold:

- (i) $A_t(\cdot)$ is \mathcal{F}_t -measurable for all $t \geq 0$,
- (ii) there exists a set $\Lambda \in \mathcal{F}_\infty = \sigma(\bigcup_{t \geq 0} \mathcal{F}_t)$ such that $\mathbb{P}_x(\Lambda) = 1$ for all $x \in E$, $\theta_t \Lambda \subset \Lambda$ for all $t > 0$, and for each $\omega \in \Lambda$, $A_\cdot(\omega)$ is a real-valued continuous function satisfying the following: $A_0(\omega) = 0$, $A_t(\omega) = A_\zeta(\omega)$ for $t \geq \zeta$, and $A_{t+s}(\omega) = A_t(\omega) + A_s(\theta_t \omega)$ for $t, s \geq 0$.

It is known that there is a one-to-one correspondence between $S_1(X)$ and the family of PCAF's (Revuz correspondence) as follows: for each $\mu \in S_1(X)$, there exists a unique PCAF $\{A_t\}_{t \geq 0}$ such that for any non-negative Borel function f on E and γ -excessive function h ($\gamma > 0$), it holds that

$$\int_E f(x)h(x)\mu(dx) = \lim_{t \downarrow 0} \frac{1}{t} \mathbb{E}_{h,m} \left[\int_0^t f(X_s) dA_s \right],$$

where $\mathbb{E}_{h,m}[\cdot] = \int_E \mathbb{E}_x[\cdot] h(x) \mu(dx)$ (see [12, Theorem 5.1.7] for example). We denote by A^μ the PCAF corresponding to $\mu \in S_1(X)$. We write the fine support of $\mu \in S_1(X)$ as F , that is,

$$F := \{x \in E : \mathbb{P}_x(\tau = 0) = 1\}, \quad \tau = \inf\{t > 0 : A_t^\mu > 0\}.$$

For $\mu \in S_1(X)$, denote $\check{X} = (\Omega, \check{X}_t, \check{\zeta}, \mathbb{P}_x)$ the time changed process of X with respect to the PCAF A^μ , that is,

$$\check{X}_t = X_{\tau_t}, \quad \tau_t = \inf\{s > 0 : A_s^\mu > t\}, \quad \check{\zeta} = A_\zeta^\mu.$$

Note that \check{X} is a μ -symmetric Hunt process on F . Write the α -order resolvent of \check{X} by

$$\check{R}_\alpha f(x) = \mathbb{E}_x \left[\int_0^\infty e^{-\alpha t} f(\check{X}_t) dt \right], \quad f \in \mathcal{B}_b(F), x \in F,$$

where $\mathcal{B}_b(F)$ is the set of bounded Borel functions on F .

We note that \check{X} also satisfies the absolute continuity condition (2.1):

Lemma 3.1. *Let μ be a measure in $S_1(X)$. Then the time changed process \check{X} satisfies the absolute continuity condition (2.1).*

Proof. Suppose $\mu(N) = 0$. By the definition of $S_1(X)$, we can take a sequence $\{E_n\}_{n=1}^\infty$ of Borel sets increasing to E such that $1_{E_n} \cdot \mu \in S_{00}(X)$ for each n and

$$\mathbb{P}_x \left(\lim_{n \rightarrow \infty} \sigma_{E \setminus E_n} \geq \zeta \right) = 1, \quad \text{for all } x \in E. \quad (3.1)$$

For each n , the Revuz correspondence (see [12, Theorem 5.1.6] for example) implies that

$$\mathbb{E}_x \left[\int_0^\infty e^{-\alpha t} 1_{N \cap E_n}(X_t) dA_t^\mu \right] = R_\alpha[1_{N \cap E_n} \cdot \mu](x) = 0$$

for all $\alpha > 0$ and $x \in E$. By letting $\alpha \downarrow 0$, we have

$$\mathbb{E}_x \left[\int_0^\infty 1_{N \cap E_n}(X_t) dA_t^\mu \right] = 0.$$

Since the inclusion $\{t < \sigma_{E \setminus E_n}\} \subset \{X_t \in E_n \cup \{\partial\}\}$ holds, the above equality and (3.1) give that

$$\begin{aligned} \mathbb{E}_x \left[\int_0^\infty 1_N(X_t) dA_t^\mu \right] &= \lim_{n \rightarrow \infty} \mathbb{E}_x \left[\int_0^{\sigma_{E \setminus E_n}} 1_N(X_t) dA_t^\mu \right] \\ &\leq \lim_{n \rightarrow \infty} \mathbb{E}_x \left[\int_0^\infty 1_{N \cap E_n}(X_t) dA_t^\mu \right] = 0. \end{aligned}$$

Now, we have for every $x \in F$

$$\begin{aligned} \check{R}_\alpha 1_N(x) &= \mathbb{E}_x \left[\int_0^\infty e^{-\alpha t} 1_N(\check{X}_t) dt \right] \\ &\leq \mathbb{E}_x \left[\int_0^\infty 1_N(\check{X}_t) dt \right] = \mathbb{E}_x \left[\int_0^\infty 1_N(X_t) dA_t^\mu \right] = 0, \end{aligned}$$

which concludes the absolute continuity condition for \check{X} (see [12, Theorem 4.2.4] for example). \square

The next proposition will play a key role later to prove Theorem 4.7 (i), one of our main results. Roughly it means that, if μ is L^p -Dynkin with respect to the time changed process, then μ is L^p -Dynkin with respect to the original process.

Proposition 3.2. *Let $p \in (1, \infty)$ and let μ be a measure in $S_1(X)$. Then, $\mu \in \mathcal{D}^1(X)$ and $\mu \in \mathcal{D}^p(\check{X})$ imply $\mu \in \mathcal{D}^p(X)$.*

To prove this, we first consider the transient version of Proposition 3.2.

Definition 3.3. When $(\mathcal{E}, \mathcal{F})$ is transient, μ is said to be *Green-bounded* (in symbols $\mu \in \mathcal{D}_0(X)$) if

$$\sup_{x \in E} \int_E r_0(x, y) \mu(dy) < \infty, \quad (3.2)$$

where $r_0(x, y) = \lim_{\alpha \downarrow 0} r_\alpha(x, y)$.

Lemma 3.4. *Suppose $(\mathcal{E}, \mathcal{F})$ is transient. Let $p \in (1, \infty)$ and μ be a smooth measure in $S_1(X)$. Then, $\mu \in \mathcal{D}_0(X)$ and $\mu \in \mathcal{D}^p(\check{X})$ imply $\mu \in \mathcal{D}^p(X)$.*

Proof of Lemma 3.4. Denote $\check{r}_\alpha(x, y)$ the α -order resolvent kernel of \check{X} . First, we claim that for all $x \in F$,

$$\check{r}_0(x, y) = r_0(x, y) \quad \text{for } \mu\text{-a.e. } y \in F. \quad (3.3)$$

Take a sequence $\{E_n\}_{n=1}^\infty$ of Borel sets increasing to E as in the definition of $\mu \in S_1(X)$. Let f be a non-negative Borel function on F . By a similar argument as in the proof of Lemma 3.1, we have

$$\mathbb{E}_x \left[\int_0^\infty 1_{\bigcup_{n=1}^\infty E_n}(X_t) f(X_t) dA_t^\mu \right] = R_0[f \cdot \mu](x)$$

for all $x \in E$. The left-hand side of the above equation is equal to $\mathbb{E}_x \left[\int_0^\infty f(X_t) dA_t^\mu \right]$. Indeed, by the inclusion $\{t < \sigma_{E \setminus E_n}\} \subset \{X_t \in E_n \cup \{\partial\}\}$ we have

$$\begin{aligned} \mathbb{E}_x \left[\int_0^\infty f(X_t) dA_t^\mu \right] &= \lim_{n \rightarrow \infty} \mathbb{E}_x \left[\int_0^{\sigma_{E \setminus E_n}} f(X_t) dA_t^\mu \right] \\ &\leq \mathbb{E}_x \left[\int_0^\infty 1_{\bigcup_{n=1}^\infty E_n}(X_t) f(X_t) dA_t^\mu \right]. \end{aligned}$$

The converse inequality is trivial. Hence, (3.3) follows from the definition of 0-order resolvent

$$R_0[f \cdot \mu](x) = \int_F r_0(x, y) f(y) \mu(dy)$$

for $x \in E$ and from the equality

$$\mathbb{E}_x \left[\int_0^\infty f(X_t) dA_t^\mu \right] = \mathbb{E}_x \left[\int_0^\infty f(X_t^\mu) dt \right] = \int_F \check{r}_0(x, y) \mu(dy)$$

for $x \in F$ which is obtained from the change of variables.

Next, we recall the resolvent equation

$$\check{r}_0(x, y) = \check{r}_\alpha(x, y) + \alpha \int_F \check{r}_0(x, z) \check{r}_\alpha(z, y) \mu(dz), \quad \alpha > 0, x, y \in F.$$

A similar calculation as (2.7) gives that, for $x \in F$

$$\begin{aligned} &\left(\int_F \check{r}_0(x, y)^p \mu(dy) \right)^{\frac{1}{p}} \\ &\leq \left(1 + \alpha \sup_{x \in F} \int_F \check{r}_0(x, y) \mu(dy) \right) \left(\sup_{x \in F} \int_F \check{r}_\alpha(x, y)^p \mu(dy) \right)^{\frac{1}{p}}. \end{aligned}$$

By combining this with (3.3), we have

$$\begin{aligned} &\left(\sup_{x \in F} \int_F r_0(x, y)^p \mu(dy) \right)^{\frac{1}{p}} \\ &\leq \left(1 + \alpha \sup_{x \in E} \int_E r_0(x, y) \mu(dy) \right) \left(\sup_{x \in F} \int_F \check{r}_\alpha(x, y)^p \mu(dy) \right)^{\frac{1}{p}}. \end{aligned} \quad (3.4)$$

The right-hand side of (3.4) is finite because of the assumptions $\mu \in \mathcal{D}_0(X)$ and $\mu \in \mathcal{D}^p(\check{X})$.

Now we will show that

$$\sup_{x \in E} \int_F r_0(x, y)^p \mu(dy) = \sup_{z \in F} \int_F r_0(z, y)^p \mu(dy). \quad (3.5)$$

Let $x \in F$. Define the 0-order hitting distribution $H_F^0(x, dz)$ by

$$H_F^0(x, A) = \mathbb{E}_x[1_A(X_{\sigma_F}); \sigma_F < \infty], \quad \text{for } x \in E, \quad A \in \mathcal{B}(E).$$

Then we can see that

$$r_0(x, y) = \int_F r_0(z, y) H_F^0(x, dz) \quad \text{q.e. } y \in E.$$

Indeed, for a non-negative Borel function f on E we have from the strong Markov property,

$$\begin{aligned} & \int_E r_0(x, y) f(y) m(dy) \\ &= \mathbb{E}_x \left[\int_0^\infty f(X_t) dt \right] = \mathbb{E}_x \left[\int_{\sigma_F}^\infty f(X_t) dt \right] \\ &= \mathbb{E}_x [R_0 f(X_{\sigma_F}); \sigma_F < \infty] = \int_E \left(\int_F r_0(z, y) H_F^0(x, dz) \right) f(y) m(dy) \end{aligned}$$

and hence the equality holds for m -a.e. $x \in E$. The desired equality for q.e. $x \in E$ follows from the fact that the functions on both hand sides are 0-excessive in $y \in E$. By applying Hölder's inequality to the measure $H_F^0(x, dz)$, we have

$$\begin{aligned} \int_E r_0(x, y)^p \mu(dy) &= \int_F r_0(x, y)^{p-1} \left(\int_F r_0(z, y) H_F^0(x, dz) \right) \mu(dy) \\ &= \int_F \int_F r_0(x, y)^{p-1} r_0(z, y) \mu(dy) H_F^0(x, dz) \\ &\leq \int_F \left(\int_F r_0(x, y)^p \mu(dy) \right)^{\frac{p-1}{p}} \left(\int_F r_0(z, y)^p \mu(dy) \right)^{\frac{1}{p}} H_F^0(x, dz) \end{aligned}$$

and the right-hand side is bounded from above by

$$\left(\int_F r_0(x, y)^p \mu(dy) \right)^{\frac{p-1}{p}} \left(\sup_{z \in F} \int_F r_0(z, y)^p \mu(dy) \right)^{\frac{1}{p}}$$

because of $H_F^0(x, F) \leq 1$. Hence we obtain (3.5).

By combining (3.4) with (3.5), we have

$$\begin{aligned} & \left(\sup_{x \in E} \int_F r_\alpha(x, y)^p \mu(dy) \right)^{\frac{1}{p}} \\ & \leq \left(\sup_{x \in E} \int_F r_0(x, y)^p \mu(dy) \right)^{\frac{1}{p}} \\ & \leq \left(1 + \alpha \sup_{x \in F} \int_F r_0(x, y) \mu(dy) \right) \left(\sup_{x \in F} \int_F \tilde{r}_\alpha(x, y)^p \mu(dy) \right)^{\frac{1}{p}}, \end{aligned}$$

which completes the proof. \square

We now prove Proposition 3.2.

Proof of Proposition 3.2. Let $Y = X^{(1)}$ be the 1-subprocess of X defined in Remark 2.9. We can find that μ is in $S_1(Y)$. We can also find that the assumptions $\mu \in \mathcal{D}(X)$ and $\mu \in \mathcal{D}^p(\check{X})$ imply $\mu \in \mathcal{D}_0(Y)$ and $\mu \in \mathcal{D}^p(\check{Y})$. Since Y is transient, Lemma 3.4 gives that $\mu \in \mathcal{D}^p(Y)$. The conclusion follows from the equality $\mathcal{D}^p(X) = \mathcal{D}^p(Y)$, which has already been noted in Remark 2.9. \square

4 Main results

In this section, we give relations between the L^p -Dynkin classes and the Sobolev embeddings as we introduced in (1.3) and (1.4). We also give variants of such relations corresponding to the L^p -Kato class and the L^p -Kato class with order δ , respectively.

4.1 L^p -Kato implies the Sobolev embedding

In this section, we first give an L^p -version of the Stollmann-Voigt inequality and some variants in Theorem 4.1. As a consequence, Corollary 4.2 proves the assertion (1.3) introduced in Section 1, that is, for a measure $\mu \in \mathcal{D}^1(X)$ and $1 \leq p \leq p'$, $\mu \in \mathcal{D}^{p'}(X)$ implies that $(\mathcal{F}, \mathcal{E}_1)$ is continuously embedded into $L^{2p}(E; \mu)$. We also give a Rellich-Kondrachov type compact embedding theorem (Corollary 4.3).

Theorem 4.1. *Let $p \in [1, \infty)$ and $\mu \in \mathcal{D}^p(X)$.*

(i) *It holds that*

$$\|u\|_{L^{2p}(E; \mu)}^2 \leq \left(\sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy) \right)^{\frac{1}{p}} \mathcal{E}_\alpha(u, u) \quad (4.1)$$

for any $u \in \mathcal{F}$ and $\alpha > 0$. In particular, the Hilbert space $(\mathcal{F}, \mathcal{E}_1)$ is continuously embedded into $L^{2p}(E; \mu)$.

(ii) *If $\mu \in \mathcal{K}^p(X)$, then it holds that*

$$\|u\|_{L^{2p}(E; \mu)}^2 \leq \varepsilon \mathcal{E}_1(u, u) + K(\varepsilon) \|u\|_{L^2(E; m)}^2 \quad (4.2)$$

for any $u \in \mathcal{F}$ and $\varepsilon > 0$, where K is a positive function on $(0, \infty)$ satisfying $\varepsilon^{-1}K(\varepsilon) \uparrow \infty$ as $\varepsilon \downarrow 0$.

(iii) *If $\mu \in \mathcal{K}^{p, \theta}(X)$ for some $\theta \in (0, 1]$, then (4.2) holds for a function $K(\varepsilon) = A\varepsilon^{-\frac{1-\theta}{\theta}}$, where A is a positive constant.*

In particular, it holds that

$$\|u\|_{L^{2p}(E; \mu)} \leq B \sqrt{\mathcal{E}_1(u, u)^{(1-\theta)}} \|u\|_{L^2(E; m)}^\theta \quad (4.3)$$

for any $u \in \mathcal{F}$, where B is another positive constant.

Note that (4.2) is similar to the notion of compactly boundedness in [29], and that (4.3) is so-called the interpolation type inequality.

Proof of Theorem 4.1. We first assume $\mu \in \mathcal{D}^p(X)$ and prove (4.1). The case $p = 1$ is exactly the Stollmann-Voigt inequality (see [24, 25]), so we assume $p > 1$. By the regularity of $(\mathcal{E}, \mathcal{F})$, it suffices to prove (4.1) for $u \in \mathcal{F} \cap C_0(E)$, where $C_0(E)$ is the set of continuous functions on E with compact support. Fix $u \in \mathcal{F} \cap C_0(E)$ and $\alpha > 0$. Define a finite measure ν on E by $\nu(dy) = u^{2p-2}(y)\mu(dy)$. By Hölder's inequality, we have for $x \in E$,

$$R_\alpha \nu(x) = \int_E r_\alpha(x, y) u^{2p-2}(y) \mu(dy) \leq \left(\sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy) \right)^{\frac{1}{p}} \left(\int_E u^{2p} d\mu \right)^{\frac{p-1}{p}}.$$

By applying the inequality (4.1) with $p = 1$ and $\nu \in S_1(X)$, we have

$$\begin{aligned} \int_E u^{2p} d\mu &= \int_E u^2 d\nu \\ &\leq \|R_\alpha \nu\|_\infty \mathcal{E}_\alpha(u, u) \\ &\leq \left(\sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy) \right)^{\frac{1}{p}} \left(\int_E u^{2p} d\mu \right)^{\frac{p-1}{p}} \mathcal{E}_\alpha(u, u), \end{aligned}$$

which concludes (4.1).

We next assume $\mu \in \mathcal{K}^p(X)$ and prove (4.2). Recall the notation $\gamma(\alpha)$ introduced in (2.11). We may assume $\gamma(\alpha) > 0$ for all $\alpha > 0$. Indeed, if $\gamma(\alpha_0) = 0$ for some $\alpha_0 > 0$, then $R_{\alpha_0}\mu(x) = 0$ for all $\alpha \geq \alpha_0$ and $x \in E$, and hence $\mu = 0$. In this case (4.2) clearly holds for $K(\varepsilon) \equiv 1$.

Note that the function $\gamma(\cdot)$ is continuous. Indeed, for $0 < \alpha_0 < \alpha < \beta$, the monotonicity of $\gamma(\cdot)$ and (2.7) imply that

$$0 \leq \gamma(\alpha) - \gamma(\beta) \leq \frac{\beta}{\alpha} \gamma(\beta) - \gamma(\beta) = \frac{\gamma(\beta)}{\alpha} (\beta - \alpha) \leq \frac{\gamma(\alpha_0)}{\alpha_0} (\beta - \alpha).$$

Define the right continuous inverse γ^{-1} of γ by

$$\gamma^{-1}(\varepsilon) := \inf\{\alpha > 0 : \gamma(\alpha) \leq \varepsilon\}, \quad \varepsilon > 0.$$

We have $\gamma^{-1}(\varepsilon) \uparrow \infty$ as $\varepsilon \downarrow 0$ because of the assumption $\mu \in \mathcal{K}^p(X)$. We also have $\gamma(\gamma^{-1}(\varepsilon)) \leq \varepsilon$ because of the right continuity of γ^{-1} . By substituting $\alpha = \gamma^{-1}(\varepsilon)$ in (4.1), we obtain (4.2) with $K(\varepsilon) = \varepsilon \gamma^{-1}(\varepsilon)$.

Finally, we assume $\mu \in \mathcal{K}^{p,\theta}(X)$ for $\theta \in (0, 1]$ and prove (4.3). By the definition of $\mathcal{K}^{p,\theta}(X)$, there exists a constant $C > 0$ such that $\gamma(\alpha + 1) \leq C(\alpha + 1)^{-\theta} \leq C\alpha^{-\theta}$ for all $\alpha > 0$. Then, for $u \in \mathcal{F}$ we have from (4.1)

$$\|u\|_{L^{2p}(E;\mu)}^2 \leq \gamma(\alpha + 1) \mathcal{E}_{\alpha+1}(u, u) \leq C\alpha^{-\theta} \mathcal{E}_1(u, u) + C\alpha^{1-\theta} \|u\|_{L^2(E;m)}^2.$$

By substituting $\alpha = C^{\frac{1}{\theta}} \varepsilon^{-\frac{1}{\theta}}$ in the right-hand side of the above inequality, we obtain (4.2) with $K(\varepsilon) = C^{\frac{1}{\theta}} \varepsilon^{-\frac{1-\theta}{\theta}}$. The interpolation inequality (4.3) follows by taking infimum over $\varepsilon > 0$. \square

By combining Theorem 4.1 (i) with Hölder's inequality, we have the following corollary.

Corollary 4.2. *Let $p' \in [1, \infty)$. Then, for any measure $\mu \in \mathcal{D}^1(X) \cap \mathcal{D}^{p'}(X)$ the Hilbert space $(\mathcal{F}, \mathcal{E}_1)$ is continuously embedded into $L^{2p}(E; \mu)$ for all $1 \leq p \leq p'$.*

At the end of this section, we give a Rellich-Kondrachov type compact embedding theorem. The following corollary is a generalization of [27, Corollary 4.5], where the statement is proved for $p = 1$ and $\mu = m$.

Corollary 4.3. *Assume X satisfies*

- (resolvent strong Feller property) $R_1(\mathcal{B}_b(E)) \subset C_b(E)$, where $C_b(E)$ is the set of bounded continuous functions on E , and
- (tightness) for any $\varepsilon > 0$, there exists a compact set $K \subset E$ such that $\sup_{x \in E} R_1 1_{K^c}(x) < \varepsilon$.

Let $p \in [1, \infty)$ and suppose $\mu \in \mathcal{K}^p(X)$. Then the Hilbert space $(\mathcal{F}, \mathcal{E}_1)$ is compactly embedded into $L^2(E; m)$ and $L^{2p}(E; \mu)$.

Proof. Suppose $\{u_n\}_{n=1}^\infty \subset \mathcal{F}$ is bounded in $(\mathcal{F}, \mathcal{E}_1)$. By [27, Corollary 4.5], $(\mathcal{F}, \mathcal{E}_1)$ is compactly embedded into $L^2(E; m)$. (We remark that the irreducibility assumption is not needed to prove [27, Corollary 4.5].) Hence we can take $u \in L^2(E; m)$ and a subsequence $\{u_{n(k)}\}_{k=1}^\infty$ such that $u_{n(k)}$ converges to u in $L^2(E; m)$ as $k \rightarrow \infty$.

Recall the notation $\gamma(\alpha)$ introduced in (2.11). By (4.1), we have

$$\begin{aligned} & \|u_{n(k)} - u_{n(l)}\|_{L^{2p}(E;\mu)}^2 \\ & \leq \gamma(\alpha) \|u_{n(k)} - u_{n(l)}\|_{\mathcal{E}_\alpha}^2 \\ & = \gamma(\alpha) \mathcal{E}(u_{n(k)} - u_{n(l)}, u_{n(k)} - u_{n(l)}) + \alpha \gamma(\alpha) \|u_{n(k)} - u_{n(l)}\|_{L^2(E;m)}^2 \\ & \leq 4\gamma(\alpha) \sup_n \|u_n\|_{\mathcal{E}_1}^2 + \alpha \gamma(\alpha) \|u_{n(k)} - u_{n(l)}\|_{L^2(E;m)}^2. \end{aligned}$$

By letting $k, l \rightarrow \infty$, and then $\alpha \rightarrow \infty$, we find that $\{u_{n(k)}\}$ is Cauchy in $L^{2p}(E; \mu)$, and this completes the proof. \square

Remark 4.4. By Corollary 2.8, assumption (A5) of [21] is equivalent to the reference measure m belonging to the L^p -Kato class. We can also see that assumption (A4) of [21] is used only to show the fact that $(\mathcal{F}, \mathcal{E}_1)$ is compactly embedded into $L^{2p}(E; m)$ in the proof of Proposition 3.2 of [21]. Hence, by Corollary 4.3 we may drop the assumption (A4) in [21].

Example 4.5 (Killed Brownian motion in a domain D). Suppose $D \subset \mathbb{R}^d$ be a domain with a smooth boundary satisfying

$$\lim_{x \in D, |x| \rightarrow \infty} m(D \cap B(x, 1)) = 0,$$

where m is the Lebesgue measure on \mathbb{R}^d . Let ∂ be a point added to D so that $D_\partial := D \cup \{\partial\}$ is the one-point compactification of D . A killed Brownian motion X in D is the process given by

$$X_t = \begin{cases} B_t, & t < \tau_D, \\ \partial, & t \geq \tau_D, \end{cases}$$

where B is a Brownian motion on \mathbb{R}^d and $\tau_D = \inf\{t > 0 : B_t \notin D\}$ is the exit time of B from D . Its Dirichlet form is $(\frac{1}{2}D, H_0^1(D))$, where $H_0^1(D)$ is the Sobolev space with zero boundary values. It is known that X satisfies the resolvent strong Feller property and the tightness property (see [28, Lemma 3.3] for example).

Hence, by combining Example 2.13 with Corollary 4.3, we can see that $H_0^1(D)$ is compactly embedded into $L^{2p}(D)$ for $p \in [1, \infty)$ with $d - p(d - 2) > 0$. This is exactly the classical Rellich-Kondrachov embedding theorem.

4.2 Sobolev embedding implies L^p -Kato

In this section, we prove (1.4) introduced in Section 1, that is, for a measure $\mu \in \mathcal{D}^1(X)$ and $1 \leq p < p'$, if $(\mathcal{F}, \mathcal{E}_1)$ is continuously embedded in $L^{2p'}(E; \mu)$, then $\mu \in \mathcal{D}^p(X)$. Furthermore, we give variants of (1.4) which give a sufficient condition for μ belonging to $\mathcal{K}^p(X)$ or $\mathcal{K}^{p, \delta}(X)$ for a suitable δ .

We first consider the case $\mu = m$ and consider the assertion that \mathcal{E}_1 is replaced by \mathcal{E} :

Lemma 4.6. *Let $p' \in (1, \infty)$ and suppose that the following Sobolev type inequality holds: there exists a constant $S > 0$ such that*

$$\|u\|_{L^{2p'}(E; m)}^2 \leq S\mathcal{E}(u, u) \quad \text{for all } u \in \mathcal{F}. \quad (4.4)$$

Then $m \in \mathcal{K}^{p, \delta}(X)$ for any $p \in [1, p')$ with $\delta = 1 - \frac{p'}{p'-1} \frac{p-1}{p}$.

Proof. First, by [30], (4.4) implies the ultra-contractivity, that is, there exists $C > 0$ such that

$$\|P_t\|_{L^1 \rightarrow L^\infty} \leq Ct^{-\frac{p'}{p'-1}} \quad \text{for all } t > 0,$$

where $\|\cdot\|_{L^q \rightarrow L^r}$ is the operator norm from $L^q(E; m)$ to $L^r(E; m)$. By Jensen's inequality, we then have

$$\|P_t\|_{L^{\frac{p}{p-1}} \rightarrow L^\infty} \leq C^{\frac{p-1}{p}} t^{-\frac{p'}{p'-1} \frac{p-1}{p}} \quad \text{for all } t > 0,$$

when $p \in (1, p')$. Since the transition kernel P_t is L^∞ -contractive, the above inequality also holds when $p = 1$, where we use the convention $1/0 = \infty$.

Fix $\alpha > 0$. For any non-negative Borel function f with $f \in L^1(E; m) \cap L^\infty(E; m)$, we have

$$\begin{aligned} R_\alpha f(x) &= \int_0^\infty e^{-\alpha t} P_t f(x) dt \\ &\leq \int_0^\infty e^{-\alpha t} \|P_t f\|_{L^\infty(E; m)} dt \\ &\leq C^{\frac{p-1}{p}} \|f\|_{L^{\frac{p}{p-1}}(E; m)} \int_0^\infty e^{-\alpha t} t^{-\frac{p'}{p'-1} \frac{p-1}{p}} dt \end{aligned}$$

$$= C \frac{p-1}{p} \|f\|_{L^{\frac{p}{p-1}}(E;m)} \Gamma(\delta) \alpha^{-\delta}$$

for m -a.e. $x \in E$, where Γ is the Gamma function and $\delta = 1 - \frac{p'}{p'-1} \frac{p-1}{p} > 0$. Since $R_\alpha f = \int_E r_\alpha(\cdot, y) f(y) m(dy)$ is α -excessive and the absolute continuity condition (2.1) holds, we have for every $x \in E$,

$$\begin{aligned} R_\alpha f(x) &= \lim_{\varepsilon \downarrow 0} e^{-\alpha\varepsilon} \mathbb{E}_x[R_\alpha f(X_\varepsilon)] = \lim_{\varepsilon \downarrow 0} e^{-\alpha\varepsilon} \int_E p_\varepsilon(x, y) R_\alpha f(y) m(dy) \\ &\leq C \frac{p-1}{p} \|f\|_{L^{\frac{p}{p-1}}(E;m)} \Gamma(\delta) \alpha^{-\delta}. \end{aligned} \quad (4.5)$$

Now, fix $x \in E$, $M > 0$ and a compact set $K \subset E$. By applying (4.5) with $f = 1_K(\cdot)(r_\alpha(x, \cdot) \wedge M)^{p-1}$, we have

$$\begin{aligned} \int_K (r_\alpha(x, y) \wedge M)^p m(dy) &\leq \int_K (r_\alpha(x, y) \wedge M)^{p-1} r_\alpha(x, y) m(dy) \\ &\leq C \frac{p-1}{p} \left(\int_K (r_\alpha(x, y) \wedge M)^p m(dy) \right)^{\frac{p-1}{p}} \Gamma(\delta) \alpha^{-\delta}, \end{aligned}$$

which means that

$$\left(\int_K (r_\alpha(x, y) \wedge M)^p m(dy) \right)^{\frac{1}{p}} \leq C \frac{p-1}{p} \Gamma(\delta) \alpha^{-\delta}.$$

Let $M \uparrow \infty$ and $K \uparrow E$. Hence we have the conclusion $m \in \mathcal{K}^{p,\delta}(X)$ by the dominated convergence theorem. \square

The following is our second main theorem, which states the opposite of Theorem 4.1 in some sense.

Theorem 4.7. *Let $p' \in (1, \infty)$ and let μ be a measure in $\mathcal{D}^1(X)$.*

(i) *Suppose the following Sobolev type inequality holds: there exists a constant $S > 0$ such that*

$$\|u\|_{L^{2p'}(E;\mu)}^2 \leq S \mathcal{E}_1(u, u) \quad (4.6)$$

for all $u \in \mathcal{F}$. Then $\mu \in \mathcal{D}^p(X)$ for any $p \in [1, p']$.

(ii) *Suppose the following Sobolev type inequality holds: there exists a function $K : (0, \infty) \rightarrow (0, \infty)$ with $\varepsilon^{-1}K(\varepsilon) \uparrow \infty$ as $\varepsilon \downarrow 0$, such that*

$$\|u\|_{L^{2p'}(E;\mu)}^2 \leq \varepsilon \mathcal{E}_1(u, u) + K(\varepsilon) \|u\|_{L^2(E;m)}^2 \quad (4.7)$$

for any $u \in \mathcal{F}$ and $\varepsilon > 0$. Then $\mu \in \mathcal{K}^p(X)$ for any $p \in (1, p']$.

(iii) *Suppose the following Sobolev type inequality holds: there exist constants $A > 0$ and $\theta \in (0, 1]$ such that*

$$\|u\|_{L^{2p'}(E;\mu)} \leq A \sqrt{\mathcal{E}_1(u, u)}^{(1-\theta)} \|u\|_{L^2(E;m)}^\theta \quad (4.8)$$

for any $u \in \mathcal{F}$. Then $\mu \in \mathcal{K}^{p,\theta(1-\delta)}(X)$ for any $p \in (1, p')$ with $\delta = 1 - \frac{p'}{p'-1} \frac{p-1}{p}$.

Remark 4.8.

1. In claim (i), when $p = p'$ the result does not hold in general. See Example 4.9 below.
2. Regarding claims (ii) and (iii), our proofs do not work when $p = 1$ and $p = p'$, but there are examples that support that the conclusion also holds for such p . Indeed, in [1, Theorem 4.9], Aizenman and Simon proved that for a Brownian motion on \mathbb{R}^d , the inequality of the type (4.7) with $p' = 1$ and $K(\varepsilon) = A \exp\{e^{B\varepsilon^{-\alpha}}\}$ implies that μ belongs to $\mathcal{K}^1(X)$. For the case $1 < p = p'$, see Example 4.9 below.

3. Regarding claim (iii), it can be seen that the exponent $\theta(1 - \delta)$ is appropriate by the following direct calculations: Let $q > 1$, $\rho \in (0, 1]$ and $\mu \in \mathcal{K}^{q,\rho}(X)$. For $1 < p < p' < q$, set $\theta' = \frac{q}{q-1} \frac{p'-1}{p'}$ and $\delta = 1 - \frac{p'}{p'-1} \frac{p-1}{p}$. Note that $\frac{1}{p'} = 1 - \theta' + \frac{\theta'}{q}$ and $\frac{1}{p} = \delta + \frac{1-\delta}{p'}$. We also set $\theta = \rho\theta'$. Then, Hölder's inequality and Theorem 4.1 (i) and (iii) imply that

$$\begin{aligned} \|u\|_{L^{2p'}(E;\mu)} &\leq \|u\|_{L^2(E;\mu)}^{1-\theta'} \|u\|_{L^{2q}(E;\mu)}^{\theta'} \\ &\leq \left(\gamma(1)\sqrt{\mathcal{E}_1(u, u)}\right)^{1-\theta'} \left(B\sqrt{\mathcal{E}_1(u, u)}^{1-\rho} \|u\|_{L^2(E;m)}^\rho\right)^{\theta'} \\ &= \gamma(1)^{1-\theta'} B^{\theta'} \sqrt{\mathcal{E}_1(u, u)}^{1-\theta} \|u\|_{L^2(E;m)}^\theta, \end{aligned}$$

that is, the assumption (4.9) holds. On the other hand, Hölder's inequality also implies that

$$\begin{aligned} &\left(\sup_{x \in E} \int_E r_\alpha(x, y)^p \mu(dy)\right)^{\frac{1}{p}} \\ &\leq \left(\sup_{x \in E} \int_E r_\alpha(x, y) \mu(dy)\right)^\delta \left(\sup_{x \in E} \int_E r_\alpha(x, y)^{p'} \mu(dy)\right)^{\frac{1-\delta}{p'}} \\ &\leq \left(\sup_{x \in E} \int_E r_\alpha(x, y) \mu(dy)\right)^{\delta+(1-\theta')(1-\delta)} \left(\sup_{x \in E} \int_E r_\alpha(x, y)^q \mu(dy)\right)^{\frac{\theta'(1-\delta)}{q}} \\ &= O(\alpha^{-\rho\theta'(1-\delta)}) \end{aligned}$$

as $\alpha \rightarrow \infty$, that is, the conclusion $\mu \in \mathcal{K}^{p,\theta(1-\delta)}(X)$ holds.

Proof of Theorem 4.7. We first prove (i). Let $X^{(1)}$ be the 1-subprocess of X . As we have seen in the proof of Proposition 3.2, it holds that $\mu \in \mathcal{S}_1(X^{(1)})$. We also have $\mu \in \mathcal{D}^1(X^{(1)})$.

Denote Y the time changed process of $X^{(1)}$ with respect to the PCAF of $X^{(1)}$ with Revuz measure μ and denote its Dirichlet form as $(\mathcal{E}^Y, \mathcal{F}^Y)$. Since the Dirichlet form of $X^{(1)}$ is $(\mathcal{E}_1, \mathcal{F})$, (4.6) implies that

$$\|u\|_{L^{2p'}(\mu)}^2 \leq S\mathcal{E}^Y(u, u) \quad \text{for all } u \in \mathcal{F}^Y.$$

Let $p \in [1, p')$. By Lemma 4.6 we have $\mu \in \mathcal{D}^p(Y)$, and then by Proposition 3.2 we have $\mu \in \mathcal{D}^p(X^{(1)})$. The conclusion follows from Remark 2.9.

We next prove (ii). Let $\beta > 0$ and denote Y the time changed process of the β -subprocess $X^{(\beta)}$ with respect to the PCAF of $X^{(\beta)}$ with Revuz measure μ . The α -order resolvent of Y can be written as

$$R_{\alpha,A}^\beta f(x) = \mathbb{E}_x \left[\int_0^\infty e^{-\alpha A_t^\mu - \beta t} f(X_t) dA_t^\mu \right], \quad \alpha > 0, x \in F.$$

By Lemma 3.1, Y satisfies the absolute continuity condition (2.1). Denote the α -order resolvent kernel of Y as $r_\alpha^\beta(x, y)$. Note that for all $x \in F$ and $\alpha, \beta > 0$, it holds that

$$r_\beta(x, y) = r_\alpha^\beta(x, y) + \alpha \int_F r_\beta(x, z) r_\alpha^\beta(z, y) \mu(dz) \quad \text{for } \mu\text{-a.e. } y \in F$$

(see [12, (6.4.11)] for example). By an argument similar to that leading to (2.7), we have

$$\begin{aligned} &\left(\sup_{x \in F} \int_E r_\beta(x, y)^p \mu(dy)\right)^{\frac{1}{p}} \\ &\leq \left(1 + \alpha \sup_{x \in E} \int_E r_\beta(x, y) \mu(dy)\right) \left(\sup_{z \in F} \int_F r_\alpha^\beta(z, y)^p \mu(dy)\right)^{\frac{1}{p}}. \end{aligned} \tag{4.9}$$

By a similar argument as to obtain (3.5), we also have

$$\sup_{x \in E} \int_E r_\beta(x, y)^p \mu(dy) = \sup_{x \in F} \int_E r_\beta(x, y)^p \mu(dy). \tag{4.10}$$

Define the process Z by $Z_t := Y_{\varepsilon^{p'}t}$. Z is an $\varepsilon^{-p'}$ -symmetric Hunt process on F , and its α -order resolvent kernel can be written as $r_{\varepsilon^{-p'}\alpha}^\beta(x, y)$. Denote by $(\mathcal{E}^Z, \mathcal{F}^Z)$ the associated Dirichlet form on $L^2(F; \varepsilon^{-p'}\mu)$.

Now, by setting $\beta = \varepsilon^{-1}K(\varepsilon)$, (4.7) implies that

$$\|u\|_{L^{2p'}(E; \varepsilon^{-p'}\mu)}^2 \leq \mathcal{E}^Z(u, u) \quad \text{for all } u \in \mathcal{F}^Z.$$

By the same argument as the proof of Lemma 4.6, there exists a constant $C > 0$, which is independent of ε and β by virtue of [30], such that

$$\left(\sup_{x \in F} \int_F r_{\varepsilon^{-p'}\alpha}^\beta(x, y)^p \varepsilon^{-p'}\mu(dy) \right)^{\frac{1}{p}} \leq C^{\frac{p-1}{p}} \Gamma(\delta) \alpha^{-\delta}$$

for all $\alpha > 0$. In particular, by substituting $\alpha = \varepsilon^{p'}$, we have

$$\left(\sup_{x \in F} \int_F r_1^\beta(x, y)^p \mu(dy) \right)^{\frac{1}{p}} \leq C^{\frac{p-1}{p}} \Gamma(\delta) \varepsilon^{\frac{p'}{p} - p'\delta} = C^{\frac{p-1}{p}} \Gamma(\delta) \varepsilon^{1-\delta},$$

where we used the relation $\frac{1}{p} = \delta + \frac{1-\delta}{p'}$.

Take $\varepsilon > 0$ sufficiently small so that $\beta > 1$. By combining the above inequality with (4.9), (4.10) and the monotonicity $r_\beta(x, y) \leq r_1(x, y)$, we have

$$\left(\sup_{x \in E} \int_E r_\beta(x, y)^p \mu(dy) \right)^{\frac{1}{p}} \leq \left(1 + \sup_{x \in E} \int_E r_1(x, y) \mu(dy) \right) C^{\frac{p-1}{p}} \Gamma(\delta) \varepsilon^{1-\delta}, \quad (4.11)$$

hence we obtain the desired conclusion $\mu \in \mathcal{K}^p(X)$.

We finally prove (iii). By the assumption (4.8), It holds that the inequality (4.7) with $K(\varepsilon) = A^{\frac{2}{\theta}} \varepsilon^{-\frac{1-\theta}{\theta}}$. By setting $\beta = \varepsilon^{-1}K(\varepsilon) = A^{\frac{2}{\theta}} \varepsilon^{-\frac{1}{\theta}}$, we have $\varepsilon = A^2 \beta^{-\theta}$ and then by (4.11)

$$\begin{aligned} & \left(\sup_{x \in E} \int_E r_\beta(x, y)^p \mu(dy) \right)^{\frac{1}{p}} \\ & \leq \left(1 + \sup_{x \in E} \int_E r_1(x, y) \mu(dy) \right) C^{\frac{p-1}{p}} \Gamma(\delta) A^{2(1-\delta)} \beta^{-\theta(1-\delta)}. \end{aligned}$$

Therefore we have the desired conclusion $\mu \in \mathcal{K}^{p, \theta(1-\delta)}(X)$. \square

Example 4.9 (Brownian motion). We continue with Example 2.13. As in Section 1, the classical Sobolev embedding theorem on \mathbb{R}^d gives that $H^1(\mathbb{R}^d)$ is continuously embedded into $L^{2p}(\mathbb{R}^d)$ for $p \in [1, \infty)$ with $d - p(d - 2) \geq 0$. By combining this with Theorem 4.7 (i), the Lebesgue measure on \mathbb{R}^d is in $\mathcal{D}^p(X)$ for $p \in [1, \infty)$ with $d - p(d - 2) > 0$.

In particular, when $d \geq 3$, by setting $p^* = d/(d - 2)$ the critical Sobolev embedding theorem gives that $H^1(\mathbb{R}^d)$ is continuously embedded into $L^{2p}(\mathbb{R}^d)$ for $p \in [1, p^*)$, but the Lebesgue measure on \mathbb{R}^d does not belong to $\mathcal{D}^{p^*}(X)$. This is an example of Remark 4.8.1.

We now assume $d \geq 1$. We also assume that $d - p'(d - 2) \geq 0$ and $1 < p < p'$. The classical Gagliardo-Nirenberg interpolation inequality says that, by setting $\theta = 1 - \frac{d(p'-1)}{2p'} = \frac{d-p'(d-2)}{2p'}$, there exists a positive constant C such that

$$\|u\|_{L^{2p'}(\mathbb{R}^d)} \leq C \|\nabla u\|_{L^2(\mathbb{R}^d)}^{1-\theta} \|u\|_{L^2(\mathbb{R}^d)}^\theta$$

holds for all $u \in H^1(\mathbb{R}^d)$. By applying Theorem 4.7 (iii) and then by letting $p' \downarrow p$, we can see that the Lebesgue measure on \mathbb{R}^d belongs to $\mathcal{K}^{p, \frac{d-p(d-2)}{2p} - \varepsilon}(X)$ for all $\varepsilon \in (0, \frac{d-p(d-2)}{2p})$, which is a slightly weaker condition than the conclusion of Example 2.13. By combining with the Gagliardo-Nirenberg inequality, we can also see that the fact that the Lebesgue measure on \mathbb{R}^d belongs to $\mathcal{K}^{p, \frac{d-p(d-2)}{2p}}(X)$, which is an example of Remark 4.8.2 for $1 < p = p'$.

5 Application: Continuity of the intersection measure in time

In this section, we give an application of the L^p -Kato class to the continuity of the intersection measure in time. Throughout this section, we assume that $p \geq 2$ is an integer and that the reference measure m is in the L^p -Dynkin class. Let $X^{(1)}, \dots, X^{(p)}$ be independent Hunt processes with the same law as X . We write $\zeta^{(1)}, \dots, \zeta^{(p)}$ as their life times and write $x_0^{(1)}, \dots, x_0^{(p)}$ as their starting points, respectively.

First, we review the construction of the intersection measure. For detail, see [7, 21] for example. Fix bounded Borel sets $J^{(1)}, \dots, J^{(p)} \subset [0, \infty)$ and write $J = \prod_{i=1}^p J^{(i)}$. For each $\varepsilon > 0$, we define the approximated (mutual) intersection measure $\ell_{J, \varepsilon}^{\text{IS}}$ of $X^{(1)}, \dots, X^{(p)}$ with respect to the (multi-parameter) time interval J by

$$\langle \ell_{J, \varepsilon}^{\text{IS}}, f \rangle = \int_E f(x) \left[\prod_{i=1}^p \int_{J^{(i)}} p_\varepsilon(x, X_s^{(i)}) ds \right] m(dy)$$

for $f \in \mathcal{B}_b(E)$, where, for convenience we regard $p_\varepsilon(x, X_s^{(i)}) = 0$ when $s \geq \zeta^{(i)}$. Then, there exists a random measure ℓ_J^{IS} on E such that, $\ell_{J, \varepsilon}^{\text{IS}}$ converges vaguely to ℓ_J^{IS} in $\mathcal{M}(E)$ and that

$$\lim_{\varepsilon \rightarrow 0} \mathbb{E} [|\langle f, \ell_{J, \varepsilon}^{\text{IS}} \rangle - \langle f, \ell_J^{\text{IS}} \rangle|^k] = 0$$

for any integer $k \geq 1$ and $f \in C_0(E)$, where $\mathcal{M}(E)$ is the set of Radon measures on E equipped with the vague topology. We call the limit ℓ_J^{IS} as the (mutual) intersection measure of $X^{(1)}, \dots, X^{(p)}$ with respect to J . For $\mathbf{t} = (t_1, \dots, t_p) \in [0, \infty)^p$ we simply denote the approximated intersection measure and the intersection measure with respect to $[\mathbf{0}, \mathbf{t}] := \prod_{i=1}^p [0, t_i]$ as $\ell_{\mathbf{t}, \varepsilon}^{\text{IS}}$ and $\ell_{\mathbf{t}}^{\text{IS}}$, respectively.

The intersection measure ℓ_J^{IS} enjoys the so-called Le Gall's moment formula: for any $f \in \mathcal{B}_b(E)$ with compact support and for any integer $k \geq 1$, it holds that

$$\begin{aligned} & \mathbb{E} [\langle f, \ell_J^{\text{IS}} \rangle^k] \\ &= \int_{E^k} f(x_1) \cdots f(x_k) \prod_{i=1}^p \left\{ \sum_{\sigma \in \mathfrak{S}_k} \int_{(J^{(i)})^k} \prod_{j=1}^k p_{s_j - s_{j-1}}(x_{\sigma(j-1)}, x_{\sigma(j)}) \right. \\ & \quad \left. ds_1 \cdots ds_k \right\} m(dx_1) \cdots m(dx_k), \end{aligned} \tag{5.1}$$

where $(J^{(i)})^k_{<} := \{(s_1, \dots, s_k) \in (J^{(i)})^k; s_1 < \dots < s_k\}$ and \mathfrak{S}_k is the set of permutations of $\{1, \dots, k\}$. For convenience we set $\sigma(0) = 0$ for $\sigma \in \mathfrak{S}_k$ and set $x_0 = x_0^{(i)}$.

The goal of this section is to realize $\{\ell_{\mathbf{t}}^{\text{IS}} : \mathbf{t} \in [0, \infty)^p\}$ as a measure-valued continuous stochastic process. That is,

Theorem 5.1. *Assume $m \in \mathcal{K}^{p, \delta}(X)$ for some $\delta \in (0, 1]$. Then it holds that*

- (i) *the $\mathcal{M}(E)$ -valued process $\{\ell_{\mathbf{t}}^{\text{IS}} : \mathbf{t} \in [0, \infty)^p\}$ has a continuous modification,*
- (ii) *for any $f \in \mathcal{B}_b(E)$, the real-valued process $\{\langle f, \ell_{\mathbf{t}}^{\text{IS}} \rangle : \mathbf{t} \in [0, \infty)^p\}$ has a modification whose paths are locally γ -Hölder continuous of every order $\gamma \in (0, \delta)$.*

Proof. Our proof is based on that of [7, Lemma 2.2.4]. First, we claim the following estimate: for any integer $k \geq 1$, positive constant $T > 0$ and $\mathbf{s}, \mathbf{t} \in [0, T]^p$, it holds that

$$\mathbb{E} [|\langle f, \ell_{\mathbf{t}}^{\text{IS}} \rangle - \langle f, \ell_{\mathbf{s}}^{\text{IS}} \rangle|^k] \leq (k!)^p \left(2^p \|f\|_\infty (\eta(T) + 1)^p \sup_{0 < t \leq pT} \{t^{-\delta} \eta(t)\} \right)^k |\mathbf{t} - \mathbf{s}|^{\delta k}, \tag{5.2}$$

where $\eta(T)$ is introduced in (2.12) and $|\mathbf{t} - \mathbf{s}|$ is the Euclidean distance between \mathbf{t} and \mathbf{s} in $[0, \infty)^p$.

Fix $\mathbf{s}, \mathbf{t} \in [0, T]^p$. Define the family $\{J_l\}_{l=1}^{2^p-1}$ consisting of products of intervals as

$$\begin{aligned} & \{J_l\}_{l=1}^{2^p-1} \\ &= \left\{ \prod_{i=1}^p [a_i, b_i] \mid [a_i, b_i] = [0, s_i \wedge t_i] \text{ or } [s_i \wedge t_i, s_i \vee t_i], 1 \leq i \leq p \right\} \setminus \left\{ \prod_{i=1}^p [0, s_i \wedge t_i] \right\}. \end{aligned}$$

Then we have $[\mathbf{0}, \mathbf{t}] \Delta [\mathbf{0}, \mathbf{s}] \subset \bigcup_l J_l$ and then

$$|\langle f, \ell_{\mathbf{t}}^{\text{IS}} \rangle - \langle f, \ell_{\mathbf{s}}^{\text{IS}} \rangle| \leq \sum_{l=1}^{2^p-1} \langle |f|, \ell_{J_l}^{\text{IS}} \rangle, \quad \mathbb{P}\text{-a.s.}$$

By Le Gall's moment formula, we have

$$\begin{aligned} & \mathbb{E}[\langle |f|, \ell_{\varepsilon, J_l}^{\text{IS}} \rangle^k] \\ &= \int_{E^k} |f|(x_1) \cdots |f|(x_k) \prod_{i=1}^p \left\{ \sum_{\sigma \in \mathfrak{S}_k} H_{a_i, b_i}^{(i)}(x_{\sigma(1)}, \dots, x_{\sigma(k)}) \right\} m(dx_1) \cdots m(dx_k), \end{aligned}$$

where

$$\begin{aligned} & H_{a_i, b_i}^{(i)}(x_1, \dots, x_p) \\ &:= \int_E \int_{[0, \infty)^p} \mathbf{1}_{\{\sum_{j=1}^k r_j \leq b_i - a_i\}} \prod_{j=1}^k p_{r_j}(x_{j-1}, x_j) dr_1 \cdots dr_k \nu_{a_i}^{(i)}(dx_0) \end{aligned}$$

and $\nu_{a_i}^{(i)}(dx_0) = \mathbb{P}_{x_0^{(i)}}(X_{a_i}^{(i)} \in dx_0)$. Note that

$$H_{a_i, b_i}^{(i)}(x_1, \dots, x_p) \leq \int_E \left[\prod_{j=1}^k \int_0^{b_i - a_i} p_s(x_{j-1}, x_j) ds \right] \nu_{a_i}^{(i)}(dx_0)$$

and then $\|H_{a_i, b_i}^{(i)}\|_{L^p(E^k, m^{\otimes k})} \leq \eta(b_i - a_i)^k$. We have by Hölder's inequality

$$\begin{aligned} \mathbb{E}[\langle |f|, \ell_{\varepsilon, J_l}^{\text{IS}} \rangle^k] &\leq (k!)^p \|f\|_{\infty}^k \prod_{i=1}^p \|H_{a_i, b_i}^{(i)}\|_{L^p(E^k, m^{\otimes k})} \\ &\leq (k!)^p \|f\|_{\infty}^k \prod_{i=1}^p \eta(b_i - a_i)^k \leq (k!)^p \|f\|_{\infty}^k \eta(T)^{(p-1)k} \eta(|\mathbf{t} - \mathbf{s}|)^k. \end{aligned}$$

In the last inequality, we used the fact that $[a_i, b_i] = [s_i \wedge t_i, s_i \vee t_i]$ holds for at least one i because of the definition of J_l . Therefore

$$\begin{aligned} \mathbb{E}[|\langle f, \ell_{\mathbf{t}}^{\text{IS}} \rangle - \langle f, \ell_{\mathbf{s}}^{\text{IS}} \rangle|^k] &\leq \left\{ \sum_{l=1}^{2^p-1} \mathbb{E}[\langle |f|, \ell_{J_l}^{\text{IS}} \rangle^k]^{1/k} \right\}^k \\ &\leq (k!)^p \left(2^p \|f\|_{\infty} (\eta(T) + 1)^p \eta(|\mathbf{t} - \mathbf{s}|) \right)^k. \end{aligned}$$

Taking account of Remark 2.12.1, the desired estimate (5.2) follows from $|\mathbf{t} - \mathbf{s}| \leq pT$ and the assumption $m \in \mathcal{K}^{p, \delta}(X)$.

Now, $(\delta k - p)/k$ increases to δ as $k \rightarrow \infty$. By applying Kolmogorov's continuity theorem (see [16, Theorem 3.23] for example) to (5.2), we can find that the process $[0, T]^p \ni \mathbf{t} \mapsto \langle f, \ell_{\mathbf{t}}^{\text{IS}} \rangle \in \mathbb{R}$ has a continuous modification whose paths are γ -Hölder continuous of every order $\gamma \in (0, \delta)$. Since $T > 0$ is arbitrary, we obtain (ii).

To prove (i), take a dense subset $\{\phi_n\}_{n=1}^{\infty}$ in $C_0^+(E)$, the family of non-negative continuous functions with compact support equipped with the uniform metric. Then $\mathcal{M}(E)$ is homeomorphic to a subset of \mathbb{R}^{∞} by the mapping $\mathcal{M}(E) \ni \mu \mapsto \{\langle \mu, \phi_n \rangle\}_{n=1}^{\infty} \in \mathbb{R}^{\infty}$, and hence the process $[0, \infty)^p \ni \mathbf{t} \mapsto \ell_{\mathbf{t}}^{\text{IS}} \in \mathcal{M}(E)$ has a continuous version by (ii). Therefore we complete the proof. \square

Acknowledgements

We would like to thank Professor Takashi Kumagai, Professor Masayoshi Takeda and an anonymous referee for helpful discussions and comments. We also wish to thank Professor Kazuhiro Kuwae for pointing out some mistakes in the first draft, which finally improves our results. This work was

supported by JSPS KAKENHI Grant Number JP18J21141.

References

- [1] M. Aizenman and B. Simon. Brownian motion and Harnack inequality for Schrödinger operators. *Comm. Pure Appl. Math.*, 35(2):209–273, 1982.
- [2] S. Albeverio, P. Blanchard, and Z. M. Ma. Feynman-Kac semigroups in terms of signed smooth measures. In *Random partial differential equations (Oberwolfach, 1989)*, volume 102 of *Internat. Ser. Numer. Math.*, pages 1–31. Birkhäuser, Basel, 1991.
- [3] D. Bakry, T. Coulhon, M. Ledoux, and L. Saloff-Coste. Sobolev inequalities in disguise. *Indiana Univ. Math. J.*, 44(4):1033–1074, 1995.
- [4] M. T. Barlow. Diffusions on fractals. In *Lectures on probability theory and statistics (Saint-Flour, 1995)*, volume 1690 of *Lecture Notes in Math.*, pages 1–121. Springer, Berlin, 1998.
- [5] M. T. Barlow and E. A. Perkins. Brownian motion on the Sierpiński gasket. *Probab. Theory Related Fields*, 79(4):543–623, 1988.
- [6] E. A. Carlen, S. Kusuoka, and D. W. Stroock. Upper bounds for symmetric Markov transition functions. *Ann. Inst. H. Poincaré Probab. Statist.*, 23(2, suppl.):245–287, 1987.
- [7] X. Chen. *Random walk intersections*, volume 157 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2010.
- [8] Z.-Q. Chen and T. Kumagai. Heat kernel estimates for stable-like processes on d -sets. *Stochastic Process. Appl.*, 108(1):27–62, 2003.
- [9] A. Dvoretzky, P. Erdős, and S. Kakutani. Double points of paths of Brownian motion in n -space. *Acta Sci. Math. Szeged*, 12(Leopoldo Fejer et Frederico Riesz LXX annos natis dedicatus, Pars B):75–81, 1950.
- [10] A. Dvoretzky, P. Erdős, and S. Kakutani. Multiple points of paths of Brownian motion in the plane. *Bull. Res. Council Israel*, 3:364–371, 1954.
- [11] A. Dvoretzky, P. Erdős, S. Kakutani, and S. J. Taylor. Triple points of Brownian paths in 3-space. *Proc. Cambridge Philos. Soc.*, 53:856–862, 1957.
- [12] M. Fukushima, Y. Oshima, and M. Takeda. *Dirichlet forms and symmetric Markov processes*, volume 19 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, extended edition, 2011.
- [13] M. Fukushima and T. Uemura. Capacitary bounds of measures and ultracontractivity of time changed processes. *J. Math. Pures Appl. (9)*, 82(5):553–572, 2003.
- [14] B. Güneysu. $RCD^*(K, N)$ spaces and the geometry of multi-particle Schrödinger semigroups. *Int. Math. Res. Not. IMRN*, 2020.
- [15] V. A. Kaimanovich. Dirichlet norms, capacities and generalized isoperimetric inequalities for Markov operators. *Potential Anal.*, 1(1):61–82, 1992.
- [16] O. Kallenberg. *Foundations of modern probability*. Probability and its Applications (New York). Springer-Verlag, New York, second edition, 2002.
- [17] W. König and C. Mukherjee. Large deviations for Brownian intersection measures. *Comm. Pure Appl. Math.*, 66(2):263–306, 2013.

- [18] K. Kuwae and M. Takahashi. Kato class measures of symmetric Markov processes under heat kernel estimates. *J. Funct. Anal.*, 250(1):86–113, 2007.
- [19] J.-F. Le Gall. Some properties of planar Brownian motion. In *École d’Été de Probabilités de Saint-Flour XX—1990*, volume 1527 of *Lecture Notes in Math.*, pages 111–235. Springer, Berlin, 1992.
- [20] P. Li and S.-T. Yau. On the parabolic kernel of the Schrödinger operator. *Acta Math.*, 156(3-4):153–201, 1986.
- [21] T. Mori. Large deviations for intersection measures of some Markov processes. *Math. Nachr.*, 293(3):533–553, 2020.
- [22] L. Saloff-Coste. *Aspects of Sobolev-type inequalities*, volume 289 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2002.
- [23] M. Schechter. *Spectra of partial differential operators*. North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1971. North-Holland Series in Applied Mathematics and Mechanics, Vol. 14.
- [24] Y. Shiozawa and M. Takeda. Variational formula for Dirichlet forms and estimates of principal eigenvalues for symmetric α -stable processes. *Potential Anal.*, 23(2):135–151, 2005.
- [25] P. Stollmann and J. Voigt. Perturbation of Dirichlet forms by measures. *Potential Anal.*, 5(2):109–138, 1996.
- [26] K.-T. Sturm. On the geometry of metric measure spaces. II. *Acta Math.*, 196(1):133–177, 2006.
- [27] M. Takeda. Compactness of symmetric Markov semigroups and boundedness of eigenfunctions. *Trans. Amer. Math. Soc.*, 372(6):3905–3920, 2019.
- [28] M. Takeda, Y. Tawara, and K. Tsuchida. Compactness of Markov and Schrödinger semi-groups: a probabilistic approach. *Osaka J. Math.*, 54(3):517–532, 2017.
- [29] N. S. Trudinger. Linear elliptic operators with measurable coefficients. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3)*, 27:265–308, 1973.
- [30] N. T. Varopoulos. Hardy-Littlewood theory for semigroups. *J. Funct. Anal.*, 63(2):240–260, 1985.
- [31] J. A. Yan. A formula for densities of transition functions. In *Séminaire de Probabilités, XXII*, volume 1321 of *Lecture Notes in Math.*, pages 92–100. Springer, Berlin, 1988.