

Composition Estimation via Shrinkage

Chong Gu
Purdue University

Abstract

In this note, we explore a simple approach to composition estimation, using penalized likelihood density estimation on a nominal discrete domain. Practical issues such as smoothing parameter selection and the use of prior information are investigated in simulations, and a theoretical analysis is attempted. The method has been implemented in a pair of R functions for use by practitioners.

Keywords: Cross-validation; Density estimation; Shrinkage.

1 Introduction

A composition refers to the proportions of a set of parts that make up a whole. For example, the relative abundance of bacterial genera in some microbiome can be represented by a vector of proportions summing up to 1. The analysis of compositional data (Aitchison 1986) found applications in many fields including metagenomics (cf. Xing, Liu, and Zhong 2017). Our task here is to estimate composition using empirical data.

Cao, Zhang, and Li (2020) had excellent discussions of the challenges posed by composition estimation and the many approaches attempted in the literature. Samples of microbiome are probed for the number of occurrences of bacterial genera, say, and a central issue is the prevalence of zero counts due to limitations of the probing technologies. Zero proportions are not among acceptable answers, and the challenge is to estimate the proportions associated with zero counts in some reasoned manner.

Given multiple samples believed to have similar composition patterns, one may form an empirical composition matrix with columns adding up to 1 but containing entries of zeros, then share information among columns to produce estimates with desirable properties; Cao et al. (2020) proposed and illustrated an approach doing just that, using as estimates some low-rank approximation of the empirical composition matrix.

In this note, we explore a simple approach to composition estimation. We take as input a matrix of raw counts, possibly with many 0's, and return a composition matrix of positive entries with the columns summing up to 1. The method is actually designed to work with one column, with information shared by other columns helping to improve performance; prior information from other sources, if available, could be equally helpful.

We shall employ penalized likelihood density estimation on a nominal discrete domain for the task. The method has long been developed and used with success in practice, but primarily on continuous domains. We now take a serious look at how it performs on a discrete domain. We shall first set the notation and outline the method, then investigate important practical issues such as smoothing parameter selection and the effective use of prior information. A pair of R functions are available for use by practitioners. Some theoretical analysis is also attempted to better understand how the method works.

2 Method

Consider a multinomial sample $\mathbf{k} \sim \text{Multinomial}(n; \mathbf{p})$, where $\mathbf{p} = (p_1, \dots, p_m)^T$, $\sum_y p_y = 1$, $\mathbf{k} = (k_1, \dots, k_m)^T$, $\sum_y k_y = n$. Our task is to estimate \mathbf{p} . When m is large and some of the p_y 's are small, $k_y = 0$ is not uncommon, and the simple maximum likelihood estimate $\hat{p}_y = k_y/n$ is undesirable.

The estimation of \mathbf{p} can be cast as the estimation of probability density $f(y)$ on domain $\mathcal{Y} = \{1, \dots, m\}$. An approach is via the penalized likelihood method, minimizing

$$-\frac{1}{n} \sum_y \left\{ k_y \eta(y) - \log \int_{\mathcal{Y}} e^{\eta(y)} \right\} + \frac{\lambda}{2} J(\eta), \quad (1)$$

where $f(y) = e^{\eta(y)} / \int_{\mathcal{Y}} e^{\eta(y)}$, $J(\eta)$ is a roughness penalty, and the smoothing parameter λ controls the trade-off between goodness-of-fit and smoothness. See, e.g., Silverman (1982) and Gu and Qiu (1993). With y nominal, a standard choice for $J(\eta)$ is $\sum_y (\eta(y) - \bar{\eta})^2$, where $\bar{\eta} = m^{-1} \sum_y \eta(y)$, which ensures invariance with respect to permutations of elements in \mathcal{Y} . This is estimation by shrinkage.

Through the specification of $\int_{\mathcal{Y}} g(y)$, one may incorporate possible prior information concerning p_y . With $\int_{\mathcal{Y}} g(y) = \sum_y w_y g(y)$, the density $f(y)$ is relative to the base measure $\{w_y\}$ on \mathcal{Y} , $p_y = w_y e^{\eta(y)} / \sum_x w_x e^{\eta(x)}$. Absent prior information, one simply sets $w_y \propto 1$, but if there is reason to suggest that (p_y/\tilde{p}_y) 's are near a constant for some \tilde{p}_y , say, one may want to use $w_y \propto \tilde{p}_y$. A nearly uniform density (relative to a base measure) is easier to estimate.

With multiple samples $\mathbf{k}_x \sim \text{Multinomial}(n_x; \mathbf{p}_x)$, $\mathbf{p}_x = (p_{x,1}, \dots, p_{x,m})^T$, $\sum_y p_{x,y} = 1$, $\mathbf{k}_x = (k_{x,1}, \dots, k_{x,m})^T$, $\sum_y k_{x,y} = n_x$, where \mathbf{p}_x 's are believed to be close to each other, one may use the collapsed data $\mathbf{k} = \sum_x \mathbf{k}_x$ to estimate a \tilde{p}_y with $w_y \propto 1$, then use $w_y \propto \tilde{p}_y$ for the estimation of individual \mathbf{p}_x 's.

3 Density Estimation on Discrete Domain: Practice

We now look at how (1) performs on a discrete domain. Some theoretical analysis is to be found in Section 5. Computation is straightforward. Our primary interests here are two fold, to check on the effectiveness of smoothing parameter selection by cross-validation, and to verify the benefit of using $w_y \propto \tilde{p}_y$ when prior information is available.




Figure 1: Density Estimation on Discrete Domain: Left: Relative efficacy $L(\lambda_o)/L(\lambda_v)$ with $w_y \propto 1$ (wider boxes) and $w_y \propto \tilde{p}_y$ (thinner boxes). Center and Right: \mathbf{p} (solid) and cross-validated $\hat{\mathbf{p}}$ (faded) with $w_y \propto 1$ (center) and $w_y \propto \tilde{p}_y$ (right); $n = 192$.

The cross-validation technique for the selection of λ in (1) as developed in Gu and Wang (2003) proved to be effective on continuous domains, and plays a central role behind the `ssden` facility in the R package `gss` (Gu 2014); technical details are to be found in Gu (2013, Sect. 7.3). We now explore its performance on a discrete domain via simple simulation.

Take $m = 100$. We first generate $Z_y \sim N(0, 1)$, then produce 50 sets of $Z_{x,y} = Z_y + z_{x,y}$ where $z_{x,y} \sim N(0, 1/4)$. One then has 50 \mathbf{p}_x 's with $p_{x,y} \propto e^{Z_{x,y}}$. Now split $N = 10000 = \sum_x n_x$ total size to these 50 multinomial distributions in a random but slightly uneven fashion, with n_x ranging from 104 to 314. With the collapsed data of size 10k, one has a cross-validated estimate \tilde{p}_y .

For each of the 50 samples, four estimates were calculated, a pair each with $w_y \propto 1$ and $w_y \propto \tilde{p}_y$. For each pair, one estimate was with λ minimizing the cross-validation score (as implemented in `ssden`) at λ_v , another with λ minimizing the Kullback-Leibler divergence $L(\lambda) = \text{KL}(\mathbf{p}, \hat{\mathbf{p}}) = \sum_y p_y \log p_y / \hat{p}_y$ at λ_o ; the subscript x is omitted in the formula, but the 50 samples were generated from 50 \mathbf{p}_x 's with generally different sizes n_x .

Ratios of $L(\lambda_o)/L(\lambda_v)$ are summarized in the boxplots in the left half of the left frame in Figure 1. The sizes $n_x \in [103, 314]$ are very low for $m = 100$, yet even with $w_y \propto 1$, $L(\lambda_o)$'s were in the range (0.080, 0.257) with the median at 0.138; the behavior of cross-validation was dichotomous, succeeding 28 times and failing 22. The method is capable, only if one can tune it properly by selecting the right λ in practice. Using $w_y \propto \tilde{p}_y$, $L(\lambda_o)$'s moved to the range (0.042, 0.100) with the median at 0.064, and cross-validation performed reliably; $L(\lambda_v)$'s had a range (0.044, 0.153) with the median at 0.080. Repeating the exercise with a more reasonable total size $N = 25k$, for $n_x \in [252, 761]$, parallel boxplots are shown in the right half of the frame; the range of $L(\lambda_v)$'s for $w_y \propto \tilde{p}_y$ are now (0.033, 0.087), with the median at 0.047.

The cross-validated estimates using a sample of $n = 192$ are shown in the center and right frames in Figure 1, with p_y 's sorted and plotted in solid. The estimates \hat{p}_y 's in faded,

vertically aligned with the p_y 's, are invariant to permutations of y , but the sorting cuts down on clutter. There are 39 $k_y = 0$ in the sample. The $w_y \propto 1$ fit has $\text{KL}(\mathbf{p}, \hat{\mathbf{p}}) = 1.244$ (really bad); the layers of \hat{p}_y 's from below in the center frame correspond to $k_y = 0, 1, \dots$. The $w_y \propto \tilde{p}_y$ fit has $\text{KL}(\mathbf{p}, \hat{\mathbf{p}}) = 0.108$, compared to $\text{KL}(\mathbf{p}, \tilde{\mathbf{p}}) = 0.126$.

4 Miscellaneous

Computation and Software

On $\mathcal{Y} = \{1, \dots, m\}$, basis functions (i.e., the likes of unit vectors) are largely independent of each other, so one needs to entertain $m - 1$ coefficients numerically, one each for all but one $y \in \mathcal{Y}$; the one less is due to a side condition needed on $\eta(y)$ to ensure a one-to-one mapping $f(y) = e^{\eta(y)} / \int_{\mathcal{Y}} e^{\eta(y)}$. When m is large, the $O(m^3)$ execution time could be demanding.

With $w_y \propto 1$, $\hat{\eta}_y$'s associated with $k_y = 0$ are all equal by symmetry, as seen in the center frame of Figure 1, so one may use that as the baseline, and exclude bases associated with $k_y = 0$ to save some computation.

A pair of R functions have been added to the `gss` package to facilitate the practical use of the proposed method.

One may use `sscomp(x, wt)` to perform density estimation on a nominal discrete domain, which takes k_y in \mathbf{x} ; $k_y = 0$ does need to be included, as the length of \mathbf{x} gives m . The default $w_y \propto 1$ can be overridden via `wt`. The function is simply a stripped down version of `ssden`, tailor-made for use in the current setting. It returns the cross-validated estimate $\hat{\mathbf{p}}$ as an $m \times 1$ matrix.

With data in a matrix of $k_{x,y}$'s of m rows, one may use `sscomp2(x)`, which collapses the columns to estimate $\tilde{\mathbf{p}}$ using `sscomp` with the default $w_y \propto 1$, calls `sscomp` with $w_y \propto \tilde{p}_y$ to estimate $\hat{\mathbf{p}}_x$ for each column, then returns the $\hat{\mathbf{p}}_x$'s in a matrix matching the input.

Remarks

Working with the log density $\eta(y)$, one guarantees $p_y > 0$, however small they might be. The domain \mathcal{Y} enters (1) via $\int_{\mathcal{Y}} e^{\eta(y)}$, so is part of data in the setting.

As seen above in the empirical study of Section 3, and also later in the theoretical analysis of Section 5, w_y 's mimicking the “shape” of p_y 's allow (1) to perform better. This resembles the mechanism behind importance sampling. With a single sample, one may use other sources of prior information in lieu of $\tilde{\mathbf{p}}$, if available.

While k_y 's should be non-negative integers according to our multinomial formulation, the algorithm and the R code work with any non-negative numbers. The same ratios k_y/n represent the same empirical pattern reflecting p_y , but a larger $n = \sum_y k_y$ makes cross-validation to pick a smaller λ , tilting \hat{p}_y closer to k_y/n .

5 Density Estimation on Discrete Domain: Theory

We now attempt some theoretical analysis concerning density estimation via (1) on a nominal discrete domain. The theory developed in the literature on continuous domains (Silverman 1982; Cox and O'Sullivan 1990; Gu and Qiu 1993) does not apply here, as conditions concerning eigenvalues of the quadratic roughness functional $J(\eta)$ no longer hold.

For a one-to-one mapping $p_y = w_y e^{\eta_y} / \sum_x w_x e^{\eta_x}$, we impose a side condition $\sum_y \eta_y = 0$. It follows that $J(\eta) = \sum_y \eta_y^2$, and (1) can be written as

$$-\frac{1}{n} \sum_y k_y \eta_y + \log \sum_y w_y e^{\eta_y} + \frac{\lambda}{2} \sum_y \eta_y^2. \quad (2)$$

Linear Approximation

Denoting by $p_{0,y} = w_y e^{\eta_{0,y}} / \sum_x w_x e^{\eta_{0,x}}$ the true probabilities, substituting $\log \sum_y w_y e^{\eta_y}$ in (2) by its quadratic approximation at $\eta_{0,y}$, and dropping terms not involving η_y , one has

$$\begin{aligned} & - \sum_y (\tilde{k}_y - p_{0,y}) \eta_y + \frac{1}{2} \left\{ \sum_y p_{0,y} (\eta_y - \eta_{0,y})^2 - \left(\sum_y p_{0,y} (\eta_y - \eta_{0,y}) \right)^2 \right\} + \frac{\lambda}{2} \sum_y \eta_y^2 \\ & = -(\tilde{\mathbf{k}} - \mathbf{p}_0)^T \boldsymbol{\eta} + \frac{1}{2} (\boldsymbol{\eta} - \boldsymbol{\eta}_0)^T (P_0 - \mathbf{p}_0 \mathbf{p}_0^T) (\boldsymbol{\eta} - \boldsymbol{\eta}_0) + \frac{\lambda}{2} \boldsymbol{\eta}^T \boldsymbol{\eta}, \end{aligned} \quad (3)$$

where $\tilde{k}_y = k_y/n$, $P_0 = \text{diag}(p_{0,1}, \dots, p_{0,m})$; for the quadratic approximation, write $A(\alpha) = \log \sum_y w_y e^{\alpha(\eta_y - \eta_{0,y}) + \eta_{0,y}}$, differentiate with respect to α , then $A(1) \approx A(0) + A'(0) + \frac{1}{2} A''(0)$.

We shall first analyze the minimizer of (3), which is linear in \tilde{k}_y , then bridge it with the minimizer of (2). Differentiating (3) with respect to $\boldsymbol{\eta}$ and setting the gradient to 0, the minimizer $\tilde{\boldsymbol{\eta}}$ satisfies

$$(\lambda I + P_0 - \mathbf{p}_0 \mathbf{p}_0^T) \tilde{\boldsymbol{\eta}} = (\tilde{\mathbf{k}} - \mathbf{p}_0) + (P_0 - \mathbf{p}_0 \mathbf{p}_0^T) \boldsymbol{\eta}_0.$$

It follows that

$$\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0 = (\lambda I + P_0 - \mathbf{p}_0 \mathbf{p}_0^T)^{-1} ((\tilde{\mathbf{k}} - \mathbf{p}_0) - \lambda \boldsymbol{\eta}_0).$$

Error Bounds

Write $B(\alpha) = \sum_y h_y w_y e^{\alpha(\eta_{1,y} - \eta_{0,y}) + \eta_{0,y}} / \sum_y w_y e^{\alpha(\eta_{1,y} - \eta_{0,y}) + \eta_{0,y}}$. It is clear that $B(1) - B(0) = \mathbf{h}^T (\mathbf{p}_1 - \mathbf{p}_0)$, where $p_{1,y} = w_y e^{\eta_{1,y}} / \sum_x w_x e^{\eta_{1,x}}$. A Taylor approximation $B(1) \approx B(0) + B'(0)$ yields $\mathbf{h}^T (\mathbf{p}_1 - \mathbf{p}_0) \approx \mathbf{h}^T (P_0 - \mathbf{p}_0 \mathbf{p}_0^T) (\boldsymbol{\eta}_1 - \boldsymbol{\eta}_0)$, so

$$(\mathbf{p}_1 - \mathbf{p}_0)^T (\boldsymbol{\eta}_1 - \boldsymbol{\eta}_0) \approx (\boldsymbol{\eta}_1 - \boldsymbol{\eta}_0)^T (P_0 - \mathbf{p}_0 \mathbf{p}_0^T) (\boldsymbol{\eta}_1 - \boldsymbol{\eta}_0) = V(\boldsymbol{\eta}_1 - \boldsymbol{\eta}_0); \quad (4)$$

the left-hand-side is $\text{KL}(\mathbf{p}_0, \mathbf{p}_1) + \text{KL}(\mathbf{p}_1, \mathbf{p}_0)$, the symmetrized Kullback-Leibler, and the right-hand-side quadratic proxy is a weighted mean square error of log probability, with

$p_{0,y}$'s as the weights. Note that $V(\boldsymbol{\eta} + C) = V(\boldsymbol{\eta})$, invariant to the side condition imposed on $\boldsymbol{\eta}$. We shall try to bound

$$(\lambda J + V)(\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0) = (\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0)^T (\lambda I + P_0 - \mathbf{p}_0 \mathbf{p}_0^T) (\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0).$$

By Cauchy-Schwartz, one can treat the bias term involving $\boldsymbol{\eta}_0$ and the variance term involving $(\tilde{\mathbf{k}}_r - \mathbf{p}_0)$ separately.

For the variance term, note that $E[(\tilde{\mathbf{k}} - \mathbf{p}_0)(\tilde{\mathbf{k}} - \mathbf{p}_0)^T] = n^{-1}(P_0 - \mathbf{p}_0 \mathbf{p}_0^T)$, so

$$\begin{aligned} E[(\tilde{\mathbf{k}} - \mathbf{p}_0)^T (\lambda I + P_0 - \mathbf{p}_0 \mathbf{p}_0^T)^{-1} (\lambda I + P_0 - \mathbf{p}_0 \mathbf{p}_0^T) (\lambda I + P_0 - \mathbf{p}_0 \mathbf{p}_0^T)^{-1} (\tilde{\mathbf{k}} - \mathbf{p}_0)] \\ = \frac{1}{n} \text{tr}((\lambda I + P_0 - \mathbf{p}_0 \mathbf{p}_0^T)^{-1} (P_0 - \mathbf{p}_0 \mathbf{p}_0^T)) = \frac{1}{n} \sum_y \frac{\rho_y}{\lambda + \rho_y} < \frac{1}{n\lambda}, \end{aligned}$$

where ρ_y 's are the eigenvalues of $P_0 - \mathbf{p}_0 \mathbf{p}_0^T$, $\sum_y \rho_y < 1$. For the bias term,

$$\lambda^2 \boldsymbol{\eta}_0^T (\lambda I + P_0 - \mathbf{p}_0 \mathbf{p}_0^T)^{-1} (\lambda I + P_0 - \mathbf{p}_0 \mathbf{p}_0^T) (\lambda I + P_0 - \mathbf{p}_0 \mathbf{p}_0^T)^{-1} \boldsymbol{\eta}_0 \leq \lambda \boldsymbol{\eta}_0^T \boldsymbol{\eta}_0.$$

Putting thing together, one has

$$(\lambda J + V)(\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0) = O_p(\lambda \boldsymbol{\eta}_0^T \boldsymbol{\eta}_0 + (n\lambda)^{-1}). \quad (5)$$

When $(w_y/p_{0,y})$'s are near a constant, $\boldsymbol{\eta}_0^T \boldsymbol{\eta}_0$ is small.

These bounds may not be sharp, but in case they are, the optimal rate also depends on how $\boldsymbol{\eta}_0^T \boldsymbol{\eta}_0$ grows with m . If $\boldsymbol{\eta}_0^T \boldsymbol{\eta}_0 \asymp m$, then the optimal rate is $(\lambda J + V)(\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0) = O_p(\sqrt{m/n})$ achieved at $\lambda \asymp (mn)^{-1/2}$; m is allowed to grow with n , but the growth rate should be slower than $O(n)$ to guarantee consistency.

Approximation Error

Set $\eta_y = \hat{\eta}_y + \alpha h_y$ in (2), where $\hat{\eta}_y$'s minimize (2) and h_y 's are arbitrary. Differentiating with respect to α and setting the derivative at $\alpha = 0$ to 0, one has

$$-\frac{1}{n} \sum_y k_y h_y + \sum_y \hat{p}_y h_y + \lambda \sum_y \hat{\eta}_y h_y = -\tilde{\mathbf{k}}^T \mathbf{h} + \hat{\mathbf{p}}^T \mathbf{h} + \lambda \hat{\boldsymbol{\eta}}^T \mathbf{h} = 0, \quad (6)$$

where $\hat{p}_y = w_y e^{\hat{\eta}_y} / \sum_x w_x e^{\hat{\eta}_x}$. Setting $\boldsymbol{\eta} = \tilde{\boldsymbol{\eta}} + \alpha \mathbf{h}$ in (3), differentiating with respect to α , then setting the derivative at $\alpha = 0$ to 0, one has

$$-(\tilde{\mathbf{k}} - \mathbf{p}_0)^T \mathbf{h} + (\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0)^T (P_0 - \mathbf{p}_0 \mathbf{p}_0^T) \mathbf{h} + \lambda \tilde{\boldsymbol{\eta}}^T \mathbf{h} = 0. \quad (7)$$

Subtracting (7) from (6) and setting $\mathbf{h} = \hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}}$, some algebra yields

$$\begin{aligned} \lambda(\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}})^T (\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}}) + (\hat{\mathbf{p}} - \tilde{\mathbf{p}})^T (\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}}) \\ = (\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0)^T (P_0 - \mathbf{p}_0 \mathbf{p}_0^T) (\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}}) - (\tilde{\mathbf{p}} - \mathbf{p}_0)^T (\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}}). \end{aligned} \quad (8)$$

Using the mean value theorem in the arguments leading to (4),

$$(\hat{\mathbf{p}} - \tilde{\mathbf{p}})^T(\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}}) = (\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}})^T(P_1 - \mathbf{p}_1\mathbf{p}_1^T)(\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}})$$

where $P_1 = \text{diag}(p_{1,1}, \dots, p_{1,m})$, $\mathbf{p}_1 = (p_{1,1}, \dots, p_{1,m})^T$ correspond to a convex combination $\boldsymbol{\eta}_1$ of $\hat{\boldsymbol{\eta}}$ and $\tilde{\boldsymbol{\eta}}$. Likewise,

$$(\tilde{\mathbf{p}} - \mathbf{p}_0)^T(\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}}) = (\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0)^T(P_2 - \mathbf{p}_2\mathbf{p}_2^T)(\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}}).$$

Assuming $\mathbf{a}^T(P_i - \mathbf{p}_i\mathbf{p}_i^T)\mathbf{b}$, $i = 1, 2$ be bounded by multiples of $\mathbf{a}^T(P_0 - \mathbf{p}_0\mathbf{p}_0^T)\mathbf{b}$ from below and above, one has from (8),

$$(\lambda J + c_1 V)(\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}}) \leq c_2 |(\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0)^T(P_0 - \mathbf{p}_0\mathbf{p}_0^T)(\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}})|$$

for some $0 < c_1, c_2 < \infty$. By Cauchy-Schwartz, $V(\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}}) \leq (c_2/c_1)^2 V(\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0)$, and in turn $\lambda J(\hat{\boldsymbol{\eta}} - \tilde{\boldsymbol{\eta}}) \leq c_3 V(\tilde{\boldsymbol{\eta}} - \boldsymbol{\eta}_0)$ for some $0 < c_3 < \infty$. Simple manipulation further propagates the rate of (5) to $(\lambda J + V)(\hat{\boldsymbol{\eta}} - \boldsymbol{\eta}_0)$.

References

- Aitchison, J. (1986). *The Statistical Analysis of Compositional Data*. London: Chapman and Hall.
- Cao, Y., A. Zhang, and H. Li (2020). Multisample estimation of bacterial composition matrix in metagenomics data. *Biometrika* 107, 75–92.
- Cox, D. D. and F. O’Sullivan (1990). Asymptotic analysis of penalized likelihood and related estimators. *Ann. Statist.* 18, 124–145.
- Gu, C. (2013). *Smoothing Spline ANOVA Models* (2nd ed.). New York: Springer-Verlag.
- Gu, C. (2014). Smoothing spline ANOVA models: R package `gss`. *J. Stat. Softw.* 58(5), 1–25.
- Gu, C. and C. Qiu (1993). Smoothing spline density estimation: Theory. *Ann. Statist.* 21, 217–234.
- Gu, C. and J. Wang (2003). Penalized likelihood density estimation: Direct cross-validation and scalable approximation. *Statist. Sin.* 13, 811–826.
- Silverman, B. W. (1982). On the estimation of a probability density function by the maximum penalized likelihood method. *Ann. Statist.* 10, 795–810.
- Xing, X., J. S. Liu, and W. Zhong (2017). MetaGen: reference-free learning with multiple metagenomic samples. *Genome Biol.* 18(187).