

Using clinical trial registries to inform Copas selection model for publication bias in meta-analysis

Ao Huang , Sho Komukai, Tim Friede , Satoshi Hattori *

Abstract

Prospective registration of study protocols in clinical trial registries is a useful way to minimize the risk of publication bias in meta-analysis, and several clinical trial registries are available nowadays. However, they are mainly used as a tool for searching studies and information submitted to the registries has not been utilized as efficiently as it could. In addressing publication bias in meta-analyses, sensitivity analysis with the Copas selection model is a more objective alternative to widely-used graphical methods such as the funnel-plot and the trim-and-fill method. Despite its ability to quantify the potential impact of publication bias, a drawback of the model is that some parameters not to be specified. This may result in some difficulty in interpreting the results of the sensitivity analysis. In this paper, we propose an alternative inference procedure for the Copas selection model by utilizing information from clinical trial registries. Our method provides a simple and accurate way to estimate all unknown parameters in the Copas selection model. A simulation study revealed that our proposed method resulted in smaller biases and more accurate confidence intervals than existing methods. Furthermore, two published meta-analyses had been re-analysed to demonstrate how to implement the proposed method in practice.

Keywords: Clinical trial registry; Meta-analysis; Publication bias; Research synthesis; Selection model

1 Introduction

Publication bias is one of the most important concerns in systematic reviews and meta-analyses. Part of the issue is that authors and scientific journals are likely to publish studies with statistically significant results than inconclusive studies[1]. To be statistically significant, the studies with small sample size need to show larger treatment effects, which is known as the small-study effect[2]. Thus, this may force us to analyze selective samples from the population of interest, and the standard meta-analysis techniques may suffer from this selection bias, which is referred to as publication bias.

The funnel plot is a simple and widely-used graphical tool to check for publication bias, which is defined by plotting the estimate of the effect size such as the log-odds ratio or the log-hazard ratio as the horizontal axis and the related measure of precision such as the square root of the sample size or the inverse of the standard error as the vertical axis. One can address the publication bias by examining asymmetry of the funnel plot visually and more formal statistical evaluation of the funnel-plot asymmetry can be made with a regression based test[3, 4] or a rank-based test [5, 6]. The trim-and-fill method is a nonparametric method to adjust publication bias utilizing the funnel-plot asymmetry[7].

All these methods are simple to apply and are widely used in practice. However, publication bias is not the only reason for asymmetry of the funnel plot; Egger et al.[3] listed different potential reasons for the funnel-plot asymmetry including the between-study heterogeneity and

* *correspondence to:* Satoshi Hattori, Institute for Open and Transdisciplinary Research Initiatives, Osaka University; email: hattoris@biostat.med.osaka-u.ac.jp

others. Then, detection of the publication bias by the funnel plot may be subjective and the trim-and-fill method may not work well[8, 9]. Alternatively, several sensitivity analysis methods have been developed. Copas and colleagues introduced the Heckman-type parametric selection model, which was originally proposed in the econometrics literature[10], to describe a selective publication process in meta-analysis. In this paper, this selection model is referred as the Copas selection model[11, 12]. The sensitivity analysis with the Copas selection model has several advantages over existing alternative sensitivity analysis methods[13, 14, 15]; a user-friendly *R* package *metasens* is available for the Copas-Shi selection model[11, 12], and its performance was empirically examined[16, 17]. Thus, we focus on this selection model in this paper.

The Copas selection model consists of a pair of models; one is for the outcome, such as the log-odds ratio or the log-hazard ratio reported in publications, and the other is a model for the latent variable describing a selective publication process. It's hard to estimate all the parameters in the Copas selection model, and thus a series of papers by Copas and colleagues took a sensitivity analysis approach; some parameters were fixed as sensitivity parameters in a certain range, and the impact on the estimate of interest was studied. However, it's not easy to define an appropriate range for the sensitivity parameters.

Prospective registration of study protocols in clinical trial registries is a useful non-statistical approach for publication bias. The International Committee of Medical Journal Editors (ICMJE) initiated a trials-registration policy as a condition for publication in its member journals in July 2005[18], which promoted the development and utilization of the clinical trial registries. ClinicalTrials.gov (<https://clinicaltrials.gov/ct2/home>) is one of the early examples which was established in 2000 and is nowadays a very popular registry with 334,954 registered studies in April 2019. World Health Organization's (WHO) International Clinical Trials Registry Platform (ICTRP) (<http://apps.who.int/trialsearch/>) is another well-known clinical trials registry which was constructed after the initiation of ICMJE's policy and has been supported by the WHO. FDA (US Food and Drug Administration) database (<https://www.accessdata.fda.gov/scripts/cder/daf/>) contains information about products approved by the FDA for human use in the United States. Besides, EU clinical trials register (<https://www.clinicaltrialsregister.eu/ctr-search/search>) and ISRCTN (<https://www.isrctn.com/>) are also well-developed and widely used in recent years as well as many national registries that often also provide lay summaries in the local language[19, 20].

Baudard et al. [21] investigated 14 meta-analyses and showed that by incorporating the information from clinical trial registries, the effect size estimates might change by 29%. It indicated that searching clinical trial registries may be useful to reduce the bias due to selective publication. On the other hand, although clinical trial registries provide potentially useful information of the registered clinical trials (e.g., study design, target sample size, development phase), they are used only as a tool to search for studies[22]. Information on unpublished studies provided by clinical trial registries is not formally incorporated in the analyses. To the best of knowledge, only two papers mentioned the use of clinical trial registries in statistical inference. Matsuoka et al. [23] proposed a publication bias detection method by utilizing the planned sample size from World Health Organization's (WHO) International Clinical Trials Registry Platform (ICTRP). Mavridis et al. [24]utilized FDA database to construct a prior distribution

of parameters in their Bayesian Copas-type model for sensitivity analysis in network meta-analysis.

In this paper, we improve statistical inference for the Copas-Shi selection model utilizing the information from clinical trial registries. To be specific, utilizing the planned sample size of unpublished studies, which are provided by most clinical trial registries, we propose a method that fit by the maximum likelihood method. All the unknown parameters can be estimated from data, and statistical inference can be conducted according to the standard maximum likelihood theory. In addition to the confidence interval with the normal quantile according to the maximum likelihood theory, we also propose two modified confidence intervals aiming improvement of performance with the small number of studies. Through numerical studies, we observed that our method outperformed existing methods with smaller biases and coverage probabilities closer to the nominal level. Thus, by utilizing the information of unpublished studies from clinical trial registries, inference based on the Copas selection model can be improved.

The remainder of this paper is organized as follows. In Section 2, we introduce two motivating meta-analyses. In Section 3, we briefly review the sensitivity analysis method of Copas and Shi[11, 12], and then introduce our new inference procedure by utilizing clinical trial registries. In Section 4, we conduct a series of simulation studies motivated by the tiotropium study to evaluate the performance of our proposed method in comparison to other competitive approaches. In Section 5, we revisit the two studies in Section 2 to illustrate the proposed method. Finally, we discuss some limitations and further research in Section 6.

2 Motivating examples

2.1 Tiotropium study

Karner et al.[25]conducted a Cochrane review of tiotropium versus placebo in COPD. The primary outcome was COPD exacerbations, which was defined as a complex of respiratory events or symptoms (new onset or an increase in at least one of cough, sputum, dyspnoea or wheeze) that lasted at least three days and required treatment with antibiotics and/or systemic corticosteroids. In each study, the odds ratio for the event of one or more exacerbations was used to compare the treatment and placebo groups. In the meta-analysis by Karner et al.[25], 22 studies were analyzed using a random-effects model, and the pooled odds ratio was reported as 0.78 with a 95% confidence interval of [0.70, 0.87].

To assess the potential risk of publication bias, Karner et al.[25]conducted the Egger’s test with the p-value of 0.22 and concluded no serious concern of publication bias. However, our additional application of the Macaskill’s regression test[4] gave a p-value of 0.06, suggesting the funnel-plot asymmetry. Then, publication bias might be a concern. Karner et al. [25]made their efforts to search the studies with results submitted on ClinicalTrials.gov. We conducted a more comprehensive search of multiple registries, including ClinicalTrials.gov, World Health Organization’s (WHO) International Clinical Trial Registry, EU clinical trials register, FDA database, and ISRCTN platform. We used the following terms: *Tiotropium* or *Spiriva* or *HandiHaler* or *Respimat* for the search. They were the same as used in the meta-analysis of

Karner et al. [25], and we also set primary completion date before February 28, 2012, which aimed to be consistent with the original meta-analysis. We detected 10 related studies on these registries, among which 8 came from ClinicalTrials.gov, 1 came from EU clinical trials register, and 1 came from World Health Organization’s (WHO) International Clinical Trials Registry. Two of them reported the outcome measure and were incorporated with published studies, and 8 studies without results were deemed as unpublished studies. Finally, we got the tiotropium dataset with 24 published studies and 8 unpublished studies with information of sample size only (see Table S1 in Web-appendix A).

2.2 Clopidogrel study

The second example is about the comparison of high and standard maintenance-dose clopidogrel on major adverse cardiovascular/cerebrovascular events (MACE/ MACCE) by Chen et al.[26]. The original meta-analysis included 12 studies and reported pooled odds ratio as 0.6 based on the fixed-effect model with a 95% CI of [0.43, 0.83], and no significant bias was concluded with the Egger’s test ($p=0.25$). However, we rerun the Macaskill’s test, which suggested funnel-plot asymmetry with a P value of 0.02. We took the procedure as given in the first example and identified 3 unpublished studies from ClinicalTrials.gov by following terms: *Clopidogrel* or *Plavix* or *Iscover*, with completion date before August 31, 2013. The resulting dataset was presented in Table S2 in Web-appendix A.

3 Inference procedures for the Copas selection model

3.1 Brief review of the sensitivity analysis method by Copas and Shi

In order to evaluate the potential impact of publication bias on estimation of the treatment effect, Copas and Shi [11] introduced a sensitivity analysis method. Suppose we conduct a meta-analysis of N studies. Let $\hat{\theta}_i$ be the estimated treatment effect of the i th study such as the log-odds ratio or the log-hazard ratio, and s_i be an estimate of its standard error. We assume that $(\hat{\theta}_i, s_i)$ is available for $i = 1, 2, \dots, N$. To integrate the treatment effects over the studies, we consider the standard random-effects model with

$$\hat{\theta}_i = \theta_i + \sigma_i \epsilon_i, \quad \theta_i \sim N(\theta, \tau^2), \quad \epsilon_i \sim N(0, 1), \quad (1)$$

where θ_i is the study-specific treatment effect of the i th study, which is a random-effect following a normal distribution with mean θ and between-study variance τ^2 , and σ_i^2 is the true within-study variance of the outcome $\hat{\theta}_i$. The random elements θ_i and ϵ_i are assumed to be independent.

To describe the selective publication process, Copas and Shi [11, 12] considered the following model

$$Y_i = \alpha_0 + \alpha_1/s_i + \delta_i, \quad \delta_i \sim N(0, 1), \quad \text{corr}(\epsilon_i, \delta_i) = \rho, \quad (2)$$

where $\text{corr}(\epsilon_i, \delta_i)$ denotes the correlation coefficient between ϵ_i and δ_i . The i th study is assumed to be published if and only if $Y_i > 0$. The term $\alpha_0 + \alpha_1/s_i$ models the small study effect. That is, studies of high precision are more likely to be published. The correlation coefficient ρ is responsible for describing the influence of the outcome $\hat{\theta}_i$ on likelihood of publication. Copas

and Shi[11] considered to make inference by maximizing the likelihood function conditional on the study being published. The conditional likelihood function is given by

$$\begin{aligned}
L_{obs}(\theta, \tau, \rho, \alpha_0, \alpha_1; s_i) &= \sum_{i=1}^N \left[\log f(\hat{\theta}_i | Y_i > 0, s_i) \right] \\
&= \sum_{i=1}^N \left[-\frac{1}{2} \log(\tau^2 + \sigma_i^2) - \frac{(\hat{\theta}_i - \theta)^2}{2(\tau^2 + \sigma_i^2)} - \log \Phi(\alpha_0 + \alpha_1/s_i) + \log \Phi(v_i) \right], \quad (3)
\end{aligned}$$

where $f(\cdot | Y_i > 0, s_i)$ is the probability density function of $\hat{\theta}_i$ conditional on $Y_i > 0$ and s_i . $\Phi(\cdot)$ is the cumulative distribution function of the standard normal distribution, and $v_i = \left\{ \alpha_0 + \alpha_1/s_i + \rho\sigma_i(\hat{\theta}_i - \theta)/(\tau^2 + \sigma_i^2) \right\} / \sqrt{1 - \rho^2\sigma_i^2/(\tau^2 + \sigma_i^2)}$. In meta-analysis literatures, s_i is usually regarded as being equal to σ_i and known. This is relevant approximately at least when the number of subjects is sufficiently large. Copas and Shi[11, 12] took care of underestimation of σ_i due to publication bias, and then they utilized the quantity $var(\theta_i | s_i, Y_i > 0)$ to estimate σ_i^2 in inference of their selection model. As argued by Copas and Shi[11, 12], the log-likelihood function (3) may take its maximum over a very flat plateau, and then it may be computationally very challenging to estimate all the parameters simultaneously. Thus Copas and Shi[11, 12] proposed that the vector (α_0, α_1) was fixed as sensitivity parameters, and (θ, τ, ρ) was estimated by maximizing the conditional log-likelihood function (3). One can examine how the estimate $\hat{\theta}$ for the overall treatment effect θ would be influenced by the assumed selective publication process. Since we do not know the true value of (α_0, α_1) , we should apply this method with various choices for the parameter vector (α_0, α_1) in a certain range. According to the formula by Copas and Shi[11], one can translate (α_0, α_1) into the expected numbers of unpublished studies by $M = \sum_{i=1}^N \{1 - P(Y_i > 0 | s_i)\} / P(Y_i > 0 | s_i)$, which is more interpretable. Then, one can evaluate how biased the estimate of the treatment effect θ is with a certain number of unpublished studies behind. Although such a consideration provides nice insights about robustness of meta-analysis results, it is still unclear what range of the number of unpublished studies is sufficient to be considered.

3.2 Proposed inference procedure utilizing clinical trial registries

In addition to the N published studies, we suppose that there are M unpublished studies identified by searching clinical trial registries. Without loss of generality, the first N studies are published and the last M studies are unpublished. Items required at registration are not necessarily common among clinical trial registries. For example, FDA database provides the sample size of each experimental group in its summary review PDF document, but other clinical trial registries may not give it. However, all the clinical trial registries we referred provide the planned total sample size (not separately by the experimental groups) of each study. Thus, we suppose the triple $(n_i, \hat{\theta}_i, s_i)$ is available for $i=1,2,\dots,N$ (published studies), and only n_i is available for $i=N+1,\dots,N+M$ (unpublished studies). A binary variable $D_i = I(Y_i > 0)$ denotes the publication status (published/unpublished) of each study. Instead of the model (2), we employ an alternative selection model

$$Y_i = \alpha_0 + \alpha_1 \sqrt{n_i} + \delta_i, \quad \delta_i \sim N(0, 1), \quad (4)$$

which was used by Copas[27]. For the models (1) and (4), we introduce the maximum likelihood approach as follows.

Considering the likelihood function of the published and the unpublished studies, the log likelihood is given by

$$\begin{aligned}
L_{full}(\theta, \tau, \rho, \alpha_0, \alpha_1) &= \sum_{i=1}^N \left[\log f(\hat{\theta}_i | Y_i > 0; n_i) + \log P(Y_i > 0; n_i) \right] \\
&+ \sum_{i=N+1}^{N+M} [\log P(Y_i < 0; n_i)] \\
&= \sum_{i=1}^N \left[-\frac{1}{2} \log(\tau^2 + \sigma_i^2) - \frac{(\hat{\theta}_i - \theta)^2}{2(\tau^2 + \sigma_i^2)} + \log \Phi(\tilde{v}_i) \right] \\
&+ \sum_{i=N+1}^{N+M} \{ \log[1 - \Phi(\alpha_0 + \alpha_1 \sqrt{n_i})] \}, \tag{5}
\end{aligned}$$

where $\tilde{v}_i = \left\{ \alpha_0 + \alpha_1 \sqrt{n_i} + \rho \sigma_i (\hat{\theta}_i - \theta) / (\tau^2 + \sigma_i^2) \right\} / \sqrt{1 - \rho^2 \sigma_i^2 / (\tau^2 + \sigma_i^2)}$. Following the treatment often made in meta-analysis literature, we suppose that $\sigma_i = s_i$. By maximizing this log-likelihood function (5), we can estimate all the parameters $(\theta, \tau, \rho, \alpha_0, \alpha_1)$ simultaneously, and the resulting maximum likelihood estimator is denoted by $(\hat{\theta}, \hat{\tau}, \hat{\rho}, \hat{\alpha}_0, \hat{\alpha}_1)$. Maximization of the log-likelihood function can be implemented with the standard non-linear optimization techniques. Here, we employ the constraint optimizer L-BFGS-B method in the `nlmnb()` function in *R* (package `stats`, version 3.6.2) to do this. Statistical inference can be conducted following the standard maximum likelihood theory.

4 Simulation study

4.1 Methods to be compared

Simulation studies were carried out to evaluate the performance of our new inference procedure. We investigated whether or not the proposed method outperforms the Copas sensitivity analysis method. For reference, we also compared it with standard random-effects meta-analysis using the restricted maximum likelihood (REML) estimation, which did not account for selective publication process. In addition to the standard REML confidence interval based on normal quantiles, the Hartung and Knapp method[28, 29] was applied using *t*-quantiles and rescaled standard errors, which is referred to as REML.KnHa in the following. As outlined in Section 3, the Copas method is a sensitivity analysis approach resulting in multiple estimates under several settings of the number of unpublished studies. For comparison with other methods, we took the estimate of the treatment effect with the smallest number of unpublished studies of the p-value of the goodness-of-fit test larger than 0.1[11], which is presented in the output of *Copas* function in *metasens* package[30]. For the REML, we used the commonly used *metafor* package with method option equal to “REML” and test equal to “knha” to obtain the CI with the Hartung and Knapp method.

4.2 Data generation

We mimicked meta-analyses motivated by the tiotropium study. Suppose each study aimed to compare two groups of the treatment group and the control group with respect to a binary outcome. The log-odds ratio was used as the summary measure of the treatment effect between two groups. Let $\hat{\theta}_i$ in the model (1) denoted the empirical log-odds ratio of the i th study. We set $\theta = -0.25$ and $\tau=0.05, 0.15$ or 0.3 . The total number of studies including published and unpublished was set as 15, 25, 50 or 100. We generated datasets according to the models (1) and (4) as follows. At first, we generated $\theta_i \sim N(\theta, \tau^2)$, which was the true log-odds ratio of the i th study. Next, we generated individual participant data of the two groups with the log-odds ratio of θ_i . We set the true event rate of the control group from the uniform distribution $U(0.2, 0.9)$, and then set the event rate of treatment group satisfying the log-odds ratio of θ_i . Similarly to Kuss[31], the total sample size of each study was generated from $LN(5, 1)$, the log-normal distribution with the location parameter 5 and scale parameter 1, and the minimum sample size was restricted to 20 patients (values below 20 were rounded up to 20). Subjects were allocated to the two groups with probability of 0.5. With the generated individual participant data, we calculated an empirical log odds ratio $\hat{\theta}_i$ and its standard error s_i . Y_i in the model (4) was generated according to the conditional distribution,

$$Y_i|\hat{\theta}_i \sim N(\alpha_0 + \alpha_1\sqrt{n_i} + \rho\sigma_i(\hat{\theta}_i - \theta)/(\tau^2 + \sigma_i^2), 1 - \rho^2\sigma_i^2/(\tau^2 + \sigma_i^2)) \quad (6)$$

and set $D_i = I(Y_i > 0)$, which entailed us to generate $(\hat{\theta}_i, Y_i)$ from the joint distribution defined by the models (1) and (4). The parameters (α_0, α_1) were set based on our consideration for the publication rate of a study with the minimum sample size of 20 and that with a large sample size of 500, which were denoted by P_{20} and P_{500} , respectively. Set $P_{500} = 0.99$, reflecting our belief that a study with 500 patients had sufficiently large probability to be published, and P_{20} was set to 0.1, 0.3 or 0.5, which represented our different concern to the publication rate of a small study with sample size of 20. Thus according to the model (4), we could derive the parameters (α_0, α_1) by solving the equations $P_{20} = \Phi(\alpha_0 + \alpha_1\sqrt{20})$ and $P_{500} = \Phi(\alpha_0 + \alpha_1\sqrt{500})$. Then we got three pairs of (α_0, α_1) as $(-2.18, 0.20)$, $(-1.24, 0.16)$ and $(-0.58, 0.13)$, which resulted in 40%, 27% and 19% unpublished studies on average in the simulated meta-analyses. The parameter of ρ was set to -0.4 or -0.8, and for each ρ we investigated 3 scenarios of τ (0.05, 0.15 or 0.3). For each scenario, we generated 1000 simulated meta-analyses.

In Figure 1, we presented funnel plots of randomly selected simulated meta-analyses under different settings of (ρ, τ) when the total number of published and unpublished studies was 50. The filled and open circles represent published and unpublished studies, respectively, and we observed our simulation setting successfully simulated the selective publication process often observed in practice. The bold vertical line represents the true θ and the dashed vertical line is the REML estimate with published studies. They were certainly different and thus publication bias had certain impacts in simulated data.

4.3 Simulation results

The properties of each method were assessed by evaluating biases and standard errors of the estimates for θ and average lengths and coverage probabilities of two-tailed 95% confidence

intervals of θ . Following the standard maximum likelihood theory, we constructed the two-tailed 95% confidence interval for the estimates of θ with our new proposal by $\hat{\theta} \pm Z_{1-\frac{\alpha}{2}} \times \text{S.E.}(\hat{\theta})$, where $Z_{1-\frac{\alpha}{2}}$ denoting the $1 - \frac{\alpha}{2}$ quantile of the standard normal distribution, and $\text{S.E.}(\hat{\theta})$ is calculated by the inverse of the Fisher information matrix. The results for this confidence interval were denoted by MLE(N) in the summary table. All these methods were implemented with non-linear optimization techniques, thus we also summarized the number of converged cases (*NOC*) in each scenario. For our method, the non-convergence was defined as a failure in optimization with negative hessian matrix or unsuccessful convergence based on the convergence indicator in the output of *nlminb* function. For the Copas sensitivity analysis method, we followed the rule of *Copas* function[16, 30].

Results of the simulation studies with $(\alpha_0, \alpha_1) = (-2.18, 0.2)$ were reported in Table 1 ($\rho = -0.4$) and Table 2 ($\rho = -0.8$), respectively. The datasets for Tables 1 and 2 had around 40% unpublished studies on average. In almost all the simulated datasets, the log-likelihood was successfully maximized and then gave estimates. In all the scenarios, the REML had considerable biases. With $\rho = -0.8$ (Table 2), biases were larger, reflecting that $\rho = -0.8$ modeled a more selective publication process. Both the Copas method and our new proposal decreased biases and ours had the smallest biases in most scenarios. With larger number of studies ($N=50$ and 100), the maximum likelihood estimator tended to have very negligible biases, and the confidence intervals based on the maximum likelihood theory (MLE(N)) had empirical coverage probabilities close to the nominal level of 95%. With smaller number of the studies ($N=15$ and 25), our method successfully decreased biases. However, coverage probabilities might not be close to the nominal level and were not satisfactory when between-study heterogeneity was moderate or substantial ($\tau = 0.15$ and 0.3). We demonstrated results of the simulation studies with $(\alpha_0, \alpha_1) = (-1.24, 0.16)$ in Tables S3 and S4, and that with $(\alpha_0, \alpha_1) = (-0.58, 0.13)$ in Tables S5 and S6 in Web-appendix B. Findings were very similar to discussed above based on Tables 1 and 2.

4.4 Modified confidence intervals

To improve the coverage probabilities with smaller number of studies, we introduced two modified confidence intervals. As suggested by Follmann and Proschan[32], an alternative confidence interval $\hat{\theta} \pm t_{(N-1);(1-\frac{\alpha}{2})} \times \text{S.E.}(\hat{\theta})$ may be used, where $t_{(N-1);(1-\frac{\alpha}{2})}$ denoting the $1 - \frac{\alpha}{2}$ quantile of Student-*t* distribution with $N - 1$ degree of freedom. Further modification was also considered by mimicking the idea of the improved estimator of variance which was proposed by Knapp and Hartung[33] in the meta-regression context; we modified the standard error of $\hat{\theta}$ by $\text{S.E.}(\hat{\theta})^\# = \text{Max}(\text{S.E.}(\hat{\theta})_{MLE}, \text{S.E.}(\hat{\theta})_{REML-KnHa})$, where $\text{S.E.}(\hat{\theta})_{REML-KnHa}$ was the estimator of standard error with HK-adjusted REML method, reflecting our belief that the standard error of our proposal should be no less than the estimator of methods without considering the selective publication process for the additional uncertainty of unpublished studies. That is, a more conservative confidence interval was calculated by $\hat{\theta} \pm t_{(N-1);(1-\frac{\alpha}{2})} \times \text{S.E.}(\hat{\theta})^\#$. We applied these two confidence intervals to the simulated meta-analyses, results were also shown in Tables 1 and 2 and Tables S3-S6 in Web-appendix B, referring as MLE(T) and MLE(*SE*[#]), respectively. We observed that these two confidence

intervals $\text{MLE}(T)$ and $\text{MLE}(SE^\sharp)$ gave certain improvement. In particular, $\text{MLE}(SE^\sharp)$ had much improved empirical coverage probabilities with $N=15$.

5 Re-analysis of two motivating examples with clinical trial registries

In this section, we applied our new inference procedure to the two case studies of the tiotropium study and clopidogrel study introduced in Section 2. To present data of unpublished studies as well as that of published studies, we propose to use a modified funnel-plot adding information of the planned sample size of unpublished studies with horizontal lines passing the y-axis at $\sqrt{n_i}$. The modified funnel-plots for tiotropium study and clopidogrel study were presented in Figures 2 and 3, respectively. We could observe that for both of the two motivation cases, all the unpublished studies concentrated on the lower part of the funnel-plot. To the data, we applied our proposed method, as well as other related methods used in the simulation section, and the results for the two motivation cases were summarized in Tables 3 and 4, respectively.

As presented in Subsection 2.1, for the tiotropium study, we identified 24 published and 8 unpublished studies. By applying the linear mixed-effect model for the log-odds ratios with the REML method, the integrated odds ratio was obtained as 0.768 with a two-tailed 95% CI of [0.697, 0.847] based on the normal approximation. The Knapp and Hartung modification gave a similar CI of [0.691, 0.854]. According to the sensitivity analysis by the Copas model, one can observe that as increasing the number of unpublished studies, the odds ratio increased up to 0.811, which was corresponding to the case of 13 unpublished studies. The overall treatment effect was significantly different from unity even with 13 unpublished studies. Suppose we have 8 published studies behind which was same with our findings from the clinical trial registries, Copas sensitivity analysis method gave the pooled odds ratio as 0.803 with a 95% CI of [0.717, 0.898]. Our proposed method gave the integrated estimate of 0.787 with two-tailed 95% confidence intervals [0.710, 0.873] and [0.706, 0.878] based on the standard normal quantile and t-quantile, respectively. In this case, the standard error from our method was larger than the standard error from REML with the Knapp and Hartung modification, so the modified confidence interval $\text{MLE}(SE^\sharp)$ was same with $\text{MLE}(T)$. To summary, all the methods provided statistically significant effect of tiotropium, indicating that selective publication might have not been much of an issue here.

For the clopidogrel study which we introduced in Subsection 2.2, with only 3 unpublished studies identified from the clinical trial registries, the estimates with and without consideration of publication bias were considerably different. By applying the REML with 12 published studies, the estimator of the pooled odds ratio was 0.579 and with two-tailed 95% CIs of [0.375, 0.892] and [0.385, 0.871] based on the normal quantile and the Knapp and Hartung modification, respectively. Both of them supported the conclusion that the high maintenance-dose clopidogrel significantly reduced the incidence of MACE/MACCE in comparison to the standard-maintenance-dose clopidogrel. However, the sensitivity analysis by the Copas model showed that as the number of unpublished studies increased up to 4, the statistical significance of the High-maintenance-dose clopidogrel disappeared, and as high as 39% change in estimates of odds ratio could be observed with 6 unpublished studies. Our proposed method gave the

integrated estimate of 0.692, and a two-tailed 95% CI based on the normal approximation was given as [0.496, 0.967], whose upper bound was very close to 1. The modified CIs introduced in Subsection 4.4 could be obtained as [0.476, 1.007] (MLE(T)) and [0.460, 1.041] (MLE(SE^\sharp)), respectively. These results suggested that the significant effect of the high maintenance-dose clopidogrel might be marginal and should be interpreted with caution.

6 Discussion

By utilizing the information on unpublished studies obtained from the clinical trial registries, we proposed a new inference procedure for the Copas selection model, which provides more objective evaluation of publication bias than the widely-used funnel plot and trim-and-fill method[16, 17]. All the unknown parameters in the Copas selection model can be estimated from data with the proposed method. It resolved the issue of the sensitivity analysis approach that some unknown parameters had to be fixed in a certain range, and then gave a more insightful interpretation. Recently, Ning et al. [34]proposed a method to estimate all the unknown parameters. Whereas their method strongly relied on an imputation based on funnel-plot symmetry, our method does not.

Here, we would like to address some issues in clinical trial registries. Firstly, different clinical trial registries may give inconsistent information. A comparison of results between ClinicalTrials.gov and FDA database by Schwartz et al.[35]showed the planned sample size registered on the clinical trial registries were not always consistent with the final sample size for statistical analysis. Besides, as acknowledged by Fleming et al.[36] the status of the studies might be inconsistent between EUCTR and ClinicalTrials.gov, the studies registered on EUCTR which marked with “completed” may marked with “ongoing” in ClinicalTrials.gov. Secondly, no clinical trial registry could cover all the related studies[37, 38, 39]. Thirdly, it is hard to integrate multiple clinical trial registries automatically since unique identifier of a study is not available among registries (e.g. World Health Organization’s (WHO) International Clinical Trial Registry Platform (ICTRP)). Due to these issues of clinical trial registries, comprehensive search of multiple clinical trial registries was recommended in practice.

Clinical trial registries provide various information other than sample size. ClinicalTrials.gov’s search result contains 26 items in the list and analysis result if the investigator submitted, ICTRP contains 40 items in the result file, and FDA database provide much more details about the registered clinical trials. As acknowledged by several cross-sectional studies, information such as funding type (government, industry or academic) and region may possibly play important roles in the publication status[40]. In addition, information of ongoing studies are also available, although we did not use them in this paper. How to take a further utilization of the clinical trial registry would deserve more attention.

Acknowledgments

The last author’s research was partly supported by Grant-in-Aid for Challenging Exploratory Research (16K12403) and for Scientific Research (16H06299, 18H03208) from the Ministry of Education, Science, Sports and Technology of Japan.

References

- [1] Thornton A, Lee P. Publication bias in meta-analysis: its causes and consequences. *Journal of Clinical Epidemiology* 2000; 53(2): 207–216.
- [2] Sterne JA, Gavaghan D, Egger M. Publication and related bias in meta-analysis: power of statistical tests and prevalence in the literature. *Journal of Clinical Epidemiology* 2000; 53(11): 1119–1129.
- [3] Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; 315(7109): 629–634.
- [4] Macaskill P, Walter SD, Irwig L. A comparison of methods to detect publication bias in meta-analysis. *Statistics in Medicine* 2001; 20(4): 641–654.
- [5] Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994: 1088–1101.
- [6] Schwarzer G, Antes G, Schumacher M. A test for publication bias in meta-analysis with sparse binary data. *Statistics in Medicine* 2007; 26(4): 721–733.
- [7] Duval S, Tweedie R. A nonparametric “trim and fill” method of accounting for publication bias in meta-analysis. *Journal of the American Statistical Association* 2000; 95(449): 89–98.
- [8] Terrin N, Schmid CH, Lau J, Olkin I. Adjusting for publication bias in the presence of heterogeneity. *Statistics in Medicine* 2003; 22(13): 2113–2126.
- [9] Peters JL, Sutton AJ, Jones DR, Abrams KR, Rushton L. Performance of the trim and fill method in the presence of publication bias and between-study heterogeneity. *Statistics in Medicine* 2007; 26(25): 4544–4562.
- [10] Heckman JJ. Sample selection bias as a specification error. *Econometrica* 1979; 47(1): 153–61.
- [11] Copas J, Shi JQ. Meta-analysis, funnel plots and sensitivity analysis. *Biostatistics* 2000; 1(3): 247–262.
- [12] Copas J, Shi JQ. A sensitivity analysis for publication bias in systematic reviews. *Statistical Methods in Medical Research* 2001; 10(4): 251–265.
- [13] Copas JB. A likelihood-based sensitivity analysis for publication bias in meta-analysis. *Journal of the Royal Statistical Society: Series C (Applied Statistics)* 2013; 62(1): 47–66.
- [14] Copas J, Jackson D. A bound for publication bias based on the fraction of unpublished studies. *Biometrics* 2004; 60(1): 146–153.
- [15] Henmi M, Copas JB, Eguchi S. Confidence intervals and P-values for meta-analysis with publication bias. *Biometrics* 2007; 63(2): 475–482.

- [16] Carpenter JR, Schwarzer G, Rücker G, Künstler R. Empirical evaluation showed that the Copas selection model provided a useful summary in 80% of meta-analyses. *Journal of Clinical Epidemiology* 2009; 62(6): 624–631.
- [17] Schwarzer G, Carpenter J, Rücker G. Empirical evaluation suggests Copas selection model preferable to trim-and-fill method for selection bias in meta-analysis. *Journal of Clinical Epidemiology* 2010; 63(3): 282–288.
- [18] DeAngelis CD, Drazen JM, Frizelle FA, et al. Clinical trial registration: a statement from the International Committee of Medical Journal Editors. *Archives of Dermatology* 2005; 141(1): 76–77.
- [19] Viergever RF, Li K. Trends in global clinical trial registration: an analysis of numbers of registered clinical trials in different parts of the world from 2004 to 2013. *BMJ open* 2015; 5(9): e008932.
- [20] Ogino D, Takahashi K, Sato H. Characteristics of clinical trial websites: information distribution between ClinicalTrials.gov and 13 primary registries in the WHO registry network. *Trials* 2014; 15(1): 428.
- [21] Baudard M, Yavchitz A, Ravaud P, Perrodeau E, Boutron I. Impact of searching clinical trial registries in systematic reviews of pharmaceutical treatments: methodological systematic review and reanalysis of meta-analyses. *BMJ* 2017; 356: j448.
- [22] Jones CW, Keil LG, Weaver MA, Platts-Mills TF. Clinical trials registries are underutilized in the conduct of systematic reviews: a cross-sectional analysis. *Systematic Reviews* 2014; 3(1): 126.
- [23] Matsuoka N, Horio H, Hamada C. Improvement of Statistical Power to Detect Publication Bias in Meta-analysis Using the Clinical Trial Registration System. *Japanese Journal of Biometrics* 2011; 32(1): 13–31.
- [24] Mavridis D, Sutton A, Cipriani A, Salanti G. A fully Bayesian application of the Copas selection model for publication bias extended to network meta-analysis. *Statistics in Medicine* 2013; 32(1): 51–66.
- [25] Karner C, Chong J, Poole P. Tiotropium versus placebo for chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews* 2014(7).
- [26] Chen Y, Zhang Y, Tang Y, Huang X, Xie Y. High-maintenance-dose clopidogrel in patients undergoing percutaneous coronary intervention: a systematic review and meta-analysis. *PloS ONE* 2013; 8(10).
- [27] Copas J. What works?: Selectivity models and meta-analysis. *Journal of the Royal Statistical Society: Series A (Statistics in Society)* 1999; 162(1): 95–109.
- [28] Hartung J, Knapp G. On tests of the overall treatment effect in meta-analysis with normally distributed responses. *Statistics in Medicine* 2001; 20(12): 1771–1782.

- [29] Hartung J, Knapp G. A refined method for the meta-analysis of controlled clinical trials with binary outcome. *Statistics in Medicine* 2001; 20(24): 3875–3889.
- [30] Carpenter J, Rücker G, Schwarzer G. copas: An R package for fitting the Copas selection model. *The R Journal* 2009; 2: 31–6.
- [31] Kuss O. Statistical methods for meta-analyses including information from studies without any events—add nothing to nothing and succeed nevertheless. *Statistics in Medicine* 2015; 34(7): 1097–1116.
- [32] Follmann DA, Proschan MA. Valid inference in random effects meta-analysis. *Biometrics* 1999; 55(3): 732–737.
- [33] Knapp G, Hartung J. Improved tests for a random effects meta-regression with a single covariate. *Statistics in Medicine* 2003; 22(17): 2693–2710.
- [34] Ning J, Chen Y, Piao J. Maximum likelihood estimation and EM algorithm of Copas-like selection model for publication bias correction. *Biostatistics* 2017; 18(3): 495–504.
- [35] Schwartz LM, Woloshin S, Zheng E, Tse T, Zarin DA. ClinicalTrials.gov and Drugs@FDA: a comparison of results reporting for new drug approval trials. *Annals of Internal Medicine* 2016; 165(6): 421–430.
- [36] Fleminger J, Goldacre B. Prevalence of clinical trial status discrepancies: a cross-sectional study of 10,492 trials registered on both ClinicalTrials.gov and the European Union Clinical Trials Register. *PloS ONE* 2018; 13(3).
- [37] Glanville JM, Duffy S, McCool R, Varley D. Searching ClinicalTrials.gov and the International Clinical Trials Registry Platform to inform systematic reviews: what are the optimal search approaches?. *Journal of the Medical Library Association: JMLA* 2014; 102(3): 177.
- [38] Tse T, Fain KM, Zarin DA. How to avoid common problems when using ClinicalTrials.gov in research: 10 issues to consider. *BMJ* 2018; 361: k1452.
- [39] Adam GP, Springs S, Trikalinos T, et al. Does information from ClinicalTrials.gov increase transparency and reduce bias? Results from a five-report case series. *Systematic Reviews* 2018; 7(1): 59.
- [40] Loder E, Loder S, Cook S. Characteristics and publication fate of unregistered and retrospectively registered clinical trials submitted to The BMJ over 4 years. *BMJ Open* 2018; 8(2): e020037.




Figure 1: *Funnel plots of randomly selected published (filled circles)/unpublished (open circles) studies in simulated datasets with $N + M = 50$ under different settings of τ and ρ*




Figure 2: A funnel-plot for the tiotropium study being modified by adding information of the planned sample size of 8 unpublished studies with horizontal lines passing the y-axis at $\sqrt{n_i}$.




Figure 3: A funnel-plot for the clopidogrel study being modified by adding information of the planned sample size of 3 unpublished studies with horizontal lines passing the y-axis at $\sqrt{n_i}$.

Table 1: Simulation results for estimation of θ with $(\alpha_0, \alpha_1, \rho) = (-2.18, 0.2, -0.4)$ and $\tau = 0.05, 0.15$ or 0.30
(40% unpublished studies on average)

Scenario	Method	$N + M = 15$				$N + M = 25$				$N + M = 50$				$N + M = 100$			
		AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC
$\tau = 0.05$	REML	-0.285(0.094)	0.955	0.377	997	-0.282(0.067)	0.941	0.275	994	-0.280(0.046)	0.902	0.187	991	-0.277(0.032)	0.868	0.129	988
	REML.KnHa	-0.285(0.094)	0.944	0.413	997	-0.282(0.067)	0.931	0.286	994	-0.280(0.046)	0.892	0.186	991	-0.277(0.032)	0.861	0.128	988
	Copas	-0.273(0.099)	0.949	0.363	991	-0.267(0.075)	0.925	0.269	983	-0.264(0.058)	0.891	0.186	986	-0.255(0.048)	0.837	0.132	981
	MLE(N)	-0.257(0.112)	0.904	0.368	968	-0.251(0.077)	0.930	0.289	995	-0.249(0.054)	0.935	0.202	998	-0.248(0.035)	0.955	0.140	999
	MLE(T)	-0.257(0.112)	0.950	0.446	968	-0.251(0.077)	0.950	0.319	995	-0.249(0.054)	0.945	0.211	998	-0.248(0.035)	0.956	0.143	999
	MLE(SE^\ddagger)	-0.257(0.112)	0.952	0.483	968	-0.251(0.077)	0.957	0.330	995	-0.249(0.054)	0.947	0.213	998	-0.248(0.035)	0.957	0.144	999
$\tau = 0.15$	REML	-0.286(0.112)	0.920	0.420	999	-0.288(0.079)	0.913	0.310	993	-0.286(0.056)	0.893	0.216	994	-0.283(0.04)	0.854	0.153	997
	REML.KnHa	-0.286(0.112)	0.932	0.481	999	-0.288(0.079)	0.928	0.333	993	-0.286(0.056)	0.903	0.223	994	-0.283(0.04)	0.852	0.155	997
	Copas	-0.268(0.128)	0.873	0.392	984	-0.271(0.091)	0.879	0.299	986	-0.265(0.072)	0.862	0.215	993	-0.256(0.059)	0.825	0.156	996
	MLE(N)	-0.258(0.130)	0.849	0.403	963	-0.255(0.091)	0.899	0.325	990	-0.252(0.067)	0.921	0.237	995	-0.249(0.044)	0.943	0.171	1000
	MLE(T)	-0.258(0.130)	0.913	0.491	963	-0.255(0.091)	0.928	0.358	990	-0.252(0.067)	0.927	0.247	995	-0.249(0.044)	0.950	0.174	1000
	MLE(SE^\ddagger)	-0.258(0.13)	0.938	0.541	963	-0.255(0.091)	0.942	0.374	990	-0.252(0.067)	0.932	0.250	995	-0.249(0.044)	0.950	0.175	1000
$\tau = 0.30$	REML	-0.294(0.149)	0.905	0.544	999	-0.297(0.107)	0.899	0.414	999	-0.299(0.075)	0.888	0.296	1000	-0.295(0.054)	0.867	0.210	1000
	REML.KnHa	-0.294(0.149)	0.923	0.651	999	-0.297(0.107)	0.924	0.457	999	-0.299(0.075)	0.906	0.311	1000	-0.295(0.054)	0.881	0.216	1000
	Copas	-0.273(0.173)	0.852	0.508	967	-0.282(0.121)	0.866	0.412	982	-0.274(0.093)	0.861	0.294	993	-0.262(0.075)	0.843	0.223	996
	MLE(N)	-0.262(0.168)	0.827	0.529	947	-0.261(0.126)	0.876	0.442	984	-0.253(0.092)	0.915	0.334	994	-0.249(0.062)	0.948	0.244	999
	MLE(T)	-0.262(0.168)	0.884	0.643	947	-0.261(0.126)	0.900	0.488	984	-0.253(0.092)	0.921	0.349	994	-0.249(0.062)	0.958	0.249	999
	MLE(SE^\ddagger)	-0.262(0.168)	0.918	0.717	947	-0.261(0.126)	0.926	0.513	984	-0.253(0.092)	0.937	0.356	994	-0.249(0.062)	0.958	0.250	999
	True	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-

N,number of total studies;AVE, mean value of estimates;SD,standard error of estimates; CP, 95%confidence interval coverage probability; LOCI,length of confidence interval;NOC,number of converged cases.

Table 2: Simulation results for estimation of θ with $(\alpha_0, \alpha_1, \rho) = (-2.18, 0.2, -0.8)$ and $\tau = 0.05, 0.15$ or 0.30
(40% unpublished studies on average)

Scenario	Method	$N + M = 15$				$N + M = 25$				$N + M = 50$				$N + M = 100$			
		AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC
$\tau = 0.05$	REML	-0.320(0.097)	0.913	0.376	998	-0.313(0.068)	0.881	0.272	992	-0.307(0.046)	0.789	0.186	992	-0.304(0.033)	0.644	0.128	983
	REML.KnHa	-0.320(0.097)	0.917	0.404	998	-0.313(0.068)	0.874	0.275	992	-0.307(0.046)	0.769	0.182	992	-0.304(0.033)	0.615	0.125	983
	Copas	-0.291(0.105)	0.906	0.365	982	-0.276(0.090)	0.862	0.267	969	-0.249(0.081)	0.773	0.190	933	-0.220(0.066)	0.664	0.138	929
	MLE(N)	-0.264(0.100)	0.918	0.352	971	-0.254(0.071)	0.942	0.264	995	-0.250(0.050)	0.946	0.186	1000	-0.249(0.033)	0.956	0.130	1000
	MLE(T)	-0.264(0.100)	0.968	0.426	971	-0.254(0.071)	0.957	0.291	995	-0.250(0.050)	0.954	0.194	1000	-0.249(0.033)	0.958	0.132	1000
	MLE(SE^\ddagger)	-0.264(0.100)	0.973	0.466	971	-0.254(0.071)	0.962	0.307	995	-0.250(0.050)	0.965	0.200	1000	-0.249(0.033)	0.959	0.134	1000
$\tau = 0.15$	REML	-0.323(0.107)	0.870	0.400	995	-0.322(0.080)	0.841	0.303	993	-0.319(0.056)	0.745	0.213	993	-0.316(0.041)	0.603	0.151	997
	REML.KnHa	-0.323(0.107)	0.886	0.450	995	-0.322(0.080)	0.860	0.321	993	-0.319(0.056)	0.759	0.218	993	-0.316(0.041)	0.611	0.152	997
	Copas	-0.286(0.130)	0.840	0.380	979	-0.278(0.110)	0.794	0.295	964	-0.249(0.096)	0.725	0.215	957	-0.215(0.078)	0.660	0.163	926
	MLE(N)	-0.266(0.116)	0.888	0.369	973	-0.259(0.084)	0.909	0.293	991	-0.254(0.061)	0.924	0.216	996	-0.249(0.041)	0.937	0.157	998
	MLE(T)	-0.266(0.116)	0.934	0.443	973	-0.259(0.084)	0.939	0.323	991	-0.254(0.061)	0.934	0.226	996	-0.249(0.041)	0.939	0.160	998
	MLE(SE^\ddagger)	-0.266(0.116)	0.954	0.493	973	-0.259(0.084)	0.950	0.348	991	-0.254(0.061)	0.942	0.235	996	-0.249(0.041)	0.944	0.162	998
$\tau = 0.30$	REML	-0.346(0.141)	0.832	0.517	999	-0.343(0.109)	0.832	0.406	998	-0.342(0.076)	0.748	0.292	1000	-0.339(0.054)	0.604	0.208	1000
	REML.KnHa	-0.346(0.141)	0.878	0.608	999	-0.343(0.109)	0.874	0.447	998	-0.342(0.076)	0.775	0.306	1000	-0.339(0.054)	0.613	0.213	1000
	Copas	-0.308(0.170)	0.791	0.483	955	-0.297(0.138)	0.809	0.395	960	-0.268(0.116)	0.765	0.299	965	-0.224(0.099)	0.729	0.236	951
	MLE(N)	-0.285(0.164)	0.822	0.484	942	-0.270(0.120)	0.875	0.406	982	-0.259(0.085)	0.917	0.314	985	-0.251(0.058)	0.943	0.225	986
	MLE(T)	-0.285(0.164)	0.877	0.581	942	-0.270(0.120)	0.900	0.447	982	-0.259(0.085)	0.929	0.328	985	-0.251(0.058)	0.946	0.229	986
	MLE(SE^\ddagger)	-0.285(0.164)	0.912	0.659	942	-0.270(0.120)	0.930	0.484	982	-0.259(0.085)	0.946	0.340	985	-0.251(0.058)	0.951	0.233	986
	True	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-

N, number of total studies; AVE, mean value of estimates; SD, standard error of estimates; CP, 95% confidence interval coverage probability; LOCI, length of confidence interval; NOC, number of converged cases.

Table 3: Summary of the statistical analysis for publication bias evaluation of tiotropium study:
 # implies expected number of unpublished studies in the sensitivity analysis with the Copas selection model

Description	Method	#	OR	95% CI	P-value
No adjustment	REML	-	0.768	[0.697, 0.847]	<.000
	REML-KnHa	-	0.768	[0.691, 0.854]	<.000
Sensitivity analysis	Copas	1	0.779	[0.707, 0.858]	<.000
		5	0.795	[0.712, 0.886]	<.000
		8	0.803	[0.717, 0.898]	<.000
		11	0.808	[0.720, 0.907]	<.000
		13	0.811	[0.721, 0.912]	0.001
Proposed	MLE(N)	-	0.787	[0.710, 0.873]	<.000
	MLE(T)	-	0.787	[0.706, 0.878]	<.000

Table 4: Summary of the statistical analysis for publication bias evaluation of clopidogrel study:
 # implies expected number of unpublished studies in the sensitivity analysis with the Copas selection model

Description	Method	#	OR	95% CI	P-value
No adjustment	REML	-	0.579	[0.375, 0.892]	0.013
	REML-KnHa	-	0.579	[0.385, 0.871]	0.013
Sensitivity analysis	Copas	2	0.690	[0.501, 0.952]	0.024
		3	0.708	[0.514, 0.975]	0.034
		4	0.740	[0.537, 1.020]	0.066
		6	0.805	[0.503, 1.288]	0.365
		8	0.782	[0.437, 1.397]	0.406
Proposed	MLE(N)	-	0.692	[0.496, 0.967]	0.031
	MLE(T)	-	0.692	[0.476, 1.007]	0.054
	MLE($SE^\#$)	-	0.692	[0.460, 1.041]	0.073

Web-appendix to “Using clinical trial registries to inform Copas selection model for publication bias in meta-analysis”

Abstract

In Web-appendix A, we provide the datasets of the tiotropium study and the clopidogrel study with information of published/unpublished studies. In Web-appendix B we present the results of simulation studies with $(\alpha_0, \alpha_1) = (-1.24, 0.16)$ and $(-0.58, 0.13)$, respectively.

Web-appendix A: Tables for the datasets of the tiotropium study and the clopidogrel study

In this appendix, we presented the datasets of the tiotropium study and the clopidogrel study.

In Table S1, the dataset of the tiotropium study was presented. Studies from No.1 to No.22 came from the original meta-analysis by Karner et al.[1] which we introduced in Subsection 2.1. By searching with multiple clinical trial registries, we further identified 10 studies with the same search terms: *Tiotropium* or *Spiriva* or *HandiHaler* or *Respimat* as the original meta-analysis. The identified 10 studies referred to the studies as from No.23 to No.32. Recall that D_i is an indicator of publication status; $D_i = 1$ if published and $D_i = 0$ otherwise. Among these ten studies, study No.23 and No.24 reported estimated treatment effects in clinical trial registries. Therefore, they had the same indicator $D_i = 1$ as the original study No.1 to No.22. The rest of them (from No.25 to No.32) were deemed as unpublished studies with the indicator $D_i = 0$. The column of “Status” denotes the status of publication and registration. Studies marked with “published & regisitered” means we can identify these studies from both online literature database and clinical trial registries, while those marked with “published” or “registered” means they can only be identified from the online literature database or clinical trial registries.

In Table S2, the dataset of the clopidogrel study was presented. Similarly to the dataset of the tiotropium study, studies with $D_i = 1$ (from No.1 to No.12) came from the original meta-analysis by Chen et al. [2]. The studies with $D_i = 0$ were identified from the clinical trial registries.

Table S1: Tiotropium dataset

No.	Study	Tiotropium		Placebo		n_i	$\log OR_i$	s_i	D_i	Status
		Events	Total	Events	Total					
1	Bateman 2010a	685	1989	842	2002	3991	-0.33	0.07	1	published & registered
2	Bateman 2010b	495	1337	288	653	1990	-0.29	0.10	1	published & registered
3	Beeh 2006	180	1236	80	403	1639	-0.37	0.15	1	published & registered
4	Brusasco 2003	129	402	156	400	802	-0.30	0.15	1	published
5	Casaburi 2002	198	550	156	371	921	-0.25	0.14	1	published
6	Chan 2007	268	608	125	305	913	0.13	0.14	1	published & registered
7	Cooper 2010	112	260	102	259	519	0.15	0.18	1	published & registered
8	Covelli 2005	9	100	12	96	196	-0.37	0.47	1	published & registered
9	Dusser 2006	213	500	272	510	1010	-0.43	0.13	1	published & registered
10	Freeman 2007	19	200	35	195	395	-0.73	0.30	1	published & registered
11	Johansson 2008	2	107	4	117	224	-0.62	0.88	1	published & registered
12	Magnussen 2008	13	228	26	244	472	-0.67	0.35	1	published & registered
13	Moita 2008	6.00	147	6	164	311	0.11	0.59	1	published & registered
14	NCT00144326	11	123	12	127	250	-0.06	0.44	1	registered
15	Niewoehner 2005	255	914	296	915	1829	-0.21	0.10	1	published & registered
16	Powrie 2007	30	69	47	73	142	-0.84	0.34	1	published & registered
17	Sun 2007	0	30	2	30	60	-1.66	1.57	1	published
18	Tashkin 2008	2001	2987	2049	3006	5993	-0.05	0.06	1	published & registered
19	Tonnel 2008	101	266	130	288	554	-0.30	0.17	1	published & registered
20	Trooster 2011	11	238	24	219	457	-0.94	0.38	1	published & registered
21	Verkinde 2006	10	46	8	54	100	0.47	0.52	1	published
22	Voshaar 2008	43	360	21	181	541	0.03	0.28	1	published & registered
23	NCT01202188	85	480	60	232	712	-0.48	0.19	1	registered
24	NCT00929110	80	266	107	266	532	-0.45	0.18	1	registered
25	NCT00668772					207			0	registered
26	NCT00662740					220			0	registered
27	NCT00157235					234			0	registered
28	NCT00274521					108			0	registered
29	NCT02172287					623			0	registered
30	NCT02173691					584			0	registered
31	2007-001946-42-ES					844			0	registered
32	2005-000650-79					389			0	registered

Table S2: Clopidogrel dataset

No.	Study	High Dose		Standard Dose		n_i	$\log OR_i$	s_i	D_i	Status
		Events	Total	Events	Total					
1	Aradi 2012	1	36	8	38	74	-2.23	1.09	1	published & registered
2	DOUBLE 2010	0	24	1	24	48	-1.14	1.66	1	published
3	EFFICIENT 2011	2	47	8	47	94	-1.53	0.82	1	published & registered
4	GRAVITAS 2011	25	1109	25	1105	2214	-0.00	0.29	1	published & registered
5	Gremmel 2011	1	21	2	23	44	-0.64	1.26	1	published
6	Han 2009	4	403	9	410	813	-0.81	0.61	1	published
7	Ren LH 2012	6	46	10	55	101	-0.39	0.56	1	published
8	Roghani 2011	4	205	2	195	400	0.65	0.87	1	published
9	Tousek 2011	1	30	2	30	60	-0.73	1.25	1	published
10	VASP-02 2008	0	58	1	62	120	-1.05	1.64	1	published & registered
11	von Beckerath 2007	1	31	1	29	60	-0.07	1.44	1	published
12	Wang 2011	14	150	30	156	306	-0.84	0.35	1	published
13	NCT01069302					106			0	registered
14	NCT01371058					350			0	registered
15	NCT01102439					82			0	registered

Web-appendix B: Tables for simulation results with additional settings

In this appendix, we presented the results of simulation studies with $(\alpha_0, \alpha_1)=(-1.24,0.16)$ (Table S3 and Table S4) and $(-0.58,0.13)$ (Table S5 and Table S6), which resulted in the 27% and 19% unpublished studies on average, respectively. The findings of these two settings were similar with the results in Tables 1 and 2. We observed that our method had the smallest biases in all the scenarios in comparison to the REML and the Copas sensitivity analysis method. With larger number of studies ($N=50$ and 100), the confidence intervals of our method based on the normal approximation had empirical coverage probabilities close to the nominal level of 95%. With smaller number of studies ($N=15$ and 25), the modified confidence intervals could be useful to improve the coverage probabilities of our method when between-study variance was considerable.

References

- [1] Karner C, Chong J, Poole P. Tiotropium versus placebo for chronic obstructive pulmonary disease. *Cochrane database of systematic reviews* 2014(7).
- [2] Chen Y, Zhang Y, Tang Y, Huang X, Xie Y. High-maintenance-dose clopidogrel in patients undergoing percutaneous coronary intervention: a systematic review and meta-analysis. *PloS one* 2013; 8(10).

Table S3: Simulation results for estimation of θ with $(\alpha_0, \alpha_1, \rho) = (-1.24, 0.16, -0.4)$ and $\tau = 0.05, 0.15$ or 0.30
(27% unpublished studies on average)

Scenario	Method	$N + M = 15$				$N + M = 25$				$N + M = 50$				$N + M = 100$			
		AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC
$\tau = 0.05$	REML	-0.280(0.083)	0.957	0.353	991	-0.276(0.063)	0.942	0.262	997	-0.275(0.044)	0.915	0.179	988	-0.272(0.031)	0.891	0.124	989
	REML.KnHa	-0.280(0.083)	0.954	0.370	991	-0.276(0.063)	0.945	0.265	997	-0.275(0.044)	0.901	0.177	988	-0.272(0.031)	0.882	0.122	989
	Copas	-0.270(0.092)	0.944	0.344	994	-0.268(0.066)	0.943	0.257	992	-0.263(0.054)	0.896	0.178	991	-0.257(0.040)	0.870	0.126	989
	MLE(N)	-0.261(0.096)	0.927	0.351	972	-0.253(0.072)	0.932	0.272	994	-0.250(0.053)	0.930	0.191	999	-0.249(0.034)	0.952	0.135	999
	MLE(T)	-0.261(0.096)	0.954	0.402	972	-0.253(0.072)	0.948	0.293	994	-0.250(0.053)	0.935	0.198	999	-0.249(0.034)	0.954	0.138	999
	MLE(SE^\ddagger)	-0.261(0.096)	0.965	0.422	972	-0.253(0.072)	0.953	0.300	994	-0.250(0.053)	0.937	0.200	999	-0.249(0.034)	0.954	0.138	999
$\tau = 0.15$	REML	-0.287(0.099)	0.920	0.390	994	-0.280(0.074)	0.919	0.293	990	-0.281(0.053)	0.900	0.205	993	-0.278(0.038)	0.878	0.145	997
	REML.KnHa	-0.287(0.099)	0.925	0.426	994	-0.280(0.074)	0.923	0.307	990	-0.281(0.053)	0.908	0.209	993	-0.278(0.038)	0.880	0.146	997
	Copas	-0.276(0.109)	0.898	0.371	992	-0.267(0.085)	0.894	0.285	993	-0.263(0.070)	0.861	0.203	996	-0.259(0.049)	0.863	0.148	998
	MLE(N)	-0.264(0.120)	0.871	0.379	973	-0.255(0.085)	0.913	0.304	995	-0.252(0.064)	0.912	0.225	1000	-0.250(0.042)	0.941	0.163	1000
	MLE(T)	-0.264(0.120)	0.916	0.435	973	-0.255(0.085)	0.931	0.328	995	-0.252(0.064)	0.921	0.233	1000	-0.250(0.042)	0.944	0.165	1000
	MLE(SE^\ddagger)	-0.264(0.120)	0.934	0.466	973	-0.255(0.085)	0.946	0.340	995	-0.252(0.064)	0.924	0.235	1000	-0.250(0.042)	0.945	0.166	1000
$\tau = 0.30$	REML	-0.286(0.134)	0.917	0.505	998	-0.293(0.103)	0.906	0.389	999	-0.291(0.072)	0.895	0.277	1000	-0.288(0.051)	0.882	0.197	1000
	REML.KnHa	-0.286(0.134)	0.940	0.570	998	-0.293(0.103)	0.917	0.417	999	-0.291(0.072)	0.912	0.287	1000	-0.288(0.051)	0.887	0.200	1000
	Copas	-0.271(0.150)	0.872	0.479	985	-0.275(0.119)	0.868	0.378	996	-0.271(0.089)	0.864	0.274	997	-0.265(0.065)	0.866	0.201	998
	MLE(N)	-0.261(0.153)	0.859	0.503	962	-0.264(0.119)	0.885	0.413	985	-0.254(0.087)	0.930	0.313	995	-0.250(0.059)	0.942	0.228	999
	MLE(T)	-0.261(0.153)	0.905	0.577	962	-0.264(0.119)	0.905	0.445	985	-0.254(0.087)	0.943	0.324	995	-0.250(0.059)	0.943	0.232	999
	MLE(SE^\ddagger)	-0.261(0.153)	0.928	0.619	962	-0.264(0.119)	0.922	0.460	985	-0.254(0.087)	0.948	0.327	995	-0.250(0.059)	0.946	0.232	999
	True	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-

N, number of total studies; AVE, mean value of estimates; SD, standard error of estimates; CP, 95% confidence interval coverage probability; LOCI, length of confidence interval; NOC, number of converged cases.

Table S4: Simulation results for estimation of θ with $(\alpha_0, \alpha_1, \rho) = (-1.24, 0.16, -0.8)$ and $\tau = 0.05, 0.15$ or 0.30
(27% unpublished studies on average)

Scenario	Method	$N + M = 15$				$N + M = 25$				$N + M = 50$				$N + M = 100$			
		AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC
$\tau = 0.05$	REML	-0.304(0.084)	0.932	0.345	989	-0.302(0.063)	0.901	0.258	991	-0.297(0.043)	0.840	0.177	985	-0.295(0.031)	0.712	0.123	982
	REML.KnHa	-0.304(0.084)	0.912	0.348	989	-0.302(0.063)	0.881	0.251	991	-0.297(0.043)	0.811	0.168	985	-0.295(0.031)	0.669	0.116	982
	Copas	-0.281(0.097)	0.912	0.336	985	-0.274(0.081)	0.888	0.254	983	-0.259(0.069)	0.824	0.179	958	-0.234(0.058)	0.728	0.130	969
	MLE(N)	-0.262(0.088)	0.933	0.331	981	-0.255(0.068)	0.931	0.252	994	-0.251(0.047)	0.939	0.177	998	-0.249(0.031)	0.952	0.123	1000
	MLE(T)	-0.262(0.088)	0.958	0.379	981	-0.255(0.068)	0.952	0.271	994	-0.251(0.047)	0.945	0.184	998	-0.249(0.031)	0.956	0.125	1000
	MLE(SE^\sharp)	-0.262(0.088)	0.963	0.398	981	-0.255(0.068)	0.955	0.280	994	-0.251(0.047)	0.950	0.186	998	-0.249(0.031)	0.956	0.126	1000
$\tau = 0.15$	REML	-0.317(0.100)	0.902	0.380	994	-0.310(0.074)	0.870	0.286	989	-0.307(0.052)	0.800	0.201	992	-0.305(0.038)	0.664	0.142	994
	REML.KnHa	-0.317(0.100)	0.899	0.404	994	-0.310(0.074)	0.869	0.292	989	-0.307(0.052)	0.793	0.200	992	-0.305(0.038)	0.658	0.140	994
	Copas	-0.290(0.118)	0.874	0.366	989	-0.274(0.100)	0.832	0.280	988	-0.256(0.088)	0.760	0.201	978	-0.231(0.070)	0.690	0.151	964
	MLE(N)	-0.274(0.116)	0.884	0.361	968	-0.259(0.080)	0.906	0.282	995	-0.254(0.057)	0.918	0.207	993	-0.250(0.038)	0.938	0.148	999
	MLE(T)	-0.274(0.116)	0.928	0.414	968	-0.259(0.080)	0.925	0.303	995	-0.254(0.057)	0.928	0.214	993	-0.250(0.038)	0.941	0.151	999
	MLE(SE^\sharp)	-0.274(0.116)	0.939	0.444	968	-0.259(0.080)	0.937	0.318	995	-0.254(0.057)	0.938	0.218	993	-0.250(0.038)	0.943	0.152	999
$\tau = 0.30$	REML	-0.322(0.135)	0.878	0.482	1000	-0.328(0.099)	0.862	0.377	999	-0.327(0.070)	0.797	0.272	1000	-0.324(0.051)	0.668	0.193	1000
	REML.KnHa	-0.322(0.135)	0.899	0.537	1000	-0.328(0.099)	0.877	0.401	999	-0.327(0.070)	0.808	0.279	1000	-0.324(0.051)	0.675	0.194	1000
	Copas	-0.290(0.156)	0.824	0.459	968	-0.288(0.130)	0.804	0.369	985	-0.270(0.106)	0.779	0.275	991	-0.239(0.091)	0.726	0.213	990
	MLE(N)	-0.272(0.149)	0.851	0.464	975	-0.270(0.109)	0.882	0.384	986	-0.259(0.079)	0.931	0.294	994	-0.251(0.054)	0.946	0.211	991
	MLE(T)	-0.272(0.149)	0.893	0.531	975	-0.270(0.109)	0.906	0.413	986	-0.259(0.079)	0.939	0.304	994	-0.251(0.054)	0.950	0.215	991
	MLE(SE^\sharp)	-0.272(0.149)	0.917	0.578	975	-0.270(0.109)	0.924	0.433	986	-0.259(0.079)	0.947	0.311	994	-0.251(0.054)	0.952	0.217	991
	True	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-

N,number of total studies;AVE, mean value of estimates;SD,standard error of estimates; CP, 95%confidence interval coverage probability; LOCI,length of confidence interval;NOC,number of converged cases.

Table S5: Simulation results for estimation of θ with $(\alpha_0, \alpha_1, \rho) = (-0.58, 0.13, -0.4)$ and $\tau = 0.05, 0.15$ or 0.30
(19% unpublished studies on average)

Scenario	Method	$N + M = 15$				$N + M = 25$				$N + M = 50$				$N + M = 100$			
		AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC
$\tau = 0.05$	REML	-0.274(0.082)	0.960	0.346	994	-0.268(0.063)	0.941	0.254	989	-0.271(0.043)	0.932	0.176	987	-0.268(0.031)	0.911	0.122	985
	REML.KnHa	-0.274(0.082)	0.945	0.357	994	-0.268(0.063)	0.934	0.254	989	-0.271(0.043)	0.922	0.172	987	-0.268(0.031)	0.903	0.119	985
	Copas	-0.265(0.088)	0.951	0.337	996	-0.261(0.067)	0.928	0.250	997	-0.261(0.051)	0.908	0.174	992	-0.258(0.036)	0.898	0.122	995
	MLE(N)	-0.261(0.095)	0.928	0.343	969	-0.253(0.071)	0.929	0.260	996	-0.253(0.050)	0.934	0.186	998	-0.250(0.033)	0.944	0.132	1000
	MLE(T)	-0.261(0.095)	0.954	0.386	969	-0.253(0.071)	0.940	0.277	996	-0.253(0.050)	0.943	0.192	998	-0.250(0.033)	0.951	0.134	1000
	MLE(SE^\ddagger)	-0.261(0.095)	0.961	0.403	969	-0.253(0.071)	0.942	0.283	996	-0.253(0.050)	0.944	0.194	998	-0.250(0.033)	0.951	0.134	1000
$\tau = 0.15$	REML	-0.274(0.096)	0.940	0.378	995	-0.269(0.074)	0.929	0.286	986	-0.275(0.052)	0.912	0.200	991	-0.272(0.037)	0.904	0.142	997
	REML.KnHa	-0.274(0.096)	0.932	0.406	995	-0.269(0.074)	0.933	0.295	986	-0.275(0.052)	0.917	0.203	991	-0.272(0.037)	0.904	0.142	997
	Copas	-0.264(0.103)	0.916	0.363	995	-0.258(0.082)	0.905	0.278	993	-0.263(0.064)	0.883	0.203	999	-0.260(0.044)	0.892	0.143	996
	MLE(N)	-0.256(0.108)	0.905	0.371	959	-0.251(0.082)	0.898	0.294	994	-0.255(0.061)	0.917	0.216	999	-0.251(0.041)	0.937	0.159	1000
	MLE(T)	-0.256(0.108)	0.935	0.418	959	-0.251(0.082)	0.922	0.314	994	-0.255(0.061)	0.925	0.223	999	-0.251(0.041)	0.938	0.162	1000
	MLE(SE^\ddagger)	-0.256(0.108)	0.949	0.442	959	-0.251(0.082)	0.931	0.323	994	-0.255(0.061)	0.927	0.225	999	-0.251(0.041)	0.938	0.162	1000
$\tau = 0.30$	REML	-0.280(0.125)	0.917	0.480	997	-0.284(0.100)	0.914	0.375	999	-0.283(0.069)	0.909	0.268	1000	-0.280(0.050)	0.896	0.191	1000
	REML.KnHa	-0.280(0.125)	0.927	0.531	997	-0.284(0.100)	0.924	0.398	999	-0.283(0.069)	0.919	0.277	1000	-0.280(0.050)	0.898	0.193	1000
	Copas	-0.264(0.143)	0.880	0.460	989	-0.272(0.109)	0.881	0.366	998	-0.269(0.082)	0.880	0.266	998	-0.264(0.059)	0.879	0.193	1000
	MLE(N)	-0.261(0.142)	0.878	0.472	961	-0.265(0.113)	0.889	0.390	982	-0.257(0.083)	0.926	0.297	996	-0.252(0.058)	0.931	0.220	1000
	MLE(T)	-0.261(0.142)	0.904	0.531	961	-0.265(0.113)	0.908	0.416	982	-0.257(0.083)	0.932	0.306	996	-0.252(0.058)	0.936	0.223	1000
	MLE(SE^\ddagger)	-0.261(0.142)	0.922	0.564	961	-0.265(0.113)	0.916	0.428	982	-0.257(0.083)	0.937	0.309	996	-0.252(0.058)	0.937	0.223	1000
	True	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-

N,number of total studies;AVE, mean value of estimates;SD,standard error of estimates; CP, 95%confidence interval coverage probability; LOCI,length of confidence interval;NOC,number of converged cases.

Table S6: Simulation results for estimation of θ with $(\alpha_0, \alpha_1, \rho) = (-0.58, 0.13, -0.8)$ and $\tau = 0.05, 0.15$ or 0.30
(19% unpublished studies on average)

Scenario	Method	$N + M = 15$				$N + M = 25$				$N + M = 50$				$N + M = 100$			
		AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC	AVE(SD)	CP	LOCI	NOC
$\tau = 0.05$	REML	-0.297(0.083)	0.939	0.340	989	-0.289(0.058)	0.929	0.253	990	-0.288(0.042)	0.870	0.173	985	-0.286(0.030)	0.791	0.120	980
	REML.KnHa	-0.297(0.083)	0.917	0.332	989	-0.289(0.058)	0.917	0.240	990	-0.288(0.042)	0.841	0.162	985	-0.286(0.030)	0.750	0.112	980
	Copas	-0.281(0.090)	0.932	0.333	995	-0.271(0.072)	0.913	0.250	989	-0.264(0.061)	0.854	0.174	974	-0.248(0.052)	0.774	0.123	962
	MLE(N)	-0.267(0.090)	0.932	0.327	976	-0.255(0.066)	0.941	0.249	995	-0.253(0.047)	0.936	0.173	999	-0.250(0.031)	0.956	0.120	1000
	MLE(T)	-0.267(0.090)	0.958	0.368	976	-0.255(0.066)	0.958	0.266	995	-0.253(0.047)	0.941	0.178	999	-0.250(0.031)	0.958	0.122	1000
	MLE(SE^\ddagger)	-0.267(0.090)	0.966	0.382	976	-0.255(0.066)	0.962	0.272	995	-0.253(0.047)	0.944	0.180	999	-0.250(0.031)	0.958	0.123	1000
$\tau = 0.15$	REML	-0.295(0.097)	0.913	0.368	988	-0.293(0.072)	0.902	0.279	987	-0.296(0.051)	0.839	0.195	989	-0.294(0.037)	0.756	0.138	992
	REML.KnHa	-0.295(0.097)	0.915	0.381	988	-0.293(0.072)	0.899	0.278	987	-0.296(0.051)	0.834	0.192	989	-0.294(0.037)	0.733	0.134	992
	Copas	-0.276(0.109)	0.877	0.359	988	-0.267(0.094)	0.849	0.272	986	-0.263(0.078)	0.802	0.195	990	-0.248(0.061)	0.754	0.143	973
	MLE(N)	-0.265(0.105)	0.883	0.352	963	-0.256(0.081)	0.904	0.277	993	-0.257(0.057)	0.916	0.200	1000	-0.251(0.038)	0.932	0.144	1000
	MLE(T)	-0.265(0.105)	0.915	0.396	963	-0.256(0.081)	0.930	0.296	993	-0.257(0.057)	0.924	0.206	1000	-0.251(0.038)	0.936	0.146	1000
	MLE(SE^\ddagger)	-0.265(0.105)	0.929	0.417	963	-0.256(0.081)	0.936	0.305	993	-0.257(0.057)	0.932	0.209	1000	-0.251(0.038)	0.940	0.146	1000
$\tau = 0.30$	REML	-0.310(0.129)	0.887	0.470	994	-0.312(0.096)	0.885	0.362	998	-0.312(0.068)	0.838	0.262	1000	-0.309(0.049)	0.761	0.186	1000
	REML.KnHa	-0.310(0.129)	0.901	0.513	994	-0.312(0.096)	0.891	0.379	998	-0.312(0.068)	0.840	0.266	1000	-0.309(0.049)	0.756	0.186	1000
	Copas	-0.284(0.149)	0.847	0.450	982	-0.286(0.119)	0.851	0.356	994	-0.273(0.097)	0.803	0.261	996	-0.256(0.077)	0.779	0.195	994
	MLE(N)	-0.278(0.145)	0.848	0.459	961	-0.272(0.108)	0.887	0.375	986	-0.261(0.077)	0.920	0.277	990	-0.252(0.052)	0.942	0.201	994
	MLE(T)	-0.278(0.145)	0.889	0.517	961	-0.272(0.108)	0.901	0.401	986	-0.261(0.077)	0.932	0.285	990	-0.252(0.052)	0.950	0.204	994
	MLE(SE^\ddagger)	-0.278(0.145)	0.911	0.550	961	-0.272(0.108)	0.916	0.414	986	-0.261(0.077)	0.936	0.289	990	-0.252(0.052)	0.950	0.205	994
	True	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-	-0.250	-	-	-

N,number of total studies;AVE, mean value of estimates;SD,standard error of estimates; CP, 95%confidence interval coverage probability; LOCI,length of confidence interval;NOC,number of converged cases.