

No signed graph with the nullity

$$\eta(G, \sigma) = |V(G)| - 2m(G) + 2c(G) - 1^*$$

Yong Lu[†], Jingwen Wu

School of Mathematics and Statistics, Jiangsu Normal University,
Xuzhou, Jiangsu 221116, People's Republic of China.

E-mail: luyong@jsnu.edu.com

Abstract

Let $G^\sigma = (G, \sigma)$ be a signed graph and $A(G, \sigma)$ be its adjacency matrix. Denote by $m(G)$ the matching number of G . Let $\eta(G, \sigma)$ be the nullity of (G, σ) . He et al. [Bounds for the matching number and cyclomatic number of a signed graph in terms of rank, Linear Algebra Appl. 572 (2019), 273–291] proved that

$$|V(G)| - 2m(G) - c(G) \leq \eta(G, \sigma) \leq |V(G)| - 2m(G) + 2c(G),$$

where $c(G)$ is the dimension of cycle space of G . Signed graphs reaching the lower bound or the upper bound are respectively characterized by the same paper. In this paper, we will prove that no signed graphs with nullity $|V(G)| - 2m(G) + 2c(G) - 1$. We also prove that there are infinite signed graphs with nullity $|V(G)| - 2m(G) + 2c(G) - s$, ($0 \leq s \leq 3c(G), s \neq 1$) for a given $c(G)$.

Key Words: Signed graph, nullity, matching number, the dimension of cycle space.

AMS Subject Classification (2010): 05C35; 05C50.

1 Introduction

Let $G = (V(G), E(G))$ be a simple graph, where $V(G)$ is the vertex set of G and $E(G)$ is the edge set of G . Let $V(G) = \{v_1, v_2, \dots, v_n\}$. Then the *adjacency matrix* $A(G)$ of G is the symmetric $n \times n$ matrix with entries $A(i, j) = 1$ (or written as $a_{ij} = 1$) if and only if $v_i v_j \in E(G)$ and zeros elsewhere. The *nullity* (resp. *rank*) of G , denoted by $\eta(G)$ (resp. $r(G)$), is the multiplicity of 0 (resp. non-zero) eigenvalue of $A(G)$. A *pendant vertex* is defined as a vertex with degree one. The unique neighbour of the pendant vertex is called *quasi-pendant vertex*. If a cycle of G contains only a vertex of degree three, and the remaining vertices are of degree two, then the cycle is called a *pendant cycle*.

*This work is supported by the National Natural Science Foundations of China (No. 11901253), the Natural Science Foundation for Colleges and Universities in Jiangsu Province of China (No. 19KJB110009), and the Science Foundation of Jiangsu Normal University (No. 18XLRX021).

[†]Corresponding author.

Assume that $V_1 \subseteq V(G)$ and $V_1 \neq \emptyset$. Let $G[V_1]$ be the *induced subgraph* of G , whose vertex set is V_1 and edge set is the set of those edges of G that have both ends in V_1 . Denoted by $G - V_1$ the induced subgraph obtained from G by deleting each vertex in V_1 and its incident edges. For convenience, if $V_1 = \{x\}$, we write $G - x$ for $G - \{x\}$. For the induced subgraph H_1 of G , $H_1 + x$ is defined as the subgraph of G induced by the vertex set $V(H_1) \cup \{x\}$.

The *matching number* of G , denoted by $m(G)$, is the size of a maximum matching of G . Let M be a maximum matching of G , if there exists the edge $e \in M$ such that the vertex $v \in e$, v is called *M -saturated*, otherwise, v is called *M -unsaturated*. An *M -alternating path* of G is defined as a path whose edges are alternately in the edge sets $E \setminus M$ and M . An *M -augmenting path* is defined as a path whose starting vertex and ending vertex are M -unsaturated.

The *distance* between u and v , denote by $d(u, v)$, is the length of the shortest path from the vertex u to v . The length of the shortest cycle of G is called the *girth* of G , denote by $g(G)$. Let $c(G)$ be the *dimension of cycle space* of G , $c(G) = |E(G)| - |V(G)| + \theta(G)$, where $\theta(G)$ is the number of connected components of G . If the cycles (if any) of G are pairwise vertex-disjoint, the acyclic graph T_G is obtained from contracting each cycle of G into a vertex, which is called *cyclic vertex*. Let W_G (resp. U) be the vertex set consisting of all cyclic vertices (resp. all non-cyclic vertices) in T_G , $V(T_G) = W_G \cup U$. Furthermore, the graph is obtained from T_G by deleting all cyclic vertices, denoted by $[T_G]$.

A *signed graph* $G^\sigma = (G, \sigma)$ consists of a simple graph G with edge set E and a mapping $\sigma : E \rightarrow \{+, -\}$. G is called the *underlying graph* of (G, σ) . The *adjacency matrix* of (G, σ) , denoted by $A(G, \sigma) = a_{ij}^\sigma = \sigma(v_i v_j) a_{ij}$, where $a_{ij} \in A(G)$. We use $r(G, \sigma)$ to denote the *rank* of a signed graph (G, σ) .

Denote by C_n^σ a signed cycle of order n . The *sign* $sgn(C_n^\sigma)$ of C_n^σ is defined as $\prod_{e \in E(C_n^\sigma)} \sigma(e)$. If $sgn(C_n^\sigma) = +$ (or $sgn(C_n^\sigma) = -$), then C_n^σ is said to be *positive* (or *negative*). If all the cycles of (G, σ) are positive, then (G, σ) is *balanced*, and *unbalanced* otherwise.




Figure 1: $\infty(p, 1, q)$, $\infty(p, l, q)$ and $\theta(p, l, q)$.

A connected graph G is called *bicyclic* if $|E(G)| = |V(G)| + 1$. The *base* of a bicyclic graph G is the unique connected bicyclic subgraph of G containing no pendant vertices.

Let $\infty(p, l, q)$ (see Figure 1) be the graph obtained from C_p , C_q and P_l by identifying v with v_1 and u with v_l , respectively. A bicyclic graph G is called an ∞ -graph, if G contains $\infty(p, l, q)$ as its base. Let P_{p+2} , P_{q+2} , P_{l+2} be three paths, where $\min\{p, l, q\} \geq 0$ and at most one of p, l, q is 0. Let $\theta(p, l, q)$ (see Figure 1) be the graph obtained from P_{p+2} , P_{l+2} and P_{q+2} by identifying the three initial vertices and terminal vertices. The bicyclic graph containing $\theta(p, l, q)$ as its base is called a θ -graph.

A connected graph G is called *tricyclic* if $|E(G)| = |V(G)| + 2$. As shown in Figure 2, there are eight types of bases of tricyclic graph, denoted by T_i , $i = 1, \dots, 8$. The tricyclic graph G can be obtained from T_i by attaching trees to its vertices.

Collatz et al. [4] had wanted to obtain all graphs of order n with $\eta(G) > 0$. Until today, this problem is also unsolved. In recent years, the investigation on the nullity (or




Figure 2: T_1 – T_8

rank) of simple graphs is an significant topic in the theory of graph spectra. In 2014, Wang and Wong [15] investigated the relationship between the nullity of the simple graph and the matching number, and proved that $|V(G)| - 2m(G) - c(G) \leq \eta(G) \leq |V(G)| - 2m(G) + 2c(G)$. In 2015, Song, Song and Tam [13] characterized the graphs with the nullity $\eta(G) = |V(G)| - 2m(G) + 2c(G)$. In 2016, Wang [14] characterized the graphs with nullity $\eta(G) = |V(G)| - 2m(G) - c(G)$. In 2019, Li and Guo [8] proved that there is no graph with nullity $\eta(G) = |V(G)| - 2m(G) + 2c(G) - 1$, and for fixed $c(G)$, infinitely many connected graphs with nullity $\eta(G) = |V(G)| - 2m(G) + 2c(G) - s$, where $0 \leq s \leq 3c(G)$ and $s \neq 1$, are also constructed.

In 2018, Lu, Wang and Zhou [10] proved that $\eta(G) - 2c(G) \leq \eta(G, \sigma) \leq \eta(G) + 2c(G)$ for an unbalanced signed graph with $c(G) \geq 1$, and all corresponding extremal graphs are characterized. In 2019, He, Hao and Lai [6] proved that for every signed graph, $|V(G)| - 2m(G) - c(G) \leq \eta(G, \sigma) \leq |V(G)| - 2m(G) + 2c(G)$. Moreover, signed graphs reaching the lower bound or the upper bound are respectively characterized in [6]. In this paper, we prove that no signed graph with nullity $\eta(G, \sigma) = |V(G)| - 2m(G) + 2c(G) - 1$, and for fixed $c(G)$, infinitely many connected signed graphs with nullity $\eta(G, \sigma) = |V(G)| - 2m(G) + 2c(G) - s$, where $0 \leq s \leq 3c(G)$ and $s \neq 1$, are also constructed.

Our main results are the following three theorems.

Theorem 1.1. *Suppose (G, σ) is an unbalanced signed unicyclic graph with an unique cycle C_q^σ , then*

- (1) $\eta(G, \sigma) = |V(G)| - 2m(G) - 1$, if $q \equiv 1 \pmod{2}$, $m([T_G]) = m(T_G)$ (Lemma 4.3 [6]);
- (2) $\eta(G, \sigma) = |V(G)| - 2m(G) + 2$, if $q \equiv 2 \pmod{4}$, $m([T_G]) = m(T_G)$ (Lemma 5.3 [6]);

(3) $\eta(G, \sigma) = |V(G)| - 2m(G)$, otherwise.

Theorem 1.2. For any unbalanced signed graph (G, σ) ,

$$\eta(G, \sigma) \neq |V(G)| - 2m(G) + 2c(G) - 1.$$

Theorem 1.3. Let (G, σ) be an unbalanced signed graph. If $c(G)$ is fixed, then there exists infinitely connected unbalanced signed graphs, such that $\eta(G, \sigma) = |V(G)| - 2m(G) + 2c(G) - s$, where $0 \leq s \leq 3c(G)$, $s \neq 1$.

2 Preliminaries

Lemma 2.1. [2] A matching M of G is a maximum matching if and only if G contains no M -augmenting path.

Lemma 2.2. [11] Let G be a graph with a pendant vertex u , and v is the unique quasi-pendant vertex of u . Then $m(G) = 1 + m(G - v) = 1 + m(G - u - v)$.

Lemma 2.3. [3] Let G be a simple graph. Then $m(G) - 1 \leq m(G - v) \leq m(G)$, for any vertex $v \in V(G)$.

Lemma 2.4. [11] Let G be a graph obtained by joining a vertex of an even cycle C by an edge to a vertex y of a connected graph H . Then $m(G) = m(C) + m(H)$.

Lemma 2.5. [8] For any undirected graph G , $\eta(G) \neq |V(G)| - 2m(G) + 2c(G) - 1$.

Lemma 2.6. [16] Let G be a graph with $x \in V(G)$,

- (1) $c(G - x) = c(G)$ if x lies outside any cycle of G ;
- (2) $c(G - x) \leq c(G) - 1$ if x lies on a cycle of G ;
- (3) $c(G - x) \leq c(G) - 2$ if x is a common vertex of distinct cycles of G ;
- (4) if the cycles of G are pairwise vertex-disjoint, then $c(G)$ precisely equals the number of cycles in G .

Lemma 2.7. [1] Let x be a vertex of (G, σ) . Then $\eta(G, \sigma) - 1 \leq \eta(G - x, \sigma) \leq \eta(G, \sigma) + 1$.

Lemma 2.8. [5] Let (G, σ) be a signed graph containing a pendant vertex u , $uv \in E(G)$ and $(H, \sigma) = (G, \sigma) - u - v$. Then $\eta(G, \sigma) = \eta(H, \sigma)$.

Lemma 2.9. [6] Let (G, σ) be a connected signed graph. Then $|V(G)| - 2m(G) - c(G) \leq \eta(G, \sigma) \leq |V(G)| - 2m(G) + 2c(G)$.

Lemma 2.10. [5] Let P_n^σ be a signed path. Then,

$$\eta(P_n^\sigma) = \begin{cases} 0, & \text{if } n \equiv 0 \pmod{2}; \\ 1, & \text{if } n \equiv 1 \pmod{2}. \end{cases}$$

Lemma 2.11. [5] Let C_p^σ be a signed cycle. Then,

$$\eta(C_p^\sigma) = \begin{cases} 0, & \text{if } p \equiv 1(\text{mod } 2); \\ 0, & \text{if } p \equiv 0(\text{mod } 4) \text{ and } \sigma(C_p) = -; \\ 0, & \text{if } p \equiv 2(\text{mod } 4) \text{ and } \sigma(C_p) = +; \\ 2, & \text{if } p \equiv 0(\text{mod } 4) \text{ and } \sigma(C_p) = +; \\ 2, & \text{if } p \equiv 2(\text{mod } 4) \text{ and } \sigma(C_p) = -. \end{cases}$$

Lemma 2.12. [7] Let (G, σ) be a signed graph. Then (G, σ) is balanced if and only if $(G, \sigma) \sim (G, +)$.

Lemma 2.13. [10] Let (G, σ) be a signed graph, if $(G_1, \sigma), (G_2, \sigma), \dots, (G_t, \sigma)$ are all the connected components of (G, σ) . Then $\eta(G, \sigma) = \sum_{i=1}^t \eta(G_i, \sigma)$.

Lemma 2.14. [6] Let (G, σ) be a connnected signed graph. Then $\eta(G, \sigma) = |V(G)| - 2m(G) + 2c(G)$ if and only if all the following conditions hold for (G, σ) :

- (1) the cycles (if any) of (G, σ) are pairwise vertex-disjoint.
- (2) for each cycle (if any) C_q^σ of (G, σ) , either $q \equiv 0(\text{mod } 4)$ and $\sigma(C_q) = +$ or $q \equiv 2(\text{mod } 4)$ and $\sigma(C_q) = -$.
- (3) $m(T_G) = m([T_G])$.

3 Proof of Theorem 1.1

In this section, we will proved that there is no unbalanced signed unicyclic graph with the nullity $\eta(G, \sigma) = |V(G)| - 2m(G) + 2c(G) - 1$. In order to proof Theorem 1.1, we need to introduce some definitions and lemmas.

In simple graph, we introduced the concept of elementary graph, either the complete graph with order two K_2 or the cycle C_q , $q \geq 3$. If the graph whose each component is elementary graph, the graph is called the basic graph. The set of signed graphs, denoted by \mathbf{U}_i , is the set of all basic graphs with i vertices in (G, σ) . For any basic graph $(u, \sigma) \in \mathbf{U}_i$, $p(u)$, $c(u)$ and $s(u)$ are defined as the number of components, the number of cycles and the number of the negative edges in the cycle of (u, σ) , respectively.

Definition 3.1. Let G be an unicyclic graph with an unique cycle C_q . Then,

- (1) E_1 : The edge set of G between C_q and $[T_G]$.
- (2) F_1 : The matching set of G with $m(G)$ edges.
- (3) F_2 : The matching set of $[T_G]$ with $m([T_G])$ edges.
- (4) F'_1 : The matching set of G with $m(G)$ edges, each of which has at least one edge in E_1 .
- (5) F''_1 : The matching set of G with $m(G)$ edges, and $M \cap E_1 = \phi$ for all $M \in F_1$.

By Definition 3.1, we have $F_1 = F_1' \cup F_1''$. We can obtain the following corollary:

Corollary 3.2. *Let C_q be an even cycle. Then,*

- (1) *If $F_1' = \phi$, the maximum matching of G is the union of a maximum matching of C_q and a maximum matching of $G - C_q$. Then $|F_1| = |F_1''| = 2|F_2|$.*
- (2) *If $F_1' \neq \phi$, then $|F_1| = |F_1'| + |F_1''| > 2|F_2|$.*

Lemma 3.3. [9] *Let (G, σ) be a signed graph on n vertices and $P_{(G, \sigma)}(\lambda) = \sum_{i=0}^n a_i \cdot \lambda^{n-i}$ be a characteristic polynomial of $A(G, \sigma)$. Then $a_i = \sum_{(u, \sigma) \in \mathbf{U}_1} (-1)^{p(u)+s(u)} \cdot 2^{c(u)}$, $i \in \{1, 2, \dots, n\}$.*

Lemma 3.4. [6] *Let (G, σ) be a connected unbalanced signed unicyclic graph whose unique cycle is C_q^σ . Then $\eta(G, \sigma) = |V(G)| - 2m(G) + 2$ if and only if $q \equiv 2 \pmod{4}$ and $m(T_G) = m([T_G])$.*

Lemma 3.5. [6] *Let (G, σ) be an unicyclic signed graph which contains the unique signed cycle C_q^σ . Then $\eta(G, \sigma) = |V(G)| - 2m(G) - 1$ if and only if q is odd and $m(T_G) = m([T_G])$.*

Lemma 3.6. [12] *Let G be a graph with at least one cycle. Suppose that all cycles of G are pairwise vertex-disjoint and each cycle is odd, then $m(T_G) = m([T_G])$ if and only if $m(G) = \sum_{C \in \mathcal{L}(G)} m(C) + m([T_G])$, where $\mathcal{L}(G)$ denotes the set of all cycles in G .*

By Lemma 3.6, we can obtain the following corollary.

Corollary 3.7. *Let G be an unicyclic graph with the unique cycle C_q , q is odd. Then $m(T_G) = m([T_G])$ if and only if $m(G) = m([T_G]) + m(C_q)$.*

Next, for the unicyclic graph with an even cycle, similar to Lemma 3.6, we give the following lemma.

Lemma 3.8. *Let G be an unicyclic graph with a cycle C_q , q is even. Then $m(T_G) = m([T_G])$ if and only if $m(G) = m(C_q) + m([T_G])$ and $M \cap E_1 = \phi$ for all $M \in F_1$.*

Proof. Suppose that $G - C_q$ has p connected components, says S_1, S_2, \dots, S_p . Since $[T_G] = G - C_q$, then S_1, S_2, \dots, S_p are also all connected components of $[T_G]$. Assume that the vertex v_c of T_G is correspond to the cycle C_q of G , which implies that $W_G = \{v_c\}$ in T_G .

In T_G , let \widetilde{S}_k ($k = 1, \dots, p$) be the subgraph induced by $V(S_k)$ and all vertices adjacent to it. Similarly, in G , let \widehat{S}_k be the subgraph induced by $V(S_k)$ and all vertices adjacent to it. Since G is an unicyclic graph, then there exists at most an edge between $V(S_k)$ and $V(C_q)$, thus for all $k = 1, \dots, p$, the subgraph \widetilde{S}_k and \widehat{S}_k are isomorphic.

Necessity: Note that $m(T_G) = m([T_G])$, and $m([T_G]) = \sum_{k=1}^p m(S_k)$. Then each maximum matching of $[T_G]$ is the maximum matching of T_G , denoted by M_T the maximum matching of $[T_G]$, thus $|M_T| = \sum_{k=1}^p m(S_k)$. We can prove $m(S_k) = m(\widetilde{S}_k)$ for $k = 1, \dots, p$. Otherwise, there exists $m(S_{k_0}) < m(\widetilde{S}_{k_0})$, where $k_0 \in \{1, \dots, p\}$. For M_T , replace the maximum matching of S_{k_0} to the maximum matching of \widetilde{S}_{k_0} , we obtained a matching of T_G , denoted by M_T' . Then $|M_T'| = m(\widetilde{S}_{k_0}) + \sum_{k \neq k_0} m(\widetilde{S}_k) > m(S_{k_0}) + \sum_{k \neq k_0} m(S_k) = |M_T|$.

This contradicts $|M_T| = m(T_G) \geq |M_T'|$. Hence $m(S_k) = m(\widetilde{S}_k)$, for all $k = 1, \dots, p$. Since the subgraph \widetilde{S}_k and \widehat{S}_k are isomorphic, then $m(\widehat{S}_k) = m(S_k) = m(\widetilde{S}_k)$, for all $k = 1, \dots, p$.

Let M_1 be a maximum matching of G . Then $m(G) = |M_1| = |M_1 \cap E(C_q)| + \sum_{k=1}^p |M_1 \cap E(\widehat{S}_k)| \leq m(C_q) + \sum_{k=1}^p m(\widehat{S}_k) = m(C_q) + \sum_{k=1}^p m(S_k)$. By $m(T_G) = m([T_G])$, we have $m(G) \geq m(C_q) + \sum_{k=1}^p m(S_k)$.

Thus

$$m(G) = m(C_q) + \sum_{k=1}^p m(S_k),$$

that is,

$$m(G) = m(C_q) + m([T_G]).$$

Next, we need to prove $M \cap E_1 = \phi$ for any maximum matching M of G , $M \in F_1$. Suppose on the contrary, there exists $M_2 \in F_1$ such that $M_2 \cap E_1 \neq \phi$. Let $w \in E_1 \cap M_2$, where $v \in V(C_q)$, $u \in V(S_{k_1})$, and $k_1 \in \{1, \dots, p\}$. Since C_q has a perfect matching, which shows that $m(C_q - v) = m(C_q) - 1 < m(C_q)$. In \widehat{S}_{k_1} , u is a quasi-pendant vertex, by Lemma 2.2, $m(S_{k_1} - u) = m(\widehat{S}_{k_1} - u - v) = m(\widehat{S}_{k_1}) - 1 = m(S_{k_1}) - 1$. Then, $|M_2| = m(C_q - v) + (1 + m(\widehat{S}_{k_1} - u - v)) + \sum_{k \neq k_1} |M_2 \cap \widehat{S}_k| < m(C_q) + \sum_{k=1}^p m(\widehat{S}_k) = m(C_q) + \sum_{k=1}^p m(S_k) = m(G)$. This contradicts $m(G) = |M_2|$, thus $M \cap E_1 = \phi$ for all $M \in F_1$.

Sufficiency: Note that $m(G) = m(C_q) + \sum_{k=1}^p m(S_k)$. We assume that $m(T_G) \neq m([T_G])$, that is, $m(T_G) = m([T_G]) + 1$ (G is an unicyclic graph). Which implies that each maximum matching of T_G must cover the vertex v_c of W_G . There exists a maximum matching M_T'' of T_G such that the edge $wv_c \in M_T''$, and the vertex w is adjacent to v_0 the vertex of C_q^σ in G . Since q is even, then $m(C_q - v_0) = m(C_q) - 1$. Assume that M_3 is a matching of G , which is the union of M_T'' and a maximum matching of $C_q - v_0$. Then $|M_3| = m(T_G) + m(C_q - v_0) = m([T_G]) + 1 + m(C_q) - 1 = m(G)$, which implies that M_3 is a maximum matching of G and $M_3 \cap E_1 \neq \phi$, a contradiction. Thus $m(T_G) = m([T_G])$. \square

Next, we give Lemmas 3.9 and 3.10 by applying the characteristic polynomial of the signed graph.

Lemma 3.9. *Let (G, σ) be an unbalanced signed unicyclic graph, which contains the unique cycle C_q^σ . If q is odd and $m(T_G) \neq m([T_G])$, then $\eta(G, \sigma) = |V(G)| - 2m(G)$.*

Proof. Let $|V(G)| = n$, $m(G) = m$ and $m(G - C_q) = l$. By Lemma 3.3, we have the characteristic polynomial of (G, σ) , $P_{(G, \sigma)}(\lambda) = \sum_{i=0}^n a_i \cdot \lambda^{n-i}$, $a_i = \sum_{(u, \sigma) \in \mathbf{U}_i} (-1)^{p(u)+s(u)} \cdot 2^{c(u)}$, for any $i \in \{1, 2, \dots, n\}$. Since q is odd, then $m(C_q) = \frac{q-1}{2}$, and C_q^σ has not a perfect matching. Note that $m(T_G) \neq m([T_G])$, by Corollary 3.7, we have $m(G) > m([T_G]) + m(C_q) = l + \frac{q-1}{2}$, that is, $2m + 1 > q + 2l$. If $i > 2m$, then (G, σ) contains no basic subgraphs with i vertices and $a_i = 0$. Thus $P_{(G, \sigma)}(\lambda) = \lambda^n + a_1 \lambda^{n-1} + \dots + a_{2m} \lambda^{n-2m} = \lambda^{n-2m} (\lambda^{2m} + a_1 \lambda^{2m-1} + \dots + a_{2m})$. Therefore $\eta(G, \sigma) \geq |V(G)| - 2m(G)$.

Next, we need to prove $a_{2m} \neq 0$. Note that $a_{2m} = \sum_{(u, \sigma) \in \mathbf{U}_{2m}} (-1)^{p(u)+s(u)} \cdot 2^{c(u)}$, since C_q^σ is an odd cycle, the basic subgraphs of (G, σ) with $2m$ vertices contain no cycle. Then $s(u) = c(u) = 0$, $p(u) = m$, we have $a_{2m} = \sum_{(u, \sigma) \in \mathbf{U}_{2m}} (-1)^m \neq 0$. Thus $\eta(G, \sigma) = |V(G)| - 2m(G)$. \square

Lemma 3.10. *Let (G, σ) be an unbalanced signed unicyclic graph with the unique cycle C_q^σ , and q is even. If $m(T_G) = m([T_G])$ and $q \equiv 0 \pmod{4}$; or $m(T_G) \neq m([T_G])$, then $\eta(G, \sigma) = |V(G)| - 2m(G)$.*

Proof. Let $F_3 = \{u | u = C_q \cup M, M \in F_2\}$, $m(G) = m$ and $m(G - C_q) = l$. C_q^σ has odd number of negative edges. Note that G is a bipartite graph. By Lemma 3.3, then

the characteristic polynomial of (G, σ) can be expressed by $P_{(G, \sigma)}(\lambda) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} b_i \cdot \lambda^{n-2i}$, $b_i = \sum_{(u, \sigma) \in \mathbf{U}_{2i}} (-1)^{p(u)+s(u)} \cdot 2^{c(u)}$, for any $i \in \{1, 2, \dots, \lfloor \frac{n}{2} \rfloor\}$. By the similar proof as in Lemma 3.9, if $i > m$, then (G, σ) contains no basic subgraphs with $2i$ vertices and $b_i = 0$. Hence

$$P_{(G, \sigma)}(\lambda) = \lambda^n + b_1 \lambda^{n-2} + \dots + b_m \lambda^{n-2m} = \lambda^{n-2m} (\lambda^{2m} + b_1 \lambda^{2m-2} + \dots + b_m).$$

Therefore $\eta(G, \sigma) \geq |V(G)| - 2m(G)$.

Next, in order to get the result $\eta(G, \sigma) = |V(G)| - 2m(G)$, we need to prove $b_m \neq 0$.

(1). $m(T_G) = m([T_G])$, and $q \equiv 0 \pmod{4}$.

By Lemma 3.8, we have $m(G) = m([T_G]) + m(C_q)$. Note that $m = \frac{q}{2} + l$. Since $2m = q + 2l$ and the definition of \mathbf{U}_{2m} , by the similar proof as in Lemma 3.9, we have $\mathbf{U}_{2m} = F_1^\sigma \cup F_3^\sigma$. If $(u, \sigma) \in F_1^\sigma$, then $p(u) = m$, $s(u) = c(u) = 0$. If $(u, \sigma) \in F_3^\sigma$, then $p(u) = l + 1$, $c(u) = 1$. Let $s(u) = 2a + 1$, $a \in \mathbf{Z}^+$ (since C_q^σ has an odd number of negative edges), then

$$b_m = \sum_{(u, \sigma) \in F_1^\sigma} (-1)^m + \sum_{(u, \sigma) \in F_3^\sigma} (-1)^{l+1+2a+1} \cdot 2^1 = (-1)^m \cdot |F_1^\sigma| + (-1)^l \cdot 2 \cdot |F_3^\sigma| = (-1)^l ((-1)^{m-l} |F_1^\sigma| + 2 \cdot |F_3^\sigma|). \text{ Since } m - l = \frac{q}{2} \text{ and } q \equiv 0 \pmod{4}, \text{ we have } m - l \equiv 0 \pmod{2}. \text{ Thus } b_m = (-1)^l (|F_1^\sigma| + 2 \cdot |F_3^\sigma|) \neq 0.$$

(2). $m(T_G) \neq m([T_G])$.

Equivalently, $m(T_G) = m([T_G]) + 1$, then each maximum matching of T_G must cover the unique cyclic vertex. By Lemma 3.8, we mainly discuss the following two cases.

Case 1. $m(G) \neq m(C_q^\sigma) + m([T_G])$, that is, $m(G) > m(C_q^\sigma) + m([T_G])$, $m > \frac{q}{2} + l$, $2m > q + 2l$. Then the basic subgraphs with $2m$ vertices have no cycle. Which shows that $F_3^\sigma = \phi$ and $\mathbf{U}_{2m} = F_1^\sigma$. Then $p(u) = m$ and $s(u) = c(u) = 0$ for any $(u, \sigma) \in \mathbf{U}_{2m}$. Thus $b_m = \sum_{(u, \sigma) \in F_1^\sigma} (-1)^m = (-1)^m |F_1^\sigma| \neq 0$.

Case 2. $m(G) = m(C_q^\sigma) + m([T_G])$ and $M \cap E_1 \neq \phi$, $\exists M \in F_1$ (equivalently, $F_1' \neq \phi$). Note that $m = l + \frac{q}{2}$, there exists some basic subgraphs with $2m$ vertices, such that contain C_q^σ as a subgraph. Then $\mathbf{U}_{2m} = F_1^\sigma \cup F_3^\sigma$. If $(u, \sigma) \in F_1^\sigma$, then $p(u) = m$, $s(u) = c(u) = 0$. If $(u, \sigma) \in F_3^\sigma$, we have $p(u) = l + 1$, $c(u) = 1$. Let $s(u) = 2a + 1$, $a \in \mathbf{Z}^+$ (since C_q^σ has an odd number of negative edges). Then

$$b_m = \sum_{(u, \sigma) \in F_1^\sigma} (-1)^m + \sum_{(u, \sigma) \in F_3^\sigma} (-1)^{l+1+2a+1} \cdot 2^1 = (-1)^m \cdot |F_1^\sigma| + (-1)^l \cdot 2 \cdot |F_3^\sigma| = (-1)^l ((-1)^{m-l} |F_1^\sigma| + 2 \cdot |F_3^\sigma|). \text{ Since } q \text{ is even, we mainly discuss the following two subcases.}$$

Subcase 2.1. If $q \equiv 0 \pmod{4}$, then $m - l = \frac{q}{2} \equiv 0 \pmod{2}$. Thus

$$b_m = (-1)^l \cdot (|F_1^\sigma| + 2 \cdot |F_3^\sigma|) \neq 0$$

Subcase 2.2. If $q \equiv 2 \pmod{4}$, then $m - l = \frac{q}{2} \equiv 1 \pmod{2}$. Thus

$b_m = (-1)^{l+1} (|F_1^\sigma| - 2 \cdot |F_3^\sigma|) = (-1)^{l+1} (|F_1^\sigma| - 2 \cdot |F_2^\sigma|)$, $|F_3| = |F_2|$. Note that $F_1' \neq \phi$, by Corollary 3.2, we have $|F_1| > 2|F_2|$. Hence $b_m \neq 0$,

Based on the above conclusions, we have $b_m \neq 0$. Thus $\eta(G, \sigma) = |V(G)| - 2m(G)$. \square

By Lemmas 3.4, 3.5, 3.9 and 3.10, we obtained Theorem 1.1.

4 Proof of Theorem 1.2

In this section, we mainly investigate the unbalanced signed graphs. In order to proof Theorem 1.2, we need to introduce some definitions and lemmas.

Let G be the graph with some pendant vertices, and G contains at least a cycle. For any pendant vertex u , we will give definitions of two types of pendant vertex u :

Definition 4.1. (1) If the quasi-pendant vertex of u does not lie on the cycle, then u is of Type I.

(2) If the quasi-pendant vertex of u lies on the cycle, then u is of Type II.

Lemma 4.2. Let (G, σ) be an unbalanced signed graph with a pendant vertex u , and v is adjacent to u . If u is of Type I, then $\eta(G, \sigma) = |V(G)| - 2m(G) + 2c(G) - s$ if and only if $\eta(G - u - v, \sigma) = |V(G - u - v)| - 2m(G - u - v) + 2c(G - u - v) - s$. ($0 \leq s \leq 3c(G)$).

Proof. By Lemmas 2.2 and 2.6,

$$\begin{aligned} |V(G)| &= |V(G - u - v)| + 2, \\ m(G) &= m(G - u - v) + 1, \\ c(G) &= c(G - u - v). \end{aligned}$$

By Lemma 2.8,

$$\eta(G - u - v, \sigma) = \eta(G, \sigma) = |V(G)| - 2m(G) + 2c(G) - s = |V(G - u - v)| + 2 - 2(m(G - u - v) + 1) + 2c(G - u - v) - s = |V(G - u - v)| - 2m(G - u - v) + 2c(G - u - v) - s. \quad \square$$

Lemma 4.3. Let (G, σ) be an unbalanced signed graph with a pendant vertex u , and v is adjacent to u . If u is of Type II, then $\eta(G, \sigma) \leq |V(G)| - 2m(G) + 2c(G) - 2$.

Proof. We prove the result by contradiction. From Lemma 2.9, suppose on the contrary, there exists some unbalanced signed graphs (H, σ) with the nullity $\eta(H, \sigma) = |V(H)| - 2m(H) + 2c(H) - s$, $s = 0, 1$. Since v lies on the cycle of (H, σ) , by Lemmas 2.2 and 2.6, we have

$$\begin{aligned} |V(H)| &= |V(H - u - v)| + 2, \\ m(H) &= m(H - u - v) + 1, \\ c(H) &\geq c(H - u - v) + 1. \end{aligned}$$

By Lemma 2.8, we have $\eta(H - u - v, \sigma) = \eta(H, \sigma) = |V(H)| - 2m(H) + 2c(H) - s \geq |V(H - u - v)| + 2 - 2(m(H - u - v) + 1) + 2c(H - u - v) + 2 - s \geq |V(H - u - v)| - 2m(H - u - v) + 2c(H - u - v) + 1$. Which contradicts Lemma 2.9. Therefore, for any unbalanced signed graph (G, σ) with a pendant vertex u of Type II, then $\eta(G, \sigma) \leq |V(G)| - 2m(G) + 2c(G) - 2$. \square

Lemma 4.4. Let (G, σ) be an unbalanced signed graph without pendant vertices. If $\eta(G, \sigma) \neq |V(G)| - 2m(G) + 2c(G)$, $c(G) \geq 2$, then there exists a vertex x on the cycle in (G, σ) such that $\eta(G - x, \sigma) \neq |V(G - x)| - 2m(G - x) + 2c(G - x)$.

Proof. (1). $g(G) = 3$, $c(G) \geq 2$.

Denote by C_q^σ the cycle of (G, σ) and $q = 3$. Since $c(G) \geq 2$, there exists a vertex x on another cycle in (G, σ) such that C_q^σ is a subgraph of $(G - x, \sigma)$. Which shows that $(G - x, \sigma)$ does not satisfy in Lemma 2.14(2), then

$$\eta(G - x, \sigma) \neq |V(G - x)| - 2m(G - x) + 2c(G - x).$$

(2). $g(G) \geq 4$, $c(G) \geq 2$.

Since $\eta(G, \sigma) \neq |V(G)| - 2m(G) + 2c(G)$, (G, σ) does not satisfy at least one of the three conditions in Lemma 2.14.

Case 1. If (G, σ) does not satisfy Lemma 2.14(1).

Note that (G, σ) contains at least two vertex-joint cycles, called C_k^σ and C_s^σ ($k, s \geq 4$). Let $G[C_k^\sigma, C_s^\sigma]$ be the graph induced by $V(C_k^\sigma)$ and $V(C_s^\sigma)$.

Subcase 1.1. $c(G) = 2$.

Since (G, σ) contains no pendant vertices, then (G, σ) is the union of an ∞ -graph (or a θ -graph) and some isolated vertices (if any), which implies that $G[C_k^\sigma, C_s^\sigma]$ is either $\infty^\sigma(p, 1, q)$ or $\theta^\sigma(p, l, q)$.

Subcase 1.1.1. $G[C_k^\sigma, C_s^\sigma] = \theta^\sigma(1, 1, 1)$.

Since (G, σ) is unbalanced, (G, σ) contains at least an unbalanced cycle, says C_q^σ , and $q = 4$. Let x be a vertex of another cycle in (G, σ) , $x \notin V(C_q^\sigma)$. Then C_q^σ is a subgraph of $(G - x, \sigma)$. Which shows that $(G - x, \sigma)$ does not satisfy Lemma 2.14(2), then

$$\eta(G - x, \sigma) \neq |V(G - x)| - 2m(G - x) + 2c(G - x).$$

Subcase 1.1.2. $G[C_k^\sigma, C_s^\sigma] \neq \theta^\sigma(1, 1, 1)$.

As shown in Figure 1, there exists a vertex x on the cycle in (G, σ) , such that $(G - x, \sigma)$ contains a pendant vertex with Type II. By Lemma 4.3, we can get that

$$\eta(G - x, \sigma) \neq |V(G - x)| - 2m(G - x) + 2c(G - x).$$

Subcase 1.2. $c(G) \geq 3$.

Subcase 1.2.1. In (G, σ) , there exists a vertex x on the cycle such that $x \notin G[C_k^\sigma, C_s^\sigma]$.

Which implies that $G[C_k^\sigma, C_s^\sigma]$ is a subgraph of $(G - x, \sigma)$. Then $(G - x, \sigma)$ does not satisfy Lemma 2.14(1), hence

$$\eta(G - x, \sigma) \neq |V(G - x)| - 2m(G - x) + 2c(G - x).$$

For example, as shown in Figure 2, the signed graphs with $T_i (i = 1, \dots, 4)$ as a underlying graph, which contains a vertex x on the cycle and $x \notin G[C_k^\sigma, C_s^\sigma]$.

Subcase 1.2.2. In (G, σ) , let x be a vertex on any cycle, then $x \in G[C_k^\sigma, C_s^\sigma]$.

Which implies that any cycle of (G, σ) is the subgraph of $G[C_k^\sigma, C_s^\sigma]$. Since $c(G) \geq 3$, (G, σ) contains one of eight types of bases of tricyclic graphs as a underlying subgraph. As shown in Figure 2, the graph T_j can be regard as two vertex-joint cycles, where $j = 5, \dots, 8$. Which implies that the tricyclic graphs T_j is a underlying subgraph of $G[C_k^\sigma, C_s^\sigma]$. As shown in Figure 2, there exists a vertex x of T_j such that $T_j - x$ also contains two vertex-joint cycles. Hence there exists a vertex x on the cycle of (G, σ) such that $(G - x, \sigma)$ does not satisfy Lemma 2.14(1), then

$$\eta(G - x, \sigma) \neq |V(G - x)| - 2m(G - x) + 2c(G - x).$$

Case 2. (G, σ) satisfies Lemma 2.14(1) but does not satisfy Lemma 2.14(2).

Subcase 2.1. There exists at least a balanced cycle in (G, σ) .

Subcase 2.1.1. There exists a balanced cycle in (G, σ) , says C_p^σ and $p \neq 0 \pmod{4}$. Let x be a vertex on another cycle and $x \notin V(C_p^\sigma)$ (since all cycles are pairwise vertex-disjoint), then C_p^σ is the subgraph of $(G - x, \sigma)$. Which shows that $(G - x, \sigma)$ does not satisfy Lemma 2.14(2), then

$$\eta(G - x, \sigma) \neq |V(G - x)| - 2m(G - x) + 2c(G - x).$$

Subcase 2.1.2. The length of all balanced graph is the multiple of 4. Then (G, σ) contains at least an unbalanced cycle in (G, σ) , says C_q^σ and $q \neq 2 \pmod{4}$. Let x be a vertex

of the balanced cycle and $x \notin V(C_q^\sigma)$ (Since all cycles are pairwise vertex-disjoint), then C_q^σ is the subgraph of $(G - x, \sigma)$. Which shows that $(G - x, \sigma)$ does not satisfy Lemma 2.14(2), then

$$\eta(G - x, \sigma) \neq |V(G - x)| - 2m(G - x) + 2c(G - x).$$

Case 2.2. Cycles of (G, σ) are unbalanced.

(G, σ) contains at least an unbalanced cycle, says C_q^σ and $q \neq 2(\text{mod } 4)$. Let x be a vertex of another unbalanced cycle in (G, σ) and $x \notin V(C_q^\sigma)$ (all cycles are pairwise vertex-disjoint). Since $c(G) \geq 2$, then C_q^σ is the subgraph of $(G - x, \sigma)$. Which shows that $(G - x, \sigma)$ does not satisfy Lemma 2.14(2), then

$$\eta(G - x, \sigma) \neq |V(G - x)| - 2m(G - x) + 2c(G - x).$$

Case 3. (G, σ) satisfies (1) and (2) of Lemma 2.14, but dose not satisfy (3) of Lemma 2.14.

Note that $m(T_G) \neq m([T_G])$, that is, $m(T_G) \geq m([T_G]) + 1$.

If $E(T_G) = \phi$, then (G, σ) is the union of some vertex-disjoint signed cycles and isolated vertices. Hence $m(T_G) = m([T_G]) = 0$, a contradiction. Therefore, we only consider $E(T_G) \neq \phi$. In T_G , each maximum matching must cover at least a pendant vertex. Otherwise, there exists an M -augmenting path in G , which contradicts Lemma 2.1. Let u be a pendant vertex of T_G . Since (G, σ) contains no pendant vertices, we have $u \in W_G$. Suppose that C_q^σ the pendant cycle of (G, σ) is correspond to the vertex u of T_G . Let v be the unique vertex with degree three in C_q^σ , and let x be a vertex on the cycle C_q^σ . Then T_{G-x} is the graph obtained from T_G and $C_q^\sigma - x$ by identifying u and v as a vertex, we mainly discuss the following two subcases.

Subcase 3.1. Each maximum matching of T_G cover all pendant vertices.

Let x be a vertex on the cycle C_q^σ , and x is adjacent to v . Noted that C_q^σ is an even cycle, then $C_q^\sigma - v - x$ is an odd path and has a perfect matching. By the definition of T_G , which shows that the maximum matching of T_{G-x} is the union of the maximum matching of T_G and the maximum matching of $C_q^\sigma - v - x$. Then

$$m(T_{G-x}) = m(T_G) + m(C_q^\sigma - v - x).$$

Hence each maximum matching of T_{G-x} must cover some vertices in W_{G-x} , we have $m(T_{G-x}) \neq m([T_{G-x}])$. Which shows that $(G - x, \sigma)$ does not satisfy Lemma 2.14(3), then

$$\eta(G - x, \sigma) \neq |V(G - x)| - 2m(G - x) + 2c(G - x).$$

Subcase 3.2. There exists some maximum matchings of T_G , denoted by $M_i(T_G)$ ($i = 1, \dots, r$), such that the pendant edge $wv \notin M_i(T_G)$, where $r \geq 1$ and v is a pendant vertex of T_G .

Let y be a vertex on the cycle C_p^σ and $d(v_0, y) = 2$, where C_p^σ the cycle of (G, σ) is correspond to v the vertex of T_G , and v_0 is the unique vertex with degree three in C_p^σ . By the definition of T_{G-y} , which shows that the maximum matching of T_{G-y} is the union of $M_i(T_G)$ ($i \in \{1, \dots, r\}$) and the maximum matching of $C_p^\sigma - y$. Then

$$m(T_{G-y}) = m(T_G) + m(C_p^\sigma - y).$$

For any $i \in \{1, \dots, r\}$, then $M_i(T_G)$ must cover some vertices in W_{G-y} by Lemma 2.1. Which implies that each maximum matching of T_{G-y} must cover some vertices in W_{G-y} , we have $m(T_{G-y}) \neq m([T_{G-y}])$. Which shows that $(G - y, \sigma)$ does not satisfy Lemma 2.14(3), then $\eta(G - y, \sigma) \neq |V(G - y)| - 2m(G - y) + 2c(G - y)$. \square

Lemma 4.5. *Let (G, σ) be an unbalanced signed graph without pendant vertices. Then*

$$\eta(G, \sigma) \neq |V(G)| - 2m(G) + 2c(G) - 1.$$

Proof. We apply induction on $c(G)$ to prove the result.

If $c(G) = 1$, then $G = C_k \cup (n - k)K_1$ ($3 \leq k \leq n$). By Theorem 1.1, we have

$$\eta(G, \sigma) \neq |V(G)| - 2m(G) + 2c(G) - 1.$$

If $c(G) \geq 2$, assume that the conclusion is true for $c(G) \leq k$. Next, we need to prove the conclusion is true for $c(G) = k + 1$. Suppose on the contrary, there exists some unbalanced signed graphs (H, σ) with $c(H) = k + 1$ such that $\eta(H, \sigma) = |V(H)| - 2m(H) + 2c(H) - 1$.

Let x be any vertex on the cycle of (H, σ) . For the signed graph $(H - x, \sigma)$, by Lemmas 2.3 and 2.6, we have

$$\begin{aligned} |V(H)| &= |V(H - x)| + 1, \\ m(H) &\leq m(H - x) + 1, \\ c(H) &\geq c(H - x) + 1. \end{aligned}$$

By Lemma 2.7, we have $\eta(H - x, \sigma) + 1 \geq \eta(H, \sigma) = |V(H)| - 2m(H) + 2c(H) - 1 \geq |V(H - x)| + 1 - 2(m(H - x) + 1) + 2(c(H - x) + 1) - 1 = |V(H - x)| - 2m(H - x) + 2c(H - x)$. By Lemma 2.9, then

$$\eta(H - x, \sigma) = |V(H - x)| - 2m(H - x) + 2c(H - x) - s, \quad s = 0, 1$$

If $(H - x, \sigma)$ is balanced, by Lemmas 2.5 and 2.12, we have $\eta(H - x, \sigma) \neq |V(H - x)| - 2m(H - x) + 2c(H - x) - 1$. We mainly consider that $(H - x, \sigma)$ is unbalanced in the following cases.

Case 1. $(H - x, \sigma)$ contains no pendant vertices.

In this case, $(H - x, \sigma)$ is also unbalanced and $c(H - x) \leq c(H) - 1 = k$. Applying the induction hypothesis to $(H - x, \sigma)$, then $\eta(H - x, \sigma) \neq |V(H - x)| - 2m(H - x) + 2c(H - x) - 1$.

Case 2. $(H - x, \sigma)$ contains some pendant vertices.

Subcase 2.1. $(H - x, \sigma)$ contains at least a pendant vertex of Type II.

By Lemma 4.3, we have $\eta(H - x, \sigma) \leq |V(H - x)| - 2m(H - x) + 2c(H - x) - 2$.

Subcase 2.2. All the pendant vertices of $(H - x, \sigma)$ are of Type I.

Suppose that $(H - x, \sigma)$ contains p pendant vertices. For the pendant vertices of Type I, by using Lemma 2.8 repeatedly, after p steps, we obtain a subgraph (H_1, σ) of (H, σ) . If (H_1, σ) contains no pendant vertices or at least a pendant vertex of Type II, then (H_1, σ) is as we required and we are done. Otherwise, in (H_1, σ) , for the pendant vertex of Type I, we continue to use Lemma 2.8 repeatedly, we obtain a subgraph (H_2, σ) of (H_1, σ) . If (H_2, σ) contains no pendant vertices or at least a pendant vertex of Type II, then (H_2, σ) is as we required and we are done. Otherwise, repeating the above steps until we obtain an unbalanced sign graph (H_0, σ) that meets the requirements.

(a). (H_0, σ) contains at least a pendant vertex of Type II.

By Lemma 4.3, we have

$$\eta(H_0, \sigma) \leq |V(H_0)| - 2m(H_0) + 2c(H_0) - 2.$$

Next, by Lemma 4.2,

$$\eta(H_0, \sigma) \leq |V(H_0)| - 2m(H_0) + 2c(H_0) - 2$$

if and only if

$$\eta(H - x, \sigma) \leq |V(H - x)| - 2m(H - x) + 2c(H - x) - 2.$$

(b). (H_0, σ) contains no pendant vertices.

Since $c(H_0) = c(H - x) \leq c(H) - 1 = k$ and H_0^σ is unbalanced, applying the induction hypothesis to H_0^σ , then $\eta(H_0, \sigma) \neq |V(H_0)| - 2m(H_0) + 2c(H_0) - 1$.

By Lemma 4.2,

$$\eta(H_0, \sigma) \neq |V(H_0)| - 2m(H_0) + 2c(H_0) - 1$$

if and only if

$$\eta(H - x, \sigma) \neq |V(H - x)| - 2m(H - x) + 2c(H - x) - 1.$$

Based on the above results. Let x be any vertex on the cycle of (H, σ) . For the nullity $\eta(H - x, \sigma)$ of signed graph $(H - x, \sigma)$, either $\eta(H - x, \sigma) \neq |V(H - x)| - 2m(H - x) + 2c(H - x) - 1$ or $\eta(H - x, \sigma) \leq |V(H - x)| - 2m(H - x) + 2c(H - x) - 2$.

If $\eta(H - x, \sigma) \neq |V(H - x)| - 2m(H - x) + 2c(H - x) - 1$, then $\eta(H - x, \sigma) = |V(H - x)| - 2m(H - x) + 2c(H - x)$. On the other hand, since $\eta(H, \sigma) \neq |V(H)| - 2m(H) + 2c(H)$, by Lemma 4.4, there exists a vertex y on the cycle of (H, σ) such that $\eta(H - x, \sigma) \neq |V(H - x)| - 2m(H - x) + 2c(H - x)$, a contradiction.

If $\eta(H - x, \sigma) \leq |V(H - x)| - 2m(H - x) + 2c(H - x) - 2$, which contradicts equation $\eta(H - x, \sigma) = |V(H - x)| - 2m(H - x) + 2c(H - x) - s$, $s = 0, 1$.

Therefore, for any unbalanced signed graph (G, σ) without pendant vertices, $\eta(G, \sigma) \neq |V(G)| - 2m(G) + 2c(G) - 1$. \square

The Proof of Theorem 1.2: Noted that the unbalanced signed graph (G, σ) contains at least an unbalanced cycle, and $c(G) \geq 1$.

Case 1. (G, σ) has no pendant vertices, we can obtained the result by Lemma 4.5.

Case 2. (G, σ) has some pendant vertices.

By the definitions of two types of pendant vertices, we mainly discuss the following two subcases.

Subcase 2.1. All pendant vertices are of Type I.

Suppose that (G, σ) contains p pendant vertices. For the pendant vertex of Type I, by using Lemma 2.8 repeatedly, after p steps, we obtain a subgraph (G_1, σ) of (G, σ) . If (G_1, σ) contains no pendant vertices or at least a pendant vertex of Type II, then (G_1, σ) is as we required and we are done. Otherwise, in (G_1, σ) , for the pendant vertex of Type I, we continue to use Lemma 2.8 repeatedly, we obtain a subgraph (G_2, σ) of (G_1, σ) . If (G_2, σ) contains no pendant vertices or at least a pendant vertex of Type II, then (G_2, σ) is as we required and we are done. Otherwise, repeating the above steps until we obtain an unbalanced sign graph (G_0, σ) that meets the requirements.

(a). (G_0, σ) contains no pendant vertices.

By Lemma 4.5, we have $\eta(G_0, \sigma) \neq |V(G_0)| - 2m(G_0) + 2c(G_0) - 1$.

(b). (G_0, σ) contains at least a pendant vertex of Type II.

By Lemma 4.3, we have

$$\eta(G_0, \sigma) \leq |V(G_0)| - 2m(G_0) + 2c(G_0) - 2.$$

Hence

$$\eta(G_0, \sigma) \neq |V(G_0)| - 2m(G_0) + 2c(G_0) - 1.$$

Next, by Lemma 4.2, we have

$\eta(G_0, \sigma) \neq |V(G_0)| - 2m(G_0) + 2c(G_0) - 1$ if and only if $\eta(G, \sigma) \neq |V(G)| - 2m(G) + 2c(G) - 1$.

Subcase 2.2. There exists at least a pendant vertex of Type II.

By Lemma 4.3, we have $\eta(G, \sigma) \neq |V(G)| - 2m(G) + 2c(G) - 1$. \square

Combining with Lemmas 2.5, 2.12 and Theorem 1.2, we can obtained the following corollary.

Corollary 4.6. *For any signed graph (G, σ) , then*

$$\eta(G, \sigma) \neq |V(G)| - 2m(G) + 2c(G) - 1.$$

5 Proof of Theorem 1.3

In order to proof Theorem 1.3, we need to introduce some definitions.

Let $K_{1,c+1}^\sigma$ be a signed star, x be the center vertex of $K_{1,c+1}^\sigma$ and y_0, \dots, y_c be pendant vertices of $K_{1,c+1}^\sigma$, respectively. $O_1^\sigma, \dots, O_{l_1}^\sigma$ be a balanced cycle of size 3, $O_{l_1+1}^\sigma, \dots, O_{l_1'}^\sigma$ be an unbalanced cycle of size 6. Let $O_{l_1'+1}^\sigma, \dots, O_c^\sigma$ be the signed graph with the pendant vertex of Type II, which obtained from C_4^+ a balanced cycle of size 4 by attaching K_2^σ . Next, we construct a new signed graph G^σ by identifying y_i with a vertex of O_i^σ , y_j with a pendant vertex of O_j^σ , where $i = 1, \dots, l_1'$, $j = l_1' + 1, \dots, c$.

The Proof of Theorem 1.3: According to the definition of G^σ , let $l_2 = l_1' - l_1$, $l_3 = c - l_1'$. In this case, there exists l_1 balanced cycles C_3^+ , l_2 unbalanced cycles C_6^- and l_3 balanced graph F^σ , which is obtained from C_4^+ a balanced cycles of size 4 by attaching K_2^σ . Which implies that y_0 is the unique pendant vertex of G^σ . Then

$$|V(G^\sigma)| = 3l_1 + 6l_2 + 5l_3 + 2,$$

$$m(G^\sigma) = l_1 + 3l_2 + 2l_3 + 1,$$

$$c(G^\sigma) = l_1 + l_2 + l_3.$$

By Lemmas 2.8 and 2.13, we obtain that $\eta(G^\sigma) = \eta(G^\sigma - x - y_0) = 2l_2 + l_3 = |V(G^\sigma)| - 2m(G^\sigma) + 2c(G^\sigma) - (3l_1 + 2l_3)$. Since $l_i \geq 0$, $3l_1 + 2l_3$ can take over any integer between 0 and $3c(G)$ expect for 1. \square

Combining with Lemma 2.12 and Theorem 1.3, we can obtained the following corollary.

Theorem 5.1. *For any signed graph (G, σ) , if $c(G)$ is fixed, then there exists infinitely connected signed graphs, such that $\eta(G, \sigma) = |V(G)| - 2m(G) + 2c(G) - s$, where $0 \leq s \leq 3c(G)$, $s \neq 1$.*

References

- [1] J.H. Bevis, K.K. Blount, G.J. Davis, The rank of a graph after vertex addition, Linear Algebra Appl. 265 (1997) 55–69.
- [2] J.A. Bondy, U.S.R. Murty, Graph Theory with Applications, Elsevier, New York, 1976.

- [3] Chen Chen, Jing Huang, Shuchao Li, On the relation between the H -rank of a mixed graph and the matching number of its underlying graph, *Linear Multilinear Algebra*. 66 (2018) 1853–1869.
- [4] L. Collatz, U. Sinogowitz, Spektren endlicher grafen, *Abh. Math. Sem. Univ. Hamburg*. 21 (1957) 63–77.
- [5] Yizheng Fan, Yue Wang, Yi Wang, A note on the nullity of unicyclic signed graphs, *Linear Algebra Appl.* 438 (2013) 1193–1200.
- [6] Shengjie He, Rong-Xia Hao, Hong-Jian Lai, Bounds for the matching number and cyclomatic number of a signed graph in terms of rank, *Linear Algebra Appl.* 572 (2019), 273–291.
- [7] Yaoping Hou, Jiongsheng Li, On the Laplacian eigenvalues of signed graphs, *Linear and Multilinear Algebra*. 51 (2003) 21–30.
- [8] Xin Li, Jiming Guo, No graph with nullity $\eta(G) = |V(G)| - 2m(G) + 2c(G) - 1$, *Discrete Math.* 268 (2019) 130–136.
- [9] Yu Liu, Lihua You, Further results on the nullity of signed graphs, *J. Appl. Math.* 1 (2014) 8pp.
- [10] Yong Lu, Ligong Wang, Qiannan Zhou, The rank of a signed graph in terms of the rank of its underlying graph, *Linear Algebra Appl.* 538 (2018), 166–186.
- [11] Xiaobin Ma, Dein Wong, Fenglei Tian, Skew-rank of an oriented graph in terms of matching number, *Linear Algebra Appl.* 495 (2016) 242–255.
- [12] S. Rula, An Chang, Yirong Zheng, The extremal graphs with respect to their nullity, *J. Inequal. Appl.* 2016 (2016) 71, 13pp.
- [13] Yazhi Song, Xiaoqiu Song, Bit-Shun Tam, A characterization of graphs G with nullity $V(G) - 2m(G) + 2c(G)$, *Linear Algebra Appl.* 465 (2015) 363–365
- [14] Long Wang, Characterization of graphs with given order, given size and given matching number that minimize nullity, *Discrete Math.* 339 (2016) 1574–1582.
- [15] Long Wang, Dein Wong, Bounds for the matching number, the edge chromatic number and the independence number of a graph in terms of rank, *Discrete Appl. Math.* 166 (2014) 276–281.
- [16] Dein Wong, Xiaobin Ma, Fenglei Tian, Relation between the skew-rank of an oriented graph and the rank of its underlying graph, *European J. Combin.* 54 (2016) 76–86.