

Random matrix model for the QCD Kondo effect

Takuya Kanazawa

Research and Development Group, Hitachi, Ltd., Kokubunji, Tokyo 185-8601, Japan

We propose a novel random matrix model that describes the QCD Kondo effect. The model correctly implements both the chiral symmetry of light quarks and the $SU(2)$ spin symmetry of heavy quarks. We take the large- N limit with N the matrix size and show that the model has three phases: the pure Kondo phase with no chiral condensate, the pure chirally broken phase with no Kondo condensate, and the coexistence phase. The model predicts that the pairing form of the Kondo condensate in the coexistence phase is significantly altered compared to the pure Kondo phase. For each of these phases we rigorously derive the low-energy effective theory of Nambu-Goldstone modes and obtain compact closed expressions for the partition function with external sources. We also include a chiral chemical potential into the model and examine the vacuum structure.

I. INTRODUCTION

In certain alloys with dilute magnetic impurities, the electrical resistivity shows a minimum at nonzero temperature, a phenomenon famously known as the Kondo effect [1–4]. The antiferromagnetic coupling between conduction electrons and an impurity gets stronger as the temperature lowers, and eventually leads to a complete screening of the magnetic moment of the impurity. This is a classical example of asymptotic freedom. The prerequisites of the Kondo effect are (i) nonzero density of states of conduction electrons, (ii) a non-Abelian interaction, and (iii) quantum loop effects. Later the Kondo problem was investigated for materials which have the density of states that vanishes at the Fermi energy as a power law, $\rho(E) \propto |E - E_F|^r$. This is called the pseudogap Kondo problem and is relevant to materials hosting quasiparticles with relativistic dispersions, including graphene, Dirac/Weyl semimetals, and d -wave superconductors [5–16]. In these systems, there is a quantum phase transition at finite coupling from the unscreened free moment phase to the strongly coupled Kondo phase. This is analogous to chiral symmetry breaking in the QCD vacuum [17, 18].

Recently the Kondo effect in relativistic nuclear and quark matter has been discussed [19–29]. In such systems the impurity is either a single heavy quark (charm and bottom) or a hadron including a heavy quark (e.g., D and B mesons). The non-Abelian symmetry pertinent to the QCD Kondo effect is the $SU(2)$ isospin symmetry and the $SU(3)$ color symmetry. (Precisely speaking, as first pointed out in [24], the Kondo effect in QCD is a relativistic analog of the *overscreened Kondo effect* [30–33], in which the low-energy physics is governed by a non-Fermi-liquid fixed point. The overscreening is typical for multichannel systems where there are more conduction electrons than are necessary to exactly screen the impurity’s spin.) The Kondo scale in quark matter is estimated to be $10 \sim 100$ MeV [23, 26] while that in normal metals is of order 10K, so they are different by a factor of 10^{11} . It is quite impressive that the same physical mechanism is operative at such drastically different scales. Since heavy flavors are phenomenologically

important as a probe of hot and dense matter in heavy ion collisions and compact stars [34–37], we expect the QCD Kondo effect to play a role there. However it must be noted that the QCD Kondo effect is always subject to competitions with other kind of condensates [24, 26]. At low baryon density the chiral condensate is dominant, while at asymptotically high baryon density the diquark condensate is dominant. Therefore in these two regions the QCD Kondo effect is strongly suppressed. Studies [20, 21, 29] have shown that a chemical potential, an external magnetic field, and a chiral chemical potential all serve as a catalyst of the QCD Kondo effect; however, at the same time they also catalyze chiral symmetry breaking and color superconductivity, so the fate of the QCD Kondo effect is, to say the least, elusive. Nevertheless it is interesting in its own right as a crossroad of condensed matter physics and high energy physics.

Random matrix theory (RMT) is a powerful framework to analyze statistical properties of interacting many-body systems [38, 39]. RMT has been applied to QCD to derive universal microscopic correlation functions of the Dirac eigenvalues at zero [40–44] and nonzero chemical potential [45–55], as well as to map out the QCD phase diagram [56–62] (see [63–67] for reviews). The matching between RMT and lattice simulation data provides a clean way to determine low-energy constants in chiral perturbation theory [68, 69]. Furthermore, since RMT is analytically tractable and yet captures the essence of QCD, it has long been used as a testing ground of newly developed methods to solve the sign problem in QCD, such as the complex Langevin algorithm [70–72], the subset method [73], and the Lefschetz thimbles [74].

In this paper, we aim to build the first random matrix model for the QCD Kondo effect. Since the guiding principle of RMT is the correct matching of symmetries, one must first understand symmetries of the QCD Kondo effect. Let us consider heavy quarks Q that are uniformly distributed in the medium of N_f light flavors ψ . As usual, there is chiral symmetry $U(1)_V \times U(1)_A \times SU(N_f)_L \times SU(N_f)_R$ for ψ . By contrast, it makes no sense to talk about chiral symmetry for Q . Instead, the relevant symmetry of Q is $U(1)_Q \times SU(2)_H$, which is called the heavy quark symmetry (HQS) [34–

37].[75] This $SU(2)_H$ symmetry originates from the fact that when the mass m_Q of Q is large, interactions that flip the spin of Q are suppressed by $1/m_Q$. Now, according to the putative QCD Kondo effect, the heavy-light condensate $\langle \bar{\psi}Q \rangle$ forms [76] and triggers spontaneous symmetry breaking [23, 25]

$$U(1)_V \times U(1)_A \times SU(N_f)_L \times SU(N_f)_R \times U(1)_Q \times SU(2)_H \\ \rightarrow U(1)_{V+Q} \times U(1)_{A+H} \times SU(N_f-1)_L \times SU(N_f-1)_R. \quad (1)$$

Here we of course assumed that there is no other condensate that interferes with the Kondo effect. In (1), $U(1)_{A+H}$ represents the dynamical locking of $U(1)_A$ and an Abelian subgroup of $SU(2)_H$. It is dubbed as chiral-HQS locking in [23, 25] and is a hallmark of the QCD Kondo effect.

In this paper, we construct a random matrix model that correctly implements both the chiral symmetry of light quarks and the heavy quark symmetry of impurities. By taking the large- N limit with N the matrix size, we show for $N_f = 1$ that there appear three distinct phases:

- Pure Kondo phase: $\langle \bar{\psi}Q \rangle \neq 0$ and $\langle \bar{\psi}\psi \rangle = 0$.
- Pure chirally broken phase: $\langle \bar{\psi}Q \rangle = 0$ and $\langle \bar{\psi}\psi \rangle \neq 0$.
- Coexistence phase: $\langle \bar{\psi}Q \rangle \neq 0 \neq \langle \bar{\psi}\psi \rangle$.

One can go to each phase by deliberately tuning a parameter of the model that controls the relative strength of interactions for $\bar{\psi}\psi$ and $\bar{\psi}Q$. For the pure Kondo phase, we show that the symmetry breaking pattern coincides with (1). In particular, the chiral-HQS locking is faithfully realized. For the coexistence phase, we reveal that the pairing form of the Kondo condensate is drastically modified due to the influence of the chiral condensate, which has not been known before. We will also include the chiral chemical potential and examine how the Kondo condensate is deformed. Summarizing, this work opens up a novel way to study quark matter with heavy flavors in an analytically tractable setup.

This paper is organized as follows. In section II we define the new random matrix model and clarify symmetries. In section III the phase structure of the model for $N \gg 1$ is investigated. In section IV we add external source terms and derive the effective theory of Nambu-Goldstone modes. In section V we introduce a chiral chemical potential into the model and study the behavior of Kondo condensates. In section VI we conclude.

II. THE MATRIX MODEL

The random matrix model we propose is defined by the partition function

$$Z = \int dW \int dV \exp[-N\eta \text{Tr}(W^\dagger W) - N \text{Tr}(V^\dagger V)] \\ \times \prod_{f=1}^{N_f} \det(\mathcal{D} + m_f \mathbb{1}_{2N}) \times \det^2(-V^\dagger + \lambda \mathbb{1}_N), \quad (2)$$

where W and V are $N \times N$ complex matrices, η and λ are parameters with $\eta > 0$, and the Dirac operator for light quarks is given by a non-Hermitian $2N \times 2N$ matrix

$$\mathcal{D} \equiv \begin{pmatrix} 0 & W + V \\ -W^\dagger + V & 0 \end{pmatrix}. \quad (3)$$

In the following, we focus on $N_f = 1$ in the chiral limit for technical simplicity. Let us introduce light quark fields $\psi_{i\alpha}$ and $\bar{\psi}_{i\alpha}$ ($i = R, L$) as well as heavy quark fields with spin $Q_{\uparrow\alpha}, Q_{\downarrow\alpha}, \bar{Q}_{\uparrow\alpha}, \bar{Q}_{\downarrow\alpha}$ where α runs from 1 to N . [77] With the aid of these Grassmann variables, we obtain

$$Z = \int d(\bar{\psi}\psi\bar{Q}Q) \int dW \int dV \\ \times \exp[-N\eta \text{Tr}(W^\dagger W) - N \text{Tr}(V^\dagger V) \\ + \bar{\psi}_{R\alpha}(W + V)_{\alpha\beta}\psi_{R\beta} + \bar{\psi}_{L\alpha}(-W^\dagger + V)_{\alpha\beta}\psi_{L\beta} \\ + \bar{Q}_{\uparrow\alpha}(-V^\dagger + \lambda \mathbb{1}_N)_{\alpha\beta}Q_{\uparrow\beta} + \bar{Q}_{\downarrow\alpha}(-V^\dagger + \lambda \mathbb{1}_N)_{\alpha\beta}Q_{\downarrow\beta}]. \quad (4)$$

Evidently the model has the symmetry $U(1)_V \times U(1)_A$ for ψ and $U(1)_Q \times SU(2)_H$ for Q .

Let us integrate out the Gaussian matrices and perform the Hubbard-Stratonovich transformation to make taking the large- N limit easier. For the W part we have

$$\int dW \exp(-N\eta W_{\alpha\beta}^* W_{\alpha\beta} + \bar{\psi}_{R\alpha} W_{\alpha\beta} \psi_{R\beta} - \bar{\psi}_{L\beta} W_{\alpha\beta}^* \psi_{L\alpha}) \\ \propto \exp\left(-\frac{1}{N\eta} \bar{\psi}_{R\alpha} \psi_{R\beta} \bar{\psi}_{L\beta} \psi_{L\alpha}\right) \quad (5) \\ \propto \int_{\mathbb{C}} d\sigma \exp(-N\eta|\sigma|^2 + \bar{\psi}_{R\alpha} \psi_{L\alpha} \sigma + \bar{\psi}_{L\alpha} \psi_{R\alpha} \sigma^*). \quad (6)$$

The V part reads

$$\int dV \exp[-NV_{\alpha\beta}^* V_{\alpha\beta} + (\bar{\psi}_{R\alpha} \psi_{R\beta} + \bar{\psi}_{L\alpha} \psi_{L\beta}) V_{\alpha\beta} \\ - (\bar{Q}_{\uparrow\beta} Q_{\uparrow\alpha} + \bar{Q}_{\downarrow\beta} Q_{\downarrow\alpha}) V_{\alpha\beta}^*] \\ \propto \exp\left[-\frac{1}{N} (\bar{\psi}_{R\alpha} \psi_{R\beta} + \bar{\psi}_{L\alpha} \psi_{L\beta}) (\bar{Q}_{\uparrow\beta} Q_{\uparrow\alpha} + \bar{Q}_{\downarrow\beta} Q_{\downarrow\alpha})\right] \quad (7) \\ \propto \int_{\mathbb{C}^4} dK \exp[-N \text{Tr}(K^\dagger K) + \bar{\psi}_{R\alpha} K_{R\uparrow} Q_{\uparrow\alpha} \\ + \bar{Q}_{\uparrow\alpha} K_{R\uparrow}^* \psi_{R\alpha} + \bar{\psi}_{L\alpha} K_{L\uparrow} Q_{\uparrow\alpha} + \bar{Q}_{\uparrow\alpha} K_{L\uparrow}^* \psi_{L\alpha} \\ + \bar{\psi}_{R\alpha} K_{R\downarrow} Q_{\downarrow\alpha} + \bar{Q}_{\downarrow\alpha} K_{R\downarrow}^* \psi_{R\alpha} + \bar{\psi}_{L\alpha} K_{L\downarrow} Q_{\downarrow\alpha} \\ + \bar{Q}_{\downarrow\alpha} K_{L\downarrow}^* \psi_{L\alpha}], \quad (8)$$

where the notation $K \equiv \begin{pmatrix} K_{R\uparrow} & K_{R\downarrow} \\ K_{L\uparrow} & K_{L\downarrow} \end{pmatrix}$ was used. Collecting everything, we obtain

$$Z \propto \int d(\bar{\psi}\psi\bar{Q}Q) \int_{\mathbb{C}} d\sigma \int_{\mathbb{C}^4} dK \\ \times \exp[-N\eta|\sigma|^2 - N \text{Tr}(K^\dagger K)]$$

$$\times \exp \left[\begin{pmatrix} \bar{\psi}_R \\ \bar{\psi}_L \\ \bar{Q}_\uparrow \\ \bar{Q}_\downarrow \end{pmatrix}_\alpha \begin{pmatrix} 0 & \sigma & K \\ \sigma^* & 0 & \\ K^\dagger & \lambda & 0 \\ 0 & \lambda & \lambda \end{pmatrix} \begin{pmatrix} \psi_R \\ \psi_L \\ Q_\uparrow \\ Q_\downarrow \end{pmatrix}_\alpha \right] \quad (9)$$

$$\propto \int_{\mathbb{C}} d\sigma \int_{\mathbb{C}^4} dK \exp[-N\eta|\sigma|^2 - N\text{Tr}(K^\dagger K)] \times \det^N \left[KK^\dagger - \lambda \begin{pmatrix} 0 & \sigma \\ \sigma^* & 0 \end{pmatrix} \right] \quad (10)$$

$$= \int_{\mathbb{C}} d\sigma \int_{\mathbb{C}^4} dK z(\sigma, K)^N, \quad (11)$$

where we defined (with rescaling $\sigma \rightarrow \sigma/\sqrt{\eta}$)

$$z(\sigma, K) \equiv e^{-|\sigma|^2 - \text{Tr}(K^\dagger K)} \det \left[KK^\dagger - \xi \begin{pmatrix} 0 & \sigma \\ \sigma^* & 0 \end{pmatrix} \right], \quad (12)$$

$$\xi \equiv \frac{\lambda}{\sqrt{\eta}}. \quad (13)$$

Note that this is an exact rewriting of the original partition function and so far no approximation has been made. The parameter ξ controls the strength of chiral symmetry breaking interactions (weaker for small $|\xi|$ and stronger for larger $|\xi|$). The phase diagram of the model as a function of ξ is the subject of the next section.

III. PHASE STRUCTURE AT LARGE N

For $N \gg 1$, the ground state of the model is determined from the saddle point analysis of the function $z(\sigma, K)$ in (12). Through a $U(1)_A$ rotation $K \rightarrow \exp(i\theta\sigma_3)K$ the phase of σ can be arbitrarily tuned, so one can assume $\sigma \geq 0$ without loss of generality. For numerics, we adopt the parametrization

$$KK^\dagger = \begin{pmatrix} a & c + id \\ c - id & b \end{pmatrix} \quad (14)$$

with

$$a, b, c, d \in \mathbb{R}, \quad a \geq 0, b \geq 0, ab \geq c^2 + d^2. \quad (15)$$

Then

$$z(\sigma, K) = e^{-\sigma^2 - a - b} \det \begin{pmatrix} a & c + id - \xi\sigma \\ c - id - \xi\sigma & b \end{pmatrix} \quad (16)$$

$$= e^{-\sigma^2 - a - b} [ab - (c - \xi\sigma)^2 - d^2]. \quad (17)$$

Suppose we rotate the vector $(c, d) \in \mathbb{R}^2$ with σ, a, b and $c^2 + d^2$ fixed. Since $ab - (c - \xi\sigma)^2 - d^2 = ab - c^2 - d^2 - \xi^2\sigma^2 + 2c\xi\sigma = 2 \begin{pmatrix} c \\ d \end{pmatrix} \cdot \begin{pmatrix} \xi\sigma \\ 0 \end{pmatrix} + \text{const.}$, it is obvious that $ab - (c - \xi\sigma)^2 - d^2$ is extremized if and only if the vector (c, d) is parallel to $(\xi\sigma, 0)$. Thus $d = 0$ at the saddle




FIG. 1. The minimum of the free energy (18) for varying ξ . b is not plotted because $a = b$ for all ξ .

point. Taking this observation into account, we find that the task is to minimize the free energy

$$f(a, b, c, \sigma) \equiv \sigma^2 + a + b - \log |ab - (c - \xi\sigma)^2| \quad (18)$$

under the conditions $\sigma \geq 0, a \geq 0, b \geq 0$ and $ab \geq c^2$. We numerically solved this problem. The result is plotted in Figure 1. Here is a brief summary of the phase structure.

Domain	Phase	Condensates
$0 \leq \xi \leq 0.5557$	Pure Kondo phase	$\langle K \rangle = \mathbb{1}_2, \langle \sigma \rangle = 0$
$0.5557 \leq \xi \leq 1$	Coexistence phase	$\langle K \rangle \neq 0, \langle \sigma \rangle \neq 0$
$\xi \geq 1$	Pure chirally broken phase	$\langle K \rangle = 0, \langle \sigma \rangle = 1$

In the pure Kondo phase at $\xi \leq 0.5557$, $KK^\dagger = \mathbb{1}_2$ in the ground state. Therefore, up to unitary rotations $K \rightarrow KU$ with $U \in U(2)$, we have $K = \mathbb{1}_2$, i.e., $\langle K_{R\uparrow} \rangle = \langle K_{L\downarrow} \rangle = 1$ and $\langle K_{R\downarrow} \rangle = \langle K_{L\uparrow} \rangle = 0$. Hence the symmetry breaking $U(1)_V \times U(1)_Q \rightarrow U(1)_{V+Q}$ occurs. Furthermore, recalling that K transforms under $U(1)_A \times SU(2)_H$ as

$$K \rightarrow e^{i\theta\sigma_3} K u \quad \text{with } e^{i\theta\sigma_3} \in U(1)_A \text{ and } u \in SU(2)_H, \quad (19)$$

it follows that $K = \mathbb{1}_2$ spontaneously breaks $U(1)_A \times SU(2)_H$ down to $U(1)_{A+H}$. This is the random matrix incarnation of the chiral-HQS locking advocated in [23, 25].

In the coexistence phase at $0.5557 \leq \xi \leq 1$, both the Kondo condensate and the chiral condensate are nonzero. Analytically one can show

$$a = \frac{4 - \xi^2 - \xi\sqrt{\xi^2 + 8}}{8}, \quad (20)$$

$$\sigma = \frac{\xi + \sqrt{\xi^2 + 8}}{4}. \quad (21)$$

Specifically, we have $KK^\dagger = a \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}$. There are infinitely many K that fulfill this form due to unitary invariance ($K \rightarrow KU$). Let us mention two particular choices. The first one is $\langle K \rangle = \sqrt{a} \begin{pmatrix} 1 & 0 \\ -1 & 0 \end{pmatrix}$. Namely,

$$\langle K_{R\uparrow} \rangle = -\langle K_{L\uparrow} \rangle = \sqrt{a} \quad \text{and} \quad \langle K_{R\downarrow} \rangle = \langle K_{L\downarrow} \rangle = 0, \quad (22)$$

implying that only up spin component of Q participates in the Kondo effect. The second one is $\langle K \rangle = \sqrt{\frac{a}{2}} \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix}$. This is a more symmetric choice, since all components equally contribute to the Kondo effect:

$$\langle K_{R\uparrow} \rangle = \langle K_{R\downarrow} \rangle = -\langle K_{L\uparrow} \rangle = -\langle K_{L\downarrow} \rangle = \sqrt{\frac{a}{2}}. \quad (23)$$

The relative minus sign of $\langle K_R \rangle$ and $\langle K_L \rangle$ in (23) arises due to the assumption that $\sigma \geq 0$. For either choice of K , one can conclude that a nonzero chiral condensate leads to breakdown of the chiral-HQS locking.

Finally, the phase for $\xi \geq 1$ is an ordinary chiral symmetry broken phase with no Kondo effect. The only spontaneously broken symmetry is $U(1)_A$.

IV. LOW-ENERGY EFFECTIVE THEORY

The procedure to derive low-energy effective theory of soft fluctuations in random matrix models is established (see e.g., [40, 78]). Following this route, in this section we examine the large- N behavior of the partition function in the presence of external fields. In addition to quark masses, we shall insert the source term for the Kondo condensates

$$j_R \bar{\psi}_R Q_\uparrow + j_L \bar{\psi}_L Q_\downarrow + j_R^* \bar{Q}_\uparrow \psi_R + j_L^* \bar{Q}_\downarrow \psi_L. \quad (24)$$

Inserting this into (2) we find the fermion determinant (for $N_f = 1$) to be

$$\det \begin{pmatrix} W+V & m & j_R & 0 \\ m^* & -W^\dagger+V & 0 & j_L \\ j_R^* & 0 & -V^\dagger+\lambda & 0 \\ 0 & j_L^* & 0 & -V^\dagger+\lambda \end{pmatrix} = \begin{cases} \det[(W+V)(W^\dagger-V) + |m|^2] \times \det^2(-V^\dagger+\lambda) \\ \quad \text{for } j_{R/L} \rightarrow 0, \text{ and} \\ \det[(W+V)(V^\dagger-\lambda) + |j_R|^2] \\ \times \det[(-W^\dagger+V)(V^\dagger-\lambda) + |j_L|^2] \quad \text{for } m \rightarrow 0. \end{cases} \quad (25)$$

The sources m and $j_{L/R}$ are similar to a magnetic field applied to a ferromagnet. Generally, in a phase with

spontaneously broken symmetry, an infinitesimal external perturbation determines the orientation of a macroscopically large condensate. In QCD, the quark mass, however small it is, determines the orientation of the chiral condensate. Recalling that the fermion determinant in QCD has the form $\det(\not{D} + m)$, such a phenomenon is possible only if \not{D} has a macroscopically large number of eigenvalues in the vicinity of zero. This is the content of the Banks-Casher relation [79]. In our matrix model, m couples to the matrix $(W+V)(W^\dagger-V)$ and j to $(W+V)(V^\dagger-\lambda)$ and $(-W^\dagger+V)(V^\dagger-\lambda)$. Therefore, in a phase with nonzero chiral/Kondo condensates, these matrices must have a macroscopically large number of eigenvalues near the origin. The level fluctuations on the scale of a mean level spacing in this domain are believed to be universal [38, 64]. One method to probe this regime in RMT is to directly diagonalize these matrices and derive spectral functions analytically. However, this direction seems difficult if not impossible for the current matrix model because W and V are intertwined in a peculiar way. Therefore we shall turn to another method: to derive the effective theory of the partition function coupled to external fields. It uncovers fruitful information about statistical properties of the underlying operator coupled to an external field [80].

Our point of departure is (9) with source terms included and σ rescaled ($\sigma \rightarrow \sigma/\sqrt{\eta}$):

$$Z \propto \int_{\mathbb{C}} d\sigma \int_{\mathbb{C}^4} dK \exp[-N|\sigma|^2 - N \text{Tr}(K^\dagger K)] \times \det^N \left(\begin{array}{cc|c} 0 & \sigma + \sqrt{\eta}m & K + J \\ \sigma^* + \sqrt{\eta}m^* & 0 & \\ \hline K^\dagger + J^\dagger & & \xi \mathbb{1}_2 \end{array} \right) \quad (26)$$

with $J \equiv \text{diag}(j_R, j_L)$. We will take the large- N limit with $m \sim j_{L/R} \sim O(1/N)$ and define the microscopic variables

$$\hat{m} \equiv Nm, \quad \hat{j}_{L/R} \equiv Nj_{L/R} \quad \text{and} \quad \hat{J} \equiv NJ. \quad (27)$$

We begin with the pure chirally broken phase with $\langle \sigma \rangle = 1$ and $\langle K \rangle = 0$. Substituting $\sigma = e^{-i\phi}$ and $K = 0$ into (26), one can straightforwardly derive the large- N behavior

$$Z \propto \int_0^{2\pi} d\phi \exp[\sqrt{\eta}(\hat{m} e^{i\phi} + \hat{m}^* e^{-i\phi})] \quad (28)$$

$$\propto I_0(2\sqrt{\eta}|\hat{m}|). \quad (29)$$

Here ϕ is the Nambu-Goldstone mode of $U(1)_A$, and I_0 is the modified Bessel function of the first kind. The light and heavy quarks decouple.

Next we consider the pure Kondo phase in which $\langle \sigma \rangle = 0$ and $\langle K \rangle = \mathbb{1}_2$. Substituting $\sigma = 0$ and $K = U$ with $U \in U(2)$ into (26) we readily obtain

$$Z \propto \int_{U(2)} dU \exp[\text{Tr}(\hat{J}U^\dagger + U\hat{J}^\dagger)]. \quad (30)$$

Interestingly enough, this coincides exactly with the finite-volume partition function of two-flavor QCD in the ε -regime [40, 80]. The role of the quark mass matrix is now played by \hat{J} . This integral can be analytically carried out [81, 82] and yields (for real \hat{J})

$$Z \propto \frac{\hat{j}_L I_0(2\hat{j}_R) I_1(2\hat{j}_L) - \hat{j}_R I_0(2\hat{j}_L) I_1(2\hat{j}_R)}{\hat{j}_L^2 - \hat{j}_R^2}. \quad (31)$$

Finally we turn to the coexistence phase. This is by far

the most complicated case. We denote the ground state as $\sigma_0 \equiv \langle \sigma \rangle$ and $K_0 \equiv \langle K \rangle$ [cf. (20) and (21)]. The five Nambu-Goldstone modes can be introduced as

$$\sigma = e^{2i\phi} \sigma_0, \quad (32)$$

$$K = e^{i\phi\sigma_3} K_0 U, \quad (33)$$

with $U \in U(2)$. From (26) we then get

$$Z \propto \oint d\phi \int_{U(2)} dU \det^N \left(\begin{array}{c|c} 0 & e^{2i\phi} \sigma_0 + \sqrt{\eta} m \\ \hline e^{-2i\phi} \sigma_0 + \sqrt{\eta} m^* & 0 \end{array} \middle| \begin{array}{c} e^{i\phi\sigma_3} K_0 U + J \\ \xi \mathbb{1}_2 \end{array} \right) \quad (34)$$

$$\approx \oint d\phi \int_{U(2)} dU \exp \left\{ -\sqrt{\eta} \xi \text{Tr} \left[R^{-1} \begin{pmatrix} 0 & \hat{m} e^{-2i\phi} \\ \hat{m}^* e^{2i\phi} & 0 \end{pmatrix} \right] + 2 \text{Re Tr} \left(R^{-1} \hat{J} e^{-i\phi\sigma_3} U^\dagger K_0^\dagger \right) \right\}, \quad (35)$$

where

$$R \equiv K_0 K_0^\dagger - \xi \sigma_0 \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (36)$$

Notice that in the second term of the exponent, $e^{-i\phi\sigma_3}$ can be absorbed into U . It leads to factorization of the partition function. Using $K_0 = \sqrt{a} \begin{pmatrix} 1 & 0 \\ -1 & 0 \end{pmatrix}$, we obtain

$$Z = \oint d\phi \exp [\sqrt{\eta} C_1(\xi) (\hat{m} e^{-2i\phi} + \hat{m}^* e^{2i\phi})] \times \int_{U(2)} dU \exp \left\{ C_2(\xi) \text{Re Tr} \left[\begin{pmatrix} \hat{j}_R & -\hat{j}_L \\ 0 & 0 \end{pmatrix} U^\dagger \right] \right\} \quad (37)$$

where

$$C_1(\xi) \equiv \frac{\xi + \sqrt{\xi^2 + 8}}{4}, \quad (38)$$

$$C_2(\xi) \equiv \sqrt{\frac{4 - \xi^2 - \xi \sqrt{\xi^2 + 8}}{2}}. \quad (39)$$

As a quick consistency check, notice that $C_1 \rightarrow 1$ and $C_2 \rightarrow 0$ as $\xi \rightarrow 1$, ensuring that the effective theory for the pure chirally broken phase (28) is smoothly recovered at the phase transition $\xi = 1$.

The integral (37) can be carried out exactly [81, 82] and yields

$$Z \propto I_0(2\sqrt{\eta} C_1(\xi) |\hat{m}|) \frac{I_1 \left(C_2(\xi) \sqrt{|\hat{j}_R|^2 + |\hat{j}_L|^2} \right)}{\sqrt{|\hat{j}_R|^2 + |\hat{j}_L|^2}}. \quad (40)$$

V. CHIRAL CHEMICAL POTENTIAL

The chiral chemical potential $\mu_5 \bar{\psi} \gamma_0 \gamma_5 \psi$ introduces a net imbalance of chiralities. Its use is motivated by the chiral magnetic effect [83, 84] in which a current is induced by a magnetic field in a chirally asymmetric matter. The interplay of the chiral chemical potential and the QCD Kondo effect was analyzed in an NJL-type model in [29]. Here we discuss it in the context of our matrix model. For simplicity, we switch off the interaction part of W , which amounts to neglecting chiral symmetry breaking entirely. By slightly modifying (9), we get the partition function

$$Z = \int d(\bar{\psi} \psi \bar{Q} Q) \int_{\mathbb{C}^4} dK \exp[-N \text{Tr}(K^\dagger K)] \times \exp \left[\begin{pmatrix} \bar{\psi}_R \\ \bar{\psi}_L \\ \bar{Q}_\uparrow \\ \bar{Q}_\downarrow \end{pmatrix}_\alpha \begin{pmatrix} \rho & 0 & K \\ 0 & -\rho & \\ K^\dagger & \lambda & 0 \\ 0 & \lambda & \end{pmatrix} \begin{pmatrix} \psi_R \\ \psi_L \\ Q_\uparrow \\ Q_\downarrow \end{pmatrix}_\alpha \right] \quad (41)$$

where ρ represents the chiral chemical potential. Integrating out fermions yields

$$Z \propto \int_{\mathbb{C}^4} dK \left\{ e^{-\text{Tr}(K^\dagger K)} \det(K K^\dagger - \mu_5 \sigma_3) \right\}^N \quad (42)$$

with

$$\mu_5 \equiv \rho \lambda. \quad (43)$$

Again we employ the parametrization (14). As $\det(K K^\dagger - \mu_5 \sigma_3)$ depends on c and d through the combination $c^2 + d^2$ one can set $d = 0$ without loss of generality. The task is to minimize the free energy

$$f_5(a, b, c) \equiv a + b - \log |(a - \mu_5)(b + \mu_5) - c^2| \quad (44)$$




FIG. 2. The Kondo condensates for varying μ_5 .

under the conditions $a \geq 0, b \geq 0$ and $ab \geq c^2$. We solved this problem numerically and found that $c = 0$ for all μ_5 . The condensates are then obtained as $\langle K_{R\uparrow} \rangle = \sqrt{a}$ and $\langle K_{L\downarrow} \rangle = \sqrt{b}$. The numerical results are plotted in Figure 2 and are summarized below.

Domain	$\langle K_{R\uparrow} \rangle$	$\langle K_{L\downarrow} \rangle$
$-1 \leq \mu_5 \leq -\mu_5^C$	$\sqrt{1 + \mu_5}$	0
$-\mu_5^C \leq \mu_5 \leq \mu_5^C$	$\sqrt{1 + \mu_5}$	$\sqrt{1 - \mu_5}$
$\mu_5^C \leq \mu_5 \leq 1$	0	$\sqrt{1 - \mu_5}$
$ \mu_5 > 1$	0	0

There are phase transitions at $\mu_5 = \pm 1$ and $\pm\mu_5^C$, where

$$\mu_5^C \equiv 0.27846 \dots \quad (45)$$

is the solution of $e^{1+x}x = 1$. As expected on physical grounds, μ_5 lifts the degeneracy of left-handed and right-handed Kondo condensates. At large μ_5 the condensates go away, which we think is an artifact of the model, similar to the UV cutoff effects in NJL models.

VI. CONCLUSIONS AND OUTLOOK

In this work, we have constructed a new random matrix model that captures the essence of the QCD Kondo effect. The model has correct symmetries of QCD with a heavy flavor. We solved the model for $N_f = 1$ in the large- N limit with N the matrix size, and classified the patterns of spontaneous symmetry breaking. The model has three phases that arise due to the competition of chiral symmetry breaking and the QCD Kondo effect. In the Kondo phase with no chiral condensate, we have shown that the $U(1)_A$ symmetry of light quarks and the $SU(2)$ heavy quark spin symmetry are locked to the diagonal $U(1)$ subgroup, as expected for the QCD Kondo phase. In the phase where the Kondo effect and chiral symmetry breaking coexist, we have shown that the pairing form of the Kondo condensate is drastically modified. This is an important prediction of our model. Moreover, for each of the three phases we derived the low-energy effective theory of Nambu-Goldstone modes and obtained closed formulas for the partition function. We hope this work contributes to better analytical understanding of the QCD phase diagram with heavy quarks.

There are various future directions of research.

- The analysis in this paper is limited to $N_f = 1$. Extension to $N_f > 1$ is a challenging open problem.
- The axial anomaly was ignored in this work. How to incorporate anomaly into our random matrix model is not clear yet.
- Our prediction for the pairing form of the Kondo condensate in the coexistence phase should be checked in more realistic NJL-type models.
- As is well known, there are three symmetry classes in chiral RMT, corresponding to quarks in a complex/real/pseudoreal representation of the gauge group [85]. The matrix model in this work corresponds to QCD with quarks in a complex representation. Extension to the other two classes (the chiral orthogonal ensemble and the chiral symplectic ensemble) will be very interesting.

Work in some of these directions is in progress.

[1] J. Kondo, Resistance Minimum in Dilute Magnetic Alloys, *Prog. Theor. Phys.* **32**, 37 (1964).
[2] A. C. Hewson, *The Kondo Problem to Heavy Fermions*, Cambridge Studies in Magnetism (Cambridge University Press, 1993).
[3] P. Coleman, Heavy Fermions: electrons at the edge of magnetism, *Handbook of Magnetism and Advanced Magnetic Materials*, 95 (2006), [arXiv:cond-mat/0612006](https://arxiv.org/abs/cond-mat/0612006).
[4] A. C. Hewson and J. Kondo, Kondo effect, *Scholarpedia* **4**, 7529 (2009).
[5] D. Withoff and E. Fradkin, Phase transitions in gapless

Fermi systems with magnetic impurities, *Phys. Rev. Lett.* **64**, 1835 (1990).
[6] C. R. Cassanello and E. Fradkin, Kondo effect in flux phases, *Phys. Rev. B* **53**, 15079 (1996).
[7] C. Gonzalez-Buxton and K. Ingersent, Renormalization-group study of Anderson and Kondo impurities in gapless Fermi systems, *Phys. Rev. B* **57**, 14254 (1998).
[8] L. Fritz and M. Vojta, Phase transitions in the pseudogap Anderson and Kondo models: Critical dimensions, renormalization group, and local-moment criticality, *Phys. Rev. B* **70**, 214427 (2004).

- [9] L. Fritz, S. Florens, and M. Vojta, Universal crossovers and critical dynamics of quantum phase transitions: A renormalization group study of the pseudogap Kondo problem, *Phys. Rev. B* **74**, 144410 (2006).
- [10] J.-H. Chen, L. Li, W. G. Cullen, E. D. Williams, and M. S. Fuhrer, Tunable Kondo effect in graphene with defects, *Nature Physics* **7**, 535 (2011).
- [11] L. Fritz and M. Vojta, The physics of Kondo impurities in graphene, *Reports on Progress in Physics* **76**, 032501 (2013).
- [12] A. Principi, G. Vignale, and E. Rossi, Kondo effect and non-Fermi-liquid behavior in Dirac and Weyl semimetals, *Phys. Rev. B* **92**, 041107 (2015).
- [13] A. K. Mitchell and L. Fritz, Kondo effect in three-dimensional Dirac and Weyl systems, *Phys. Rev. B* **92**, 121109 (2015).
- [14] J.-H. Sun, D.-H. Xu, F.-C. Zhang, and Y. Zhou, Magnetic impurity in a Weyl semimetal, *Phys. Rev. B* **92**, 195124 (2015).
- [15] T. Yanagisawa, Kondo Effect in Dirac Systems, *J. Phys. Soc. Japan* **84**, 074705 (2015).
- [16] L. Li, J.-H. Sun, Z.-H. Wang, D.-H. Xu, H.-G. Luo, and W.-Q. Chen, Magnetic states and Kondo screening in Weyl semimetals with chiral anomaly, *Phys. Rev. B* **98**, 075110 (2018).
- [17] Y. Nambu and G. Jona-Lasinio, Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. I, *Phys. Rev.* **122**, 345 (1961).
- [18] Y. Nambu and G. Jona-Lasinio, Dynamical Model of Elementary Particles Based on an Analogy with Superconductivity. II, *Phys. Rev.* **124**, 246 (1961).
- [19] S. Yasui and K. Sudoh, Heavy-quark dynamics for charm and bottom flavor on the Fermi surface at zero temperature, *Phys. Rev. C* **88**, 015201 (2013), [arXiv:1301.6830 \[hep-ph\]](#).
- [20] K. Hattori, K. Itakura, S. Ozaki, and S. Yasui, QCD Kondo effect: quark matter with heavy-flavor impurities, *Phys. Rev. D* **92**, 065003 (2015), [arXiv:1504.07619 \[hep-ph\]](#).
- [21] S. Ozaki, K. Itakura, and Y. Kuramoto, Magnetically induced QCD Kondo effect, *Phys. Rev. D* **94**, 074013 (2016), [arXiv:1509.06966 \[hep-ph\]](#).
- [22] S. Yasui, Kondo effect in charm and bottom nuclei, *Phys. Rev. C* **93**, 065204 (2016), [arXiv:1602.00227 \[hep-ph\]](#).
- [23] S. Yasui, K. Suzuki, and K. Itakura, Kondo phase diagram of quark matter, *Nucl. Phys. A* **983**, 90 (2019), [arXiv:1604.07208 \[hep-ph\]](#).
- [24] T. Kanazawa and S. Uchino, Overscreened Kondo effect, (color) superconductivity and Shiba states in Dirac metals and quark matter, *Phys. Rev. D* **94**, 114005 (2016), [arXiv:1609.00033 \[cond-mat.str-el\]](#).
- [25] S. Yasui, K. Suzuki, and K. Itakura, Topology and stability of the Kondo phase in quark matter, *Phys. Rev. D* **96**, 014016 (2017), [arXiv:1703.04124 \[hep-ph\]](#).
- [26] K. Suzuki, S. Yasui, and K. Itakura, Interplay between chiral symmetry breaking and the QCD Kondo effect, *Phys. Rev. D* **96**, 114007 (2017), [arXiv:1708.06930 \[hep-ph\]](#).
- [27] T. Kimura and S. Ozaki, Conformal field theory analysis of the QCD Kondo effect, *Phys. Rev. D* **99**, 014040 (2019), [arXiv:1806.06486 \[hep-ph\]](#).
- [28] K. Hattori, X.-G. Huang, and R. D. Pisarski, Emergent QCD Kondo effect in two-flavor color superconducting phase, *Phys. Rev. D* **99**, 094044 (2019), [arXiv:1903.10953 \[hep-ph\]](#).
- [29] D. Suenaga, K. Suzuki, Y. Araki, and S. Yasui, Kondo effect driven by chirality imbalance, (2019), [arXiv:1912.12669 \[hep-ph\]](#).
- [30] P. Nozieres and A. Blandin, Kondo effect in real metals, *J. Phys. France* **41**, 193 (1980).
- [31] I. Affleck, A Current Algebra Approach to the Kondo Effect, *Nucl. Phys. B* **336**, 517 (1990).
- [32] I. Affleck and A. W. Ludwig, Critical theory of overscreened Kondo fixed points, *Nucl. Phys. B* **360**, 641 (1991).
- [33] O. Parcollet, A. Georges, G. Kotliar, and A. Sengupta, Overscreened multichannel SU(N) Kondo model: Large-N solution and conformal field theory, *Phys. Rev. B* **58**, 3794 (1998), [arXiv:cond-mat/9711192](#).
- [34] M. Neubert, Heavy quark symmetry, *Phys. Rept.* **245**, 259 (1994), [arXiv:hep-ph/9306320](#).
- [35] M. Neubert, Heavy quark effective theory, *Subnucl. Ser.* **34**, 98 (1997), [arXiv:hep-ph/9610266](#).
- [36] A. V. Manohar and M. B. Wise, *Heavy Quark Physics*, Cambridge Monographs on Particle Physics, Nuclear Physics and Cosmology (Cambridge University Press, 2000).
- [37] A. Hosaka, T. Hyodo, K. Sudoh, Y. Yamaguchi, and S. Yasui, Heavy Hadrons in Nuclear Matter, *Prog. Part. Nucl. Phys.* **96**, 88 (2017), [arXiv:1606.08685 \[hep-ph\]](#).
- [38] T. Guhr, A. Muller-Groeling, and H. A. Weidenmuller, Random matrix theories in quantum physics: Common concepts, *Phys. Rept.* **299**, 189 (1998), [arXiv:cond-mat/9707301](#).
- [39] G. Akemann, J. Baik, and P. Di Francesco, *The Oxford Handbook of Random Matrix Theory*, Oxford Handbooks in Mathematics (Oxford University Press, 2011).
- [40] E. V. Shuryak and J. Verbaarschot, Random matrix theory and spectral sum rules for the Dirac operator in QCD, *Nucl. Phys. A* **560**, 306 (1993), [arXiv:hep-th/9212088](#).
- [41] J. Verbaarschot and I. Zahed, Spectral density of the QCD Dirac operator near zero virtuality, *Phys. Rev. Lett.* **70**, 3852 (1993), [arXiv:hep-th/9303012](#).
- [42] J. J. M. Verbaarschot, The Spectrum of the Dirac operator near zero virtuality for $N(c) = 2$ and chiral random matrix theory, *Nucl. Phys. B* **426**, 559 (1994), [arXiv:hep-th/9401092](#).
- [43] G. Akemann, P. Damgaard, U. Magnea, and S. Nishigaki, Universality of random matrices in the microscopic limit and the Dirac operator spectrum, *Nucl. Phys. B* **487**, 721 (1997), [arXiv:hep-th/9609174](#).
- [44] P. H. Damgaard and S. M. Nishigaki, Universal spectral correlators and massive Dirac operators, *Nucl. Phys. B* **518**, 495 (1998), [arXiv:hep-th/9711023](#).
- [45] J. C. Osborn, Universal results from an alternate random matrix model for QCD with a baryon chemical potential, *Phys. Rev. Lett.* **93**, 222001 (2004), [arXiv:hep-th/0403131](#).
- [46] G. Akemann, J. Osborn, K. Splittorff, and J. Verbaarschot, Unquenched QCD Dirac operator spectra at nonzero baryon chemical potential, *Nucl. Phys. B* **712**, 287 (2005), [arXiv:hep-th/0411030](#).
- [47] J. Osborn, K. Splittorff, and J. Verbaarschot, Chiral symmetry breaking and the Dirac spectrum at nonzero chemical potential, *Phys. Rev. Lett.* **94**, 202001 (2005), [arXiv:hep-th/0501210](#).
- [48] G. Akemann, The Complex Laguerre symplectic ensemble of non-Hermitian matrices, *Nucl. Phys. B* **730**, 253

- (2005), [arXiv:hep-th/0507156](#).
- [49] T. Kanazawa, T. Wettig, and N. Yamamoto, Chiral random matrix theory for two-color QCD at high density, *Phys. Rev. D* **81**, 081701 (2010), [arXiv:0912.4999 \[hep-ph\]](#).
- [50] G. Akemann, T. Kanazawa, M. Phillips, and T. Wettig, Random matrix theory of unquenched two-colour QCD with nonzero chemical potential, *JHEP* **03**, 066, [arXiv:1012.4461 \[hep-lat\]](#).
- [51] T. Kanazawa, T. Wettig, and N. Yamamoto, Singular values of the Dirac operator in dense QCD-like theories, *JHEP* **12**, 007, [arXiv:1110.5858 \[hep-ph\]](#).
- [52] T. Kanazawa and T. Wettig, Stressed Cooper pairing in QCD at high isospin density: effective Lagrangian and random matrix theory, *JHEP* **10**, 055, [arXiv:1406.6131 \[hep-ph\]](#).
- [53] T. Kanazawa, Color-flavor locking and 2SC pairing in random matrix theory, (2020), [arXiv:2004.04483 \[hep-th\]](#).
- [54] T. Kanazawa, Chiral random matrix theory for single-flavor spin-one Cooper pairing, (2020), [arXiv:2004.13680 \[hep-th\]](#).
- [55] T. Kanazawa, Chiral random matrix theory for colorful quark-antiquark condensates, (2020), [arXiv:2005.08471 \[hep-th\]](#).
- [56] A. M. Halasz, A. Jackson, R. Shrock, M. A. Stephanov, and J. Verbaarschot, On the phase diagram of QCD, *Phys. Rev. D* **58**, 096007 (1998), [arXiv:hep-ph/9804290](#).
- [57] B. Vanderheyden and A. Jackson, Random matrix model for chiral symmetry breaking and color superconductivity in QCD at finite density, *Phys. Rev. D* **62**, 094010 (2000), [arXiv:hep-ph/0003150](#).
- [58] S. Pepin and A. Schafer, QCD at high baryon density in a random matrix model, *Eur. Phys. J. A* **10**, 303 (2001), [arXiv:hep-ph/0010225](#).
- [59] B. Vanderheyden and A. Jackson, Random matrix study of the phase structure of QCD with two colors, *Phys. Rev. D* **64**, 074016 (2001), [arXiv:hep-ph/0102064](#).
- [60] B. Klein, D. Toublan, and J. Verbaarschot, The QCD phase diagram at nonzero temperature, baryon and isospin chemical potentials in random matrix theory, *Phys. Rev. D* **68**, 014009 (2003), [arXiv:hep-ph/0301143](#).
- [61] B. Klein, D. Toublan, and J. Verbaarschot, Diquark and pion condensation in random matrix models for two color QCD, *Phys. Rev. D* **72**, 015007 (2005), [arXiv:hep-ph/0405180](#).
- [62] B. Vanderheyden and A. Jackson, Random matrix models for phase diagrams, *Rept. Prog. Phys.* **74**, 102001 (2011), [arXiv:1105.1291 \[hep-ph\]](#).
- [63] J. J. M. Verbaarschot, Universal behavior in Dirac spectra, in *NATO Advanced Study Institute on Confinement, Duality and Nonperturbative Aspects of QCD* (1997) pp. 343–378, [arXiv:hep-th/9710114](#).
- [64] J. Verbaarschot and T. Wettig, Random matrix theory and chiral symmetry in QCD, *Ann. Rev. Nucl. Part. Sci.* **50**, 343 (2000), [arXiv:hep-ph/0003017](#).
- [65] J. Verbaarschot, QCD, chiral random matrix theory and integrability, in *Application of random matrices in physics. Proceedings, NATO Advanced Study Institute, Les Houches, France, June 6-25, 2004* (2005) pp. 163–217, [arXiv:hep-th/0502029](#).
- [66] G. Akemann, Matrix Models and QCD with Chemical Potential, *Int. J. Mod. Phys. A* **22**, 1077 (2007), [arXiv:hep-th/0701175](#).
- [67] T. Kanazawa, *Dirac Spectra in Dense QCD* (Springer Japan, 2013).
- [68] M. Berbenni-Bitsch, S. Meyer, A. Schafer, J. Verbaarschot, and T. Wettig, Microscopic universality in the spectrum of the lattice Dirac operator, *Phys. Rev. Lett.* **80**, 1146 (1998), [arXiv:hep-lat/9704018](#).
- [69] H. Fukaya *et al.* (JLQCD), Two-flavor lattice QCD simulation in the epsilon-regime with exact chiral symmetry, *Phys. Rev. Lett.* **98**, 172001 (2007), [arXiv:hep-lat/0702003](#).
- [70] A. Mollgaard and K. Splittorff, Complex Langevin Dynamics for chiral Random Matrix Theory, *Phys. Rev. D* **88**, 116007 (2013), [arXiv:1309.4335 \[hep-lat\]](#).
- [71] K. Nagata, J. Nishimura, and S. Shimasaki, Gauge cooling for the singular-drift problem in the complex Langevin method - a test in Random Matrix Theory for finite density QCD, *JHEP* **07**, 073, [arXiv:1604.07717 \[hep-lat\]](#).
- [72] J. Bloch, J. Glesaaen, J. Verbaarschot, and S. Zafeiropoulos, Complex Langevin Simulation of a Random Matrix Model at Nonzero Chemical Potential, *JHEP* **03**, 015, [arXiv:1712.07514 \[hep-lat\]](#).
- [73] J. Bloch, A subset solution to the sign problem in random matrix simulations, *Phys. Rev. D* **86**, 074505 (2012), [arXiv:1205.5500 \[hep-lat\]](#).
- [74] F. Di Renzo and G. Eruzzi, Thimble regularization at work: from toy models to chiral random matrix theories, *Phys. Rev. D* **92**, 085030 (2015), [arXiv:1507.03858 \[hep-lat\]](#).
- [75] In the presence of N_h flavors of heavy quarks, the heavy quark symmetry is enlarged to $U(1)_Q \times SU(2N_h)$. However, in this paper we only consider a single heavy flavor.
- [76] This is a relativistic analog of the hybridization of impurity spin and conduction electrons in the Coqblin-Schreiffer model [86, 87].
- [77] We suppress the spinor index of ψ because it is not necessary for the description of symmetry breaking. Minimalism is a basic tenet of RMT.
- [78] A. M. Halasz and J. Verbaarschot, Effective Lagrangians and chiral random matrix theory, *Phys. Rev. D* **52**, 2563 (1995), [arXiv:hep-th/9502096](#).
- [79] T. Banks and A. Casher, Chiral Symmetry Breaking in Confining Theories, *Nucl. Phys.* **B169**, 103 (1980).
- [80] H. Leutwyler and A. V. Smilga, Spectrum of Dirac operator and role of winding number in QCD, *Phys. Rev. D* **46**, 5607 (1992).
- [81] A. Jackson, M. Sener, and J. Verbaarschot, Finite volume partition functions and Itzykson-Zuber integrals, *Phys. Lett. B* **387**, 355 (1996), [arXiv:hep-th/9605183](#).
- [82] A. Balantekin, Character expansions, Itzykson-Zuber integrals, and the QCD partition function, *Phys. Rev. D* **62**, 085017 (2000), [arXiv:hep-th/0007161](#).
- [83] D. E. Kharzeev, L. D. McLerran, and H. J. Warringa, The Effects of topological charge change in heavy ion collisions: 'Event by event P and CP violation', *Nucl. Phys. A* **803**, 227 (2008), [arXiv:0711.0950 \[hep-ph\]](#).
- [84] K. Fukushima, D. E. Kharzeev, and H. J. Warringa, The Chiral Magnetic Effect, *Phys. Rev. D* **78**, 074033 (2008), [arXiv:0808.3382 \[hep-ph\]](#).
- [85] J. J. M. Verbaarschot, Spectrum of the QCD Dirac operator and chiral random matrix theory, *Phys. Rev. Lett.* **72**, 2531 (1994), [arXiv:hep-th/9401059](#).
- [86] N. Read and D. M. Newns, On the solution of the Coqblin-Schreiffer Hamiltonian by the large-N expansion

technique, *J. Phys. C: Solid State Phys.* **16**, 3273 (1983).
[87] N. Bickers, Review of techniques in the large- N expan-

sion for dilute magnetic alloys, *Rev. Mod. Phys.* **59**, 845 (1987).