

# The Effects of Stringent Interventions for Coronavirus Pandemic

Ting Tian<sup>a,1</sup>, Wenxiang Luo<sup>a,1</sup>, Yukang Jiang<sup>a,1</sup>, Minqiong Chen<sup>a</sup>, Canhong Wen<sup>b</sup>, Wenliang Pan<sup>a</sup>, and Xueqin Wang<sup>b,2</sup>

<sup>a</sup>*School of Mathematics, Sun Yat-sen University*

<sup>b</sup>*School of Management, University of Science and Technology of China*

## Abstract

The pandemic of COVID-19 has caused severe public health consequences around the world. Many interventions of COVID-19 have been implemented. It is of great public health and societal importance to evaluate the effects of interventions in the pandemic of COVID-19. In this paper, with help of synthetic control method, regression discontinuity and a Susceptible-Infected and infectious without isolation-Hospitalized in isolation-Removed (SIHR) model, we evaluate the horizontal and longitudinal effects of stringent interventions implemented in Wenzhou, a representative urban city of China, where stringent interventions were enforced to curb its own epidemic situation with rapidly increasing newly confirmed cases. We found that there were statistically significant treatment effects of those stringent interventions which reduced the cumulative confirmed cases of COVID-19. Those reduction effects would increase over time. Also, if the stringent interventions were delayed by 2 days or mild interventions were implemented instead, the expected number of cumulative confirmed cases would have been nearly 2 times or 5 times of the actual number. The effects of stringent interventions are significant in mitigating the epidemic situation of COVID-19. The slower the interventions were implemented, the more severe the epidemic would have been, and the stronger the interventions would have been required.

---

<sup>1</sup>Ting Tian, Wenxiang Luo and Yukang Jiang contributed equally to this article

<sup>2</sup>Xueqin Wang: wangxq20@ustc.edu.cn.

**Keywords:** COVID-19, treatment effect, regression discontinuity, synthetic control method, SIHR model

## 1. INTRODUCTION

The novel coronavirus disease (COVID-19) as an acute infection has rapidly spread over 200 countries in the world since December 2019 ([World Health Organization, 2020](#)). There were 2,804,799 reported cumulative confirmed cases and 193,711 cumulative deaths around the world and 84,341 cumulative confirmed cases and 4,643 cumulative deaths from China until April 26 , 2020 ([National Health Commission of the People’s Republic of China, 2020b](#)). These numbers are rapidly arising daily. This coronavirus family includes the virus that caused to Severe Acute Respiratory Syndrome (SARS) in 2003 and another that caused to Middle East Respiratory Syndrome (MERS) in 2012 ([Zhu et al., 2020](#)). Early cases of COVID-19 suggested that it might not be as severe as SARS-CoV and MERS-CoV, but the rapid increase in confirmed cases and the evidence of human-to-human transmission indicated that this coronavirus was more contagious than both SARS-CoV and MERS-CoV ([Wang et al., 2020](#); [Guan et al., 2020](#); [Chan et al., 2020](#)).

Compared with the outbreak of SARS, the Chinese government mounted responses rapidly to contain COVID-19 by reducing the risk of personal exposure and transmission ([Tu et al., 2020](#)). As the city with the largest number of confirmed cases in Zhejiang province ([National Health Commission of the People’s Republic of China, 2020a](#)), where it activated the first-level public health emergency response on January 23, 2020, Wenzhou had suspended all inter-provincial and inter-city shuttle buses, and chartered buses, and carried out temperature testing for passengers of buses arriving in Wenzhou as its own policies to deal with the epidemic situation since January 27, 2020 ([Wenzhou Municipal People’s Government, 2020a](#)). These strict control measures were implemented across Wenzhou, where one family member was assigned to go out for necessity purchase of each household every two days since February 1, 2020 ([Wenzhou Municipal People’s Government, 2020b](#)). Thus, Wenzhou had adopted the stringent intervention policies, including both traffic blockade inside and outside the city, and the strictly closed management of the community, to intensively prevent and control of the epidemic situation of COVID-19. In addition, an industrial city of Guangdong province, which activated the first-level public health emergency response on

January 23, 2020 ([Health Commission of Guangdong Province, 2020](#)), Shenzhen adopted closely following the close contact of confirmed cases and isolating them on the same time. Comparatively, Shenzhen implemented relatively mild but early interventions.

This paper concentrated on the strict intervention policies of COVID-19 that started on February 1, 2020 in Wenzhou and evaluated the treatment effects of those strict interventions, providing insights into the effects of the strict interventions. We also assessed what could have happened in Wenzhou if the stringent interventions were delayed by 0 to 5 days or if mild interventions were implemented instead. We accomplished this through simulation by using the time-varying Susceptible-Infected and infectious without isolation-Hospitalized in isolation-Removed (SIHR) model ([Tan et al., 2020](#)). Finally, we simulated the likely outcomes if relatively mild interventions as used in Shenzhen, China, were implemented in lieu of the stringent interventions.

In the next section, we describe the classical causal effects methods and dynamic transmission models of COVID-19 in detail. The corresponding analysis of epidemic data in Wenzhou are illustrated in Section 3. Section 4 discusses the effects of stringent interventions of COVID-19.

## 2. METHODOLOGIES

### 2.1 Data collection

We compiled a total of 504 confirmed cases ([Health Commission of Wenzhou, 2000](#)) from January 21, 2020 when the confirmed cases were first reported, to February 25, 2020 when no newly reported confirmed cases had been reported for a week, from the official website of the health commission of Wenzhou, and obtained the residential population of Wenzhou from the 2019 statistical yearbook of the Zhejiang Provincial Bureau of Statistics ([Statistics Bureau of Zhejiang Province, 2019](#)). The total confirmed cases of Shenzhen during the same period were collected. The cases per 100,000 individuals were calculated by the prevalence of COVID-19 multiplying 100,000 individuals in Wenzhou. We searched the daily number of cumulative confirmed cases in each city of China using the R package "nCov2019" ([Wu](#)

et al., 2020), and collected the number of cumulative confirmed cases from the top 100 cities for the epidemic of COVID-19 (except Hubei province) on February 25, 2020. The corresponding population density and GDP per capita statistics of these 100 cities were obtained from their latest statistical yearbooks.

## 2.2 Synthetic Control Method

To identify the treatment effects of the highly stringent interventions implemented on February 1, 2020, we compared Wenzhou with cities without those interventions. The synthetic control method was used to construct a data-driven control-group in the absence of those interventions in Wenzhou (Abadie and Gardeazabal, 2003; Abadie et al., 2010). The observed cases per 100,000 for Wenzhou (active intervention city) at the time  $t$  with interventions were denoted by  $Y_{1t}^1$ , and cases per 100,000 for cities  $j$  at the time  $t$  without interventions by  $Y_{jt}^0$ . The difference of  $Y_{1t}^1 - Y_{1t}^0$  was the treatment effect of the interventions on the cases per 100,000 for the active intervention city in the post-interventions period. Note that  $Y_{1t}^0$  is counterfactual for the active intervention city in the post-interventions period. Reliable estimates of cases per 100,000 for a synthetic intervention city were used to derive  $Y_{1t}^1 - Y_{1t}^0$ , where a synthetic intervention city resembled the active intervention city by possible pre-interventions characteristics. We selected the GDP per-capita (Zhang et al., 2020), population density (Oto-Peralías, 2020), and three-day cases per 100,000 before the implementation of interventions to construct a resembling city of Wenzhou. The effects of unobserved factors varied over time and were controlled by the linear combination of cases per 100,000 of pre-interventions (Abadie et al., 2010). Thus, the treatment effect of interventions in Wenzhou was formulated as (Abadie et al., 2011):

$$Y_{1t}^1 - Y_{1t}^0 = Y_{1t}^1 - \sum_2^{J+1} w_j^* Y_{jt}^0. \quad (2.1)$$

To determine  $J$ , the number of cities demographically similar to Wenzhou, we selected the top 100 cities of cumulative confirmed cases on February 25, 2020 in China (except Hubei province) and used hierarchical clustering (Ward Jr, 1963) to group those cities based on population density and GDP per capita. A homogenous group including Wenzhou and

other  $J$  cities were grouped together. Thus, those  $J$  cities could be considered as highly resembling to Wenzhou. The sum of weights for each of  $J$  cities was 1 and the values of weights in Equation (2.1) were determined by pre-interventions characteristics. The placebo tests were used to identify the magnitude of the treatment effect (i.e. the causal effect of interventions) (Abadie et al., 2010).

### 2.3 Regression Discontinuity

For the strict intervention policies of Wenzhou, we vertically compared the difference in the epidemic situation of COVID-19 before and after the implementation of such intervention policies. Regression Discontinuity (RD) (Thistlethwaite and Campbell, 1960; Hahn et al., 2001; Imbens and Lemieux, 2008) was used to evaluate whether the implementation of strict intervention policies of Wenzhou on February 1, 2020 had a significant effect on its own COVID-19 epidemic. The outcome of these stringent interventions was formulated as:

$$\log y_t = \alpha_0 + \alpha_1 h_t + \alpha_2(t - c) + \alpha_3 h_t(t - c)^k + \varepsilon_t \quad (2.2)$$

where the logarithmic transformation of the daily number of cumulative confirmed cases as a response variable. The values of  $t$  include 1 to 36, representing the number of days starting from January 21, 2020 as a rate variable in Equation (2.2) of RD, for example, the date of February 1 was denoted as 12 of  $t$ .  $h$  is an indicator variable for the implementation of the strict intervention policies in Wenzhou, given a value of 1 as the implementation of the policies but a value of 0 as no implementation of the policies. This indicates that the values are 0 and 1 before and after February 1, 2020 (including February 1 itself), respectively and the subscript of  $t$  indicates the number of days as defined before.  $c$  is the number of days corresponding to the cutoff-point whose value is 12, representing the policy implemented on February 1, 2020. The values of  $k$  could be denoted as the  $k^{th}$  order of interaction effect between rate variable and policy. We compared the  $k^{th}$  order of interaction effect item in the Equation (2.2) by AIC.

In RD, to determine whether  $c$  is an obvious jump point, the plot of both the rate variable and response variable was used to identify their relationship (Calonico et al., 2015a).

Different numbers of bins could be used to divide both the left-hand side and right-hand side of the cutoff-point to small intervals. Thus, the scatter plot of both median values of the rate variable and mean values of the response variable could be drawn (Calonico et al., 2015b). The actual rate variables are weighted uniformly in each bin, and the fitted curves of both the left-hand side and right-hand side are used to examine whether the cutoff-point can be considered as a threshold.

#### 2.4 Susceptible-Infected and infectious without isolation-Hospitalized in isolation-Removed (SIHR) model

The dynamic system of SIHR with four classes: Susceptible ( $S$ ), infected and infectious without isolation ( $I$ ), hospitalized in isolation ( $H$ ), removed ( $R$ ) (Li, 2018; Hsieh et al., 2004; Tan et al., 2020) was defined as:

$$\begin{aligned}
\frac{dS(t)}{dt} &= -\alpha \frac{I(t)}{N} S(t), \\
\frac{dI(t)}{dt} &= \alpha \frac{I(t)}{N} S(t) - \beta I(t), \\
\frac{dH(t)}{dt} &= \beta I(t) - \gamma H(t), \\
\frac{dR(t)}{dt} &= \gamma H(t), \\
N &= S(t) + I(t) + H(t) + R(t),
\end{aligned} \tag{2.3}$$

where the transmission rate  $\alpha$  is the average rate of being infected given contact over unit time,  $1/\beta$  is the mean of the incubation period,  $1/\gamma$  is the mean of the hospitalization period. To consider the effects of interventions, we introduced the time-varying transmission rate, which was defined as (Tan et al., 2020):

$$\alpha_{\alpha_0, d, m}(t) = \frac{\alpha_0}{1 + \exp(\lambda_m(t - d - m/2))}, \tag{2.4}$$

where  $\alpha_0$  denotes the maximum transmission rate of COVID-19 during the early outbreak,  $d$  represents the time when the interventions start to be effective and the transmission rate starts to decline,  $m$  is the duration of a process where the epidemic is to nearly vanish,  $\lambda_m$  is selected as  $(2 \log((1 - \varepsilon)/\varepsilon))/m$  and  $\varepsilon$  is specified to be 0.01. The smaller the values of  $d$

and  $m$ , the earlier effectiveness and the stronger intensity of interventions were implemented, respectively. We simulated the likely outcomes of delaying stringent interventions from 0 to 5 days by changing the values of  $d$  and of mild interventions on the same starting time of Wenzhou policies by changing the value of  $m$  to Shenzhen policies.

### 3. CASE STUDY

#### 3.1 Synthetic Control Method Analysis

We selected the top 100 cities outside Hubei province according to the cumulative confirmed cases on February 25, 2020 (**Table 1** in the Supplementary Materials). Based on the per capita GDP and population density of each city, these cities (including Wenzhou) were clustered into 4 groups using hierarchical clustering. Among them, a homogenous group consisted of Wenzhou and other 45 cities (**Figure 1** in the Supplementary Materials), where Taizhou of Zhejiang province implemented similar policies to Wenzhou and was excluded from our analysis. The remaining 44 cities formed a counterfactual city resembling Wenzhou (**Table 2** in the Supplementary Materials).

Before the strict interventions were implemented on February 1, 2020, the trends of actual cases per 100,000 in both Wenzhou and "synthetic Wenzhou" were highly similar, suggesting that this synthetic city can be used to estimate the "counterfactual" results of Wenzhou. After 2 days of the implementation of the policies, the growth rate of actual cases per 100,000 in Wenzhou remarkably slower than that of "synthetic Wenzhou". The number of cases per 100,000 in "Synthetic Wenzhou" on February 25, 2020 was 10.32, which is 1.69 times the actual Wenzhou (6.08) (**Figure 1**). In other words, Wenzhou had not implemented the strict interventions on February 1, 2020, the number of cumulative confirmed cases would have reached 954 on February 25, 2020, i.e., the epidemic of COVID-19 in Wenzhou would have expanded to approximately 1.7 times.






Figure 1: **The trends of COVID-19 cases per 100,000 between Wenzhou and Synthetic Wenzhou from January 21 to February 25, 2020.** The black dashed line indicated the time of implementation of the intervention policies in Wenzhou. The grey dashed line indicated 2 days delay of implementation of the intervention policies in Wenzhou.

A placebo test was performed to determine the significance level of the difference in the trends of COVID-19 cases per 100,000. To this end, we plotted the gap curves between Wenzhou and synthetic Wenzhou by in turn exchanging Wenzhou and one of each 44 cities in the homogenous group of Wenzhou. The gap of COVID-19 cases per 100,000 between Wenzhou and our synthetic Wenzhou was the largest compared to the rest, i.e. the negative

effect of the intervention policies on COVID-19 per 100,000 in Wenzhou was the lowest of all. For those 44 cities, the probability of having a gap for Wenzhou under a random permutation of the control measures was 5%, which is conventionally regarded as statistically significant. This suggested that the effect of the implementation of the policies in Wenzhou was significantly different from the implementation of the policies in the remaining 44 cities, indicating that the strict interventions of Wenzhou might have significantly reduced the COVID-19 cases per 100,000 (**Figure 2**).




Figure 2: **The permutation test of the treatment effect of the implementation of the policies in Wenzhou and other 44 control regions.** All grey curves represent the placebo tests of COVID-19 cases per 100,000 gaps between a control city (an arbitrary random city of 44 cities) and synthetic control city (a combination of remaining 43 cities and Wenzhou), and the blue curve represents the placebo test of COVID-19 cases per 100,000 gap between Wenzhou and synthetic Wenzhou.

### 3.2 Regression Discontinuity Analysis

The values of AIC for each regression discontinuity model for different  $k^{th}$  order of interaction effects between rate variable and policy were shown in **Table 1**.

Table 1: The values of AIC for each model with different  $k^{th}$  order of interaction effects

Models	$df$	AIC
No interaction	4	97.344
First order	5	-28.861
$Time^2$ without interaction	5	44.057
Second order	6	-40.846
$Time^3$ without interaction	6	-34.639
Third order	7	-42.019

Looking at **Table 1**, the model included item  $Policy*Time^3$  ( $h_t(t - c)^3$ ) had the lowest value of AIC but it had more parameters ( $df = 7$ ). Also, the slopes of the fitted curves between the left-hand side and the right-hand side of the cutoff-point were different (**Figure 3**). The points lied on the right-hand side of quadratic curve. Thus, RD could be used to examine the treatment effect of interventions implemented in Wenzhou and the model included the second order of interaction effects between policy and rate variable was reasonable. The coefficient of interventions is -0.350 (p-value: 0.003) and the interaction effect between interventions and time is -0.438 (p-value: <0.001) indicating that there was a significant treatment effect of highly stringent interventions implemented in Wenzhou on February 1, 2020 (**Table 2**).

Table 2: The coefficients of variables in the RD and their corresponding p-values

Variables	Coefficients	p-values
Intercept	5.958	<0.001
<i>Policy</i> ( $h_t$ )	-0.350	0.003
<i>Time</i> ( $(t - c)$ )	0.511	<0.001
<i>Policy*Time</i> ( $h_t(t - c)$ )	-0.438	<0.001
<i>Policy*Time</i> <sup>2</sup> ( $h_t(t - c)^2$ )	-0.002	0.001
Adjusted $R^2$	0.994	




Figure 3: **The plot of both rate variable and response variable.** The green fitted curves of relationship between median of time within each bin and mean of log(cumulative confirmed cases) within each bin, the black dashed vertical line represents the time of cutoff-point (with value 12)

### 3.3 Simulation of stringent interventions delay or mild interventions instead

Our simulation projected that the expected cumulative confirmed cases would be 1.84 times of the actual cases for a 2-day delay, 2.45 times for a 3-day delay, 3.26 times for a 4-day delay and 4.30 times for a 5-day delay on February 25, 2020. The corresponding 95% credible interval (CI) for these projected numbers of cases were presented in **Table 3**. The

full simulation results from January 21 to February 25, 2020 are presented in **Figure 2** of the Supplementary Materials. According to **Table 3**, the expected cumulative confirmed cases for 0-day delay (i.e. the predicted cumulative confirmed cases) was very close to the actual cumulative confirmed cases of Wenzhou. The expected cumulative confirmed cases were 925 with corresponding 95% CI (571,1547) for the 2-day delay, 1233 with corresponding 95% CI (702,2233) for the 3-day delay, 1644 with corresponding 95% CI (851,3171) for the 4-day delay, and 2167 with corresponding 95% CI (1034, 4578) for the 5-day delay. The observed cumulative confirmed cases had been stable since February 17, 2020, however, the expected cumulative confirmed continued to increase if the stringent interventions were delayed. Based on **Figure 1**, if the stringent interventions were delayed by 2 days or more, the epidemic of COVID-19 in Wenzhou could have been more severe than that in synthetic Wenzhou (954 cumulative confirmed cases on February 25, 2020). If the mild interventions as those implemented in Shenzhen were implemented in Wenzhou on February 1, 2020, the expected number of cumulative confirmed cases would have been 2319 with corresponding 95% CI (1145,5189) on February 25, 2020 (**Figure 4**).

Table 3: The likely outcome of intervention implemented delay in different days including the expected cumulative confirmed cases with corresponding 95% CI from February 16 to February 25, 2020.

Date	Actual	0 day	1 day	2 days	3 days	4 days	5 days
02/16	503	514(361,749)	678(453,1059)	892(553,1490)	1178(674,2128)	1556 (810,2998)	2034 (978,4264)
02/17	504	517 (363,754)	683 (456,1066)	899 (557,1502)	1190 (680,2151)	1576 (819,3036)	2064 (990,4330)
02/18	504	519 (364,757)	686 (458,1072)	905 (560,1513)	1200 (685,2170)	1591 (826,3067)	2087 (1000,4387)
02/19	504	521 (365,760)	689 (460,1077)	910 (563,1521)	1209 (690,2185)	1604 (832,3092)	2106 (1008,4435)
02/20	504	522 (366,763)	692 (462,1081)	914 (565,1528)	1215 (693,2198)	1615 (837,3112)	2122 (1015,4472)
02/21	504	524 (367,764)	694 (463,1084)	917 (567,1533)	1220 (696,2208)	1624 (841,3129)	2134 (1021,4502)
02/22	504	525 (368,766)	695 (464,1087)	919 (568,1538)	1224 (698,2217)	1630 (845,3143)	2145 (1025,4527)
02/23	504	525 (368,767)	697(465,1089)	922 (570,1542)	1228 (700,2224)	1636 (847,3154)	2153 (1029,4548)
02/24	504	526 (369,769)	698 (465,1091)	923 (570,1545)	1230 (701,2229)	1640 (850,3164)	2161 (1032,4564)
02/25	504	526 (369,769)	698 (466,1092)	925 (571,1547)	1233 (702,2233)	1644 (851,3171)	2167 (1034,4578)






Figure 4: **The expected number of cumulative confirmed cases for mild interventions implemented on February 1, 2020 in Wenzhou.** The points represent the actual cumulative confirmed cases of Wenzhou.

#### 4. DISCUSSION

We used daily reported cumulative confirmed case data from January 21 to February 25, 2020 in Wenzhou. By using the synthetic control method, the trend of COVID-19 cases per 100,000 for the synthetic Wenzhou as a control group of no intervention were compared with those for Wenzhou as a treatment group with interventions. The COVID-19 cases per

100,000 in Wenzhou were significantly lower than those in synthetic Wenzhou after February 3, 2020. This indicated that the implementation of the strict interventions on February 1, 2020 in Wenzhou had a significant effect on controlling the epidemic of COVID-19.

By using regression discontinuity analysis, we also concluded that the implementation of strict intervention policies had a significant treatment effect on the epidemic in Wenzhou. Moreover, the statistical treatment effects were evaluated. That is, since the intervention policies were implemented in Wenzhou (corresponding coefficient of policy : -0.350), the growth rate of the reported cumulative confirmed cases of COVID-19 were reduced, and this "reduction" effect would increase over time (corresponding coefficient of interaction between time and policy: -0.438). Conversely, if the policies were not implemented, the reported cumulative confirmed cases of COVID-19 would increase over time (corresponding coefficient of time: 0.511).

The horizontal and longitudinal comparisons were made to examine the treatment effects of the strict intervention policies implemented in Wenzhou, including the suspension of public transportation in the city, the closure of highway junctions, and strict community access control. Since January 21, 2020, compared with other cities outside Hubei province, the number of reported cumulative confirmed cases in Wenzhou had been consistently in the top two places, i.e., its COVID-19 epidemic situation was relatively severe. If strict intervention policies were not implemented, the outbreak of COVID-19 would be expected to expand to 1.7 times. Therefore, it can be concluded from the results of the two methods that the strict intervention policies in Wenzhou, where the epidemic was severe, had significantly suppressed its epidemic situation.

Based on our simulation, if the stringent interventions were delayed by a few days, such as 2 days, the epidemic situation would have been remarkably more severe, where the expected number of cumulative confirmed cases would have been nearly 2 times of the actual number of cases on February 25, 2020. If the mild interventions were implemented in lieu of the stringent interventions, the expected number of cumulative confirmed cases would have been 4.60 times of the actual number of cases on February 25, 2020. Overall, if the stringent

interventions were delayed or the mild interventions were implemented instead, though on the same day, the expected cumulative confirmed cases would have continued to increase while the actual epidemic situation was under control in the Wenzhou.

SUPPLEMENTARY MATERIALS

Table 1: Top 100 cities of cumulative confirmed cases until 25 February (except Hubei province), and their corresponding population density and GDP per-capita

Province	City	Cumulative confirmed cases	Population density	GDP per-capita (RMB)
Chongqing	Chongqing	576	376.43083	65933
Zhejiang	Wenzhou	504	763.832	65055
Guangdong	Shenzhen	417	6484	189568
Beijing	Beijing	400	7794.8	192957.1
Guangdong	Guangzhou	346	2005	155491
Shanghai	Shanghai	336	3814	126634
Henan	Xinyang	274	342.0925	36951.12
Shandong	Jining	258	745.7689	58972
Hunan	Changsha	242	689.9653	136920
Jiangxi	Nanchang	229	749.1894	95116.22
Heilongjiang	Harbin	198	204.4821	48345.92
Anhui	Hefei	174	706.5906	116352.2
Zhejiang	Hangzhou	169	518.958	140180
Anhui	Bengbu	160	640.6788	50662
Henan	Zhengzhou	157	1361.268	101352.1
Zhejiang	Ningbo	157	835.575	132603
Hunan	Yueyang	156	385.9844	59165
Anhui	Fuyang	155	1058.312	21700
Henan	Nanyang	155	376.4511	35554.64
Zhejiang	Taizhou	146	610.846	79541
Sichuan	Chengdu	143	1121.066	86911
Henan	Zhumadian	139	466.5252	33773.42
Tianjin	Tianjin	135	2596.509	118944
Jiangxi	Xinyu	130	373.411	86791
Jiangxi	Shangrao	123	298.7281	36899
Shaanxi	Xi'an	120	989.6814	86000
Jiangxi	Jiujiang	118	260.1498	55141.93
Anhui	Bozhou	108	625.3881	24388.01
Jiangxi	Yichun	106	297.4825	39268.42
Hunan	Shaoyang	102	353.9426	24178
Guangdong	Zhuhai	98	1111.758	159428
Guangdong	Dongguan	97	3404.544	98939
Jiangsu	Nanjing	93	1280.735	152886
Henan	Shangqiu	91	684.3516	32669.96
Jiangsu	Suzhou	87	1238.501	173765
Guangdong	Foshan	84	2043.873	127691
Hong Kong	Hong Kong	84	6733.915	336147

Continued on next page

Table 1 – continued from previous page

Province	City	Cumulative confirmed cases	Population density	GDP per-capita (RMB)
Anhui	Anqing	83	388.8447	37243.79
Hunan	Changde	82	320.3518	58160
Hunan	Zhuzhou	80	357.0236	65442
Jiangsu	Xuzhou	79	748.1513	76915
Jiangxi	Ganzhou	76	219.4027	40212
Henan	Zhoukou	76	725.6292	30820.63
Hunan	Loudi	76	484.3908	39249
Jiangxi	Fuzhou	72	215.0821	34156.95
Fujian	Fuzhou	71	645	102037
Sichuan	Garz	69	7.772442	22097
Anhui	Lu'an	69	338.3381	24638.49
Guangdong	Zhongshan	66	1838.889	110585
Jiangsu	Huai'an	66	491.0269	73204
Guangdong	Huizhou	62	431.25	85418
Shandong	Qingdao	60	674.295	128459
Hunan	Yiyang	59	363.4552	39937
Hebei	Tangshan	58	586.1787	82692
Henan	Pingdingshan	58	797.921	42586
Henan	Xinxiang	57	700.7014	43696.44
Guangxi	Nanning	55	328.0617	59259
Jiangsu	Wuxi	55	1420.899	174270
Zhejiang	Jinhua	55	512.155	73428
Fujian	Putian	55	685.5792	77325
Hainan	Sanya	54	393.8903	63046.1
Yunnan	Kunming	53	326.1905	76387
Henan	Anyang	53	924.4508	46443.24
Heilongjiang	Shuangyashan	52	62.64733	35527
Jiangsu	Changzhou	51	978.6368	149275
Shandong	Linyi	49	617.9618	44534
Jiangsu	Lianyungang	48	593.5653	61332
Hebei	Cangzhou	48	563.0002	48226
Hunan	Hengyang	48	473.1156	42163
Shandong	Jinan	47	912.3639	106302
Shandong	Yantai	47	521.5907	110231
Fujian	Quanzhou	47	790.1907	97614
Heilongjiang	Suihua	47	149.0159	29625.32
Heilongjiang	Jixi	46	74.9566	27639.35
Jilin	Changchun	45	373.3042	86465
Zhejiang	Jiaxing	45	1119.11	103858
Shandong	Weifang	44	580.8031	65721

Continued on next page

Table 1 – continued from previous page

Province	City	Cumulative confirmed cases	Population density	GDP per-capita (RMB)
Guangxi	Beihai	44	525.6818	73074
Hunan	Yongzhou	43	242.9526	33035
Heilongjiang	Qiqihar	43	119.6062	23676.17
Sichuan	Dazhou	42	342.6378	28066
Zhejiang	Shaoxing	42	608.1652	107853
Anhui	Suzhou	41	580.5048	28693.98
Jiangsu	Nantong	40	692.9567	115320
Hunan	Huaihua	40	180.6035	30449
Sichuan	Nanchong	39	512.8576	28516
Henan	Xuchang	39	888.1906	63987.61
Hainan	Haikou	39	973.3623	56055.67
Hunan	Chenzhou	39	244.7258	50482
Anhui	Ma'anshan	38	577.1796	82075
Shandong	Weihai	38	488.1835	128774
Shandong	Liaocheng	38	696.5371	51935
Shandong	Dezhou	37	560.7567	58252
Shanxi	Jinzhong	37	206.2	42916
Jiangsu	Taizhou	37	801.0181	110180
Hunan	Xiangtan	36	572.2733	75609
Gansu	Lanzhou	36	284.3636	73042
Guizhou	Guiyang	36	606.9471	78449
Shandong	Tai'an	35	726.7105	64714
Henan	Luohe	35	1020.252	46318

Table 2: The pool of cities in the group of Wenzhou and their corresponding weights

Province	City	Population density	GDP per-capita (RMB)	Weight
Chongqing	Chongqing	376.43083	65933	0
Shandong	Jining	745.7689	58972	0
Jiangxi	Nanchang	749.1894	95116.22	0
Anhui	Hefei	706.5906	116352.2	0
Henan	Zhengzhou	1361.268	101352.1	0
Hunan	Yueyang	385.9844	59165	0
Sichuan	Chengdu	1121.066	86911	0
Tianjin	Tianjin	2596.509	118944	0
Jiangxi	Xinyu	373.411	86791	0.69
Shaanxi	Xi'an	989.6814	86000	0
Jiangxi	Jiujiang	260.1498	55141.93	0
Guangdong	Dongguan	3404.544	98939	0
Hunan	Changde	320.3518	58160	0
Hunan	Zhuzhou	357.0236	65442	0
Jiangsu	Xuzhou	748.1513	76915	0
Fujian	Fuzhou	645	102037	0
Guangdong	Zhongshan	1838.889	110585	0
Jiangsu	Huai'an	491.0269	73204	0
Guangdong	Huizhou	431.25	85418	0
Hebei	Tangshan	586.1787	82692	0
Guangxi	Nanning	328.0617	59259	0
Zhejiang	Jinhua	512.155	73428	0
Fujian	Putian	685.5792	77325	0
Hainan	Sanya	393.8903	63046.1	0.31
Yunnan	Kunming	326.1905	76387	0
Jiangsu	Lianyungang	593.5653	61332	0
Shandong	Jinan	912.3639	106302	0
Shandong	Yantai	521.5907	110231	0
Fujian	Quanzhou	790.1907	97614	0
Jilin	Changchun	373.3042	86465	0
Zhejiang	Jiaxing	1119.11	103858	0
Shandong	Weifang	580.8031	65721	0
Guangxi	Beihai	525.6818	73074	0
Zhejiang	Shaoxing	608.1652	107853	0
Jiangsu	Nantong	692.9567	115320	0
Henan	Xuchang	888.1906	63987.61	0
Hainan	Haikou	973.3623	56055.67	0
Anhui	Ma'anshan	577.1796	82075	0
Shandong	Dezhou	560.7567	58252	0
Jiangsu	Taizhou	801.0181	110180	0
Hunan	Xiangtan	572.2733	75609	0

Continued on next page

**Table 2 – continued from previous page**

<b>Province</b>	<b>City</b>	<b>Population density</b>	<b>GDP per-capita (RMB)</b>	<b>Weight</b>
Gansu	Lanzhou	284.3636	73042	0
Guizhou	Guiyang	606.9471	78449	0
Shandong	Tai'an	726.7105	64714	0

---






Figure 1: A dendrogram tree of clustering results




Figure 2: The expected number of cumulative confirmed cases for stringent interventions if they were delayed by different days. The points represent the actual cumulative confirmed cases of Wenzhou.

## REFERENCES

- Abadie, A., Diamond, A., and Hainmueller, J. (2010), “Synthetic control methods for comparative case studies: Estimating the effect of Californias tobacco control program,” *Journal of the American statistical Association*, 105(490), 493–505.
- Abadie, A., Diamond, A., and Hainmueller, J. (2011), “Synth: An r package for synthetic control methods in comparative case studies,” *Journal of Statistical Software*, 42(13).
- Abadie, A., and Gardeazabal, J. (2003), “The economic costs of conflict: A case study of the Basque Country,” *American economic review*, 93(1), 113–132.
- Calonico, S., Cattaneo, M. D., and Titiunik, R. (2015a), “Optimal data-driven regression discontinuity plots,” *Journal of the American Statistical Association*, 110(512), 1753–1769.
- Calonico, S., Cattaneo, M. D., and Titiunik, R. (2015b), “rdrobust: An r package for robust nonparametric inference in regression-discontinuity designs,” *R Journal*, 7(1), 38–51.
- Chan, J. F. W., Yuan, S., Kok, K. H., To, K. K. W., Chu, H., Yang, J., Xing, F., Liu, J., Yip, C. C.-Y., Poon, R. W.-S. et al. (2020), “A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster,” *The Lancet*, 395(10223), 514–523.
- Guan, W., Ni, Z., Hu, Y., Liang, W., Ou, C., He, J., Liu, L., Shan, H., Lei, C., Hui, D. S. et al. (2020), “Clinical characteristics of coronavirus disease 2019 in China,” *New England journal of medicine*, 382(18), 1708–1720.
- Hahn, J., Todd, P., and Van der Klaauw, W. (2001), “Identification and estimation of treatment effects with a regression-discontinuity design,” *Econometrica*, 69(1), 201–209.
- Health Commission of Guangdong Province (2020), “First-level public health emergency response . [accessed 2020 March 31].”  
**URL:** [http://wsjkw.gd.gov.cn/zwyw\\_yqxx/content/post\\_2878895.html](http://wsjkw.gd.gov.cn/zwyw_yqxx/content/post_2878895.html)
- Health Commission of Wenzhou (2000), “Outbreak of COVID-19 in Wenzhou.”

- Hsieh, Y.-H., Lee, J.-Y., and Chang, H.-L. (2004), “SARS epidemiology modeling,” *Emerging infectious diseases*, 10(6), 1165.
- Imbens, G. W., and Lemieux, T. (2008), “Regression discontinuity designs: A guide to practice,” *Journal of econometrics*, 142(2), 615–635.
- Li, M. Y. (2018), *An introduction to mathematical modeling of infectious diseases*, Vol. 2 Springer.
- National Health Commission of the People’s Republic of China (2020a), “Distribution of COVID-19 cases in China. [accessed 2020 April 26],”  
**URL:** <http://2019ncov.chinacdc.cn/2019-nCoV/index.html>
- National Health Commission of the People’s Republic of China (2020b), “Distribution of COVID-19 cases in the world. [accessed 2020 April 26],”  
**URL:** <http://2019ncov.chinacdc.cn/2019-nCoV/global.html>
- Oto-Peralías, D. (2020), “Regional correlations of COVID-19 in Spain,” , .
- Statistics Bureau of Zhejiang Province (2019), *Zhejiang Statistical Yearbook 2019*.
- Tan, J., Jiang, Y., Tian, T., and Wang, X. (2020), “P-SIHR Probabilistic Graphical Model: an Applicable Model of COVID-19 in Estimating the Number of Infectious Individuals,” *Acta Mathematicae Applicatae Sinica*, 43(2).
- Thistlethwaite, D. L., and Campbell, D. T. (1960), “Regression-discontinuity analysis: An alternative to the ex post facto experiment.,” *Journal of Educational psychology*, 51(6), 309.
- Tu, W., Tang, H., Chen, F. et al. (2020), “Epidemic Update and Risk Assessment of 2019 Novel Coronavirus? China, January 28, 2020,” *China CDC Weekly*, 2(6), 83–86.
- Wang, C., Horby, P. W., Hayden, F. G., and Gao, G. F. (2020), “A novel coronavirus outbreak of global health concern,” *The Lancet*, 395(10223), 470–473.
- Ward Jr, J. H. (1963), “Hierarchical grouping to optimize an objective function,” *Journal of the American statistical association*, 58(301), 236–244.

Wenzhou Municipal People’s Government (2020a), “Notice of the Leading Group for the Prevention and Control of COVID-19 in Wenzhou (No. 3) . [accessed 2020 March 31],”.

**URL:** [http://www.wenzhou.gov.cn/art/2020/1/28/art\\_1219304\\_41858870.html](http://www.wenzhou.gov.cn/art/2020/1/28/art_1219304_41858870.html)

Wenzhou Municipal People’s Government (2020b), “Notice of the Leading Group for the Prevention and Control of COVID-19 in Wenzhou (No. 6) . [accessed 2020 March 31],”.

**URL:** [http://www.wenzhou.gov.cn/art/2020/2/2/art\\_1219304\\_41867466.html](http://www.wenzhou.gov.cn/art/2020/2/2/art_1219304_41867466.html)

World Health Organization (2020), “Rolling updates on coronavirus disease (COVID-19). [accessed 2020 May 14],”.

**URL:** <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen>

Wu, T., Ge, X., Yu, G., and Hu, E. (2020), “Open-source analytics tools for studying the COVID-19 coronavirus outbreak,” *medRxiv*, .

Zhang, Y., Tian, H., Zhang, Y., and Chen, Y. (2020), “Is the epidemic spread related to GDP? Visualizing the distribution of COVID-19 in Chinese Mainland,” *arXiv preprint arXiv:2004.04387*, .

Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R. et al. (2020), “A novel coronavirus from patients with pneumonia in China, 2019,” *New England Journal of Medicine*, .