

A note on a conjecture of star chromatic index for outerplanar graphs*

Xingchao Deng^a, Qingye Yao^b, Yanbing Zhang^c and Xudong Cui^d

^{a b c d}College of Mathematical Science, Tianjin Normal University

Tianjin City, 300071, P. R. China

Abstract

A *star edge coloring* of a graph G is a proper edge coloring of G without bichromatic paths or cycles of length four. The *star chromatic index*, $\chi_{st}(G)$, of G is the minimum number k for which G has a star edge coloring by k colors. In [2], L. Bezegová et al. conjectured that $\chi'_{st}(G) \leq \lfloor \frac{3\Delta}{2} \rfloor + 1$ when G is an outerplanar graph with maximum degree $\Delta \geq 3$. In this paper we obtained that $\chi'_{st}(G) \leq \Delta + 6$ when G is an 2-connected outerplanar graph with diameter 2 or 3. If G is an 2-connected outerplanar graph with maximum degree 5, then $\chi_{st}(G) \leq 9$.

Keywords: star chromatic index; diameter; outerplanar graph; maximal outerplanar graph

1 Introduction

All graphs in this paper are simple and undirected. For a graph G , we use $V(G)$ and $E(G)$ to denote the vertex and edge set, $|V(G)|$ and $|E(G)|$ to denote the number of vertices and edges respectively. The *star coloring* of a graph is a proper coloring and the vertices of any two color classes of the star colored graph induce a star forest. In 1973, Grünbaum initiated the star coloring in [5]. A proper edge coloring of a graph G , which has no 2-edge colored 4-length paths or cycles, is a *star edge coloring* of G . The minimum number of colors needed for a star edge coloring of G is called the *star chromatic index* of it, denoted by $\chi'_{st}(G)$. In 2008, Xincheng Liu and Kai Deng introduced the star edge coloring in [7]. Obviously, a star edge coloring of a graph G is a star coloring of the line graph of G . A *strong edge coloring* of a graph G , denoted by $\chi'_s(G)$, is a proper edge coloring so that no edge can be adjacent to two edges with the same color in which every color class gives an induced matching. The definition was introduced by Fouquet and Jolivet (1983) to solve a problem involving radio networks and their frequencies. It is easy to know that the strong edge coloring of a graph G is a star edge coloring of it, then $\chi'_{st}(G) \leq \chi'_s(G)$. The diameter of a graph G , denoted by $diam(G)$, is the maximum distance of any two vertices in the graph G . We use the symbol $G_1 \vee G_2$ to denote the join graph of G_1, G_2 . A path with n vertices is denoted by P_n . A graph with n vertices is denoted by G_n .

If a graph G can be drawn in the plane so that its edges intersect only at their ends, then G is called a planar graph. In particular, when all vertices of a planar graph G lie on a same face, then G is an outerplanar graph. A minor of a graph G is obtained by deleting some vertices, edges or contracting

*This work is supported by the Science & Technology development Fund of Tianjin Education Commission for Higher Education, China (No. 2019KJ090).

^aCorresponding author. E-mail: dengyuqiu1980@126.com , xcdeng@mail.tjnu.edu.cn

some edges of the graph G . For the graphs G and H such that no minor of G is isomorphic to H , then G is called a H minor free graph.

Recently, Dvořák and Mohar have studied the star edge coloring of complete graphs and 3-regular graphs. They obtained some results and proposed a conjecture shown as follows:

Theorem 1.1 [4] Let K_n be a complete graph with n vertices, then

$$2n(1 + o(1)) \leq \chi'_{st}(K_n) \leq n \frac{2^{2\sqrt{2}(1+o(1))\sqrt{\log n}}}{(\log n^{\frac{1}{4}})} O\left(\frac{\log \Delta}{\log \log \Delta}\right)^2.$$

Theorem 1.2 [4] If G is a graph with the maximum degree Δ , then

$$\chi'_{st}(G) \leq \chi'_{st}(K_{\Delta+1}) \cdot O\left(\frac{\log \Delta}{\log \log \Delta}\right)^2,$$

therefore,

$$\chi'_{st}(G) \leq \Delta \cdot 2^{O(1)\sqrt{\log \Delta}}.$$

Theorem 1.3 [4]

(a) When G is a subcubic graph, then $\chi'_{st}(G) \leq 7$.

(b) If G is a simple cubic graph, then $\chi'_{st}(G) \geq 4$, and the equality holds if and only if it covers the graph of the 3-cubic.

Conjecture 1.4 [4] Let G be a subcubic graph, then $\chi'_{st}(G) \leq 6$.

Bezegová et al. have given some results and a conjecture on outerplanar graph in [2] as follows:

Theorem 1.5 [2] If G is an outerplanar graph, then

(1) $\chi'_{st}(G) \leq \lfloor 1.5\Delta \rfloor + 12$.

(2) If $\Delta(G) \leq 3$, then $\chi'_{st}(G) \leq 5$.

Conjecture 1.6 [2] If G is an outerplanar graph with the maximum degree Δ , then $\chi'_{st}(G) \leq \lfloor 1.5\Delta \rfloor + 1$.

The following theorems 1.7-1.10 give some results on the star chromatic indexes of outerplanar graphs.

Theorem 1.7 [9] When G is a maximal outerplanar graph with maximum degree $\Delta = 4$, then $\chi'_{st}(G) = 6$.

Theorem 1.8 [6] Let G be a maximal outerplanar graph with n vertices, then $6 \leq \chi'_{st}(G) \leq n - 1$.

Theorem 1.9 [6] If G is a maximal outerplanar graph with maximum degree Δ and $n(n \geq 8)$ vertices, then $\chi'_{st}(F_{\Delta+1}) \leq \chi'_{st}(G) \leq n$.

Y. Q. Wang et al. obtained the following result by edge partition method.

Theorem 1.10 [17] If G is an outerplanar graph with the maximum degree Δ , then $\chi'_{st}(G) \leq \lfloor 1.5\Delta \rfloor + 5$.

Lemma 1.11 [11] A graph G is an outerplanar graph if and only if G is K_4 and $K_{2,3}$ minor free.

In [1], one has proved that almost all graphs have diameter 2, so it is important to discuss the star edge coloring of outerplanar graphs with diameters 2 or 3.

For the sake of narrative, let $\zeta_n^d = \{G_n | G_n \text{ is an } 2\text{-connected outerplanar graph with diameter } d \text{ and } n \text{ vertices}\}$. In this paper, we give some upper bound based on maximum degree of graphs $G_n (\in \zeta_n^2 \cup \zeta_n^3)$.

2 $\chi'_{st}(G_n (\in \zeta_n^2))$

First of all, we recall the star chromatic indexes of three basic graphs [9].

Lemma 2.1 [9] For the path P_n with $(n \geq 5)$ vertices, we have $\chi'_{st}(P_n) = 3$.

Lemma 2.2 [9] Let C_n be a cycle with n vertices, then

$$\chi'_{st}(C_n) = \begin{cases} 4, & n = 5, \\ 3, & \text{else.} \end{cases} \quad (1)$$

Lemma 2.3 [9] F_{n+1} is a Fan graph with $n + 1$ vertices, i.e. $F_{n+1} = K_1 \vee P_n$, then

$$\chi'_{st}(F_{n+1}) = \begin{cases} n + 2, & n = 4, \\ n + 1, & n = 2, 3, 5, 6, \\ n, & n \geq 7. \end{cases} \quad (2)$$

Definition 2.4 When G is a planar graph, we say that G is a maximal planar graph if G is a nonplanar graph when adding an edge for any two nonadjacent vertices.

Definition 2.5 Let G be an outerplanar graph. When adding an edge for any two nonadjacent vertices, the resulting graph is a nonouterplanar graph, then G is called a maximal outerplanar graph.

We know that any outerplanar graph can become a maximal outerplanar graph by adding edges from the definition of the maximal outerplanar graph, and the diameter of the resulting graph will not be increased. Thus, we only need to study the star chromatic index of maximal outerplanar graph with diameter 1 and diameter 2 in order to study the upper bound of the star chromatic index of 2-connected outerplanar graph with diameter 2. Since only K_3 has diameter 1 and the connected subgraph with diameter 2 of K_3 is P_3 , which has no star edge coloring. Therefore, we wouldn't consider the maximal outerplanar with diameter 1.

In [3], L. Beineke and R. Pippert gave a method to construct the maximal outerplanar graph shown as follows:

(a) K_3 is a maximal outerplanar graph;

(b) Let G_1 be a maximal outerplanar graph embedded in a plane with vertices lying on the exterior face F_1 and G_2 be the resulting graph by adding a new vertex which is adjacent to some two vertices of an edge on F_1 . Then G_2 is a maximal outerplanar graph;

(c) Any maximal outerplanar graph can be obtained by the processes (a)(b).

Theorem 2.6 Let $G_n \in \zeta_n^2$, then

$$\Delta(G_n) \leq \chi'_{st}(G_n) \leq \begin{cases} n+1, & n=5, \\ n, & n=4,6,7, \\ \Delta(G_n), & n \geq 8. \end{cases} \quad (3)$$

Proof: By the above construction method, we know that maximal outerplanar graphs G_n with diameter 2 have the following two cases: (1) $G_n = F_n (n \geq 4)$, (2) $G_n = G_6^1$.




Fig. 1. G_6^1

For the case (1), the Lemma 2.3 shows that the theorem is correct. In the case (2), F_5 is a vertex-induced subgraph of G_6^1 , then $\chi'_{st}(G_6^1) \geq \chi'_{st}(F_5) = 6$. It is easy to check that the coloring of G_6^1 shown in Fig. 1 is a star 6-edge coloring.

According to the definition of the star edge coloring, we have $\chi'_{st}(G_n) \geq \Delta$. For $n \geq 8$, we know that $\Delta(G_n) = n - 1$. \square

3 $\chi'_{st}(G_n (\in \zeta_n^3))$

By the definition of maximal outerplanar graphs and Lemma 1.11, we only need to study the star chromatic index of the maximal outerplanar graph with diameter 2 or 3 for studying the upper bound of the star chromatic index of G_n . Let $\xi_n^d = \{H_n | H_n \text{ are the 2-connected maximal outerplanar graph with diameter } d \text{ and } n \text{ vertices}\}$. Let n be an integer, for any G_n , there is some $H_n \in \xi_n^2 \cup \xi_n^3$ such that G_n is a subgraph of H_n . The following two cases are discussed according to the diameter of H_n .

3.1 $H_n \in \xi_n^2$

Theorem 3.1 Let $H_n \in \xi_n^2$ and G_n is an edge-include subgraph of H_n . When $H_n = F_n$, then we have

$$\Delta(G_n) \leq \chi'_{st}(G_n) \leq \Delta(G_n) + 3.$$

If $H_n = G_6^1$, we obtained that

$$\chi'_{st}(G_n) = 4. \quad (4)$$

Proof: In the following, we prove the theorem by two cases based on H_n with diameter 2.




Fig. 2. G_6^1

3.1.1 $H_n = G_6^1$

We delete any two edges of $\{v_0v_2, v_0v_3, v_2v_3\}$ in H_n , then the diameter of resulting graph is 3. In the sense of isomorphism, we only consider the graph G_6^1 shown in Fig. 2, which is obtained by deleting the edges v_0v_2, v_0v_3 in H_n . Since the edges $v_1v_2, v_2v_5, v_5v_3, v_3v_4$ must be colored by at least three different colors, and there are no bichromatic 4-length path in G_6^1 , moreover the color of v_2v_3 is different from the colors of $v_1v_2, v_2v_3, v_5v_3, v_3v_4$, thus we obtain that $\chi'_{st}(G_6^1) \geq 4$. It is easy to check that the coloring of Fig. 2 is a star edge coloring and only uses four colors, then $\chi'_{st}(G_6^1) = 4$.

3.1.2 $H_n = F_n$

Let $H_n = F_n$, then $n \geq 6$ (since $\text{diam}(C_5)=2$). We discuss according to the value of n as follows.

Case 1. If $n = 6$, the star edge coloring of H_6 is shown in Fig. 3. When we delete the edge(s) which is(are) not in the exterior face of H_6 , the resulting graph has diameter 3. Let the set of deleting edge(s) to be D . We can obtain that $\chi'_{st}(G_6) = \chi'_{st}(H_6) - |D| \leq 6 - |D| = \Delta(G_6) + 1$.

Case 2. When $n = 7$, the star edge coloring of H_n is shown in Fig. 3. Similarly, when we delete the edge(s) D in the interior face of H_7 , the diameter of the resulting graph becomes 3. Therefore, we obtain that $\chi'_{st}(G_7) \leq \chi'_{st}(C_7) + |D| = 3 + |D| = \Delta(G_7) + 1$.




Fig. 3. F_6 and F_7

Case 3. If $n \geq 8$, to ensure $\text{diam}(G_n) = 3$, by enumeration, we can obtain that the deleted edges D ($|D| \leq 3$). Therefore, we can get the formula as follows, $\chi'_{st}(G_n) \leq \chi'_{st}(H_n) = \Delta(H_n) \leq \Delta(G_n) + 3$.

Finally, when G_n is an edge induced subgraph of $H_n = F_n (\in \xi_n^2)$, we obtain that $\Delta(G_n) \leq \chi'_{st}(G_n) \leq \Delta(G_n) + 3$. \square

3.2 $H_n \in \xi_n^3$

We firstly discuss the property of H_n in the following.

Theorem 3.2 If $H_n \in \xi_n^3$, then H_n has G_6^1 or G_6^2 (shown in the Fig. 4) as a vertex induced subgraph.

Proof: By the construction method of maximal outerplanar graph in [3], maximal outerplanar graph of order 5 is F_5 under the meaning of isomorphism, thus F_5 must be a vertex induced subgraph of $H_n(n \geq 5)$. Assume that the maximum degree vertex of F_5 is v_0 and other vertices are v_1, v_2, v_3, v_4 . We add a vertex v_5 to F_5 in order to obtain a big maximal outerplanar graph: if v_5 is adjacent to $v_i, v_{i+1}(i \in \{1, 2, 3\})$, the theorem is proven; if v_5 is adjacent to v_0, v_1 or v_0, v_4 , we obtain a Fan F_6 , then, we continue to add vertex, there is some vertex v_j which is adjacent to $v_i, v_{i+1}(i \in \{1, 2, 3, \dots, j-1\})$, otherwise, the diameter of the resulting graph is 2. \square




Fig. 4. G_6^2




Fig. 5. $H_n(G_6^1 \subseteq H_n)$

Theorem 3.3 Let G be a connected graph. When G has two adjacent vertices $v_0, v_1 \in G$ and the distance between v_0 and any other vertex of G is less than $diam(G)$, then adding some vertices v_2, v_3, \dots, v_n and edges $v_1v_2, v_2v_3, \dots, v_{n-1}v_n, v_0v_2, v_0v_3, \dots, v_0v_n$, we obtain a new graph G' with $diam(G') = diam(G)$.

By theorem 3.3, we know that $v_0v_1, v_1v_2, v_2v_3, v_3v_4, v_4v_0$ can be extended to be Fans as shown in Fig. 5. Assume that $\Delta(H_n) = \Delta$, then $\Delta = \max\{d(v_0), d(v_2), d(v_3)\}$. Especially, the graph of $d(v_0) = d(v_2) = d(v_3) = \Delta$ is denoted by $H_{n\Delta}$, we give a lower bound of $\chi'_{st}(H_{n\Delta})$ in the following theorem.

Theorem 3.4 $\chi'_{st}(H_{n\Delta}) \geq \Delta(H_{n\Delta}) + 2$.

Proof: We consider the subgraph $G_{n\Delta}$ (shown as the Fig. 6) of $H_{n\Delta}$. We firstly prove $\chi'_{st}(G_{n\Delta}) \geq \Delta + 1$. It is easy to know that $\chi'_{st}(G_{n\Delta}) \geq \Delta$ by the definition of star edge coloring. Assume that $\chi'_{st}(G_{n\Delta}) = \Delta$, since $d(v_0) = \Delta$, there are some vertex $v'_0 \in N(v_0)$ such that $c(v_0v'_0) = c(v_2v_3)$. Similarly, there are some vertex $v'_3 \in N(v_3)$ such that $c(v_3v'_3) = c(v_0v_2)$, thus the path $v'_0 - v_0 - v_2 - v_3 - v'_3$ is 2-edge colored, which contradicts to the definition of star edge coloring. Therefore, we obtain that $\chi'_{st}(H_{n\Delta}) \geq \chi'_{st}(G_{n\Delta}) \geq \Delta + 1$.

Now, we assume that $\chi'_{st}(G_{n\Delta}) = \Delta + 1$. Without loss of generality, let $c(v_0v_1) = 1, c(v_0v_0^{(1)}) = 2, c(v_0v_0^{(2)}) = 3, \dots, c(v_0v_0^{(\Delta-4)}) = \Delta - 3, c(v_0v_4) = \Delta - 2, c(v_0v_3) = \Delta - 1, c(v_0v_2) = \Delta, c(v_2v_3) = \Delta + 1$ (If there are some vertex $v_{x_1} \in \{v_1, v_0^{(1)}, v_0^{(2)}, \dots, v_0^{(\Delta-4)}, v_4\}$, such that $c(v_0v_{x_1}) = c(v_2v_3)$, then for any vertex $v_y \in \{v_1, v_2^{(1)}, v_2^{(2)}, \dots, v_2^{(\Delta-4)}, v_5\}$, we have $c(v_2v_y) \neq c(v_0v_3)$). The coloring scheme is isomorphic to the form: $c(v_0v_1) = 1, c(v_0v_0^{(1)}) = 2, c(v_0v_0^{(2)}) = 3, \dots, c(v_0v_0^{(\Delta-4)}) = \Delta - 3, c(v_0v_4) = \Delta - 2, c(v_0v_3) = \Delta - 1, c(v_0v_2) = \Delta, c(v_2v_3) = \Delta + 1$.

Let $\{c(v_3v_4), c(v_3v_5), c(v_3v_3^{(1)}), c(v_3v_3^{(2)}), \dots, c(v_3v_3^{(\Delta-4)})\} = Q$. Firstly, we prove that $\Delta - 2 \notin Q$: since $c(v_3v_4) \neq \Delta + 1$ or $\Delta - 2$, there is some vertex $v_{x_2} \in \{v_2, v_1, v_0^{(1)}, v_0^{(2)}, \dots, v_0^{(\Delta-4)}\}$ such that $c(v_0v_{x_2}) = c(v_3v_4)$, which forces that there is a 4-path colored by $\Delta - 2, c(v_3v_4)$ in G_{n_Δ} when $\Delta - 2 \in Q$. The coloring is not proper.

Since $\{\Delta - 2, \Delta - 1, \Delta + 1\} \notin Q$ and $|Q| = \Delta - 2$, $Q = \{1, 2, \dots, \Delta - 3, \Delta\}$. Similarly, we have $\{1, \Delta, \Delta + 1\} \notin \{c(v_2v_1), c(v_2v_5), c(v_2v_2^{(1)}), c(v_2v_2^{(2)}), \dots, c(v_2v_2^{(\Delta-4)})\}$. Since $\Delta \in Q$, $\Delta - 1 \notin \{c(v_2v_1), c(v_2v_5), c(v_2v_2^{(1)}), c(v_2v_2^{(2)}), \dots, c(v_2v_2^{(\Delta-4)})\}$, a contradiction. \square

Theorem 3.5 If $H_n \in \xi_n^3$, and $G_n \in \zeta_n^3$, then $\Delta(G_n) \leq \chi'_{st}(G_n) \leq \Delta(G_n) + 6$; when $G_n \in \xi_n^3$, we have $\Delta(G_n) \leq \chi'_{st}(G_n) \leq \Delta(G_n) + 4$.

Proof: We prove the theorem through the following two cases 3.5.1 and 3.5.2.

Case 3.5.1 H_n has G_6^1 as an induced subgraph.

We get the Fig. 7 (denoted by H'_{n_Δ}) by adding edges $v_1v_0^{(1)}, v_0^{(i)}v_0^{(i+1)}, v_2^{(1)}v_1, v_2^{(j)}v_2^{(j+1)}, v_2^{(\Delta-4)}v_5, v_5v_3^{(1)}, v_3^{(k)}v_3^{(k+1)}, v_3^{(\Delta-4)}v_4, v_0^{(\Delta-4)}v_4$ in G_{n_Δ} where $1 \leq i, j, k \leq \Delta - 3$. In the following, we discuss the upper bound of $\chi'_{st}(H'_{n_\Delta})$ in terms of Δ .




Fig. 6. G_{n_Δ}




Fig. 7. H'_{n_Δ}

- (1) When $\Delta = 5$, the Fig. 8(a) gives a star edge coloring, then $\Delta + 2 = 7 \leq \chi'_{st}(H'_{n_\Delta}) \leq 9 = \Delta + 4$.
- (2) If $\Delta = 6$, a star edge coloring is given in the Fig. 8(b), we obtain that $\Delta + 2 = 8 \leq \chi'_{st}(H'_{n_\Delta}) \leq 9 = \Delta + 3$.
- (3) For the case $\Delta = 7$, it is easy to check that the coloring of the Fig. 8(c) is a star edge coloring, thus $\Delta + 2 = 9 \leq \chi'_{st}(H'_{n_\Delta}) \leq 11 = \Delta + 4$.
- (4) $\Delta = 8$, we consider the star edge coloring of the Fig. 8(d), thus $\Delta + 2 = 10 \leq \chi'_{st}(H'_{n_\Delta}) \leq 11 = \Delta + 3$.
- (5) When $\Delta \geq 9$, we consider the following coloring function $c : c(v_0v_1) = 1, c(v_0v_4) = \Delta - 2, c(v_2v_1) = 2, c(v_2v_5) = \Delta + 2, c(v_3v_5) = 1, c(v_3v_4) = \Delta + 2, c(v_0v_2) = \Delta, c(v_0v_3) = \Delta - 1, c(v_2v_3) = \Delta + 1, c(v_0v_0^{(l)}) = l + 1, c(v_2v_2^{(l)}) = l + 2, c(v_3v_3^{(l)}) = l + 1, c(v_1v_0^{(1)}) = \Delta + 3, c(v_0^{(m-1)}v_0^{(m)}) = m + 3, c(v_0^{(\Delta-5)}v_0^{(\Delta-4)}) = \Delta + 1, c(v_0^{(\Delta-4)}v_4) = \Delta + 3, c(v_1v_2^{(1)}) = 5, c(v_2^{(n-1)}v_2^{(n)}) = n + 4, c(v_2^{(\Delta-5)}v_2^{(\Delta-4)}) = 2, c(v_2^{(\Delta-4)}v_5) = 3, c(v_5v_3^{(1)}) = 4, c(v_3^{(p-1)}v_3^{(p)}) = p + 3, c(v_3^{(\Delta-6)}v_3^{(\Delta-5)}) = \Delta + 2, c(v_3^{(\Delta-5)}v_3^{(\Delta-4)}) = \Delta, c(v_3^{(\Delta-4)}v_4) = 2$, where $l, m, p \in \mathbb{Z}_+$ and $m \in [2, \Delta - 5], n \in [2, \Delta - 5], p \in [2, \Delta - 6]$. As an example, the Fig.8(e) gives a star edge coloring of H'_{n_9} .

Next, we prove that the coloring c is a star edge coloring: the paths $v_1 - v_2^{(1)} - v_2^{(2)} - \dots - v_2^{(\Delta-5)} -$

$v_2^{(\Delta-4)} - v_5, v_5 - v_3^{(1)} - v_3^{(2)} - \dots - v_3^{(\Delta-5)} - v_3^{(\Delta-4)} - v_4, v_1 - v_0^{(1)} - v_0^{(2)} - \dots - v_0^{(\Delta-5)} - v_0^{(\Delta-4)} - v_4$ are denoted by P_2, P_3, P_0 , respectively. By the Theorem 1.4 of [5], we know that the Fans $P_0 \vee v_0, P_2 \vee v_2, P_3 \vee v_3$ has no bichromatic 4-paths(cycles).

Now, assume P is a bichromatic 3-path(cycle) of H'_{n_Δ} . If the path(cycle) P contains v_0v_2, v_0v_3 or v_2v_3 as an edge, then all the bichromatic 3-path(cycle) P of H'_{n_Δ} are $v_2^{(i)} - v_2 - v_3 - v_3^{(i+1)}, v_1 - v_2 - v_3 - v_3^{(1)}, v_5 - v_2 - v_3 - v_4, v_2^{(i)} - v_2 - v_0 - v_0^{(i+1)}, v_1 - v_2 - v_0 - v_0^{(1)}, v_2^{(5)} - v_2 - v_0 - v_4, v_0^{(j)} - v_0 - v_3 - v_3^{(j)}, v_1 - v_0 - v_3 - v_5$, where $i, j \in \mathbb{Z}_+$ and $i \in [1, \Delta - 5], j \in [1, \Delta - 4]$. It is easy to check that any one of the bichromatic 3-path(cycle) can't be extended to become bichromatic 4-paths(cycles), thus H'_{n_Δ} has no bichromatic 4-paths(cycles) containing v_0v_2, v_0v_3 or v_2v_3 as an edge.

If the path(cycle) P contains $v_1v_0^{(1)}, v_1v_2^{(1)}, v_4v_0^{(\Delta-4)}, v_4v_3^{(\Delta-4)}, v_5v_2^{(\Delta-4)}$ or $v_5v_3^{(1)}$ as an edge, then all the bichromatic 3-path(cycle) P of H'_{n_Δ} are (the path P containing v_0v_2, v_0v_3 or v_2v_3 as an edge can be discussed similarly) $v_2^{(1)} - v_1 - v_2 - v_2^{(3)}, v_1 - v_2^{(1)} - v_2 - v_2^{(3)}, v_2^{(1)} - v_1 - v_0 - v_0^{(4)}, v_2^{(\Delta-4)} - v_5 - v_2 - v_2^{(1)}, v_5 - v_2^{(\Delta-4)} - v_2 - v_2^{(1)}, v_2^{(\Delta-4)} - v_5 - v_3 - v_3^{(2)}, v_3^{(1)} - v_5 - v_3 - v_3^{(3)}, v_5 - v_3^{(1)} - v_3 - v_3^{(3)}, v_3^{(1)} - v_5 - v_2 - v_2^{(2)}, v_3^{(\Delta-4)} - v_4 - v_3 - v_3^{(1)}, v_4 - v_3^{(\Delta-4)} - v_3 - v_3^{(1)}, v_3^{(\Delta-4)} - v_4 - v_0 - v_0^{(1)}$. It is easy to check that any one of the bichromatic 3-path(cycle) can't be extended to become bichromatic 4-paths(cycles), thus the graph H'_{n_Δ} has no bichromatic 4-paths(cycles) containing $v_1v_0^{(1)}, v_1v_2^{(1)}, v_4v_0^{(\Delta-4)}, v_4v_3^{(\Delta-4)}, v_5v_2^{(\Delta-4)}$ or $v_5v_3^{(1)}$ as an edge.

In the following, we consider the paths P containing $v_0v_1, v_1v_2, v_2v_5, v_5v_3, v_3v_4$ or v_4v_0 as an edge. All the bichromatic 3-path(cycle) P of H'_{n_Δ} are (the path P containing $v_0v_2, v_0v_3, v_2v_3, v_1v_0^{(1)}, v_1v_2^{(1)}, v_4v_0^{(5)}, v_4v_3^{(5)}, v_5v_2^{(5)}$ or $v_5v_3^{(1)}$ as an edge can be discussed similarly) $v_1 - v_2 - v_2^{(\Delta-4)} - v_2^{(\Delta-5)}, v_1 - v_2 - v_2^{(\Delta-5)} - v_2^{(\Delta-4)}, v_2 - v_1 - v_0 - v_0^{(1)}, v_2 - v_5 - v_3 - v_4, v_4 - v_3 - v_3^{(\Delta-6)} - v_3^{(\Delta-5)}, v_4 - v_3 - v_3^{(\Delta-5)} - v_3^{(\Delta-6)}, v_4 - v_0 - v_0^{(\Delta-6)} - v_0^{(\Delta-5)}, v_4 - v_0 - v_0^{(\Delta-5)} - v_0^{(\Delta-6)}$. One can find that any one of the bichromatic 3-path(cycle) can't be extended to become bichromatic 4-paths(cycles), thus the graph H'_{n_Δ} has no bichromatic 4-paths(cycles) containing $v_0v_1, v_1v_2, v_2v_5, v_5v_3, v_3v_4$ or v_4v_0 as an edge.

Let H''_{n_Δ} be the subgraph by deleting $v_0v_2, v_0v_3, v_2v_3, v_1v_0^{(1)}, v_1v_2^{(1)}, v_4v_0^{(\Delta-4)}, v_4v_3^{(\Delta-4)}, v_5v_2^{(\Delta-4)}, v_5v_3^{(1)}, v_0v_1, v_1v_2, v_2v_5, v_5v_3, v_3v_4$ and v_4v_0 in H'_{n_Δ} . The graph H''_{n_Δ} is not connected and has three connected components, which are vertex-induced subgraphs of Fans $P_0 \vee v_0, P_2 \vee v_2$ and $P_3 \vee v_3$, thus H''_{n_Δ} doesn't contain bichromatic 4-paths(cycles), therefore H'_{n_Δ} has no bichromatic 4-paths(cycles).

So overall, we obtain that $\Delta + 2 \leq \chi'_{st}(H_{n_\Delta}) \leq \Delta + 3$ when $\Delta \geq 9$. When $\Delta = 5$ or 7 , we proved that




Fig. 8. Star edge coloring of H'_{n_Δ} ($\Delta = 5, 6, \dots, 9$)

$\chi'_{st}(H_{n_\Delta}) \leq \Delta + 4$. In the end, we have $\Delta \leq \chi'_{st}(H_n) \leq \chi'_{st}(H_{n_\Delta}) \leq \Delta + 3$ when $\Delta \geq 8$ or $\Delta = 6$. \square

Case 3.5.2 If H_n does not contain G_6^1 , then H_n must contain G_6^2 as an edge induced subgraph.

By the construction method of maximal outerplanar graph in [3], it is easy to check that there is no vertex adjacent to v_0, v_4 or v_2, v_3 , otherwise H_n must contain an edge induced subgraph G_6^1 . Since H_n has diameter three, H_n has two cases shown as Fig. 9.




Fig. 9. $H_n(G_6^2 \subseteq H_N)$

Case 1: As shown in the left of Fig. 9, we discuss the case $d(v_0) = d(v_3) = d(v_4) = \Delta$ (denoted by H_{n_Δ}). We consider the upper bound of $\chi'_{st}(H_{n_\Delta})$ in terms of Δ .

(1) If $\Delta = 4$, it is easy to check that the coloring of Fig. 10(a) is a star edge coloring with $\Delta + 2$ colors, then $\chi'_{st}(H_{n_\Delta}) \leq 6 = \Delta + 2$. Since F_5 is an edge induced subgraph and $\chi'_{st}(F_5) = \Delta + 2$, thus $\chi'_{st}(H_{n_\Delta}) = \Delta + 2$;

(2) The case of $\Delta = 5$, the Fig. 10(b) give a star edge coloring with $\Delta + 3$ colors, therefore $\chi'_{st}(H_{n_\Delta}) \leq 8 = \Delta + 3$;

(3) For $\Delta = 6$, we consider the Fig. 10(c) which has a star edge coloring with $\Delta + 3$ colors, thus $\chi'_{st}(H_{n_\Delta}) \leq 9 = \Delta + 3$;

(4) When $\Delta \geq 7$, we consider the following coloring function $c : c(v_0v_1) = \Delta - 3, c(v_0v_2) = \Delta - 2, c(v_0v_3) = \Delta + 1, c(v_0v_4) = \Delta + 2, c(v_3v_5) = \Delta - 1, c(v_4v_5) = \Delta + 4, c(v_0v_0^{(l)}) = l, c(v_3v_3^{(m)}) = m + 2, c(v_4v_4^{(p)}) = p + 2, c(v_1v_2) = \Delta + 3, c(v_2v_3) = \Delta, c(v_3v_4) = \Delta + 3, c(v_0^{(l)}v_0^{(l+1)}) = l + 3, c(v_3^{(m)}v_3^{(m+1)}) = m, c(v_4^{(p)}v_4^{(p+1)}) = p, c(v_3^{(\Delta-4)}v_5) = \Delta - 4, c(v_5v_4^{(1)}) = \Delta$, where $l, m, p \in \mathbb{Z}_+$ and $l, m \in [l, \Delta - 4], p \in [l, \Delta - 3]$.

We prove the coloring function c is a star edge coloring: the paths $v_0^{(1)} - v_0^{(2)} - \dots - v_0^{(\Delta-4)} - v_1 - v_2 - v_3 - v_4, v_3^{(1)} - v_3^{(2)} - \dots - v_3^{(\Delta-4)} - v_5, v_5 - v_4^{(1)} - v_4^{(2)} - \dots - v_4^{(\Delta-4)} - v_4^{(\Delta-3)}$ are denoted by P_0, P_3, P_4 respectively. By the theorem 1.4 of [5], we know that the Fans $P_0 \vee v_0, P_3 \vee v_3, P_4 \vee v_4$ have no bichromatic paths or cycles of length four. Assume P is a path(cycle) of H_{n_Δ} containing v_0v_3, v_3v_4, v_2v_3 or $v_5v_4^{(1)}$ and is a bichromatic path(cycle) with length three. All bichromatic 3-path(cycle) of H_{n_Δ} are $v_1 - v_2 - v_3 - v_4, v_2 - v_3 - v_5 - v_4^{(1)}, v_0 - v_2 - v_3 - v_3^{(3)}, v_3^{(i)} - v_3 - v_4 - v_4^{(i)}$ ($1 \leq i \leq \Delta - 4$), $v_5 - v_3 - v_4 - v_4^{(\Delta-3)}, v_4 - v_4^{(1)} - v_5 - v_3^{(3)}$. By the coloring function c , we know that all the paths(cycles) can't be become bichromatic 4-paths(cycles) by extending, thus H_{n_Δ} does not contain bichromatic 4-paths(cycles) including v_0v_3, v_3v_4, v_2v_3 or $v_5v_4^{(1)}$. Since $c(v_0v_4) = \Delta + 2, c(v_4v_5) = \Delta + 4$, and $\Delta + 2, \Delta + 4$ appear only once in H_{n_Δ} , deleting the edges $v_3v_4, v_2v_3, v_0v_3, v_0v_4, v_4v_5$ and $v_5v_4^{(1)}$, we obtain a subgraph denoted by H'_{n_Δ} . The subgraph H'_{n_Δ} is not connected and has three connected components, which are vertex induced Fans $P_0 \vee v_0, P_3 \vee v_3$ and $P_4 \vee v_4$, thus H'_{n_Δ} has no bichromatic 4-paths and cycles. Therefore, H_{n_Δ} has no bichromatic 4-paths and cycles. We obtain that $\chi'_{st}(H_{n_\Delta}) \leq \Delta + 4$. As an example, the Fig. 10(d) gives the coloring function of H_{n_7} .

Thus $\Delta \leq \chi'_{st}(H_n) \leq \Delta + 4$.




Fig. 10. Case 1: Star edge coloring of H_{n_Δ} ($\Delta = 4, 5, 6, 7$)

Case 2: This case of diagram is shown in Fig. 11, we discuss the graph with $d(v_2) = d(v_3) = \Delta$ (denoted by $H_{n_\Delta}^2$). We consider the upper bound of $\chi'_{st}(H_{n_\Delta}^2)$ based on the value of Δ .

- (1) When $\Delta = 4$, we consider the coloring of the Fig. 11(a), then $\chi'_{st}(H_{n_\Delta}^2) \leq 11 = \Delta + 3$.
- (2) If $\Delta = 5$, by the coloring of the Fig. 11(b), we know that $\chi'_{st}(H_{n_\Delta}^2) \leq \Delta + 3$, moreover $\chi'_{st}(H_{n_\Delta}^2) \geq \chi'_{st}(F_5) = \Delta + 2$, thus $\Delta + 2 = 7 \leq \chi'_{st}(H_{n_\Delta}^2) \leq 8 = \Delta + 3$.
- (3) When $\Delta = 6$, the Fig. 11(c) gives its a star edge coloring with $\Delta + 3$ colors, thus $\chi'_{st}(H_{n_\Delta}^2) \leq 9 = \Delta + 3$.
- (4) For $\Delta = 7$, considering the coloring of the Fig. 11(d), we obtain that $\chi'_{st}(H_{n_\Delta}^2) \leq 10 = \Delta + 3$.
- (5) The case of $\Delta = 8$, a star edge coloring is shown in the Fig. 11(e), thus $\chi'_{st}(H_{n_\Delta}^2) \leq 11 = \Delta + 3$.
- (6) When $\Delta = 9$, we consider the coloring of the Fig. 11(f), then $\chi'_{st}(H_{n_\Delta}^2) \leq 11 = \Delta + 2$.
- (7) $\Delta = 10$, the Fig. 11(g) gives its a star edge coloring with $\Delta + 1$, thus $\chi'_{st}(H_{n_\Delta}^2) \leq 12 = \Delta + 2$.
- (8) When $\Delta \geq 10$, we consider the following coloring function $c : c(v_0v_1) = \Delta + 2, c(v_0v_2) = \Delta - 1, c(v_0v_3) = \Delta + 1, c(v_0v_4) = 5, c(v_1v_2) = \Delta, c(v_2v_3) = \Delta - 2, c(v_3v_4) = \Delta, c(v_1v_5) = \Delta + 1, c(v_2v_5) = 1, c(v_3v_6) = \Delta - 3, c(v_4v_6) = \Delta + 2, c(v_2v_2^{(l)}) = l + 1, c(v_3v_3^{(m)}) = m, c(v_5v_2^{(1)}) = 4, c(v_2^{(p)}v_2^{(p+1)}) = p + 4, c(v_2^{(\Delta-6)}v_2^{(\Delta-5)}) = 1, c(v_2^{(\Delta-5)}v_2^{(\Delta-4)}) = 2, c(v_3^{(q)}v_3^{(q+1)}) = q + 3, c(v_3^{(\Delta-5)}v_3^{(\Delta-4)}) = 1, c(v_3^{(\Delta-4)}v_4) = 2$, where $l, m, p, q \in \mathbb{Z}_+$ and $l, m \in [l, \Delta - 4], p, q \in [l, \Delta - 6]$.

We prove the coloring function c is a star edge coloring in the case(8): the paths $v_5 - v_2^{(1)} - v_2^{(2)} - \dots - v_2^{(\Delta-5)} - v_2^{(\Delta-4)}, v_3^{(1)} - v_3^{(2)} - \dots - v_3^{(\Delta-5)} - v_3^{(\Delta-4)} - v_6$ are denoted by P_2, P_3 respectively. By the theorem 1.4 of [5], we know that the Fans $P_2 \vee v_2$ and $P_3 \vee v_3$ have no bichromatic paths and cycles of length four. Assume the bichromatic path(cycle) P containing an edge $v_0v_1, v_0v_2, v_0v_3, v_0v_4, v_1v_5, v_4v_6, v_1v_2$ or v_3v_4 in $H_{n_\Delta}^2$. All the bichromatic 3-path(cycle) P are $v_5 - v_1 - v_0 - v_3, v_6 - v_4 - v_0 - v_1, v_1 - v_2 - v_3 - v_4, v_2^{(4)} - v_2 - v_0 - v_4, v_3^{(5)} - v_3 - v_0 - v_4$ in $H_{n_\Delta}^2$. It is easy to check that any one of the above paths P can't become bichromatic 4-path(cycle) by extending, then the graph $H_{n_\Delta}^2$ have no bichromatic 4-paths(cycles) including $v_0v_1, v_0v_2, v_0v_3, v_0v_4, v_1v_5, v_4v_6, v_1v_2$ or v_3v_4 . Since $c(v_2v_3) = \Delta - 2$, and $\Delta - 2$ appear only once in $H_{n_\Delta}^2$, thus $H_{n_\Delta}^2$ have no bichromatic 4-paths(cycles) including the edge v_2v_3 . By deleting the edges $v_0v_1, v_0v_2, v_0v_3, v_0v_4, v_1v_5, v_4v_6, v_1v_2, v_3v_4$ and v_2v_3 , we obtain the subgraph $H_{n_\Delta}^{2'}$. The graph $H_{n_\Delta}^{2'}$ is not connected and has two connected components, which are Fans $P_2 \vee v_2$ and $P_3 \vee v_3$, thus $H_{n_\Delta}^{2'}$ have no bichromatic 4-paths(cycles). In summary, $H_{n_\Delta}^2$ have no bichromatic 4-paths(cycles). We obtain that $\chi'_{st}(H_{n_\Delta}^2) \leq \Delta + 2$.

Thus, $\Delta \leq \chi'_{st}(H_n) \leq \chi'_{st}(H_{n_\Delta}^2) \leq \Delta + 2$.

According to the Cases 3.5.1 and 3.5.2, when $H_n \in \xi_n^3$, we obtain that $\Delta(H_n) \leq \chi'_{st}(H_n) \leq \Delta(H_n) + 4$. In the document [8], a classification of $G_n (\in \zeta_n^3)$ has been given. On the basis of the classification, we find that for every $G_n \in \zeta_n^3$ there exists a $H_n \in \xi_n^3$ such that G_n is a subgraph of H_n and $\Delta(H_n) \leq \Delta(G_n) + 2$. Therefore, we can obtain that $\Delta(G_n) \leq \chi'_{st}(G_n) \leq \chi'_{st}(H_n) \leq \Delta(H_n) + 4 \leq \Delta(G_n) + 6$. \square




Fig. 11. Case 2: the star edge coloring of $H_{n\Delta}^2$ ($\Delta = 5, \dots, 10$)

4 Outerplanar graphs with $\Delta = 5$

Theorem 4.1 Let G be a 2-connected outerplanar graph with n vertices and $\Delta(G) = 5$, then $\chi'_{st}(G) \leq \lfloor 1.5\Delta \rfloor + 2 = \Delta + 4 = 9$.

Proof. Since G has n vertices and $\Delta(G) = 5$, obviously $n \geq 6$. When $n = 6$, G is isomorphic to F_6 . By Lemma 2.3, we know that $\chi'_{st}(F_6) = 6 < 9$. Since maximal outerplanar graphs with maximal degree 5 can be drawn as Fig. 12 when the order is going infinite and the F_6 s between E and F form a cycle. Inspired by the document [14], we give a star 9-edge coloring as Fig. 12 showing. If $n > 6$, any outerplanar graph with maximum degree 5 can become a maximal outerplanar graph with maximum degree 5 by adding edges, moreover it is a subgraph of Fig. 12. Since Fig. 12 has a 9 star edge coloring, we obtain that any 2-connected outerplanar graph with $\Delta(G) = 5$ has star chromatic index at most 9.




Fig. 12. A 9 star edge coloring

5 Concluding remarks

In this paper, we discuss the star chromatic index of outerplanar graphs with small diameter inspired by the Conjecture 1.5. Otherwise, we find that the upper bound of star chromatic index of outerplanar graphs with diameter 2 or 3 is $\Delta + 6$, which is much less than $\lfloor \frac{3\Delta}{2} \rfloor + 1$ when Δ increases infinitely. Theorem 1.5 tells us that $\chi'_{st}(G) \leq 5$ when G is a subcubic outerplanar graph. By Theorem 1.7, we know that outerplanar graphs with $\Delta = 4$ have star chromatic index 6. Moreover, Theorem 4.1 obtains that 2-connected outerplanar graphs have star chromatic indexes no more than 9 when $\Delta = 5$. So overall, these results make us sure that the following conjecture is true.

Conjecture 4.1 If G is a 2-connected outerplanar graph with the maximum degree $\Delta \geq 6$, then $\chi'_{st}(G) \leq \Delta + 6$.

Conjecture 4.2 If G is a 2-connected maximal outerplanar graph with the maximum degree $\Delta \geq 6$, then $\chi'_{st}(G) \leq \Delta + 4$.

In [2], Bezegová et al. proved that $\chi'_{st}(T) \leq \lfloor \frac{3\Delta}{2} \rfloor$, when T be a tree with maximum degree Δ and every tree with a Δ -vertex whose all neighbors are Δ -vertices achieves the upper bound. We propose the following open question inspired by Bezegová's and our results.

Problem 4.2 If G is a 2-connected graph with linear size and maximum degree Δ , then $\chi'_{st}(G) \leq \Delta + c$ (c is a constant)?

References

- [1] B. Bollbás, W. Fulton, A. Katok, F. Kirwan, P. Sarnak, Random graphs(Second Edition), Cambridge University Press, England 2001.
- [2] L. Bezegová, B. Lužar, M. Mockovčiaková, R. Soták, R. škrekovski, Star edge coloring of some classes of graphs, J. Graph Theory 81 (2016)73-82 .
- [3] L. Beineke and R. Pippert, A census of ball and disk dissections, Graph Theory and Applications (Proc. Conf., Western Michigan Univ., Kalamazoo, Mich., 1972), 25-40, Lect. Notes. Math., 303, Springer, 1972.
- [4] Z. Dvořák, B. Mohar, R. Šámal, Star chromatic index, J. Graph Theory 72 (2013)313-326.
- [5] B. Grünbaum, Acyclic colorings of planar graphs, Israel J. Math. 14 (1973), 390-408.
- [6] K. Deng , S. L. Tian, Star edge coloring of maximal outer plane graphs, Applied Mathematics A Journal of Chinese Universities 26(4) (2011)489-494.
- [7] B. X. -S. Liu, K. Deng, An upper bound on the star chromatic index of graphs with $\Delta \geq 7$, J. Lanzhou Univ (Nat Sci) 44 (2008)94-95.
- [8] X. Deng, H. Song, G. Su, R. Tian, Rainbow connection number of biconnected outerplanar graphs with small diameters. Advances in mathematics(China), 2018, 47(03)373-382.
- [9] X. S. Liu , K. Deng , Star edge coloring of graphs, Master thesis (2007).
- [10] M. Molloy, B. Reed , A bound on the strong chromatic index of a graph, J. Comb. Theory Ser. B 69 (1997)103-109 .
- [11] G. Chartrand, F. Harary, Planar permutation graphs, Ann. Inst. H. Poincaré B. 3 (1967)433-438.
- [12] G. Hopkins, W. Staton, Outerplanarity without topology, Bull. Inst. Comb. Appl. 21 (1997)112-116.

- [13] H. Lei, Y. Shi, Z. Song, Star chromatic index of subcubic multigraph, *J. Graph Theory* 88(4) (2017)1-11.
- [14] H. Lei, Y. Shi, Z. Song, T. Wang, Star 5-edge-colorings of subcubic multigraphs, *Discrete Mathematics* 341(4) (2018)950-956.
- [15] S. Mitchell, Algorithms on trees and maximal outerplanar graphs: design, complexity analysis, and data structures study, PhD Thesis, University of Virginia, Charlottesville, VA, USA, 1977.
- [16] D. B. West, Introduction to graph theory (second edition), Prentice Hall, NJ, 2001.
- [17] Y. Q. Wang, W. F. Wang, Y. Wang, Edge-partition and star chromatic index, *Applied Mathematics and Computation*, 333 (2018)480-489.