

On the Borel Submonoid of a Symplectic Monoid

Mahir Bilen Can, Hayden Houser, Corey Wolfe

Department of Mathematics, Tulane University,
6823 St. Charles Ave, New Orleans, LA, 70118,

January 20, 2022

Abstract

In this article, we study the Bruhat-Chevalley-Renner order on the complex symplectic monoid MSp_n . After showing that this order is completely determined by the Bruhat-Chevalley-Renner order on the linear algebraic monoid of $n \times n$ matrices M_n , we focus on the Borel submonoid of MSp_n . By using this submonoid, we introduce a new set of type B set partitions. We determine their count by using the “folding” and “unfolding” operators that we introduce. We show that the Borel submonoid of a rationally smooth reductive monoid with zero is rationally smooth. Finally, we analyze the nilpotent subsemigroups of the Borel semigroups of M_n and MSp_n . We show that, contrary to the case of MSp_n , the nilpotent subsemigroup of the Borel submonoid of M_n is irreducible.

Keywords: Symplectic monoid, Renner monoid, Borel submonoid, rationally smooth, set partitions, (un)folding
MSC: 20M32, 20G99, 06A99

1 Introduction

Let M be a complex reductive monoid with unit group G , and let B be a Borel subgroup in G . Then we have a square of inclusions as in the following diagram

$$\begin{array}{ccc} \overline{B} & \hookrightarrow & M \\ \uparrow & & \uparrow \\ B & \hookrightarrow & G \end{array}$$

where \overline{B} is the Zariski closure of B in M ; we will call \overline{B} the *Borel submonoid* determined by B . Although its combinatorics and geometry are relatively less explored compared to that of the ambient reductive monoid, the Borel submonoid is a very important object for

the study of the representation theory of M [11, Theorem 3.4]. In the special case of the linear algebraic monoid of $n \times n$ matrices, the $B \times B$ -orbits in \overline{B} are parametrized by the set partitions of $\{1, \dots, n\}$, providing a gateway to an uncharted domain for combinatorialists, see [7]. In this regard, our goal in this paper is to present first combinatorial results, whose analogous versions are obtained in [7], for the Borel submonoid of a “symplectic monoid” that we will define next.

Let l be a positive integer, and set $n := 2l$. The set of all $l \times l$ matrices with entries from \mathbb{C} will be denoted by M_l . We let J denote the $n \times n$ matrix, $J = \begin{bmatrix} 0 & J_l \\ -J_l & 0 \end{bmatrix}$, where J_l is the unique antidiagonal $l \times l$ permutation matrix, that is,

$$J_l = \begin{bmatrix} 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 1 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 1 & \cdots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \end{bmatrix}.$$

The *symplectic group* is defined by $Sp_n := \{A \in GL_n : A^\top J A = J\}$. This is the group of linear automorphisms of \mathbb{C}^n that preserve the skew-bilinear form that is defined by J . (Once we fix the even integer $n = 2l$, in the sequel, it will be convenient for us to denote Sp_n by G .) Let us denote the central extension of Sp_n in GL_n by GSp_n . This is the smallest reductive subgroup of GL_n that contains both of the subgroups G and the group of invertible scalar matrices $\{cI_n : c \in \mathbb{C}^*\}$, where I_n denotes the $n \times n$ identity matrix. The Zariski closure of GSp_n in M_n is called the *n -th symplectic monoid*. Such monoids were first considered by Grigor’ev [13]. The following concrete description of the n -th symplectic monoid, which we will denote by MSp_n , is due to Doty [10, Proposition 4.3]: $MSp_n := \{A \in M_n : A^\top J A = A J A^\top = cJ, c \in \mathbb{C}\}$. Basic geometric ingredients (the Renner monoid, the cross section lattice, and a cell decomposition) of MSp_n are described explicitly by Li and Renner in [16]. An in-depth analysis of the rational points of MSp_n over finite fields, including some fascinating combinatorial formulas about its Renner monoids, are described by Cao, Lei, and Li in [9]. To describe the main results of our paper, next, we will briefly review the Renner monoid of MSp_n in relation with the rook monoid.

To keep our notation simple, let us denote by B the Borel subgroup consisting of the upper triangular matrices in GSp_n . The *natural action* of $B \times B$ on MSp_n is defined by $(b_1, b_2) \cdot x = b_1 x b_2^{-1}$, where $b_1, b_2 \in B$, $x \in MSp_n$. This action has finitely many orbits [21, 16] and moreover the orbits are parametrized by a finite inverse semigroup, $MSp_n = \bigsqcup_{\sigma \in \mathcal{R}_G} B \sigma B$. The finite inverse semigroup \mathcal{R}_G is called the *symplectic Renner monoid*; it is the symplectic version of the *rook monoid* \mathcal{R}_n , which consists of 0/1 square matrices of size n with at most one 1 in each row and each column. In fact, \mathcal{R}_G is a submonoid of \mathcal{R}_n . The elements of \mathcal{R}_n are called *rooks*, and we will call the elements of \mathcal{R}_G the *symplectic rooks*. The *Bruhat-Chevalley-Renner order* on \mathcal{R}_n is defined by

$$\sigma \leq \tau \iff B_n \sigma B_n \subseteq \overline{B_n \tau B_n} \tag{1.1}$$

for $\sigma, \tau \in \mathcal{R}_n$. (We will introduce the most general form of the Bruhat-Chevalley-Renner order in the preliminaries section.) An explicit combinatorial description of \leq , in the spirit of Deodhar's criteria, is obtained in [8]. By using this explicit characterization of \leq , it is shown in [6] that (\mathcal{R}_n, \leq) is a graded, bounded, EL-shellable poset.

The first main observation in our paper, Theorem 3.9, states that, for $\sigma, \tau \in \mathcal{R}_G$, we have

$$\sigma \leq \tau \text{ in } \mathcal{R}_G \iff \sigma \leq \tau \text{ in } \mathcal{R}_n.$$

An important family of subposets of \mathcal{R}_n are defined as follows. Let k be an integer in $\{0, 1, \dots, n\}$, and let

$$\mathcal{B}_n(k) := \{\sigma \in \mathcal{R}_n : \sigma \text{ is upper triangular and } \text{rank}(\sigma) = k\}$$

and

$$\mathcal{B}_n := \bigsqcup_{k=0}^n \mathcal{B}_n(k).$$

Then \mathcal{B}_n parametrizes the $B_n \times B_n$ -orbits in the Borel monoid \overline{B}_n . Actually, \mathcal{B}_n is the lower interval $[0, 1] = \{x \in \mathcal{R}_n : x \leq 1\}$ in \mathcal{R}_n . Therefore, \mathcal{B}_n is also EL-shellable. Generalizing this observation, in [7], joint with Cherniavsky, the first author showed that each poset $(\mathcal{B}_n(k), \leq)$ ($k \in \{0, 1, \dots, n\}$) is a graded, bounded, EL-shellable poset. In fact, it turns out that $(\mathcal{B}_n(k), \leq)$ is a union of $\binom{n}{k}$ maximal subintervals all of which have the same minimum element. An important combinatorial aspect of this development is that, as a set, $\mathcal{B}_n(k)$ is in bijection with the set partitions of $\{1, \dots, n+1\}$ with k blocks. In particular, the cardinality $|\mathcal{B}_n(k)|$ is given by the Stirling numbers of the second kind, $S(n+1, k)$. In our second main result, we obtain similar results for the rank k elements of the Borel submonoid \overline{B} in MSp_n . We should mention that the type BC analogs of the set partitions with respect to "refinement order" is well known [19]. For a more recent study of their combinatorial properties, see [1].

We will denote by \mathcal{B}_G the submonoid of all upper triangular elements in the symplectic Renner monoid \mathcal{R}_G . In other words, $\mathcal{B}_G = \{x \in \mathcal{R}_G : x \leq 1\} = [0, 1]$ in \mathcal{R}_G . The k -th symplectic Stirling poset, denoted by $\mathcal{B}_G(k)$, is the subposet defined by

$$\mathcal{B}_G(k) := \{x \in \mathcal{B}_G : \text{rank}(x) = k\}. \tag{1.2}$$

In Theorem 4.6, we prove that the k -th symplectic Stirling poset is a graded bounded poset with unique minimum element, and with $\binom{l}{k} 2^k$ maximal elements, all of which are rank k diagonal idempotents. It is now a natural question to find the cardinality of each of the posets $\mathcal{B}_G(k)$. We answer this question in Theorem 5.13. It turns out that

$$|\mathcal{B}_G(k)| = \sum_{a+b+c=k} 2^{a+c} 3^b \binom{l}{b} S(l+1, l+1-a) S(l+1, l+1-c),$$

where $(a, b, c) \in \mathbb{Z}_{\geq 0}^3$. Here, for integers $s, t \in \mathbb{Z}$ such that $s \leq t$, $S(s, t)$ stands for the (s, t) -th Stirling numbers of the second kind.

Reductive monoids are regular in the semigroup theory sense. Geometrically, the only smooth reductive monoids with one-dimensional center are the monoids of $n \times n$ matrices [20]. A complex algebraic variety of dimension n is called *rationally smooth* if for every $x \in X$, the local cohomology groups $H^i(X, X \setminus \{x\})$ are zero for $i \neq 2n$, and $H^{2n}(X, X \setminus \{x\}) = \mathbb{Q}$. It turns out that the rationally smooth reductive monoids have rich combinatorial and geometric structures [22, 23]. Their classification has been completed by Renner [22, 24]. In particular, MSp_n is a rationally smooth monoid. Gonzales showed that the rationally smooth reductive monoids are GKM manifolds, see [12]. This means that the relevant (equivariant) cohomological data for such a monoid can be recovered from the knowledge of torus invariant points and curves alone.

In Theorem 6.3, we show that, if M is a rationally smooth reductive monoid with zero, and \overline{B} is a Borel submonoid in M , then \overline{B} is rationally smooth as well. Although we do not exploit this information here, we can now use Renner's H -polynomials for computing the intersection cohomology Poincaré polynomials of many Borel submonoids. In particular, this idea is applicable to the case of MSp_n . We plan to revisit this topic in a future paper.

We now describe the organization of our paper. In Section 2 we briefly summarize some basic properties of the symplectic groups and monoids. Section 3 is devoted to the study of the Bruhat-Chevalley-Renner order on MSp_n . In this section we prove our first result, Theorem 3.9. Empowered by the concrete description of the partial order, we begin our study of the Borel submonoid of MSp_n in Section 4. In particular, in this section, we give a count of the number of rank k elements of the symplectic upper triangular roots, Theorem 5.13. The purpose of Section 6 is to show that the Borel submonoids of rationally smooth reductive monoids with zeros are rationally smooth, Theorem 6.3. In the final part of our paper, we return to our study of the symplectic monoids. We observe that, unlike the case of the monoid of $n \times n$ matrices, the subsemigroup of nilpotent elements of the Borel submonoid of MSp_n is not irreducible for $n \geq 2$.

Acknowledgements. The authors thank Yonah Cherniavsky and Zhenheng Li. The first author is partially supported by a grant from the Louisiana Board of Regents.

2 Preliminaries

In this section we will review the basic ingredients of our objects.

2.1 Symplectic groups.

Let n be a positive integer of the form $n = 2l$ for some $l \in \mathbb{Z}$. Let us denote by H (resp. G) the special linear group SL_n (resp. the symplectic group Sp_n). Then $G \subseteq H$ with equality if $n = 2$. We will denote by T_H and B_H the maximal diagonal torus and the Borel subgroup consisting of upper triangular matrices in H , respectively. Then the intersections $T_G := G \cap T_H$ and $B_G := G \cap B_H$ are, respectively, the maximal diagonal torus and a Borel subgroup containing T_G in G .

Let $\theta : H \rightarrow H$ denote the following involutory automorphism:

$$\theta(A) = J(A^\top)^{-1}J^{-1} \quad A \in H.$$

Then the fixed subgroup of θ in H is G . In other words, we have $H^\theta = G$. Also, it is easy to verify that $B_G = B_H^\theta$ and that $T_G = T_H^\theta$. With this choice of T_H and B_H , we know that the normalizer of T_H in H , that is, $N_H(T_H)$ is equal to the $n \times n$ monomial matrices in H , and the elements of the Weyl group, $W_H := N_H(T_H)/T_H$, are represented by the permutation matrices of size n . We will denote W_H by S_n . The *one-line notation* of an element w of S_n is the sequence (w_1, \dots, w_n) , where $w_i = w(i)$ for $i \in \{1, \dots, n\}$. In this notation, the Weyl group of (G, T_G) has a convenient description as the fixed point subgroup of the induced involution, $\theta : S_n \rightarrow S_n$ which is defined by

$$\theta(w) := (n+1-w_n, n+1-w_{n-1}, \dots, n+1-w_1) \quad w \in S_n.$$

In other words, we have

$$W_G = \{w \in S_n : \theta(w) = w\}.$$

By working with the root system corresponding to (G, B_G, T_G) , one knows that (W_G, S_G) , as a Coxeter group, is generated by

$$S_G = \{r_i r_{n-i} : 1 \leq i \leq l-1\} \cup \{r_l\},$$

where r_j ($j \in \{1, \dots, n-1\}$) denotes the simple transposition $r_j = (j, j+1)$ in S_n . Let us define s_1, \dots, s_l by setting

$$s_j := \begin{cases} r_j r_{n-j} & \text{if } j \in \{1, \dots, l-1\}; \\ r_l & \text{if } j = l. \end{cases} \quad (2.1)$$

In this notation, the Coxeter-Dynkin diagram of (G, B_G, T_G) can be depicted as in Figure 2.1. This labeling is consistent with the labeling that is given in [4].

$$s_1 \text{ --- } s_2 \text{ --- } \dots \text{ --- } s_{l-1} \text{ === } s_l$$

Figure 2.1: The Coxeter-Dynkin diagram of type C_l .

2.2 Symplectic monoids.

Let M be a reductive monoid with unit group G . Then, by definition, G is a connected reductive group. Let B be a Borel subgroup in G , and let T be a maximal torus of G that is contained in B . Then the Weyl group of G is given by $W := N_G(T)/T$. The Bruhat-Chevalley decomposition of G is the finite decomposition $G = \bigsqcup_{w \in W} B\dot{w}B$. Likewise, the $B \times B$ -orbits in M are parametrized by a finite inverse semigroup, which is called the *Renner*

monoid of M ; it is defined as the quotient $R := \overline{N_G(T)}/T$, where $\overline{N_G(T)}$ denotes the Zariski closure of $N_G(T)$ in M . Then the *Bruhat-Chevalley-Renner decomposition* of M is given by

$$M = \bigsqcup_{r \in R} BrB.$$

The dot on r indicates that we are choosing a representative of r from $\overline{N_G(T)}$. In general, it is not true that R is a submonoid of M . An excellent survey of the Renner monoids of classical monoids can be found in [15].

Notation 2.2. Let G denote, as before, the symplectic group Sp_n . Then the Renner monoid of the symplectic monoid MSp_n will be denoted by \mathcal{R}_G . The Weyl group of G will be denoted by W_G .

1) Let θ denote the involution that we introduced before, that is, $\theta(i) = n - i + 1$ for $i \in \{1, \dots, n\}$. A subset $S \subseteq \{1, \dots, n\}$ is called an *admissible subset* if $\theta(S) \cap S = \emptyset$. For $i, j \in \{1, \dots, n\}$, let E_{ij} denote the (i, j) -th elementary matrix. Then the (k, l) -th entry of E_{ij} is 1 if $(k, l) = (i, j)$, and it is 0 if $(k, l) \neq (i, j)$. In [16, Theorem 3.1.8], it is shown that

$$\mathcal{R}_G = \left\{ \sum_{i \in I, w \in W_G} E_{i, wi} : I \text{ is admissible} \right\}. \quad (2.3)$$

2) An *injective partial transformation on $\{1, \dots, n\}$* is an injective map $f : D \rightarrow R$, where $D = D(f)$ and $R = R(f)$ are two subsets from $\{1, \dots, n\}$ with equal cardinalities.

Definition 2.4. The set of all injective partial transformations on $\{1, \dots, n\}$ is called the *rook monoid*; we will denote it by \mathcal{R}_n . As we mentioned before, \mathcal{R}_n is the Renner monoid of M_n .

In [16, Theorem 3.1.10], Li and Renner show that

$$\mathcal{R}_G = \{x \in \mathcal{R}_n : D(x) \text{ and } R(x) \text{ are admissible, and } x \text{ is singular}\} \cup W_G. \quad (2.5)$$

3) For $k \in \{0, 1, \dots, n\}$, let e_k denote the diagonal idempotent, $e_k := E_{11} + E_{22} + \dots + E_{kk}$. Also, let e_0 denote the $n \times n$ 0-matrix. The cross-section lattice of \mathcal{R}_G is then given by

$$\Lambda_G := \{e_0, e_1, e_2, \dots, e_l, e_n\}. \quad (2.6)$$

(Notice the jump in the indices of the last two idempotents e_l to e_n . This is not a typo!) In this notation, the Renner monoid of MSp_n is given by

$$\mathcal{R}_G = \bigsqcup_{e_i \in \Lambda} W_G e_i W_G. \quad (2.7)$$

Since e_n is the identity element, the subset $W_G e_n W_G$ is equal to W_G . Therefore, the rank of a singular element in \mathcal{R}_G is at most l .

3 The Bruhat-Chevalley-Renner order

Let M be a reductive monoid with unit group G . Let B be a Borel subgroup in G , and let T be a maximal torus in G such that $T \subseteq B$. Let us denote the Renner monoid of M by R . The *Bruhat-Chevalley-Renner order* on R is the following partial order: for $x, y \in R$,

$$x \leq_{BCR} y \iff BxB \subseteq \overline{ByB},$$

where the bar over ByB stands for the Zariski closure in M . Whenever it is clear from the context, we will omit writing the subscript BCR in \leq_{BCR} . Note that the restriction of \leq to W is known as the Bruhat-Chevalley order on W , which is defined by the same formulation,

$$x \leq y \iff BxB \subseteq \overline{ByB} \text{ for } x, y \in W,$$

where the bar over ByB stands for the Zariski closure in G .

The Weyl group W is a graded poset with the rank function $\ell_W : W \rightarrow \mathbb{Z}_{\geq 0}$ defined by

$$\ell_W(w) = \dim BwB - \dim B \text{ for } w \in W.$$

Note that W is a Coxeter group and it has a system of Coxeter generators, denoted by S . For $w \in W$, $\ell_W(w)$ can also be defined as the minimal number of simple reflections s_{i_1}, \dots, s_{i_r} from S with $w = s_{i_1} \cdots s_{i_r}$. A subgroup that is generated by a subset $I \subset S$ will be denoted by W_I and it will be called a *parabolic subgroup of W* . For $I \subseteq S$, we will denote by D_I the following set:

$$D_I := \{x \in W : \ell_W(xw) = \ell_W(x) + \ell_W(w) \text{ for all } w \in W_I\}. \quad (3.1)$$

The *type-map*, $\lambda : \Lambda \rightarrow 2^S$, is defined by $\lambda(e) := \{s \in S : se = es\}$ for $e \in \Lambda$. The containment ordering between $G \times G$ -orbit closures in M is transferred via λ to a sublattice of the Boolean lattice on S . Associated with $\lambda(e)$ are the following sets: $\lambda_*(e) := \cap_{f \leq e} \lambda(f)$ and $\lambda^*(e) := \cap_{f \geq e} \lambda(f)$. We define the subgroups

$$W(e) := W_{\lambda(e)}, \quad W_*(e) := W_{\lambda_*(e)}, \quad W^*(e) := W_{\lambda^*(e)}.$$

Then we have

1. $W(e) = \{a \in W : ae = ea\}$,
2. $W^*(e) = \cap_{f \geq e} W(f)$,
3. $W_*(e) = \cap_{f \leq e} W(f) = \{a \in W : ae = ea = e\}$.

We know from [18, Chapter 10] that $W(e)$, $W^*(e)$, and $W_*(e)$ are parabolic subgroups of W , and furthermore, we know that $W(e) \cong W^*(e) \times W_*(e)$. If $W(e) = W_I$ and $W_*(e) = W_K$ for some subsets $I, K \subset S$, then we define $D(e) := D_I$ and $D_*(e) := D_K$.

Theorem/Definition (Pennell-Putcha-Renner): For every $x \in WeW$ there exist elements $a \in D_*(e), b \in D(e)$, which are uniquely determined by x , such that

$$x = aeb^{-1}. \quad (3.2)$$

The decomposition of x in (3.2) will be called the *standard form of x* . Let e, f be two elements from Λ . It is proven in [17] that if $x = aeb^{-1}$ and $y = cfd^{-1}$ are two elements in standard form in R , then

$$x \leq y \iff e \leq f, a \leq cw, w^{-1}d^{-1} \leq b^{-1} \quad \text{for some } w \in W(f)W(e). \quad (3.3)$$

We will occasionally write $D(e)^{-1}$ to denote the set $\{b^{-1} : b \in D(e)\}$.

3.1 Deodhar's criteria.

Our goal in this section is to present a practical description of the Bruhat-Chevalley-Renner order on the rook monoid.

Let us denote by B_n the Borel subgroup of invertible upper triangular matrices in M_n . The Renner monoid \mathcal{R}_n is a graded poset with the following rank function [21]:

$$\ell(x) = \dim(B_n x B_n), \quad x \in \mathcal{R}_n.$$

There is a combinatorial formula for computing the values of ℓ [8].

We represent elements of \mathcal{R}_n by n -tuples. For $x = (x_{ij}) \in \mathcal{R}_n$ we define the sequence (x_1, \dots, x_n) by

$$x_j = \begin{cases} 0 & \text{if the } j\text{-th column consists of zeros,} \\ i & \text{if } x_{ij} = 1. \end{cases} \quad (3.4)$$

By abuse of notation, we denote both the matrix and the sequence (x_1, \dots, x_n) by x . For example, the associated sequence of the partial permutation matrix

$$x = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

is $x = (3, 0, 4, 0)$.

Next, we define a useful partial order on finite sets of integers. Let $\{i_1, \dots, i_k\}$ and $\{j_1, \dots, j_k\}$ be two equinumerous sets of integers such that $i_1 < \dots < i_k$ and $j_1 < \dots < j_k$. We will write

$$\{i_1, \dots, i_k\} \leq \{j_1, \dots, j_k\} \iff i_1 \leq j_1, i_2 \leq j_2, \dots, i_k \leq j_k. \quad (3.5)$$

Let $x = (x_1, \dots, x_n)$ be an element from \mathcal{R}_n . For $i \in \{1, \dots, n\}$, we define

$$\tilde{x}(i) := \{x_1, \dots, x_i\}.$$

In this notation, the main result of [8] is as follows.

Theorem 3.6. *Let $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ be two elements from \mathcal{R}_n . Then $x \leq_{BCR} y$ if and only if for every $i \in \{1, \dots, n-1\}$ we have $\tilde{x}(i) \leq \tilde{y}(i)$.*

Example 3.7. Let $x = (3, 1, 5, 2, 4)$ and $y = (5, 2, 4, 3, 1)$ be two elements from \mathcal{R}_5 . Since

$$\begin{aligned}\tilde{x}(1) &= \{3\} \leq \{5\} = \tilde{y}(1), \\ \tilde{x}(2) &= \{1, 3\} \leq \{2, 5\} = \tilde{y}(2), \\ \tilde{x}(3) &= \{1, 3, 5\} \leq \{2, 4, 5\} = \tilde{y}(3), \\ \tilde{x}(4) &= \{1, 2, 3, 5\} \leq \{2, 3, 4, 5\} = \tilde{y}(4),\end{aligned}$$

we see that $x \leq_{BCR} y$.

Theorem 3.6 is quite useful for explicit computations.

3.2 The Bruhat-Chevalley-Renner order on \mathcal{R}_G .

As before, let n be an even number, $n = 2l$, $l \in \mathbb{Z}_+$. Recall that W_G denotes the Weyl group of Sp_n . Then W_G is the centralizer in \mathcal{S}_n of the involution $\theta = (1, n)(2, n-1) \cdots (l, l+1)$. As a Coxeter group, W_G has type BC_l . In [2, Corollary 8.1.9], it is shown that for two elements u and v from W_G ,

$$u \leq v \text{ in } W_G \iff u \leq v \text{ in } W_H = \mathcal{S}_n. \quad (3.8)$$

We will extend (3.8) to the Renner monoid of MSp_n .

Theorem 3.9. *Let x any y be two elements from \mathcal{R}_G . Then*

$$x \leq y \text{ in } \mathcal{R}_G \iff x \leq y \text{ in } \mathcal{R}_n.$$

Proof. We write x and y in their standard form, $x = aeb^{-1}$ and $y = cfd^{-1}$, where $a \in D_*(e)$, $b \in D(e)$, $c \in D_*(f)$, $d \in D(f)$. Of course, a, b, c , and d are elements of W_G . By (3.3) we know that

$$x \leq y \text{ in } \mathcal{R}_G \iff e \leq f, a \leq cw, w^{-1}d^{-1} \leq b^{-1},$$

for some $w \in W_G(f)W_G(e)$. The idempotents of \mathcal{R}_G are in \mathcal{R}_n , hence, the Bruhat-Chevalley-Renner order on them is the one that is induced from \mathcal{R}_n . It follows from (3.8) that the relations $a \leq cw$ and $w^{-1}d^{-1} \leq b^{-1}$ hold in W_G if and only if they hold in $\mathcal{S}_n \subseteq \mathcal{R}_n$. Therefore, the relation $x \leq y$ holds in \mathcal{R}_G if and only if it holds in \mathcal{R}_n . \square

Corollary 3.10. *Let $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ be two elements from \mathcal{R}_G . Then $x \leq_{BCR} y$ if and only if for every $i \in \{1, \dots, n-1\}$ we have $\tilde{x}(i) \leq \tilde{y}(i)$.*

Proof. It follows from Theorem 3.9 that $x \leq_{BCR} y$ in \mathcal{R}_G if and only if $x \leq_{BCR} y$ in \mathcal{R}_n . The rest of the proof follows from Theorem 3.6. \square

4 The Borel Submonoid of MSp_n

We will follow our convention that if the even integer $n = 2l$ is fixed, then G stands for Sp_n . We will denote by B_G , as before, the Borel subgroup in Sp_n that is defined by

$$B_G := B_n \cap Sp_n.$$

The Borel subgroup of the unit group GSp_n of MSp_n is given by $B := \mathbb{C}^* B_G$, and the corresponding Borel submonoid of MSp_n is the Zariski closure of B in MSp_n . Evidently, B is a connected, hence irreducible, algebraic group. Then \overline{B} is an irreducible $B \times B$ -variety. The orbits of $B \times B$ are parametrized by $x \in \mathcal{R}_G$ such that $x \leq 1$. Indeed, we have

$$\mathcal{B}_G = \mathcal{B}_n \cap \mathcal{R}_G := \{x \in \mathcal{R}_G : x \leq 1\}.$$

We depict the Bruhat-Chevalley-Renner order on \mathcal{B}_{Sp_n} in Figure 4.1.

We are going to reformulate the description of \mathcal{B}_G in two different ways.

1. It is observed in [17, Lemma 2.3] that an element r from \mathcal{R}_G satisfies $r \leq 1$ if and only if $a \leq b$, where aeb^{-1} is the standard form of r . Thus, we have

$$\mathcal{B}_G = \{aeb^{-1} : aeb^{-1} \text{ is in standard form, } e \in \Lambda_G, a \in D_*(e), b \in D(e), \text{ and } a \leq b\}. \quad (4.1)$$

2. Let $D(x)$ (resp. $R(x)$) denote the domain (resp. range) of an element $x \in \mathcal{R}_n$. The data of $D(x)$ and $R(x)$ are not enough to recover x . One needs to know the (bijective) assignment between them to uniquely determine x . Let $D(x)$ and $R(x)$ be as follows:

$$D(x) = \{i_1, \dots, i_k\} \quad \text{and} \quad R(x) = \{j_1, \dots, j_k\},$$

where

$$x(i_t) = j_t \quad \text{for } t \in \{1, \dots, k\}.$$

We will assume that the entries of $D(x)$ are listed in the increasing order as in $1 \leq i_1 < \dots < i_k \leq n$, however, we may not have the same ordering on the corresponding elements of $R(x)$.

Notice that in order for x to be ≤ 1 in the Bruhat-Chevalley-Renner order its matrix representation has to have all of its nonzero entries on or above the main diagonal. Since $D(x)$ gives the column indices of the nonzero entries in x , and since $R(x)$ gives the row indices of the nonzero entries in x , we see that

$$x \leq 1 \iff i_t \geq j_t \quad \text{for every } t \in \{1, \dots, k\}.$$

By (2.5), for the elements x in $\mathcal{R}_G \setminus \{1\}$, both of the subsets $D(x), R(x) \subseteq \{1, \dots, n\}$ are admissible.




Figure 4.1: Bruhat-Chevalley-Renner order on \mathcal{B}_{Sp_4} .

Question 4.2. What is the cardinality of \mathcal{B}_G ? By the second item, our problem is equivalent to the question of finding, for every $k \in \{1, \dots, l\}$, the number of pairs of admissible subsets

$$I = \{i_1, \dots, i_k\} \quad \text{and} \quad J = \{j_1, \dots, j_k\}$$

such that

1. $1 \leq i_1 < \dots < i_k \leq n$ and $J \subset \{1, \dots, n\}$; there are no order constraints on the elements of J .
2. $i_t \geq j_t \geq 1$ for every $t \in \{1, \dots, k\}$.

It turns out that the number of admissible subsets of $\{1, \dots, n\}$ has a pleasant formula. We begin with a simple lemma.

Lemma 4.3. *Let n and k be two positive integers such that $1 \leq k \leq n$. Assume that n is an even number, $n = 2l$. Let $A_{n,k}$ denote the set of admissible subsets of $\{1, \dots, n\}$ with k elements. If $a_{n,k}$ denotes the cardinality of $A_{n,k}$, then*

$$a_{n,k} = \sum_{r=0}^k \binom{l}{r} \binom{l-r}{k-r} = \binom{l}{k} 2^k.$$

Proof. Clearly, by the pigeon-hole principle, if $k > l$, then $A_{n,k} = \emptyset$. Also, in this case, $\binom{l-r}{k-r}$ is 0 for every $r \in \{0, \dots, k\}$, hence, $a_{n,k} = 0$. Therefore, we will assume that $k \leq l$.

Let $A = \{i_1, \dots, i_k\}$ ($i_1 < \dots < i_k$) be an element from $A_{n,k}$. The entries of A satisfy the inequalities

$$1 \leq i_1 < \dots < i_r \leq l < i_{r+1} < \dots < i_k \leq n.$$

We will determine the number of such A . Clearly, the first r entries, i_1, \dots, i_r , can be chosen in $\binom{l}{r}$ ways. Then the remaining entries, i_{r+1}, \dots, i_k , cannot be contained in the set $\{\theta(i_s) = n - i_s + 1 : s \in \{1, \dots, r\}\}$. In other words, $\{i_{r+1}, \dots, i_k\} \subseteq \{l+1, \dots, 2l\} \setminus \{\theta(i_s) = n - i_s + 1 : s \in \{1, \dots, r\}\}$. Then, the number of possibilities for $\{i_{r+1}, \dots, i_k\}$ is given by $\binom{l-r}{k-r}$. Therefore, in total, we have $\sum_{r=0}^k \binom{l}{r} \binom{l-r}{k-r}$ possibilities for A . This finishes the proof of the first equality. To prove the second equality, we manipulate the summation as follows:

$$\sum_{r=0}^k \binom{l}{r} \binom{l-r}{k-r} = \sum_{r=0}^k \frac{l!}{(l-r)!r!} \frac{(l-r)!}{(l-k)!(k-r)!} = \sum_{r=0}^k \frac{l!}{r!(l-k)!(k-r)!}. \quad (4.4)$$

Let us multiply and divide each summand in the last sum in (4.4) by $k!$. Then by reorganizing the terms we get

$$\sum_{r=0}^k \frac{l!}{(l-k)!k!} \frac{k!}{r!(k-r)!} = \sum_{r=0}^k \binom{l}{k} \binom{k}{r} = \binom{l}{k} \sum_{r=0}^k \binom{k}{r} = \binom{l}{k} 2^k.$$

This finishes the proof. □

Since the empty set is admissible, we set $a_{n,0} = 1$.

Corollary 4.5. *The total number of admissible subsets of $\{1, \dots, n\}$, that is, $|\cup_{k=0}^l A_{n,k}|$, is equal to 3^l .*

Proof. We will determine the number $\sum_{k=0}^l a_{n,k} = \sum_{k=0}^l \binom{l}{k} 2^k$. But, by the binomial theorem, $f(2) = \sum_{k=0}^l \binom{l}{k} 2^k$, where $f(x) = (1+x)^l$. \square

Theorem 4.6. *The k -th symplectic Stirling poset $\mathcal{B}_G(k)$ is a graded bounded poset with a unique minimum element. There are $\binom{l}{k} 2^k$ maximal elements in $\mathcal{B}_G(k)$.*

Proof. If $k = n$ (resp. $k = 0$), then $\mathcal{B}_G(k) = \{id\}$ (resp. $\mathcal{B}_G(k) = \{\mathbf{0}\}$), hence, in these cases there is nothing to prove. We proceed with the assumption that $1 \leq k \leq l$. Since \mathcal{B}_G is equal to the intersection $\mathcal{R}_G \cap \mathcal{B}_n$, we have

$$\mathcal{B}_G(k) = \mathcal{B}_n(k) \cap \mathcal{R}_G \text{ for } k \in \{1, \dots, l\}.$$

Notice that the rook $id(k) := (0, \dots, 0, 1, 2, \dots, k)$ is a symplectic rook. In fact, $id(k)$ is the unique minimum of $\mathcal{B}_n(k)$. It follows from Theorem 3.9 that $id(k)$ is the unique minimum element in $\mathcal{B}_G(k)$ as well. Next, we will show that $\mathcal{B}_G(k)$ is a graded poset. To this end, it will suffice to show that every maximal element of $\mathcal{B}_G(k)$ has the same rank. In [7, Lemma 5.1], it is shown that the maximal elements of $\mathcal{B}_G(k)$ are given by the diagonal idempotents of rank k in \mathcal{R}_n . Clearly, any diagonal idempotent of rank k whose domain and range are admissible subsets in $\{1, \dots, n\}$ are contained in $\mathcal{B}_G(k)$. But for a diagonal matrix, the domain and the range agree, therefore, the number of diagonal idempotents in $\mathcal{B}_G(k)$ is equal to the number of admissible subsets in $\{1, \dots, n\}$. This number is equal to $\binom{l}{k} 2^k$ by Lemma 4.3. Next, we will show that there are no other maximal elements in $\mathcal{B}_G(k)$. Towards a contradiction, let $a = (a_1, \dots, a_n)$ be a maximal element in $\mathcal{B}_G(k)$ which is not a diagonal idempotent. Since a is an upper triangular rook, we know that $a_i \leq i$ for every $i \in \{1, \dots, n\}$. Let $i \in \{1, \dots, n\}$ be the smallest index such that $0 < a_i \not\leq i$. Let j denote a_i . Then we know that $a_j = 0$. Let b denote the rook matrix that is obtained by interchanging a_i and a_j . It is easy to verify that b is an element of $\mathcal{B}_G(k)$ such that $a < b$. This contradicts with our assumption that a is a maximal element. Hence, the proof of our theorem is finished. \square

Remark 4.7. For each $d \in \{1, \dots, l\}$, by (3.5), we have a very special poset structure on $A_{n,d}$. It is easily seen from [14, Section 6.1.1] that $(A_{n,d}, \leq)$ is isomorphic to the Bruhat-Chevalley order on the Grassmann variety G/P_d , where $G = Sp_n$ and P_d is the maximal parabolic subgroup corresponding to the set of simple generators $S_G \setminus \{s_d\}$ in W_G .

Proposition 4.8. *Let Λ_G be the cross section lattice of G as in (2.6). If e is an element from $\Lambda_G \setminus \{1\}$, then $W_G(e)$ is a maximal parabolic subgroup in W_G . Conversely, any maximal parabolic subgroup of W_G is obtained this way.*

Proof. The cross section lattice Λ_G is part of the cross section lattice of the monoid M_n . It is easy to verify that, if the matrix rank of e is d , then the centralizer of e in $W_H = S_n$ is the maximal parabolic subgroup generated by the set $\{r_1, \dots, r_{n-1}\} \setminus \{r_d\}$. Let s_1, \dots, s_l denote, as defined in (2.1), the simple Coxeter generators for W_G . Now it is easy to check that, for every $j \in \{1, \dots, l\} \setminus \{d\}$, we have

$$s_j e = e s_j, \quad \text{hence, } W_G(e) = \langle s_j : j \in \{1, \dots, l\} \setminus \{d\} \rangle.$$

Our second assertion is now easy to verify. This finishes the proof. \square

Next, we will compute the stabilizer of an element e from $\Lambda_G \setminus \{1\}$.

Proposition 4.9. *Let e be an element from $\Lambda_G \setminus \{1\}$. If the matrix rank of e is d , where $1 \leq d < l$, then $(W_G)_*(e)$ is generated by the simple Coxeter generators s_{d+1}, \dots, s_l . If $d = l$, then $(W_G)_*(e) = \{1\}$.*

Proof. Once again, the proof will follow from the corresponding computation that is performed in the rook monoid (the Renner monoid of M_n). In that case, by explicitly computing the matrix products $r_j e$ ($j \in \{1, \dots, n-1\}$), one sees that $r_j e = e r_j = e$ if and only if $j \in \{d+1, \dots, n-1\}$. It follows immediately from this observation that $s_j = r_j r_{n-j}$ stabilizes e if and only if $j \in \{d+1, \dots, l\}$. It also follows that if the rank of e is l , then s_l does not stabilize e , hence, $(W_G)_*(e) = \{1\}$. This finishes the proof. \square

5 Folding, unfolding

Let n be a positive integer. A collection S_1, \dots, S_r of subsets of the set $S := \{1, \dots, n\}$ is said to be a *set partition* of S if S_i 's ($i = 1, \dots, r$) are mutually disjoint and $\cup_{i=1}^r S_i = S$. In this case, the S_i 's are called the blocks of the partition. The collection of all set partitions of S is denoted by Π_n . We will often drop set parentheses and commas and just put vertical bars between blocks. If S_1, \dots, S_k are the blocks of a set partition π from Π_n , then the *standard form* of π is defined as $S_1|S_2|\dots|S_k$, where we assume that $\min S_1 < \dots < \min S_k$ and the elements of each block are listed in increasing order. For example, $\pi = 136|2459|78$ is a set partition from Π_9 . Set partitions can be visualized by using ‘‘arc-diagrams’’ which we will define next.

A linearly ordered poset is called a *chain*. We will identify chains by their Hasse diagrams; we draw a Hasse diagram by placing the smallest entry on the left and connecting the vertices by arcs. For example, in Figure 5.1, we have the chain on 9 vertices, where each arc represents a covering relation.



Figure 5.1: A chain on 9 vertices.

Definition 5.1. A *labeled chain* is a chain whose vertices are labeled by distinct numbers. An *arc-diagram on n vertices* is a disjoint union of labeled chains where the labels are from $\{1, \dots, n\}$ and each label $i \in \{1, \dots, n\}$ is used exactly once. We depict an example in Figure 5.2.




Figure 5.2: An arc-diagram on 9 vertices

Clearly, in an arc-diagram subchains represents the blocks of the corresponding set partition. We know from [3, Lemma 1.17] that the number of set partitions of $S = \{1, \dots, n\}$ into k blocks, denoted by $S(n, k)$, and called the (n, k) -th *Stirling number of the second kind*, is given by the formula $S(n, k) = \frac{1}{k!} \sum_{i=1}^k (-1)^i \binom{k}{i} (k-i)^n$. The recurrence formula for the Stirling numbers of the second kind is well-known:

$$S(l+1, k) = S(l, k-1) + kS(l, k),$$

where

$$S(l, k) = \begin{cases} 1 & \text{if } l = k = 0; \\ 0 & \text{if } l > 0 \text{ and } k = 0; \\ 0 & \text{if } l < 0 \text{ or } k < 0 \text{ or } l < k. \end{cases}$$

Let \mathcal{B}_n denote the submonoid of \mathcal{R}_n such that if $x \in \mathcal{B}_n$, then x is an upper triangular matrix. The subsemigroup of nilpotent elements in \mathcal{B}_n will be denoted by $\mathcal{B}_n^{\text{nil}}$. For each A in \mathcal{B}_n , there exists a unique $(n+1) \times (n+1)$ nilpotent matrix, \tilde{A} , which is obtained from A by appending to it a column and a row of zeros as follows:

$$A \mapsto \tilde{A} := \begin{bmatrix} 0 & & & \\ \vdots & A & & \\ 0 & \dots & 0 & \end{bmatrix} \in \mathcal{B}_{n+1} \quad (A \in \mathcal{B}_n). \quad (5.2)$$

In this notation, it is easily verified that (5.2) defines a set bijection $\mathcal{B}_n \longrightarrow \mathcal{B}_{n+1}^{\text{nil}}$. There is a simple bijection between $\mathcal{B}_{n+1}^{\text{nil}}$ and Π_{n+1} which is defined as follows: the matrix corresponding to the set partition A has an entry equal to 1 in row i and column j if and only if (i, j) is an arc of A . Therefore, for $k \in \{1, \dots, n+1\}$, we have

$$S(n+1, k) = |\{A \in \mathcal{B}_n : \text{rank} A = n+1-k\}|. \quad (5.3)$$

It follows from the bijections above that the number of elements of \mathcal{B}_n is given by the summation $b_{n+1} := \sum_{k=0}^{n+1} S(n+1, k)$, which is called the $(n+1)$ th Bell number. As a convention, we set $b_0 = 1$ and $b_1 = 1$.

It is easy to check that the number of elements of \mathcal{R}_n of rank k is given by

$$|\{A \in \mathcal{R}_n : \text{rank}(A) = k\}| = \binom{n}{k} \frac{n!}{(n-k)!}. \quad (5.4)$$

We will express this cardinality by using Stirling numbers of the second kind.

Every element A of \mathcal{R}_n has a triangular decomposition in \mathcal{R}_n ,

$$A = A_l + A_d + A_u, \quad (5.5)$$

where A_l is a strictly lower triangular matrix, A_d is a diagonal matrix, and A_u is a strictly upper triangular matrix.

Proposition 5.6. *Let $S_{a,b,c}(n)$ denote the number of elements $A \in \mathcal{R}_n$ such that $\text{rank}(A_l) = a$, $\text{rank}(A_d) = b$, and $\text{rank}(A_u) = c$, where A_l, A_d , and A_u are as in (5.5). Then we have*

$$\begin{aligned} \binom{n}{k} \frac{n!}{(n-k)!} &= \sum_{a+b+c=k} S_{a,b,c}(n) \\ &= \sum_{a+b+c=k} \binom{n}{b} S(n+1, n+1-a) S(n+1, n+1-c). \end{aligned}$$

Proof. The number of strictly upper triangular elements of rank k in \mathcal{R}_n is equal to the number of strictly lower triangular rank k elements in \mathcal{R}_n . Now the proof of the first equality follows from the equality in (5.4) and the uniqueness of the triangular decomposition in (5.5). The proof of the second equality follows from (5.3) together with the fact that there are exactly $\binom{n}{b}$ diagonal 0/1 matrices of rank b . \square

We proceed with the assumption that n is an even number of the form $n = 2l$, $l \in \mathbb{Z}_+$. In the sequel, we will count the number of elements of \mathcal{B}_G , where $G = Sp_n$, by a technique that we call *unfolding*. But before that we want to demonstrate that the elements of \mathcal{R}_G behave well under “folding”. We already mentioned the result of Li and Renner [16, Theorem 3.1.10], which states that, if an element A from \mathcal{R}_G is singular, then both of the domain and the range of A are admissible subsets in $\{1, \dots, n\}$. Furthermore, the elements of $D(A)$ correspond to the indices of the nonzero columns of A , and the elements of $R(A)$ correspond to the indices of the nonzero rows of A . This shows that A can be folded vertically as well as horizontally. We will demonstrate what we mean here by an example.

Example 5.7. In this example, we fold an element of \mathcal{R}_{Sp_8} horizontally from top to bottom:

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \longrightarrow \begin{bmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Example 5.8. In this example, we fold the matrix of the previous example vertically from left to right:

$$\begin{bmatrix} 1 & 0 & 0 & 0 & \hline 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \longrightarrow \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Definition 5.9. We will denote the horizontal folding operation from top to bottom by F_{TB} . Likewise, we will denote the vertical folding operation from left to right by F_{LR} .

Clearly, the operations F_{TB} and F_{LR} can be composed. In fact, they commute,

$$F_{TB}F_{LR} = F_{LR}F_{TB}. \quad (5.10)$$

Let F denote the composition of the folding operators as in (5.10). We will refer to F by the *folding map*.

Proposition 5.11. *The folding map is a surjective map from \mathcal{R}_{Sp_n} onto the rook monoid \mathcal{R}_l . Furthermore, the restricted folding map, $F|_{\mathcal{B}_G}$, which we will denote by F' , is surjective as well.*

Proof. If we show that F' is surjective, then the surjectivity of F will follow. To this end, let A be an element from \mathcal{R}_l , and let $A = A_l + A_d + A_u$ be its triangular decomposition. Recall from the introduction that J_l denotes the $l \times l$ permutation matrix with 1's on its anti-diagonal. Now we define an $n \times n$ matrix B as follows: $B := \begin{bmatrix} \mathbf{0} & \tilde{A}_l \\ \mathbf{0} & A_d + A_u \end{bmatrix}$, where $\mathbf{0}$ is the $l \times l$ 0 matrix, and $\tilde{A}_l := J_l A_l$. In other words, \tilde{A}_l is the matrix whose i th row is the $(l - i + 1)$ th row of A_l . Since the indices of the nonzero columns of B are contained in the set

$\{l+1, \dots, n\}$, the domain of B is an admissible subset in $\{1, \dots, n\}$. It is also easy to check that the set of row indices of B is an admissible subset of $\{1, \dots, n\}$. Clearly, B is upper triangular matrix, therefore, $B \in \mathcal{B}_G$. Finally, by its construction, the image of B under F' is equal to A , $F'(B) = F(B) = A$. This finishes the proofs of our assertions. \square

We are now ready to count the number of elements of \mathcal{B}_G by “unfolding” the elements of \mathcal{R}_l first horizontally from bottom to top, and then vertically from right to left. We will demonstrate our count by an example.

Example 5.12. We will compute the preimage of $J_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ under the restricted folding map $F' : \mathcal{B}_{Sp_4} \rightarrow \mathcal{R}_2$. Equivalently, we will determine the set $F_{LR}^{-1}(F_{TB}^{-1}(J_2)) \cap \mathcal{B}_{Sp_4}$. Since we are looking for the upper triangular elements in the preimage, the lower halves of the 4×2 matrices in $F_{TB}^{-1}(J_2)$ must be upper triangular. The following matrices are the possibilities:

$$\begin{array}{ccc}
 \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ \hline 0 & 1 \\ 0 & 0 \end{bmatrix} & & \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ \hline 0 & 0 \\ 0 & 0 \end{bmatrix} \\
 & \searrow^{F_{TB}} \quad \swarrow_{F_{TB}} & \\
 & \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} &
 \end{array}$$

Let A_1 denote the 4×2 matrix that is on the top-left position, and let A_2 denote the 4×2 matrix that is on the top-right position. Then, the following two matrices are mapped onto A_1 by F_{LR} :

$$\begin{bmatrix} 0 & 0 & \color{red}{|} & 1 & 0 \\ 0 & 0 & \color{red}{|} & 0 & 0 \\ 0 & 0 & \color{red}{|} & 0 & 1 \\ 0 & 0 & \color{red}{|} & 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 & \color{red}{|} & 0 & 0 \\ 0 & 0 & \color{red}{|} & 0 & 0 \\ 0 & 0 & \color{red}{|} & 0 & 1 \\ 0 & 0 & \color{red}{|} & 0 & 0 \end{bmatrix} \xrightarrow{F_{LR}} \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$$

Likewise, the following two matrices are folded onto A_2 by F_{LR} :

$$\begin{bmatrix} 0 & 1 & \color{red}{|} & 0 & 0 \\ 0 & 0 & \color{red}{|} & 0 & 1 \\ 0 & 0 & \color{red}{|} & 0 & 0 \\ 0 & 0 & \color{red}{|} & 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 & \color{red}{|} & 1 & 0 \\ 0 & 0 & \color{red}{|} & 0 & 1 \\ 0 & 0 & \color{red}{|} & 0 & 0 \\ 0 & 0 & \color{red}{|} & 0 & 0 \end{bmatrix} \xrightarrow{F_{LR}} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$$

We find that, in total, there are four matrices that fold onto J_2 .

We are now ready to present a formula for the number of elements of \mathcal{B}_G that lie in the preimage of the folding map F' .

Theorem 5.13. *The number of elements of rank k in \mathcal{B}_G is given by the formula*

$$\sum_{a+b+c=k} 2^{a+c} 3^b \binom{l}{b} S(l+1, l+1-a) S(l+1, l+1-c), \quad (5.14)$$

where $(a, b, c) \in \mathbb{Z}_{\geq 0}^3$.

Proof. Let A be an element from \mathcal{R}_l with the triangular decomposition $A = A_l + A_d + A_u$. Then it is easy to verify that

$$|F'^{-1}(A) \cap \mathcal{B}_{S_{p_n}}| = |F'^{-1}(A_l) \cap \mathcal{B}_{S_{p_n}}| |F'^{-1}(A_d) \cap \mathcal{B}_{S_{p_n}}| |F'^{-1}(A_u) \cap \mathcal{B}_{S_{p_n}}|. \quad (5.15)$$

Let us denote the matrix ranks of $A_l, A_d,$ and A_u by $a, b,$ and $c,$ respectively. Then we denote the three factors on the right hand side of (5.15) by the notation $f_a(A), f_b(A),$ and $f_u(A),$ respectively. Our choice of the subscripts for f_a, f_b, f_c will be clarified in the next two paragraphs.

As it was shown for the special case of J_2 in Example 5.12, if B is an element from $F'^{-1}(A),$ then the lower $l \times l$ half of the $2l \times l$ matrix $F_{LR}(B)$ must be an upper triangular matrix. In other words, when we unfold A to a $2l \times l$ matrix, all of the nonzero entries of A_l are moved into the upper $l \times l$ portion of the resulting matrix, hence, there is a unique $2l \times l$ matrix A'_l such that $F_{TB}(A'_l) = A_l.$ Moreover, for every subset of the nonzero entries of $A'_l,$ there exists a unique A''_l in \mathcal{B}_G such that $F_{LR}(A''_l) = A_l.$ It follows from these arguments that

$$f_a(A) = |F_{LR}^{-1}(A'_l)| = \binom{a}{0} + \binom{a}{1} + \cdots + \binom{a}{a} = 2^a. \quad (5.16)$$

A similar argument shows that

$$f_c(A) = \binom{c}{0} + \binom{c}{1} + \cdots + \binom{c}{c} = 2^c. \quad (5.17)$$

We now consider the possible unfolding of the diagonal matrix $A_d.$ Recall that the rank of A_d is $b.$ For every s element subset of the set of nonzero entries of $A_d,$ there exists a unique $2l \times l$ matrix A'_d such that $F_{TB}(A'_d) = A_d$ and $\text{rank}(A'_d) = s.$ Likewise, for every r element subset of the set of nonzero entries of $A'_d,$ there exists a unique $2l \times 2l$ matrix A''_d such that $F_{LR}(A''_d) = A'_d$ and $\text{rank}(A''_d) = r.$ In total, there exist $\sum_{s=0}^b \sum_{r=0}^s \binom{s}{r} \binom{b}{s}$ elements in the preimage $F'^{-1}(A_d).$ But this double sum has a closed form:

$$f_b(A) = \sum_{s=0}^b \sum_{r=0}^s \binom{s}{r} \binom{b}{s} = 3^b. \quad (5.18)$$

By combining (5.16), (5.17), and (5.18), we find that $|F'^{-1}(A) \cap \mathcal{B}_{S_{p_n}}| = 2^a 2^c 3^b = 2^{a+c} 3^b,$ which actually depends only on the ranks of the matrices $A_l, A_d,$ and $A_c.$ Our formula now follows from Proposition 5.6. \square

Remark 5.19. It is easy to check that $\mathcal{B}_{Sp_n}(1)$ is equal to $\mathcal{B}_n(1)$. The Hasse diagram of $(\mathcal{B}_n(1), \leq)$ is a fishnet, see [7, Figure 1.9]. By Theorem 3.9, we know that, for every $k \in \{1, \dots, l\}$, $(\mathcal{B}_{Sp_n}(k), \leq)$ is a subposet of $(\mathcal{B}_n(k), \leq)$. However, unless $k = 1$, the inclusion map $\mathcal{B}_{Sp_n}(k) \hookrightarrow \mathcal{B}_n(k)$ does not preserve the interval structure. Indeed, already for $k = 2$, the cardinalities of $\mathcal{B}_{Sp_4}(2)$ and $\mathcal{B}_4(2)$ are different.

6 Rationally Smooth Borel Submonoids

Let M be a reductive monoid with zero. Let G denote its unit group, which is a connected reductive group. Then M is called a *semisimple monoid* if G has a one-dimensional center. The classification of semisimple (smooth) monoids is due to Renner [20]. It turns out that a semisimple monoid M is smooth if and only if M is isomorphic to the monoid of $n \times n$ matrices, M_n , for some $n \in \mathbb{Z}_+$. Note that the situation for general reductive monoids is not very different; a reductive monoid with zero is smooth if and only if M is of the form

$$M = \left(G_0 \times \prod_{i=1}^r M_{n_i} \right) / Z,$$

where Z is a finite central torus that does not intersect the unit group of $\prod_{i=1}^r M_{n_i}$, and G_0 is a semisimple subgroup [25, Section 11]. The semisimple monoids whose cohomological properties are as good as one hopes for are identified by Renner also [22]. They are called “rationally smooth” monoids.

Let X be a complex algebraic variety with $\dim X = n$, and let x be a point in X . The variety X is called *rationally smooth at x* if there exists an open neighborhood U of x such that for all $y \in U$, the following holds:

$$H^m(X, X \setminus \{y\}) = \begin{cases} \{0\} & \text{if } m \neq 2n; \\ \mathbb{Q} & \text{if } m = 2n. \end{cases}$$

X is called *rationally smooth* if it is rationally smooth at every point x in X . The classification as well as various characterizations of rationally smooth reductive monoids is given in [22, 24].

We will now adapt another result of Renner [24, Theorem 2.2] in our setting.

Lemma 6.1. *Let X and Y be two (normal) Borel submonoids of the (normal) reductive monoids M and N , respectively. Assume that both of M and N have zero elements, and assume that there exists a finite dominant morphism of algebraic monoids $g : M \rightarrow N$. Under these assumptions, X is rationally smooth if and only if Y is rationally smooth.*

Proof. By abuse of notation, we will denote the restriction $g|_X$ by g also. By our assumptions, the algebraic monoids X and Y have zero elements, denoted by 0_X and 0_Y , respectively. Let B denote the Borel subgroup contained in X , and let T be a maximal torus contained in B . Then 0_X is the unique closed $B \times B$ orbit in X , hence, $X \setminus \{0\}$ is (rationally) smooth. Let $\mu : \mathbb{C}^* \times M \rightarrow M$ be a central (in B) one-parameter subgroup action on M such

that $\lim_{t \rightarrow 0} \mu(t, x) = 0_X$ for every $x \in M$. Then the quotients $\mathbb{P}_X := (X \setminus \{0_X\})/\mathbb{C}^*$ and $\mathbb{P}_M := (M \setminus \{0_M\})/\mathbb{C}^*$ are projective $T \times T$ varieties such that $\mathbb{P}_X \subsetneq \mathbb{P}_M$. Furthermore, \mathbb{P}_X and \mathbb{P}_M are rationally smooth. Similarly, we have the rationally smooth, projective $T' \times T'$ varieties $\mathbb{P}_Y \subsetneq \mathbb{P}_N$, where T' is the maximal torus in $g(T) \subset Y$.

By result of Brion [5, Lemma 1.3] we know that X (resp. Y) is rationally smooth if and only if the Euler characteristic of \mathbb{P}_X (resp. the Euler characteristic of \mathbb{P}_Y) is equal to the number of $T \times T$ fixed points in \mathbb{P}_X (resp. the number of $T' \times T'$ fixed points in \mathbb{P}_Y). Let us denote by $C(M)$ (resp. by $C(N)$) the closed $G \times G$ -orbit in \mathbb{P}_M (resp. the closed $G' \times G'$ -orbit in \mathbb{P}_N), where G (resp. G') is the unit group of M (resp. of N). Since the $T \times T$ fixed points in \mathbb{P}_X lie in the closed intersection $\mathbb{P}_X \cap C(M)$, and since $g|_{C(M)} : C(M) \rightarrow C(N)$ is a bijection, we see that the Euler characteristics of X and Y are equal. In particular, X is rationally smooth if and only if Y is rationally smooth. \square

Two reductive monoids M and N are said to be *equivalent* if there exists a reductive monoid L with two finite dominant morphisms $L \rightarrow M$ and $L \rightarrow N$. If M and N are equivalent monoids, then we will write $M \sim_0 N$. It is easy to verify that \sim_0 is an equivalence relation. Let M be a reductive monoid with zero. According to [24, Theorem 2.4],

$$M \text{ is rationally smooth} \iff M \sim_0 \prod_{i=1}^r M_{n_i}. \quad (6.2)$$

We are now ready to prove the main result of this section.

Theorem 6.3. *Let M be a rationally smooth reductive monoid with zero. Let B be a Borel subgroup in M and let \overline{B} denote the corresponding Borel submonoid. Then \overline{B} is a rationally algebraic semigroup.*

Proof. Since M is rationally smooth, we know from (6.2) that there exists a reductive monoid L admitting two finite dominant morphisms:

$$\begin{array}{ccc} & L & \\ f \swarrow & & \searrow g \\ M & & \prod_{i=1}^r M_{n_i} \end{array}$$

Without loss of generality, we may assume that L has a zero. Let B_L be a Borel subgroup of L . As f and g are finite and dominant morphisms, they are surjective. In particular, the subgroups $f(B_L)$ and $g(B_L)$ are Borel subgroups in M and $\prod_{i=1}^r M_{n_i}$, respectively. We set $X := \overline{f(B_L)}$ and $Y := \overline{g(B_L)}$. Then X and Y are Borel submonoids in M and $\prod_{i=1}^r M_{n_i}$, respectively. Since Y is (rationally) smooth, by Lemma 6.1, so is $\overline{B_L}$. Once again by using Lemma 6.1, we see that X is rationally smooth. This finishes the proof of our theorem. \square

As an application of Theorem 6.3, we consider the symplectic monoid MSp_n . By the Renner's classification of rationally smooth simple group embeddings [23, Corollary 3.5], MSp_n is a rationally smooth semisimple monoid. Therefore, by Theorem 6.3, its Borel submonoid is rationally smooth.

7 Final Remarks: Nilpotent Subsemigroups

In this section we will contrast some properties of the Borel submonoids of M_n and MSp_n . We begin with a general observation.

Lemma 7.1. *Let M be a reductive monoid with the Bruhat-Chevalley-Renner decomposition $M = \sqcup_{r \in R} B\dot{r}B$, where B is a Borel subgroup in M , and R is the Renner monoid of M . If an element r from R satisfies the following two properties, then every element of the orbit $B\dot{r}B$, where \dot{r} is a representative of r in $\overline{N_G(T)}$, is nilpotent:*

- (1) r is nilpotent in R , that is, $r^m = 0$ for some $m \in \mathbb{Z}_+$;
- (2) $r \preceq 1$.

Proof. Since M is a linear algebraic monoid, it admits an embedding into M_n for a sufficiently large positive integer n . By conjugating with an element of GL_n , we assume that B is contained in the upper triangular Borel submonoid of M_n . Clearly, if we can prove our assertion for $M = M_n$ and $B = B_n$, then the general case will follow. In this case, the Renner monoid is given by the rook monoid \mathcal{R}_n , and we can identify \mathcal{R}_n as a submonoid of M_n . An element r from \mathcal{R}_n satisfies the two properties in our hypotheses if and only if it is a strictly upper triangular rook. But the product of an upper triangular matrix with a strictly upper triangular matrix is strictly upper triangular. Therefore, any element of $B\dot{r}B$ is strictly upper triangular, hence, nilpotent. This finishes the proof of our assertion. \square

We should note that we cannot replace any of the two requirements in Lemma 7.1. Indeed, if r is not nilpotent, then any of its representatives \dot{r} , which is contained in $B\dot{r}B$, is not nilpotent. For the second item, we consider the matrices $r = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$ and $b_2 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$.

Then, we have $rb_2 = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}$, which is not nilpotent. Evidently, the set of nilpotent elements in a Borel submonoid is a closed subset.

Definition 7.2. The subvariety $\overline{B}^{nil} := \{x \in \overline{B} : x^m = 0 \text{ for some } m \in \mathbb{Z}_+\}$ will be called the *nilpotent semigroup* of \overline{B} .

Corollary 7.3. *Let B be a Borel subgroup in a reductive monoid M with zero. Then the nilpotent semigroup of \overline{B} is a $B \times B$ -stable algebraic subsemigroup of M .*

Proof. By definition, \overline{B}^{nil} is defined by the polynomial relations $x^m = 0$ ($m \in \mathbb{Z}_+$), therefore, it is a closed subset of \overline{B} . By Lemma 7.1, we know that \overline{B}^{nil} is $B \times B$ -stable. In fact, the proof of this lemma shows that \overline{B}^{nil} is a semigroup. \square

Next, we will show that \overline{B}_n^{nil} is an irreducible variety.

Proposition 7.4. *The nilpotent semigroup of M_n is an irreducible algebraic semigroup of dimension $\binom{n}{2}$.*

Proof. By Lemma 7.1, we know that $\overline{B_n}^{nil} = \bigsqcup B_n r B_n$, where the union is over all strictly upper triangular rooks in \mathcal{R}_n . It is easy to check that

1. $r_0 := (0, 1, 2, \dots, n-1)$ is a strictly upper triangular rook;
2. if r is a strictly upper triangular rook, then $r \leq r_0$.

These two conditions imply that $\overline{B_n r_0 B_n} = \overline{B_n}^{nil}$. It is easy to check that $\ell(r_0) = 1 + 2 + \dots + (n-1) = \binom{n}{2}$. Since the orbit $B_n r_0 B_n$ is an irreducible variety, so is $\overline{B_n}^{nil}$. Thus, in light of Corollary 7.3, the proofs of our assertions are finished. \square

Unfortunately the nice situation as in Proposition 7.4 does not hold for the nilpotent semigroup of $\overline{B_{Sp_n}}$. It turns out that $\overline{B_{Sp_n}}^{nil}$ has many irreducible components in varying dimensions. See Figure 7.1 for an example.




Figure 7.1: The Hasse diagram of $\overline{B_{Sp_4}}^{nil}$.

References

- [1] Eli Bagno, Riccardo Biagioli, and David Garber. *Some identities involving second kind Stirling numbers of types B and D* . *Electron. J. Combin.*, 26(3):Paper 3.9, 20, 2019.
- [2] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter groups*, volume 231 of *Graduate Texts in Mathematics*. Springer, New York, 2005.
- [3] Miklós Bóna. *Combinatorics of permutations*. Discrete Mathematics and its Applications (Boca Raton). CRC Press, Boca Raton, FL, second edition, 2012. With a foreword by Richard Stanley.
- [4] Nicolas Bourbaki. *Lie groups and Lie algebras. Chapters 4–6*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2002. Translated from the 1968 French original by Andrew Pressley.
- [5] M. Brion. *Rational smoothness and fixed points of torus actions*. *Transform. Groups*, 4(2-3):127–156, 1999. Dedicated to the memory of Claude Chevalley.
- [6] Mahir Bilen Can. *The rook monoid is lexicographically shellable*. *European J. Combin.*, 81:265–275, 2019.
- [7] Mahir Bilen Can and Yonah Cherniavsky. *Stirling Posets*. *Israel Journal of Mathematics*, 2020. <https://doi.org/10.1007/s11856-020-2004-1>.
- [8] Mahir Bilen Can and Lex E. Renner. *Bruhat-Chevalley order on the rook monoid*. *Turkish J. Math.*, 36(4):499–519, 2012.
- [9] You’an Cao, Jie Lei, and Zhenheng Li. *The symplectic monoid*. *Comm. Algebra*, 42(12):5425–5453, 2014.
- [10] Stephen Doty. *Polynomial representations, algebraic monoids, and Schur algebras of classical type*. *J. Pure Appl. Algebra*, 123(1-3):165–199, 1998.
- [11] Stephen Doty. *Representation theory of reductive normal algebraic monoids*. *Trans. Amer. Math. Soc.*, 351(6):2539–2551, 1999.
- [12] Richard Gonzales. *Rational smoothness, cellular decompositions and GKM theory*. *Geom. Topol.*, 18(1):291–326, 2014.
- [13] D. Ju. Grigor’ev. An analogue of the Bruhat decomposition for the closure of the cone of a classical Chevalley group series. *Dokl. Akad. Nauk SSSR*, 257(5):1040–1044, 1981.
- [14] Venkatramani Lakshmibai and Komaranapuram N. Raghavan. *Standard monomial theory*, volume 137 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2008. Invariant theoretic approach, Invariant Theory and Algebraic Transformation Groups, 8.

- [15] Zhenheng Li, Zhuo Li, and You'an Cao. [Algebraic monoids and Renner monoids](#). In *Algebraic monoids, group embeddings, and algebraic combinatorics*, volume 71 of *Fields Inst. Commun.*, pages 141–187. Springer, New York, 2014.
- [16] Zhenheng Li and Lex E. Renner. [The Renner monoids and cell decompositions of the symplectic algebra](#). *Internat. J. Algebra Comput.*, 13(2):111–132, 2003.
- [17] Edwin A. Pennell, Mohan S. Putcha, and Lex E. Renner. [Analogue of the Bruhat-Chevalley order for reductive monoids](#). *J. Algebra*, 196(2):339–368, 1997.
- [18] Mohan S. Putcha. *Linear algebraic monoids*, volume 133 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1988.
- [19] Victor Reiner. [Non-crossing partitions for classical reflection groups](#). *Discrete Math.*, 177(1-3):195–222, 1997.
- [20] Lex E. Renner. [Classification of semisimple algebraic monoids](#). *Trans. Amer. Math. Soc.*, 292(1):193–223, 1985.
- [21] Lex E. Renner. [Analogue of the Bruhat decomposition for algebraic monoids](#). *J. Algebra*, 101(2):303–338, 1986.
- [22] Lex E. Renner. [The \$H\$ -polynomial of a semisimple monoid](#). *J. Algebra*, 319(1):360–376, 2008.
- [23] Lex E. Renner. [Descent systems for Bruhat posets](#). *J. Algebraic Combin.*, 29(4):413–435, 2009.
- [24] Lex E. Renner. [Rationally smooth algebraic monoids](#). *Semigroup Forum*, 78(3):384–395, 2009.
- [25] D. A. Timashëv. [Equivariant compactifications of reductive groups](#). *Mat. Sb.*, 194(4):119–146, 2003.