

Accounting for the large radial tension in Morris-Thorne wormholes

Peter K.F. Kuhfittig
 Department of Mathematics
 Milwaukee School of Engineering
 Milwaukee, Wisconsin 53202-3109, USA

It is well known that a Morris-Thorne wormhole can only be held open by violating the null energy condition, physically realizable by the use of “exotic matter.” Unfortunately, even a small or moderately-sized wormhole would have a radial tension equal to that of the interior of a massive neutron star. So outside a neutron-star setting, such an outcome is problematical at best, calling for more than an appeal to exotic matter whose introduction had a completely different objective and with possibly different outcomes. The purpose of this paper is to account for the enormous radial tension in three ways: (1) directly invoking noncommutative geometry, an offshoot of string theory, (2) appealing to noncommutative geometry in conjunction with $f(R)$ modified gravity, and (3) determining the possible effect of a small extra spatial dimension.

I. INTRODUCTION

Wormholes are handles or tunnels in spacetime connecting widely separated regions of our Universe or even different universes. In many ways, wormholes are as good a prediction of Einstein’s theory as black holes, but they are subject to some severe restrictions.

Morris and Thorne [1] proposed the following static and spherically symmetric line element for a wormhole space-time:

$$ds^2 = -e^{2\Phi(r)} dt^2 + \frac{dr^2}{1 - b(r)/r} + r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (1)$$

using units in which $c = G = 1$. Here $\Phi = \Phi(r)$ is called the *redshift function*, which must be everywhere finite to prevent an event horizon. The function $b = b(r)$ is called the *shape function* since it determines the spatial shape of the wormhole when viewed, for example, in an embedding diagram. For the shape function we must have $b(r_0) = r_0$, where $r = r_0$ is the radius of the *throat* of the wormhole, and $\lim_{r \rightarrow \infty} b(r)/r = 0$. Another important requirement is the *flare-out condition* at the throat: $b'(r_0) < 1$, while $b(r) < r$ near the throat. The flare-out condition can only be met by violating the null energy condition (NEC)

$$T_{\alpha\beta} k^\alpha k^\beta \geq 0 \quad (2)$$

for all null vectors k^α , where $T_{\alpha\beta}$ is the stress-energy tensor. Matter that violates this condition is referred to as “exotic” in Ref. [1]. In particular, for the radial outgoing null vector $(1, 1, 0, 0)$, the violation reads $T_{\alpha\beta} k^\alpha k^\beta = \rho + p_r < 0$. Here $T^t_t = -\rho(r)$ is the energy density, $T^r_r = p_r(r)$ the radial pressure, and $T^\theta_\theta = T^\phi_\phi = p_t(r)$ the lateral pressure.

Next, let us state the Einstein field equations:

$$\rho(r) = \frac{b'}{8\pi r^2}, \quad (3)$$

$$p_r(r) = \frac{1}{8\pi} \left[-\frac{b}{r^3} + 2 \left(1 - \frac{b}{r} \right) \frac{\Phi'}{r} \right], \quad (4)$$

and

$$p_t(r) = \frac{1}{8\pi} \left(1 - \frac{b}{r} \right) \left[\Phi'' - \frac{b'r - b}{2r(r - b)} \Phi' + (\Phi')^2 + \frac{\Phi'}{r} - \frac{b'r - b}{2r^2(r - b)} \right]. \quad (5)$$

We also need to recall that the radial tension $\tau(r)$ is the negative of the radial pressure $p_r(r)$.

Having established the need for exotic matter, Morris and Thorne go on to state that the Einstein field equations can be rearranged to yield τ : reintroducing c and G for now, this is given by

$$\tau(r) = \frac{b(r)/r - 2[r - b(r)]\Phi'(r)}{8\pi Gc^{-4}r^2}. \quad (6)$$

From this condition it follows that the radial tension at the throat is

$$\tau = \frac{1}{8\pi Gc^{-4}r_0^2} \approx 5 \times 10^{41} \frac{\text{dyn}}{\text{cm}^2} \left(\frac{10 \text{ m}}{r_0} \right)^2. \quad (7)$$

In particular, for $r_0 = 3$ km, τ has the same magnitude as the pressure at the center of a massive neutron star. Such a situation has already been considered. According to Ref. [2], the enormous pressure near the center could cause a wormhole to form, provided that there is a core of quark matter.

To see how these ideas fit together, let us return to the null vector $(1, 1, 0, 0)$ and observe that $\rho + p_r < 0$ can be written $\tau - \rho > 0$. In the nongeometrized units of Eq. (6), the inequality becomes $\tau > \rho c^2$, possibly leading to an enormous radial tension at the throat. On the other hand, if ρ is extremely small, then τ can be quite small, as well. An even more revealing case, the zero-density solution corresponding to a constant shape function [see Eq. (3)], is discussed by Visser [3]. Similarly, if r_0 is extremely large, then, according to Eq. (6), τ may again be small. So while exotic matter ensures the violation of the NEC, a large radial tension is not a defining property. In particular, if we are not dealing with the interior of a neutron star, then the excessive radial tension becomes hard to explain. The usual, often unstated, assumption is that the extreme conditions are due to exotic matter. This assumption now takes on the appearance of merely giving the mystery a name.

The purpose of the paper is to account for the enormous radial tension in three ways: (1) directly invoking noncommutative geometry, itself an offshoot of string theory, (2) appealing to noncommutative geometry in conjunction with $f(R)$ modified gravity, and (3) analyzing the effect of a small extra spatial dimension.

II. NONCOMMUTATIVE GEOMETRY

There is no doubt that string theory has become ever more influential, as exemplified by the realization that coordinates may become noncommutative operators on a D -brane [4, 5]. The result, noncommutative geometry, can help eliminate the divergences that normally occur in general relativity. The reason is that noncommutativity replaces point-like objects by smeared objects, as a result of which spacetime can be encoded in the commutator $[\mathbf{x}^\mu, \mathbf{x}^\nu] = i\theta^{\mu\nu}$, where $\theta^{\mu\nu}$ is an antisymmetric matrix that determines the fundamental cell discretization of spacetime in the same way that Planck's constant \hbar discretizes phase space [6].

A natural way to model the smearing effect is by means of a Gaussian distribution of minimal length $\sqrt{\beta}$ instead of the Dirac delta function [6–10]. An equally effective way, discussed in Refs. [11–13], is to assume that the energy density of the static and spherically symmetric and particle-like gravitational source has the form

$$\rho_\beta(r) = \frac{\mu_1 \sqrt{\beta}}{\pi^2 (r^2 + \beta)^2}, \quad (8)$$

where μ_1 is a constant. Eq. (8) can be interpreted to mean that the gravitational source causes the mass μ_1 of a particle to be diffused throughout a region of linear dimension $\sqrt{\beta}$ due to the uncertainty; so $\sqrt{\beta}$ has units of length. Following Ref. [6], Eq. (8) likewise leads to the mass distribution

$$m(r) = \int_0^r 4\pi(r')^2 \rho(r') dr' = \frac{2M}{\pi} \left(\tan^{-1} \frac{r}{\sqrt{\beta}} - \frac{r\sqrt{\beta}}{r^2 + \beta} \right), \quad (9)$$

where M is now the total mass of the source. Observe that $m(0) = 0$ but $m(r)$ rapidly rises to M as r increases.

According to Ref. [6], noncommutative geometry is an intrinsic property of spacetime and does not depend on any particular features such as curvature. Moreover, the relationship between the radial pressure and energy density is given by

$$p_r = -\rho_\beta. \quad (10)$$

The reason is that the source is a self-gravitating droplet of anisotropic fluid of density ρ_β and the radial pressure is needed to prevent a collapse to a matter point.

In Ref. [6], the line element describing a black hole is obtained by solving the Einstein field equations with the Gaussian distribution as a matter source. Momentarily returning to Eq. (8), by using this form instead of the Gaussian distribution, the line element describing a black hole becomes

$$ds^2 = - \left[1 - \frac{(4M/\pi) \left(\tan^{-1} \frac{r}{\sqrt{\beta}} - \frac{r\sqrt{\beta}}{r^2 + \beta} \right)}{r} \right] dt^2 + \left[1 - \frac{(4M/\pi) \left(\tan^{-1} \frac{r}{\sqrt{\beta}} - \frac{r\sqrt{\beta}}{r^2 + \beta} \right)}{r} \right]^{-1} dr^2 + r^2 (d\theta^2 + \sin^2\theta d\phi^2). \quad (11)$$

As $\beta \rightarrow 0$, we recover the Schwarzschild line element.

Regarding the event horizon, due to the uncertainty, we are now dealing with a smeared surface instead of a smeared particle. These ideas carry over to wormholes in the sense that the throat also becomes a smeared surface.

III. NONCOMMUTATIVE-GEOMETRY INSPIRED WORMHOLES

A. The shape function

Since we are now concerned with the effects of noncommutative geometry, we have to return to Ref. [6] to discuss the length scales. It is stated that, at first glance, there appears to be a need to modify the 4D Einstein action to incorporate the noncommutative effects. We are dealing here with a very small scale which is definitely needed to account for the large radial tension at the throat of any moderately-sized Morris-Thorne wormhole. At the same time, however, we are dealing with a geometric structure over an underlying manifold which affects gravity in a subtle and indirect way: noncommutativity influences the mass-energy and momentum distribution and propagation [14–16], while the energy-momentum density determines the spacetime curvature. It is concluded in Ref. [6] that in general relativity, the effects of noncommutativity can be taken into account by keeping the standard form of the Einstein tensor on the left-hand side of the field equations and introducing the modified energy-momentum tensor as a source on the right-hand side. This leads to the conclusion, also discussed in Ref. [17], that the length scales need not be microscopic, which, in turn, allows us to obtain the shape function by means of Eqs. (3) and (8), but it does not prevent us from examining the region near the throat. The shape function is given by

$$b(r) = \frac{8M\sqrt{\beta}}{\pi} \int_{r_0}^r \frac{(r')^2 dr'}{[(r')^2 + \beta]^2} + r_0 = \frac{4M\sqrt{\beta}}{\pi} \left(\frac{1}{\sqrt{\beta}} \tan^{-1} \frac{r}{\sqrt{\beta}} - \frac{r}{r^2 + \beta} - \frac{1}{\sqrt{\beta}} \tan^{-1} \frac{r_0}{\sqrt{\beta}} + \frac{r_0}{r_0^2 + \beta} \right) + r_0. \quad (12)$$

According to Ref. [18], $b = b(r)$ has the usual properties of a shape function: $b(r_0) = r_0$, $0 < b'(r_0) < 1$, and $\lim_{r \rightarrow \infty} b(r)/r = 0$.

Based on the above discussion, we are allowed to use Eq. (8) to obtain a wormhole solution on a macroscopic scale. The fact remains, however, that $\rho_\beta(r)$ in Eq. (8) describes a particle-like gravitational source near the origin. To study the energy density farther away from the origin, some additional assumptions will have to be made. That is the topic of the next subsection.

B. The throat surface

As noted in Sec. II, we assume that the throat $r = r_0$ is a smeared surface. The best way to describe such a surface is to imagine that the smeared particle at $r = 0$ is replaced by a set of smeared particles on the surface, which is precisely what causes the surface to be smeared. Now let us consider the same outward radial direction for the particle and surface, as shown in Fig. 1. The two labeled line segments were drawn with the same length and direction to show the correspondence between the particle at $r = 0$ and the particle on the surface. As a consequence, it is reasonable to assume that the energy density ρ_s of the smeared surface is a simple translation of Eq. (8) in the r -direction:

$$\rho_s = \frac{\mu_2 \sqrt{\beta}}{\pi^2 [(r - r_0)^2 + \beta]^2}, \quad (13)$$

where μ_2 is the mass of the surface. Moreover, according to Ref. [6], we are dealing here with a kind of quantum pressure, i.e., an outward push induced by noncommuting coordinate quantum fluctuations, which allow us to go from




FIG. 1: Section of a sphere of radius $r = r_0$.

point-particle to surface. We also need to recall that r_0 is a relatively moderate throat radius, such as the 3-km radius discussed in Sec. I. Finally, Eq. (10) carries over to the surface, as long as we confine ourselves to the outward radial direction.

Since the tension τ is the negative of p_r , the violation of the NEC, $p_r + \rho_s < 0$, now becomes $\tau - \rho_s > 0$ at the throat. The condition $p_r = -\rho_s$ implies that we are right on the edge of violating the NEC, i.e., $\tau - \rho_s = 0$. In particular, when $r = r_0$,

$$\rho_s = \frac{\mu_2}{\pi^2} \frac{1}{\beta^{3/2}}, \quad (14)$$

but we still have $\tau - \rho_s = 0$. However, since the throat is a smeared surface, we can only assert that $r \approx r_0$. So by Eq. (13), ρ_s is reduced in value and we obtain the desired

$$\tau - \rho_s > 0. \quad (15)$$

Finally, Eq. (14) implies that ρ_s and hence τ are extremely large near the throat. This is our main result, to be confirmed later by other means.

To check its plausibility, let us return to Sec. I, Eq. (7), and recall that for a throat size of 10 m, $\tau \approx 5 \times 10^{41}$ dyn/cm². Suppose μ_2 has the rather minute value 10^{-10} g. Applying Eq. (14), we have

$$\tau = \rho_s c^2 = \frac{\mu_2}{\pi^2} (\sqrt{\beta})^{-3} c^2 = 5 \times 10^{41} \frac{\text{dyn}}{\text{cm}^2}. \quad (16)$$

To satisfy this relationship, the value $\sqrt{\beta} = 10^{-11}$ cm is sufficient. Since $\sqrt{\beta}$ may be much smaller, we can accommodate even larger values of τ .

IV. CONNECTION TO $f(R)$ MODIFIED GRAVITY

Modified gravitational theories, including $f(R)$ modified gravity, have been invoked to explain various phenomena in general relativity. In this section we present a heuristic argument to show that $f(R)$ modified gravity cannot provide a satisfactory explanation for the extreme radial tension at the throat without appealing to noncommutative geometry.

First we need to recall that in $f(R)$ gravity, the Ricci scalar R in the Einstein-Hilbert action $S_{\text{EH}} = \int \sqrt{-g} R dx^4$ is replaced by a nonlinear function $f(R)$. Next, let us state the gravitational field equations in the form used by Lobo and Oliveira [19]:

$$\rho(r) = F(r) \frac{b'(r)}{r^2}, \quad (17)$$

$$p_r(r) = -F(r)\frac{b(r)}{r^3} + F'(r)\frac{rb'(r) - b(r)}{2r^2} - F''(r)\left[1 - \frac{b(r)}{r}\right], \quad (18)$$

and

$$p_t(r) = -\frac{F'(r)}{r}\left[1 - \frac{b(r)}{r}\right] + \frac{F(r)}{2r^3}[b(r) - rb'(r)]; \quad (19)$$

here $F = \frac{df}{dR}$. It is also assumed that $\Phi(r) \equiv \text{constant}$, so that $\Phi'(r) \equiv 0$. Otherwise, according to Ref. [19], the analysis becomes intractable. Fortunately, the redshift function does not have to be specified in our discussion.

Returning to the field equations, at $r = r_0$, we have

$$\rho(r_0) = F(r_0)\frac{b'(r_0)}{r_0^2} \quad (20)$$

and

$$\tau(r_0) = -p_r(r_0) = F(r_0)\frac{b(r_0)}{r_0^3} - F'(r_0)\frac{r_0b'(r_0) - b(r_0)}{2r_0^2}. \quad (21)$$

To produce a large radial tension, it is sufficient for $F'(r_0)$ to be large and positive. $F(r_0)$ itself can be quite small and still meet the flare-out condition $b'(r_0) = r_0^2\rho(r_0)/F(r_0) < 1$. The result is a large tension, while retaining the small $\rho(r_0)$.

The real question is whether the positive conclusion from $f(R)$ gravity can be considered satisfactory. While $F(r_0)$ does not need to meet any special requirements, the very large $F'(r_0)$ demanded seems rather artificial, even unphysical. Fortunately, we can still fall back on our previous noncommutative-geometry theory.

To that end, we need to refer to Ref. [20], which discusses the connection between noncommutative geometry and $f(R)$ modified gravity. According to Ref. [19], the Ricci scalar R is given by $R = 2b'(r)/r$. So in the vicinity of the throat, $R \approx 2\alpha/r^2$, for $\alpha < 1$. To be consistent with the notation in Ref. [20], we will assume the following form for the Ricci scalar:

$$R(r) = \frac{2\alpha}{r^2}, \quad (22)$$

where α is a constant. It is shown in Ref. [20] that the noncommutative-geometry background yields the form

$$f(R) = \frac{2M_{r_0}}{\pi^2\beta^{3/2}} \ln\left(1 + \frac{\beta R}{2\alpha}\right), \quad (23)$$

where M_{r_0} is the mass of a spherical shell of radius $r = r_0$, which we can take to be the surface of the throat. Given that $\beta R/2\alpha = \beta/r^2 < 1$, the Maclaurin expansion now yields

$$f(R) \approx \frac{2M_{r_0}}{\pi^2\beta^{3/2}} \left[\frac{\beta R}{2\alpha} - \frac{1}{2} \left(\frac{\beta R}{2\alpha} \right)^2 \right] < \frac{2M_{r_0}}{\pi^2\beta^{3/2}} \frac{\beta R}{2\alpha} = \frac{M_{r_0}}{\pi^2\alpha\sqrt{\beta}} R. \quad (24)$$

So

$$f'(R) < \frac{M_{r_0}}{\pi^2\alpha\sqrt{\beta}}. \quad (25)$$

This result agrees with the more convenient assumption that $f(R)$ has the approximate form

$$f(R) = \frac{M_{r_0}}{\pi^2\alpha\sqrt{\beta}} R^{1-\varepsilon}, \quad \varepsilon \ll 1. \quad (26)$$

To show this, observe that for $\varepsilon = 0$, $(1 - \varepsilon)R^{-\varepsilon} = 1$, while $\frac{\partial}{\partial \varepsilon}(1 - \varepsilon)R^{-\varepsilon} < 0$. So for $\varepsilon > 0$, $(1 - \varepsilon)R^{-\varepsilon} < 1$. We now have

$$f'(R) = \frac{M_{r_0}}{\pi^2\alpha\sqrt{\beta}}(1 - \varepsilon)R^{-\varepsilon} < \frac{M_{r_0}}{\pi^2\alpha\sqrt{\beta}}, \quad (27)$$

in agreement with Inequality (25).

Combining these observations, we have

$$F(R) = F = \frac{df}{dR} = \frac{M_{r_0}}{\pi^2 \alpha \sqrt{\beta}} (1 - \varepsilon) R^{-\varepsilon},$$

$$\frac{dF}{dR} = \frac{M_{r_0}}{\pi^2 \alpha \sqrt{\beta}} (1 - \varepsilon) (-\varepsilon) R^{-\varepsilon-1},$$

and

$$\frac{dR}{dr} = -\frac{4\alpha}{r^3}.$$

The result is

$$\frac{dF}{dr} = \frac{dF}{dR} \frac{dR}{dr} = \frac{M_{r_0}}{\pi^2 \alpha \sqrt{\beta}} (1 - \varepsilon) (-\varepsilon) R^{-\varepsilon-1} \left(-\frac{4\alpha}{r^3} \right) > 0$$

near the throat $r = r_0$. So dF/dr is large as long as $\sqrt{\beta} \ll M_{r_0}$. We conclude that $F'(r_0)$ can indeed be large and positive due to the noncommutative-geometry background.

The connection between noncommutative geometry and $f(R)$ modified gravity can be viewed from a broader perspective. From Eq. (17) with $b'(r) = 2\alpha$,

$$F(r) = \frac{r^2}{b'(r)} \rho(r) = \frac{r^2}{2\alpha} \rho(r) = \frac{1}{2\alpha/r^2} \frac{\mu_1 \sqrt{\beta}}{\pi^2 (r^2 + \beta)^2}, \quad (28)$$

while $r(R) = \sqrt{2\alpha/R}$ from Eq. (22). Substituting in Eq. (28), we get

$$F(R) = \frac{\mu_1 \sqrt{\beta}}{\pi^2} \frac{1}{R \left(\frac{2\alpha}{R} + \beta \right)^2}. \quad (29)$$

Since $F = \frac{df}{dR}$, Eq. (29) yields

$$f(R) = \int_0^R \frac{\mu_1 \sqrt{\beta}}{\pi^2} \frac{1}{R' \left(\frac{2\alpha}{R'} + \beta \right)^2} dR' = \frac{\mu_1 \sqrt{\beta}}{\pi^2} \frac{(\beta R + 2\alpha) \ln(\beta R + 2\alpha) - \beta R}{\beta^2 (\beta R + 2\alpha)} + C. \quad (30)$$

Lobo and Oliveira [19] go on to show that the matter threading the wormhole satisfies the null energy condition, while the effective energy-momentum tensor arising from the modified gravity theory leads to a violation, thereby supporting the wormhole.

The result is of interest to us because Eq. (30) is obtained from noncommutative geometry and therefore provides a motivation for the actual choice of the function $f(R)$.

V. THE EXTRA SPATIAL DIMENSION

In discussing an extra spatial dimension, we are going to follow Ref. [21], which starts with a generic line element in Schwarzschild coordinates:

$$ds^2 = -e^{2\Phi(r)} dt^2 + e^{2\lambda(r)} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (31)$$

Here it is assumed that from symmetry considerations, the fifth component has the form $e^{2\mu(r,l)} dl^2$, where l is the extra coordinate. In a wormhole setting, we naturally assume that $e^{2\lambda(r)} = 1 - b(r)/r$, resulting in the form

$$ds^2 = -e^{2\Phi(r)} dt^2 + \frac{dr^2}{1 - b(r)/r} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) + e^{2\mu(r,l)} dl^2, \quad (32)$$

again assuming that $c = G = 1$.

Ref. [21] is concerned primarily with the null energy condition. As already noted, the purpose of this paper is to account for the extreme radial tension at the throat.

Since Ref. [21] uses an orthonormal frame, we need to introduce the following notations for the components of the stress-energy tensor: $T_{00} = \rho$, the energy density, $T_{11} = p_r$, the radial pressure, and $\tau = -p_r$, the radial tension. Still following Ref. [21], we choose an orthonormal basis $\{e_{\hat{a}}\}$ which is dual to the following 1-form basis:

$$\theta^0 = e^{\Phi(r)} dt, \quad \theta^1 = [1 - b(r)/r]^{-1/2} dr, \quad \theta^2 = r d\theta, \quad \theta^3 = r \sin \theta d\phi, \quad \theta^4 = e^{\mu(r,l)} dl. \quad (33)$$

The resulting 1- and 2-forms and the derivations of the Riemann curvature and Ricci tensors given in Ref. [21]. For convenience, let us restate the components of the Ricci tensor in the orthonormal frame:

$$R_{00} = -\frac{1}{2} \frac{d\Phi(r)}{dr} \frac{rb' - b}{r^2} + \frac{d^2\Phi(r)}{dr^2} \left(1 - \frac{b}{r}\right) + \left[\frac{d\Phi(r)}{dr}\right]^2 \left(1 - \frac{b}{r}\right) + \frac{2}{r} \frac{d\Phi(r)}{dr} \left(1 - \frac{b}{r}\right) + \frac{d\Phi(r)}{dr} \frac{\partial\mu(r,l)}{\partial r} \left(1 - \frac{b}{r}\right), \quad (34)$$

$$R_{11} = \frac{1}{2} \frac{d\Phi(r)}{dr} \frac{rb' - b}{r^2} - \frac{d^2\Phi(r)}{dr^2} \left(1 - \frac{b}{r}\right) - \left[\frac{d\Phi(r)}{dr}\right]^2 \left(1 - \frac{b}{r}\right) + \frac{rb' - b}{r^3} - \frac{\partial^2\mu(r,l)}{\partial r^2} \left(1 - \frac{b}{r}\right) + \frac{1}{2} \frac{\partial\mu(r,l)}{\partial r} \frac{rb' - b}{r^2} - \left[\frac{\partial\mu(r,l)}{\partial r}\right]^2 \left(1 - \frac{b}{r}\right), \quad (35)$$

$$R_{22} = R_{33} = -\frac{1}{r} \frac{d\Phi(r)}{dr} \left(1 - \frac{b}{r}\right) + \frac{1}{2} \frac{rb' - b}{r^3} + \frac{b}{r^3} - \frac{1}{r} \frac{\partial\mu(r,l)}{\partial r} \left(1 - \frac{b}{r}\right), \quad (36)$$

$$R_{44} = -\frac{d\Phi(r)}{dr} \frac{\partial\mu(r,l)}{\partial r} \left(1 - \frac{b}{r}\right) - \frac{\partial^2\mu(r,l)}{\partial r^2} \left(1 - \frac{b}{r}\right) + \frac{1}{2} \frac{\partial\mu(r,l)}{\partial r} \frac{rb' - b}{r^2} - \left[\frac{\partial\mu(r,l)}{\partial r}\right]^2 \left(1 - \frac{b}{r}\right) - \frac{2}{r} \frac{\partial\mu(r,l)}{\partial r} \left(1 - \frac{b}{r}\right). \quad (37)$$

Since we are primarily interested in the radial tension and pressure at the throat, where $b(r_0) = r_0$, we may assume that $1 - b/r = 0$ in all these expressions.

To study the radial tension $\tau(r) = -p_r(r) = -T_{11}$, we start with the Einstein field equations in the orthonormal frame:

$$G_{\alpha\beta} = R_{\alpha\beta} - \frac{1}{2} R g_{\alpha\beta} = 8\pi T_{\alpha\beta}; \quad (38)$$

thus

$$g_{\alpha\beta} = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}. \quad (39)$$

From $G_{11} = 8\pi T_{11}$ we obtain

$$8\pi p_r(r) = R_{11} - \frac{1}{2} R g_{11}, \quad (40)$$

where the Ricci scalar is given by

$$R = R^i_i = -R_{00} + R_{11} + R_{22} + R_{33} + R_{44}. \quad (41)$$

It follows that

$$16\pi p_r(r) = R_{00} + R_{11} - R_{22} - R_{33} - R_{44} \quad (42)$$

and, as can be readily checked,

$$16\pi p_r(r) = -\frac{1}{2r^2} \frac{\partial \mu(r, l)}{\partial r} [rb'(r) - b(r)] - \frac{2b(r)}{r^3}. \quad (43)$$

We conclude that if $\partial \mu(r, l)/\partial r < 0$, Eq. (43) is a tension whose magnitude depends on $\partial \mu(r, l)/\partial r$. In principle, then, $\tau(r)$ can be extremely large.

Our final task is to show that the model can be made consistent with observation. More precisely, we wish the extra dimension to be extremely small or even curled up, as in string theory. To avoid a needless complication, we would also like the extra dimension to be essentially fixed from the four-dimensional perspective. For this purpose, the form chosen, $e^{2\mu(r, l)} dl^2$, is particularly convenient: assume that $\mu(r, l)$ is negative and large in absolute value. This causes the proper distance $e^{\mu(r, l)} dl$ and hence the fifth component to be extremely small, even negligible; the outcome is thereby consistent with observation. Now consider

$$\frac{\partial}{\partial r} e^{\mu(r, l)} = e^{\mu(r, l)} \frac{\partial \mu(r, l)}{\partial r}, \quad (44)$$

we already know that $|\partial \mu(r, l)/\partial r|$ can be extremely large. At the same time, $e^{\mu(r, l)}$ can be extremely small; in fact, given the above assumptions, $e^{\mu(r, l)}$ can be close to Planck size. Observationally, then,

$$\frac{\partial}{\partial r} e^{\mu(r, l)} \approx 0. \quad (45)$$

Since there is no discernable change in the r -direction, the fifth dimension is essentially fixed from the four-dimensional perspective.

VI. CONCLUSION

This paper starts by recalling that a Morris-Thorne wormhole can only be held open by the use of exotic matter whose sole function is to ensure that the NEC is violated. We have seen, however, that even a moderately-sized wormhole with $r_0 = 3$ km has a radial tension equal to that of the interior of a massive neutron star. Attributing this outcome to exotic matter merely gives the mystery a name since exotic matter was introduced for a completely different reason and with possibly different outcomes. The purpose of this paper is to account for the large radial tension in three ways: (1) directly appealing to noncommutative geometry, itself an offshoot of string theory, (2) invoking noncommutative geometry in conjunction with $f(R)$ modified gravity, and (3) determining the possible effect of a small extra spatial dimension. All three models help confirm that whenever we are dealing with this extreme regime, the effects of string theory cannot be neglected [22].

-
- [1] M.S. Morris, K.S. Thorne, Am. J. Phys. **56**, 395 (1988).
 - [2] P.K.F. Kuhfittig, Adv. Math. Phys. **2013** 630196 (2012).
 - [3] M. Visser, Lorentzian Wormholes: From Einstein to Hawking, American Institute of Physics, New York, 1995) page 146.
 - [4] E. Witten, Nucl. Phys. B **460** 335 (1996).
 - [5] N. Seiberg, E. Witten, J. High Energy Phys. **9909**, 032 (1999).
 - [6] P. Nicolini, A. Smailagic, E. Spallucci, Phys. Lett. B **632**, 547 (2006).
 - [7] P. Nicolini, E. Spallucci, Class. Quant. Grav. **27**, 015010 (2010).
 - [8] M. Rinaldi, Class. Quant. Grav. **28**, 105022 (2011).
 - [9] F. Rahaman, P.K.F. Kuhfittig, S. Ray, S. Islam, Phys. Rev. D **86**, 106101 (2012).
 - [10] P.K.F. Kuhfittig, Int. J. Pure Appl. Math. **89**, 401 (2013).
 - [11] J. Liang, B. Liu, Europhys. Lett. **100**, 30001 (2012).
 - [12] K. Nozari, S.H. Mehdipour, Class. Quant. Grav. **25** 175015 (2008).
 - [13] P.K.F. Kuhfittig, Adv. High Energy Phys. **2012**, 462493 (2012).
 - [14] A. Smailagic, E. Spallucci, Phys. Rev. D **65**, 107701 (2002).
 - [15] A. Smailagic, E. Spallucci, J. Phys. A **36**, L-467 (2003).

- [16] A. Smailagic, E. Spallucci, *J. Phys. A* **36**, L-517 (2003).
- [17] S. Rani, A. Jawad, *Adv. High Energy Phys.* **2016**, 7815242 (2016).
- [18] P.K.F. Kuhfittig, *Int. J. Mod. Phys. D* **24**, 1550023 (2015).
- [19] F.S.N. Lobo, M.A. Oliveira, *Phys. Rev. D* **80** 104012 (2009).
- [20] P.K.F. Kuhfittig, *New Horizons in Mathematical Physics (NHMP)* **2**, 62 (2018).
- [21] P.K.F. Kuhfittig, *Phys. Rev. D* **98**, 064041 (2018).
- [22] L. Susskind, *Phys. Rev. Lett.* **71**, 2367 (1993).