

Dark energy as generalised superfluid excitations

Elías Castellanos^{1,*} and Celia Escamilla-Rivera^{2,†}

¹Mesoamerican Centre for Theoretical Physics,

Universidad Autónoma de Chiapas, Carretera Zapata Km. 4,

Real del Bosque (Terán), Tuxtla Gutiérrez 29040, Chiapas, México.

²Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México,
Circuito Exterior C.U., A.P. 70-543, México D.F. 04510, México.

In this paper we present a generic form of the cosmological Equation of State derived from the formalism of Statistical Mechanics, assuming that a cosmic acceleration scenario can be interpreted as a system of quasi-particles. By considering a generalised superfluidity approach, in which the energy spectrum can be modulated via a parameter s , a negative equation of state arises as excitations associated with an *exotic* superfluid system. We show that for $s \sim 0.293$, $w \simeq -1$, which can be related to the standard equation of state for the Λ CDM model.

PACS numbers: 98.80.-k, 98.80.Es, 05.20.-y, 03.75.Kk

FROM STATISTICAL MECHANICS TO STANDARD FLRW COSMOLOGY

The observed dynamics of our universe is often described in terms of the FLRW metric. Even though, the adoption of this convenient geometry in Einstein's equations does not give information about the *specific* form for the Equation of State (EoS) of the effective cosmological fluid, $P = w\rho_E$. Here, the total energy density in the universe is denoted by ρ_E and the pressure by P . The standard concordance model, Λ CDM, fits in this scenario where the fluid can be identified into matter (including baryonic and cold dark matter) ρ_m , radiation ρ_r , and a not so well understood dark energy ρ_Λ . Towards this direction, many attempts have been done starting from the simple relation above described by a barotropic EoS (notice that here we call w as the EoS parameter) to some complex ones with an explicit relation between P and ρ_E . All of them to achieve the current cosmic acceleration. As it is standard, to obtain this behaviour we require an energy density with significant negative pressure at late times. This means that the evolution ratio between the pressure and energy density is negative, i.e. $w(z) = P/\rho_E < 0$. All reasonable fitting dark energy models available in the literature are in agreement at this point [1–6].

The evolution Friedmann equation for a spatially flat universe

$$\left(\frac{H(z)}{H_0}\right)^2 \propto [\Omega_m(1+z)^3 + \Omega_\Lambda f(z)], \quad (1)$$

where H_0 is the Hubble parameter and the index m denotes the matter components related to radiation, barionic and cold dark matter. From this equation it is possible to compute the EoS, $w(z)$. This can be done when the current value for the dark energy density is written as $\rho_{0(\Lambda)} = \rho_\Lambda(z)f^{-1}(z)$, with

$f(z) = \exp\left[3\int_0^z \frac{1+w(\tilde{z})}{1+\tilde{z}} d\tilde{z}\right]$. By modelling $w(z)$, we can give directly an entire evolution description of Eq. (1), as e.g., in the case of quintessence models $w = \text{constant}$, the solution for $f(z)$ is $f(z) = (1+z)^{3(1+w)}$.

If we consider the case of the Cosmological Constant Λ ($w = -1$) we obtain $f = 1$. Other cases explore a dark energy density ρ_Λ with varying and non-varying $w(z)$ [7–10], just to cite a few.

Notice that for a given functional form of $f(z)$, the contribution of the dark energy density to $H(z)$, goes to more negative values of $w(z)$. This is an impact in the evolution of dark energy on the dynamical age of the universe. As we mentioned, to get a dark energy model with late-time negative pressure we can think in two frameworks: on one hand, a quintessence model which shows a wide application in tracker the slow roll condition of scalar fields¹ and demands a constant EoS [11], a references therein. In numerical terms, according to Planck 2018 [12], the dark energy EoS parameter for a flat universe is $w = -1.006 \pm 0.045$, which is consistent with Λ . On the other hand, for kinessence models, the EoS is a function of z and several dark energy models with different parameterisations of $w(z)$ has been discussed in the literature [13].

According to the latter ideas, several types of *fluid* are used for cosmological viable scenarios, as:

- Non-relativistic matter: An ordinary non-relativistic matter EoS as $w = 0$ (i.e cold dust), correspond to a diluted fluid with $\rho \propto a^{-3} = V_3^{-1}$, where V_3 is the volume in 3 space-like dimensions.
- Ultra-relativistic matter: An ultra-relativistic matter EoS given by $w = 1/3$ (e.g. radiation, or matter in the very early universe), correspond to a fluid as $\rho \propto a^{-4}$. In an expanding universe, the energy density decreases faster than the volume expansion, because radiation has

*Electronic address: ecastellanos@mctp.mx

†Electronic address: celia.escamilla@nucleares.unam.mx

¹ An associated particle can be identified with a boson with zero spin.

a momentum and, by the de Broglie hypothesis a wavelength is redshifted.

- Cosmic accelerated inflation: This case can be characterised by a dark energy EoS. In the simplest scenario, this correspond to the EoS of Λ with $w = -1$. The solution of the conservation equation for the scale factor is not valid and $a \propto e^{Ht}$. In general, the expansion of the universe is accelerating for any EoS $w < -1/3$.
- Hypothetical fluids: In this case, a hypothetical phantom energy is considered to be with a EoS $w < -1$, and we will have a Big Rip. Using current data, it is impossible to distinguish between phantom $w < -1$ and non-phantom $w \geq -1$.

We should mention that the first two cases can be derived from first principles, since that can be deduced from quantum and statistical mechanics. The dark energy and phantom-like fluids cannot be derived as the latter. From the standard model conception, it seems that in our universe exist only two types of particles: bosons and fermions². The nature of this difference relies basically on some intrinsic property, the so-called spin. As a consequence, the statistics, i.e., how these particles occupy a single quantum energy state, depends on their quantum nature. Furthermore, it is well known that the system under consideration displays a significant departure from their classical behaviour at certain critical temperature, depending on whether particles are the constituents of the system (bosons or fermions). In other words, the quantum behaviour is relevant, and under these circumstances and the quantum properties of the corresponding particles must be taken into account. Bosons obey Bose–Einstein statistics and fermions obey Fermi–Dirac statistics. Summarising, the corresponding properties of both systems are quite different when quantum properties are taken into account, the bosons do not fulfill the Pauli’s exclusion principle, while fermions do.

Thus, according to the concepts described above, it can be demonstrated that the occupation number in the Grand Canonical Ensemble can be expressed as³

$$n(\epsilon_p)_{\pm} = \frac{1}{\exp \beta(\epsilon_p - \mu) \pm 1}, \quad (2)$$

where (+) stands for fermions and (−) for bosons. As usual, $\beta = 1/\kappa T$, being κ the Boltzmann’s constant and T the temperature. Additionally, ϵ_p is the corresponding single-particle energy spectrum and μ is the chemical potential. Thus, we

have to deduce, through the dynamics, what is the specific functional form of the energy spectrum for a single particle to analyse the occupation number Eq. (2), which contains relevant thermodynamical information of the system. At this point, it is important to mention that the corresponding EoS contains, in fact, the aforementioned relevant thermodynamical information, if one knows the dynamical properties of the system, i.e., if these properties can be deduced from the experiment or a microscopic model of matter. In other words, the analytical form of the EoS depends on the substance under consideration, and its deduction is dependent on the experiment and/or a microscopic model of matter [14].

Let us calculate the EoS for some thermodynamical systems, by using all the information given above. For this goal, let us appeal to Ref. [15] to exemplify the method. We assume that the semiclassical single-particle energy spectrum for a given system can be expressed as $\epsilon = p^s/\alpha$, where p is the corresponding momentum, s is some positive real number and α is a constant with an adequate dimensions. The system under consideration is ideal, i.e., interactions among its constituents are neglected. To generalise, our system can be situated in n space-like dimensions. Under these circumstances it is straightforward to show that the EoS is given by [15]

$$P = \frac{s}{n} \left(\frac{U}{V_n} \right), \quad (3)$$

where, P is the pressure, U is the internal energy, and V_n is the hyper volume under consideration. Thus, we can define the energy density for any space-like dimension as $\rho_\epsilon = (U/V_n)$.

By using the general relation Eq. (3), we are able to recover the standard results in 3 space-like dimensions (V_3):

- Non-relativistic system: EoS of the form $P = \frac{2}{3}(U/V_3) = \frac{2}{3}\rho_\epsilon$ corresponding to a single-particle energy spectrum $\epsilon = \frac{p^2}{2m}$.
- Ultra-relativistic system: the EoS in this case is of the form $P = \frac{1}{3}(U/V_3) = \frac{1}{3}\rho_\epsilon$ which corresponds to the single-particle energy spectrum $\epsilon = cp$, with c the speed of light.

Let us remark that the properties of the system, and in particular the EoS, strongly depends on the functional form of the single-particle energy spectrum. Due to the functional form of the single particle energy spectrum $\epsilon \sim p^s$, we can obtain an infinite set of EoS in 3 space-like dimensions, depending on the value of the parameter s . However, the last assertion can lead to *exotic* single-particle energy spectra which can be related, in principle, to the kinetic energy of each particle.

Finally, notice that in the case of radiation the EoS is given by $P = \frac{1}{3}(U/V_3) = \frac{1}{3}\rho_\epsilon$. Nevertheless, in this case, we have to make some additional assumptions because for these systems, the total number of particles is not conserved, and consequently, the corresponding chemical potential must be set equal to zero [15]. Moreover, let us remark that the EoS for

² Dark matter and dark energy could be in a different scheme. However, we assume here that even in this case there are only bosons and fermions in the universe.

³ Defined also as the partition function. In this work we assume the equivalence among different ensembles, i.e., our system lies in the thermodynamic limit. In consequence, the choice of another ensemble is irrelevant for the approach followed in the present work since it predicts the same physics.

radiation (together with the corresponding EoS for dust $P = 0$) can be obtained also from first principles without deeper conceptual complications.

Finally, we focus on the following scenarios for different values of the parameter s . For instance, notice that in 3 space-like dimensions when $s = 3$, the relation between the pressure and the energy density is such that $P = (U/V_3) = \rho_\epsilon$, which is a consequence of a semiclassical single-particle energy spectrum of the form $\epsilon = p^3/\alpha$. Notice that an EoS of the form $P = -(U/V_3) = -\rho_\epsilon$ (in the case of dark energy) implies a single-particle energy spectrum $\epsilon = p^{-3}/\alpha$ by using Eq. (3). The last assumption has no physical meaning, since a dispersion relation $\sim p^{-3}$, leads to a negative number of micro-states due to the above analysis is only valid for real positive values of s and it is unable to reproduce the standard EoS for dark energy.

SUPERFLUID EXCITATIONS AS DARK ENERGY EOS

To put in context the system under consideration, let us first analyse some thermodynamical conditions. For this goal, we will describe the nature of the system associated with our model. The system under consideration is a picture of a condensate of some generic bosons (or scalars) and its corresponding elementary excitations. In this aim we calculate the condensation temperature in order to analyse the possibilities to get a condensed state. We start with the semiclassical single-particle energy spectrum $\epsilon = p^s/\alpha$, in three space-like dimensions for a real positive number s .

In this scenario the condensation temperature T_c is given by

$$T_c = \frac{1}{\kappa} \left[\left(\frac{s \hbar^3 \Gamma(3/s)}{4\pi \alpha^{3/s} \zeta(3/s)} \right) n \right]^{s/3}, \quad (4)$$

where, $n = N/V$ is the density of particles, κ is the Boltzmann's constant, \hbar is the Planck's constant, $\Gamma(3/s)$ is the Gamma Function and $\zeta(3/s)$ is the Riemann Zeta Function. Let us remark that for specific values of the parameter s in the range of $0 < s < 3$ the condensation temperature is well defined in the thermodynamic limit. In other words, the temperature of the system in which we are able to describe the associated excitations corresponds at least to temperatures $T < T_c$ which is well defined for values of the parameter s between $0 < s < 3$. This last assertion restrings the onset of condensation and consequently the value of the parameter s , at least in the thermodynamic limit by fundamental physics.

In this line of thought, we propose that a generalised Landau's roton spectrum for dark energy in 3 space-like dimensions that can be written as follows

$$\epsilon_p = \Delta + \frac{(p - p_0)^s}{\alpha}, \quad (5)$$

where s is a real positive number. This energy spectrum is a generalisation of the spectrum associated with superfluidity behaviour of liquid Helium II when $s = 2$, or the so-called

Landau's roton spectrum [15, 16]. In Eq. (5), $\Delta = \epsilon_{p=p_0}$ is some gap energy. The term p_0 , can be interpreted as the momentum near to the extremal of the energy spectrum Eq. (5), i.e., has the behaviour of a quasi-particle. Excitations with momenta close to p_0 are referred to as rotons in the usual non-relativistic case, i.e., for $s = 2$, when the chemical potential μ is set to be zero. We must mention here that a typical superfluid behaviour of the system at low momenta has a linear dispersion relation $\epsilon_p = cp$ with c the speed of sound when $s = 2$ see for instance [15, 16]. This linear dispersion relation at low momenta is caused by the presence of interactions within the system that create elementary excitations related somehow to the so-called Bogoliubov excitations [17]. Thus in the case $s = 2$, the excitation spectrum at the low-energy limit ($p \rightarrow 0$) is identical to that of a phonon, where the sound velocity c is given by $c = \frac{\hbar}{m} \sqrt{4\pi a n}$, being a the s -wave scattering length and n the corresponding density. Excitations with momenta in the linear region are called phonons; those with momenta close to the minimum are called rotons. Excitations with momenta near the maximum are called maxons. However, when $0 < s < 3$ with $s \neq 2$ at very low momenta this situation should change according to the specific value of the parameter s , see Appendix A.

We compute the relation between the pressure P and the internal energy U associated with Eq. (5) near to p_0 , in other words, the corresponding EoS. Consequently, we obtain ⁴

$$\frac{PV}{\kappa T} = \frac{4\pi V}{h^3} \int_0^\infty e^{-\left[\Delta + \frac{(p-p_0)^s}{\alpha}\right]/\kappa T} p^2 dp \simeq \bar{N}, \quad (6)$$

where \bar{N} is the *equilibrium* number of quasi-particles that we call for our purposes *generalised rotons*.

To solve the integral Eq. (6) we take, as was mentioned above $s > 0$, together with the assumption that for low temperatures, i.e., for temperatures of interest, the minimal value of the occupation number $n(\epsilon_p)$ is at most $\exp(-\Delta/\kappa T)$ which is larger than unity. Thus, we can express the occupation number Eq. (2) for bosons as $n(\epsilon_p) \approx \exp(-\beta\epsilon_p)$. With the change of variables $p = p_0 + (\alpha\kappa T)^{1/s}x$, we get from Eq. (6)

$$\frac{PV}{\kappa T} = \frac{4\pi V}{h^3} p_0^2 (\alpha\kappa T)^{1/s} e^{-\Delta/\kappa T} \times \int_{-\infty}^\infty e^{-s} \left(1 + \frac{(\alpha\kappa T)^{1/s}}{p_0} x + \frac{(\alpha\kappa T)^{2/s}}{p_0^2} x^2 \right), \quad (7)$$

without losing generality we are able to obtain

$$\frac{PV}{\kappa T} = \frac{4\pi V}{h^3 s} (\alpha\kappa T)^{1/s} p_0^2 e^{-\Delta/\kappa T} f(s) \quad (8)$$

⁴ From this point forward, the volume V is assumed to be in 3 space-like dimensions.

where

$$f(s) = 3 - \Gamma\left(\frac{1}{s}\right) - \frac{(\alpha\kappa T)^{1/s}}{p_0} \Gamma\left(\frac{2}{s}\right) - \frac{(\alpha\kappa T)^{2/s}}{p_0} \Gamma\left(\frac{3}{s}\right), \quad (9)$$

and $\Gamma(\mu)$ is the Gamma function. To achieve the corresponding internal energy U , we employ the useful thermodynamic functions $A = -PV$ (Helmholtz free energy) and $S = -\left(\frac{\partial A}{\partial T}\right)_V$ (entropy), with $U = A + TS$.

At this point it is important to mention that the phonon contributions to the various thermodynamic properties of the system become rather unimportant near to p_0 ; then, the rotons are the only excitations that need to be considered. Thus, according to our model the particles that form the condensed phase does not contribute to the cosmological energy density and pressure. After straightforward calculations, we obtain for the energy density ρ_E

$$\begin{aligned} \frac{U}{V} \equiv \rho_E &= \frac{4\pi}{h^3 s^2} e^{-(\Delta/\kappa T)} p_0^2 (\alpha\kappa T)^{1/s} \left\{ 3(\Delta s + \kappa T) \right. \\ &\quad \left. - (\alpha\kappa T)^{1/s} \left[2(\Delta s + s\kappa T) \Gamma\left(\frac{1}{s}\right) - (\alpha\kappa T)^{1/s} \right. \right. \\ &\quad \times \left(-(\Delta s + 3s\kappa T) \Gamma\left(\frac{2}{s}\right) \right. \\ &\quad \left. \left. \left. + (\alpha\kappa T)^{1/s} (\Delta s + 4\kappa T) \Gamma\left(\frac{3}{s}\right) \right) \right] \right\}. \quad (10) \end{aligned}$$

Multiplying and dividing by $s(\kappa T f(s))$ we get

$$P = s\rho_E \left(\frac{f(s)}{g(s)} \right), \quad (11)$$

where $g(s)$ is given by

$$\begin{aligned} g(s) &= \frac{1}{\kappa T} \left\{ 3(\Delta s + \kappa T) - (\alpha\kappa T)^{1/s} \left[2(\Delta s + s\kappa T) \right. \right. \\ &\quad \times \left. \Gamma\left(\frac{1}{s}\right) - (\alpha\kappa T)^{1/s} \left(-(\Delta s + 3s\kappa T) \Gamma\left(\frac{2}{s}\right) \right. \right. \\ &\quad \left. \left. \left. + (\alpha\kappa T)^{1/s} (\Delta s + 4\kappa T) \Gamma\left(\frac{3}{s}\right) \right) \right] \right\}. \quad (12) \end{aligned}$$

It must be mentioned that at $T = 0$, the above results tend to zero and clearly, there are not *generalised rotons* in the system. Notice also that for even values of the parameter s the term $s \left(\frac{f(s)}{g(s)} \right)$ is always positive. The interesting scenarios for the EoS Eq. (11) are given for odd values of the parameter s , together with values between $0 < s < 1$, in which the right hand term of Eq. (11) becomes negative. This term seems to be relevant in the search for negative contributions on the energy density of the EoS Eq. (11) near to p_0 , that we interpreted as dark energy.

We define our EoS Eq. (11) in terms of w and s as

$$w(s) \equiv \frac{P}{\rho_E} = s \left(\frac{f(s)}{g(s)} \right), \quad (13)$$

with units of $[T] = \text{Kelvin}$, $[\Delta] = \text{Joules}$, $\alpha = \text{seconds/meter}$, $[p_0] = \text{kilograms meters/seconds}$ and, $\kappa = 1.3 \times 10^{-23} \text{Joules/Kelvin}$.

In Figure (1) we plot Eq. (13) in function of the parameter s at low temperatures. We have three relevant regions which depict the behaviour of the system depending on the value of s . Specifically, when s is large enough we have that $w(s) \rightarrow 0$, i.e., the system behaves as dust. We obtain that for values $s \gtrsim 0.293$ the system behaves as a *normal fluid*, in the sense that the energy density is positive. The interesting scenario seems to be placed for values $s \lesssim 0.293$ in which the energy density becomes negative. For $s \lesssim 0.293$ the system behaves as phantom fluid with $w(s) < -1$. For $s \sim 0.293$, the value of $w(s)$ approaches to -1 in a range of low temperatures according to our approximation. Using the facts described above, we can rewrite Eq. (5) with $s \sim 0.293$ as $\epsilon_p \sim (p - p_0)^{0.293}$. Therefore, in our approach the excitations (or quasi-particles) that we call *generalised rotons*, close enough to p_0 of the corresponding energy spectrum Eq. (5), are able to mimic dark energy when $s \sim 0.293$.

At this point we must mention that an optimal path to fix the free parameters of our model is to evaluate Eq. (5) by considering the initial value $p_0 \geq 0$, in order to obtain initially the case $\epsilon \propto p^s/\alpha$. Once we consider our ansatz Eq. (5) we select a maximum value around the CMB temperature ($T = 2.8\text{K}$) (see also Figure 1; which in fact must be smaller than the corresponding critical temperature T_c Eq. (4)), where the expression inside the parenthesis in Eq. (5) is valid and the values for α does not change significantly. It seems to be that according to temperatures of order of the CMB, Eq. (4) predicts around 10^{143} *condensed particles* in the universe⁵. For this case, the only possible initial value is $\alpha = 1.1$ and then $p_0 \lesssim 10^{-12}$ by using the precedent values. We must mention here that we are not able to obtain the corresponding values of the free parameters from first principles or from fundamental physics⁶. At this level the justification for the election of these specific values for the free parameters is precisely that we are not able to get an EoS of the form $P = -\rho_E$ outside the range of vales presented above. Additionally, the relevant quantities for our analysis will be rescaled for an interval between $p : \{0, 1\}$. Clearly, we must confront the predictions of the model also with observations in order to get some insight to bound the corresponding free parameters. These topics deserve deeper analysis that we will present elsewhere.

⁵ We consider a current expansion rate that indicates a critical density of the universe about $9 \times 10^{-27} \text{kgm}^{-3}$. However, this density is the total density of both matter and energy.

⁶ However, we can fix an interval for the parameter s obtained in Eq. (4).




FIG. 1: *Top left*: Plot of $w(s)$ in terms of $f(s)$ and $g(s)$ at different low temperatures in Kelvins (K) indicated inside the box plot. The EoS units considered here are normalised by κ . The values taking into account were: $\alpha = 1.1$, $\Delta = 0.1$, $\kappa = 1.3 \times 10^{-23}$ and $p_0 = 1.0$. It is interesting to notice that all the graphs intersect when $s \sim 0.293$, which corresponds to $w(s) \simeq -1$. *Top right*: Plot of $w(s)$ at the order of the CMB temperature (blue curve) $T \approx 2.8$ K. Notice that the intersection value of s remains unchanged. *Bottom*: The pressure from Eq. (8). We consider first order terms in $f(s)$ and the obtained value $s = 0.293$. The pressure was normalised in units of $\xi = 10^{-13} e^{\Delta/\kappa} h^3 \kappa^{-1}$. Clearly, while the temperature tends to zero the pressure vanishes, which corresponds to an EoS as dust ($w \rightarrow 0$).

DISCUSSION

In this paper, we present an EoS described by a generic Landau's roton spectrum in 3 space-like dimensions, which can reproduce dark energy characteristics in certain laboratory conditions. To achieve this condition, we define a free parameter s that modulates the $(p - p_0)$ term that appears in Eq. (5), which value can be used to obtain direct information of the corresponding EoS for the superfluid. We obtain that the value $s \sim 0.293$, reproduces $w(s) \simeq -1$. This result leads to a scenario that mimics a Λ CDM EoS that can be eventually extended (and perhaps tested) with superfluidity in the laboratory. These type of *fractional powers* of s , are not an unusual topic in modern physics, e.g., they are related to properties of fractional derivatives and consequently, with *fractional kinetics* [18–21]. Even more, they can be used in several physical applications, see for instance [20–23] and references therein. For instance, in Ref. [18] it was analysed a fractional Schrödinger-type wave equation in 1-dimension for a free particle. In this case, the corresponding energy spectrum is given by $\epsilon_k = \eta(\hbar k)^{2\gamma}$, where γ takes real positive

values and η is defined as $\eta \equiv \frac{1}{2}(m^{2\gamma-1}c^{2(\gamma-1)})^{-1}$. It is noteworthy to mention that this result is in agreement with our approach when $s = 2\gamma$.

According to the results obtained in the Appendix A, the system behaves as $p^{1/6}$, in which we interpreted the term $\sim \alpha^{1/2} \sqrt{\frac{2U_0 N}{V}}$ as the *speed of sound* for low momenta up to some maximal value p_m . Additionally, it is seem to be that there is a region between p_m and p_0 in which we do not have any information about the functional form of ϵ_p . In fact, if we assume the usual behaviour, this region must be formed by a combination of phonons and rotons. Thus, it could be interesting to explore the system in this interval of values for the momenta for low temperatures (or even a scenario of dynamical s) in order to extract information of the system and set at least bounds for the free parameters of the model and eventually compare with observations. These issues are out of the scope of this work, however will be explored in future works. Finally, it is remarkable that our proposal gives insights that can eventually be applied at cosmological scales to understand the nature of dark energy in the universe.

Acknowledgments: EC acknowledges the receipt of the grant from the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy. CE-R acknowledges the Royal Astronomical Society as FRAS 10147, PAPIIT Project IA100220 and would like to acknowledge networking support by the COST Action CA18108.

-
- [1] A. G. Riess *et al.* [Supernova Search Team], *Astron. J.* **116** (1998), 1009-1038 doi:10.1086/300499 [arXiv:astro-ph/9805201 [astro-ph]].
- [2] P. M. Garnavich *et al.* [Supernova Search Team], *Astrophys. J.* **509** (1998), 74-79 doi:10.1086/306495 [arXiv:astro-ph/9806396 [astro-ph]].
- [3] J. L. Tonry *et al.* [Supernova Search Team], *Astrophys. J.* **594** (2003), 1-24 doi:10.1086/376865 [arXiv:astro-ph/0305008 [astro-ph]].
- [4] T. Padmanabhan, *Phys. Rept.* **380** (2003), 235-320 doi:10.1016/S0370-1573(03)00120-0 [arXiv:hep-th/0212290 [hep-th]].
- [5] P. Peebles and B. Ratra, *Rev. Mod. Phys.* **75** (2003), 559-606 doi:10.1103/RevModPhys.75.559 [arXiv:astro-ph/0207347 [astro-ph]].
- [6] R. Caldwell, R. Dave and P. J. Steinhardt, *Phys. Rev. Lett.* **80** (1998), 1582-1585 doi:10.1103/PhysRevLett.80.1582 [arXiv:astro-ph/9708069 [astro-ph]].
- [7] A. Shafieloo, V. Sahni and A. A. Starobinsky, *Phys. Rev. D* **86** (2012), 103527 doi:10.1103/PhysRevD.86.103527 [arXiv:1205.2870 [astro-ph.CO]].
- [8] M. Chevallier and D. Polarski, *Int. J. Mod. Phys. D* **10** (2001), 213-224 doi:10.1142/S0218271801000822 [arXiv:gr-qc/0009008 [gr-qc]].
- [9] C. Escamilla-Rivera, *Galaxies* **4** (2016) no.3, 8 doi:10.3390/galaxies4030008 [arXiv:1605.02702 [astro-ph.CO]].
- [10] T. Clifton, [arXiv:gr-qc/0610071 [gr-qc]].
- [11] S. Nesseris and L. Perivolaropoulos, *Phys. Rev. D* **70** (2004), 043531 doi:10.1103/PhysRevD.70.043531 [arXiv:astro-ph/0401556 [astro-ph]].
- [12] N. Aghanim *et al.* [Planck 2018], [arXiv:1807.06209 [astro-ph.CO]].
- [13] V. Sahni, T. D. Saini, A. A. Starobinsky and U. Alam, *JETP Lett.* **77** (2003), 201-206 doi:10.1134/1.1574831 [arXiv:astro-ph/0201498 [astro-ph]].
- [14] Leopoldo García-Colín Scherer, *Introducción a la Termodinámica Clásica*, 4a ed, Ed. Trillas (2005).
- [15] R.K. Pathria, *Statistical Mechanics*, Butterworth Heineman, Oxford (1996).
- [16] C. J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases*, Cambridge University Press, Cambridge (2002).
- [17] M. Ueda, *Fundamentals and New Frontiers of Bose-Einstein Condensation* (World Scientific, Singapore) 2010.
- [18] R. Herrmann, arXiv:math-ph/0510099v4, (2006).
- [19] F. Riewe, *Phys. Rev E*, **55** 3 (1996).
- [20] G.M. Zaslavsky, *Physics Reports* **371** 461-580, (2002).
- [21] M. Dalir, *Applied Mathematical Sciences*, **4** 21, 1021-1032, (2010).
- [22] R. Hilfer, *Fractional Calculus in Physics*, World Scientific Publishing Co. Pte. Ltd, (2000).
- [23] K. A. Lazopoulos and A. K. Lazopoulos, *J Mech Behav Mater*, 26(1-2): 43-54, (2017).

[24] D. A. McQuarry, *Statistical Mechanics*, University Science Books, (2000).

[25] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series and Products*, Academic Press, New York, (1980).

Appendix A

Let us analyse the corresponding excitation spectrum at the low-energy limit $p \ll p_0$, i.e., the phonon-like behaviour, if there exist, for the specific value $s \sim 1/3$. Let us start with a N-body Hamiltonian in three dimensions where we have just introducing $p = \hbar k$

$$\hat{H} = \sum_{\vec{k}=0} \left[\frac{\hbar^s k^s}{\alpha} \right] \hat{a}_{\vec{k}}^\dagger \hat{a}_{\vec{k}} + \frac{U_0}{2V} \sum_{\vec{k}=0} \sum_{\vec{p}=0} \sum_{\vec{q}=0} \hat{a}_{\vec{p}}^\dagger \hat{a}_{\vec{q}}^\dagger \hat{a}_{\vec{p}+\vec{k}} \hat{a}_{\vec{q}-\vec{k}}, \quad (14)$$

where the $U_0 = 4\pi\hbar^2 a/m$ is the corresponding interaction potential with a the scattering length. Additionally, $\hat{a}_{\vec{k}}^\dagger$, $\hat{a}_{\vec{k}}$ are the corresponding creation and annihilation operators that satisfy the usual canonical commutation relations for bosons. We assume that for temperatures $T < T_c$ (see Eq. (4))

$$N_0 \approx N, \quad \sum_{\vec{k} \neq 0} N_{\vec{k}} \ll N, \quad (15)$$

being N the total number of particles, $N_{\vec{k}}$ the number of particles in the excited states, and N_0 the number of particles in the ground state. Keeping terms up to second order in \hat{a}_0 , the Hamiltonian (14) becomes

$$\begin{aligned} \hat{H} = & \frac{U_0 N^2}{2V} + \sum_{\vec{k} \neq 0} \left[\frac{\hbar^s k^s}{\alpha} + \frac{U_0 N}{V} \right] \hat{a}_{\vec{k}}^\dagger \hat{a}_{\vec{k}} \\ & + 2\hat{a}_0^\dagger \hat{a}_0 \sum_{\vec{k} \neq 0} \hat{a}_{\vec{k}}^\dagger \hat{a}_{\vec{k}} + \hat{a}_0^{\dagger 2} \sum_{\vec{k} \neq 0} \hat{a}_{\vec{k}} \hat{a}_{-\vec{k}} + \hat{a}_0^2 \sum_{\vec{k} \neq 0} \hat{a}_{\vec{k}}^\dagger \hat{a}_{-\vec{k}}^\dagger. \end{aligned} \quad (16)$$

Finally, in the same order of the approximation we assume that $\hat{a}_0^\dagger \hat{a}_0$, \hat{a}_0^2 , $\hat{a}_0^{\dagger 2} = N$. Using these facts, the Hamiltonian (14) can be re-expressed as follows

$$\begin{aligned} \hat{H} = & \frac{U_0 N^2}{2V} + \sum_{\vec{k} \neq 0} \left[\frac{\hbar^s k^s}{\alpha} + \frac{U_0 N}{V} \right] \hat{a}_{\vec{k}}^\dagger \hat{a}_{\vec{k}} \\ & + \sum_{\vec{k} \neq 0} \frac{U_0 N}{2V} \left[\hat{a}_{\vec{k}}^\dagger \hat{a}_{-\vec{k}}^\dagger + \hat{a}_{\vec{k}} \hat{a}_{-\vec{k}} \right]. \end{aligned} \quad (17)$$

In order to obtain the ground state energy associated with our system, let us diagonalize the above Hamiltonian by introducing the so-called Bogoliubov transformations [15, 16].

After some algebra, the following diagonalized Hamiltonian can be obtained

$$\begin{aligned} \hat{H} = & \frac{U_0 N^2}{2V} + \sum_{\vec{k} \neq 0} \sqrt{\epsilon_k \left(\epsilon_k + \frac{2U_0 N}{V} \right)} \hat{b}_{\vec{k}}^\dagger \hat{b}_{\vec{k}} \\ & + \sum_{\vec{k} \neq 0} \left\{ -\frac{1}{2} \left[\frac{U_0 N}{V} + \epsilon_k - \sqrt{\epsilon_k \left(\epsilon_k + \frac{2U_0 N}{V} \right)} \right] \right\}, \end{aligned} \quad (18)$$

where $\epsilon_k = \frac{\hbar^s k^s}{\alpha}$. The last summation in the Hamiltonian (18) diverges as $(U_0 N/V)^2/2\epsilon_k$, as can be seen by performing an expansion of the last term in Eq. (18) for large k .

The pseudo-potential $U_P(\mathbf{r})$ can be expressed as follows

$$U_P(\mathbf{r}) = U_0 \delta(\mathbf{r}) \frac{\partial}{\partial r} r, \quad (19)$$

Notice that in our case, the divergence is basically of order α/k^s .

The 3-D Fourier transform in spherical coordinates can be written as [24, 25]

$$V(r) = \alpha \left(\frac{2}{\pi} \right)^{1/2} \int_0^{p_m} \frac{k \sin(kr)}{(\hbar k)^s r} dk. \quad (20)$$

Thus, the Fourier transform upon $1/k^s$ can be expressed generically as follows

$$V(r) = \frac{\sqrt{\frac{2}{\pi}} p_m^{3-s} {}_1F_2 \left(\frac{3}{2} - \frac{s}{2}; \frac{3}{2}, \frac{5}{2} - \frac{s}{2}; -\frac{1}{4} p_m^2 r^2 \right)}{3-s}, \quad (21)$$

where ${}_1F_2$ is a generalized hypergeometric function valid for $0 < s < 3$. The term p_m is a maximum momenta, i.e., a cut-off, included in order to avoid the divergence of $V(r)$, clearly small enough compared with p_0 . Fortunately, ${}_1F_2 \rightarrow \frac{3p_m^{s/3}}{4\sqrt{2\pi}}$ when $r \rightarrow 0$, i.e., is regular at the origin. Consequently, the pseudo potential is just a simple delta-function potential $U_P(r) = U_0 \delta(r)$ for $0 < s < 3$, see for instance Ref. [17]

It is noteworthy to mention that even in this situation the action of the operator (19), removes the divergence in the Fourier transform (21). Thus, by subtracting the divergent term $(U_0 N/V)^2/2\epsilon_k$ in the corresponding Hamiltonian (18), we safely remove the divergent behavior.

Finally, our Hamiltonian now is given by

$$\begin{aligned} \hat{H} = & \frac{U_0 N^2}{2V} + \sum_{\vec{k} \neq 0} \sqrt{\epsilon_k \left(\epsilon_k + \frac{2U_0 N}{V} \right)} \hat{b}_{\vec{k}}^\dagger \hat{b}_{\vec{k}} \\ & + \sum_{\vec{k} \neq 0} \left\{ -\frac{1}{2} \left[\frac{U_0 N}{V} + \epsilon_k - \sqrt{\epsilon_k \left(\epsilon_k + \frac{2U_0 N}{V} \right)} \right. \right. \\ & \left. \left. - \left(\frac{U_0 N}{V} \right)^2 \frac{1}{2\epsilon_k} \right] \right\}. \end{aligned} \quad (22)$$

Thus, in one hand, the ground state energy is well defined for values $0 < s < 3$ compatible with the result Eq. (4). On the other hand, from the N -body Hamiltonian (22), we are able to recognize the energy of the Bogoliubov excitations

$$E_k = \sqrt{\epsilon_k \left(\epsilon_k + \frac{2U_0 N}{V} \right)}. \quad (23)$$

The long-wavelength limit, $k \rightarrow 0$, associated with the above expression, leads to the following dispersion relation

$$E_p = \frac{p^{s/2}}{\alpha^{1/2}} \sqrt{\frac{2U_0 N}{V}} \left[1 + \frac{p^s}{4\alpha U_0 N} \right] \quad (24)$$

Notice that in the case $s = 2$ we recover the linear phonon-like dispersion relation for the excitations, i.e., $E_k \sim p$.

However, in the case $s \sim 1/3$ we obtain

$$E_p = \frac{p^{1/6}}{\alpha^{1/2}} \sqrt{\frac{2U_0 N}{V}} \left[1 + \frac{p^{1/3}}{4\alpha U_0 N} \right] \quad (25)$$

which basically behaves as $E_k \sim p^{1/6}$, for low momenta valid up to some value p_m .