
CONFIDENCE INTERVAL FOR OFF-POLICY EVALUATION FROM DEPENDENT SAMPLES VIA BANDIT ALGORITHM: APPROACH FROM STANDARDIZED MARTINGALES

A PREPRINT

Masahiro Kato

CyberAgent Inc.

Tokyo, Japan

masahiro_kato@cyberagent.co.jp

January 3, 2022

ABSTRACT

This study addresses the problem of *off-policy evaluation* (OPE) from *dependent samples* obtained via the *bandit algorithm*. The goal of OPE is to evaluate a new policy using historical data obtained from *behavior policies* generated by the bandit algorithm. Because the bandit algorithm updates the policy based on past observations, the samples are not *independent and identically distributed* (i.i.d.). However, several existing methods for OPE do not take this issue into account and are based on the assumption that samples are i.i.d. In this study, we address this problem by constructing an estimator from a *standardized martingale difference sequence*. To standardize the sequence, we consider using *evaluation data* or *sample splitting* with *two-step estimation*. This technique produces an estimator with asymptotic normality without restricting a class of behavior policies. In an experiment, the proposed estimator performs better than existing methods, which assume that the behavior policy converges to a time-invariant policy.

1 Introduction

The *multi-armed bandit* (MAB) problem is one of the sequential decision-making problems, which is applied in various applications, such as ad-optimization, personalized medicine, search engines, and recommendation systems. In the MAB problem, we choose an action at a period and observe the outcome. To reduce the cost of searching for better policies, the evaluation of a new policy using historical data obtained from past trials has gathered great attention (Beygelzimer & Langford, 2009; Li et al., 2010; Athey & Wager, 2017). This framework is called *off-policy evaluation* (OPE) (Dudík et al., 2011; Wang et al., 2017; Narita et al., 2019; Bibaut et al., 2019; Kallus & Uehara, 2019a; Oberst & Sontag, 2019). Although several methods for OPE have been proposed, existing studies often presume that the samples are *independent and identically distributed* (i.i.d.). However, in the MAB problem algorithm, the policy is usually updated based on past observations, and the samples are not i.i.d. due to the policy updating process. In this case, the consistency and asymptotic normality of the existing methods are not guaranteed. In particular, the asymptotic normality is critical because it guarantees the \sqrt{n} -consistency and is needed for the confidence interval in hypothesis testing to determine whether the new policy is better than the existing policy. Thus, the motivation for establishing a novel method for performing OPE from dependent samples is strong.

Several previous studies consider OPE from dependent samples (der Laan & Mark, 2008; Hadad et al., 2019; Kallus & Uehara, 2019b; Kato et al., 2020a). The strategies for deriving asymptotic normality can be classified into three approaches. In the first approach, under the assumption that the policy used in past trials converges to the time-invariant policy in probability, the asymptotic normality is derived using the theories related to *martingales* (der Laan & Mark, 2008; Hadad et al., 2019; Kato et al., 2020a). In the second approach, the policy is assumed to be batch updated, where, although the policy is updated using past observations, the sample size under a fixed policy is sufficient (Hahn et al., 2011). In the third approach, both the stationarity and the conditions of *mixingales* (Kosorok,

2008), which requires the independence of time-separated samples, are assumed (Kallus & Uehara, 2019b). However, existing methods have several drawbacks. With the first approach, the assumption that the policy converges to a time-invariant policy sometimes does not hold. For example, if the algorithms of the MAB problem change during past trials, it is difficult to justify this assumption. The second and third approaches cannot be applied in cases where the policy is sequentially updated, and the conditions of mixingales do not hold, respectively.

To overcome these drawbacks, in this study, we propose a novel estimator with the asymptotic normality, which does not assume policy convergence, the existence of batches, and the independence of time-separated samples. To accomplish this goal, we refocus on the first approach. In the first approach, we use a *martingale difference sequence* (MDS) for constructing an estimator. In this approach, the convergence of the policy used in past trials to the time-invariant policy in probability is assumed for using *central limit theorem* (CLT) of an MDS, which requires that the variance of an MDS becomes asymptotically constant. In this paper, instead of assuming the convergence of the policy, we consider constructing an MDS with asymptotically constant variance by *standardization* using an estimator of the variance. To implement the standardization, we assume access to evaluation (test) data or conduct *sample splitting*. Through this technique, we can guarantee the asymptotic normality of the proposed estimator with fewer assumptions.

Contributions and Organization of this Paper: This paper has three main contributions. First, this paper provides a theoretical solution for causal inference from dependent samples obtained via bandit feedback, which is more realistic than the situation that has been often considered in the existing work. Second, the proposed estimator achieves the asymptotic normality with fewer assumptions compared with existing solutions. Third, the estimator also achieves a lower mean squared error (MSE) in experiments using benchmark datasets.

Additional Related Work: There are various studies for OPE under the assumption that samples are i.i.d. (Dudík et al., 2011; Wang et al., 2017; Narita et al., 2019; Bibaut et al., 2019; Kallus & Uehara, 2019a; Oberst & Sontag, 2019), but there are fewer studies focus on the case in which samples are not i.i.d. When the policy converges, der Laan & Mark (2008), Hadad et al. (2019), and Kato et al. (2020a) proposed using an estimator based on an MDS. When including the covariates for OPE, der Laan & Mark (2008) mainly suggested using targeted maximum likelihood estimation. On the other hand, Hadad et al. (2019) and Kato et al. (2020a) proposed using an estimator based on the doubly robust or augmented IPW estimator. Kallus & Uehara (2019b) dealt with the application of reinforcement learning and proposed an estimator constructed from dependent samples. In their study, they proposed *time cross-fitting*, which is a variant of cross-fitting of double/debiased machine learning (Chernozhukov et al., 2018). The theoretical guarantee is based on mixingale theory. For obtaining the asymptotic normality, conducting the standardization is also proposed by Luedtke & van der Laan (2016). Luedtke & van der Laan (2016) tries to solve a similar problem in the batch policy updating. In addition to this difference of policy updating, there are the following two points different from us. First, in their solution, they require estimators of f^* and e^* to satisfy some conditions like the Donsker condition. However, in our estimator, we do not impose such restrictions on estimators of f^* and e^* except for the boundedness. This difference is critical because, in practice, as Chernozhukov et al. (2018) pointed out, the estimators of f^* and e^* usually do not satisfy the Donsker condition when we use modern methods such as the Lasso and Random forest regression. Second, for obtaining an estimator with the asymptotic normality under sequential policy updating, we propose using evaluation data or sample splitting.

2 Problem Setting

In this section, we introduce our problem setting.

2.1 Data Generating Process

Let A_t be the action taking variable in $\mathcal{A} = \{1, 2, \dots, K\}$, X_t be the *covariate* observed by the decision maker when choosing an action, and \mathcal{X} be the domain of covariate. Let us denote a random variable of a reward at a period t as a function $Y_t : \mathcal{A} \rightarrow \mathbb{R}$. Let $\mathbb{1}[\cdot]$ be an indicator function. In this paper, we have access to a set of *historical data*, $\mathcal{S}_T = \{(X_t, A_t, Y_t)\}_{t=1}^T$, with the following data generating process (DGP):

$$\{(X_t, A_t, Y_t)\}_{t=1}^T \sim p(x)p_t(a | x, \Omega_{t-1})p(y | a, x), \quad (1)$$

where $Y_t = \sum_{a=1}^K \mathbb{1}[A_t = a]Y_t(a)$, $p(x)$ denote the density of the covariate X_t , $p_t(a | x, \Omega_{t-1})$ denote the probability of assigning an action A_t conditioned on a covariate X_t , $p(y | a, x)$ denote the density of an outcome Y_t conditioned on A_t and X_t , and $\Omega_{t-1} \in \mathcal{M}_{t-1}$ denotes the history defined as $\Omega_{t-1} = \{X_{t-1}, A_{t-1}, Y_{t-1}, \dots, X_1, A_1, Y_1\}$ with the domain \mathcal{M}_{t-1} . We assume that $p(x)$ and $p(y | a, x)$ are invariant across periods, but $p_t(a | x, \Omega_{t-1})$ can take different value across periods. Besides, we allow the decision maker to change $p_t(a | x, \Omega_{t-1})$ based on past observations.

In this case, samples $\{(X_t, A_t, Y_t)\}_{t=1}^T$ are correlated across periods, i.e., samples are not i.i.d. Here, we introduce *behavior policies*, which determine the probability $p_t(a | x, \Omega_{t-1})$. Let a behavior policy $\pi_t : \mathcal{A} \times \mathcal{X} \times \mathcal{M}_{t-1} \rightarrow (0, 1)$ in the t -th period be a function of a covariate X_t , an action A_t , and a history Ω_{t-1} . In this study, we assume that $\pi_t(a | x, \Omega_{t-1}) = p_t(a | x, \Omega_{t-1})$.

Remark 1 (Evaluation Data). In Section 4.3, we also assume that there is access to *evaluation data*, $\mathcal{E}_N = \{X_i\}_{i=1}^N$. In Remark 7, we relax this assumption by introducing the *sample splitting*.

2.2 Off-Policy Evaluation

Under the DGP defined in the previous subsection, we consider estimating the value of an *evaluation policy* using samples obtained under the behavior policies. Let an evaluation policy $\pi^e : \mathcal{A} \times \mathcal{X} \rightarrow (0, 1)$ be a function of a covariate X_t and an action A_t ; this evaluation policy can be considered as the probability of taking the action A_t conditional on the covariate X_t . We are interested in estimating the expected reward from a given pre-specified evaluation policy $\pi^e(a | x)$. Then, we define the expected reward under π^e as $R(\pi^e) := \mathbb{E} \left[\sum_{a=1}^K \pi^e(a, X_t) Y_t(a) \right]$. We also denote $R(\pi^e)$ as θ_0 . The goal of this study is to estimate $R(\pi^e)$ using dependent samples under a sequentially-updated policy. To identify $R(\pi^e)$, we make the following assumptions on the policies and the outcomes.

Assumption 1. The behavior policies $\pi_t(a | \cdot, \Omega_{t-1})$ and the evaluation policy $\pi^e(a | \cdot)$ are known¹. Additionally, the evaluation policy is deterministic.

Assumption 2. There exist C_1 and C_2 such that $\frac{\pi^e(a|x)}{\pi_t(a|x, \Omega_{t-1})} \leq C_1$ and $|Y_t| \leq C_2$.

The deterministic evaluation policy looks restrictive. However, the optimal policy that maximizes the expected reward is a deterministic policy when the policy is not restricted. Therefore, when we find the optimal policy from a set of policies, such as *off-policy learning* (Zhao et al., 2012; Kitagawa & Tetenov, 2018; Zhou et al., 2018; Chernozhukov et al., 2019), this assumption can be accommodated. Thus, restricting the policy class to the deterministic policy class would be reasonable in practice. In fact, most of the existing methods follow this way.

Remark 2 (Existing Methods for OPE). We review three types of standard estimators of $R(\pi^e)$ under the case where $p_1(a | x) = p_2(a | x) = \dots = p_T(a | x) = p(a | x)$ in the DGP defined in (1). The first estimator is an inverse probability weighting (IPW) estimator given by $\frac{1}{T} \sum_{t=1}^T \sum_{a=1}^K \frac{\pi^e(a|X_t) \mathbb{1}[A_t=a] Y_t}{p(a|X_t)}$ (Horvitz & Thompson, 1952; Rubin, 1987; Hirano et al., 2003). Although this estimator is unbiased when the behavior policy is known, it has a high variance. The second estimator is a direct method (DM) estimator $\frac{1}{T} \sum_{t=1}^T \sum_{a=1}^K \hat{f}_T(a, X_t)$, where $\hat{f}_T(a, X_t)$ is an estimator of $f^*(a, X_t)$ (Hahn, 1998). This estimator is known to be weak against model misspecification for $f^*(a, X_t)$. The third estimator is a doubly robust estimator defined as $\frac{1}{T} \sum_{t=1}^T \sum_{a=1}^K \left(\frac{\pi^e(a|X_t) \mathbb{1}[A_t=a] (Y_t - \hat{f}_T(a, X_t))}{p(a|X_t)} + \pi^e(a | X_t) \hat{f}_T(a, X_t) \right)$ (Robins et al., 1994; Chernozhukov et al., 2018). Under certain conditions, it is known that this estimator achieves the lower bound (a.k.a semiparametric lower bound), which is the lower bound of the asymptotic MSE of OPE, among regular \sqrt{T} -consistent estimators (van der Vaart, 1998, Theorem 25.20).

Remark 3 (Unconfoundedness). Existing methods often make an assumption called unconfoundedness, that is, they assume that the outcomes $(Y_t(0), Y_t(1), \dots, Y_t(K))$ and the action A_t are conditionally independent on X_t for identification. In the DGP, we can obtain a similar result from the assumption that an action A_t is chosen according to probability $p_t(a | x, \Omega_{t-1})$.

Remark 4 (Semiparametric Lower Bound). The lower bound of the variance can be defined for an estimator under some posited models of the DGP. If this posited model is a parametric model, then the lower bound is equal to the Cramér-Rao lower bound. When this posited model is a non- or semiparametric model, the corresponding lower bound can still be defined (Bickel et al., 1998). Narita et al. (2019) shows that the semiparametric lower bound of the DGP (1) under $p_1(a | x, \Omega_0) = p_2(a | x, \Omega_1) = \dots = p_T(a | x, \Omega_{T-1}) = p(a | x)$ is $\mathbb{E} \left[\sum_{a=1}^K \left\{ \frac{(\pi^e(a|X_t))^2 v^*(a, X_t)}{p(a|X_t)} + (\pi^e(a | X_t) f^*(a, X_t) - \theta_0)^2 \right\} \right]$.

Notations: Let a be an action in \mathcal{A} . We denote $\mathbb{E}[Y_t(a) | x]$, $\mathbb{E}[Y_t^2(a) | x]$, and $\text{Var}(Y_t(a) | x)$ as $f^*(a, x)$, $e^*(a, x)$, and $v^*(a, x)$, respectively. Let \mathcal{F} be the domain of $f^*(a, x)$. Let $\hat{f}_t(a, x)$, $\hat{e}_t(a, x)$, and $\hat{\theta}_t$ be estimators of $f^*(a, x)$, $e^*(a, x)$, and θ_0 constructed from Ω_t , respectively. Let $\mathcal{N}(\mu, \text{var})$ be the normal distribution with mean μ and variance var .

¹The proposed method requires the function of the polices, not only $\pi_t(a | X_t, \Omega_{t-1})$ for a specific X_t .

2.3 Policy Updating

In the DGP defined in Section 2.1, we allow the behavior policy $\pi_t(a \mid x, \Omega_{t-1})$ to be updated based on past observations. For example, when applying some algorithms proposed in the MAB problem, we usually optimize the probability to maximize the cumulative reward using past observations. When the behavior policy $\pi_t(a \mid x, \Omega_{t-1})$ depends on past observations, the samples are not i.i.d. In this case, we cannot apply the standard methods for OPE, which assume that the samples are i.i.d. To clarify this problem, we classify the patterns of policy updating into two scenarios: *sequential policy updating* and *batch policy updating*. In sequential policy updating, the policy can be updated each period. In batch policy updating, the policy is updated after observing some samples (Hahn et al., 2011; Narita et al., 2019). In this study, we consider a problem using sequential policy updating.

3 OPE under a Converging Policy

In this section, we review OPE when a policy converges to the time-invariant policy in probability. As der Laan & Mark (2008), Hadad et al. (2019) and Kato et al. (2020a) proposed, in such a situation, we can construct an estimator with the asymptotic normality by using an MDS (Hall et al., 2014). To conduct OPE when the behavior policy is known, we can use the following score function $\psi : \Theta \times \mathcal{X} \times \mathcal{A} \times \mathbb{R} \times \Pi_t \times \Pi \times \mathcal{F} \rightarrow \mathbb{R}$ (Chernozhukov et al., 2018):

$$\psi_t(\theta; x, k, y, \pi_t, \pi^e, f) = \sum_{a=1}^K \left\{ \frac{\pi^e(a \mid x) \mathbb{1}[A_t = a] \{y - f(a, x)\}}{\pi_t(a \mid x, \Omega_{t-1})} + \pi^e(a \mid x) f(a, x) \right\} - \theta, \quad (2)$$

where Θ , Π_t , and Π are spaces of a parameter θ , a behavior policies π_t , and an evaluation policy π^e . Then, for the true policy value θ_0 , the score function with $\theta = \theta_0$, $\psi_t(\theta_0; X_t, A_t, Y_t, \pi_t, \pi^e, \hat{f}_{t-1})$, satisfies $\mathbb{E}[\psi_t(\theta_0; X_t, A_t, Y_t, \pi_t, \pi^e, \hat{f}_{t-1})] = 0$. As we explain in Remark 2, there are several other candidate estimators, including the IPW estimator. However, the estimator based on the score function defined in (2) achieves the lowest asymptotic variance when π_t is fixed among periods (Bickel et al., 1998; Scharfstein et al., 1999). Using $\{\psi_t(\theta_0; X_t, A_t, Y_t, \pi_t, \pi^e, \hat{f}_{t-1})\}_{t=1}^T$, we consider constructing an MDS $\{z_t\}_{t=1}^T$ that satisfies the conditions of the following CLT for an MDS.

Proposition 1. [CLT for an MDS, Hamilton (1994), Proposition 7.9, p. 194] Let $\{z_t\}_{t=1}^\infty$ be an MDS. Suppose that (a) $\mathbb{E}[z_t^2] = \nu_t^2 > 0$ with $(1/T) \sum_{t=1}^T \nu_t^2 \rightarrow \nu^2 > 0$; (b) $\mathbb{E}[|z_t|^r] < \infty$ for some $r > 2$; (c) $(1/T) \sum_{t=1}^T z_t^2 \xrightarrow{P} \nu^2$. Then $\sqrt{T} \frac{1}{T} \sum_{t=1}^T z_t \xrightarrow{d} \mathcal{N}(0, \nu^2)$.

When Y_t , π^e/π_t , and f_{t-1} are bounded, we can easily show that the condition (b) holds (Assumption 2). The remaining task is to show that conditions (a) and (c) hold.

Kato et al. (2020a) constructed $\check{z}_t = \psi_t(\theta_0; X_t, A_t, Y_t, \pi_t, \pi^e, \hat{f}_{t-1})$, where $\{\check{z}_t\}_{t=1}^T$ is an MDS. Then, by assuming $\pi_t(a \mid x, \Omega_{t-1}) \xrightarrow{P} \alpha(a \mid x)$ and some regularity conditions, Kato et al. (2020a) proved that the sequence $\{\check{z}_t\}_{t=1}^T$ satisfies the conditions (a) and (c) with $\nu^2 = \mathbb{E} \left[\left(\sum_{a=1}^K \left\{ \frac{\pi^e(a \mid x) \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\alpha(a \mid X_t)} + \pi^e(a \mid X_t) f^*(a, X_t) \right\} - \theta_0 \right)^2 \right]$. Thus, by the assumption $\pi_t(a \mid x, \Omega_{t-1}) \xrightarrow{P} \alpha(a \mid x)$, the variance of h_t converges to time-invariant value, and this property is the motivation of the assumption proposed by Kato et al. (2020a). Then, under $\pi_t(a \mid x, \Omega_{t-1}) \xrightarrow{P} \alpha(a \mid x)$, the estimator proposed defined as $\hat{\theta}_T^{\text{A2IPW}} = \frac{1}{T} \sum_{t=1}^T \sum_{a=1}^K \left\{ \frac{\pi^e(a \mid X_t) \mathbb{1}[A_t = a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a \mid X_t, \Omega_{t-1})} + \pi^e(a \mid x) \hat{f}_{t-1}(a, X_t) \right\}$ can be proved to be consistent and has the asymptotic normality (Hadad et al., 2019; Kato et al., 2020a)². We refer to this estimator as *Adaptive AIPW* (A2IPW). We can also define the corresponding IPW estimator and refer to it as *adaptive IPW* (AdaIPW) estimator.

Remark 5 (Efficiency of A2IPW). The asymptotic variance of the A2IPW estimator matches the semiparametric lower bound under a time-invariant policy α (Kato et al., 2020a).

²They assume the almost sure convergence of the policy for showing the mean convergence of the variance, but we can relax it to the convergence in probability by the boundedness of the MDS (Loeve, 1977).

4 OPE from Dependent Samples under Sequential Policy Updating

In this section, we discuss a strategy for constructing an estimator without assuming the convergence of the policy. Then, we show a method for constructing an estimator with the asymptotic normality.

4.1 Strategy for OPE from Dependent Samples under Sequential Policy Updating

However, there are various applications in which we cannot assume that $\pi_t(a | x, \Omega_{t-1}) \xrightarrow{P} \alpha(a | x)$. Therefore, we have a strong motivation for constructing an estimator with asymptotic normality without assuming that $\pi_t(a | x, \Omega_{t-1}) \xrightarrow{P} \alpha(a | x)$. Our strategy is to construct an MDS $\{\tilde{z}_t\}_{t=1}^T$ satisfying conditions (a) and (c) by standardization of \check{z}_t as $\tilde{z}_t = \left(\sqrt{\text{Var}(\check{z}_t | \Omega_{t-1})}\right)^{-1} \check{z}_t$, where $\text{Var}(\check{z}_t | \Omega_{t-1})$ is the conditional variance of \check{z}_t . The standardization means that the variance of \tilde{z}_t becomes 1. Then, we can easily show that the conditions (a) and (b) hold. Thus, if we know the variance $\text{Var}(\check{z}_t | \Omega_{t-1})$, we can construct an MDS with constant variance by the standardization.

Remark 6 (Lindeberg Condition). The proposed strategy can be interpreted using the Lindeberg condition. For all $\epsilon > 0$ and an MDS $\{z_t\}_{t=1}^T$ such that $\sum_{t=1}^T \mathbb{E}[z_t^2 | \Omega_{t-1}] / \sum_{t=1}^T \mathbb{E}[z_t^2] \xrightarrow{P} 1$, the Lindeberg condition is $\left(\sum_{t=1}^T \mathbb{E}[z_t^2]\right)^{-1} \sum_{t=1}^T \mathbb{E}\left[z_t^2 \mathbb{1}\left[|z_t| \geq \epsilon \sqrt{\sum_{t=1}^T \mathbb{E}[z_t^2]}\right]\right] \xrightarrow{P} 0$ as $T \rightarrow \infty$ (Brown, 1971). The first condition $\sum_{t=1}^T \mathbb{E}[z_t^2 | \Omega_{t-1}] / \sum_{t=1}^T \mathbb{E}[z_t^2] \xrightarrow{P} 1$ restricts the class of an MDS to an MDS with asymptotically time-invariant variance. Here, we note that $\sigma_t^2 = \mathbb{E}[z_t^2 | \Omega_{t-1}]$. The strategy of this paper can be intuitively interpreted as a transformation of the original sequence \check{z}_t into such an MDS with asymptotically time-invariant variance.

4.2 OPE with a Weighted Average Estimator: Case with Known Variance

Then, we propose an estimator based on the MDS $\{\tilde{z}_t\}_{t=1}^T$ and refer to the estimator as a *weighted Average Adaptive Augmented IPW* (A3IPW) estimator. For $\{\hat{f}_t\}_{t=0}^{T-1}$ and $\sigma_t^2 = \text{Var}(\check{z}_t | \Omega_{t-1})$, we define the A3IPW estimator as $\hat{\theta}_T^{\text{A3IPW}} = \left(\sum_{t=1}^T \frac{1}{\sqrt{\sigma_t^2}}\right)^{-1} \sum_{t=1}^T \frac{1}{\sqrt{\sigma_t^2}} \sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t) \mathbb{1}[A_t=a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\}$. Then, we can show that the estimator has the asymptotic normality as follows.

Theorem 1. *Suppose that there exists C_2 such that $|\hat{f}_t| \leq C_2$. Then, under Assumption 2, $\left(\sum_{t=1}^T \frac{1}{\sqrt{\sigma_t^2}}\right) \sqrt{T} \left(\hat{\theta}_T^{\text{A3IPW}} - \theta_0\right) \xrightarrow{d} \mathcal{N}(0, 1)$ as $T \rightarrow \infty$.*

Proof. For $\tilde{z}_t = \frac{1}{\sqrt{\sigma_t^2}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t) \mathbb{1}[A_t=a] \{y - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right)$, $\{\tilde{z}_t\}_{t=1}^T$ is an MDS because $\mathbb{E}[\tilde{z}_t | \Omega_{t-1}] = \frac{1}{\sqrt{\sigma_t^2}} \mathbb{E}[\check{z}_t | \Omega_{t-1}] = 0$. Because $\text{Var}\left(\tilde{z}_t / \sqrt{\sigma_t^2}\right) = \mathbb{E}[\mathbb{E}[(\check{z}_t - \mathbb{E}[\check{z}_t])^2 | \Omega_{t-1}] / \sigma_t^2] = \mathbb{E}[\mathbb{E}[\check{z}_t^2 | \Omega_{t-1}] / \sigma_t^2] = \mathbb{E}[\text{Var}(\check{z}_t | \Omega_{t-1}) / \sigma_t^2] = 1$, we have $(1/T) \sum_{t=1}^T \text{Var}(\tilde{z}_t) = 1$. Here, we used $\mathbb{E}[\check{z}_t | \Omega_{t-1}] = \mathbb{E}[\check{z}_t] = 0$. Besides, we can show that \tilde{z}_t is bounded from assumptions, and $(1/T) \sum_{t=1}^T \tilde{z}_t^2 \xrightarrow{P} 1$ from the weak law of large numbers of Proposition 5 in Appendix A. Then, we can apply CLT for an MDS. \square

4.3 OPE with a Weighted Average Estimator: Case with Unknown Variance

In general, the variance σ_t^2 is unknown and needed to be replaced with an estimator. However, we cannot use the samples used for constructing π_t and \hat{f}_{t-1} to estimate σ_t^2 due to the dependency. For this reason, we assume access to a dataset only with covariates, $\mathcal{E}_N = \{X_i\}_{i=1}^N$, which is independent from the historical data \mathcal{S}_T . Because \mathcal{E}_N is independent from $\mathcal{S}_T = \{(X_t, A_t, Y_t)\}_{t=1}^T$, we can approximate σ_t^2 by applying the law of large numbers to the sample average $g'_{t,N}$, defined as

$$\frac{1}{N} \sum_{i=1}^N \sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_i))^2 (\hat{e}_{t-1}(a, X_i) - \hat{f}_{t-1}^2(a, X_i))}{\pi_t(a | X_i, \Omega_{t-1})} + \left(\pi^e(a | X_i) \hat{f}_{t-1}(a, X_i) - \hat{\theta}_{t-1}\right)^2 \right\}.$$

The consistency of $g'_{t,N}$ to the variance of $\psi_t(\theta_0; X_t, A_t, Y_t, \pi_t, \pi^e, f^*)$ is showed as follows.

Theorem 2. Suppose that, for all $x \in \mathcal{X}$ and $a \in \mathcal{A}$, $\hat{f}_{t-1}(a, x) - f^*(a, x) \xrightarrow{P} 0$, $\hat{e}_{t-1}(a, x) - e^*(a, x) \xrightarrow{P} 0$ and $\hat{\theta}_t \xrightarrow{P} \theta_0$ as $t \rightarrow \infty$. Then, for $\pi_t \in \Pi$, we have $g'_{t,(1-r)T} - \sigma_t^{*2} \xrightarrow{P} 0$ as $t \rightarrow \infty$ and $N \rightarrow \infty$, where

$$\sigma_t^{*2} = \mathbb{E} \left[\sum_{a=1}^K \left\{ \frac{(\pi^e(a|X_t))^2 \nu^*(a, X_t)}{\pi_t(a|X_t, \Omega_{t-1})} + \left(\pi^e(a | X_t) f^*(a, X_t) - \theta_0 \right)^2 \right\} \right].$$

In practice, we need to avoid having the estimator of σ_t^{*2} be less than or equal to 0. Thus, instead of $g'_{t,(1-r)T}$, we use the estimator $g_{t,N}$, defined as $g_{t,(1-r)T} = \max\{g'_{t,N}, \epsilon\}$, where $\epsilon > 0$ is the lower bound of the variance σ_t^{*2} . The estimator $g_{t,N}$ is consistent, as is $g'_{t,N}$. The value ϵ is assumed to be known, and we can use any sufficiently small value for ϵ because it is simply a technical term. The proof is shown in Appendix B. Then, using $g_{t,N}$, we define the *Feasible A3IPW* (FA3IPW) estimator $\hat{\theta}_T^{\text{FA3IPW}}$ as $\left(\sum_{t=1}^T \frac{1}{\sqrt{g_{t,N}}} \right)^{-1} \sum_{t=1}^T \sum_{a=1}^K \frac{1}{\sqrt{g_{t,N}}} \left\{ \frac{\pi^e(a|X_t) \mathbb{1}[A_t=a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\}$. Similarly, we can also define the corresponding IPW estimator and refer to it as *Feasible weighted Average Adaptive IPW estimator* (FA2aIPW).

In some applications, we can obtain the dataset $\mathcal{E}_N = \{X_i\}_{i=1}^N$ as evaluation (test) data, which is the target of a new policy (Kato et al., 2020b). On the other hand, by introducing a sample splitting, we can relax the assumption as the following remark.

Remark 7 (Sample Splitting). To relax the access to \mathcal{E}_N , we consider splitting $\{(X_t, A_t, Y_t)\}_{t=1}^T$ into $\{(X_t, A_t, Y_t)\}_{t=1}^{\lfloor rT \rfloor}$ and $\{(X_t, A_t, Y_t)\}_{t=\lfloor rT \rfloor+1}^T$, where $0 < r < 1$. In the following discussion, for simplicity, we assume that rT is an integer (i.e., $\lfloor rT \rfloor = rT$). The samples $\{X_t\}_{t=rT}^T$ and $\{(X_t, A_t, Y_t)\}_{t=1}^{rT}$ are independent. Thus, to approximate σ_t^2 , we can apply the law of large numbers to the sample average $g_{t,(1-r)T} = \max\{g'_{t,(1-r)T}, \epsilon\}$, where $g'_{t,(1-r)T}$ is defined as $\frac{1}{(1-r)T} \sum_{s=rT}^T \sum_{a=1}^K \left\{ \frac{(\pi^e(a|X_s))^2 (\hat{e}_{t-1}(a, X_s) - \hat{f}_{t-1}^2(a, X_s))}{\pi_t(a|X_s, \Omega_{t-1})} + \left(\pi^e(a | X_s) \hat{f}_{t-1}(a, X_s) - \hat{\theta}_{t-1} \right)^2 \right\}$. More details on sample splitting are provided in Appendix D.

4.4 Asymptotic Properties of a FA3IPW Estimator

Let $\hat{z}_t = \sum_{t=1}^T \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t) \mathbb{1}[A_t=a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a | X_t) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right)$. We can show that \hat{z}_t is an MDS as $\mathbb{E}[\hat{z}_t | \Omega_{t-1}] = \frac{1}{\sqrt{g_{t,N}}} \mathbb{E}[\hat{z}_t | \Omega_{t-1}] = 0$. Based on this property, the following theorems gives us the consistency and asymptotic normality of $\hat{\theta}_T^{\text{FA3IPW}}$. First, we show the consistency of $\hat{\theta}_T^{\text{FA3IPW}}$, i.e., $\hat{\theta}_T^{\text{FA3IPW}} \xrightarrow{P} \theta_0$.

Theorem 3 (Consistency of the Proposed Estimator). Suppose that there exist C_2 and C_3 such that $|\hat{f}_t| \leq C_2$ and $0 < 1/\sqrt{g_{t,N}} \leq C_3$. Then, under Assumptions 2, $\hat{\theta}_T^{\text{FA3IPW}} \xrightarrow{P} \theta_0$.

Proof. From the boundedness, we can apply the law of large numbers for an MDS (Proposition 5 in Appendix A). Therefore, $\frac{1}{T} \sum_{t=1}^T \hat{z}_t \xrightarrow{P} 0$. \square

The following theorem shows the asymptotic normality of $\hat{\theta}_T^{\text{FA3IPW}}$. The proof is in Appendix C.

Theorem 4 (Asymptotic Distribution of the Proposed Estimator). Suppose that

- (i) there exist C_2 and C_3 such that $|\hat{f}_t| \leq C_2$ and $0 < 1/\sqrt{g_{t,N}} \leq C_3$;
- (ii) Pointwise convergence in probability of \hat{f}_t , i.e., $\forall x \in \mathcal{X}$, $\hat{f}_t(a, x) - f^*(a, x) \xrightarrow{P} 0$ as $t \rightarrow \infty$;
- (iii) $g_{t,N} - \sigma_t^{*2} \xrightarrow{P} 0$ as $t \rightarrow \infty$ and $N \rightarrow \infty$.

Then, under Assumptions 2, $\left(\frac{1}{\sqrt{T}} \sum_{t=1}^T \frac{1}{\sqrt{g_{t,N}}} \right) \left(\hat{\theta}_T^{\text{FA3IPW}} - \theta_0 \right) \xrightarrow{d} \mathcal{N}(0, 1)$ as $T, N \rightarrow \infty$.

Algorithm 1 TSFA3IPW Estimation

For $t = 1, 2, \dots, T$, let $q_t = \sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t)\mathbb{1}[A_t=a]\{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a|X_t)\hat{f}_{t-1}(a, X_t) \right\}$.

Initialization: Let \tilde{g}_t be 1 or some other positive value for $t = 1, 2, \dots, T - 1$.

Obtain $\{\tilde{\theta}_t\}_{t=1}^{T-1}$ using $\{\tilde{g}_t\}_{t=1}^{T-1}$ by $\tilde{\theta}_t = \left(\sum_{s=1}^t \frac{1}{\sqrt{g_s}}\right)^{-1} \sum_{s=1}^t \frac{1}{\sqrt{g_s}} q_s$.

Obtain $\{g_{t,N}\}_{t=1}^T$ using $\{\tilde{\theta}_t\}_{t=1}^{T-1}$, an appropriate value $\hat{\theta}_0^{(1)}$ such as $\hat{\theta}_0^{(1)} = 0$, and \mathcal{E}_N .

Obtain $\hat{\theta}_T^{\text{TSFA3IPW}}$ using $\{g_{t,N}\}_{t=1}^T$ by $\hat{\theta}_T^{\text{TSFA3IPW}} = \left(\sum_{s=1}^T \frac{1}{\sqrt{g_{s,N}}}\right)^{-1} \sum_{s=1}^T \frac{1}{\sqrt{g_{s,N}}} q_s$.

Remark 8 (Difference of Assumption between the Consistency and Asymptotic Normality). The asymptotic normality requires $g_{t,N} - \sigma_t^{*2} \xrightarrow{P} 0$. However, we can obtain the consistency $\hat{\theta}_T^{\text{A3IPW}} \xrightarrow{P} \theta_0$ without assuming $g_{t,N} - \sigma_t^{*2} \xrightarrow{P} 0$. From this property, we can obtain the consistent estimator of the variance, σ_t^{*2} , which requires the consistent estimator of the policy value, θ_0 .

Remark 9 (Heuristic Stabilization). The proposed estimator requires a sequence of $\{f_t\}_{t=0}^{T-1}$. Because the early estimator of f_t may be unstable due to the insufficient amount of training data, some heuristic stabilizations may improve its performance. Hadad et al. (2019) proposed a similar estimator and suggested using a weight sequence for stabilization. In the setting of Kato et al. (2020a), they can change $p_t(a|x, \Omega_{t-1})$ and proposed adjusting the probability to stabilize their proposed estimator. In this study, we take a simple approach by suggesting discarding some samples at first and constructing the proposed estimator without them. We call this estimator the *stabilized FA3IPW* (SFA3IPW) estimator. In some experiments, this approach improves the estimator's performance.

4.5 Efficiency of A3IPW Estimator

In the problem setting, it is not easy to discuss an efficiency of an estimator. For simplicity, we consider the lower bound of the variance of an estimator within a class of estimators such that $\hat{\theta}_T = \sum_{t=1}^T w_t \tilde{\psi}_t(X_t, A_t, Y_t, \pi_t)$, where $\tilde{\psi}$ is a score function and w_t is a weight satisfying $\sum_{t=1}^T w_t = 1$. Let us note that we usually use $w_t = 1/T$ in the case where samples are i.i.d. First, we consider the variance of $\tilde{\psi}_t$. In OPE, the lowest variance is given as $\mathbb{E} \left[\sum_{a=1}^K \left\{ \frac{(\pi^e(a|X_t))^2 v^*(a, X_t)}{\pi_t(a|X_t, \Omega_{t-1})} + \left(\pi^e(a|X_t) f^*(a, X_t) - \theta_0 \right)^2 \right\} \right]$. In the proposed estimator, each element of $\{\tilde{\psi}_t\}_{t=1}^T$ achieves this variance. In addition, it is well known that for a sequence of positive values $\phi_t = \mathbb{E}[\tilde{\psi}_t^2 | \Omega_{t-1}]$, the weight $w_t = \frac{1}{\sqrt{\phi_t}} / \sum_{i=1}^T \frac{1}{\sqrt{\phi_i}}$ minimizes the variance of $\hat{\theta}_T$ when the covariances among the score functions are 0. When $\{\phi_t\}_{t=1}^T$ is an MDS, the covariance is 0. Thus, the proposed estimator uses the weights and scores that minimize the variance for an estimator with a form that satisfies $\sum_{t=1}^T w_t \tilde{\psi}_t(X_t, A_t, Y_t, \pi_t)$.

5 Main Algorithm: Two-step FA3IPW Estimation

Under the assumption of $g_{t,N} - \sigma_t^{*2} \xrightarrow{P} 0$, the proposed estimator $\hat{\theta}_T^{\text{A3IPW}}$ has the asymptotic normality. However, to estimate $\sigma_t^{*2} = \text{Var}(\tilde{z}_t | \Omega_{t-1})$, a consistent estimator of θ_0 is required, which is why we use $g_{t,N}$. However, we can obtain a consistent estimator of θ_0 without imposing the assumption that $g_{t,N} - \sigma_t^{*2} \xrightarrow{P} 0$. Based on these properties, we propose *two-step estimation*. First, using arbitrary appropriate values for $g_{t,N} = \tilde{g}_t$ for $t = 1, 2, \dots, T - 1$, such as $\tilde{g}_t = 1$, we obtain the initial estimates, $\{\tilde{\theta}_t\}_{t=1}^{T-1}$. Then, using $\tilde{\theta}_t \xrightarrow{P} \theta_0$, we construct $g_{t,N} - \sigma_t^{*2} \xrightarrow{P} 0$. Finally, we can estimate the next estimator of θ_0 , $\hat{\theta}_T^{\text{TSFA3IPW}}$, with asymptotic normality from Theorem 4. We summarize the process of *two-step FA3IPW* (TSFA3IPW) estimation in Algorithm 1.

Remark 10 (Estimation of Functions f_t^* and e_t^*). Several nonparametric regression estimators can be proved to be consistent for dependent samples via certain bandit algorithms (Yang & Zhu, 2002).

Table 1: Experimental results under the RW policy. The method with the lowest MSE is in bold.

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.037	0.193	0.064	0.254	0.048	0.219	0.032	0.178	0.128	0.358	0.101	0.313	0.029	0.163	0.045	0.210
SFA3IPW	0.037	0.192	0.061	0.243	0.051	0.225	0.029	0.169	0.142	0.376	0.096	0.308	0.033	0.178	0.042	0.206
DM	0.101	0.168	0.456	0.171	0.281	0.179	0.209	0.122	0.489	0.143	0.413	0.153	0.100	0.158	0.109	0.165
AdaIPW	0.068	0.261	0.089	0.297	0.069	0.263	0.063	0.251	0.129	0.359	0.133	0.360	0.089	0.242	0.070	0.252
A2IPW	0.038	0.193	0.063	0.250	0.049	0.220	0.035	0.185	0.119	0.345	0.100	0.314	0.032	0.164	0.052	0.226
FA2daIPW	0.068	0.261	0.090	0.301	0.074	0.273	0.057	0.238	0.140	0.373	0.133	0.358	0.075	0.237	0.059	0.236

Table 2: Experimental results under the UCB policy. The method with the lowest MSE is in bold.

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.041	0.201	0.095	0.308	0.098	0.306	0.026	0.160	0.452	0.657	0.077	0.274	3.484	1.798	0.066	0.257
SFA3IPW	0.042	0.198	0.071	0.263	0.126	0.353	0.029	0.167	0.488	0.679	0.070	0.262	5.248	2.240	0.077	0.275
DM	0.022	0.089	0.218	0.114	0.245	0.168	0.195	0.130	0.489	0.144	0.401	0.141	0.112	0.201	0.107	0.172
AdaIPW	0.078	0.275	0.099	0.314	0.188	0.423	0.051	0.223	0.891	0.929	0.101	0.315	11.424	2.560	0.072	0.267
A2IPW	0.043	0.206	0.084	0.290	0.097	0.304	0.025	0.158	0.854	0.909	0.083	0.286	2.363	1.400	0.060	0.245
FA2daIPW	0.072	0.263	0.124	0.351	0.173	0.398	0.052	0.225	0.481	0.680	0.094	0.304	11.031	2.628	0.079	0.281

6 Experiments

In this section, using benchmark datasets, we demonstrate the effectiveness of the proposed FA3IPW estimator and the FA3IPW estimator with discarding some early samples for stabilization (SFA3IPW estimator in Remark 9). Following Dudík et al. (2011) and Farajtabar et al. (2018), we evaluate the proposed estimators using classification datasets by transforming them into contextual bandit data. From the LIBSVM repository, we use the satImage, pendigits, mnist, dna, letter, sensorless, connect-4, and covtype datasets³. First, we create an adaptive policy π_t^a by following a random walk and perform upper confidence bound (UCB) algorithms in the MAB problem. Then, we construct a behavior policy as a mixture of π_t^a and the uniform random policy π_t^u , defined as $\pi_t = 0.7\pi_t^a + 0.3\pi_t^u$ (Kallus & Uehara, 2019a). When we use the RW policy, the policy π_t^a does not converge. In contrast, when we use the UCB policy to generate π_t^a , the policy converges to a time-invariant policy. To construct an evaluation policy, we create a deterministic policy π_d by training a logistic regression classifier on historical data. Then, we construct the evaluation policy π^e as a mixture of π^d and the uniform random policy π^u , defined as $\pi^e = 0.7\pi^d + 0.3\pi^u$. Through experiments, the behavior policy π_t is assumed to be known. More details are in Appendix E.

We compare the MSEs of six estimators, the FA3IPW, SFA3IPW, DM, AdaIPW, A2IPW, and FA2daIPW estimators. For the FA3IPW, SFA3IPW, and FA2daIPW estimators, we applied the two-step estimation. In each experiment, we have access to historical data with $T = 1000$ and evaluation data with $N = 1000$. With the SFA3IPW estimator, we discard the first 1000 samples of the historical data. When estimating f^* and e^* , we use the Nadaraya-Watson regression (NW) estimator, which is used in Yang & Zhu (2002). The resulting MSEs and their standard deviations (SDs) over 20 replications of each experiment are shown in Tables 1 and 2. In the experimental results, when the RW policy is used, the FA3IPW and SFA3IPW estimators show preferable performances compared with the A2IPW estimator. This is because the performance of the A2IPW estimator is guaranteed only when the policy converges to a time-invariant policy. On the other hand, when the UCB policy is used, the A2IPW estimator also works well as we can expect. From the results, we recommend using the FA3IPW estimator when the policy does not converge. More importantly, in the case, we can construct confidence intervals from the FA3IPW estimator, but cannot construct it from the A2IPW estimator. In Appendix E, we show the experimental results with other settings.

7 Conclusion

This study presented solutions for causal inference from dependent samples obtained via bandit feedback. By utilizing evaluation data or sample splitting, we can standardize the variance of an MDS. Then, we can apply the CLT for an MDS to obtain the asymptotic normality of the proposed estimator. In experiments, the proposed estimators showed theoretically expected performances.

³<https://www.csie.ntu.edu.tw/~cjlin/libsvmtools/datasets/>

Broader Impact

In various applications, such as A/B testing and clinical trials, the problem of causal inference from dependent samples arises. Because this study promotes the use of OPE for those applications by presenting an effective solution to this problem, the ethical issues related to such applications will become more critical. For example, in clinical trials, the proposed method also should take the criteria proposed by the US Food and Drug Administration (FDA, 2019) into account.

Besides, although this study solved an essential problem with OPE, the limitations of methods for OPE should still be recognized. The existing methods for OPE try to find causality only from data without requiring domain-specific knowledge, but they sometimes lead to inappropriate decision making by producing incorrect results. When finding causality in the real world, it might be desirable to utilize domain-specific knowledge combined with the results obtained from machine learning methods.

References

- Athey, S. and Wager, S. Efficient policy learning. *arXiv preprint arXiv:1702.02896*, 2017.
- Beygelzimer, A. and Langford, J. The offset tree for learning with partial labels. In *KDD*, pp. 129–138, 2009.
- Bibaut, A., Malenica, I., Vlassis, N., and Van Der Laan, M. More efficient off-policy evaluation through regularized targeted learning. In *ICML*, volume 97, pp. 654–663. PMLR, 2019.
- Bickel, P. J., Klaassen, C. A. J., Ritov, Y., and Wellner, J. A. *Efficient and Adaptive Estimation for Semiparametric Models*. Springer, 1998.
- Brown, B. M. Martingale central limit theorems. *Ann. Math. Statist.*, 42(1):59–66, 02 1971.
- Chernozhukov, V., Chetverikov, D., Demirer, M., Duflo, E., Hansen, C., Newey, W., and Robins, J. Double/debiased machine learning for treatment and structural parameters. *Econometrics Journal*, 21:C1–C68, 2018.
- Chernozhukov, V., Demirer, M., Lewis, G., and Syrgkanis, V. Semi-parametric efficient policy learning with continuous actions. In *NeurIPS*. 2019.
- Chu, W., Li, L., Reyzin, L., and Schapire, R. Contextual bandits with linear payoff functions. In *AISTATS*, volume 15 of *Proceedings of Machine Learning Research*, pp. 208–214, 2011.
- der Laan, V. and Mark, J. I. The construction and analysis of adaptive group sequential designs. 2008.
- Dudík, M., Langford, J., and Li, L. Doubly Robust Policy Evaluation and Learning. In *ICML*, pp. 1097–1104, 2011.
- Farajtabar, M., Chow, Y., and Ghavamzadeh, M. More robust doubly robust off-policy evaluation. *ICML*, pp. 1447–1456, 2018.
- FDA. Adaptive designs for clinical trials of drugs and biologics: Guidance for industry. Technical report, U.S. Department of Health and Human Services Food and Drug Administration (FDA), Center for Drug Evaluation and Research (CDER), Center for Biologics Evaluation and Research (CBER), 2019.
- Greene, W. *Econometric Analysis*. Pearson Education, 2003.
- Hadad, V., Hirshberg, D. A., Zhan, R., Wager, S., and Athey, S. Confidence intervals for policy evaluation in adaptive experiments, 2019.
- Hahn, J. On the role of the propensity score in efficient semiparametric estimation of average treatment effects. *Econometrica*, 66:315–331, 1998.
- Hahn, J., Hirano, K., and Karlan, D. Adaptive experimental design using the propensity score. *Journal of Business and Economic Statistics*, 29(1):96–108, 2011.
- Hall, P., Heyde, C., Birnbaum, Z., and Lukacs, E. *Martingale Limit Theory and Its Application*. Communication and Behavior. Elsevier Science, 2014.
- Hamilton, J. *Time series analysis*. Princeton Univ. Press, 1994.
- Hirano, K., Imbens, G. W., and Ridder, G. Efficient estimation of average treatment effects using the estimated propensity score. *Econometrica*, 71(4):1161–1189, 2003.
- Horvitz, D. G. and Thompson, D. J. A generalization of sampling without replacement from a finite universe. *Journal of the American Statistical Association*, 47(260):663–685, 1952.
- Kallus, N. and Uehara, M. Intrinsically efficient, stable, and bounded off-policy evaluation for reinforcement learning. In *NeurIPS*, pp. 3320–3329. 2019a.

- Kallus, N. and Uehara, M. Efficiently breaking the curse of horizon: Double reinforcement learning in infinite-horizon processes. *arXiv preprint arXiv:1909.05850*, 2019b.
- Kato, M., Ishihara, T., Honda, J., and Narita, Y. Adaptive experimental design for efficient treatment effect estimation: Randomized allocation via contextual bandit algorithm. *arXiv:2002.05308*, 2020a.
- Kato, M., Uehara, M., and Yasui, S. Off-policy evaluation and learning for external validity under a covariate shift, 2020b.
- Kitagawa, T. and Tetenov, A. Who should be treated? empirical welfare maximization methods for treatment choice. *Econometrica*, 86:591–616, 2018.
- Kosorok, M. R. *Introduction to Empirical Processes and Semiparametric Inference*. Springer Series in Statistics. Springer New York, New York, NY, 2008.
- Li, L., Chu, W., Langford, J., and Schapire, R. E. A contextual-bandit approach to personalized news article recommendation. In *WWW*, pp. 661–670, 2010.
- Loeve, M. *Probability Theory*. Graduate Texts in Mathematics. Springer, 1977.
- Luedtke, A. R. and van der Laan, M. J. Statistical inference for the mean outcome under a possibly non-unique optimal treatment strategy. *Annals of statistics*, 44(2):713–742, Apr 2016.
- Narita, Y., Yasui, S., and Yata, K. Efficient counterfactual learning from bandit feedback. *AAAI*, 2019.
- Oberst, M. and Sontag, D. Counterfactual off-policy evaluation with gumbel-max structural causal models. In *ICML*, volume 97, pp. 4881–4890, 2019.
- Robins, J. M., Rotnitzky, A., and Zhao, L. P. Estimation of regression coefficients when some regressors are not always observed. *Journal of the American Statistical Association*, 89:846–866, 1994.
- Rubin, D. B. *Multiple Imputation for Nonresponse in Surveys*. Wiley, New York, 1987.
- Scharfstein, D., Rotnitzky, A., and Robins, J. M. Adjusting for nonignorable dropout using semi-parametric models. *Journal of the American Statistical Association*, 94:1096–1146, 1999.
- van der Vaart, A. W. *Asymptotic statistics*. Cambridge University Press, Cambridge, UK, 1998.
- Wang, Y.-X., Agarwal, A., and Dudik, M. Optimal and adaptive off-policy evaluation in contextual bandits. In *Proceedings of the 34th International Conference on Machine Learning*, pp. 3589–3597, 2017.
- Yang, Y. and Zhu, D. Randomized allocation with nonparametric estimation for a multi-armed bandit problem with covariates. *Ann. Statist.*, 30(1):100–121, 02 2002. doi: 10.1214/aos/1015362186.
- Zhao, Y., Zeng, D., Rush, A. J., and Kosorok, M. R. Estimating individualized treatment rules using outcome weighted learning. *Journal of the American Statistical Association*, 107:1106–1118, 2012.
- Zhou, Z., Athey, S., and Wager, S. Offline multi-action policy learning: Generalization and optimization. *arxiv preprint arxiv:1810.04778*, 2018.

A Preliminaries

A.1 Mathematical Tools

Proposition 2. [Slutsky Theorem, Greene (2003), Theorem D. 16 1, p. 1117] If $a_n \xrightarrow{d} a$ and $b_n \xrightarrow{p} b$, then

$$a_n b_n \xrightarrow{d} ba.$$

Definition 1. [Uniformly Integrable, Hamilton (1994), p. 191] A sequence $\{A_t\}$ is said to be uniformly integrable if for every $\epsilon > 0$ there exists a number $c > 0$ such that

$$\mathbb{E}[|A_t| \cdot \mathbb{1}[|A_t| \geq c]] < \epsilon$$

for all t .

Proposition 3. [Sufficient Conditions for Uniformly Integrable, Hamilton (1994), Proposition 7.7, p. 191] (a) Suppose there exist $r > 1$ and $M < \infty$ such that $\mathbb{E}[|A_t|^r] < M$ for all t . Then $\{A_t\}$ is uniformly integrable. (b) Suppose there exist $r > 1$ and $M < \infty$ such that $\mathbb{E}[|b_t|^r] < M$ for all t . If $A_t = \sum_{j=-\infty}^{\infty} h_j b_{t-j}$ with $\sum_{j=-\infty}^{\infty} |h_j| < \infty$, then $\{A_t\}$ is uniformly integrable.

Proposition 4 (L^r Convergence Theorem, Loeve (1977)). Let $0 < r < \infty$, suppose that $\mathbb{E}[|a_n|^r] < \infty$ for all n and that $a_n \xrightarrow{p} a$ as $n \rightarrow \infty$. The following are equivalent:

- (i) $a_n \rightarrow a$ in L^r as $n \rightarrow \infty$;
- (ii) $\mathbb{E}[|a_n|^r] \rightarrow \mathbb{E}[|a|^r] < \infty$ as $n \rightarrow \infty$;
- (iii) $\{|a_n|^r, n \geq 1\}$ is uniformly integrable.

A.2 Martingale Limit Theorems

Proposition 5. [Weak Law of Large Numbers for Martingale, Hall et al. (2014)] Let $\{S_n = \sum_{i=1}^n X_i, \Omega_t, t \geq 1\}$ be a martingale and $\{b_n\}$ a sequence of positive constants with $b_n \rightarrow \infty$ as $n \rightarrow \infty$. Then, writing $X_{ni} = X_i \mathbb{1}[|X_i| \leq b_n]$, $1 \leq i \leq n$, we have that $b_n^{-1} S_n \xrightarrow{p} 0$ as $n \rightarrow \infty$ if

- (i) $\sum_{i=1}^n P(|X_i| > b_n) \rightarrow 0$;
- (ii) $b_n^{-1} \sum_{i=1}^n \mathbb{E}[X_{ni} | \Omega_{t-1}] \xrightarrow{p} 0$, and;
- (iii) $b_n^2 \sum_{i=1}^n \{\mathbb{E}[X_{ni}^2] - \mathbb{E}[\mathbb{E}[X_{ni} | \Omega_{t-1}]]^2\} \rightarrow 0$.

Remark 11. The weak law of large numbers for martingale holds when the random variable is bounded by a constant.

B Proof of Theorem 2

Proof. The variance σ_t^{2*} can be calculated as follows:

$$\begin{aligned} \sigma_t^{*2} &= \mathbb{E} \left[\left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | X_t) f^*(a, X_t) \right\} - \theta_0 \right)^2 \right] \\ &= \mathbb{E} \left[\left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t(a) - f^*(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | X_t) f^*(a, X_t) \right\} - \theta_0 \right)^2 \right] \\ &= \mathbb{E} \left[\sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_t))^2 (e^*(a, X_t) - (f^*(a, X_t))^2)}{\pi_t(a | X_t, \Omega_{t-1})} + (\pi^e(a | X_t) f^*(a, X_t) - \theta_0)^2 \right\} \right]. \end{aligned}$$

Therefore, what we want to show is

$$\begin{aligned} & \frac{1}{N} \sum_{i=1}^N \sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_i))^2 (\hat{e}_{t-1}(a | X_i) - \hat{f}_{t-1}^2(a | X_i))}{\pi_t(a | X_i, \Omega_{t-1})} + \left(\pi^e(a | X_i) \hat{f}_{t-1}(a, X_i) - \hat{\theta}_{t-1} \right)^2 \right\} \\ & - \mathbb{E} \left[\sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_t))^2 (e^*(a, X_t) - (f^*(a, X_t))^2)}{\pi_t(a | X_t, \Omega_{t-1})} + \left(\pi^e(a | X_t) f^*(a, X_t) - \theta_0 \right)^2 \right\} \right] \\ & \xrightarrow{P} 0. \end{aligned}$$

From the assumptions, $\hat{f}_{t-1}(a, x) - f^*(a, x) \xrightarrow{P} 0$, $\hat{e}_{t-1}(a, x) - e^*(a, x) \xrightarrow{P} 0$, and $\hat{\theta}_{t-1} - \theta_0 \xrightarrow{P} 0$ as $t \rightarrow \infty$,

$$\begin{aligned} & \frac{1}{N} \sum_{t=1}^N \sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_i))^2 (\hat{e}_{t-1}(a | X_i) - \hat{f}_{t-1}^2(a | X_i))}{\pi_t(a | X_i, \Omega_{t-1})} + \left(\pi^e(a | X_i) \hat{f}_{t-1}(a, X_i) - \hat{\theta}_{t-1} \right)^2 \right\} \\ & - \frac{1}{N} \sum_{i=1}^N \sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_i))^2 (e^*(a, X_i) - (f^*(a, X_i))^2)}{\pi_t(a | X_i, \Omega_{t-1})} + \left(\pi^e(a | X_i) f^*(a, X_i) - \theta_0 \right)^2 \right\} \\ & \xrightarrow{P} 0 \text{ as } t \rightarrow \infty. \end{aligned}$$

Here, from the weak law of large numbers, as $N \rightarrow \infty$,

$$\begin{aligned} & \frac{1}{N} \sum_{i=1}^N \sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_i))^2 (e^*(a, X_i) - (f^*(a, X_i))^2)}{\pi_t(a | X_i, \Omega_{t-1})} + \left(\pi^e(a | X_i) f^*(a, X_i) - \theta_0 \right)^2 \right\} \\ & = \frac{1}{N} \sum_{i=1}^N \sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_i))^2 \nu^*(a, X_i)}{\pi_t(a | X_i, \Omega_{t-1})} + \left(\pi^e(a | X_i) f^*(a, X_i) - \theta_0 \right)^2 \right\} \\ & \xrightarrow{P} \mathbb{E} \left[\sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_i))^2 \nu^*(a, X_i)}{\pi_t(a | X_i, \Omega_{t-1})} + \left(\pi^e(a | X_i) f^*(a, X_i) - \theta_0 \right)^2 \right\} \right]. \end{aligned}$$

Therefore, as $t \rightarrow \infty$ and $N \rightarrow \infty$,

$$\begin{aligned} & \frac{1}{N} \sum_{i=1}^N \sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_i))^2 (\hat{e}_{t-1}(a | X_i) - \hat{f}_{t-1}^2(a | X_i))}{\pi_t(a | X_i, \Omega_{t-1})} + \left(\pi^e(a | X_i) \hat{f}_{t-1}(a, X_i) - \hat{\theta}_{t-1} \right)^2 \right\} \\ & - \mathbb{E} \left[\sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_i))^2 \nu^*(a, X_i)}{\pi_t(a | X_i, \Omega_{t-1})} + \left(\pi^e(a | X_i) f^*(a, X_i) - \theta_0 \right)^2 \right\} \right] \xrightarrow{P} 0. \end{aligned}$$

□

C Proof of Theorem 4

Proof. We have

$$\sqrt{T}(\hat{\theta}_T - \theta_0) = \left(\frac{1}{T} \sum_{t=1}^T \frac{1}{\sqrt{g_{t,N}}} \right)^{-1} \sqrt{T} \frac{1}{T} \sum_{t=1}^T \hat{z}_t,$$

where note that

$$\hat{z}_t = \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right).$$

If we show

$$\sqrt{T} \frac{1}{T} \sum_{t=1}^T \hat{z}_t \xrightarrow{d} \mathcal{N}(0, 1) \text{ as } T \rightarrow \infty, \quad (3)$$

then we can show that

$$\left(\frac{1}{T} \sum_{t=1}^T \frac{1}{\sqrt{g_{t,N}}} \right) \sqrt{T}(\hat{\theta}_T - \theta_0) = \sqrt{T} \frac{1}{T} \sum_{t=1}^T \hat{z}_t \xrightarrow{d} \mathcal{N}(0, 1).$$

To show (3), we use the central limit theorem for an MDS (Proposition 1 in Appendix A). Because $\{\hat{z}_t\}_{t=1}^T$ is an MDS, we check the following three conditions of Proposition 1:

- (a) $\mathbb{E}[\hat{z}_t^2] = \nu_t^2 > 0$ with $(1/T) \sum_{t=1}^T \nu_t^2 \rightarrow \nu^2 > 0$;
- (b) $\mathbb{E}[|\hat{z}_t|^r] < \infty$ for some $r > 2$;
- (c) $(1/T) \sum_{t=1}^T \hat{z}_t^2 \xrightarrow{P} \nu^2$.

Step 1: Check of Condition (a)

We have

$$\begin{aligned} \mathbb{E}[\hat{z}_t^2] &= \mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right] \\ &= \mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right] \\ &\quad - \mathbb{E} \left[\left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right] \\ &\quad + \mathbb{E} \left[\left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right]. \end{aligned}$$

Because the last term is 1, We have

$$\begin{aligned} &\mathbb{E}[\hat{z}_t^2] - 1 \\ &= \mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right] \\ &\quad - \mathbb{E} \left[\left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right]. \end{aligned}$$

First, we show that the first and second terms vanishes asymptotically. We have

$$\begin{aligned} &\mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right] \\ &\quad - \mathbb{E} \left[\left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right] \\ &\leq \mathbb{E} \left[\left| \left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{y - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right. \right. \\ &\quad \left. \left. - \left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right| \right]. \end{aligned}$$

From the boundedness of each variable in \hat{z}_t , there exist absolute constants \tilde{C}_1 and \tilde{C}_2 such that

$$\begin{aligned} &\leq \mathbb{E} \left[\left| \left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{y - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right. \right. \\ &\quad \left. \left. - \mathbb{E} \left[\left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right] \right| \right] \\ &\leq \mathbb{E} \left[\left| \frac{1}{\sqrt{g_{t,N}}} - \frac{1}{\sqrt{\sigma_t^{*2}}} \right| \tilde{C}_1 + \mathbb{E} \left[\left| \hat{f}_{t-1}(a, X_t) - f^*(a, X_t) \right| \right] \tilde{C}_2 \right]. \end{aligned}$$

From the continuous mapping theorem and assumption that $g_{t,N} - \sigma_t^{*2} \xrightarrow{P} 0$ as $t \rightarrow \infty$ and $N \rightarrow \infty$, we have $1/\sqrt{g_{t,N}} - 1/\sqrt{\sigma_t^{*2}} \xrightarrow{P} 0$ as $t \rightarrow \infty$ and $N \rightarrow \infty$ from the continuous mapping theorem. Then, if $1/\sqrt{g_{t,N}}$ is uniformly integrable, we can prove $\mathbb{E}[|1/\sqrt{g_{t,N}} - 1/\sqrt{\sigma_t^{*2}}|] \rightarrow 0$ as $t \rightarrow \infty$ using L^r -convergence theorem (Proposition 4 in Appendix A).

Because $\hat{f}_{t-1}(a, x)$ is independent from X_t , we have $\mathbb{E}[\hat{f}_{t-1}(a, X_t) | X_t = x] = \mathbb{E}[\hat{f}_{t-1}(a, x)]$. For $\mathbb{E}[\hat{f}_{t-1}(a, x)]$ and fixed x , from the point convergence of $\hat{f}_{t-1}(a, x)$, we have

$$\mathbb{E} \left[\left| \hat{f}_{t-1}(a, X_t) - \mathbb{E}[f^*(a, X_t)] \right| | X_t = x \right] = \mathbb{E} \left[\left| \hat{f}_{t-1}(a, x) - \mathbb{E}[f^*(a, x)] \right| \right] \rightarrow 0.$$

Then, from the Lebesgue's dominated convergence theorem, for $\mathbb{E}[\hat{f}_{t-1}(a, X_t) | X_t = x]$, which pointwisely converges to $\mathbb{E}[f^*(a, X_t) | X_t = x]$, we can show that

$$\mathbb{E}_{X_t} \left[\left| \mathbb{E}[\hat{f}_{t-1}(a, X_t) | X_t] - \mathbb{E}[f^*(a, X_t) | X_t] \right| \right] \rightarrow 0.$$

We can show that $1/\sqrt{g_{t,N}}$ and \hat{f}_{t-1} are uniformly integrable from the boundedness of $1/\sqrt{g_{t,N}}$ and \hat{f}_{t-1} (Proposition 3 in Appendix A). Then, as $t \rightarrow \infty$ and $T \rightarrow \infty$,

$$\begin{aligned} &\mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right. \\ &\quad \left. - \mathbb{E} \left[\left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right] \right] \\ &\leq \mathbb{E} \left[\left| \frac{1}{\sqrt{g_{t,N}}} - \frac{1}{\sqrt{\sigma_t^{*2}}} \right| \tilde{C}_1 + \mathbb{E} \left[\left| \hat{f}_{t-1}(a, X_t) - f^*(a, X_t) \right| \right] \tilde{C}_2 \right] \rightarrow 0. \end{aligned}$$

Therefore, for any $\epsilon > 0$, there exists $\tilde{t} > 0$ and $\tilde{T} > 0$ such that, for $T > \max\{\tilde{T}, \tilde{t}/r\}$,

$$\begin{aligned} &\mathbb{E}[\hat{z}_t^2] - 1 \\ &= \frac{1}{T} \sum_{t=1}^T \left(\mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a | X_t) \mathbb{1}[A_t = a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right. \right. \\ &\quad \left. \left. - \mathbb{E} \left[\left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{(\pi^e(a | X_t))^2 \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\pi_t(a | X_t, \Omega_{t-1})} + \pi^e(a | x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right] \right] \right) \\ &\leq \tilde{t}/T + \epsilon. \end{aligned}$$

Thus, we have $(1/T) \sum_{t=1}^T \nu_t^2 - 1 \leq \tilde{t}/T + \epsilon \rightarrow 0$ as $T \rightarrow \infty$.

Step 2: Check of Condition (b)

We can prove this condition using the assumption of the boundedness of the random variables.

Step 3: Check of Condition (c)

To show $(1/T) \sum_{t=1}^T \hat{z}_t^2 \xrightarrow{P} \nu^2$, we show (i) $(1/T) \sum_{t=1}^T \hat{z}_t^2 - (1/T) \sum_{t=1}^T \mathbb{E}[\hat{z}_t^2 | \Omega_{t-1}] \xrightarrow{P} 0$; (ii) $(1/T) \sum_{t=1}^T \mathbb{E}[\hat{z}_t^2 | \Omega_{t-1}] - \nu^2 \xrightarrow{P} 0$, where $\nu^2 = \mathbb{E} \left[\left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{(\pi^e(a|X_t))^2 \mathbb{1}_{[A_t=a]} \{Y_t - f^*(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a|x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right] = 1$.

Proof of (i): We can prove (i) directly applying law of large numbers for an MDS to an MDS $\phi_t = \hat{z}_t^2 - \mathbb{E}[\hat{z}_t^2 | \Omega_{t-1}]$. The conditions for using the law of large numbers are satisfied by the boundedness of components of \hat{z}_t .

Proof of (ii) Next, we show that

$$\frac{1}{T} \sum_{t=1}^T \mathbb{E}[\hat{z}_t^2 | \Omega_{t-1}] - \nu^2 \xrightarrow{P} 0.$$

From Markov's inequality, for $\varepsilon > 0$, we have

$$\mathbb{P} \left(\left| \frac{1}{T} \sum_{t=1}^T \mathbb{E}[\hat{z}_t^2 | \Omega_{t-1}] - \nu^2 \right| \geq \varepsilon \right) \leq \frac{\mathbb{E} \left[\left| \frac{1}{T} \sum_{t=1}^T \mathbb{E}[\hat{z}_t^2 | \Omega_{t-1}] - \nu^2 \right| \right]}{\varepsilon} \leq \frac{\frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[\left| \mathbb{E}[\hat{z}_t^2 | \Omega_{t-1}] - \nu^2 \right| \right]}{\varepsilon}.$$

Then, we consider showing $\mathbb{E} \left[\left| \mathbb{E}[\hat{z}_t^2 | \Omega_{t-1}] - \nu^2 \right| \right] \rightarrow 0$. Here, we have

$$\begin{aligned} & \mathbb{E} \left[\left| \mathbb{E}[\hat{z}_t^2 | \Omega_{t-1}] - \nu^2 \right| \right] \\ &= \mathbb{E} \left[\left| \mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t) \mathbb{1}_{[A_t=a]} \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a|x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right. \right. \right. \\ & \quad \left. \left. \left. - \left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t) \mathbb{1}_{[A_t=a]} \{Y_t - f^*(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a|x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right| \Omega_{t-1} \right] \right| \right] \\ &= \mathbb{E} \left[\left| \mathbb{E} \left[\mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t) \mathbb{1}_{[A_t=a]} \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a|x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right. \right. \right. \right. \right. \\ & \quad \left. \left. \left. - \left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t) \mathbb{1}_{[A_t=a]} \{Y_t - f^*(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a|x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right| X_t, \Omega_{t-1} \right] \right| \Omega_{t-1} \right] \right| \right]. \end{aligned}$$

Then, by using Jensen's inequality,

$$\begin{aligned} & \mathbb{E} \left[\left| \mathbb{E}[\hat{z}_t^2 | \Omega_{t-1}] - \nu^2 \right| \right] \\ & \leq \mathbb{E} \left[\left| \mathbb{E} \left[\mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t) \mathbb{1}_{[A_t=a]} \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a|x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right. \right. \right. \right. \right. \\ & \quad \left. \left. \left. - \left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t) \mathbb{1}_{[A_t=a]} \{Y_t - f^*(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a|x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right| \Omega_{t-1} \right] \right] \right| \right] \\ & = \mathbb{E} \left[\left| \mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t) \mathbb{1}_{[A_t=a]} \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a|x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right. \right. \right. \right. \\ & \quad \left. \left. \left. - \left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a|X_t) \mathbb{1}_{[A_t=a]} \{Y_t - f^*(a, X_t)\}}{\pi_t(a|X_t, \Omega_{t-1})} + \pi^e(a|x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right| X_t, \Omega_{t-1} \right] \right] \right| \right]. \end{aligned}$$

Because \hat{f}_{t-1} and $g_{t,N}$ are constructed from Ω_{t-1} ,

$$\begin{aligned} & \mathbb{E} \left[\left| \mathbb{E} [z_t^2 \mid \Omega_{t-1}] - \nu^2 \right| \right] \\ & \leq \mathbb{E} \left[\left| \mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a \mid X_t) \mathbb{1}[A_t = a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a \mid X_t, \Omega_{t-1})} + \pi^e(a \mid x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right. \right. \right. \\ & \quad \left. \left. \left. - \left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a \mid X_t) \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\pi_t(a \mid X_t, \Omega_{t-1})} + \pi^e(a \mid x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \mid X_t, \hat{f}_{t-1}, g_{t,N} \right] \right| \right]. \end{aligned}$$

From the results of Step 1, there exist \tilde{C}_1 and \tilde{C}_2 such that

$$\begin{aligned} & \mathbb{E} \left[\left| \mathbb{E} [z_t^2 \mid \Omega_{t-1}] - \nu^2 \right| \right] \\ & \leq \mathbb{E} \left[\left| \mathbb{E} \left[\left\{ \frac{1}{\sqrt{g_{t,N}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a \mid X_t) \mathbb{1}[A_t = a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a \mid X_t, \Omega_{t-1})} + \pi^e(a \mid x) \hat{f}_{t-1}(a, X_t) \right\} - \theta_0 \right) \right\}^2 \right. \right. \right. \\ & \quad \left. \left. \left. - \left\{ \frac{1}{\sqrt{\sigma_t^{*2}}} \left(\sum_{a=1}^K \left\{ \frac{\pi^e(a \mid X_t) \mathbb{1}[A_t = a] \{Y_t - f^*(a, X_t)\}}{\pi_t(a \mid X_t, \Omega_{t-1})} + \pi^e(a \mid x) f^*(a, X_t) \right\} - \theta_0 \right) \right\}^2 \mid X_t, \hat{f}_{t-1}, g_{t,N} \right] \right| \right] \\ & \leq \mathbb{E} \left[\left| \frac{1}{\sqrt{g_{t,N}}} - \frac{1}{\sqrt{\sigma_t^{*2}}} \right| \tilde{C}_1 + \mathbb{E} \left[\left| \hat{f}_{t-1}(a, X_t) - f^*(a, X_t) \right| \right] \tilde{C}_2 \right]. \end{aligned}$$

Then, from L^r convergence theorem, by using point convergence and the boundedness of $g_{t,N}$ and \hat{f}_{t-1} , we have $\mathbb{E} \left[\left| \mathbb{E} [z_t^2 \mid \Omega_{t-1}] - \nu^2 \right| \right] \rightarrow 0$ as $t \rightarrow \infty$ and $N \rightarrow \infty$. Therefore, as $T \rightarrow \infty$ and $N \rightarrow \infty$,

$$\mathbb{P} \left(\left| \frac{1}{T} \sum_{t=1}^T \mathbb{E} [z_t^2 \mid \Omega_{t-1}] - \nu^2 \right| \geq \varepsilon \right) \leq \frac{\frac{1}{T} \sum_{t=1}^T \mathbb{E} \left[\left| \mathbb{E} [z_t^2 \mid \Omega_{t-1}] - \nu^2 \right| \right]}{\varepsilon} \rightarrow 0.$$

As a conclusion,

$$\frac{1}{T} \sum_{t=1}^T z_t^2 - \nu^2 = \frac{1}{T} \sum_{t=1}^T (z_t^2 - \mathbb{E} [z_t^2 \mid \Omega_{t-1}] + \mathbb{E} [z_t^2 \mid \Omega_{t-1}] - \nu^2) \xrightarrow{P} 0.$$

Conclusion

From the results of Steps 1–3, we showed the theorem. \square

Remark 12 (Donsker Propeties). Readers might feel that functions need to satisfy some assumptions such as Donsker propeties (Chernozhukov et al., 2018).

D Details of Sample Splitting

As well as Theorem 2, we prove the consistency of $g_{t,(1-r)T}$ in the following theorem.

Corollary 1. *Suppose that, for all $x \in \mathcal{X}$ and $k \in \mathcal{A}$, $\hat{f}_{t-1}(a, x) - f^*(a, x) \xrightarrow{P} 0$, $\hat{e}_{t-1}(a, x) - e^*(a, x) \xrightarrow{P} 0$ and $\hat{\theta}_t \xrightarrow{P} \theta_0$ as $t \rightarrow \infty$. Then, for $\pi_t \in \Pi$, we have $g'_{t,(1-r)T} - \sigma_t^{*2} \xrightarrow{P} 0$ as $t \rightarrow \infty$ and $T \rightarrow \infty$, where*

$$\sigma_t^{*2} = \mathbb{E} \left[\sum_{a=1}^K \left\{ \frac{(\pi^e(a \mid X_t))^2 \nu^*(a, X_t)}{\pi_t(a \mid X_t, \Omega_{t-1})} + (\pi^e(a \mid X_t) f^*(a, X_t) - \theta_0)^2 \right\} \right].$$

Then, using $g_{t,(1-r)T}$, we define a FA3IPW estimator with sample splitting $\hat{\theta}_{rT}^{\text{FA3IPW-SS}}$ as $\left(\sum_{t=1}^{rT} \frac{1}{\sqrt{g_{t,(1-r)T}}} \right)^{-1} \sum_{t=1}^{rT} \frac{1}{\sqrt{g_{t,(1-r)T}}} \left\{ \frac{\pi^e(a \mid X_t) \mathbb{1}[A_t = a] \{Y_t - \hat{f}_{t-1}(a, X_t)\}}{\pi_t(a \mid X_t, \Omega_{t-1})} + \pi^e(a \mid x) \hat{f}_{t-1}(a, X_t) \right\}$. The asymptotic distribution can be obtained from Theorem 1 as follows.

Corollary 2 (Asymptotic Distribution of the Proposed Estimator). *Suppose that*

Table 3: Specification of datasets

Dataset	the number of samples	Dimension	the number of classes
satimage	4,435	35	6
pendigits	7,496	16	10
mnist	60,000	780	10
dna	2,000	180	3
letter	15,000	16	26
sensorless	58,509	48	11
connect-4	67,557	126	3
covtype	581,012	54	7

- (i) there exist C_2 and C_3 such that $|\hat{f}_t| \leq C_2$ and $0 < 1/\sqrt{g_{t,(1-r)T}} \leq C_3$;
- (ii) Pointwise convergence in probability of \hat{f}_t , i.e., $\forall x \in \mathcal{X}$, $\hat{f}_t(a, x) - f^*(a, x) \xrightarrow{P} 0$ as $t \rightarrow \infty$;
- (iii) $g_{t,(1-r)T} - \sigma_t^{*2} \xrightarrow{P} 0$ as $t \rightarrow \infty$ and $T \rightarrow \infty$.

Then, under Assumptions 2, $\left(\frac{1}{\sqrt{rT}} \sum_{t=1}^{rT} \frac{1}{\sqrt{g_{t,(1-r)T}}}\right) \left(\hat{\theta}_{rT}^{\text{FA3IPW}} - \theta_0\right) \xrightarrow{d} \mathcal{N}(0, 1)$ as $T \rightarrow \infty$.

Thus, we can obtain an estimator with asymptotic normality without assuming the access to \mathcal{E}_N .

E Details of Experiments

In this section, we show the details and additional results with different settings of Section 6. First, we show the description of the datasets in Table. All datasets are downloaded from <https://www.csie.ntu.edu.tw/~cjlin/libsvmtools/datasets/>. For these datasets, in the experiment shown in Section 6, we use a policy $\pi_t = 0.7\pi_t^a + 0.3\pi_t^u$ with 2,000 samples. We use the UCB and RW policy for the adaptive policy. The UCB algorithm follows the LinUCB proposed by Chu et al. (2011). In the random walk policy, we update the policy as follows:

- (i) At the first period, the policy π_1^a is given from the uniform distribution with a normalization for becoming $\sum_{k=1}^K \pi_1^a(k | X_1) = 1$.
- (ii) At a period t , we update the policy as $\pi_{t+1}^a(k | X_{t+1}) = \pi_t^a(k | X_t) + \eta_{k,t}$, where $\eta_{k,t}$ is a random variable drawn from the normal distribution.
- (iii) For satisfying the definition of the probability, we normalize $\pi_{t+1}^a(k | X_{t+1})$ to become $\sum_{k=1}^K \pi_{t+1}^a(k | X_{t+1}) = 1$.
- (iv) Iterate (ii) and (iii) until the trial ends.

For simplicity, we do not use context in the RW policy. For nonparametric regression, we only use the Nadaraya-Watson regression in Section 6. The evaluation policy is fixed at $\pi^e = 0.7\pi^d + 0.3\pi^u$. The policy π_d is a prediction of the logistic regression trained by all samples in \mathcal{S}_T . If the logistic regression predicts a class $a \in \mathcal{A}$ as the true label given X_t , the policy becomes $\pi_d(a, X_t) = 1$ and $\pi_d(a', X_t) = 0$ for $a' \neq a$.

In Table 4, we use a policy $\pi_t = 0.7\pi_t^a + 0.3\pi_t^u$ with $T = 1,000$ and $N = 1,000$. For nonparametric regression, we use the K-nearest neighbor regression.

In Table 5, we use a policy $\pi_t = 0.7\pi_t^a + 0.3\pi_t^u$ with $T = 500$ and $N = 1,000$. For nonparametric regression, we use the Nadaraya-Watson regression.

In Table 6, we use a policy $\pi_t = 0.7\pi_t^a + 0.3\pi_t^u$ with $T = 500$ and $N = 1,000$. For nonparametric regression, we use the K-nearest neighbor regression.

In Table 7, we use a policy $\pi_t = 0.5\pi_t^a + 0.5\pi_t^u$ with $T = 1,000$ and $N = 1,000$. For nonparametric regression, we use the Nadaraya-Watson regression.

In Table 8, we use a policy $\pi_t = 0.5\pi_t^a + 0.5\pi_t^u$ with $T = 1,000$ and $N = 1,000$. For nonparametric regression, we use the K-nearest neighbor regression.

Table 4: Experimental results under the RW policy (the upper table) and UCB policy (the lower table) under a policy $\pi_t = 0.7\pi_t^a + 0.3\pi_t^u$ with $T = 1,000$ and $N = 1,000$, and the K-nearest neighbor regression. The method with the lowest MSE is in bold.

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.024	0.155	0.049	0.217	0.078	0.280	0.017	0.131	0.187	0.422	0.071	0.261	0.022	0.132	0.044	0.208
SFA3IPW	0.023	0.148	0.058	0.239	0.075	0.274	0.021	0.144	0.170	0.408	0.067	0.259	0.023	0.140	0.050	0.221
DM	0.051	0.147	0.212	0.192	0.460	0.190	0.368	0.147	0.437	0.154	0.271	0.187	0.123	0.151	0.073	0.188
AdaIPW	0.042	0.203	0.075	0.267	0.099	0.315	0.042	0.204	0.189	0.423	0.117	0.341	0.073	0.205	0.058	0.242
A2IPW	0.024	0.155	0.048	0.216	0.078	0.279	0.018	0.135	0.173	0.408	0.074	0.270	0.025	0.133	0.047	0.216
FA2daIPW	0.042	0.204	0.074	0.265	0.099	0.314	0.039	0.196	0.203	0.434	0.109	0.328	0.060	0.205	0.055	0.234

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.061	0.246	0.100	0.316	0.130	0.359	0.036	0.183	0.223	0.471	0.077	0.266	4.627	1.969	0.049	0.222
SFA3IPW	0.062	0.246	0.126	0.344	0.156	0.393	0.032	0.169	0.164	0.405	0.092	0.293	4.150	1.918	0.052	0.226
DM	0.014	0.117	0.020	0.127	0.401	0.158	0.358	0.164	0.452	0.162	0.256	0.195	0.135	0.108	0.075	0.183
AdaIPW	0.089	0.296	0.129	0.350	0.197	0.438	0.080	0.272	0.312	0.556	0.106	0.309	12.278	2.869	0.086	0.292
A2IPW	0.067	0.259	0.098	0.313	0.128	0.355	0.037	0.184	0.193	0.439	0.076	0.266	4.100	1.882	0.042	0.206
FA2daIPW	0.084	0.287	0.125	0.348	0.207	0.452	0.079	0.271	0.381	0.614	0.101	0.300	10.234	2.581	0.090	0.296

Table 5: Experimental results under the RW policy (upper table) and UCB policy (lower table) under a policy $\pi_t = 0.7\pi_t^a + 0.3\pi_t^u$ with $T = 500$ and $N = 1,000$, and the Nadaraya-Watson regression. The method with the lowest MSE is in bold.

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.052	0.215	0.116	0.338	0.094	0.296	0.040	0.198	0.195	0.441	0.097	0.291	0.031	0.173	0.072	0.267
SFA3IPW	0.045	0.202	0.099	0.314	0.134	0.353	0.048	0.220	0.267	0.516	0.127	0.347	0.041	0.201	0.067	0.254
DM	0.096	0.148	0.466	0.207	0.404	0.231	0.245	0.197	0.469	0.162	0.384	0.178	0.117	0.206	0.127	0.217
AdaIPW	0.082	0.283	0.144	0.375	0.121	0.336	0.070	0.264	0.196	0.442	0.114	0.322	0.077	0.221	0.121	0.346
A2IPW	0.051	0.213	0.113	0.333	0.095	0.293	0.044	0.207	0.190	0.436	0.096	0.295	0.031	0.169	0.075	0.273
FA2daIPW	0.089	0.295	0.149	0.384	0.117	0.338	0.061	0.246	0.202	0.449	0.111	0.314	0.065	0.225	0.116	0.340

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.057	0.239	0.248	0.497	0.160	0.396	0.048	0.216	0.166	0.407	0.160	0.400	2.245	1.291	0.059	0.237
SFA3IPW	0.054	0.232	0.295	0.542	0.257	0.495	0.051	0.224	0.243	0.492	0.183	0.423	2.657	1.487	0.077	0.271
DM	0.036	0.139	0.303	0.128	0.336	0.204	0.238	0.200	0.465	0.144	0.391	0.198	0.122	0.191	0.111	0.216
AdaIPW	0.161	0.401	0.416	0.642	0.303	0.548	0.079	0.276	0.204	0.451	0.175	0.417	9.985	2.344	0.099	0.310
A2IPW	0.064	0.254	0.220	0.469	0.149	0.384	0.046	0.213	0.171	0.413	0.149	0.386	2.102	1.251	0.065	0.247
FA2daIPW	0.187	0.429	0.442	0.662	0.276	0.520	0.079	0.275	0.206	0.453	0.195	0.440	8.764	2.242	0.096	0.305

The proposed FA3IPW estimator performs well in many datasets. Especially when we use the RW policy, the performance of the estimator tends to be better than the other methods.

Table 6: Experimental results under the RW policy (upper table) and UCB policy (lower table) under a policy $\pi_t = 0.7\pi_t^a + 0.3\pi_t^u$ with $T = 500$ and $N = 1,000$, and the K-nearest neighbor regression. The method with the lowest MSE is in bold.

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.048	0.213	0.088	0.279	0.080	0.283	0.038	0.194	0.229	0.457	0.112	0.330	0.029	0.169	0.083	0.287
SFA3IPW	0.055	0.222	0.090	0.282	0.098	0.307	0.043	0.207	0.296	0.521	0.118	0.343	0.038	0.195	0.082	0.287
DM	0.068	0.178	0.292	0.235	0.545	0.199	0.371	0.167	0.435	0.154	0.305	0.256	0.131	0.170	0.088	0.239
AdaIPW	0.089	0.280	0.143	0.356	0.114	0.330	0.080	0.282	0.255	0.474	0.157	0.394	0.077	0.242	0.127	0.354
A2IPW	0.048	0.210	0.086	0.277	0.080	0.280	0.042	0.206	0.239	0.469	0.113	0.332	0.029	0.170	0.077	0.277
FA2daIPW	0.093	0.289	0.145	0.354	0.115	0.336	0.072	0.268	0.242	0.457	0.157	0.395	0.067	0.240	0.130	0.356

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.123	0.350	0.136	0.364	0.164	0.402	0.051	0.225	0.172	0.415	0.102	0.306	5.501	2.345	0.076	0.276
SFA3IPW	0.081	0.277	0.113	0.335	0.210	0.457	0.057	0.238	0.217	0.463	0.131	0.347	2.528	1.406	0.085	0.291
DM	0.026	0.153	0.088	0.144	0.509	0.151	0.376	0.165	0.435	0.185	0.277	0.255	0.131	0.163	0.090	0.216
AdaIPW	0.232	0.482	0.253	0.503	0.253	0.502	0.109	0.331	0.193	0.439	0.140	0.372	9.117	2.109	0.091	0.300
A2IPW	0.130	0.359	0.132	0.359	0.168	0.410	0.051	0.225	0.161	0.399	0.100	0.305	2.805	1.661	0.070	0.265
FA2daIPW	0.194	0.440	0.280	0.529	0.243	0.492	0.100	0.316	0.205	0.453	0.143	0.375	7.886	2.007	0.100	0.316

Table 7: Experimental results under the RW policy (upper table) and UCB policy (lower table) under a policy $\pi_t = 0.5\pi_t^a + 0.5\pi_t^u$ with $T = 1,000$ and $N = 1,000$, and the Nadaraya-Watson regression. The method with the lowest MSE is in bold.

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.043	0.207	0.064	0.253	0.056	0.237	0.023	0.151	0.105	0.296	0.062	0.237	0.032	0.150	0.048	0.219
SFA3IPW	0.040	0.199	0.066	0.257	0.055	0.231	0.028	0.166	0.103	0.319	0.076	0.270	0.031	0.152	0.052	0.228
DM	0.101	0.150	0.446	0.162	0.302	0.188	0.213	0.149	0.489	0.125	0.416	0.150	0.094	0.144	0.106	0.188
AdaIPW	0.075	0.273	0.108	0.327	0.067	0.255	0.047	0.217	0.102	0.303	0.072	0.259	0.160	0.237	0.070	0.261
A2IPW	0.043	0.208	0.065	0.255	0.057	0.238	0.023	0.153	0.098	0.290	0.059	0.234	0.037	0.155	0.050	0.224
FA2daIPW	0.077	0.276	0.106	0.320	0.068	0.260	0.046	0.214	0.109	0.309	0.074	0.261	0.120	0.231	0.070	0.260

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.041	0.203	0.096	0.310	0.155	0.384	0.028	0.167	0.203	0.447	0.130	0.353	3.280	1.811	0.057	0.236
SFA3IPW	0.042	0.204	0.092	0.303	0.112	0.334	0.028	0.166	0.214	0.462	0.102	0.306	5.639	2.371	0.070	0.263
DM	0.039	0.104	0.256	0.123	0.249	0.157	0.137	0.132	0.490	0.151	0.344	0.178	0.115	0.174	0.063	0.135
AdaIPW	0.078	0.274	0.134	0.365	0.213	0.459	0.048	0.220	0.204	0.446	0.127	0.342	4.397	1.517	0.086	0.292
A2IPW	0.041	0.201	0.113	0.335	0.149	0.378	0.028	0.167	0.193	0.429	0.112	0.321	2.062	1.427	0.051	0.223
FA2daIPW	0.083	0.284	0.138	0.371	0.225	0.470	0.057	0.239	0.217	0.465	0.151	0.382	3.792	1.355	0.096	0.307

Table 8: Experimental results under the RW policy (upper table) and UCB policy (lower table) under a policy $\pi_t = 0.5\pi_t^a + 0.5\pi_t^u$ with $T = 1,000$ and $N = 1,000$, and the K-nearest neighbor regression. The method with the lowest MSE is in bold.

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.038	0.195	0.060	0.246	0.058	0.237	0.025	0.158	0.097	0.306	0.055	0.234	0.031	0.147	0.044	0.206
SFA3IPW	0.040	0.199	0.056	0.237	0.071	0.261	0.029	0.172	0.098	0.313	0.080	0.281	0.031	0.156	0.045	0.210
DM	0.054	0.157	0.217	0.209	0.459	0.186	0.368	0.164	0.449	0.142	0.263	0.190	0.118	0.126	0.083	0.163
AdaIPW	0.053	0.230	0.098	0.313	0.091	0.301	0.054	0.232	0.107	0.326	0.071	0.267	0.137	0.185	0.074	0.251
A2IPW	0.038	0.196	0.060	0.244	0.056	0.233	0.025	0.159	0.098	0.310	0.056	0.236	0.036	0.146	0.046	0.211
FA2daIPW	0.052	0.228	0.100	0.316	0.092	0.303	0.053	0.229	0.105	0.320	0.075	0.273	0.099	0.183	0.068	0.236

Datasets Metrics	satimage		pendigits		mnist		dna		letter		sensorless		connect-4		covtype	
	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD	MSE	SD
FA3IPW	0.059	0.243	0.075	0.272	0.106	0.324	0.033	0.179	0.183	0.421	0.076	0.248	40.446	6.261	0.093	0.304
SFA3IPW	0.064	0.253	0.077	0.272	0.136	0.369	0.035	0.179	0.214	0.459	0.068	0.255	106.595	10.183	0.071	0.263
DM	0.016	0.114	0.029	0.117	0.417	0.160	0.282	0.120	0.437	0.169	0.180	0.193	0.142	0.142	0.036	0.163
AdaIPW	0.072	0.266	0.152	0.384	0.142	0.376	0.059	0.235	0.218	0.462	0.111	0.297	31.899	5.361	0.104	0.316
A2IPW	0.051	0.225	0.072	0.266	0.103	0.319	0.031	0.173	0.187	0.427	0.074	0.239	20.816	4.481	0.076	0.276
FA2daIPW	0.069	0.264	0.156	0.386	0.152	0.389	0.065	0.249	0.207	0.450	0.119	0.311	58.652	7.429	0.137	0.369