

Gravitational shock waves and scattering amplitudes

Andrea Cristofoli

Niels Bohr International Academy and Discovery Center

The Niels Bohr Institute, University of Copenhagen

Blegdamsvej 17, DK-2100 Copenhagen, Denmark

E-mail: a.cristofoli@nbi.ku.dk

ABSTRACT: We study gravitational shock waves using scattering amplitude techniques. After first reviewing the derivation in General Relativity as an ultrarelativistic boost of a Schwarzschild solution, we provide an alternative derivation by exploiting a novel relation between scattering amplitudes and solutions to Einstein field equations. We prove that gravitational shock waves arise from the classical part of a three point function with two massless scalars and a graviton. The region where radiation is localized has a distributional profile and it is now recovered in a natural way, thus bypassing the introduction of singular coordinate transformations as used in General Relativity. The computation is easily generalized to arbitrary dimensions and we show how the exactness of the classical solution follows from the absence of classical contributions at higher loops. A classical double copy between gravitational and electromagnetic shock waves is also provided and for a spinning source, using the exponential form of three point amplitudes, we infer a remarkable relation between gravitational shock waves and spinning ones, also known as gyratons. Using this property, we infer a family of exact solutions describing gravitational shock waves with spin. We then compute the phase shift of a particle in a background of shock waves finding agreement with an earlier computation by Amati, Ciafaloni and Veneziano for particles in the high energy limit. Applied to a gyraton, it provides a result for the scattering angle to all orders in spin.

KEYWORDS: Scattering Amplitudes, General Relativity

Contents

1	Introduction	1
2	The Aichelburg-Sexl metric	3
3	Gravitational shock waves from scattering amplitudes	4
4	A classical double copy for gravitational shock waves	11
5	Spinning gravitational shock waves	12
6	The scattering angle in the high energy limit	13
7	Conclusion	15
8	Acknowledgments	15

1 Introduction

The study of General Relativity using scattering amplitudes techniques is in a golden era thanks to state of the art computations for interacting black holes and the possibility to relate classical gravitational observables to scattering amplitudes [1–6]. Nowadays, the literature is vast and includes different approaches to deal with post-Newtonian and post-Minkowskian black holes [7–19], including also classical spin effects for Kerr black holes [20–36] and tidal effects [37, 38]. The existence of this literature might seem surprising, given that we are trading General Relativity for an even more complicated quantum gravitational system and its classical limit. However, the introduction of concepts such as unitarity and double-copy [39–61] has made possible not only the computation of observables relevant for LIGO/Virgo [62] but also to unravel new structures in classical field theory, proving that quantum mechanics can help us in elucidating the essence of classical physics. In fact, the EOB approach [63], which led to accurate models of gravitational wave signals for a binary system, was inspired by these ideas [64]. Along this line, this paper describes perturbative solutions in General Relativity using the scattering amplitude approach recently developed by Kosower, Maybee and O’Connell [4]. We focus on the Aichelburg-Sexl metric describing a gravitational shock

wave sourced by a massless particle [65]. Derived almost simultaneously by Aichelburg, Sexl, Penrose [66] and Bonnor [67], it has been central to our understanding of graviton dominance in high energy scattering [68], and in the past years it has been studied in different settings [69–73]. As we will see, an alternative derivation is also possible using a novel relation between perturbative solutions to Einstein’s field equations and scattering amplitudes. Several authors have conjectured a similar connection and in the case of a static massive source, it has led to the computation at second order in G_N of the Schwarzschild¹ and Kerr-Newman solution [75–78]. However, the lack of a covariant framework has made it impossible to treat more general cases such as those described by an energy momentum tensor sourced by massless particles with spin. This work is a step toward this direction. We start by first reviewing the original derivation by Aichelburg and Sexl of a gravitational shock wave employing an ultrarelativistic boost of a Schwarzschild solution. We then present a relation between classical solutions in General Relativity and the classical part of three point functions from quantum field theory. Using a massless particle coupled to a graviton, we derive the complete Aichelburg-Sexl metric as an exact solution to Einstein field equations. The region where the radiation is localized has a well known distributional profile and it is now recovered from the amplitude itself, bypassing the introduction of singular coordinate transformations as used in General Relativity. We also generalize the computation to arbitrary D dimensions finding agreement in the literature [79, 80] with the ultrarelativistic boost of the so called Tangherlini metric [81, 82]. We then extend the classical double copy for static black holes to gravitational shock waves, showing that their single copy is described by electromagnetic ones. For a spinning source, using the exponential form of three point amplitudes, we infer a remarkable relation between gravitational shock waves and spinning ones, also known as gyratons. From this, we obtain solutions describing spinning gravitational shock waves directly from the spinless case, avoiding the use of ultrarelativistic boosts on Kerr black holes [83]. To our knowledge, the existence of such a relation between exact solutions in General Relativity was previously unknown. Interestingly, this relation resembles the Newman-Janis algorithm [84] which provides a Kerr solution from a complex deformation of Schwarzschild, recently studied by Arkani-Hamed, Huang and O’Connell in [24]. We then compute the phase shift of a particle in a background of shock waves, finding agreement with earlier computations for particles in the high energy limit [85, 86]. Applied to a gyraton, it provides a result for the scattering angle valid to all orders in the spin.

We will work throughout in natural units and in mostly negative signature.

¹See also [74] for a derivation up to G_N^3 using EFT methods.

2 The Aichelburg-Sexl metric

Aichelburg and Sexl derived for the first time an exact solution to Einstein field equations describing the gravitational field generated by a massless particle [65]. Their procedure employed the use of an ultrarelativistic boost of a Schwarzschild solution, previously used by D'Eath to address the scattering of two ultrarelativistic black holes [87]. Let us review their original derivation. We start by introducing the Schwarzschild metric in isotropic coordinates [88]

$$ds^2 = \frac{(1-A)^2}{(1+A)^2} dt^2 - (1+A)^4 (dx^2 + dy^2 + dz^2) \quad , \quad A = \frac{mG_N}{2\sqrt{x^2 + y^2 + z^2}} \quad (2.1)$$

and we decompose it as

$$ds^2 = dt^2 - dx^2 - dy^2 - dz^2 + \left[\frac{(1-A)^2}{(1+A)^2} - 1 \right] dt^2 - \left[(1+A)^4 - 1 \right] (dx^2 + dy^2 + dz^2) \quad (2.2)$$

If we apply a Lorentz transformation to eq.(2.1) on the x -direction,

$$t = \frac{\bar{t} - v\bar{x}}{\sqrt{1-v^2}} \quad , \quad x = \frac{\bar{x} - v\bar{t}}{\sqrt{1-v^2}} \quad , \quad y = \bar{y} \quad , \quad z = \bar{z} \quad (2.3)$$

the previous line element changes to

$$ds^2 = d\bar{t}^2 - d\bar{x}^2 - d\bar{y}^2 - d\bar{z}^2 + \left[\frac{(1-A')^2}{(1+A')^2} - 1 \right] \frac{(d\bar{t} - v d\bar{x})^2}{1-v^2} - \left[(1+A')^4 - 1 \right] \left(\frac{(d\bar{x} - v d\bar{t})^2}{1-v^2} + d\bar{y}^2 + d\bar{z}^2 \right) \quad (2.4)$$

$$A' = \frac{mG_N \sqrt{1-v^2}}{2 \{ (\bar{x} - v\bar{t})^2 + (1-v^2)(\bar{y}^2 + \bar{z}^2) \}^{1/2}} \quad (2.5)$$

We write $m = p\sqrt{1-v^2}$ and expand (2.4-2.5) around $v = 1$ for a fixed value of p to find

$$ds^2 = d\bar{t}^2 - d\bar{x}^2 - d\bar{y}^2 - d\bar{z}^2 - \frac{4pG_N}{|\bar{t} - \bar{x}|} (d\bar{t} - d\bar{x})^2 \quad , \quad \bar{x} \neq \bar{t} \quad (2.6)$$

In order to include also the missing region given by $\bar{x} = \bar{t}$, Aichelburg and Sexl proposed a coordinate transformation which becomes singular in the limit for $v = 1$

$$\begin{aligned} x' - vt' &= \bar{x} - v\bar{t} \\ x' + vt' &= \bar{x} + v\bar{t} - 4pG_N \log \left[\sqrt{(\bar{x} - v\bar{t})^2 + (1-v^2)} - (\bar{x} - \bar{t}) \right] \end{aligned} \quad (2.7)$$

Using the following relation

$$\lim_{v \rightarrow 1} \left[\frac{1}{\sqrt{(x' - vt')^2 + (1 - v^2)\rho}} - \frac{1}{\sqrt{(x' - vt')^2 + (1 - v^2)}} \right] = -2\delta(t' - x') \log(\rho) \quad (2.8)$$

the line element assumes the usual form of an impulsive pp-wave

$$ds^2 = dt'^2 - dx'^2 - dy'^2 - dz'^2 + 4pG_N \delta(t' - x') \log(y'^2 + z'^2) (dt' - dx')^2 \quad (2.9)$$

The latter defines a global solution given by two copies of Minkowski space connected by a singularity along a light cone coordinate. Among the relevant properties of this solution we can notice that in going from eq.(2.1) to eq.(2.9) we have changed the algebraic type of the Weyl tensor from Petrov type D to the radiative type N [89], a property first discovered by Pirani [90]. Moreover, from the computation of the associated Einstein tensor we can infer that the energy momentum tensor is simply that of a massless particle, thus confirming the physical interpretation of the metric.

3 Gravitational shock waves from scattering amplitudes

The idea to perturbatively solve Einstein field equations using quantum field theory techniques dates back to a paper by Duff [76] where the Schwarzschild solution was derived up to G_N^2 order. After, several authors used known relations among off-shell scattering amplitudes and form factors so as to include quantum effects in the latter, confirming the same results for the classical part [78]. Both approaches require the knowledge of the Einstein-Hilbert action expanded around a fixed background which becomes intractable already after few iterations in the coupling $k = \sqrt{32\pi G_N}$. In order to have a better control on the complexity of the calculation, it would be desirable to relate scattering amplitudes directly to the metric tensor in the same way as these have been related to classical observables in [4]. To this end we start by considering a Riemannian manifold and a Minkowskian background. We then introduce an off-shell continuation of the second quantized solution to the linearized Einstein field equations

$$\hat{h}_{\mu\nu}(x) = \frac{k}{2} \sum_{\lambda} \int d\Phi_{off}(q) \left[\epsilon_{\mu\nu}^{\lambda}(q) \hat{a}_q^{\lambda} e^{-iq \cdot x} + (\epsilon_{\mu\nu}^{\lambda})^{\dagger}(q) (\hat{a}_q^{\lambda})^{\dagger} e^{iq \cdot x} \right] \quad (3.1)$$

To ensure the gauge dependence of the metric, the sum runs over longitudinal polarizations and the measure of integration used in [4] has been replaced with

$$d\Phi_{on}(q) = \frac{d^D q}{(2\pi)^D} 2\pi \delta(q^2) \theta(q_0) \quad \rightarrow \quad d\Phi_{off}(q) = \frac{d^D q}{(2\pi)^D} \frac{1}{q^2} \quad (3.2)$$

We have added the subscripts *on-off* to denote that every integral carrying such measure of integration will lead to an integrand with a momentum which is respectively on-shell or off-shell. In our case, eq.(3.2) ensures the off-shellness of the graviton and the fact we are not looking for radiative modes of the metric tensor. The $i\epsilon$ prescription is implicitly assumed. We then propose the following wave-function describing our system in absence of interactions

$$|\Psi_{\text{in}}\rangle = \int d\Phi_{\text{on}}(p)\phi(p) |p\rangle \otimes |0\rangle \quad , \quad d\Phi_{\text{on}}(p) = \frac{d^D p}{(2\pi)^D} \hat{\delta}^{(+)}(p^2 - m^2) \quad (3.3)$$

where $\hat{\delta}^{(+)}(p^2 - m^2) = 2\pi\delta(p^2 - m^2)\theta(p_0)$. The state $|0\rangle$ denotes the vacuum state of the gravitational field while $\phi(p)$ is a proper wave-packet describing the source. We now define the metric tensor satisfying the non linear Einstein field equations as

$$g_{\mu\nu}(x) = \eta_{\mu\nu} + h_{\mu\nu}(x) \quad , \quad h_{\mu\nu}(x) = \langle \Psi_I(t) | \hat{h}_{\mu\nu}^I(x) | \Psi_I(t) \rangle \quad (3.4)$$

The operator $\hat{h}_{\mu\nu}^I(x)$ is defined as the action of $U_{\text{int}}(+\infty, t)$ on (3.1), while the state $|\Psi_I(t)\rangle$ is defined as the evolution at time t of the initial state under $U_{\text{int}}(t, -\infty)$. We can now express (3.4) as follows

$$h_{\mu\nu}(x) = \langle \Psi_I(t) | U_{\text{int}}^\dagger(+\infty, t) \hat{h}_{\mu\nu}(x) U_{\text{int}}(+\infty, t) | \Psi_I(t) \rangle \quad (3.5)$$

$$= \langle \text{in} \Psi | U_{\text{int}}^\dagger(t, -\infty) U_{\text{int}}^\dagger(+\infty, t) \hat{h}_{\mu\nu}(x) U_{\text{int}}(+\infty, t) U_{\text{int}}(t, -\infty) | \Psi_{\text{in}} \rangle \quad (3.6)$$

$$= \langle \text{in} \Psi | S^\dagger \hat{h}_{\mu\nu}(x) S | \Psi_{\text{in}} \rangle \quad (3.7)$$

where we have introduced the S matrix of the system. In doing so, we have been able to relate the solution to the complete Einstein field equations with a plane wave operator, by encoding all non linearities in the S matrix alone. Using then $S = 1 + iT$ we can expand eq.(3.7) neglecting for the moment terms proportional to TT^\dagger

$$h_{\mu\nu}(x) = i \langle \Psi_{\text{in}} | \left(\hat{h}_{\mu\nu}(x) T - T^\dagger \hat{h}_{\mu\nu}(x) \right) | \Psi_{\text{in}} \rangle \quad (3.8)$$

From which

$$h_{\mu\nu}(x) = \frac{ik}{2} \sum_\lambda \int d\Phi_{\text{off}}(q) d\Phi_{\text{on}}(p) d\Phi_{\text{on}}(p') \phi(p) \phi^\dagger(p') \times \\ [\langle p' q^\lambda | T | p \rangle \epsilon_{\mu\nu}^\lambda(q) e^{-iq \cdot x} - \langle p' | T^\dagger | p q^\lambda \rangle (\epsilon_{\mu\nu}^\lambda)^\dagger(q) e^{iq \cdot x}] \quad (3.9)$$

$$= -k \sum_\lambda \int d\Phi_{\text{off}}(q) d\Phi_{\text{on}}(p) d\Phi_{\text{on}}(p') \text{Im} \left[\phi(p) \phi^\dagger(p') \langle p' q^\lambda | T | p \rangle \epsilon_{\mu\nu}^\lambda(q) e^{-iq \cdot x} \right] \quad (3.10)$$

Matrix elements in eq.(3.10) usually describe on-shell scattering amplitudes thanks to the covariant measures which have a Dirac delta in each integrated momentum. Having assumed instead an off-shell integration measure for gravitons, the term in eq.(3.10) won't be an on-shell scattering amplitude but an off-shell three point function given by

$$\langle p'q^\lambda | T | p \rangle = (2\pi)^D \delta^D(p - q - p') (\epsilon_{\alpha\beta}^\lambda)^\dagger(q) \mathcal{M}^{\alpha\beta}(p, p', q) \quad (3.11)$$

We now choose harmonic coordinates which amounts to requiring the following identity to hold

$$\sum_\lambda \epsilon_{\mu\nu}^\lambda(q) (\epsilon_{\alpha\beta}^\lambda)^\dagger(q) = \frac{1}{2} (\eta_{\mu\alpha} \eta_{\nu\beta} + \eta_{\mu\beta} \eta_{\nu\alpha} - \frac{2}{D-2} \eta_{\mu\nu} \eta_{\alpha\beta}) \equiv P_{\mu\nu\alpha\beta} \quad (3.12)$$

Using this, eq.(3.10) becomes

$$h_{\mu\nu}(x) = -k \int d\Phi_{off}(q) d\Phi_{on}(p) d\Phi_{on}(p') \text{Im} \left[\phi(p) \phi^\dagger(p') \times \right. \\ \left. (2\pi)^D \delta^D(p' + q - p) P_{\mu\nu\alpha\beta} \mathcal{M}^{\alpha\beta}(p, p', q) e^{-iq \cdot x} \right] \quad (3.13)$$

We now proceed by making explicit the integration measure for the source particle. Integrating over p' we obtain

$$h_{\mu\nu}(x) = -k \int d\Phi_{off}(q) \frac{d^D p}{(2\pi)^D} \frac{d^D p'}{(2\pi)^D} \hat{\delta}^{(+)}(p^2 - m^2) \hat{\delta}^{(+)}(p'^2 - m^2) \times \\ \text{Im} \left[\phi(p) \phi^\dagger(p') (2\pi)^D \delta^D(p - q - p') P_{\mu\nu\alpha\beta} \mathcal{M}^{\alpha\beta}(p, p', q) e^{-iq \cdot x} \right] \quad (3.14)$$

$$h_{\mu\nu}(x) = -k \int d\Phi_{off}(q) \frac{d^D p}{(2\pi)^D} \hat{\delta}^{(+)}(p^2 - m^2) \hat{\delta}^{(+)}((p - q)^2 - m^2) \times \\ \text{Im} \left[\phi(p) \phi^\dagger(p - q) P_{\mu\nu\alpha\beta} \mathcal{M}^{\alpha\beta}(p, p', q) e^{-iq \cdot x} \right] \quad (3.15)$$

where it is implicitly assumed that $\mathcal{M}^{\alpha\beta}(p, p', q)$ is constrained with $p' = p - q$. Making also explicit the off-shell integration measure we obtain,

$$h_{\mu\nu}(x) = -k \int \frac{d^D q}{(2\pi)^D} \frac{1}{q^2} \int \frac{d^D p}{(2\pi)^D} \hat{\delta}^{(+)}(p^2 - m^2) \hat{\delta}^{(+)}(q^2 - 2q \cdot p) \times \\ \text{Im} \left[\phi(p) \phi^\dagger(p - q) P_{\mu\nu\alpha\beta} \mathcal{M}^{\alpha\beta}(p, p', q) e^{-iq \cdot x} \right] \quad (3.16)$$

For a wave-packet sharply peaked around a given momentum p_0 ² we obtain

$$h_{\mu\nu}(x) = -k \int \frac{d^D q}{(2\pi)^D} \frac{\hat{\delta}^{(+)}(q^2 - 2q \cdot p_0)}{q^2} \text{Im}[P_{\mu\nu\alpha\beta} \mathcal{M}^{\alpha\beta}(p_0, p' = p_0 - q, q) e^{-iq \cdot x}] \quad (3.17)$$

We are thus left with a remarkable relation between the classical metric tensor satisfying Einstein field equation and three point functions with an external graviton, valid both for massive and massless sources

$$h_{\mu\nu}(x) = -k \int \frac{d^D q}{(2\pi)^D} \frac{\hat{\delta}^{(+)}(q^2 - 2q \cdot p_0)}{q^2} \text{Im}[P_{\mu\nu\alpha\beta} \mathcal{M}^{\alpha\beta}(p_0, p' = p_0 - q, q) e^{-iq \cdot x}] \quad (3.18)$$

Let us consider the massless case. Taking advantage of this covariant relation, we can proceed to explore which space-time corresponds to a three point function with an off-shell graviton and a massless source. Based on what has been discussed in Section 2, this should correspond to a gravitational shock wave. We start at tree level from the interaction of a graviton with a massless source

$$\mathcal{M}^{\mu\nu}(p_1, p_2) = \frac{ik}{2} (p_1^\mu p_2^\nu + p_2^\mu p_1^\nu - \eta^{\mu\nu} p_1 \cdot p_2) \quad (3.19)$$

$$P_{\mu\nu\alpha\beta} \mathcal{M}^{\alpha\beta}(p, q) = \frac{ik}{2} \left[2p_\mu p_\nu - p_\mu q_\nu - p_\nu q_\mu + \eta_{\mu\nu} p \cdot q \right] \quad (3.20)$$

where in the last equation we have expressed the whole contributions in terms of p^μ and q^μ , being the former the incoming momenta. The whole metric tensor depends on only two functions

$$h_{\mu\nu}(x) = -\frac{k^2}{2} \left[2p_\mu p_\nu \Theta(x) - p_\mu \Theta_\nu(x) - p_\nu \Theta_\mu(x) + \eta_{\mu\nu} p^\alpha \Theta_\alpha(x) \right] \quad (3.21)$$

$$\Theta(x) = \int \frac{d^D q}{(2\pi)^D} \hat{\delta}^{(+)}(q^2 - 2q \cdot p) \frac{\cos(q \cdot x)}{q^2} \quad (3.22)$$

$$\Theta_\mu(x) = \int \frac{d^D q}{(2\pi)^D} \hat{\delta}^{(+)}(q^2 - 2q \cdot p) \frac{q_\mu}{q^2} \cos(q \cdot x) \quad (3.23)$$

In the classical limit we implement the limit for small q by considering the integration domain where $p \gg q$. This amounts to disregarding the Heaviside theta in eq.(3.18) as well as the q^2 term in its Dirac delta

²For further details, see [4], Section 4.

$$\hat{\delta}^{(+)}(q^2 - 2q \cdot p) \rightarrow 2\pi\delta(2q \cdot p) \quad (3.24)$$

In this limit $\Theta_\mu(x)$ is vanishing being the integrand an odd and real valued function and we are thus left with the computation of $\Theta(x)$. Using then the integral representation for a Dirac delta together with the Schwinger parametrization we obtain

$$\Theta(x) = -i \int_{\mathbb{R}} ds \int_{\mathbb{R}_+} dt \int \frac{d^D q}{(2\pi)^D} e^{-iq \cdot (x-2ps) + iq^2 t} \quad (3.25)$$

The latter is a complex Gaussian integral and its computation gives

$$\Theta(x) = -\left(\frac{i}{4\pi}\right)^{\frac{D}{2}} \int_{\mathbb{R}} ds \int_{\mathbb{R}_+} dt \frac{e^{-i\frac{(x-2ps)^2}{4t}}}{t^{\frac{D}{2}}} \quad (3.26)$$

Expanding the square in the exponential,

$$\Theta(x) = -\left(\frac{i}{4\pi}\right)^{\frac{D}{2}} \int_{\mathbb{R}} ds \int_{\mathbb{R}_+} dt \frac{e^{-i\frac{(x-2ps)^2}{4t}}}{t^{\frac{D}{2}}} \quad (3.27)$$

$$= -2\pi \left(\frac{i}{4\pi}\right)^{\frac{D}{2}} \int_{\mathbb{R}_+} \frac{dt}{t^{\frac{D}{2}}} e^{-\frac{ix^2}{4t}} \delta\left(\frac{p \cdot x}{t}\right) \quad (3.28)$$

$$= -2\pi \left(\frac{i}{4\pi}\right)^{\frac{D}{2}} \delta(p \cdot x) \int_{\mathbb{R}_+} \frac{dt}{t^{\frac{D}{2}}} e^{-\frac{ix^2}{4t}} |t| \quad (3.29)$$

Changing variables to $t = \frac{1}{u}$, we get

$$\Theta(x) = -2\pi \left(\frac{i}{4\pi}\right)^{\frac{D}{2}} \delta(p \cdot x) \int_{\mathbb{R}_+} du \frac{e^{-\frac{ix^2}{4}}}{u^{\frac{6-D}{2}}} \quad (3.30)$$

At this point, we should carefully distinguish the computation in $D = 4$ from other dimensions. One can realize it by computing separately the two cases and by a comparison afterwards. We start from the case with $D = 4$,

$$\Theta(x) = \frac{1}{8\pi} \delta(p \cdot x) \int_{\mathbb{R}_+} \frac{du}{u} e^{-\frac{ix^2}{4}} \quad (3.31)$$

In order to compute this integral, we consider its partie finie (Pf) to find

$$\begin{aligned}
\Theta(x) &= \frac{1}{8\pi} \delta(p \cdot x) \text{Pf} \lim_{z \rightarrow 0} \int_z^{+\infty} \frac{du}{u} e^{-\frac{iux^2}{4}} \\
&= \frac{1}{8\pi} \delta(p \cdot x) \text{Pf} \lim_{z \rightarrow 0} \int_1^{+\infty} \frac{du}{u} e^{-\frac{izux^2}{4}} \\
&= \frac{1}{8\pi} \delta(p \cdot x) \text{Pf} \lim_{z \rightarrow 0} E_1 \left(\frac{zix^2}{4} \right)
\end{aligned} \tag{3.32}$$

where we have introduced the exponential integral $E_1(x)$. Using the Puiseux series

$$E_1(z) = -\gamma - \log z - \sum_{k=1}^{\infty} \frac{(-z)^k}{kk!} \quad , \quad |\arg(z) < \pi| \tag{3.33}$$

The result is

$$\Theta(x) = -\frac{1}{8\pi} \delta(p \cdot x) \log(|x^2|) \quad , \quad D = 4 \tag{3.34}$$

As for the case $D \neq 4$, we evaluate it by first rescaling eq.(3.30),

$$\Theta(x) = -2\pi \left(\frac{i}{4\pi} \right)^{\frac{D}{2}} \delta(p \cdot x) \left(\frac{x^2}{4} \right)^{\frac{4-D}{2}} \int_{\mathbb{R}_+} du e^{-iu} u^{\frac{D-6}{2}} \tag{3.35}$$

After a Wick rotation, the remaining integral defines a Gamma function, from which

$$\Theta(x) = \frac{\pi^{\frac{2-D}{2}} \Gamma(\frac{D-2}{2})}{4} \frac{\delta(p \cdot x)}{(x^2)^{\frac{D-4}{2}}} \quad , \quad D > 4 \tag{3.36}$$

We can thus summarize our results,³

$$\Theta(x) = \begin{cases} -\frac{1}{8\pi} \delta(p \cdot x) \log(|x^2|) & , \quad D = 4 \\ \frac{\pi^{\frac{2-D}{2}} \Gamma(\frac{D-2}{2})}{4} \frac{\delta(p \cdot x)}{(x^2)^{\frac{D-4}{2}}} & , \quad \text{otherwise} \end{cases} \tag{3.37}$$

Using eq.(3.21) we can read the metric tensor related to a three point function with a massless source and an off-shell graviton.

³One could also infer the $D = 4$ case from the following regularization $\frac{\Gamma(\frac{D-4}{2})}{x^{D-4}} \rightarrow -2 \log(x)$. This amounts to remove a divergent quantity from the metric tensor with a gauge transformation.

The final result is

$$h_{\mu\nu}(x) = \begin{cases} 4G_N p_\mu p_\nu \delta(p \cdot x) \log(|x^2|) & , \quad D = 4 \\ -8\pi^{\frac{4-D}{2}} G_N p_\mu p_\nu \frac{\Gamma(\frac{D-2}{2}) \delta(p \cdot x)}{(D-4)(x^2)^{\frac{D-4}{2}}} & , \quad \text{otherwise} \end{cases} \quad (3.38)$$

We will shortly argue that contributions from higher loops produce only divergences which are removed from the cut terms proportional to TT^\dagger in (3.7). This procedure provides an exact solution to Einstein field equations already at linear order in G_N . In $D = 4$ the line element reads

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = dt^2 - dx^2 - dy^2 - dz^2 + 4G_N \delta(p \cdot x) \log(|x^2|) p_\mu p_\nu dx^\mu dx^\nu \quad (3.39)$$

For a massless particle moving along the x direction we recover the Aichelburg-Sexl metric (2.9) for a gravitational shock wave

$$ds^2 = dt^2 - dx^2 - dy^2 - dz^2 + 4p G_N \delta(t - x) \log(y^2 + z^2) (dt - dx)^2 \quad (3.40)$$

In D dimensions, the metric is in agreement with earlier computations describing the ultrarelativistic boost of the Schwarzschild-Tangherlini metric in D dimensions [80]. As for the coordinates associated with this metric, we notice that eq.(3.38) satisfies the harmonic gauge condition, equivalent at linear order in G_N with the linear harmonic,

$$\eta^{\alpha\beta} \Gamma_{\alpha\beta}^\mu = 0 \quad \rightarrow \quad \partial_\alpha h^{\mu\alpha} = \frac{1}{2} \partial^\mu h \quad (3.41)$$

which can be easily seen to be satisfied thanks to eq.(3.38) being traceless. In $D = 4$

$$\partial_\alpha h^{\mu\alpha} = 4G_N p^\mu p^\alpha \partial_\alpha \delta(p \cdot x) \log(|x^2|) + 8G_N p^\mu p^\alpha \delta(p \cdot x) \frac{x_\mu}{x^2} = 0 \quad (3.42)$$

with the same result in higher dimensions. This is consistent with the harmonic gauge choice made in eq.(3.12). The advantages of this computation with respect to the derivation from classical General Relativity are several. The exactness of the solution already at linear in G_N can now be explained in light of the absence of classical contributions to higher loops in three point functions with two massless scalars and a graviton [2]. This is in contrast with the computation for a massive three point function where the classical part from higher loop orders is non vanishing and needed in order to reproduce the expansion of Schwarzschild in G_N [78]. Remarkably, the distributional profile emerges in a natural way from the amplitude itself, with no need to introduce singular coordinate transformations as those in eq.(2.7). As we will see, this property is more general: it is valid also for gravitational shock waves carrying a spin dependence.

4 A classical double copy for gravitational shock waves

In the previous section we have shown a relation between perturbative solutions to Einstein field equations and scattering amplitudes. The latter can be introduced also for a gauge theory as classical electromagnetism. We start by introducing the following operator for a gauge potential

$$\hat{A}^\mu(x) = \sum_\lambda \int d\Phi_{off}(q) \left[\epsilon_\mu^\lambda(q) \hat{a}_q^\lambda e^{-iq \cdot x} + (\epsilon_\mu^\lambda)^\dagger(q) (\hat{a}_q^\lambda)^\dagger e^{iq \cdot x} \right] \quad (4.1)$$

Following the same steps seen before and working in Feynman gauge we can easily derive a relation between a gauge potential $A^\mu(x)$ and three point functions in scalar QED with an external photon,

$$A^\mu(x) = \int \frac{d^D q}{(2\pi)^D} \frac{\hat{\delta}^{(+)}(q^2 - 2q \cdot p_0)}{q^2} \text{Im}[\mathcal{M}^\mu(p_0, p' = p_0 - q, q) e^{-iq \cdot x}] \quad (4.2)$$

We now consider electromagnetic shock waves [91]. We find natural to relate these to a three point amplitude of a massless scalar particle coupled to a photon,

$$\mathcal{M}^\mu = -ie(2p^\mu - q^\mu) \quad (4.3)$$

Using this, we can express the gauge potential $A^\mu(x)$ in terms of (3.22, 3.23),

$$A^\mu(x) = -e \left[2p^\mu \Theta(x) - \Theta^\mu(x) \right] \quad (4.4)$$

The final result for an electromagnetic shock wave is

$$A^\mu(x) = \begin{cases} \frac{e}{4\pi} p^\mu \delta(p \cdot x) \log(|x^2|) & , \quad D = 4 \\ -\frac{e}{2\pi} p^\mu \pi^{\frac{4-D}{2}} \frac{\Gamma\left(\frac{D-2}{2}\right) \delta(p \cdot x)}{(D-4)(x^2)^{\frac{D-4}{2}}} & , \quad \text{otherwise} \end{cases} \quad (4.5)$$

We may now consider the classical double copy procedure shown in [42] in order to construct a solution in General Relativity. This amounts to the following replacement

$$e \rightarrow 16\pi G_N \quad , \quad p^\mu \rightarrow p^\mu p^\nu \quad (4.6)$$

Remarkably, this gives the correct gravitational shock wave in arbitrary D dimensions of eq.(3.38) showing that gravitational and electromagnetic shock waves are related by a classical double copy. This fact was shown to hold for static black holes [43, 45] and then generalized to accelerating ones in [42]. Here, we have proven that a classical double copy is satisfied also by shock waves in gravity and gauge theories.

5 Spinning gravitational shock waves

Having studied in depth the relation between massless particles and gravitational shock waves, we find natural to investigate the same relation for the case of a spinning source. As we will see, this leads to a family of classical solutions also known in the literature as gyratons [92, 93]. For ease of discussion we restrict ourselves to the case $D = 4$. In order to perform the computation, we take advantage of the exponential representation of three point functions for a spinning massive particle emitting a graviton [22, 28]

$$\mathcal{M}_{\mu\nu}^S = \mathcal{M}_{\mu\nu} e^{a \cdot q} \quad , \quad a^\mu = \frac{s^\mu}{m} \quad (5.1)$$

being $\mathcal{M}_{\mu\nu}$ the associated spinless three point amplitude and s^μ the spin vector of the source. In the massless limit we treat a^μ as a constant and express the spin vector using the representation of the Pauli-Lubanski pseudovector for a massless particle [94]

$$s^\mu = \frac{1}{E} W^\mu \quad , \quad W^\mu = \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} S^{\nu\alpha} p^\beta \quad (5.2)$$

Using this, the metric tensor reads

$$h_{\mu\nu}^S(x) = -\frac{k^2}{2} \left(2p_\mu p_\nu \Theta^S(x) - p_\mu \Theta_\nu^S(x) - p_\nu \Theta_\mu^S(x) + \eta_{\mu\nu} \Theta_\alpha^S(x) p^\alpha \right) \quad (5.3)$$

$$\Theta^S(x) = \int \frac{d^4 q}{(2\pi)^4} \hat{\delta}^{(+)}(q^2 - 2q \cdot p) \frac{\cos(q \cdot x) e^{a \cdot q}}{q^2} \quad (5.4)$$

$$\Theta_\mu^S(x) = \int \frac{d^4 q}{(2\pi)^4} \hat{\delta}^{(+)}(q^2 - 2q \cdot p) \frac{q_\mu}{q^2} \cos(q \cdot x) e^{a \cdot q} \quad (5.5)$$

We can now prove a relation between spinless gravitational shock waves and gyratons. We restrict to the classical limit by considering the integration region where $p \gg q$, thus

$$\Theta^S(x) = \frac{\Theta(x - ia) + \Theta(x + ia)}{2} \quad , \quad \Theta_\mu^S(x) = \frac{\Theta_\mu(x - ia) + \Theta_\mu(x + ia)}{2} \quad (5.6)$$

We now consider the behavior of $\Theta(x)$ under a complex shift. Introducing $a = \sqrt{-a^\mu a_\mu}$ we obtain the following expression

$$\Theta(x - ia) = -\frac{1}{8\pi} \delta(p \cdot x) \log(|x^2 + a^2|) \quad (5.7)$$

Eq.(5.7) is real valued due to the absence of linear terms in $a \cdot x$. This is ensured by the Dirac delta in $p \cdot x$ and the fact that the Pauli-Lubanski pseudovector and a^μ are proportional to p^μ [94]. Thus,

$$\begin{cases} \Theta^S(x) = \Theta(x - ia) \\ \Theta_\mu^S(x) = \Theta_\mu(x - ia) \end{cases} \quad \rightarrow \quad h_{\mu\nu}^S(x) = h_{\mu\nu}(x - ia) \quad (5.8)$$

Remarkably, thanks to the exponential form of the three point amplitude we can now read the metric tensor sourced by a massless spinning source directly from the spinless case using the shift $x^\mu \rightarrow x^\mu - ia^\mu$. To our knowledge, this property between spinless shock waves and gyratons was previously unknown and it relates to the exponential form of the energy momentum tensor for linearized Kerr black holes [20], which is preserved after an ultrarelativistic boost. The line element is

$$ds^2 = dt^2 - dx^2 - dy^2 - dz^2 + 4G_N \delta(p \cdot x) \log(|(x^2 + a^2)|) p_\mu p_\nu dx^\mu dx^\nu \quad (5.9)$$

In particular, for a spinning particle moving along the x direction

$$ds^2 = dt^2 - dx^2 - dy^2 - dz^2 + 4G_N p \delta(x - t) \log(|y^2 + z^2 - a^2|) (dt - dx)^2 \quad (5.10)$$

in agreement with an earlier computation by Ferrari and Pendenza [83] describing the ultrarelativistic boost of a Kerr black hole. The derivation by a simple shift in a^μ is remarkable, since the same in classical General Relativity is much more complicated. Interestingly, this procedure resembles the Newman-Janis algorithm [84] which provides a Kerr solution from a complex deformation of Schwarzschild, this last recently studied by Arkani-Hamed, Huang and O'Connell in [24]. As for the singularity at $y^2 + z^2 = a^2$, we interpret this as the remnant of the singularity in the equatorial plane.

6 The scattering angle in the high energy limit

The computation of geodesics in a gravitational shock wave background has been explored by several authors [95, 96]. Since the whole space-time is Minkowskian up to a region defined by a null light cone coordinate, geodesics are fully determined from the net change in momentum of a particle

$$\Delta p_0^\mu = \frac{1}{2} \int_{\mathbb{R}} d\sigma \partial^\mu h_{\alpha\beta}(x(\sigma)) p_0^\alpha p_0^\beta \quad (6.1)$$

where the subscript 0 denotes the particle, while σ the affine parameter of its world-line. To leading order in G_N we assume free motion

$$x_0^\mu(\sigma) = p_0^\mu \sigma + b^\mu \quad , \quad b \cdot p_0 = 0 \quad , \quad b \cdot p = 0 \quad (6.2)$$

being b^μ a covariant impact parameter and p^μ the momentum associated with the shock wave. The resulting change of impulse in $D = 4$ reads

$$\Delta p_0^\mu = \frac{1}{2} p_0^\alpha p_0^\beta \int_{\mathbb{R}} d\sigma 8G_N p_\alpha p_\beta \left[\delta(p \cdot x_0(\sigma)) \frac{x_0^\mu(\sigma)}{x_0^2(\sigma)} \right] = \frac{4G_N p \cdot p_0}{b \cdot b} b^\mu \quad (6.3)$$

Having computed the change of momentum experienced by the particle, we can compute the associated phase shift using

$$\sin(\theta) = \frac{\Delta p_0^\mu b_\mu}{p_0 b} \quad (6.4)$$

where we have introduced $b = \sqrt{-b_\mu b^\mu}$. The result is

$$\sin(\theta) = \frac{4G_N p \cdot p_0}{p_0 b} \quad (6.5)$$

Let's now consider the massless limit,

$$p \cdot p_0 = p p_0 - \vec{p} \cdot \vec{p}_0 = 2p_0^2 \quad , \quad s = 4p_0^2 \quad (6.6)$$

If we now apply the small angle approximation we obtain

$$\theta = \frac{4G_N \sqrt{s}}{b} \quad (6.7)$$

in agreement with an earlier computation by Dray and t'Hooft [69]. Interestingly, as shown by Amati, Ciafaloni and Veneziano, the same result agrees with the leading order scattering angle between particles in the high energy limit [85]. We can generalize this result including effects to all orders in spin using as a source the metric tensor for a gyraton derived in eq.(5.8). This provides the following result

$$\Delta p^\mu = \frac{4G_N p \cdot p_0 b^\mu}{a^2 - b^2} \quad (6.8)$$

The scattering angle in the massless limit and including effects to all order in spin reads

$$\theta = \frac{2G_N \sqrt{s}}{b - a} + \frac{2G_N \sqrt{s}}{b + a} \quad (6.9)$$

Interestingly, this scattering angle in the high energy limit resembles a striking similarity with the all order in spin result by Vines [20] including the pole at $b = a$.

7 Conclusion

We have derived a relation between perturbative solutions to Einstein field equations and off-shell scattering amplitudes thanks to a covariant framework developed by Kosower, Maybee and O’Connell [4]. We have studied to which gravitational field corresponds a scattering amplitude with an off-shell graviton and two massless particles finding that the latter describes a gravitational shock wave also known as Aichelburg-Sexl metric [65]. The result has been easily generalized to arbitrary D dimensions finding agreement with previous computation of D dimensional shock waves in General Relativity [80]. The advantage of this computation are several. We have been able to avoid singular coordinate transformations which were used in General Relativity to deal with the singular behavior of the gravitational field along a light cone coordinate. Remarkably, the distributional profile emerges in a natural way from the amplitude itself, while the exactness of the classical solution at linear in G_N can now be explained in light of the absence of classical contributions at higher loops for three point functions with massless particles. We have also shown that a classical double copy is satisfied between gravitational and electromagnetic shock waves and for a spinning source, using the exponential form of three point amplitudes, we have inferred a remarkable relation between gravitational shock waves and spinning ones, also known as gyratons. Using this property, we have been able to infer solutions describing spinning gravitational shock waves directly from the spinless case, thus bypassing the derivation in General Relativity involving an ultrarelativistic boost of a Kerr black hole. We have computed the phase shift of a particle in a background of shock waves finding agreement with earlier computations for the scattering angle of particles in the high energy limit [85, 86]. Applied to a gyraton, it has provided a result to all orders in the spin.

8 Acknowledgments

I am sincerely thankful to Poul Damgaard, for his constant support and several useful conversations. I also thank Donal O’Connell, Paolo Di Vecchia and Marcello Ortaggio for useful conversations and comments on the manuscript. This work has been based partly on funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 764850 (“SAGEX”).

References

- [1] T. Damour, “High-energy gravitational scattering and the general relativistic two-body problem,” *Phys. Rev. D* **97**, no. 4, 044038 (2018) doi:10.1103/PhysRevD.97.044038 [arXiv:1710.10599 [gr-qc]].
- [2] N. J. Bjerrum-Bohr, P. H. Damgaard, G. Festuccia, L. Planté and P. Vanhove, “General Relativity from Scattering Amplitudes,” *Phys. Rev. Lett.* **121** (2018) no.17, 171601 doi:10.1103/PhysRevLett.121.171601 [arXiv:1806.04920 [hep-th]].
- [3] C. Cheung, I. Z. Rothstein and M. P. Solon, “From Scattering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion,” *Phys. Rev. Lett.* **121**, no. 25, 251101 (2018) doi:10.1103/PhysRevLett.121.251101 [arXiv:1808.02489 [hep-th]].
- [4] D. A. Kosower, B. Maybee and D. O’Connell, “Amplitudes, Observables, and Classical Scattering,” *JHEP* **02** (2019), 137 doi:10.1007/JHEP02(2019)137 [arXiv:1811.10950 [hep-th]].
- [5] Z. Bern, C. Cheung, R. Roiban, C. H. Shen, M. P. Solon and M. Zeng, “Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order,” *Phys. Rev. Lett.* **122** (2019) no.20, 201603 doi:10.1103/PhysRevLett.122.201603 [arXiv:1901.04424 [hep-th]].
- [6] Z. Bern, C. Cheung, R. Roiban, C. H. Shen, M. P. Solon and M. Zeng, “Black Hole Binary Dynamics from the Double Copy and Effective Theory,” *JHEP* **1910** (2019) 206 doi:10.1007/JHEP10(2019)206 [arXiv:1908.01493 [hep-th]].
- [7] T. Damour, “Classical and Quantum Scattering in Post-Minkowskian Gravity,” arXiv:1912.02139 [gr-qc].
- [8] A. Cristofoli, N. E. J. Bjerrum-Bohr, P. H. Damgaard and P. Vanhove, “Post-Minkowskian Hamiltonians in general relativity,” *Phys. Rev. D* **100** (2019) no.8, 084040 doi:10.1103/PhysRevD.100.084040 [arXiv:1906.01579 [hep-th]].
- [9] A. Cristofoli, P. H. Damgaard, P. Di Vecchia and C. Heissenberg, “Second-order Post-Minkowskian scattering in arbitrary dimensions,” [arXiv:2003.10274 [hep-th]].
- [10] A. Koemans Collado, P. Di Vecchia and R. Russo, “Revisiting the 2PM eikonal and the dynamics of binary black holes,” *Phys. Rev. D* **100** (2019) no.6, 066028 doi:10.1103/PhysRevD.100.066028 [arXiv:1904.02667 [hep-th]].
- [11] Y. F. Bautista and A. Guevara, “From Scattering Amplitudes to Classical Physics: Universality, Double Copy and Soft Theorems,” [arXiv:1903.12419 [hep-th]].
- [12] C. Cheung and M. P. Solon, “Classical Gravitational Scattering at $\mathcal{O}(G^3)$ from Feynman Diagrams,” [arXiv:2003.08351 [hep-th]].

- [13] Z. Bern, H. Ita, J. Parra-Martinez and M. S. Ruf, “Universality in the classical limit of massless gravitational scattering,” [arXiv:2002.02459 [hep-th]].
- [14] N. Bjerrum-Bohr, A. Cristofoli, P. H. Damgaard and H. Gomez, “Scalar-Graviton Amplitudes,” JHEP **11** (2019), 148 doi:10.1007/JHEP11(2019)148 [arXiv:1908.09755 [hep-th]].
- [15] N. E. J. Bjerrum-Bohr, A. Cristofoli and P. H. Damgaard, “Post-Minkowskian Scattering Angle in Einstein Gravity,” arXiv:1910.09366 [hep-th].
- [16] G. Kälin and R. A. Porto, “From Boundary Data to Bound States,” JHEP **01** (2020), 072 doi:10.1007/JHEP01(2020)072 [arXiv:1910.03008 [hep-th]].
- [17] G. Kälin and R. A. Porto, “Post-Minkowskian Effective Field Theory for Conservative Binary Dynamics,” [arXiv:2006.01184 [hep-th]].
- [18] J. Blümlein, A. Maier, P. Marquard, G. Schäfer and C. Schneider, “From Momentum Expansions to Post-Minkowskian Hamiltonians by Computer Algebra Algorithms,” Phys. Lett. B **801** (2020), 135157 doi:10.1016/j.physletb.2019.135157 [arXiv:1911.04411 [gr-qc]].
- [19] J. Blümlein, A. Maier, P. Marquard and G. Schäfer, “Testing binary dynamics in gravity at the sixth post-Newtonian level,” [arXiv:2003.07145 [gr-qc]].
- [20] J. Vines, “Scattering of two spinning black holes in post-Minkowskian gravity, to all orders in spin, and effective-one-body mappings,” Class. Quant. Grav. **35** (2018) no.8, 084002 doi:10.1088/1361-6382/aaa3a8 [arXiv:1709.06016 [gr-qc]].
- [21] J. Vines, J. Steinhoff and A. Buonanno, “Spinning-black-hole scattering and the test-black-hole limit at second post-Minkowskian order,” Phys. Rev. D **99** (2019) no.6, 064054 doi:10.1103/PhysRevD.99.064054 [arXiv:1812.00956 [gr-qc]].
- [22] A. Guevara, A. Ochirov and J. Vines, “Scattering of Spinning Black Holes from Exponentiated Soft Factors,” JHEP **1909** (2019) 056 doi:10.1007/JHEP09(2019)056 [arXiv:1812.06895 [hep-th]].
- [23] A. Guevara, A. Ochirov and J. Vines, “Black-hole scattering with general spin directions from minimal-coupling amplitudes,” Phys. Rev. D **100** (2019) no.10, 104024 doi:10.1103/PhysRevD.100.104024 [arXiv:1906.10071 [hep-th]].
- [24] N. Arkani-Hamed, Y. t. Huang and D. O’Connell, “Kerr black holes as elementary particles,” JHEP **2001** (2020) 046 doi:10.1007/JHEP01(2020)046 [arXiv:1906.10100 [hep-th]].
- [25] B. Maybee, D. O’Connell and J. Vines, “Observables and amplitudes for spinning particles and black holes,” JHEP **1912** (2019) 156 doi:10.1007/JHEP12(2019)156 [arXiv:1906.09260 [hep-th]].

- [26] N. Siemonsen and J. Vines, “Test black holes, scattering amplitudes and perturbations of Kerr spacetime,” arXiv:1909.07361 [gr-qc].
- [27] P. H. Damgaard, K. Haddad and A. Helset, “Heavy Black Hole Effective Theory,” JHEP **1911** (2019) 070 doi:10.1007/JHEP11(2019)070 [arXiv:1908.10308 [hep-ph]].
- [28] R. Aoude, K. Haddad and A. Helset, “On-shell heavy particle effective theories,” arXiv:2001.09164 [hep-th].
- [29] K. Haddad and A. Helset, “The double copy for heavy particles,” [arXiv:2005.13897 [hep-th]].
- [30] N. Moynihan, “Kerr-Newman from Minimal Coupling,” JHEP **01** (2020), 014 doi:10.1007/JHEP01(2020)014 [arXiv:1909.05217 [hep-th]].
- [31] M. Z. Chung, Y. T. Huang, J. W. Kim and S. Lee, “The simplest massive S-matrix: from minimal coupling to Black Holes,” JHEP **1904** (2019) 156 doi:10.1007/JHEP04(2019)156 [arXiv:1812.08752 [hep-th]].
- [32] Z. Bern, A. Luna, R. Roiban, C. H. Shen and M. Zeng, “Spinning Black Hole Binary Dynamics, Scattering Amplitudes and Effective Field Theory,” [arXiv:2005.03071 [hep-th]].
- [33] G. Kälin and R. A. Porto, “From boundary data to bound states. Part II. Scattering angle to dynamical invariants (with twist),” JHEP **02** (2020), 120 doi:10.1007/JHEP02(2020)120 [arXiv:1911.09130 [hep-th]].
- [34] M. Z. Chung, Y. T. Huang and J. W. Kim, “Classical potential for general spinning bodies,” [arXiv:1908.08463 [hep-th]].
- [35] M. Z. Chung, Y. T. Huang and J. W. Kim, “Kerr-Newman stress-tensor from minimal coupling to all orders in spin,” [arXiv:1911.12775 [hep-th]].
- [36] M. Z. Chung, Y. t. Huang, J. W. Kim and S. Lee, “Complete Hamiltonian for spinning binary systems at first post-Minkowskian order,” JHEP **05** (2020), 105 doi:10.1007/JHEP05(2020)105 [arXiv:2003.06600 [hep-th]].
- [37] C. Cheung and M. P. Solon, “Tidal Effects in the Post-Minkowskian Expansion,” [arXiv:2006.06665 [hep-th]].
- [38] D. Bini, T. Damour and A. Geralico, “Scattering of tidally interacting bodies in post-Minkowskian gravity,” Phys. Rev. D **101** (2020) no.4, 044039 doi:10.1103/PhysRevD.101.044039 [arXiv:2001.00352 [gr-qc]].
- [39] Z. Bern, J. Carrasco and H. Johansson, “New Relations for Gauge-Theory Amplitudes,” Phys. Rev. D **78** (2008), 085011 doi:10.1103/PhysRevD.78.085011 [arXiv:0805.3993 [hep-ph]].

- [40] Y. T. Huang, U. Kol and D. O’Connell, “The Double Copy of Electric-Magnetic Duality,” [arXiv:1911.06318 [hep-th]].
- [41] A. Luna, R. Monteiro, I. Nicholson and D. O’Connell, “Type D Spacetimes and the Weyl Double Copy,” *Class. Quant. Grav.* **36** (2019), 065003 doi:10.1088/1361-6382/ab03e6 [arXiv:1810.08183 [hep-th]].
- [42] A. Luna, R. Monteiro, I. Nicholson, D. O’Connell and C. D. White, “The double copy: Bremsstrahlung and accelerating black holes,” *JHEP* **06** (2016), 023 doi:10.1007/JHEP06(2016)023 [arXiv:1603.05737 [hep-th]].
- [43] R. Monteiro, D. O’Connell and C. D. White, “Black holes and the double copy,” *JHEP* **12** (2014), 056 doi:10.1007/JHEP12(2014)056 [arXiv:1410.0239 [hep-th]].
- [44] R. Monteiro, D. O’Connell and C. D. White, “Gravity as a double copy of gauge theory: from amplitudes to black holes,” *Int. J. Mod. Phys. D* **24** (2015) no.09, 1542008 doi:10.1142/S0218271815420080
- [45] A. Luna, R. Monteiro, D. O’Connell and C. D. White, “The classical double copy for Taub–NUT spacetime,” *Phys. Lett. B* **750** (2015), 272-277 doi:10.1016/j.physletb.2015.09.021 [arXiv:1507.01869 [hep-th]].
- [46] A. Luna, R. Monteiro, I. Nicholson, A. Ochirov, D. O’Connell, N. Westerberg and C. D. White, “Perturbative spacetimes from Yang-Mills theory,” *JHEP* **04** (2017), 069 doi:10.1007/JHEP04(2017)069 [arXiv:1611.07508 [hep-th]].
- [47] A. Luna, I. Nicholson, D. O’Connell and C. D. White, *JHEP* **03** (2018), 044 doi:10.1007/JHEP03(2018)044 [arXiv:1711.03901 [hep-th]].
- [48] J. J. M. Carrasco, L. Rodina, Z. Yin and S. Zekioglu, “Simple encoding of higher derivative gauge and gravity counterterms,” [arXiv:1910.12850 [hep-th]].
- [49] Z. Bern, J. J. Carrasco, M. Chiodaroli, H. Johansson and R. Roiban, “The Duality Between Color and Kinematics and its Applications,” [arXiv:1909.01358 [hep-th]].
- [50] Z. Bern, J. J. Carrasco, W. M. Chen, H. Johansson and R. Roiban, “Gravity Amplitudes as Generalized Double Copies of Gauge-Theory Amplitudes,” *Phys. Rev. Lett.* **118** (2017) no.18, 181602 doi:10.1103/PhysRevLett.118.181602 [arXiv:1701.02519 [hep-th]].
- [51] H. Johansson and A. Ochirov, “Double copy for massive quantum particles with spin,” *JHEP* **09** (2019), 040 doi:10.1007/JHEP09(2019)040 [arXiv:1906.12292 [hep-th]].
- [52] Y. F. Bautista and A. Guevara, “On the Double Copy for Spinning Matter,” [arXiv:1908.11349 [hep-th]].
- [53] W. D. Goldberger and J. Li, “Strings, extended objects, and the classical double copy,” *JHEP* **02** (2020), 092 doi:10.1007/JHEP02(2020)092 [arXiv:1912.01650 [hep-th]].

- [54] W. D. Goldberger, J. Li and S. G. Prabhu, “Spinning particles, axion radiation, and the classical double copy,” *Phys. Rev. D* **97** (2018) no.10, 105018 doi:10.1103/PhysRevD.97.105018 [arXiv:1712.09250 [hep-th]].
- [55] W. D. Goldberger and A. K. Ridgway, “Bound states and the classical double copy,” *Phys. Rev. D* **97** (2018) no.8, 085019 doi:10.1103/PhysRevD.97.085019 [arXiv:1711.09493 [hep-th]].
- [56] W. D. Goldberger, S. G. Prabhu and J. O. Thompson, “Classical gluon and graviton radiation from the bi-adjoint scalar double copy,” *Phys. Rev. D* **96** (2017) no.6, 065009 doi:10.1103/PhysRevD.96.065009 [arXiv:1705.09263 [hep-th]].
- [57] W. D. Goldberger and A. K. Ridgway, “Radiation and the classical double copy for color charges,” *Phys. Rev. D* **95** (2017) no.12, 125010 doi:10.1103/PhysRevD.95.125010 [arXiv:1611.03493 [hep-th]].
- [58] J. Plefka, C. Shi, J. Steinhoff and T. Wang, “Breakdown of the classical double copy for the effective action of dilaton-gravity at NNLO,” *Phys. Rev. D* **100** (2019) no.8, 086006 doi:10.1103/PhysRevD.100.086006 [arXiv:1906.05875 [hep-th]].
- [59] J. Plefka, C. Shi and T. Wang, “Double copy of massive scalar QCD,” *Phys. Rev. D* **101** (2020) no.6, 066004 doi:10.1103/PhysRevD.101.066004 [arXiv:1911.06785 [hep-th]].
- [60] J. Plefka, J. Steinhoff and W. Wormsbecher, *Phys. Rev. D* **99** (2019) no.2, 024021 doi:10.1103/PhysRevD.99.024021 [arXiv:1807.09859 [hep-th]].
- [61] T. Adamo and A. Ilderton, “Classical and quantum double copy of back-reaction,” [arXiv:2005.05807 [hep-th]].
- [62] A. Antonelli, A. Buonanno, J. Steinhoff, M. van de Meent and J. Vines, “Energetics of two-body Hamiltonians in post-Minkowskian gravity,” *Phys. Rev. D* **99** (2019) no.10, 104004 doi:10.1103/PhysRevD.99.104004 [arXiv:1901.07102 [gr-qc]].
- [63] A. Buonanno and T. Damour, *Phys. Rev. D* **59** (1999), 084006 doi:10.1103/PhysRevD.59.084006 [arXiv:gr-qc/9811091 [gr-qc]].
- [64] T. Damour, *Int. J. Mod. Phys. A* **23** (2008), 1130-1148 doi:10.1142/S0217751X08039992 [arXiv:0802.4047 [gr-qc]].
- [65] P. Aichelburg and R. Sexl, “On the Gravitational field of a massless particle,” *Gen. Rel. Grav.* **2** (1971), 303-312 doi:10.1007/BF00758149
- [66] R. Penrose, *General Relativity: papers in honour of J.L. Synge*, ed. L. O’Raifeartaigh (Oxford, 1972)
- [67] Bonnor, W. B., *Commun. Math. Phys.* **13** (1969) 163
- [68] G. ’t Hooft, “Graviton Dominance in Ultrahigh-Energy Scattering,” *Phys. Lett. B* **198** (1987) 61. doi:10.1016/0370-2693(87)90159-6

- [69] T. Dray and G. 't Hooft, “The Gravitational Shock Wave of a Massless Particle,” Nucl. Phys. B **253** (1985), 173-188 doi:10.1016/0550-3213(85)90525-5
- [70] D. N. Kabat and M. Ortiz, “Eikonal quantum gravity and Planckian scattering,” Nucl. Phys. B **388** (1992) 570 doi:10.1016/0550-3213(92)90627-N [hep-th/9203082].
- [71] M. Ortaggio, P. Krtous and J. Podolsky, “Ultrarelativistic boost of the black ring,” Phys. Rev. D **71** (2005), 124031 doi:10.1103/PhysRevD.71.124031 [arXiv:gr-qc/0503026 [gr-qc]].
- [72] M. Ortaggio, “Ultrarelativistic boost of spinning and charged black rings,” J. Phys. Conf. Ser. **33** (2006), 386-392 doi:10.1088/1742-6596/33/1/047 [arXiv:gr-qc/0601093 [gr-qc]].
- [73] M. Ortaggio and M. Astorino, “Ultrarelativistic boost of a black hole in the magnetic universe of Levi-Civita–Bertotti–Robinson,” Phys. Rev. D **97** (2018) no.10, 104052 doi:10.1103/PhysRevD.97.104052 [arXiv:1805.05382 [gr-qc]].
- [74] W. D. Goldberger and I. Z. Rothstein, “An Effective field theory of gravity for extended objects,” Phys. Rev. D **73** (2006), 104029 doi:10.1103/PhysRevD.73.104029 [arXiv:hep-th/0409156 [hep-th]].
- [75] A. K. Collado, P. Di Vecchia, R. Russo and S. Thomas, “The subleading eikonal in supergravity theories,” JHEP **10** (2018), 038 doi:10.1007/JHEP10(2018)038 [arXiv:1807.04588 [hep-th]].
- [76] M. Duff, Phys. Rev. D **7** (1973), 2317-2326 doi:10.1103/PhysRevD.7.2317
- [77] D. Neill and I. Z. Rothstein, “Classical Space-Times from the S Matrix,” Nucl. Phys. B **877** (2013), 177-189 doi:10.1016/j.nuclphysb.2013.09.007 [arXiv:1304.7263 [hep-th]].
- [78] N. E. J. Bjerrum-Bohr, J. F. Donoghue and B. R. Holstein, “Quantum corrections to the Schwarzschild and Kerr metrics,” Phys. Rev. D **68** (2003), 084005 doi:10.1103/PhysRevD.68.084005 [arXiv:hep-th/0211071 [hep-th]].
- [79] H. de Vega and N. G. Sanchez, “Particle Scattering at the Planck Scale and the Aichelburg-sexl Geometry,” Nucl. Phys. B **317** (1989), 731-756 doi:10.1016/0550-3213(89)90541-5
- [80] V. Ferrari, P. Pendenza and G. Veneziano ”Beam like gravitational waves and their geodesics”
- [81] R. Emparan and H. S. Reall, “Black Holes in Higher Dimensions,” Living Rev. Rel. **11** (2008), 6 doi:10.12942/lrr-2008-6 [arXiv:0801.3471 [hep-th]].
- [82] G. U. Jakobsen, “Schwarzschild-Tangherlini Metric from Scattering Amplitudes,” [arXiv:2006.01734 [hep-th]].

- [83] Valeria Ferrari, Paolo Pendenza, “Boosting the Kerr metric” *Gen Relat Gravit* **22**, 1105–1117 (1990).
- [84] E. Newman and A. Janis, *J. Math. Phys.* **6** (1965), 915-917 doi:10.1063/1.1704350
- [85] D. Amati, M. Ciafaloni and G. Veneziano, “Classical and Quantum Gravity Effects from Planckian Energy Superstring Collisions,” *Int. J. Mod. Phys. A* **3** (1988), 1615-1661 doi:10.1142/S0217751X88000710
- [86] D. Amati, M. Ciafaloni and G. Veneziano, “Higher Order Gravitational Deflection and Soft Bremsstrahlung in Planckian Energy Superstring Collisions,” *Nucl. Phys. B* **347** (1990), 550-580 doi:10.1016/0550-3213(90)90375-N
- [87] P. D’Eath, “Perturbation methods for interactions between strongly self-gravitating systems,” *Conf. Proc. C* **7606286** (1979), 249-288
- [88] S. Weinberg, “Gravitation and Cosmology,”
- [89] A. Petrov, “The Classification of spaces defining gravitational fields,” *Gen. Rel. Grav.* **32** (2000), 1661-1663 doi:10.1023/A:1001910908054
- [90] F. Pirani, “Invariant formulation of gravitational radiation theory,” *Phys. Rev.* **105** (1957), 1089-1099 doi:10.1103/PhysRev.105.1089
- [91] K. Lechner, “Electrodynamics of massless charged particles,” *J. Math. Phys.* **56** (2015) no.2, 022901 doi:10.1063/1.4906813 [arXiv:1405.4805 [hep-th]].
- [92] W. Bonnor, “Spinning null fluid in general relativity,” *Int. J. Theor. Phys.* **3** (1970), 257-266 doi:10.1007/BF00669753
- [93] V. P. Frolov, W. Israel and A. Zelnikov, “Gravitational field of relativistic gyratons,” *Phys. Rev. D* **72** (2005), 084031 doi:10.1103/PhysRevD.72.084031 [arXiv:hep-th/0506001 [hep-th]].
- [94] M. Maggiore, “A Modern introduction to quantum field theory,”
- [95] R. Steinbauer, “Geodesics and geodesic deviation for impulsive gravitational waves,” *J. Math. Phys.* **39** (1998), 2201-2212 doi:10.1063/1.532283 [arXiv:gr-qc/9710119 [gr-qc]].
- [96] A. Lecke, R. Steinbauer and R. Svarc, “The regularity of geodesics in impulsive pp-waves,” *Gen. Rel. Grav.* **46** (2014), 1648 doi:10.1007/s10714-013-1648-0 [arXiv:1310.1322 [gr-qc]].