

# General RGEs for dimensionful couplings in the $\overline{\text{MS}}$ scheme

Lohan Sartore\*

*Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes,  
CNRS/IN2P3, 53 Avenue des Martyrs, 38026 Grenoble, France*

Recently, Poole & Thomsen have presented the Renormalization Group Equations (RGEs) in the  $\overline{\text{MS}}$  scheme for dimensionless parameters of a general renormalizable gauge theory in a new formalism based on the Local Renormalization Group. In this Letter, we apply the dummy field method to the expressions in this formalism in order to derive the RGEs for the dimensionful couplings which so far have been missing in this approach. The complete set of RGEs has been implemented in a new version of the public code PyR@TE dedicated to the automatic computation of the RGEs for any general renormalizable (non-supersymmetric) gauge theory.

*Introduction.* — The knowledge of Renormalization Group Equations (RGEs) is crucial in the context of studies of gauge theories since they provide the necessary link between the couplings of the theory at different energy scales. The expressions for the dimensionless couplings RGEs of a general gauge theory were first presented almost forty years ago [1–6]. These results were later extended to the case of dimensionful parameters [7] using the so-called *dummy field method* [7, 8], which relates the  $\beta$ -functions of dimensionful parameters to those of dimensionless ones. A thorough presentation of the dummy field method is given in [9] along with a proper derivation of the general two-loop RGEs for dimensionful couplings in the  $\overline{\text{MS}}$  scheme, correcting some of the results presented in [7]. Using these general expressions by hand can be quite involved and error-prone. For this reason, they have been implemented in the Mathematica package SARAH [10] and the Python package PyR@TE [11, 12], which are both publicly available.

Recently, a new formalism was proposed [13] for the RGEs of general gauge theories that possesses many interesting features compared to the old Machacek & Vaughn formalism, both on the conceptual and practical sides. Within the framework of the Local Renormalization Group [14–16], relations can be derived that involve the coefficients appearing in the  $\beta$ -functions of gauge, Yukawa and quartic couplings at different loop orders: these are the so-called Weyl Consistency Conditions. Based on these relations and the existing results in the literature at two-loop, the authors of [13] were able to present for the first time the general expressions of the gauge coupling  $\beta$ -functions at the three-loop order. More recently, this formalism helped in the derivation of the 4-loop gauge  $\beta$ -functions in the Standard Model [17]. In the light of these successful developments, we believe that this formalism will allow in a near future further progress in the field of general RGEs.

Compared to the previous formalism, several new attractive features should be noted. First, the expressions of the  $\beta$ -functions are equally concise for gauge theories based either on a simple gauge group or a semi-simple

one. In particular, in the presence of multiple Abelian gauge factors, the effects of kinetic mixing [18, 19] are encompassed in the formalism in a systematic way. Another salient feature is the easy adaptability of the formalism to renormalization schemes other than  $\overline{\text{MS}}$ , the Weyl Consistency Conditions being scheme-independent relations. Finally, it was shown [13, 20] that the Weyl consistency conditions can help solve the ambiguities related to the treatment of  $\gamma_5$  in fermion loops starting to appear at the 3- and 4-loop order in Yukawa and gauge  $\beta$ -functions, respectively.

Currently, there is however one missing piece compared to the previous formalism, namely the expressions of the RGEs for all (renormalizable) dimensionful couplings. The aim of the present Letter is to bridge this gap providing general expressions for the  $\beta$ -functions of dimensionful couplings, obtained by properly applying the dummy field method to the formalism of Poole & Thomsen.

*Description of the formalism.* — We consider a general gauge theory containing an arbitrary set of real scalars  $\phi_a$  and Weyl fermions  $\psi_i$ . Following [13], we define the Majorana spinor

$$\Psi_i \equiv \begin{pmatrix} \psi \\ \psi^\dagger \end{pmatrix}_i. \quad (1)$$

With this definition, the most general Lagrangian density may be written (see [13] for details):

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4}G_{AB}^{-2}F_{\mu\nu}^A F^{B\mu\nu} + \frac{1}{2}(D_\mu\phi)_a(D^\mu\phi)_a \\ & + \frac{i}{2}\Psi^T \begin{pmatrix} 0 & \sigma^\mu \\ \bar{\sigma}^\mu & 0 \end{pmatrix} D_\mu\Psi \\ & - \frac{1}{2}y_{aij}\Psi_i\Psi_j\phi_a - \frac{1}{2}m_{ij}\Psi_i\Psi_j \\ & - \frac{1}{2}\mu_{ab}\phi_a\phi_b - \frac{1}{3!}t_{abc}\phi_a\phi_b\phi_c - \frac{1}{4!}\lambda_{abcd}\phi_a\phi_b\phi_c\phi_d, \end{aligned} \quad (2)$$

where  $G_{AB}$  is a matrix containing the gauge couplings associated to the (semi-)simple gauge group of the theory,

\* sartore@lpsc.in2p3.fr



The mapping (21) allows for a straightforward derivation of the individual contributions to the  $\beta$ -functions of the dimensional couplings, starting from the known expressions in the dimensionless case. For instance, the diagrams contributing to the fermion mass RGEs ( $\beta_{ij}^{(\ell)}$ ) are obtained from the individual contributions to the Yukawa couplings RGEs ( $\beta_{aij}^{(\ell)}$ ), when the external scalar field  $\phi_a$  is taken to be the dummy field  $\phi_{\hat{d}}$ . Consequently, one should in practice perform the following replacements:

$$a \rightarrow \hat{d}, \quad y_{aij} \rightarrow m_{ij}, \quad \lambda_{abcd} \rightarrow t_{abc}.$$

Similarly, starting from the quartic couplings RGEs, one may derive the  $\beta$ -functions for the trilinear and scalar mass couplings when one or two of the external scalar legs are replaced by a dummy field. This

procedure is applied on a diagrammatic basis, allowing in particular the identification of unphysical tadpole contributions which must be discarded [9].

*Results.* — We now turn to the presentation of the results. The various quantities appearing in the expressions of the  $\beta$ -functions are directly taken from [13]. Therefore, we invite the reader to refer to the definitions presented therein. We show below the whole set of RGEs obtained in the  $\overline{MS}$  scheme for the fermion mass, trilinear and scalar mass couplings. The following equations complete the list of RGEs presented in [13] for the dimensionless couplings. We use a notation where the fermion indices are made implicit. In this context,  $m$ ,  $y_a$ , and any other tensor carrying two fermion indices may be seen as matrices in the space of the fermions of the theory.

### Fermion mass $\beta$ -functions

At 1-loop:

$$\beta^{(1)} = \xi_1^{(1)} m C_2(F) + \xi_2^{(1)} y_b m y_b + \xi_3^{(1)} m \tilde{Y}_2(F). \quad (22)$$

At 2-loop:

$$\begin{aligned} \beta^{(2)} = & \xi_1^{(2)} \tilde{C}_2(F) m C_2(F) & + \xi_2^{(2)} m C_2(F) C_2(F) & + \xi_3^{(2)} m C_2(F, G) \\ & + \xi_4^{(2)} m C_2(F, S) & + \xi_5^{(2)} m C_2(F, F) & + \xi_6^{(2)} y_b T^A \tilde{y}_b m T^B G_{AB}^2 \\ & + \xi_7^{(2)} Y_2(F) \tilde{T}^A m T^B G_{AB}^2 & + \xi_8^{(2)} y_b \tilde{m} y_c [C_2(S)]_{bc} & + \xi_9^{(2)} y_b C_2(F) \tilde{m} y_b \\ & + \xi_{10}^{(2)} y_b \tilde{m} y_b C_2(F) & + \xi_{11}^{(2)} Y_2(F, C_S) m & + \xi_{12}^{(2)} Y_2(F, C_F) m \\ & + \xi_{13}^{(2)} m \tilde{Y}_2(F) C_2(F) & + \xi_{14}^{(2)} y_b \tilde{y}_c y_d t_{bcd} & + \xi_{15}^{(2)} y_b \tilde{y}_c m \tilde{y}_b y_c \\ & + \xi_{16}^{(2)} y_b \tilde{m} y_c \tilde{y}_b y_c & + \xi_{17}^{(2)} y_b \tilde{y}_c m \tilde{y}_c y_b & + \xi_{18}^{(2)} Y_4(F) m \\ & + \xi_{19}^{(2)} y_b \tilde{m} Y_2(F) y_b & + \xi_{20}^{(2)} m \tilde{Y}_2(F, Y_F) & + \xi_{21}^{(2)} y_b \tilde{m} y_c [Y_2(S)]_{bc} \\ & + \xi_{22}^{(2)} Y_2(F, Y_S) m. \end{aligned} \quad (23)$$

### Trilinear couplings $\beta$ -functions

At 1-loop:

$$\beta_{abc}^{(1)} = \tau_1^{(1)} [C_2(S)]_{ae} t_{ebc} + \tau_2^{(1)} \lambda_{abef} t_{efc} + \tau_3^{(1)} [Y_2(S)]_{ae} t_{ebc} + \tau_4^{(1)} \text{Tr} [m \tilde{y}_a y_b \tilde{y}_c]. \quad (24)$$

At 2-loop:

$$\begin{aligned} \beta_{abc}^{(2)} = & \tau_1^{(2)} (T_\phi^A T_\phi^C)_{ae} G_{AB}^2 G_{CD}^2 (T_\phi^B T_\phi^D)_{bf} t_{efc} & + \tau_2^{(2)} (T_\phi^A T_\phi^C)_{ab} G_{AB}^2 G_{CD}^2 (T_\phi^B T_\phi^D)_{ef} t_{efc} \\ & + \tau_3^{(2)} [C_2(S)]_{ae} [C_2(S)]_{bf} t_{efc} & + \tau_4^{(2)} [C_2(S)]_{ae} [C_2(S)]_{ef} t_{fbc} \\ & + \tau_5^{(2)} [C_2(S, G)]_{ae} t_{ebc} & + \tau_6^{(2)} [C_2(S, S)]_{ae} t_{ebc} \\ & + \tau_7^{(2)} [C_2(S, F)]_{ae} t_{ebc} & + \tau_8^{(2)} (T_\phi^A)_{ae} (T_\phi^B)_{bf} G_{AB}^2 \lambda_{efgh} t_{ghc} \\ & + \tau_9^{(2)} \lambda_{abef} [C_2(S)]_{fg} t_{egc} & + \tau_{10}^{(2)} [C_2(S)]_{ae} t_{efg} \lambda_{fgbc} \\ & + \tau_{11}^{(2)} [C_2(S)]_{ae} \lambda_{ebfg} t_{fgc} & + \tau_{12}^{(2)} \lambda_{aefg} \lambda_{efgh} t_{hbc} \\ & + \tau_{13}^{(2)} t_{aef} \lambda_{eghb} \lambda_{fghc} & + \tau_{14}^{(2)} \lambda_{abef} \lambda_{eghc} t_{fgh} \end{aligned}$$

$$\begin{aligned}
& +\tau_{15}^{(2)} \lambda_{abef} \lambda_{efgh} t_{ghc} & +\tau_{16}^{(2)} (T_\phi^A T_\phi^C)_{ab} G_{AB}^2 G_{CD}^2 \text{Tr} [T^D T^B \tilde{m} y_c] \\
& +\tau_{17}^{(2)} (T_\phi^A T_\phi^C)_{ab} G_{AB}^2 G_{CD}^2 \text{Tr} [T^D T^B \tilde{y}_c m] & +\tau_{18}^{(2)} [Y_2(S, C_F)]_{ae} t_{ebc} \\
& +\tau_{19}^{(2)} [C_2(S)]_{ae} [Y_2(S)]_{ef} t_{fbc} & +\tau_{20}^{(2)} \lambda_{abef} [Y_2(S)]_{fg} t_{egc} \\
& +\tau_{21}^{(2)} G_{AB}^2 \text{Tr} [m T^A \tilde{y}_a y_b T^B \tilde{y}_c] & +\tau_{22}^{(2)} G_{AB}^2 \text{Tr} [y_a T^A \tilde{m} y_b T^B \tilde{y}_c] \\
& +\tau_{23}^{(2)} [C_2(S)]_{ae} \text{Tr} [y_e \tilde{m} y_b \tilde{y}_c] & +\tau_{24}^{(2)} [C_2(S)]_{ae} \text{Tr} [y_e \tilde{y}_b m \tilde{y}_c] \\
& +\tau_{25}^{(2)} \text{Tr} [m \tilde{y}_a y_b \tilde{y}_c \tilde{C}_2(F)] & +\tau_{26}^{(2)} \text{Tr} [y_a m y_b \tilde{y}_c \tilde{C}_2(F)] \\
& +\tau_{27}^{(2)} \text{Tr} [m \tilde{y}_e y_a \tilde{y}_f] \lambda_{efbc} & +\tau_{28}^{(2)} \text{Tr} [y_a \tilde{y}_e y_b \tilde{y}_f] t_{efc} \\
& +\tau_{29}^{(2)} \text{Tr} [m \tilde{y}_a y_e \tilde{y}_f] \lambda_{efbc} & +\tau_{30}^{(2)} \text{Tr} [y_a \tilde{y}_b y_e \tilde{y}_f] t_{efc} \\
& +\tau_{31}^{(2)} [Y_4(S)]_{ae} t_{ebc} & +\tau_{32}^{(2)} [Y_2(S, Y_F)]_{ae} t_{ebc} \\
& +\tau_{33}^{(2)} \text{Tr} [m \tilde{y}_a y_b \tilde{y}_e y_c \tilde{y}_e] & +\tau_{34}^{(2)} \text{Tr} [y_a \tilde{m} y_b \tilde{y}_e y_c \tilde{y}_e] \\
& +\tau_{35}^{(2)} \text{Tr} [y_a \tilde{y}_b m \tilde{y}_e y_c \tilde{y}_e] & +\tau_{36}^{(2)} \text{Tr} [m \tilde{y}_a y_e \tilde{y}_b y_c \tilde{y}_e] \\
& +\tau_{37}^{(2)} \text{Tr} [m \tilde{y}_a y_b \tilde{y}_c Y_2(F)] & +\tau_{38}^{(2)} \text{Tr} [y_a \tilde{m} y_b \tilde{y}_c Y_2(F)] .
\end{aligned} \tag{25}$$

### Scalar mass $\beta$ -functions

At 1-loop:

$$\begin{aligned}
\beta_{ab}^{(1)} = & \sigma_1^{(1)} [C_2(S)]_{ae} \mu_{eb} & + \sigma_2^{(1)} \lambda_{abef} \mu_{ef} & + \sigma_3^{(1)} t_{aef} t_{efb} \\
& + \sigma_4^{(1)} [Y_2(S)]_{ae} \mu_{eb} & + \sigma_5^{(1)} \text{Tr} [m \tilde{m} y_a \tilde{y}_b] & + \sigma_6^{(1)} \text{Tr} [m \tilde{y}_a m \tilde{y}_b] .
\end{aligned} \tag{26}$$

At 2-loop:

$$\begin{aligned}
\beta_{ab}^{(2)} = & \sigma_1^{(2)} (T_\phi^A T_\phi^C)_{ae} G_{AB}^2 G_{CD}^2 (T_\phi^B T_\phi^D)_{bf} \mu_{ef} & + \sigma_2^{(2)} (T_\phi^A T_\phi^C)_{ab} G_{AB}^2 G_{CD}^2 (T_\phi^B T_\phi^D)_{ef} \mu_{ef} \\
& + \sigma_3^{(2)} [C_2(S)]_{ae} [C_2(S)]_{bf} \mu_{ef} & + \sigma_4^{(2)} [C_2(S)]_{ae} [C_2(S)]_{ef} \mu_{fb} \\
& + \sigma_5^{(2)} [C_2(S, G)]_{ae} \mu_{eb} & + \sigma_6^{(2)} [C_2(S, S)]_{ae} \mu_{eb} \\
& + \sigma_7^{(2)} [C_2(S, F)]_{ae} \mu_{eb} & + \sigma_8^{(2)} (T_\phi^A)_{ae} (T_\phi^B)_{bf} G_{AB}^2 \lambda_{efgh} \mu_{gh} \\
& + \sigma_9^{(2)} \lambda_{abef} [C_2(S)]_{fg} \mu_{eg} & + \sigma_{10}^{(2)} t_{aef} [C_2(S)]_{fg} t_{egb} \\
& + \sigma_{11}^{(2)} [C_2(S)]_{ae} \lambda_{ebfg} \mu_{fg} & + \sigma_{12}^{(2)} [C_2(S)]_{ae} t_{efg} t_{fgb} \\
& + \sigma_{13}^{(2)} \lambda_{aefg} \lambda_{efgh} \mu_{hb} & + \sigma_{14}^{(2)} \lambda_{aegh} \lambda_{bfg} \mu_{ef} \\
& + \sigma_{15}^{(2)} \lambda_{abef} t_{egh} t_{fg} & + \sigma_{16}^{(2)} t_{aef} t_{egh} \lambda_{bfg} \\
& + \sigma_{17}^{(2)} \lambda_{abef} \lambda_{efgh} \mu_{gh} & + \sigma_{18}^{(2)} t_{aef} \lambda_{efgh} t_{ghb} \\
& + \sigma_{19}^{(2)} (T_\phi^A T_\phi^C)_{ab} G_{AB}^2 G_{CD}^2 \text{Tr} [T^D T^B \tilde{m} m] & + \sigma_{20}^{(2)} [Y_2(S, C_F)]_{ae} \mu_{eb} \\
& + \sigma_{21}^{(2)} [C_2(S)]_{ae} [Y_2(S)]_{ef} \mu_{fb} & + \sigma_{22}^{(2)} \lambda_{abef} [Y_2(S)]_{fg} \mu_{eg} \\
& + \sigma_{23}^{(2)} t_{aef} [Y_2(S)]_{fg} t_{egb} & + \sigma_{24}^{(2)} G_{AB}^2 \text{Tr} [y_a T^A \tilde{m} m T^B \tilde{y}_b] \\
& + \sigma_{25}^{(2)} G_{AB}^2 \text{Tr} [y_a T^A \tilde{y}_b m T^B \tilde{m}] & + \sigma_{26}^{(2)} G_{AB}^2 \text{Tr} [y_a T^A \tilde{m} y_b T^B \tilde{m}] \\
& + \sigma_{27}^{(2)} [C_2(S)]_{ae} \text{Tr} [y_e \tilde{m} y_b m] & + \sigma_{28}^{(2)} [C_2(S)]_{ae} \text{Tr} [y_e \tilde{m} m y_b] \\
& + \sigma_{29}^{(2)} \text{Tr} [y_a \tilde{y}_b m \tilde{m} \tilde{C}_2(F)] & + \sigma_{30}^{(2)} \text{Tr} [y_a \tilde{m} y_b \tilde{m} \tilde{C}_2(F)] \\
& + \sigma_{31}^{(2)} \text{Tr} [y_a \tilde{m} m \tilde{y}_b \tilde{C}_2(F)] & + \sigma_{32}^{(2)} \text{Tr} [m \tilde{y}_a y_b \tilde{m} \tilde{C}_2(F)] \\
& + \sigma_{33}^{(2)} \text{Tr} [y_a \tilde{y}_e y_b \tilde{y}_f] \mu_{ef} & + \sigma_{34}^{(2)} \text{Tr} [y_a \tilde{y}_e m \tilde{y}_f] t_{efb} \\
& + \sigma_{35}^{(2)} \text{Tr} [m \tilde{y}_e m \tilde{y}_f] \lambda_{efab} & + \sigma_{36}^{(2)} \text{Tr} [y_a \tilde{y}_b y_e \tilde{y}_f] \mu_{ef}
\end{aligned}$$

$$\begin{aligned}
& +\sigma_{37}^{(2)} \text{Tr} [y_a \tilde{m} y_e \tilde{y}_f] t_{efb} & +\sigma_{38}^{(2)} \text{Tr} [m \tilde{m} y_e \tilde{y}_f] \lambda_{efab} \\
& +\sigma_{39}^{(2)} [Y_4(S)]_{ae} \mu_{eb} & +\sigma_{40}^{(2)} [Y_2(S, Y_F)]_{ae} \mu_{eb} \\
& +\sigma_{41}^{(2)} \text{Tr} [m \tilde{y}_a m \tilde{y}_e y_b \tilde{y}_e] & +\sigma_{42}^{(2)} \text{Tr} [y_a \tilde{y}_b m \tilde{y}_e m \tilde{y}_e] \\
& +\sigma_{43}^{(2)} \text{Tr} [y_a \tilde{m} m \tilde{y}_e y_b \tilde{y}_e] & +\sigma_{44}^{(2)} \text{Tr} [y_a \tilde{m} y_b \tilde{y}_e m \tilde{y}_e] \\
& +\sigma_{45}^{(2)} \text{Tr} [y_a \tilde{y}_b y_e \tilde{m} m \tilde{y}_e] & +\sigma_{46}^{(2)} \text{Tr} [y_a \tilde{m} y_e \tilde{m} y_b \tilde{y}_e] \\
& +\sigma_{47}^{(2)} \text{Tr} [y_a \tilde{m} y_e \tilde{y}_b m \tilde{y}_e] & +\sigma_{48}^{(2)} \text{Tr} [y_a \tilde{y}_b m \tilde{m} Y_2(F)] \\
& +\sigma_{49}^{(2)} \text{Tr} [y_a \tilde{m} y_b \tilde{m} Y_2(F)] & +\sigma_{50}^{(2)} \text{Tr} [y_a \tilde{m} m \tilde{y}_b Y_2(F)] \\
& +\sigma_{51}^{(2)} \text{Tr} [m \tilde{y}_a y_b \tilde{m} Y_2(F)] . & 
\end{aligned} \tag{27}$$

### Fermion mass coefficients

At 1-loop:

$$\xi_1^{(1)} = -6, \quad \xi_2^{(1)} = 2, \quad \xi_3^{(1)} = 1. \tag{28}$$

At 2-loop:

$$\begin{aligned}
\xi_1^{(2)} &= 0, & \xi_2^{(2)} &= -3, & \xi_3^{(2)} &= -\frac{97}{3}, & \xi_4^{(2)} &= \frac{11}{6}, & \xi_5^{(2)} &= \frac{5}{3}, & \xi_6^{(2)} &= 12, \\
\xi_7^{(2)} &= 0, & \xi_8^{(2)} &= 6, & \xi_9^{(2)} &= 10, & \xi_{10}^{(2)} &= 6, & \xi_{11}^{(2)} &= 9, & \xi_{12}^{(2)} &= -\frac{1}{2}, \\
\xi_{13}^{(2)} &= -\frac{7}{2}, & \xi_{14}^{(2)} &= -2, & \xi_{15}^{(2)} &= 2, & \xi_{16}^{(2)} &= 0, & \xi_{17}^{(2)} &= -2, & \xi_{18}^{(2)} &= 0, \\
\xi_{19}^{(2)} &= -2, & \xi_{20}^{(2)} &= -\frac{1}{4}, & \xi_{21}^{(2)} &= -1, & \xi_{22}^{(2)} &= -\frac{3}{4}. & & & & 
\end{aligned} \tag{29}$$

### Trilinear coefficients

At 1-loop:

$$\tau_1^{(1)} = -9, \quad \tau_2^{(1)} = 3, \quad \tau_3^{(1)} = \frac{3}{2}, \quad \tau_4^{(1)} = -12. \tag{30}$$

At 2-loop:

$$\begin{aligned}
\tau_1^{(2)} &= 6, & \tau_2^{(2)} &= 30, & \tau_3^{(2)} &= 0, & \tau_4^{(2)} &= \frac{9}{2}, & \tau_5^{(2)} &= -\frac{143}{4}, & \tau_6^{(2)} &= \frac{11}{4}, \\
\tau_7^{(2)} &= \frac{10}{4}, & \tau_8^{(2)} &= -9, & \tau_9^{(2)} &= 24, & \tau_{10}^{(2)} &= -\frac{9}{2}, & \tau_{11}^{(2)} &= -9, & \tau_{12}^{(2)} &= \frac{1}{4}, \\
\tau_{13}^{(2)} &= -3, & \tau_{14}^{(2)} &= -3, & \tau_{15}^{(2)} &= 0, & \tau_{16}^{(2)} &= -36, & \tau_{17}^{(2)} &= -36, & \tau_{18}^{(2)} &= \frac{15}{2}, \\
\tau_{19}^{(2)} &= 0, & \tau_{20}^{(2)} &= -3, & \tau_{21}^{(2)} &= 0, & \tau_{22}^{(2)} &= 0, & \tau_{23}^{(2)} &= 12, & \tau_{24}^{(2)} &= 6, \\
\tau_{25}^{(2)} &= -24, & \tau_{26}^{(2)} &= -24, & \tau_{27}^{(2)} &= 6, & \tau_{28}^{(2)} &= 6, & \tau_{29}^{(2)} &= 0, & \tau_{30}^{(2)} &= 0, \\
\tau_{31}^{(2)} &= -\frac{3}{2}, & \tau_{32}^{(2)} &= -\frac{9}{4}, & \tau_{33}^{(2)} &= 24, & \tau_{34}^{(2)} &= 12, & \tau_{35}^{(2)} &= 12, & \tau_{36}^{(2)} &= 24, \\
\tau_{37}^{(2)} &= 12, & \tau_{38}^{(2)} &= 12. & & & & & & & & 
\end{aligned} \tag{31}$$

### Scalar mass coefficients

At 1-loop:

$$\sigma_1^{(1)} = -6, \quad \sigma_2^{(1)} = 1, \quad \sigma_3^{(1)} = 1, \quad \sigma_4^{(1)} = 1, \quad \sigma_5^{(1)} = -4, \quad \sigma_6^{(1)} = -2. \quad (32)$$

At 2-loop:

$$\begin{aligned} \sigma_1^{(2)} &= 2, & \sigma_2^{(2)} &= 10, & \sigma_3^{(2)} &= 0, & \sigma_4^{(2)} &= 3, & \sigma_5^{(2)} &= -\frac{143}{6}, & \sigma_6^{(2)} &= \frac{11}{6}, \\ \sigma_7^{(2)} &= \frac{10}{6}, & \sigma_8^{(2)} &= -3, & \sigma_9^{(2)} &= 8, & \sigma_{10}^{(2)} &= 8, & \sigma_{11}^{(2)} &= -3, & \sigma_{12}^{(2)} &= -3, \\ \sigma_{13}^{(2)} &= \frac{1}{6}, & \sigma_{14}^{(2)} &= -1, & \sigma_{15}^{(2)} &= -\frac{1}{2}, & \sigma_{16}^{(2)} &= -2, & \sigma_{17}^{(2)} &= 0, & \sigma_{18}^{(2)} &= 0, \\ \sigma_{19}^{(2)} &= -12, & \sigma_{20}^{(2)} &= 5, & \sigma_{21}^{(2)} &= 0, & \sigma_{22}^{(2)} &= -1, & \sigma_{23}^{(2)} &= -1, & \sigma_{24}^{(2)} &= 0, \\ \sigma_{25}^{(2)} &= 0, & \sigma_{26}^{(2)} &= 0, & \sigma_{27}^{(2)} &= 2, & \sigma_{28}^{(2)} &= 4, & \sigma_{29}^{(2)} &= -8, & \sigma_{30}^{(2)} &= -8, \\ \sigma_{31}^{(2)} &= -4, & \sigma_{32}^{(2)} &= -4, & \sigma_{33}^{(2)} &= 2, & \sigma_{34}^{(2)} &= 4, & \sigma_{35}^{(2)} &= 1, & \sigma_{36}^{(2)} &= 0, \\ \sigma_{37}^{(2)} &= 0, & \sigma_{38}^{(2)} &= 0, & \sigma_{39}^{(2)} &= -1, & \sigma_{40}^{(2)} &= -\frac{3}{2}, & \sigma_{41}^{(2)} &= 4, & \sigma_{42}^{(2)} &= 8, \\ \sigma_{43}^{(2)} &= 8, & \sigma_{44}^{(2)} &= 4, & \sigma_{45}^{(2)} &= 4, & \sigma_{46}^{(2)} &= 4, & \sigma_{47}^{(2)} &= 4, & \sigma_{48}^{(2)} &= 4, \\ \sigma_{49}^{(2)} &= 4, & \sigma_{50}^{(2)} &= 2, & \sigma_{51}^{(2)} &= 2. \end{aligned} \quad (33)$$

*Conclusions.* — We have presented in this Letter the general expressions of the two-loop  $\overline{\text{MS}}$   $\beta$ -functions for dimensionful parameters in the formalism of Poole & Thomsen. This completes the set of dimensionless RGEs given in [13] up to the respective loop orders 3-2-2 for gauge, Yukawa and quartic couplings. The full set of RGEs in this new formalism has been implemented in a new version of the PyR@TE software, PyR@TE 3 [21, 22]. We note finally that the procedure presented in this Letter would remain valid using different renormalization schemes, both in the non-supersymmetric and supersymmetric case.

---

## ACKNOWLEDGMENTS

We are grateful to Colin Poole and Anders Eller Thomsen for many useful discussions. We would also like to thank Ingo Schienbein for his comments and support in the preparation of this Letter.

This work was supported in part by the IN2P3 project “Théorie – BSMGA”.

- 
- [1] M. E. Machacek and M. T. Vaughn, Nuclear Physics B **222**, 83 (1983).
  - [2] M. E. Machacek and M. T. Vaughn, Nuclear Physics B **236**, 221 (1984).
  - [3] M. E. Machacek and M. T. Vaughn, Nuclear Physics B **249**, 70 (1985).
  - [4] I. Jack and H. Osborn, Nuclear Physics B **207**, 474 (1982).
  - [5] I. Jack and H. Osborn, Journal of Physics A: Mathematical and General **16**, 1101 (1983).
  - [6] I. Jack and H. Osborn, Nuclear Physics B **249**, 472 (1985).
  - [7] M. Luo, H. Wang, and Y. Xiao, Physical Review D **67**, 065019 (2003).
  - [8] S. P. Martin and M. T. Vaughn, Phys. Rev. D **50**, 2282 (1994).
  - [9] I. Schienbein, F. Staub, T. Steudtner, and K. Svirina, Nuclear Physics B **939**, 1 (2019).
  - [10] F. Staub, Computer Physics Communications **185**, 1773.
  - [11] F. Lyonnet, I. Schienbein, F. Staub, and A. Wingerter, Computer Physics Communications **185**, 1130 (2014).
  - [12] F. Lyonnet and I. Schienbein, Computer Physics Communications **213**, 181 (2017).
  - [13] C. Poole and A. E. Thomsen, Journal of High Energy Physics **2019**, 10.1007/JHEP09(2019)055 (2019).
  - [14] H. Osborn, Physics Letters B **222**, 97 (1989).
  - [15] I. Jack and H. Osborn, Nuclear Physics B **343**, 647 (1990).
  - [16] H. Osborn, Nuclear Physics B **363**, 486 (1991).
  - [17] J. Davies, F. Herren, C. Poole, M. Steinhauser, and A. E. Thomsen, Physical Review Letters **124**, 071803 (2020).
  - [18] M. Luo and Y. Xiao, Physics Letters B **555**, 279 (2003).
  - [19] R. M. Fonseca, M. Malinský, and F. Staub, Physics Letters B **726**, 882 (2013).
  - [20] C. Poole and A. E. Thomsen, Physical Review Letters **123**, 041602 (2019).
  - [21] L. Sartore and I. Schienbein, Computer Physics Communications **261**, 107819 (2021).

[22] <https://github.com/lsartore/pyrate>.